CHARACTERIZATION OF COMPLEMENTING PAIRS OF $(\mathbb{Z}_{\geq 0})^n$

HUI RAO, YA-MIN YANG, AND YUAN ZHANG*

ABSTRACT. Let A, B, C be subsets of an abelian group G. A pair (A, B) is called a Cpair if $A, B \subset C$ and C is the direct sum of A and B. The $(\mathbb{Z}_{\geq 0})$ -pairs are characterized by de Bruijn in 1950 and the $(\mathbb{Z}_{\geq 0})^2$ -pairs are characterized by Niven in 1971. In this paper, we characterize the $(\mathbb{Z}_{\geq 0})^n$ -pairs for all $n \geq 1$. We show that every $(\mathbb{Z}_{\geq 0})^n$ -pair is characterized by a weighted tree if it is primitive, that is, it is not a Cartesian product of a $(\mathbb{Z}_{\geq 0})^p$ -pair and a $(\mathbb{Z}_{\geq 0})^q$ -pair of lower dimensions.

1. Introduction

Let \mathbb{Z} be the set of integers. Let $A, B, C \subset \mathbb{Z}^n$. We define $A+B = \{a+b; a \in A, b \in B\}$. We call A + B the *direct sum* of A and B and denoted by $A \oplus B$, if every element $c \in A + B$ has a unique decomposition as c = a + b with $a \in A, b \in B$. We say (A, B) is a *complementing pair* of C, or a C-pair, if $C = A \oplus B$ and $A, B \subset C$.

We are interested in \mathbb{Z}^n -pairs and \mathbb{N}^n -pairs, where we denote $\mathbb{N} = \mathbb{Z}_{\geq 0}$ for simplicity. Furthermore, a \mathbb{Z}^n -pair (A, B) is called a \mathbb{Z}^n -tiling if $\#A < \infty$, where #A denotes the cardinality of A; in this case, we call A a \mathbb{Z}^n -tile. Similarly, we define \mathbb{N}^n -tiling and \mathbb{N}^n -tile.

The characterization of \mathbb{Z} -pairs is an important problem in additive number theory. This problem is first considered by de Bruijn [2] in 1950. In 1974, Swenson [13] showed that this problem is an *NP*-hard problem. There are very few results on this problem.

To determine when a finite set $A \subset \mathbb{Z}$ is a \mathbb{Z} -tile is somehow easier. This was done by Newman [9] when #A is a power of a prime number, see also Tijdeman [15]. Coven and Meyerowitz [1] solved the problem when #A contains at most two prime factors, see also Sands [11]. If #A contains more than two prime factors, the problem is still widely open [14, 1]. There are almost no results on \mathbb{Z}^n -pairs with $n \geq 2$, except that Szegedy [12] characterized the \mathbb{Z}^d -tile T in case that #T is a prime number or #T = 4.

Date: September 22, 2020.

The work is supported by NSFS Nos. 11971195 and 11601172.

²⁰⁰⁰ Mathematics Subject Classification: 11P83, 52C22

Key words and phrases: complementing pair, power series, weighted tree.

^{*} The correspondence author.

The characterization of N-pairs is much easier and it was settled by de Bruijn [3], and was rediscovered by Vaidya [16]. Let $(n_k)_{k\geq 1}$ be a sequence of integers with $n_k \geq 2$. Any $t \in \mathbb{N}$ can be written uniquely in the form

(1.1)
$$t = d_1 + n_1 d_2 + (n_1 n_2) d_3 + \dots + (n_1 \cdots n_{L-1}) d_L,$$

where $d_j \in \{0, 1, \dots, n_j - 1\}$ for each $1 \leq j \leq L$ and $d_L > 0$. We denote the right hand side of (1.1) by $\overline{d_1 \dots d_L}$.

Proposition 1.1. (de Bruijn [3], Vaidya [16]) (i) A pair (T, \mathcal{J}) is an \mathbb{N} -pair with $\#T = \#\mathcal{J} = \infty$ if and only if there exists a sequence of integers $(n_k)_{k\geq 1}$ with $n_k \geq 2$ such that

(1.2)
$$T = \bigcup_{L=1}^{\infty} \{\overline{d_1 \dots d_L}; \ d_j = 0 \ if \ j \ is \ odd\}, \ \mathcal{J} = \bigcup_{L=1}^{\infty} \{\overline{d_1 \dots d_L}; \ d_j = 0 \ if \ j \ is \ even\},$$

or the other round.

(ii) T is an \mathbb{N} -tile if and only if there exist $L \geq 2$ and $(n_k)_{k=1}^L$ with $n_k \geq 2$ such that

$$T = \{\overline{d_1 \dots d_L}; \ d_j = 0 \ if j \ is \ odd\} \ or \ T = \{\overline{d_1 \dots d_L}; \ d_j = 0 \ if j \ is \ even\}$$

Remark 1.2. Let (T, \mathcal{J}) be an N-pair with $\#T = \#\mathcal{J} = \infty$. Long [7] made the interesting observation that $(T, -\mathcal{J})$ is a Z-pair; Eigen and Hajian [4] showed that there exists a continuum number of sets $\tilde{\mathcal{J}}$ such that $(T, \tilde{\mathcal{J}})$ is a Z-pair.

After a partial result was obtained by Hansen [5], Niven [10] characterized the \mathbb{N}^2 -pairs (See Example 1.1 in the end of this section).

The goal of the present paper is to characterize the \mathbb{N}^n -pairs for all $n \ge 1$. For simplicity, we call (T, \mathcal{J}) an \mathbb{N}^* -pair if (T, \mathcal{J}) is an \mathbb{N}^n -pair for some $n \ge 1$.

It is easy to show that if (I_1, J_1) is an \mathbb{N}^n -pair and (I_2, J_2) is an \mathbb{N}^m -pair, then $(T, \mathcal{J}) = (I_1 \times I_2, J_1 \times J_2)$ is an \mathbb{N}^{n+m} -pair, and we call (T, \mathcal{J}) the Cartesian product of the two pairs. (See Lemma 2.1.)

An \mathbb{N}^n -pair (T, \mathcal{J}) is called *primitive*, if it is not a Cartesian product of two \mathbb{N}^* -pairs with lower dimensions. All \mathbb{N} -pairs are primitive. Clearly, to characterize \mathbb{N}^* -pairs, we only need understand primitive \mathbb{N}^* -pairs.

In literature, polynomials have been used to study direct sum decomposition of abelian groups ([6, 11, 1]) or finite sets ([8]). The first idea in this paper is to use power series to handle \mathbb{N}^* -pairs. A power series with finite many variables is called a *binary power series* if its coefficients belong to {0,1}. Let $x = (x_1, \ldots, x_n)$ be an *n*-tuple variable. For $\boldsymbol{a} = (a_1, \dots, a_n) \in \mathbb{N}^n$, we denote $x^{\boldsymbol{a}} = x_1^{a_1} \cdots x_n^{a_n}$. For $A \subset \mathbb{N}^n$, we define

(1.3)
$$A(x) = \sum_{a \in A} x^a.$$

For example $\mathbb{N}(x) = \sum_{k \ge 0} x^k$ and $\mathbb{N}^2(x, y) = \mathbb{N}(x)\mathbb{N}(y)$. Clearly, (T, \mathcal{J}) is a \mathbb{N}^n -pair if and only if $T(x)\mathcal{J}(x) = \mathbb{N}^n(x)$.

We call (C, D) an *interval pair* if it is a $\{0, 1, ..., N-1\}$ -pair for some integer $N \ge 2$, and we call N the *size* of the pair (C, D).

Remark 1.3. Nathanson [8] proved that if $T \oplus S = \prod_{j=1}^{n} \{0, 1, \dots, a_j\}$ is a higher dimensional interval pair, then (T, S) is a cartesian product of one-dimensional interval pairs.

1.1. Extension process. Now, we introduce two ways to produce \mathbb{N}^* -pairs from a known \mathbb{N}^n -pair (T, \mathcal{J}) . Let $j \in \{1, \ldots, n\}$. Denote $x = (\bar{x}, x_j, \bar{x})$, where $\bar{x} = (x_1, \ldots, x_{j-1})$ and $\bar{x} = (x_{j+1}, \ldots, x_n)$.

(1) Extension of the first type. Let $N \ge 2$ and (C, D) be a $\{0, 1, \ldots, N-1\}$ -pair. Set

(1.4)
$$T^*(x) = C(x_j)T(\bar{x}, x_j^N, \bar{\bar{x}}), \quad \mathcal{J}^*(x) = D(x_j)\mathcal{J}(\bar{x}, x_j^N, \bar{\bar{x}}).$$

Then (T^*, \mathcal{J}^*) is an \mathbb{N}^n -pair, and we call it a first type extension of (T, \mathcal{J}) . (See Lemma 3.1.)

(2) Extension of the second type. Let $\delta \in \{0,1\}, m \ge 2$, and $a \in (\mathbb{N} \setminus \{0\})^m$. Set

$$L_{\boldsymbol{a}} = \mathbb{N}^m \setminus (\mathbb{N}^m + \boldsymbol{a}).$$

Denote $y = (y_1, ..., y_m)$, and define $L_a(y)$ as we did in (1.3). (For example, if a = (1, 1), then $L_a(y_1, y_2) = 1 + \sum_{k \ge 1} y_1^k + \sum_{k \ge 1} y_2^k$.) Set

(1.5)
$$T^*(\bar{x}, y, \bar{\bar{x}}) = L^{\delta}_{\boldsymbol{a}}(y)T(\bar{x}, y^{\boldsymbol{a}}, \bar{\bar{x}}), \quad \mathcal{J}^*(\bar{x}, y, \bar{\bar{x}}) = L^{1-\delta}_{\boldsymbol{a}}(y)\mathcal{J}(\bar{x}, y^{\boldsymbol{a}}, \bar{\bar{x}}).$$

Then (T^*, \mathcal{J}^*) is an \mathbb{N}^{n+m-1} -pair, and we call it a second type extension of (T, \mathcal{J}) . (See Lemma 3.2.)

Let $(T_j, \mathcal{J}_j)_{j=0}^k$ be a sequence of \mathbb{N}^* -pairs. If (T_j, \mathcal{J}_j) is a first type or second type extension of $(T_{j-1}, \mathcal{J}_{j-1})$ for $j = 1, \ldots, k$, then we say (T_k, \mathcal{J}_k) is a *finite extension* of (T_0, \mathcal{J}_0) .

An extension of an \mathbb{N}^* -pair (T, \mathcal{J}) is called an *illegal extension*, if it is a second type extension in (1.5) satisfying $(T, \mathcal{J}) = (\{0\}, \mathbb{N})$ and $\delta = 0$, or $(T, \mathcal{J}) = (\mathbb{N}, \{0\})$ and $\delta = 1$. An extension changes the primitivity if and only if it is illegal (Theorem 3.1). Our first main result is the following. **Theorem 1.1.** An \mathbb{N}^* -pair (T, \mathcal{J}) is primitive if and only if it is a finite extension of an \mathbb{N} -pair, and the extension process contains no illegal extension.

A finite extension of an \mathbb{N} -pair is primitive is proved in Section 3. The other direction of Theorem 1.1 is proved in Section 4-8, where Lemma 5.1 and Theorem 6.1 contain new techniques and new phenomena which are different from dimension one and two.

1.2. Weighted tree of an \mathbb{N}^* -pair. For a primitive \mathbb{N}^* -pair, we introduce a weighted tree to record the information of the extension process in Theorem 1.1. Indeed, a weighted tree provides a visualization of the structure of the associated \mathbb{N}^* -pair.

Let (V, Γ) be a finite tree with a root ϕ , where V is the node set and Γ is the edge set. A node is called a *top* if it has no offspring. Let V_0 denote the set of tops of (V, Γ) . In Section 9, we define a *weight* of (V, Γ) to be a quadruple

(1.6)
$$((T_0, \mathcal{J}_0), \{\boldsymbol{\delta}_v\}_{v \in V \setminus V_0}, \{\boldsymbol{\alpha}_v\}_{v \in V \setminus \{\phi\}}, \{(C_v, D_v)\}_{v \in V \setminus \{\phi\}}),$$

where (T_0, \mathcal{J}_0) is an N-pair and we call it the *initial pair*. We associate an N^{*}-pair to a weighted tree, and we call it the N^{*}-pair generated by the weighted tree. We show that

Theorem 1.2. An \mathbb{N}^* -pair (T, \mathcal{J}) is a primitive \mathbb{N}^n -pair if and only if (T, \mathcal{J}) is generated by a weighted tree with n tops.

Remark 1.4. We show that (T, \mathcal{J}) is an \mathbb{N}^* -tiling if and only if in the associated weighted tree, the initial pair is an \mathbb{N} -tiling with $\#T_0 < \infty$ and the map δ is constantly zero, or the other round (see Remark 9.2). It is folklore that an \mathbb{N}^n -tile is also a \mathbb{Z}^n -tile, so our construction may shed some light to the study of \mathbb{Z}^n -tiles with $n \geq 2$, an area is almost untouched.

In Theorem 9.1 we give an explicit formula for \mathbb{N}^* -pairs generated by weighted trees. (In two dimensional case, the formula is given by (1.7).) Actually, for each $v \in V$, we define two sets $P_v, Q_v \subset \mathbb{N}^n$, and we show that $T = \bigoplus_{v \in V} P_v$ and $\mathcal{J} = \bigoplus_{v \in V} Q_v$, which give further factorizations of T and \mathcal{J} .

Example 1.1. Niven's characterization of the \mathbb{N}^2 -pairs is the case n = 2 of Theorem 1.1. In the following we describe an extension process to generate primitive \mathbb{N}^2 -pairs.

Let (T_0, \mathcal{J}_0) be a non-trivial \mathbb{N} -pair, $\delta \in \{0, 1\}$, $\boldsymbol{a} = (a_1, a_2) \in (\mathbb{N} \setminus \{0\})^2$, and let $\Phi_x = (C_1, D_1)$ and $\Phi_y = (C_2, D_2)$ be two interval pairs with size N_1 and N_2 , respectively. Moreover, $\delta = 1$ if $T_0 = \{0\}$ and $\delta = 0$ if $\mathcal{J}_0 = \{0\}$. Regarding ϕ as a variable, we have $T_0(\phi)\mathcal{J}_0(\phi) = \mathbb{N}(\phi)$. Applying a second type extension by setting $\phi = x^{a_1}y^{a_2}$, we obtain an \mathbb{N}^2 -pair

$$T_1(x,y) = L_{\boldsymbol{a}}^{\delta}(x,y)T_0(x^{a_1}y^{a_2}), \quad \mathcal{J}_1(x,y) = L_{\boldsymbol{a}}^{1-\delta}(x,y)\mathcal{J}_0(x^{a_1}y^{a_2}).$$

Applying two first type extensions to (T_1, \mathcal{J}_1) , we obtain

(1.7)
$$T(x,y) = C_1(x)C_2(y)L_{\boldsymbol{a}}^{\delta}(x^{N_1}, y^{N_2})T_0(x^{a_1N_1}y^{a_2N_2}),$$
$$\mathcal{J}(x,y) = D_1(x)D_2(y)L_{\boldsymbol{a}}^{1-\delta}(x^{N_1}, y^{N_2})\mathcal{J}_0(x^{a_1N_1}y^{a_2N_2}).$$

As a corollary of Theorem 1.2, a pair (T, \mathcal{J}) is a primitive \mathbb{N}^2 -pair if and only if it is given by (1.7).

The paper is organized as follows. In Section 2, we prove several simple lemmas. Section 3 is devoted to extensions of \mathbb{N}^* -pairs. Section 4–Section 8 investigate the reductions of primitive \mathbb{N}^* -pairs; Theorem 1.1 is proved in Section 8. In section 9, we introduce weighted trees. Theorem 1.2 is proved in Section 10.

2. Uniqueness and primitivity

We denote $\mathbf{0}_m = (0, \ldots, 0) \in \mathbb{Z}^m$; if the dimension is implicit, then we just write **0**. Let \prec_q be the *lexicographical order* on \mathbb{Z}^n .

Lemma 2.1. (i) (Uniqueness) If (T, \mathcal{J}) and (T, \mathcal{J}') are two \mathbb{N}^n -pairs, then $\mathcal{J} = \mathcal{J}'$.

(ii) If $T = I_1 \times I_2$ and $\mathcal{J} = J_1 \times J_2$ where (I_1, J_1) is an \mathbb{N}^n -pair and (I_2, J_2) is an \mathbb{N}^m -pair, then (T, \mathcal{J}) is an \mathbb{N}^{n+m} -pair.

Proof. (i) Let us list \mathcal{J} as $(t_k)_{k\geq 1}$ by the following rule: for $s, t \in \mathcal{J}$, we list s before t if |s| < |t|; if |s| = |t|, we list s before t if $s \prec_g t$. We list \mathcal{J}' as $(t'_k)_{k\geq 1}$ by the same rule.

Clearly $t_1 = t'_1 = \mathbf{0}$. Since $T \oplus (\mathcal{J} \setminus \{t_1\}) = T \oplus (\mathcal{J}' \setminus \{t_1\})$, the minimum elements (according to the above rule) of $\mathcal{J} \setminus \{t_1\}$ and $\mathcal{J}' \setminus \{t'_1\}$ coincide, so $t_2 = t'_2$. Continue this procedure, we obtain $\mathcal{J} = \mathcal{J}'$.

(ii) Denote $x = (x_1, \ldots, x_n), y = (y_1, \ldots, y_m)$. We have $(I_1 \times I_2)(x, y) \cdot (J_1 \times J_2)(x, y) = I_1(x)J_1(x)I_2(y)J_2(y) = \mathbb{N}^n(x)\mathbb{N}^m(y) = \mathbb{N}^{n+m}(x, y)$, which proves the second assertion. \Box

Actually (T, \mathcal{J}) is non-primitive if one of T and \mathcal{J} is a Cartesian product.

Lemma 2.2. If (T, \mathcal{J}) is an \mathbb{N}^{n+m} -pair such that $T = I_1 \times I_2 \subset \mathbb{N}^n \times \mathbb{N}^m$, then \mathcal{J} can be written as $J_1 \times J_2 \subset \mathbb{N}^n \times \mathbb{N}^m$.

Proof. Denote $x = (x_1, \ldots, x_n), y = (y_1, \ldots, y_m)$. The assumptions imply that

$$I_1(x)I_2(y)\mathcal{J}(x,y) = \mathbb{N}^{n+m}(x,y).$$

Setting $y = \mathbf{0}_m$, we obtain $I_1(x)\mathcal{J}(x,\mathbf{0}_m) = \mathbb{N}^n(x)$. Let $J_1(x) = \mathcal{J}(x,\mathbf{0}_m)$, then (I_1,J_1) is an \mathbb{N}^n -pair. Similarly, let $J_2(y) = \mathcal{J}(\mathbf{0}_n, y)$, then (I_2, J_2) is an \mathbb{N}^m -pair. By Lemma 2.1(ii), $(I_1 \times I_2, J_1 \times J_2)$ is an \mathbb{N}^{n+m} -pair. Therefore $\mathcal{J} = J_1 \times J_2$ by Lemma 2.1(i). \Box

We close this section with several results about \mathbb{N} -pairs which will be needed later.

Lemma 2.3. ([3, 11]) If (T, \mathcal{J}) is an \mathbb{N} -pair, then there exists an integer $p \geq 2$, and an \mathbb{N} -pair (A, B) such that T = pA, $\mathcal{J} = \{0, 1, \dots, p-1\} \oplus pB$ or the other round.

Lemma 2.4. ([3, 11, 15]) If (C, D) is a $\{0, 1, \ldots, N-1\}$ -pair with $N \ge 2$, then there exists $p \ge 2$ and a $\{0, 1, \ldots, \frac{N}{p} - 1\}$ -pair $(\widetilde{C}, \widetilde{D})$ such that $C = p\widetilde{C}, D = p\widetilde{D} + \{0, 1, \ldots, p-1\}$ or the other round.

We denote $C_k \uparrow T$ if $C_k \subset C_{k+1}$ and $T = \bigcup_{k \ge 1} C_k$. The following lemma is a easy consequence of Proposition 1.1. We leave the simple proof to the reader.

Lemma 2.5. Let (T, \mathcal{J}) be an \mathbb{N} -pair. If $\#T = \#\mathcal{J} = \infty$, then there exists a sequence of interval pairs $\{(C_k, D_k)\}_{k\geq 1}$ such that $C_k \uparrow T$ and $D_k \uparrow \mathcal{J}$, and

$$T = C_k \oplus N_k A_k, \quad \mathcal{J} = D_k \oplus N_k B_k$$

for some $A_k, B_k \subset \mathbb{N}$, where N_k is the size of (C_k, D_k) . If $\#T < \infty$, then there exist an integer $N \ge 1$ and $D \subset \mathbb{N}$ such that $T \oplus D = \{0, 1, \dots, N-1\}$ and $\mathcal{J} = D \oplus N\mathbb{N}$.

3. Extensions of \mathbb{N}^n -pairs

In this section, we prove that a finite extension of an \mathbb{N} -pair is a primitive \mathbb{N}^* -pair.

Let $a \in \mathbb{N}^m$ with a > 0, recall that $L_a = \mathbb{N}^m \setminus (\mathbb{N}^m + a)$. Take $z \in \mathbb{N}^m$, and let k be the largest integer such that $z - ka \in \mathbb{N}^m$, then $z - ka \in L_a$. Hence

$$(3.1) L_{\boldsymbol{a}} \oplus \mathbb{N}\boldsymbol{a} = \mathbb{N}^m.$$

Especially, $L_p = \{0, 1, \dots, p-1\}$. Denote $x = (x_1, \dots, x_n)$ and $\bar{x} = (x_2, \dots, x_n)$.

Lemma 3.1. The pair (T^*, \mathcal{J}^*) defined in (1.4) is an \mathbb{N}^n -pair.

Proof. Without loss of generality, we assume j = 1 in (1.4). We have

$$T^*(x)\mathcal{J}^*(x) = C(x_1)D(x_1)T(x_1^N,\bar{x})\mathcal{J}(x_1^N,\bar{x})$$

= $(1 + x_1 + \dots + x_1^{N-1})\mathbb{N}(x_1^N)\mathbb{N}^{n-1}(\bar{x}) = \mathbb{N}^n(x).$

The lemma is proved.

Lemma 3.2. The pair (T^*, \mathcal{J}^*) defined in (1.5) is an \mathbb{N}^{n+m-1} -pair.

Proof. Without loss of generality, we assume j = 1 in (1.5). We have

$$T^*(y,\bar{x})\mathcal{J}^*(y,\bar{x}) = L_{\boldsymbol{a}}(y)T(y^{\boldsymbol{a}},\bar{x})\mathcal{J}(y^{\boldsymbol{a}},\bar{x}) = L_{\boldsymbol{a}}(y)\sum_{k\geq 0} y^{k\boldsymbol{a}}\mathbb{N}^{n-1}(\bar{x})$$
$$= \mathbb{N}^m(y)\mathbb{N}^{n-1}(\bar{x}) = \mathbb{N}^{n+m-1}(y,\bar{x}).$$

(The third equality is due to (3.1).) The lemma is proved.

In the rest of this section, we investigate the primitivity property. We say a binary power series $T(x_1, \ldots, x_n)$ with $n \ge 2$ is variable separable, if there exist an integer m < n, a permutation τ on $\{2, \ldots, n\}$, and two sets $F \in \mathbb{N}^m$, $G \in \mathbb{N}^{n-m}$ such that

(3.2)
$$T(x_1, \dots, x_n) = F(x_1, x_{\tau(2)}, \dots, x_{\tau(m)}) G(x_{\tau(m+1)}, \dots, x_{\tau(n)}).$$

The following lemma is a direct consequence of Lemma 2.2.

Lemma 3.3. An \mathbb{N}^n -pair (T, \mathcal{J}) with $n \ge 2$ is primitive if and only if T(x) is not variable separable.

By convention, for $T \subset \mathbb{N}$, we say T(x) is variable separable if and only if #T = 1.

Lemma 3.4. Let $T \subset \mathbb{N}^n$. The following three statements are equivalent to each other: (i) $T(x_1, x_2, \ldots, x_n)$ is variable separable.

- (ii) $T(yz, x_2, \ldots, x_n)$ is variable separable.
- (iii) $T(x_1^m, x_2, \ldots, x_n)$ is variable separable for some $m \ge 2$.

Proof. (i) implies (ii) and (i) implies (iii) are trivial. In the following, we prove the other direction implications.

(ii) \Rightarrow (i). Let us first consider that case n = 1. If T(yz) is variable separable, then T(yz) = F(y)G(z). Set z = 1, we obtain T(y) = F(y)G(1). Since both T(y) and F(y) are binary, we obtain that G(1) = 1. Similarly, we have F(1) = 1. Hence T(1) = 1, which means #T = 1 and T(x) is variable separable.

Now we assume $n \ge 2$. Suppose $T(yz, x_2, \ldots, x_n)$ is variable separable, then there is a permutation τ on $\{2, \ldots, n\}$, and two binary power series F and G such that either

$$T(yz, x_2, \dots, x_n) = F(y, x_{\tau(2)}, \dots, x_{\tau(k)})G(z, x_{\tau(k+1)}, \dots, x_{\tau(n)})$$
 with $k \le n$, or

$$T(yz, x_2, \dots, x_n) = F(y, z, x_{\tau(2)}, \dots, x_{\tau(k)})G(x_{\tau(k+1)}, \dots, x_{\tau(n)})$$
 with $k < n$.

Set y = 1, $z = x_1$, we obtain a variable separable factorization of $T(x_1, \ldots, x_n)$.

(iii) \Rightarrow (i). If n = 1, the implication is trivial. Now we assume that $n \ge 2$. Suppose $T(x_1^m, x_2, \ldots, x_n)$ is variable separable, then we have

(3.3)
$$T(x_1^m, x_2, \dots, x_n) = F(x_1, x_{\tau(2)}, \dots, x_{\tau(k)}) G(x_{\tau(k+1)}, \dots, x_{\tau(n)})$$

where k < n. Since the power of x_1 of any term in F must be a multiple of m, we infer that there exists $F^* \in \mathbb{N}^n$ such that

(3.4)
$$F(x_1, x_{\tau(2)}, \dots, x_{\tau(k)}) = F^*(x_1^m, x_{\tau(2)}, \dots, x_{\tau(k)}).$$

Substituting (3.4) into (3.3), and then replacing x_1^m by x_1 , we obtain a variable separable factorization of $T(x_1, \ldots, x_n)$.

Theorem 3.1. Let (T^*, \mathcal{J}^*) be an extension of (T, \mathcal{J}) . If the extension is not illegal, then (T^*, \mathcal{J}^*) is primitive if and only if (T, \mathcal{J}) is primitive.

Proof. (i) Let (T^*, \mathcal{J}^*) be a first type extension of (T, \mathcal{J}) satisfying $T^*(x) = C(x_1)T(x_1^N, \bar{x})$. If n = 1, then both (T, \mathcal{J}) and (T^*, \mathcal{J}^*) are primitive. Now we assume $n \ge 2$. By Lemma 3.4, we have

 (T^*, \mathcal{J}^*) is non-primitive $\Leftrightarrow T^*(x)$ is variable separable

 $\Leftrightarrow T(x_1^N, \bar{x}) = T^*(x)/C(x_1) \text{ is variable separable}$ $\Leftrightarrow T(x) \text{ is variable separable } \Leftrightarrow (T, \mathcal{J}) \text{ is non-primitive.}$

The theorem holds in this case.

(ii) Let (T^*, \mathcal{J}^*) be a second type extension of (T, \mathcal{J}) with parameters δ and a. Let us assume $\delta = 0$, then $T^*(y, \bar{x}) = T(y^a, \bar{x})$, where $y = (y_1, \ldots, y_m)$. Similar as above, (T^*, \mathcal{J}^*) is non-primitive is equivalent to T(x) is variable separable. If n > 1, it is equivalent to (T, \mathcal{J}) is non-primitive; if n = 1, it is equivalent to $(T, \mathcal{J}) = (\{0\}, \mathbb{N})$, which means the extension is illegal. The theorem is proved.

4. \mathbb{N}^* -pairs of pure type

Let $\mathbf{e}_1, \ldots, \mathbf{e}_n$ be the canonical basis of \mathbb{Z}^n . Let $n \geq 2$, set

(4.1)
$$L_0 = \{(a_1, \dots, a_n) \in \mathbb{N}^n; \text{ at least one } a_j \text{ is } 0\}$$

to be the union of (n-1)-faces of \mathbb{N}^n . An \mathbb{N}^n -pair (T, \mathcal{J}) with $n \geq 2$ is said to be of *pure* type, if either $L_0 \subset T$ or $L_0 \subset \mathcal{J}$. In this section, we characterize \mathbb{N}^* -pairs of pure type.

Let $a, b \in \mathbb{N}^n$. We say $a \leq b$ if $b - a \in \mathbb{N}^n$, and a < b if $b - a \in (\mathbb{N} \setminus \{0\})^n$.

Lemma 4.1. If (T, \mathcal{J}) is an \mathbb{N}^n -pair such that $L_0 \subset \mathcal{J}$, then < induces a linear order on T. We call a the germ of T, where a is the minimum of $T \setminus \{\mathbf{0}\}$ with respect to <.

Proof. Denote $a \lor b = \max\{a, b\}$, and $(a_1, \ldots, a_n) \lor (b_1, \ldots, b_n) = (a_1 \lor b_1, \ldots, a_n \lor b_n)$.

Let $\mathbf{a}, \mathbf{b} \in T$. Since $L_0 \subset \mathcal{J}, \mathbf{a} + L_0$ and $\mathbf{b} + L_0$ do not overlap. If neither $\mathbf{a} - \mathbf{b} > 0$ nor $\mathbf{b} - \mathbf{a} > 0$, then $\mathbf{c} := (\mathbf{a} - \mathbf{b}) \lor \mathbf{0} \in L_0$ and $\mathbf{c}' := (\mathbf{b} - \mathbf{a}) \lor \mathbf{0} \in L_0$. It follows that $\mathbf{b} + \mathbf{c} = \mathbf{a} + \mathbf{c}'$, a contradiction.

Lemma 4.2. Let (T, \mathcal{J}) be an \mathbb{N}^n -pair with $L_0 \subset \mathcal{J}$, and let a be the germ of T. Then (i) $L_a \subset \mathcal{J}$; (ii) for every integer $k \ge 0$, $(ka + L_a) \cap T$ contains at most one element.

Proof. (i) is obvious. Now we prove (ii). Suppose on the contrary $u, u' \in (ka + L_a) \cap T$. We assume that u' < u (Lemma 4.1). Then $u - u' \in L_a$ since $0 \le u - u' \le u - ka$. So u = (u) + 0 and u = (u') + (u - u') are two different (T, \mathcal{J}) -decompositions of u, which is a contradiction.

The following result is an extension of Lemma 2.3 (the case n = 1) and Niven [10, Lemma 5] (the case n = 2) to higher dimensions.

Theorem 4.1. Let $n \ge 2$, let (T, \mathcal{J}) be an \mathbb{N}^n -pair of pure type with $L_0 \subset \mathcal{J}$, and let **a** be the germ of T. Then there exists an \mathbb{N} -pair (A, B) such that

$$T = Aa, \quad \mathcal{J} = Ba \oplus L_a.$$

Proof. For $k \ge 0$, denote $\Omega_k = k\boldsymbol{a} + L_{\boldsymbol{a}}$, and we call it the k-th section of \mathbb{N}^n . For every integer $k \ge 0$, we claim that

(i) $T \cap \Omega_k = \{ka\}$ or \emptyset ; (ii) $\mathcal{J} \cap \Omega_k = \Omega_k$ or \emptyset .

By Lemma 4.2, $\mathbf{0}, \mathbf{a} \in T$ and $L_{\mathbf{a}} \subset \mathcal{J}$, so the claim holds for k = 0. Suppose the claim is false and let $k \ge 1$ be the smallest integer such that (i) or (ii) fails.

First, by the minimality of k, we have

$$T \cap \bigcup_{j=0}^{k-1} \Omega_j = U \boldsymbol{a}, \quad \mathcal{J} \cap \bigcup_{j=0}^{k-1} \Omega_j = V \boldsymbol{a} \oplus L_{\boldsymbol{a}}$$

where $U, V \subset \{0, 1, \dots, k-1\}$. Secondly, we assert that

$$k \boldsymbol{a} \notin T$$
 and $k \notin U + V$.

Suppose $ka \in T$, then $(ka) + (\mathbf{b})$, $\mathbf{b} \in L_a$, are the (T, \mathcal{J}) -decomposition of elements in $ka + L_a$, and this forces that (i) and (ii) hold for k, a contradiction. By the same reason, we have that $k \notin U + V$.

Let $C = T \cap \Omega_k$ and $D = \mathcal{J} \cap \Omega_k$. Then C contains at most one element. The set Ω_k is covered by

$$(T \cap \bigcup_{j=0}^{k} \Omega_j)) \oplus (\mathcal{J} \cap \bigcup_{j=0}^{k} \Omega_j) = (U\boldsymbol{a} \cup C) \oplus ((V\boldsymbol{a} + L_{\boldsymbol{a}}) \cup D).$$

Eliminating the elements apparently outside of Ω_k , we have that Ω_k is covered by $D \cup (C \oplus L_a) \cup ((U+V)a + L_a)$. Since $k \notin U + V$, we finally obtain

(4.2)
$$\Omega_k \subset (C \oplus L_a) \cup D,$$

which implies that both C and D are not empty.

Let us denote $C = \{\mathbf{c}\}$. Since $\mathbf{c} \neq k\mathbf{a}$, there exists *i* such that $\mathbf{c} - \mathbf{e}_i \in \Omega_k$, so $\mathbf{c} - \mathbf{e}_i \in D$ by (4.2). Since $\mathbf{a} \in T$ and $\mathbf{a} - \mathbf{e}_i \in \mathcal{J}$,

$$\boldsymbol{a} + (\mathbf{c} - \mathbf{e}_i) = \mathbf{c} + (\boldsymbol{a} - \mathbf{e}_i)$$

provides two (T, \mathcal{J}) -decompositions of a same point. This contradiction proves our claim.

Our claim implies that T = Aa and $\mathcal{J} = Ba + L_a$ for some $A, B \subset \mathbb{N}$. From $Aa \oplus Ba \oplus L_a = T \oplus \mathcal{J} = \mathbb{N}a \oplus L_a$, we deduce that (A, B) is an \mathbb{N} -pair. The theorem is proved. \Box

5. A key Lemma

In this section we prove a crucial lemma which will be used in Section 6.

Let $n \ge 1$ and write $z = (z_1, \ldots, z_n)$.

Lemma 5.1. Let a > 0 be a point in \mathbb{N}^n . Let $A_0, B_0 \subset \mathbb{N}$ with $0, 1 \in A_0$ and $0 \in B_0$, and denote $F_0 = A_0 a, G_0 = B_0 a \oplus L_a$. Let $\Omega \subset \mathbb{N}a$. If F_1, G_1 are two subsets of \mathbb{N}^n such that

(5.1)
$$F_0(z)G_1(z) + F_1(z)G_0(z) = L_a(z)\Omega(z),$$

then $F_1 = A_1 a$ and $G_1 = B_1 a \oplus L_a$ for some $A_1, B_1 \subset \mathbb{N}$.

Proof. The assumption (5.1) implies that

$$(F_0 \oplus G_1) \cup (F_1 \oplus G_0) = \Omega \oplus L_a$$
 and $(F_0 \oplus G_1) \cap (F_1 \oplus G_0) = \emptyset$

Denote $\Omega = (d_k a)_{k \ge 1}$, where $(d_k)_{k \ge 1}$ is an increasing sequence in \mathbb{N} . We denote $\Omega_k := d_k a + L_a$ and call it the k-th section of $\Omega + L_a$. We claim that:

Claim 1. For all $k \ge 0$, $G_1 \cap \Omega_k = \emptyset$ or Ω_k .

We prove the claim by induction. Clearly the claim holds for k = 1. Let $k \ge 2$ and assume that $G_1 \cap \Omega_j = \emptyset$ or Ω_j holds for j < k. Then

(5.2)
$$G_1 \cap (\Omega_1 \cup \cdots \cup \Omega_{k-1}) = V \boldsymbol{a} + L_{\boldsymbol{a}},$$

where $V \subset \{d_1, d_2, ..., d_{k-1}\}$. Hence, for j < k,

(5.3)
$$(F_0 \oplus G_1) \cap \Omega_j = (F_0 \oplus (\{g_0, \dots, g_h\}\boldsymbol{a} + L_{\boldsymbol{a}})) \cap \Omega_j = \emptyset \text{ or } \Omega_j.$$

Consequently, for j < k, as the complement of the right hand side of (5.3),

(5.4)
$$(F_1 \oplus G_0) \cap \Omega_j = \emptyset \text{ or } \Omega_j.$$

Let f^* be the smallest element in F_1 such that $f^* \notin \mathbb{N}a$. By (5.3) and (5.4),

(5.5)
$$f^* \notin \Omega_1 \cup \dots \cup \Omega_{k-1} \text{ (if } f^* \text{ exists)}.$$

Suppose that Claim 1 is false for k. Denote $C := G_1 \cap \Omega_k$. We assert that

$$(5.6) (F_0 \oplus G_1) \cap \Omega_k = C.$$

Notice that

$$(F_0 \oplus G_1) \cap \Omega_k = [(F_0 \oplus (\{g_0, \dots, g_h\}\boldsymbol{a} + L_{\boldsymbol{a}})) \cap \Omega_k] \cup [(F_0 \oplus C) \cap \Omega_k].$$

The second term on the right hand side is C, so the first term must be the empty set, which implies (5.6).

By (5.6) and (5.4), we have that $(F_1 \oplus G_0) \cap \Omega_k \neq \emptyset$, so there exists $f \in F_1$ such that $(f + G_0) \cap \Omega_k \neq \emptyset$. The intersection is not Ω_k implies that $f \notin \mathbb{N}a$, so we have $f \in \Omega_k$ since $f \geq f^*$. Since $F_1 \cap \Omega_k$ contains at most one element, we conclude that $f = f^*$.

Let *i* be the index in $\{1, \ldots, n\}$ such that $f - \mathbf{e}_i \in \Omega_k$. Then $f - \mathbf{e}_i$ is not covered by $f + G_0$, and hence it is not covered by $F_1 + G_0$. It follows that $f - \mathbf{e}_i \in C$. Therefore, on one hand, we have

$$\boldsymbol{a} + (f - \mathbf{e}_i) \in F_0 + C \subset F_0 + G_1;$$

on the other hand, $f + (\mathbf{a} - \mathbf{e}_i) \in F_1 + L_{\mathbf{a}} \subset F_1 + G_0$. This is a contradiction, and Claim 1 is proved. Therefore, (5.2) and (5.5) holds for all k, which imply that $G_1 = B_1 \mathbf{a} \oplus L_{\mathbf{a}}$, $F_1 = A_1 \mathbf{a}$ for some $A_1, B_1 \subset \mathbb{N}$. The lemma is proved.

6. Marginal pairs of \mathbb{N}^* -pairs

Let $j_1, \ldots, j_m (1 \le m < n)$ be a subsequence of $1, \ldots, n$. We call $Q = \mathbb{N}\mathbf{e}_{j_1} \oplus \cdots \oplus \mathbb{N}\mathbf{e}_{j_m}$ an *m*-face of \mathbb{N}^n . Define $\rho : Q \to \mathbb{N}^m$ as $\rho(\sum_{\ell=1}^m c_\ell \mathbf{e}_{j_\ell}) = (c_1, \ldots, c_m)$. We denote

(6.1)
$$T_Q = T \cap Q, \quad \mathcal{J}_Q = \mathcal{J} \cap Q$$

and call $(\rho(T_Q), \rho(\mathcal{J}_Q))$ the marginal pair induced by Q.

Lemma 6.1. Let (T, \mathcal{J}) be an \mathbb{N}^n -pair and Q be an m-face of \mathbb{N}^n . Then (T_Q, \mathcal{J}_Q) is a Q-pair. Consequently, the marginal pair $(\rho(T_Q), \rho(\mathcal{J}_Q))$ is an \mathbb{N}^m -pair.

Proof. In $T(x)\mathcal{J}(x) = \mathbb{N}^n(x)$, setting $x_k = 0$ if $k \notin \{j_1, \ldots, j_m\}$, we obtain $T_Q(x)\mathcal{J}_Q(x) = Q(x)$, which implies that $T_Q \oplus \mathcal{J}_Q = Q$. The second assertion is obvious. \Box

If (T, \mathcal{J}) is an N-pair with $\{0, 1, \dots, p-1\} \subset \mathcal{J}$ and $p \in T$, we call p the germ of T.

Theorem 6.1. Let (T, \mathcal{J}) be an \mathbb{N}^n -pair and $Q = \mathbb{N}\mathbf{e}_1 \oplus \cdots \oplus \mathbb{N}\mathbf{e}_m$ with $1 \leq m \leq n$. If the marginal pair $(\rho(T_Q), \rho(\mathcal{J}_Q))$ is of pure type in case of m > 1, then there exists an \mathbb{N}^{n-m+1} -pair $(\widetilde{T}, \widetilde{\mathcal{J}})$ such that

(6.2)
$$T(x) = \widetilde{T}(x_1^{a_1} \cdots x_m^{a_m}, x_{m+1}, \dots, x_n),$$
$$\mathcal{J}(x) = L_{\boldsymbol{a}}(x_1, \dots, x_m) \widetilde{\mathcal{J}}(x_1^{a_1} \cdots x_m^{a_m}, x_{m+1}, \dots, x_n).$$

where $\mathbf{a} = (a_1, \ldots, a_m)$ is the germ of $\rho(T_Q)$.

We note that (6.2) is a second type extension if m > 1, and is of first type if m = 1. Moreover, if m = n, then Theorem 6.1 becomes Theorem 4.1 or Lemma 2.3.

Proof. By Theorem 4.1 or Lemma 2.3, there exists an N-pair (A, B) with $1 \in A$ such that

(6.3)
$$\rho(T_Q) = A\boldsymbol{a}, \quad \rho(\mathcal{J}_Q) = B\boldsymbol{a} \oplus L_{\boldsymbol{a}},$$

(Remember that in case of $m = 1, L_{\boldsymbol{a}} = \{0, 1, \dots, \boldsymbol{a} - 1\}$.) Let $\mathbf{m} \in \mathbb{N}^{n-m}$. Define

$$T_{\mathbf{m}} = \{ u \in \mathbb{N}^m; \ (u, \mathbf{m}) \in T \}, \quad \mathcal{J}_{\mathbf{m}} = \{ v \in \mathbb{N}^m; \ (v, \mathbf{m}) \in \mathcal{J} \}.$$

Denote $\overline{\overline{x}} = (x_{m+1}, \ldots, x_n)$, then

(6.4)
$$T(x) = \sum_{\mathbf{p} \in \mathbb{N}^{n-m}} T_{\mathbf{p}}(x_1, \dots, x_m) \bar{x}^{\mathbf{p}}, \quad \mathcal{J}(x) = \sum_{\mathbf{q} \in \mathbb{N}^{n-m}} \mathcal{J}_{\mathbf{q}}(x_1, \dots, x_m) \bar{x}^{\mathbf{q}}.$$

We claim that for all $\mathbf{m} \in \mathbb{N}^{n-m}$, it holds that

(6.5)
$$\mathbb{N}^m(x_1,\ldots,x_m) = \sum_{\mathbf{p}+\mathbf{q}=\mathbf{m}} T_{\mathbf{p}}(x_1,\ldots,x_m) \mathcal{J}_{\mathbf{q}}(x_1,\ldots,x_m).$$

Clearly $\mathbb{N}^n(x) = \mathbb{N}^m(x_1, \dots, x_m) \sum_{\boldsymbol{d} \in \mathbb{N}^{n-m}} \bar{\bar{x}}^{\boldsymbol{d}}$. On the other hand, by (6.4),

$$\mathbb{N}^{n}(x) = T(x)\mathcal{J}(x) = \sum_{\boldsymbol{d}\in\mathbb{N}^{n-m}}\sum_{\mathbf{p}+\mathbf{q}=\boldsymbol{d}}T_{\mathbf{p}}(x_{1},\ldots,x_{m})\mathcal{J}_{\mathbf{q}}(x_{1},\ldots,x_{m})\bar{x}^{\boldsymbol{d}}.$$

Comparing the terms in the above two equations with the factor $\bar{x}^{\mathbf{m}}$, we obtain (6.5).

Denote $\bar{x} = (x_1, \ldots, x_m)$. We shall prove by induction that

(6.6)
$$T_{\mathbf{m}}(\bar{x}) = f_{\mathbf{m}}(\bar{x}^{a}), \quad \mathcal{J}_{\mathbf{m}}(\bar{x}) = L_{a}(\bar{x})G_{\mathbf{m}}(\bar{x}^{a}),$$

for some $f_{\mathbf{m}}, G_{\mathbf{m}} \subset \mathbb{N}$.

For $\mathbf{m} = \mathbf{0}$, $(T_{\mathbf{0}}, \mathcal{J}_{\mathbf{0}}) = (\rho(T_Q), \rho(\mathcal{J}_Q))$. In this case (6.6) holds by (6.3).

Suppose (6.6) holds if we replace \mathbf{m} by any point $\mathbf{m}' \leq \mathbf{m}$ and $\mathbf{m}' \neq \mathbf{m}$. We are going to show that (6.6) holds for \mathbf{m} .

If $\mathbf{p} + \mathbf{q} = \mathbf{m}$ and $\mathbf{p} \neq \mathbf{0}$, $\mathbf{q} \neq \mathbf{0}$, then (6.6) holds for \mathbf{p} and \mathbf{q} , so

(6.7)
$$T_{\mathbf{p}}(\bar{x})\mathcal{J}_{\mathbf{q}}(\bar{x}) = L_{\boldsymbol{a}}(\bar{x})f_{\mathbf{p}}(\bar{x}^{\boldsymbol{a}})G_{\mathbf{q}}(\bar{x}^{\boldsymbol{a}}).$$

By (6.5), we have

(6.8)
$$\sum_{\mathbf{p}+\mathbf{q}=\mathbf{m}} T_{\mathbf{p}}(\bar{x}) \mathcal{J}_{\mathbf{q}}(\bar{x}) = L_{\boldsymbol{a}}(\bar{x}) \mathbb{N}(\bar{x}^{\boldsymbol{a}}),$$

Subtracting (6.7) from (6.8), we obtain that there exists $\Omega \subset \mathbb{N}$ such that

$$T_{\mathbf{0}}(\bar{x})\mathcal{J}_{\mathbf{m}}(\bar{x}) + T_{\mathbf{m}}(\bar{x})\mathcal{J}_{\mathbf{0}}(\bar{x}) = L_{\boldsymbol{a}}(\bar{x})\Omega(\bar{x}^{\boldsymbol{a}}).$$

Hence, by Lemma 5.1, we obtain (6.6).

Finally, substituting (6.6) into (6.4), we obtain that (6.2) holds for some $\widetilde{T}, \widetilde{\mathcal{J}} \in \mathbb{N}^{n-m+1}$. It is an easy matter to show that $(\widetilde{T}, \widetilde{\mathcal{J}})$ is an \mathbb{N}^{n-m+1} -pair. The theorem is proved. \Box

7. 1-face reduction of \mathbb{N}^* -pair

Let (T, \mathcal{J}) be a primitive \mathbb{N}^n -pair. We say (T, \mathcal{J}) is of class \mathcal{F}_0 , if

$$T \cap \mathbb{N}\mathbf{e}_i = \{\mathbf{0}\} \text{ or } \mathbb{N}\mathbf{e}_i, \text{ for all } j \in \{1, \dots, n\}.$$

In this section we show that any primitive \mathbb{N}^n -pair is a finite extension of a pair of class \mathcal{F}_0 .

Denote $\bar{x} = (x_2, \ldots, x_n)$. Set $Q = \mathbb{N}\mathbf{e}_1$ and $\boldsymbol{a} = p$ in Theorem 6.1, we obtain

Lemma 7.1. Let (T, \mathcal{J}) be an \mathbb{N}^n -pair. If $p\mathbf{e}_1 \in T$ and $\{0, 1, \dots, p-1\}\mathbf{e}_1 \subset \mathcal{J}$, then there exists an \mathbb{N}^n -pair $(\widetilde{T}, \widetilde{\mathcal{J}})$ such that

$$T(x) = \widetilde{T}(x_1^p, \bar{x}), \quad \mathcal{J}(x) = (1 + x_1 + \dots + x_1^{p-1})\widetilde{\mathcal{J}}(x_1^p, \bar{x}).$$

Using the above lemma repeatedly, we get 'bigger' factors of T(x) and $\mathcal{J}(x)$.

Proposition 7.2. Let (T, \mathcal{J}) be an \mathbb{N}^n -pair. Denote $T \cap \mathbb{N}\mathbf{e}_1 = A\mathbf{e}_1, \mathcal{J} \cap \mathbb{N}\mathbf{e}_1 = B\mathbf{e}_1$. Let (C, D) be a $\{0, 1, \ldots, N-1\}$ -pair such that

$$A = C \oplus NA_1, \quad B = D \oplus NB_1$$

for an N-pair (A_1, B_1) . Then there exists an Nⁿ-pair $(\widetilde{T}, \widetilde{\mathcal{J}})$ such that

(7.1)
$$T(x) = C(x_1)\widetilde{T}(x_1^N, \bar{x}), \quad \mathcal{J}(x) = D(x_1)\widetilde{\mathcal{J}}(x_1^N, \bar{x}).$$

Proof. We prove the result by induction on N. If N = 1, the proposition holds trivially. Assume $N \ge 2$ and that the proposition holds for any pair and any positive integer less than N.

Assume $1 \in D$ without loss of generality.

If $C = \{0\}$, let h be the smallest integer such that $h \notin B$. Then by Lemma 7.1,

$$T(x) = T^*(x_1^h, \bar{x}), \quad \mathcal{J}(x) = (1 + x_1 + \dots + x_1^{h-1})\mathcal{J}^*(x_1^h, \bar{x})$$

Since $C(x_1) = 1$, $D(x_1) = 1 + x_1 + \dots + x_1^{N-1}$ and N|h, set

$$\widetilde{T}(x) = T^*(x_1^{h/N}, \bar{x}), \quad \widetilde{\mathcal{J}}(x) = \sum_{j=0}^{h/N-1} x_1^j \mathcal{J}^*(x_1^{h/N}, \bar{x}),$$

we obtain (7.1).

Now suppose $\#C \ge 2$. Let p be the smallest element of C other than 0. Then $p \ge 2$. On one hand, by Lemma 2.4, there exists a $\{0, 1, \ldots, N/p - 1\}$ -pair $(\widetilde{C}, \widetilde{D})$ such that

(7.2)
$$C(x_1) = \widetilde{C}(x_1^p), \quad D(x_1) = (1 + x_1 + \dots + x_1^{p-1})\widetilde{D}(x_1^p).$$

On the other hand, by Lemma 7.1, there exists an \mathbb{N}^n -pair (T^*, \mathcal{J}^*) such that

(7.3)
$$T(x) = T^*(x_1^p, \bar{x}), \quad \mathcal{J}(x) = (1 + x_1 + \dots + x_1^{p-1})\mathcal{J}^*(x_1^p, \bar{x}).$$

Denote $T^* \cap \mathbb{N}\mathbf{e}_1 = A^*\mathbf{e}_1$, $\mathcal{J}^* \cap \mathbb{N}\mathbf{e}_1 = B^*\mathbf{e}_1$. By (7.3), we have $A = pA^*$ and $B = \{0, 1, \dots, p-1\} + pB^*$, which together with (7.2) imply that

$$A^* = \widetilde{C} \oplus (N/p)A_1, \quad B^* = \widetilde{D} \oplus (N/p)B_1.$$

So by induction hypothesis, there exists an \mathbb{N}^n -pair $(\widetilde{T}, \widetilde{\mathcal{J}})$ such that

$$T^*(x) = \widetilde{C}(x_1)\widetilde{T}(x_1^{N/p}, \bar{x}), \quad \mathcal{J}^*(x) = \widetilde{D}(x_1)\widetilde{\mathcal{J}}(x_1^{N/p}, \bar{x}).$$

This together with (7.2) and (7.3) imply (7.1). The lemma is proved.

The following theorem is proved by Niven ([10, Lemma 4]) in the two dimensional case.

Theorem 7.1. If (T, \mathcal{J}) be a primitive \mathbb{N}^n -pair with $n \ge 2$, then either $T \cap \mathbb{N}\mathbf{e}_1$ or $\mathcal{J} \cap \mathbb{N}\mathbf{e}_1$ is finite.

Proof. Let $Q = \mathbb{N}\mathbf{e}_2 \oplus \cdots \oplus \mathbb{N}\mathbf{e}_n$. Recall that $T_Q = T \cap Q$, $\mathcal{J}_Q = \mathcal{J} \cap Q$. Write $T \cap \mathbb{N}\mathbf{e}_1 = A\mathbf{e}_1$, $\mathcal{J} \cap \mathbb{N}\mathbf{e}_1 = B\mathbf{e}_1$. Then $A \oplus B = \mathbb{N}$ and $T_Q \oplus \mathcal{J}_Q = Q$ (Lemma 6.1). Define $\rho(a_1, \ldots, a_n) = (a_2, \ldots, a_n)$.

Let (C, D) be a $\{0, 1, \dots, N-1\}$ -pair satisfying the assumptions of Proposition 7.2. Then there exists $H \subset \mathbb{N}^n$ such that

(7.4)
$$T(x) = C(x_1)H(x).$$

Setting $x_1 = 0$, we obtain $T(0, x_2, ..., x_n) = H(0, x_2, ..., x_n)$, which implies that $H \cap Q = T \cap Q = T_Q$. So $T_Q \subset H$, and by (7.4), we have

(7.5)
$$C \times \rho(T_Q) \subset T.$$

Suppose on the contrary that both A and B are infinite sets. By Lemma 2.5, there exists a sequence of interval pairs $\{C_k, D_k\}_{k=1}^{\infty}$ such that $C_k \uparrow A$ and $D_k \uparrow B$ and (C_k, D_k) satisfies the assumptions of Proposition 7.2. This together with (7.5) imply that $A \times \rho(T_Q) \subset T$. Similarly, we have $B \times \rho(\mathcal{J}_Q) \subset \mathcal{J}$.

Since $(A \times \rho(T_Q), B \times \rho(\mathcal{J}_Q))$ is an \mathbb{N}^n -pair (Lemma 2.1), the above two inclusion relations imply that $T = A \times \rho(T_Q)$ and $\mathcal{J} = B \times \rho(\mathcal{J}_Q)$, which imply (T, \mathcal{J}) is not primitive. This contradiction proves the theorem.

Theorem 7.2. Let (T, \mathcal{J}) be a primitive \mathbb{N}^n -pair. Then there exists a primitive \mathbb{N}^n -pair $(\widetilde{T}, \widetilde{\mathcal{J}})$ in \mathcal{F}_0 such that (T, \mathcal{J}) is a finite (first type) extension of $(\widetilde{T}, \widetilde{\mathcal{J}})$.

Proof. We first reduce (T, \mathcal{J}) along the variable x_1 . Denote $T \cap \mathbb{N}\mathbf{e}_1 = A\mathbf{e}_1, \mathcal{J} \cap \mathbb{N}\mathbf{e}_1 = B\mathbf{e}_1$. Then either A or B is finite (Theorem 7.1). Assume A is finite. By Lemma 2.5, there exist $D \subset B$ such that $A \oplus D = \{0, 1, \dots, N-1\}$ and $B = D + N\mathbb{N}$. Hence, by Proposition 7.2, there exists an \mathbb{N}^n -pair (T_1, \mathcal{J}_1) such that

(7.6)
$$T(x) = A(x_1)T_1(x_1^N, \bar{x}), \quad \mathcal{J}(x) = D(x_1)\mathcal{J}_1(x_1^N, \bar{x}).$$

We claim that: (i) $T_1 \cap \mathbb{N}\mathbf{e}_1 = \{\mathbf{0}\};$ (ii) $T_1 \cap \mathbb{N}\mathbf{e}_j = T \cap \mathbb{N}\mathbf{e}_j$ for $j \neq 1$.

From $T \cap \mathbb{N}\mathbf{e}_1 = A\mathbf{e}_1$ we infer that $T(x_1, \mathbf{0}) = A(x_1)$, which together with (7.6) imply that $T_1(x_1^N, \mathbf{0}) = 1$. So (i) holds. Pick j > 1, by (7.6), we have $T(\mathbf{0}, x_j, \mathbf{0}) = T_1(\mathbf{0}, x_j, \mathbf{0})$, which means that $T \cap \mathbb{N}\mathbf{e}_j = T_1 \cap \mathbb{N}\mathbf{e}_j$. This proves (ii).

Next, we reduce (T_1, \mathcal{J}_1) along the variables x_2, \ldots, x_n one by one, and finally we obtain an \mathbb{N}^n -pair $(\tilde{T}, \tilde{\mathcal{J}})$ of class \mathcal{F}_0 . The theorem is proved.

8. Proof of Theorem 1.1

The following lemma provides the last ingredient for proving Theorem 1.1.

Lemma 8.1. Let $n \ge 2$ and (T, \mathcal{J}) be a primitive \mathbb{N}^n -pair of class \mathcal{F}_0 . Then there exists an m-face Q_0 with $2 \le m \le n$ such that the marginal pair induced by Q_0 is of pure type. *Proof.* Without loss of generality, we assume that $T \cap \mathbb{N}\mathbf{e}_j = \{\mathbf{0}\}$ holds for at least one j. By rearranging the order of variables, we may assume that

$$T \cap \mathbb{N}\mathbf{e}_j = \{\mathbf{0}\}$$
 for $j = 1, \dots, M$, and $T \cap \mathbb{N}\mathbf{e}_j = \mathbb{N}\mathbf{e}_j$ otherwise,

where $1 \leq M \leq n$. Denote $Q = \bigoplus_{j=1}^{M} \mathbb{N} \mathbf{e}_j$ and $Q' = \bigoplus_{j=M+1}^{n} \mathbb{N} \mathbf{e}_j$. (If M = n, we set $Q' = \{\mathbf{0}\}$.) Then (T_Q, \mathcal{J}_Q) is a Q-pair and $(T_{Q'}, \mathcal{J}_{Q'})$ is a Q'-pair.

We claim that either $\#T_Q \ge 2$ or $\#\mathcal{J}_{Q'} \ge 2$. Suppose on the contrary that $\#T_Q = 1$ and $\#\mathcal{J}_{Q'} = 1$. Then $T_Q = \mathcal{J}_{Q'} = \{\mathbf{0}\}$, which implies that $\mathcal{J}_Q = Q \subset \mathcal{J}$ and $T_{Q'} = Q' \subset T$. It follows that $(T, \mathcal{J}) = (Q', Q)$ and it is not primitive, which contradicts our assumption.

Without loss of generality, let us assume that $\#T_Q \ge 2$. Let $x^{a'}$ be a term of $T_Q(x) - 1$ such that the number of non-zero entries of a' attains the minimum. Write $x^{a'}$ as

$$x^{a'} = x_{j_1}^{r_1} \cdots x_{j_m}^{r_m}, \quad \text{where } j_1, \dots, j_m \in \{1, \dots, M\}.$$

Then $m \ge 2$ since $T \cap \mathbb{N}\mathbf{e}_j = \{\mathbf{0}\}$ for $1 \le j \le M$. Let $Q_0 = \mathbb{N}\mathbf{e}_{j_1} \oplus \cdots \oplus \mathbb{N}\mathbf{e}_{j_m}$, which is the smallest face containing \mathbf{a}' . Let (f,g) be the marginal pair induced by Q_0 , then (f,g)is an \mathbb{N}^m -pair of pure type by the minimality of m. The lemma is proved. \Box

Proof of Theorem 1.1. By Theorem 3.1, we only need show that any primitive \mathbb{N}^* -pair is a finite extension of an \mathbb{N} -pair. Let (T, \mathcal{J}) be a primitive \mathbb{N}^n -pair. We prove by induction on the dimension of (T, \mathcal{J}) . If n = 1, the theorem is trivially true.

Assume $n \geq 2$. By Theorem 7.2, there exists a primitive \mathbb{N}^n -pair (T_1, \mathcal{J}_1) of class \mathcal{F}_0 such that (T, \mathcal{J}) is a finite extension of it. By Lemma 8.1, (T_1, \mathcal{J}_1) has an *m*-face Q_0 with $m \geq 2$ such that the marginal pair induced by Q_0 is of pure type. So by Theorem 6.1, there exists a primitive \mathbb{N}^{n-m+1} -pair $(\widetilde{T}, \widetilde{\mathcal{J}})$ such that (T_1, \mathcal{J}_1) is a second type extension of $(\widetilde{T}, \widetilde{\mathcal{J}})$. Finally, by induction hypothesis, $(\widetilde{T}, \widetilde{\mathcal{J}})$ is a finite extension of an \mathbb{N} -pair, say (T_0, \mathcal{J}_0) . Then $(T_0, \mathcal{J}_0) \to (\widetilde{T}, \widetilde{\mathcal{J}}) \to (T_1, \mathcal{J}_1) \to (T, \mathcal{J})$ indicates the extension process from (T_0, \mathcal{J}_0) to (T, \mathcal{J}) .

9. Weighted tree of \mathbb{N}^* -pair

In this section, we study weighted trees and the associated \mathbb{N}^* -pairs.

9.1. Weighted tree. A graph (V, Γ) with node set V and edge set Γ is a *tree* if it is connected and has no cycle. A *rooted tree* is a triple (V, Γ, ϕ) where (V, Γ) is a tree and $\phi \in V$; we call ϕ the *root* of the tree.

Let (V, Γ) be a finite tree with root ϕ . The *level* of v, denoted by |v|, is the length of the (unique) path joining v and ϕ . A node v is called an *offspring* of u, if there is an edge

joining u and v, and |v| = |u| + 1; meanwhile we call u the parent of v. For $u, v \in V$, we use [u, v] to denote the edge joining u and its offspring v. We call u an ancestor of v if there is a path joining u and v, and |u| < |v|. A node v is called a top if it has no offspring. We denote by V_0 the set of tops. We shall use the nodes as variables of power series.

Definition 9.1. We call the quadruple

$$(9.1) \qquad \qquad ((T_0, \mathcal{J}_0), \boldsymbol{\delta}, \boldsymbol{\alpha}, \Phi)$$

a weight of the tree (V, Γ) , where (T_0, \mathcal{J}_0) is an N-pair, called the *initial pair*,

$$\boldsymbol{\delta}: V \setminus V_0 \to \{0,1\}, \ \boldsymbol{\alpha}: V \setminus \{\phi\} \to \mathbb{N} \setminus \{0\}, \ \Phi: V \setminus \{\phi\} \to \{\text{interval pairs}\}$$

are three maps, and in addition, $\delta_{\phi} = 1$ if $T_0 = \{0\}$ and $\delta_{\phi} = 0$ if $\mathcal{J}_0 = \{0\}$.

We call $\#V - \#V_0$ the norm of the tree. Two tops are said to be equivalent, if they share the same parents. An equivalent class of V_0 is called a *branch* of V_0 .

9.2. Constructing \mathbb{N}^* -pairs from weighted trees. Now we define the pair generated by a weighted tree inductively on the norm of the tree.

Notice that $\Lambda_0 = (\{\phi\}, \emptyset)$ is the only tree with norm 0. The weight of Λ_0 only consists of the initial pair, which we denote by (T_0, \mathcal{J}_0) . We define the \mathbb{N}^* -pair associate with the weighted tree Λ_0 to be (T_0, \mathcal{J}_0) .

Let $q \ge 1$. Suppose for any weighted tree with norm less than q, we have associated an \mathbb{N}^m -pair with it, where m is the number of tops of the tree.

Let (V, Γ) be a weighted tree with norm q. Let $V_0 = \{x_1, \ldots, x_n\}$. Choose a branch $\{x_1, \ldots, x_p\}$ of V_0 and denote their parents by z_1 . Let $(\tilde{V}, \tilde{\Gamma})$ be the subtree of (V, Γ) obtained by deleting the nodes x_1, \ldots, x_p and the edges $[z_1, x_1], \ldots, [z_1, x_p]$. Then

(9.2)
$$\widetilde{V} = V \setminus \{x_1, \cdots, x_p\}, \quad \widetilde{V}_0 = \{z_1, x_{p+1}, \ldots, x_n\}.$$

We define the weight of $(\tilde{V}, \tilde{\Gamma})$ to be the restriction of the weight $((T_0, \mathcal{J}_0), \delta, \alpha, \Phi)$ on it. (Now z_1 is a top of the subtree, so δ_{z_1} is not needed in the restricted weight.)

Since $\#\widetilde{V} - \#\widetilde{V}_0 = q - 1$, by the induction hypothesis, there is an \mathbb{N}^{n-p+1} -pair $(\widetilde{T}, \widetilde{\mathcal{J}})$ associated with the weighted tree of $(\widetilde{V}, \widetilde{\Gamma})$.

Denote $\delta = \delta_{z_1}$. For j = 1, ..., p, let $a_j = \alpha_{x_j}$, $(C_j, D_j) = \Phi_{x_j}$, and let N_j be the size of (C_j, D_j) . Denote $\bar{x} = (x_{p+1}, ..., x_n)$ and $a = (a_1, ..., a_p)$.

First, applying an second type extension with parameters δ and \boldsymbol{a} to (T, \mathcal{J}) , we obtain

$$F(x) = L_{a}^{\delta}(x_{1}, \dots, x_{p})\widetilde{T}(\prod_{j=1}^{p} x_{j}^{a_{j}}, \bar{x}), \quad G(x) = L_{a}^{1-\delta}(x_{1}, \dots, x_{p})\widetilde{\mathcal{J}}(\prod_{j=1}^{p} x_{j}^{a_{j}}, \bar{x}).$$

Secondly, applying p extensions of the first type consecutively to the pair (F, G), along the variables x_1, \ldots, x_p , we obtain

(9.3)
$$T(x) = \left(\prod_{j=1}^{p} C_j(x_j)\right) L_a^{\delta}(x_1^{N_1}, \dots, x_p^{N_p}) \widetilde{T}(\prod_{j=1}^{p} x_j^{a_j N_j}, \bar{x}),$$
$$\mathcal{J}(x) = \left(\prod_{j=1}^{p} D_j(x_j)\right) L_a^{1-\delta}(x_1^{N_1}, \dots, x_p^{N_p}) \widetilde{\mathcal{J}}(\prod_{j=1}^{p} x_j^{a_j N_j}, \bar{x}).$$

We call (T, \mathcal{J}) the \mathbb{N}^* -pair generated by the weighted tree (9.1).

To show the generated pair is well-defined, we need to show that (T, \mathcal{J}) is independent of the choice of the branch $\{x_1, \ldots, x_p\}$. We will do this in Theorem 9.1.

Remark 9.2. Clearly, $\#T < \infty$ if and only if $\#T_0 < \infty$ and δ is constantly 0; a similar result holds for $\#\mathcal{J}$. Also, (T, \mathcal{J}) is of class \mathcal{F}_0 if and only if $\Phi_v = (\{\mathbf{0}\}, \{\mathbf{0}\})$ for all tops.

9.3. A closed formula of (T, \mathcal{J}) . Let (T, \mathcal{J}) be the pair defined by (9.3). We define a map $\mu : V \times V_0 \to \mathbb{N}$ as follows. Pick $v \in V$ and $x^* \in V_0$.

If v is an ancestor of x^* , let $v, v_1, \ldots, v_{k-1}, v_k = x^*$ be the path from v to x^* ; for $j = 1, \ldots, k$, let $c_j = \alpha_{v_j}$ and M_j be the size of Φ_{v_j} . We define

(9.4)
$$\mu(v, x^*) = \begin{cases} \prod_{j=1}^k c_j M_j, & \text{if } v \text{ is an ancestor of } x^*; \\ 1, & \text{if } v = x^*; \\ 0, & \text{otherwise.} \end{cases}$$

Moreover, we define $\mu(v) = (\mu(v, x_1), \dots, \mu(v, x_n)).$

For each $v \in V$, we define two power series P_v and Q_v as follows. Denote $\Phi_v = (C_v, D_v)$. If v is a top, we set

(9.5)
$$P_v(x) = C_v(v), \quad Q_v(x) = D_v(v).$$

(If $V = \{\phi\}$, we make the convention $P_{\phi}(\phi) = T_0(\phi)$ and $Q_{\phi}(\phi) = \mathcal{J}_0(\phi)$.)

If v is not a top, let z_1, \ldots, z_m be the offsprings of v. Denote $\mathbf{b} = (\boldsymbol{\alpha}_{z_1}, \ldots, \boldsymbol{\alpha}_{z_m})$ and M_j be the size of Φ_{z_j} . (If $v = \phi$, we set $C_{\phi} = T_0$ and $D_{\phi} = \mathcal{J}_0$.) We define

(9.6)
$$P_{v}(x) = C_{v}(x^{\mu(v)}) L_{\mathbf{b}}^{\boldsymbol{\delta}_{v}}(x^{M_{1}\mu(z_{1})}, \dots, x^{M_{m}\mu(z_{m})}),$$
$$Q_{v}(x) = D_{v}(x^{\mu(v)}) L_{\mathbf{b}}^{1-\boldsymbol{\delta}_{v}}(x^{M_{1}\mu(z_{1})}, \dots, x^{M_{m}\mu(z_{m})}).$$

Theorem 9.1. Let (T, \mathcal{J}) be an \mathbb{N}^n -pair generated by the tree (V, Γ) with weight (9.1). Then $P_v(x)$ and $Q_v(x)$ defined by (9.5) and (9.6) are binary power series, and

(9.7)
$$T(x) = \prod_{v \in V} P_v(x), \quad \mathcal{J}(x) = \prod_{v \in V} Q_v(x)$$

Proof. Denote $q = \#V - \#V_0$. We prove the result by induction on q. If q = 0, then $V = \{\phi\}$, and (9.7) holds by our convention. Now we assume that $q \ge 1$.

Let $(\widetilde{V}, \widetilde{\Gamma})$ be the subtree of (V, Γ) given by (9.2), and the weight is the restricted weight. According to this restricted weight, for any $v \in \widetilde{V}$ and $z \in \widetilde{V}_0 = \{z_1, x_{p+1}, \ldots, x_n\}$, we can define a map $\widetilde{\mu}(v, z)$ as (9.4), and power series $\widetilde{P}_v, \widetilde{Q}_v$ as (9.5) and (9.6).

Let $(\widetilde{T}, \widetilde{\mathcal{J}})$ be the pair generated by the weighted tree $(\widetilde{V}, \widetilde{\Gamma})$. Denote $\overline{x} = (x_{p+1}, \ldots, x_n)$. By induction hypothesis, we have

(9.8)
$$\widetilde{T}(z_1, \bar{x}) = \prod_{v \in \widetilde{V}} \widetilde{P}_v(z_1, \bar{x}).$$

For j = 1, ..., p, denote $a_j = \alpha_{x_j}$, and let N_j be the size of $\Phi_{x_j} = (C_{x_j}, D_{x_j})$. We assert that

(9.9)
$$x^{\mu(v)} = (z_1, \bar{x})^{\widetilde{\mu}(v)} \circ \left(z_1 \mapsto \prod_{j=1}^p x_j^{a_j N_j}\right), \text{ for } v \in \widetilde{V}.$$

Notice that (9.9) holds if the following holds:

(9.10)
$$\mu(v, x_j) = \begin{cases} \widetilde{\mu}(v, z_1) a_j N_j, & \text{if } 1 \le j \le p; \\ \widetilde{\mu}(v, x_j) & \text{if } p+1 \le j \le n. \end{cases}$$

If v is an ancestor of z_1 or $v = z_1$, we have $\mu(v, x_j) = \tilde{\mu}(v, z_1)\mu(z_1, x_j) = \tilde{\mu}(v, z_1)a_jN_j$, $1 \le j \le p$; if v is not an ancestor of z_1 , we have $\mu(v, x_1) = \cdots = \mu(v, x_p) = \tilde{\mu}(v, z_1) = 0$. So the first assertion of (9.10) holds. The second assertion holds since the path from v to x_j in Γ is also a path in $\tilde{\Gamma}$. This proves (9.10) and (9.9) follows.

Now, we consider the relations between P_v and \tilde{P}_v for $v \in \tilde{V}$.

If $v = z_1$, since z_1 is a top of \widetilde{V} , we have $\widetilde{P}_{z_1}(z_1, \overline{x}) = C_{z_1}(z_1)$; moreover, $\mu(z_1) = (a_1 N_1, \ldots, a_p N_p, 0, \cdots, 0)$ and $\mu(x_j) = \mathbf{e}_j$, $1 \le j \le n$. Denote $\mathbf{a} = (a_1, \ldots, a_p)$, we have

(9.11)
$$P_{z_1}(x) = C_{z_1} \left(\prod_{j=1}^p x_j^{a_j N_j} \right) L_{a}^{\delta_{z_1}}(x_1^{N_1}, \dots, x_p^{N_p}) \\ = \left(\widetilde{P}_{z_1}(z_1, \bar{x}) \circ (z_1 \mapsto \prod_{j=1}^p x_j^{a_j N_j}) \right) L_{a}^{\delta_{z_1}}(x_1^{N_1}, \cdots, x_p^{N_p}).$$

We claim that

(9.12)
$$P_v(x) = \widetilde{P}_v(z_1, \bar{x}) \circ \left(z_1 \mapsto \prod_{j=1}^p x_j^{a_j N_j}\right), \quad \text{if } v \in \widetilde{V} \setminus \{z_1\}.$$

If $v \in \{x_{p+1}, \ldots, x_n\}$, we have $P_v(x) = \widetilde{P}_v(z_1, \overline{x}) = C_v(v)$, clearly (9.12) holds.

If $v \in \widetilde{V} \setminus \{z_1, x_{p+1}, \ldots, x_n\}$, denote the offsprings of v by y_1, \cdots, y_m . Set $\mathbf{b} = (\boldsymbol{\alpha}_{y_1}, \cdots, \boldsymbol{\alpha}_{y_m})$, and let M_i be the size of Φ_{y_i} for $1 \leq i \leq m$. Since $C_v = \widetilde{C}_v$ and

 $\delta_v = \widetilde{\delta}_v$, we have

$$P_{v}(x) = C_{v}(x^{\mu(v)}) L_{\mathbf{b}}^{\delta_{v}}(x^{M_{1}\mu(y_{1})}, \dots, x^{M_{m}\mu(y_{m})}),$$

$$\widetilde{P}_{v}(z_{1}, \bar{x}) = C_{v}((z_{1}, \bar{x})^{\widetilde{\mu}(v)}) L_{\mathbf{b}}^{\delta_{v}}((z_{1}, \bar{x})^{M_{1}\widetilde{\mu}(y_{1})}, \dots, (z_{1}, \bar{x})^{M_{m}\widetilde{\mu}(y_{m})}).$$

Notice that (9.9) holds for all $v, y_1, \ldots, y_m \in \widetilde{V} \setminus \{z_1\}$, so (9.12) follows. The claim is proved. Therefore, we have

$$T(x) = \left(\prod_{j=1}^{p} C_{x_j}(x_j)\right) L_{a}^{\delta_{z_1}}(x_1^{N_1}, \dots, x_p^{N_p}) \cdot \widetilde{T}\left(\prod_{j=1}^{p} x_j^{a_j N_j}, \bar{x}\right)$$
(By (9.3))

$$= \left(\prod_{j=1}^{p} P_{x_j}(x)\right) L_{a}^{\delta_{z_1}}(x_1^{N_1}, \dots, x_p^{N_p}) \cdot \prod_{v \in \widetilde{V}} \widetilde{P}_v(z_1, \bar{x}) \circ (z_1 \mapsto \prod_{j=1}^{p} x_j^{a_j N_j})$$
(By (9.8))

$$= \left(\prod_{j=1}^{p} P_{x_{j}}(x)\right) \left(L_{a}^{\delta_{z_{1}}}(x_{1}^{N_{1}}, \dots, x_{p}^{N_{p}}) \widetilde{P}_{z_{1}}(\prod_{j=1}^{p} x_{j}^{a_{j}N_{j}}, \bar{x})\right) \left(\prod_{v \in \widetilde{V} \setminus \{z_{1}\}} P_{v}(x)\right) \quad (By \ (9.12))$$
$$= \prod P_{v}(x). \qquad (By \ (9.11).)$$

$$=\prod_{v\in V} P_v(x).$$
(By (9.11).

By symmetry, we have $\mathcal{J}(x) = \prod_{v \in V} Q_v(x)$. The theorem is proved.

10. Proof of Theorem 1.2 and an example

Proof of Theorem 1.2. Thanks to Theorem 3.1, we only need show that any primitive \mathbb{N}^* -pair (T, \mathcal{J}) can be generated by a weighted tree. If (T, \mathcal{J}) is an \mathbb{N} -pair, then it is generated by the tree $\Lambda_0 = (\{\phi\}, \emptyset)$ with initial pair $(T_0, \mathcal{J}_0) = (T, \mathcal{J})$.

Suppose (T, \mathcal{J}) is generated by a tree (V, Γ) with weight $((T_0, \mathcal{J}_0), \delta, \alpha, \Phi)$. In the following, we show that any one step extension (T^*, \mathcal{J}^*) of it can also be generated by a weighted tree. Denote the tops of (V, Γ) by x_1, \ldots, x_n , and assume the extension is along the variable x_1 . Denote $\bar{x} = (x_2, \ldots, x_n)$.

Case 1. Suppose (T^*, \mathcal{J}^*) is a first type extension of (T, \mathcal{J}) given by

$$T^*(x) = C(x_1)T(x_1^N, \bar{x}), \quad \mathcal{J}^*(x) = D(x_1)\mathcal{J}(x_1^N, \bar{x}),$$

where (C, D) is a $\{0, \ldots, N-1\}$ -pair. Denote $\Phi_{x_1} = (C_{x_1}, D_{x_1})$. We define a new weight of (V, Γ) by only modifying Φ_{x_1} to the interval pair $(C + NC_{x_1}, D + ND_{x_1})$. Then (T^*, \mathcal{J}^*) is generated by the tree (V, Γ) with this new weight.

Case 2. Suppose (T^*, \mathcal{J}^*) is a second type extension of (T, \mathcal{J}) given by

$$T^*(y,\bar{x}) = L^s_{\mathbf{b}}(y)T(y^{\mathbf{b}},\bar{x}), \quad \mathcal{J}^*(y,\bar{x}) = L^{1-s}_{\mathbf{b}}(y)\mathcal{J}(y^{\mathbf{b}},\bar{x})$$

where $y = (y_1, ..., y_m)$, $\mathbf{b} = (b_1, ..., b_m) > 0$ and $s \in \{0, 1\}$. We set $V' = V \cup \{y_1, ..., y_m\}$, and let $\Gamma' = \Gamma \cup \{[x_1, y_j]; j = 1, ..., m\}$ be the edge set. We define a



FIGURE 1. A rooted tree

weight $((T_0, \mathcal{J}_0), \delta', \alpha', \Phi')$ of (V', Γ') , which is the extension of the original weight by adding the following parameters: (i) Set $\delta'_{x_1} = s$; (ii) For $j \in \{1, \ldots, m\}$, set $\alpha'_{y_j} = b_j$ and $\Phi'_{y_j} = (\{\mathbf{0}\}, \{\mathbf{0}\})$. Then (T^*, \mathcal{J}^*) is generated by (V', Γ') with the above weight. \Box

The function a and F.					
node	y_1	x_4	x_1	x_2	x_3
α	2	3	1	1	1
C	$\{0, 2\}$	$\{0,2\}$	{0}	{0}	{0}
D	$\{0,1\}$	$\{0,1\}$	{0}	{0}	{0}
N	4	4	1	1	1

The function $\boldsymbol{\alpha}$ and $\boldsymbol{\Phi}$:

TABLE 1. We set $\boldsymbol{\delta}$ defined on $\{\phi, y_1\}$ to be constantly zero.

Example 10.1. Let (Γ, V) be a rooted tree indicated by Figure 1. The node set $V = \{\phi, y_1, x_1, x_2, x_3, x_4\}$. We set the initial N-pair to be

$$T_0(\phi) = 1 + \phi^2$$
, $\mathcal{J}_0(\phi) = (1 + \phi) \sum_{k=0}^{\infty} \phi^{4k}$;

the other parameters of the weight is given by Table 1. Set

$$V_0 = \{\phi\}, V_1 = \{\phi; y_1, x_4\}, V_2 = \{\phi; y_1, x_4; x_1, x_2, x_3\}.$$

Let $(V_j, \Gamma_j), j = 0, 1, 2$, be subtrees of (V, Γ) . The \mathbb{N}^* -tile T_1 corresponding to the subtree (V_1, Γ_1) is $T_1(y_1, x_4) = (1 + y_1^2)(1 + x_4^2)(1 + y_1^{16}x_4^{24})$. The \mathbb{N}^* -tile T_2 corresponding to the tree $(V_2, \Gamma_2) = (V, \Gamma)$ is

$$T_2(x_1, x_2, x_3, x_4) = 1 \cdot 1 \cdot 1 \cdot (1 + (x_1 x_2 x_3)^2) \cdot (1 + x_4^2) \cdot (1 + (x_1 x_2 x_3)^{16} x_4^{24}) .$$

= $P_{x_1}(x) \cdot P_{x_2}(x) \cdot P_{x_3}(x) \cdot P_{y_1}(x) \cdot P_{x_4}(x) \cdot P_{\phi}(x).$

HUI RAO, YA-MIN YANG, AND YUAN ZHANG*

References

- E. M. Coven and A. Meyercovitz, *Tiling the integers with translates of one finite set*, J. Algebra, 212 (1999), 161–174.
- [2] N. G. de Bruijn, On bases for the set of integers, Publ. Math., 1(3)(1956), 583–590.
- [3] N. G. de Bruijn, On number systems, Nieuw Arch. Wisk., 4(3)(1956), 15-17.
- [4] S. Eigen and A. Hajian, Sequences of integers and ergodic transformations, Adv. Math., 73 (1989), 256–262.
- [5] R. T. Hansen, Complementing pairs of subsets in the plane, Duke Math. J. 36(1969), 441–449.
- [6] G. Hajós, Sur la factorisation des groupes abélians, Casopis Pest. math. Fys. (3) 4 (1950), 157–162.
- [7] C. T. Long, Addition theorems for sets of integers, *Pacific J. Math.* 23(1967), 107–112.
- [8] M. B. Nathanson, Complementing sets of n-tuples of integers. Proc. Amer. Math. Soc. 34 (1972), 71-72.
- [9] D. J. Newman, Tesselation of integers. J. Number Theory 9 (1977), 107-111.
- [10] I. Niven, A characterization of complementing sets of pairs of integers. Duke Math. J. 38 (1971), 193–203.
- [11] A. D. Sands, On Keller's conjecture for certain cyclic groups, Proc. Edinburg Math. Soc. (1), 22 (1979), 17–21.
- [12] Szegedy, M. Algorithms to tile the infinite grid with finite clusters. Proceedings of the 39th Annual Symposium on the Foundations of Computer Science, FOCS 98. November8C111998, Palo Alto, CA. pp.137C145. IEEE Computer Society.
- [13] C. Swenson, Direct sum subset decompositions of Z. Pacific J. Math. 53 (1974), 629–633.
- [14] S. Szabó, A type of factorization of finite Abelian groups, Discrete Math. 54(1985), 121–124.
- [15] R. Tijdeman, Decomposition of the integer as a direc sum of two subsets, in "Number Theory(Paris, 1992-1993)," London Math. Soc. Lecture Note Ser., Vol.215, 261–276. Cambridge Univ. Press, 1995.
- [16] A. M. Vaidya, On complementing sets of nonnegative integers, Math. Mag. 39(1966), 43-44.

School of Mathematics and Statistic, Huazhong University of Science and Technology, Wuhan, 430074, China

E-mail address: hrao@mail.ccnu.edu.cn

INSTITUTE OF APPLIED MATHEMATICS, COLLEGE OF SCIENCE, HUAZHONG AGRICULTURE UNIVERSITY, WUHAN,430070, CHINA.

E-mail address: yangym09@mail.hzau.edu.cn

School of Mathematics and Statistic, Huazhong University of Science and Technology, Wuhan, 430074, China

E-mail address: yzhangccnu@sina.cn