

On factorized groups with permutable subgroups of factors

Victor S. Monakhov and Alexander A. Trofimuk

Department of Mathematics and Programming Technologies,

Francisk Skorina Gomel State University,

Gomel 246019, Belarus

e-mail: victor.monakhov@gmail.com

e-mail: alexander.trofimuk@gmail.com[†]

To the 75th anniversary of Mohamed Asaad

Abstract. The subgroups A and B of a group G are called msp-permutable, if the following statements hold: AB is a subgroup of G ; the subgroups P and Q are mutually permutable, where P is an arbitrary Sylow p -subgroup of A and Q is an arbitrary Sylow q -subgroup of B , $p \neq q$. In the present paper, we investigate groups that factorized by two msp-permutable subgroups. In particular, the supersolubility of the product of two supersoluble msp-permutable subgroups is proved.

Keywords. mutually permutable subgroups, Sylow subgroups, msp-permutable subgroups, supersoluble groups.

Mathematics Subject Classification. 20D10, 20D20.

Throughout this paper, all groups are finite and G always denotes a finite group. We use the standard notations and terminology of [3]. The notation $Y \leq X$ means that Y is a subgroup of a group X .

The subgroups A and B of a group G are called *mutually (totally) permutable*, if $UB = BU$ and $AV = VA$ (respectively, $UV = VU$) for all $U \leq A$ and $V \leq B$.

The idea of totally and mutually permutable subgroups was first initiated by M. Asaad and A. Shaalan in [1]. This direction have since been subject of an in-depth study of many authors. An exhaustive report on this matter appears in [3, chapters 4–5].

It is quite natural to consider a factorized group $G = AB$ in which certain subgroups of the factors A and B are mutually (totally) permutable. In this direction, V.S. Monakhov [7] obtained the solubility of a group $G = AB$ under the assumption that the subgroups A and B are soluble and the Carter subgroups (Sylow subgroups) of A and of B are permutable.

We introduce the following

Definition. The subgroups A and B of a group G are called msp-permutable, if the following statements hold:

- (1) AB is a subgroup of G ;
- (2) the subgroups P and Q are mutually permutable, where P is an arbitrary Sylow p -subgroup of A and Q is an arbitrary Sylow q -subgroup of B , $p \neq q$.

In the present paper, we investigate groups that factorized by two msp-permutable subgroups. In particular, the supersolubility of the product of two supersoluble msp-permutable subgroups is proved.

1 Preliminaries

In this section, we give some definitions and basic results which are essential in the sequel. A group whose chief factors have prime orders is called *supersoluble*. Recall that a *p -closed* group is a group with a normal Sylow p -subgroup and a *p -nilpotent* group is a group with a normal Hall p' -subgroup.

Denote by G' , $Z(G)$, $F(G)$ and $\Phi(G)$ the derived subgroup, centre, Fitting and Frattini subgroups of G , respectively; \mathbb{P} the set of all primes. We use E_{p^t} to denote an elementary abelian group of order p^t and Z_m to denote a cyclic group of order m . The semidirect product of a normal subgroup A and a subgroup B is written as follows: $A \rtimes B$.

The monographs [2], [5] contain the necessary information of the theory of formations. The formations of all nilpotent, p -groups and supersoluble groups are denoted by \mathfrak{N} , \mathfrak{N}_p and \mathfrak{U} , respectively. A formation \mathfrak{F} is said to be *saturated* if $G/\Phi(G) \in \mathfrak{F}$ implies $G \in \mathfrak{F}$. A *formation function* is a function f defined on \mathbb{P} such that $f(p)$ is a, possibly empty, formation. A formation \mathfrak{F} is said to be *local* if there exists a formation function f such that $\mathfrak{F} = \{G \mid G/F_p(G) \in f(p)\}$. Here $F_p(G)$ is the greatest normal p -nilpotent subgroup of G . We write $\mathfrak{F} = LF(f)$ and f is a local definition of \mathfrak{F} . By [5, Theorem IV.3.7], among all possible local definitions of a local formation \mathfrak{F} there exists a unique f such that f is integrated (i.e., $f(p) \subseteq \mathfrak{F}$ for all $p \in \mathbb{P}$) and full (i.e., $f(p) = \mathfrak{N}_p f(p)$ for all $p \in \mathbb{P}$). Such local definition f is said to be *canonical local definition* of \mathfrak{F} . By [5, Theorem IV.4.6], a formation is saturated if and only if it is local.

A subgroup H of a group G is called \mathbb{P} -subnormal in G , see [12], if either $H = G$, or there is a chain subgroups

$$H = H_0 \leq H_1 \leq \dots \leq H_n = G, \quad |H_i : H_{i-1}| \in \mathbb{P}, \quad \forall i.$$

A group G is called *w-supersoluble* (widely supersoluble), if every Sylow subgroup of G is \mathbb{P} -subnormal in G . Denote by $\mathbf{w}\mathfrak{U}$ the class of all w-supersoluble groups, see [12]. In [12, Theorem 2.7, Proposition 2.8] proved that $\mathbf{w}\mathfrak{U}$ is a subgroup-closed saturated formation and every group from $\mathbf{w}\mathfrak{U}$ has an ordered Sylow tower of supersoluble type. By [8, Theorem B], [9, Theorem 2.6], [12, Theorem 2.13], $G \in \mathbf{w}\mathfrak{U}$ if and only if G has an ordered Sylow tower of supersoluble type and every metanilpotent (biprimary) subgroup of G is supersoluble.

Denote by $\mathbf{v}\mathfrak{U}$ the class of groups all of whose primary cyclic subgroups are \mathbb{P} -subnormal. In [8, Theorem B] proved that $\mathbf{v}\mathfrak{U}$ is a subgroup-closed saturated formation and $G \in \mathbf{v}\mathfrak{U}$ if and only if G has an ordered Sylow tower of supersoluble type and every biprimary subgroup of G with a cyclic Sylow subgroup is supersoluble. It is easy to verify that $\mathfrak{U} \subseteq \mathbf{w}\mathfrak{U} \subseteq \mathbf{v}\mathfrak{U} \subseteq \mathcal{D}$. Here \mathcal{D} is the formation of all groups which have an ordered Sylow tower of supersoluble type.

If H is a subgroup of G , then $H_G = \bigcap_{x \in G} H^x$ is called *the core* of H in G . If a group G contains a maximal subgroup M with trivial core, then G is said to be *primitive* and M is its *primitivator*. A simple check proves the following lemma.

Lemma 1.1. *Let \mathfrak{F} be a saturated formation and G be a group. Assume that $G \notin \mathfrak{F}$, but $G/N \in \mathfrak{F}$ for all non-trivial normal subgroups N of G . Then G is a primitive group.*

Lemma 1.2. ([5, Theorem 15.6]) *Let G be a soluble primitive group and M is a primitivator of G . Then the following statements hold:*

- (1) $\Phi(G) = 1$;
- (2) $F(G) = C_G(F(G)) = O_p(G)$ and $F(G)$ is an elementary abelian subgroup of order p^n for some prime p and some positive integer n ;
- (3) G contains a unique minimal normal subgroup N and moreover, $N = F(G)$;
- (4) $G = F(G) \rtimes M$ and $O_p(M) = 1$.

Lemma 1.3. ([10, Lemma 2.16]) *Let \mathfrak{F} be a saturated formation containing \mathfrak{U} and G be a group with a normal subgroup E such that $G/E \in \mathfrak{F}$. If E is cyclic, then $G \in \mathfrak{F}$.*

Lemma 1.4. *Let \mathfrak{F} be a formation, G group, A and B subgroups of G such that A and B belong to \mathfrak{F} . If $[A, B] = 1$, then $AB \in \mathfrak{F}$.*

Proof. Since

$$[A, B] = \langle [a, b] \mid a \in A, b \in B \rangle = 1,$$

it follows that $ab = ba$ for all $a \in A, b \in B$. Let

$$A \times B = \{(a, b) \mid a \in A, b \in B\},$$

$$(a_1, b_1)(a_2, b_2) = (a_1a_2, b_1b_2), \quad \forall a_1, a_2 \in A, b_1, b_2 \in B$$

be the external direct product of groups A and B . Since $A \in \mathfrak{F}$, $B \in \mathfrak{F}$ and \mathfrak{F} is a formation, we have $A \times B \in \mathfrak{F}$. Let $\varphi : A \times B \rightarrow AB$ be a function with $\varphi((a, b)) = ab$. It is clear that φ is a surjection. Because

$$\begin{aligned} \varphi((a_1, b_1)(a_2, b_2)) &= \varphi((a_1a_2, b_1b_2)) = a_1a_2b_1b_2 = \\ &= a_1b_1a_2b_2 = \varphi((a_1, b_1))\varphi((a_2, b_2)), \end{aligned}$$

it follows that φ is an epimorphism. The core $\text{Ker } \varphi$ contains all elements (a, b) such that $ab = 1$. In this case $a = b^{-1} \in A \cap B \leq Z(G)$. By the Fundamental Homomorphism Theorem,

$$A \times B / \text{Ker } \varphi \cong AB.$$

Since $A \times B \in \mathfrak{F}$ and \mathfrak{F} is a formation, $A \times B / \text{Ker } \varphi \in \mathfrak{F}$. Hence $AB \in \mathfrak{F}$. \square

Lemma 1.5. ([4]) *Let a group $G = HK$ be the product of subgroups H and K . If L is normal in H and $L \leq K$, then $L \leq K_G$.*

Lemma 1.6. *Let $G = P \rtimes M$ be a primitive soluble group, where M is a primitivator of G and P is a Sylow p -subgroup of G . Let A and B be subgroups of M and $M = AB$. If $B \leq N_G(X)$ for every subgroup X of P , then the following statements hold:*

- (1) B is a cyclic group of order dividing $p - 1$;
- (2) $[A, B] = 1$.

Proof. We fix an element $b \in B$. If $x \in P$, then $x^b \in \langle x \rangle$, since $B \leq N_G(\langle x \rangle)$ by hypothesis. Hence $x^b = x^{m_x}$, where m_x is a positive integer and $1 \leq m_x \leq p$. If $y \in P \setminus \{x\}$, then

$$(xy)^b = (xy)^{m_{xy}} = x^{m_{xy}}y^{m_{xy}}, \quad (xy)^b = x^by^b = x^{m_x}y^{m_y},$$

$$x^{m_{xy}}y^{m_{xy}} = x^{m_x}y^{m_y}, \quad x^{m_{xy}-m_x} = y^{m_y-m_{xy}} = 1, \quad m_{xy} = m_x = m_y.$$

Therefore we can assume that $x^b = x^{n_b}$ for all $x \in P$, where $1 \leq n_b \leq p$ and n_b is a positive integer.

Assume that there exist $d \in B$ and $y \in P \setminus \{1\}$ such that $y^d = y$. Then $n_d = 1$ and $x^d = x$ for all $x \in P$, i.e. $d \in C_G(P) = P$ and $d = 1$. Consequently B is a group automorphism of a group of order p . Hence B is cyclic of order dividing $p - 1$.

Show that $[A, B] = 1$. We fix an element $[b^{-1}, a^{-1}] \in [A, B]$. Since P is normal in G , it follows that $x^a \in P$ for any $a \in A$ and any $x \in P$. Hence

$$\begin{aligned} x^{[b^{-1}, a^{-1}]} &= x^{bab^{-1}a^{-1}} = (x^b)^{ab^{-1}a^{-1}} = \\ &= ((x^{n_b})^a)^{b^{-1}a^{-1}} = ((x^a)^{n_b})^{b^{-1}a^{-1}} = ((x^a)^b)^{b^{-1}a^{-1}} = (x)^{abb^{-1}a^{-1}} = x. \end{aligned}$$

Therefore $[b^{-1}, a^{-1}] \in C_G(P) = P$. Since $[A, B] \leq M$, we have $[b^{-1}, a^{-1}] \in M \cap P = 1$ and $[A, B] = 1$. \square

2 Properties of msp-permutable subgroups

We will say that a group G satisfies the property:

- E_π if G has at least one Hall π -subgroup;
- C_π if G satisfies E_π and any two Hall π -subgroups of G are conjugate in G ;
- D_π if G satisfies C_π and every π -subgroup of G is contained in some Hall π -subgroup of G .

Such a group is also called an E_π -group, C_π -group, and D_π -group, respectively.

Lemma 2.1. *Let A and B be msp-permutable subgroups of G and $G = AB$.*

(1) *If N is a normal subgroup of G , then $G/N = (AN/N)(BN/N)$ is the msp-permutable product of subgroups AN/N and BN/N .*

(2) *If $A \leq H \leq G$, then H is the msp-permutable product of subgroups A and $H \cap B$.*

(3) *If $G \in D_\pi$, then there exist Hall π -subgroups G_π , A_π and B_π of G , of A and, of B , respectively, such that $G_\pi = A_\pi B_\pi$ is the msp-permutable product of subgroups A_π and B_π .*

Proof. 1. Let $p \in \pi(AN/N)$, X/N be a Sylow p -subgroup of AN/N and P be a Sylow p -subgroup of A . Then $PN/N = X/N$. Similarly, if $q \in \pi(BN/N)$ such that $q \neq p$, Y/N is a Sylow q -subgroup of BN/N and Q is a Sylow q -subgroup of B . Then $QN/N = Y/N$. By hypothesis, P and Q are mutually permutable. Hence X/N and Y/N are mutually permutable.

2. By Dedekind's identity, $H = A(H \cap B)$. Let A_q be a Sylow q -subgroup of A , R be a Sylow r -subgroup of $H \cap B$, where $q \neq r$, and B_r be a Sylow

r -subgroup of B containing R . Since $(H \cap B_r)$ is a Sylow r -subgroup of $H \cap B$ and $R \leq H \cap B_r$, it follows that $R = H \cap B_r$.

Because A_q and B_r are mutually permutable, we have $A_q U \leq G$ for every subgroup U of R .

Let V be an arbitrary subgroup of A_q . Since A_q and B_r are mutually permutable,

$$VB_r \leq G, \quad H \cap VB_r = V(H \cap B_r) = VR \leq G.$$

Hence A_q and R are mutually permutable.

3. By [3, Theorem 1.1.19], there are Hall π -subgroups G_π , A_π and B_π of G , of A , and of B , respectively, such that $G_\pi = A_\pi B_\pi$. Since A and B are msp-permutable, it follows that obviously, A_π and B_π are msp-permutable. \square

Lemma 2.2. *Let A and B be msp-permutable subgroups of G and $G = AB$. Let $p, r \in \pi(G)$, p be the greatest prime in $\pi(G)$ and r be the smallest prime in $\pi(G)$. Then the following statements hold:*

- (1) *if A and B are p -closed, then G is p -closed;*
- (2) *if A and B are r -nilpotent, then G is r -nilpotent;*
- (3) *if A and B have an ordered Sylow tower of supersoluble type, then G has an ordered Sylow tower of supersoluble type.*

Proof. 1. By [3, Theorem 1.1.19], there are Sylow p -subgroups P , P_1 and P_2 of G , of A , and of B , respectively, such that $P = P_1 P_2$. By hypothesis, P_1 is normal in A and P_2 is normal in B . Let H_1 and H_2 be Hall p' -subgroups of A and of B , respectively, and Q be a Sylow q -subgroup of H_1 , where $q \in \pi(H_1)$. Choose a chain of subgroups

$$1 = Q_0 < Q_1 < \dots < Q_{t-1} < Q_t = Q, \quad |Q_{i+1} : Q_i| = q.$$

Since A and B are msp-permutable, we have $P_2 Q_i$ is a subgroup of G for every i . Since $|P_2 Q_1 : P_2| = q$ and $p > q$, it follows that P_2 is normal in $P_2 Q_1$. Then by induction, we have that P_2 is normal in $P_2 Q$. Because q is an arbitrary prime in $\pi(H_1)$, it follows that P_2 is normal in $P_2 H_1$ and $\langle H_1, H_2 \rangle \leq N_G(P_2)$. Similarly, $\langle H_1, H_2 \rangle \leq N_G(P_1)$. Hence $P = P_1 P_2$ is normal in G .

2. Let R , R_1 and R_2 be Sylow r -subgroups of G , of A , and of B , respectively, such that $R = R_1 R_2$. Let K_1 and K_2 be Hall r' -subgroups of A and of B . Let $q \in \pi(G) \setminus \{r\}$, Q , Q_1 and Q_2 be Sylow q -subgroups of G , of A , and of B , respectively, such that $Q = Q_1 Q_2$. Choose a chain of subgroups

$$1 = V_0 < V_1 < \dots < V_{t-1} < V_t = R_1, \quad |V_{i+1} : V_i| = r.$$

Since A and B are msp-permutable, $V_i Q_2$ is a subgroup of G for every i . Since $|V_1 Q_2 : Q_2| = r$ and $q > r$, it follows that Q_2 is normal in $V_1 Q_2$. Then

by induction, we have that $R_1 \leq N_G(Q_2)$. By hypothesis, A is r -nilpotent, hence $R_1 \leq N_G(Q_1)$ and $R_1 \leq N_G(Q)$. Similarly, $R_2 \leq N_G(Q)$ and G has a r -nilpotent Hall $\{r, q\}$ -subgroup RQ . Since q is an arbitrary prime in $\pi(G) \setminus \{r\}$, it follows that G is soluble and r -nilpotent by [11, Corollary].

3. By (1), we have that a Sylow p -subgroup P is normal in G for the greatest $p \in \pi(G)$. By Lemma 2.1 (1), G/P is the product of msp-permutable subgroups AN/N and BN/N . By induction, G/N has an ordered Sylow tower of supersoluble type, hence G has an ordered Sylow tower of supersoluble type. \square

Theorem 2.1. *Let A and B be msp-permutable subgroups of G and $G = AB$. If A and B are soluble, then G is soluble.*

Proof. We use induction on the order of G and the method of proof from [7, Theorem 2]. Let $N \neq 1$ be a soluble normal subgroup of G . By Lemma 2.1 (1), G/P is the product of soluble msp-permutable subgroups AN/N and BN/N . By induction, G/N is soluble, hence G is soluble. In what follows, we assume that G contains no non-trivial soluble normal subgroups.

Since A is soluble, $U = O_s(A) \neq 1$ for some $s \in \pi(A)$. If B is an s -subgroup of G , then $G = AG_s$, $U \leq G_s$ and $U^G \leq (G_s)_G$ by Lemma 1.5, a contradiction. Hence B is not s -subgroup of G and let Q be an arbitrary Sylow q -subgroup of B , where $q \in \pi(B) \setminus \{s\}$. Since A and B are msp-permutable,

$$UQ^x = UQ^{ba} = U^a(Q^b)^a = (UQ^b)^a = (Q^bU)^a = Q^xU$$

for every $x = ba \in G$, where $b \in B$ and $a \in A$. By [6, Theorem 7.2.5], $D = U^Q \cap Q^U$ is subnormal in G . Since $U^Q \leq UQ$ and UQ is soluble, it follows that D is a soluble subnormal subgroup of G and $D = 1$. Hence

$$[U, Q] \leq [U^Q, Q^U] \leq D = 1.$$

This is true for any Sylow q -subgroup of B , therefore $[U, Q^B] = 1$.

Let $H = N_G(U)$. By Dedekind's identity, $H = A(H \cap B)$. By Lemma 2.1 (2), H is the product of soluble msp-permutable subgroups A and $H \cap B$. By induction, H is soluble. Since $[U, Q^B] = 1$, we have $Q^B \leq N_G(U) = H$. Because $G = AB = HB$, Q^B is normal in B and $Q^B \leq H$, it follows that $Q^B \leq H_G = 1$ by Lemma 1.5, a contradiction. \square

Lemma 2.3. *Let $G = G_1G_2$ be the product of msp-permutable subgroups G_1 and G_2 . If a Sylow p -subgroup P of G is normal in G and abelian, then $P \cap G_i$ is normal in G for every $i \in \{1, 2\}$.*

Proof. Assume that $i, j \in \{1, 2\}$ and $i \neq j$. It is clear that $P \cap G_i$ is a Sylow p -subgroup of G_i and $P \cap G_i = (G_i)_p$ is normal in G_i . Hence G_i has a Hall p' -subgroup $(G_i)_{p'}$. Since G_i and G_j are msp-permutable, it follows that $(G_i)_p(G_j)_{p'}$ is a subgroup of G and $(G_j)_{p'} \leq N_G((G_i)_p)$, because every subgroup of G is p -closed. By hypothesis, P is abelian, therefore $(G_i)_p$ is normal in P and

$$G_j = (G_j)_p(G_j)_{p'} = (P \cap G_j)(G_j)_{p'} \leq N_G((G_i)_p).$$

Hence $(G_i)_p$ is normal in $G = G_i G_j = G_1 G_2$ for every $i \in \{1, 2\}$. \square

3 Proof of the main theorem

Theorem 3.1. *Let \mathfrak{F} be a subgroup-closed saturated formation such that $\mathfrak{U} \subseteq \mathfrak{F} \subseteq \mathcal{D}$. Let $G = G_1 G_2$ be the product of msp-permutable subgroups G_1 and G_2 . If $G_1, G_2 \in \mathfrak{F}$, then $G \in \mathfrak{F}$.*

Proof. By Lemma 2.2 (3), G has an ordered Sylow tower of supersoluble type. Let P be a Sylow p -subgroup of G , where p is the greatest prime in $\pi(G)$. Then P is normal in G .

Assume that $G \notin \mathfrak{F}$. Let N be a non-trivial normal subgroup of G . Hence

$$G/N = (G_1 N/N)(G_2 N/N),$$

$$G_1 N/N \cong G_1/G_1 \cap N \in \mathfrak{F}, \quad G_2 N/N \cong G_2/G_2 \cap N \in \mathfrak{F}.$$

By Lemma 2.1 (1), $G_1 N/N$ and $G_2 N/N$ are msp-permutable. Consequently, G/N satisfies the hypothesis of the theorem, and by induction, $G/N \in \mathfrak{F}$. Since \mathfrak{F} is saturated, G is primitive by Lemma 1.1. Hence $\Phi(G) = 1$, $G = N \rtimes M$, where $N = C_G(N) = F(G) = O_p(G) = P$ is a unique minimal normal subgroup of G by Lemma 1.2. Therefore M is a Hall p' -subgroup of G and $M = (G_1)_{p'}(G_2)_{p'}$ for some Hall p' -subgroups $(G_1)_{p'}$ and $(G_2)_{p'}$ of G_1 and of G_2 , respectively.

Suppose that p divides $|G_1|$ and $|G_2|$. By Lemma 2.3, $P \leq G_1 \cap G_2$. Let $P_1 \leq P$ and $|P_1| = p$. Since $P \leq G_1$ and Q permutes with P_1 for every Sylow subgroup Q of $(G_2)_{p'}$, we have $P_1(G_2)_{p'} \leq G$ and $(G_2)_{p'} \leq N_G(P_1)$. Similarly, since $P \leq G_2$ and R permutes with P_1 for every Sylow subgroup R of $(G_1)_{p'}$, it follows that $P_1(G_1)_{p'} \leq G$ and $(G_1)_{p'} \leq N_G(P_1)$. Hence $M = (G_1)_{p'}(G_2)_{p'} \leq N_G(P_1)$ and P_1 is normal in G . By Lemma 1.3, $G \in \mathfrak{F}$, a contradiction.

Thus $P \leq G_1$ and G_2 is a p' -subgroup of G . By Lemma 1.6 (1), G_2 is a cyclic group of order dividing $p - 1$. Hence $G_2 \in g(p)$, where g is a

canonical local definition of a saturated formation \mathfrak{U} . Since $\mathfrak{U} \subseteq \mathfrak{F}$, we have by [5, Proposition IV.3.11], $g(p) \subseteq f(p)$, where f is a canonical local definition of a saturated formation \mathfrak{F} . Hence $G_2 \in f(p)$. Since $P \leq G_1$, it follows that $G_1 = P \rtimes (G_1)_{p'}$. Because $G_1 \in \mathfrak{F}$ and $F_p(G_1) = P$, we have $G_1/F_p(G_1) = G_1/P \cong (G_1)_{p'} \in f(p)$. By Lemma 1.6 (2), $[(G_1)_{p'}, (G_2)_{p'}] = 1$. Since $(G_1)_{p'} \in f(p)$, $(G_2)_{p'} \in f(p)$ and $f(p)$ is a formation, it follows that by Lemma 1.4, $G/P \cong M = (G_1)_{p'}(G_2)_{p'} \in f(p)$. Because $P \in \mathfrak{N}_p$, we have $G \in \mathfrak{F}$, a contradiction. The theorem is proved. \square

Corollary 3.1. *Let $G = G_1G_2$ be the product of msp-permutable subgroups G_1 and G_2 .*

1. *If $G_1, G_2 \in \mathfrak{U}$, then $G \in \mathfrak{U}$.*
2. *If $G_1, G_2 \in \text{w}\mathfrak{U}$, then $G \in \text{w}\mathfrak{U}$.*
3. *If $G_1, G_2 \in \text{v}\mathfrak{U}$, then $G \in \text{v}\mathfrak{U}$.*

References

- [1] Asaad, M., Shaalan, A.: On supersolvability of finite groups. Arch. Math. **53**, 318–326 (1989).
- [2] Ballester-Bolínches, A., Ezquerro, L.M.: Classes of Finite Groups. Springer, Dordrecht (2006).
- [3] Ballester-Bolínches, A., Esteban-Romero, R., Asaad, M.: Products of finite groups. Walter de Gruyter, Berlin (2010).
- [4] Chuniĭin, S.A.: Simplicité de groupe fini et les ordres de ses classes d'éléments conjugués. C. R. Acad. Sci. Paris. **191**, 397–399 (1930).
- [5] Doerk, K., Hawkes, T.: Finite Soluble Groups. Walter De Gruyter, Berlin, New York (1992).
- [6] Lennox, J.C., Stonehewer, S.E.: Subnormal subgroups of groups. Clarendon Press, Oxford (1987).
- [7] Monakhov, V.S.: On the solvability of a group with permutable subgroups. Mathematical Notes. **93**, 460–464 (2013).
- [8] Monakhov, V.S., Kniahina, V.N.: Finite group with \mathbb{P} -subnormal subgroups. Ricerche Mat. **62** (2), 307–323 (2013).
- [9] Monakhov, V.S.: Finite groups with abnormal and \mathfrak{U} -subnormal subgroups. Siberian Math. J. **57** (2), 352–363 (2016).

- [10] Skiba, A.N.: On weakly s -permutable subgroups of finite groups. *J. Algebra*. **315**, 192–209 (2007).
- [11] Tyutyanov, V.N., Kniahina, V.N.: Finite groups with biprimary Hall subgroups. *J. Algebra*. **443**, 430–440 (2015).
- [12] Vasil'ev, A.F., Vasil'eva, T.I., Tyutyanov, V.N.: On the finite groups of supersoluble type. *Siberian Math. J.* **51** (6), 1004–1012 (2010).