# Reversals of Rényi Entropy Inequalities under Log-Concavity

James Melbourne University of Minnesota melbo013@umn.edu Tomasz Tkocz Carnagie Mellon University ttkocz@andrew.cmu.edu

May 25, 2020

#### Abstract

We establish a discrete analog of the Rényi entropy comparison due to Bobkov and Madiman. For log-concave variables on the integers, the min entropy is within  $\log e$  of the usual Shannon entropy. Additionally we investigate the entropic Rogers-Shephard inequality studied by Madiman and Kontoyannis, and establish a sharp Rényi version for certain parameters in both the continuous and discrete cases.

## 1 Introduction

The Rényi entropy [33] is a family of entropies parameterized by an  $\alpha$  belonging to the extended non-negative real line that generalizes several important notions of uncertainty, most significantly the Shannon entropy when  $\alpha = 1$ , but also the min entropy  $\alpha = \infty$ , the collision entropy  $\alpha = 2$ , and the Hartley or max-entropy  $\alpha = 0$ .

**Definition 1.1.** For a random variable X taking countably many values with probabilities  $p_i > 0$ , and  $\alpha \in (0,1) \cup (1,\infty)$ , the Rényi entropy  $H_{\alpha}(X)$  is

$$H_{\alpha}(X) = \frac{\log \sum_{i} p_{i}^{\alpha}}{1 - \alpha},\tag{1}$$

while the remaining  $\alpha$  are defined through continuous limits.  $H_0(X) = \log |\{i : p_i > 0\}|$ ,  $H_1(X) = H(X) = -\sum_i p_i \log p_i$ , and  $H_{\infty}(X) = -\log ||p||_{\infty}$ , where  $|\cdot|$  denotes cardinality, and  $||p||_{\infty}$  denotes the  $\ell_{\infty}$  norm of the sequence  $p_i$ .

Similar definitions are put forth in the continuous setting.

**Definition 1.2.** For an  $\mathbb{R}^d$ -valued random variable Y with density function f, and  $\alpha \in (0,1) \cup (1,\infty)$ , the Rényi entropy  $h_{\alpha}(Y)$  is

$$h_{\alpha}(Y) = \frac{\log \int_{\mathbb{R}^d} f^{\alpha}(x) dx}{1 - \alpha},\tag{2}$$

while the remaining  $\alpha$  are defined through continuous limits.  $h_0(X) = \log |\{x : f(x) > 0\}|$ ,  $h_1(X) = h(X) = -\int f(x) \log f(x) dx$ , and  $h_{\infty}(X) = -\log ||f||_{\infty}$ , where  $|\cdot|$  denotes Lebesgue volume, and  $||f||_{\infty}$  denotes the  $L_{\infty}$  norm with respect to the Lebesgue measure.

The Rényi entropy can usefully be described in terms of the  $\alpha - 1$ -norm<sup>1</sup>,

$$H_{\alpha}(X) = \frac{\log \sum_{i} p_{i}^{\alpha - 1} p_{i}}{\alpha - 1} = -\log ||p||_{\alpha - 1, p}$$
(3)

where for a positive function q and measure  $\mu$  on a discrete set, and  $s \in (-\infty, 0) \cup (0, \infty)$  we define

$$||q||_{s,\mu} = \left(\sum_{i} q_i^s \mu_i\right)^{\frac{1}{s}}.$$
 (4)

Thus it follows from Jensen's inequality that  $H_p(X)$  is non-increasing in  $\alpha$ . The same argument in the continuous setting obviously holds. Thus, for  $\alpha < \beta$ 

$$H_{\beta}(X) \le H_{\alpha}(X) \tag{5}$$

$$h_{\beta}(Y) \le h_{\alpha}(Y). \tag{6}$$

It has been known since [3] (see also [2]) that inequality (6) can be reversed up to an additive constant for log-concave random variables.

**Theorem 1.1** (Bobkov-Madiman, [3]). Let Y be random vector in  $\mathbb{R}^d$  with density  $f: \mathbb{R}^d \to [0,\infty)$  such that  $f((1-t)x+ty) \geq f^{1-t}(x)f^t(y)$  for all  $t \in (0,1)$  and  $x,y \in \mathbb{R}^d$ . Then

$$h(Y) \le h_{\infty}(Y) + d. \tag{7}$$

Recently (see [11, 18]), a sharp comparison result has been also established between entropies of arbitrary orders.

**Theorem 1.2** (Fradelizi-Li-Madiman-Wang, [11, 18]). Under the assumptions of Theorem 1.1, for  $\alpha < \beta$ , we have

$$h_{\alpha}(Y) - h_{\beta}(Y) \le h_{\alpha}(Z) - h_{\beta}(Z),\tag{8}$$

where Z is a random variable with density  $e^{-\sum_{i=1}^{d} x_i}$  on  $(0, \infty)^d$ .

The key to this theorem is the following fact: for f as in Theorem 1.1, the function  $t \mapsto \log(t \int f^t)$  is concave on  $(0, +\infty)$ .

Our pursuit in this paper will be the reversal of (5) for the log-concave random variables on  $\mathbb{Z}$ . That is random variables X which are unimodal, and such that the sequence  $p_n = \mathbb{P}(X = n)$  satisfies  $p_i^2 \geq p_{i-1}p_{i+1}$ . We say that such random variables are monotone (or that their distributions are monotone) if the sequence  $(p_n)$  is monotone. We will prove the following main theorem.

**Theorem 1.3.** For  $\alpha \in (0, \infty)$ , a log-concave random variable X, and  $Z_p$  a geometric random variable with parameter p,

$$H_{\alpha}(X) - H_{\infty}(X) < \lim_{p \to 0} H_{\alpha}(Z_p) - H_{\infty}(Z_p)$$
(9)

$$= \log \alpha^{\frac{1}{\alpha - 1}}.\tag{10}$$

<sup>&</sup>lt;sup>1</sup>This is not truly a norm when  $\alpha < 2$ 

This provides a discrete analog of Bobkov and Madiman's result, with a sharp constant provided by an extremizing sequence. Let us note, this is distinct from the continuous setting where for instance, an exponential distributions extremizes  $h(X) - h_{\infty}(X)$  over log-concave distributions on  $\mathbb{R}$ . It is consequence of the proof of Theorem 1.3, that  $H_{\alpha}(X) - H_{\infty}(X) < \log \alpha^{\frac{1}{\alpha-1}}$ , for X log-concave, and thus that no such extremizer exists in the discrete setting.

The proof methods in the discrete setting are distinct as well. In the continuous setting, there are analytic and convexity tools that can be leveraged, the Prékopa-Leindler inequality, integration by parts, perspective function convexity, etc.

Though there has been significant recent interest in developing discrete versions of these continuous techniques (see [10, 16, 31, 29]), the authors are not aware of a version that could deliver Theorem 1.3. Our proof method will show that any log-concave distribution majorizes a "two-sided geometric distribution" with the same  $\infty$ -Rényi entropy. Using the Schur concavity of the Rényi entropy the problem is reduced to direct computation on these two-sided geometric distributions.

First progress on this result was made by the authors in [30] by a similar argument, which only applied however to monotone log-concave variables, and was leveraged by the concavity of Shannon entropy to deliver a sub-optimal reversal,  $H(X) - H_{\infty}(X) \leq \log 2e$  for general log-concave variables. In contrast the results here,  $H_{\alpha}(X) - H_{\infty}(X) \leq \log \alpha^{\frac{1}{\alpha-1}}$  are sharp for all  $\alpha$ , and they apply to general log-concave variables, without the additional assumption of monotonicity.

As an application we investigate a conjecture of Madiman and Kontoyannis [26], which can be regarded as an entropic analog of the Rogers-Shephard inequality from Convex Geometry. We consider general orders of the Rényi entropy, where the inequality can be considered a reversal, under the assumption of log-concavity, of the Rényi entropy power inequalities that have attracted recent attention (see [4, 5, 7, 17, 19, 20, 24, 28, 27, 32, 34]).

Let us outline the paper. In the next section we gather basic definitions and properties regarding log-concave sequences and majorization. In Section 3 we prove the desired inequalities in the case that the distribution considered is two-sided geometric. In Section 4 we show that every log-concave distribution majorizes a two-sided geometric distribution with the same min-entropy, and use the Schur concavity of the Rényi entropy to reduce the problem to the result proven in Section 3. In Section 5 we discuss a conjecture which would imply a sharp reversal of (5) for all Rényi entropies (not just with comparison to min-entropy), but only for monotone log-concave distributions. This conjecture amounts to the discrete analog of a result that does hold for all log-concave densities in  $\mathbb{R}^d$ , however as we show, in the discrete case, the result fails without the assumption of monotonicity. In Section 6, we establish Rényi versions of the Rogers-Shephard inequality in both the continuous and discrete cases.

**Acknowledgements.** We would like to thank Mokshay Madiman and Arnaud Marsiglietti for fruitful discussions throughout the development of the paper. TT's research is partially supported by NSF grant DMS-1955175.

### 2 Preliminaries

We will use the following notation for integer intervals. For  $a \leq b \in \mathbb{Z}$ ,  $[\![a,b]\!] \coloneqq \{x \in \mathbb{Z} : a \leq x \leq b\}$ ,  $[\![a,b]\!] \coloneqq \{x \in \mathbb{Z} : a \leq x \leq b\}$  and so on. We also let

 $\llbracket a, \infty \llbracket \text{ denote } \{x \in \mathbb{Z} : a \le x\}$ 

**Definition 2.1** (Log-concavity). A sequence  $f: \mathbb{Z} \to [0, \infty)$  is log-concave when it satisfies

$$f^{2}(n) \ge f(n-1)f(n+1) \tag{11}$$

for all  $n \in \mathbb{Z}$  and  $a \leq b \subseteq \{f > 0\}$  implies  $[a,b] \subseteq \{f > 0\}$  (in other words, the support of f is a contiguous interval of integers). A  $\mathbb{Z}$ -valued random variable X is log-concave when the sequence  $p_i = \mathbb{P}(X = i)$  is log-concave. We will denote the space of all log-concave probability densities on  $\mathbb{Z}$  by  $\mathcal{L}(\mathbb{Z})$ .

There have already been a few interesting information theoretic results regarding log-concavity in discrete settings, see for example [14, 15], but there is a vast mathematical literature, we mention only [36] and recall some relevant facts. The class of log-concave sequences is closed under convolution, and thus the log-concave random variables are closed under independent summation. The class is also closed under weak limits. Important examples of log-concave distributions are the Bernoulli, Binomial, Geometric, Negative Binomial, and Poisson distribution.

Let  $\ell_1(\mathbb{Z})$  denote the functions  $f: \mathbb{Z} \to \mathbb{R}$  such that  $\sum_{i \in \mathbb{Z}} |f(i)| < \infty$ .

**Definition 2.2** (Decreasing rearrangment). For a function  $f: \mathbb{Z} \to [0, \infty)$  in  $\ell_1(\mathbb{Z})$  denote the sequence  $f_i^{\downarrow}$  to be the decreasing rearrangment of f. Explicitly,  $f^{\downarrow}$  satisfies  $f_i^{\downarrow} \geq f_{i+1}^{\downarrow}$  for all i and  $f_i^{\downarrow} = f(\tau(i))$  for a bijection  $\tau: \mathbb{N} \to \mathbb{Z}$ .

**Definition 2.3** (Majorization). For  $\ell_1(\mathbb{Z})$  functions  $f, g : \mathbb{Z} \to [0, \infty)$ , we say f majorizes g and write  $f \succ g$  when

$$\sum_{i=1}^{k} f_i^{\downarrow} \ge \sum_{i=1}^{k} g_i^{\downarrow},\tag{12}$$

and equality holds in the limit with  $k \to \infty$ .

We let  $\mathcal{P}(\mathbb{Z})$  denote the space of probability densities on  $\mathbb{Z}$ .

**Definition 2.4** (Schur Convexity). A function  $\Phi : \mathcal{P}(\mathbb{Z}) \to \mathbb{R} \cup \{\infty\}$  is Schur-convex when  $f \succ g$  implies

$$\Phi(f) \ge \Phi(g). \tag{13}$$

 $\Phi$  is Schur-concave when  $-\Phi$  is Schur-convex.

**Definition 2.5** (Two-sided geometric distribution). A density function  $\varphi$  on  $\mathbb{Z}$  is a two-sided geometric distribution when there exists  $p, q \in [0, 1)$  and  $m \in \mathbb{Z}$  such that its density function  $\varphi$  can be expressed as

$$\varphi(n) = \frac{(1-p)(1-q)}{1-pq} f(n-m). \tag{14}$$

with

$$f(n) = \begin{cases} p^n & \text{for } n \ge 0\\ q^{-n} & \text{for } n \le 0. \end{cases}$$
 (15)

with the convention that  $0^0 = 1$ . We will denote the set of all such densities by  $\mathcal{E}(\mathbb{Z})$ .

For q=0 and m=0,  $\varphi$  is the usual geometric distribution with parameter 1-p, and when p=q=0 we have a point mass at m. In any case, observe that all two-sided geometric distributions are log-concave, since their support is by definition contiguous,  $\varphi^2(n)=\varphi(n+1)\varphi(n-1)$  for  $n\neq m$  and  $\varphi^2(m)=\max_n \varphi^2(n)\geq \varphi(m-1)\varphi(m+1)$ .

## 3 Two-sided geometric distributions

In this section we will derive bounds on the Rényi entropy on the two-sided geometric distribution.

**Lemma 1.** If  $\varphi$  is a density function on  $\mathbb{Z}$  with a two sided geometric distribution with parameters p and q, then for  $\alpha \in (0, \infty)$ , we have

$$H_{\alpha}(\varphi) - H_{\infty}(\varphi) < \frac{\log \alpha}{\alpha - 1}.$$
 (16)

with  $\frac{\log \alpha}{\alpha - 1}\Big|_{\alpha = 1} := \log e$ .

*Proof.* First we proceed with the case  $\alpha \neq 1$ . By direct computation,

$$H_{\alpha}(\varphi) - H_{\infty}(\varphi) = (1 - \alpha)^{-1} \log \left( \sum_{n} \varphi^{\alpha}(n) \right) + \log \|\varphi\|_{\infty}$$
(17)

$$= (1 - \alpha)^{-1} \left( \log \|\varphi\|_{\infty}^{\alpha} + \log \left( \frac{1}{1 - p^{\alpha}} + \frac{1}{1 - q^{\alpha}} - 1 \right) \right) + \log \|\varphi\|_{\infty}$$
 (18)

$$= \frac{\log \|\varphi\|_{\infty} + \log \left(\frac{1}{1-p^{\alpha}} + \frac{1}{1-q^{\alpha}} - 1\right)}{1-\alpha} \tag{19}$$

$$= \frac{\log\left(\frac{\frac{1}{1-p^{\alpha}} + \frac{1}{1-q^{\alpha}} - 1}{\frac{1}{1-p} + \frac{1}{1-q} - 1}\right)}{1 - \alpha}.$$
 (20)

Thus, it suffices to prove for  $\alpha > 1$ 

$$\frac{1}{1-p} + \frac{1}{1-q} - 1 < \alpha \left( \frac{1}{1-p^{\alpha}} + \frac{1}{1-q^{\alpha}} - 1 \right)$$
 (21)

and

$$\frac{1}{1-p} + \frac{1}{1-q} - 1 > \alpha \left( \frac{1}{1-p^{\alpha}} + \frac{1}{1-q^{\alpha}} - 1 \right)$$
 (22)

when  $\alpha \in (0,1)$ . Note that we have equality when  $\alpha = 1$ , so it suffices to show the function,

$$F(\alpha) = \alpha \left( \frac{1}{1 - p^{\alpha}} + \frac{1}{1 - q^{\alpha}} - 1 \right)$$
 (23)

is strictly increasing. Computing directly,

$$F'(\alpha) = \frac{p^{\alpha} \log p^{\alpha}}{(1 - p^{\alpha})^2} + \frac{q^{\alpha} \log q^{\alpha}}{(1 - q^{\alpha})^2} + \frac{1}{1 - p^{\alpha}} + \frac{1}{1 - q^{\alpha}} - 1.$$
 (24)

Writing  $x = p^{\alpha}$  and  $y = q^{\alpha}$  it is enough to prove

$$\frac{x\log x}{(1-x)^2} + \frac{y\log y}{(1-y)^2} + \frac{1}{1-x} + \frac{1}{1-y} > 1$$
 (25)

for  $x, y \in (0, 1)$ . To this end, we consider  $f(x) = \frac{x \log x}{(1-x)^2} + \frac{1}{1-x}$ ,  $x \in (0, 1)$  and argue that  $f(x) > \frac{1}{2}$  for  $x \in (0, 1)$ . Note that  $(1-x)^3 f'(x) = 2(1-x) + (1+x) \log x$  and the right

hand side, call it g(x), is an increasing function of x on (0,1) because  $g'(x) = \log x + \frac{1}{x} - 1 = -\log \frac{1}{x} + \frac{1}{x} - 1 > 0$ . Thus g(x) < g(1) = 0 for  $x \in (0,1)$  which shows that f'(x) < 0 for  $x \in (0,1)$ . Thus f is strictly decreasing and we get  $f(x) > f(1-) = \frac{1}{2}$  for  $x \in (0,1)$ .

When  $\alpha = 1$ , a direct computation gives

$$H(\varphi) - H_{\infty}(\varphi) = -\frac{\frac{p \log p}{(1-p)^2} + \frac{q \log q}{(1-q)^2}}{\frac{1}{1-p} + \frac{1}{1-q} - 1}$$

which is strictly less than 1 by (25).

### 4 Proof of Theorem 1.3

**Lemma 2.** For a non-negative monotone log-concave  $\ell_1(\mathbb{Z})$  function f, supported on ]n, k[], there exists a unique log-affine positive function g with infinite set of support containing ]n, k[], the same mode, maximum value, and  $\ell_1(\mathbb{Z})$  norm such that  $f \succ g$ .

Proof. Without loss of generality, let us assume that f is non-increasing and supported on [0, k[. Define for  $q \in [0, 1)$  a new function  $g_q$  to be log-affine supported on  $[0, \infty]$  and such that  $g_q(1) = f(1)$  and g(j+1) = qg(j). Note that  $\sum_j g_0(j) \leq \sum_j f(j) < \lim_{q \to 1} \sum_j g_q(j) = \infty$  and the function  $q \mapsto \sum_j g_q(j)$  is strictly increasing in q. Thus, by the intermediate value theorem, there exists a unique  $q^* \in [0,1]$  such that  $\sum_j g_{q^*}(j) = \sum_j f(j)$ . Take  $g = g_{q^*}$ , and observe that g is log-affine and satisfies  $||g||_{\infty} = ||f||_{\infty}$  by construction. Since g is log-affine and f is log-concave, for the function f-g there exists and  $f \in [1,n]$  such that  $f(k)-g(k) \geq 0$  on [1,l] and  $f(k)-g(k) \leq 0$ .

Using the notation  $\mathcal{L}_m(\mathbb{Z}) = \{ f \in \mathcal{L}(\mathbb{Z}) : H_\infty(f) = m \}$  and  $\mathcal{E}_m(\mathbb{Z}) = \{ f \in \mathcal{E}(\mathbb{Z}) : H_\infty(f) = m \}$ , we will prove the following general result, that roughly states minimizers of Schur-convex functions among log-concave densities satisfying a constraint on their maximum value, are two-sided geometric distributions.

**Theorem 4.1.** For a Schur convex function,  $\Phi : \mathcal{P}(\mathbb{Z}) \to \mathbb{R}$ , and  $f \in \mathcal{L}_m(\mathbb{Z})$  there exists  $\varphi \in \mathcal{E}_m(\mathbb{Z})$  such that

$$\Phi(f) \ge \Phi(\varphi). \tag{26}$$

Proof. We will prove the result by showing that given a log-concave density f there exists a two-sided geometric distribution  $\varphi$  with the same maximum such that  $f \succ \varphi$ . Without loss of generality, suppose that  $f(0) = ||f||_{\infty}$  and  $t \coloneqq \sum_{j=0}^{\infty} f(j) \ge \sum_{j=-\infty}^{0} f(j)$ , note  $t \ge \frac{1}{2}$ . Define  $f_+(j) = \mathbbm{1}_{[0,\infty)}(j)f(j)$  and  $f_-(j) = \mathbbm{1}_{(-\infty,0]}(j)f(j)$ , define  $a_+$  to be the log-affine function of the same maximum and majorized by  $f_+$ , while  $a_-$  to be the log-affine function of the same maximum and majorized by  $f_-$ , as supplied by Lemma 2. Note that  $a_+(j) = f(0)p^j$  for  $j \ge 0$  and  $p \in [0,1)$  while  $a_-(k) = f(0)q^{-j}$  for  $j \le 0$  for  $q \in [0,1)$ . Note that by our assumptions  $\sum_{j=0}^{\infty} a_+(j) = \sum_{j=0}^{\infty} f(j) \ge \sum_{j=-\infty}^{0} f(j) = \sum_{j=-\infty}^{0} a_-(j)$  implies  $p \ge q$ . Define

$$\varphi(j) = \begin{cases} a_{+}(j), & j > 0 \\ f(0), & j = 0 \\ a_{-}(j), & j < 0. \end{cases}$$
 (27)

By construction  $\sum_i \varphi_i = \sum_i f(i)$ . In particular,  $\varphi$  is a density function on  $\mathbb{Z}$ . Observe that  $\varphi$  is the two-sided geometric distribution majorized by f as desired, since for every n

$$\sum_{i=1}^{n} \varphi_i^{\downarrow} = f(0) + \sum_{i=1}^{j} a_+(i) + \sum_{i=1}^{n-j-1} a_-(i)$$
 (28)

$$\leq f(0) + \sum_{i=1}^{j} f_{+}(i) + \sum_{i=1}^{n-j-1} f_{-}(i)$$
(29)

$$\leq \sum_{i=1}^{n} f_i^{\downarrow} \tag{30}$$

Thus  $\varphi \prec f$  and by Schur-convexity  $\Phi(\varphi) \leq \Phi(f)$ , and thus the proof is complete.  $\square$ 

Proof of Theorem 1.3. As is well known, see [25, 13] the Rényi entropy is Schur-concave. Given  $f \in \mathcal{L}_m(\mathbb{Z})$ , by Theorem 4.1 there exists  $\varphi \in \mathcal{E}_m(\mathbb{Z})$  such that

$$H_{\alpha}(f) \le H_{\alpha}(\varphi)$$
 (31)

$$< H_{\infty}(\varphi) + \frac{\log \alpha}{\alpha - 1}$$
 (32)

$$=H_{\infty}(f)+\frac{\log\alpha}{\alpha-1},\tag{33}$$

where (32) follows from Lemma 1. This gives the strict inequality of the theorem. To show that it is attained in the limit of geometric distribution is an easy and direct computation.  $\Box$ 

# 5 Varentropy and logconcavity of auxiliary function

In analogy to the continuous case, we pose the following conjecture.

Conjecture 5.1. Let  $(x_n)_{n=1}^N$  be a finite monotone log-concave sequence. Then the function

$$F(t) = \log \left[ t \sum_{n=1}^{N} x_n^t \right]$$

is concave on  $(0, +\infty)$ .

By looking at the second derivative, this conjecture is equivalent to the statement that for every finite monotone log-concave sequence  $(y_n)_{n=1}^N$   $(y_n = x_n^t)$ , we have

$$\frac{\left(\sum y_n \log^2 y_n\right) \left(\sum y_n\right) - \left(\sum y_n \log y_n\right)^2}{\left(\sum y_n\right)^2} \le 1$$

(the left-hand side can be seen as the varentropy – see [11]).

Note that Conjecture 5.1 implies the desired sharp comparison between Rényi entropies of arbitrary two orders for monotone log-concave random variables. Indeed, suppose X is such a random variable. By an approximation argument, we can assume that the support of X is finite, say it is  $\{1, \dots, N\}$  and  $p_n = \mathbb{P}(X = n) > 0$ ,  $n = 1, \dots, N$ , is log-concave. We have

$$H_{\alpha}(X) = \frac{1}{1-\alpha} \log \left[ \sum_{n=1}^{N} p_n^{\alpha} \right] = \log \left( \alpha^{\frac{1}{\alpha-1}} \right) + F(\alpha), \tag{34}$$

where  $F(\alpha)$  is the function from Conjecture 5.1 for the sequence  $(p_n)$ . Suppose  $1 < \alpha < \beta$  and write  $\alpha = 1 - \lambda + \lambda \beta$  with  $\lambda = \frac{\alpha - 1}{\beta - 1} \in (0, 1)$ . If F was concave, then we would have

$$F(\alpha) = F((1 - \lambda) \cdot 1 + \lambda \beta) \ge (1 - \lambda)F(1) + \lambda F(\beta) = \lambda F(\beta)$$

which, by (34), becomes

$$H_{\alpha}(X) - H_{\beta}(X) \le \log\left(\frac{\alpha^{\frac{1}{\alpha - 1}}}{\beta^{\frac{1}{\beta - 1}}}\right).$$
 (35)

Proceeding in a similar way, this would also follow for  $\alpha < \beta < 1$  as well as  $\alpha < 1 < \beta$ .

As opposed to the continuous case, Conjecture 5.1 cannot hold without the monotonicity assumption: for example, for the sequence  $p=(\frac{1}{4},\frac{1}{2},1,\frac{1}{2},\frac{1}{4})$  the function F is not concave because we have F''(3)>0.0009. This is a consequence of the fact that for a symmetric geometric random variable X with  $\mathbb{P}(X=k)=p^{|k|},\ k\in\mathbb{Z}$ , and  $\alpha<\beta$ , the opposite inequality to (35) holds.

In view of what is true in the continuous case [8] we propose the following strengthening of Conjecture 5.1.

**Conjecture 5.2.** Let  $(y_n)_{n=1}^N$  be a finite positive monotone and concave sequence, that is  $y_n \ge \frac{y_{n-1}+y_{n+1}}{2}$ , 1 < n < N. Then for every  $\gamma > 0$ , the function

$$K(t) = (t + \gamma) \sum_{n=1}^{N} y_n^{t/\gamma}$$

is log-concave, that is  $\log K(t)$  is concave on  $(-\gamma, +\infty)$ .

The approach from [8] also suggests that the following stronger inequality be true: for every complex number z = u + iv with  $u > -\gamma$ , we have  $|K(z)| \ge K(u)$ . It turns out that it implies Conjecture 5.2 (see [8] for a detailed argument).

As a final remark, we also point out that Conjecture 5.1 has an application to Khinchine-type inequalities: it would furnish a large family of examples of the so-called ultra sub-Gaussian random variables, see Remark 13 in [12]. Conjecture 5.1 was verified therein for sequences of length 3.

# 6 Application to Rényi Entropic Rogers-Shephard Inequality

It is a classical theorem of Convex Geometry called the Rogers-Shephard Inequality [35] that for a convex body  $K \subseteq \mathbb{R}^d$  (a compact convex set with non-empty interior), we have

$$Vol(K - K) \le {2d \choose d} Vol(K),$$
 (36)

with equality if and only if K is the d-dimensional simplex. This can be easily rephrased as a Rényi entropic inequality if we recall the usual definition of the Rényi entropy for continuous variables. The Rogers-Shephard inequality thus says that for independent X and Y with a common density function f on  $\mathbb{R}^d$  supported on a convex body in  $\mathbb{R}^d$ ,

$$h_0(X - Y) \le h_0(X) + \log \binom{2d}{d}. \tag{37}$$

Note that by Stirling's formula  $\log {2d \choose d} \approx d \log 4$  and the best possible dimension independent c such that  $h_0(X - Y) \leq h_0(X) + d \log c$  holds is c = 4.

An entropic analog of Rogers-Shephard has been pursued in [22], where the following is conjectured.

Conjecture 6.1 (Madiman-Kontoyannis [22]). For log-concave X and Y iid  $\mathbb{R}^d$  valued random variables

$$h(X - Y) \le h(X) + d\log 2 \tag{38}$$

with the equality case given by the d-dimensional exponential product distribution.

In the same article Madiman and Kontoyannis prove that under the same hypotheses  $h(X-Y) \leq h(X) + d \log 4$ , holds. Moreover using alternate methods in [6] a bound of  $h(X-Y) \leq h(X) + d \log e$ , was obtained. In the continuous case we present the following generalization to the Rényi entropy.

**Theorem 6.1.** Let X and Y be iid log-concave random vectors in  $\mathbb{R}^d$ . If  $\alpha \in [2, \infty]$ , then

$$h_{\alpha}(X - Y) \le h_{\alpha}(X) + d\log 2,\tag{39}$$

with equality when X has exponential distribution  $\mathbb{1}_{(0,\infty)^d}(x)e^{-\sum_i x_i}$ . If  $\alpha \in [0,2]$ , then

$$h_{\alpha}(X - Y) \le h_{\alpha}(X) + d\log \alpha^{\frac{1}{\alpha - 1}}.$$
(40)

Note that when  $\alpha = 1$  we recover [6], the sharpest known bound in the Shannon case  $h(X - Y) \leq h(X) + d \log e$ . We will need the following corollary of Theorem 1.2.

Corollary 6.2 (Fradelizi-Madiman-Wang [11]). For  $0 \le \alpha \le \beta$ , and X a log-concave vector in  $\mathbb{R}^d$ , we have

$$h_{\beta}(X) \le h_{\alpha}(X) \le h_{\beta}(X) + d\log\frac{c(\alpha)}{c(\beta)}$$
 (41)

where

$$c(\alpha) = \alpha^{\frac{1}{\alpha - 1}},\tag{42}$$

with  $c(\infty) := 1, c(1) := e$ , and  $c(0) := \infty$ .

*Proof.* If Z is a random variable with density  $e^{-x}\mathbb{1}_{(0,\infty)}$ , then  $h_{\alpha}(Z) = \frac{\log \int_{0}^{\infty} e^{-\alpha x} dx}{1-\alpha} = \log \alpha^{\frac{1}{\alpha-1}}$ . Let  $Z_d$  be a random vector in  $\mathbb{R}^d$  with density  $e^{-\sum_{i} x_i} \mathbb{1}_{(0,\infty)^d}$ . By Theorem 1.2,

$$h_{\alpha}(X) \le h_{\beta}(X) + h_{\alpha}(Z_d) - h_{\beta}(Z_d) \tag{43}$$

and 
$$h_{\alpha}(Z_d) - h_{\beta}(Z_d) = d(h_{\alpha}(Z) - h_{\beta}(Z)) = d \log \frac{c(\alpha)}{c(\beta)}$$
.

Proof of Theorem 6.1. The result hinges on the following equality,

$$h_{\infty}(X - Y) = h_2(X). \tag{44}$$

Explanation: letting f denote the shared density of X and Y, the density function  $f_{X-Y}$  of X-Y is given by

$$f_{X-Y}(z) = \int_{\mathbb{R}^d} f(z-x)f(-x)dx = \int_{\mathbb{R}^d} f(z+x)f(x)dx,$$
 (45)

which is log-concave and even, thus  $||f_{X-Y}||_{\infty} = f_{X-Y}(0) = \int_{\mathbb{R}^d} f^2(x) dx$ . For  $\alpha \geq 2$ ,

$$h_{\alpha}(X - Y) \le h_{\infty}(X - Y) + d\log\frac{c(\alpha)}{c(\infty)}$$
 (46)

$$= h_2(X) + d\log\frac{c(\alpha)}{c(\infty)} \tag{47}$$

$$\leq h_{\alpha}(X) + d\log\frac{c(2)}{c(\alpha)} + d\log\frac{c(\alpha)}{c(\infty)} \tag{48}$$

$$= h_{\alpha}(X) + d\log\frac{c(2)}{c(\infty)}.$$
(49)

Since  $\frac{c(2)}{c(\infty)} = 2$ , this completes the first proof. Moreover, this result is sharp with equality for  $\mathbb{1}_{(0,\infty)^d} e^{-\sum_i x_i}$ , which can be verified by checking the d=1 case and then tensorizing. When  $\alpha < 2$ ,

$$h_{\alpha}(X - Y) \le h_{\infty}(X - Y) + d\log\frac{c(\alpha)}{c(\infty)}$$
(50)

$$= h_2(X) + d\log \alpha^{\frac{1}{\alpha - 1}} \tag{51}$$

$$\leq h_{\alpha}(X) + d\log \alpha^{\frac{1}{\alpha - 1}}. (52)$$

Remark 6.3. Identity (44) was also crucial in [17] in Li's proof of the conjectural entropic Buseman theorem (see [1, 23]) in the case of Rényi entropy of order 2. In fact, in the symmetric case (39) for  $\alpha = 2$  follows from Li's result [17]. When d = 1 and X is symmetric, Conjecture 6.1 is a special case of [1, Conjecture 1]. Alternatively, under the symmetry assumption X - Y has the same distribution as X + Y, and the result follows from the additive case [9, 38, 23] for  $\alpha \leq 1$  even when X and Y are dependent. Under stronger concavity assumptions, there are analogous results for  $\alpha \geq 1$ , see [37, 21, 20].

From Theorem 1.3, repeating the above proof mutatis mutandis, we can establish discrete analogs of the Rényi entropic Rogers-Shephard.

**Theorem 6.2.** For X and Y iid log-concave variables on  $\mathbb{Z}$ ,

$$H_{\alpha}(X - Y) - H_{\alpha}(X) < \log c(\alpha), \tag{53}$$

with

$$c(\alpha) = \begin{cases} 2\alpha^{\frac{1}{\alpha-1}}, & \text{if } \alpha \in (2, \infty], \\ \alpha^{\frac{1}{\alpha-1}}, & \text{if } \alpha \in (0, 2]. \end{cases}$$
 (54)

The limiting cases 1 and  $\infty$  are understood by the same conventions as Corollary 6.2,  $1^{\frac{1}{1-1}} := e$  and  $\infty^{\frac{1}{\infty-1}} := 1$ . In the case of the Shannon entropy this gives

$$H(X - Y) - H(X) < \log e. \tag{55}$$

When  $\alpha = 0$ ,  $H_0(X - Y) \le H_0(X) + \log 2$  holds, and is strict whenever the support of X is finite. The inequality is sharp as can be seen by taking a log-concave distribution supported on  $\{0, 1, \ldots, n\}$  for n large.

*Proof.* The only modification of the proof of Theorem 6.1 is in (48) and (49), where we use  $H_2(X) < H_{\infty}(X) + \log \alpha^{\frac{1}{1-\alpha}}$  and  $H_{\infty}(X) \le H_2(X)$ .

Observe that when  $X_p$  and  $Y_p$  are iid geometric with parameter p, and  $\alpha > 0$ , then

$$\lim_{p \to 0} H_{\alpha}(X_p - Y_p) - H_{\alpha}(X_p) = \lim_{p \to 0} \frac{1}{1 - \alpha} \log \frac{1 + (1 - p)^{\alpha}}{(2 - p)^{\alpha}} = \log 2.$$
 (56)

Thus  $c(\alpha)$  cannot be improved beyond 2 for any  $\alpha$ , and hence Theorem 6.2 is sharp for  $\alpha \in \{2, \infty\}$ .

### References

- [1] K. Ball, P. Nayar, and T. Tkocz. A reverse entropy power inequality for log-concave random vectors. *Studia Math*, 235(1):17–30, 2016.
- [2] K. Ball and V. H. Nguyen. Entropy jumps for isotropic log-concave random vectors and spectral gap. *Studia Math.*, 213(1):81–96, 2012.
- [3] S. Bobkov and M. Madiman. The entropy per coordinate of a random vector is highly constrained under convexity conditions. *IEEE Trans. Inform. Theory*, 57(8):4940–4954, August 2011.
- [4] S. G. Bobkov and G. P. Chistyakov. Bounds for the maximum of the density of the sum of independent random variables. *Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI)*, 408(Veroyatnost i Statistika. 18):62–73, 324, 2012.
- [5] S. G. Bobkov and G. P. Chistyakov. Entropy power inequality for the Rényi entropy. *IEEE Trans. Inform. Theory*, 61(2):708–714, February 2015.
- [6] S. G. Bobkov and M. Madiman. On the problem of reversibility of the entropy power inequality. In *Limit theorems in probability, statistics and number theory*, volume 42 of *Springer Proc. Math. Stat.*, pages 61–74. Springer, Heidelberg, 2013. Available online at arXiv:1111.6807.
- [7] S. G. Bobkov and A. Marsiglietti. Variants of the entropy power inequality. *IEEE Transactions on Information Theory*, 63(12):7747–7752, 2017.
- [8] J. H. E. Cohn. Some integral inequalities. The Quarterly Journal of Mathematics, 20(1):347–349, 1969.
- [9] T. M. Cover and Z. Zhang. On the maximum entropy of the sum of two dependent random variables. *IEEE Trans. Inform. Theory*, 40(4):1244–1246, 1994.

- [10] B. A. Slomka D. Halikias, B. Klartag. Discrete variants of Brunn-Minkowski type inequalities. arXiv preprint arXiv:1911.04392, 2019.
- [11] M. Fradelizi, M. Madiman, and L. Wang. Optimal concentration of information content for log-concave densities. In C. Houdré, D. Mason, P. Reynaud-Bouret, and J. Rosinski, editors, *High Dimensional Probability VII: The Cargèse Volume*, Progress in Probability. Birkhäuser, Basel, 2016. Available online at arXiv:1508.04093.
- [12] A. Havrilla and T. Tkocz. Sharp Khinchin-type inequalities for symmetric discrete uniform random variables. arXiv preprint arXiv:1912.13345, 2019.
- [13] S.-W. Ho and S. Verdú. Convexity/concavity of Rényi entropy and α-mutual information. In 2015 IEEE International Symposium on Information Theory (ISIT), pages 745–749. IEEE, 2015.
- [14] O. Johnson. Log-concavity and the maximum entropy property of the Poisson distribution. *Stochastic Process. Appl.*, 117(6):791–802, 2007.
- [15] O. Johnson, I. Kontoyiannis, and M. Madiman. Log-concavity, ultra-log-concavity, and a maximum entropy property of discrete compound Poisson measures. *Discrete Appl. Math.*, 161(9):1232–1250, 2013.
- [16] B. Klartag and J. Lehec. Poisson processes and a log-concave Bernstein theorem. Studia Math., 247(1):85–107, 2019.
- [17] J. Li. Rényi entropy power inequality and a reverse. Studia Math., 242(3):303-319, 2018.
- [18] J. Li, M. Fradelizi, and M. Madiman. Concentration of information content for convex measures. arXiv preprint arXiv:1512.01490, 2019.
- [19] J. Li, A. Marsiglietti, and J. Melbourne. Rényi entropy power inequalities for s-concave densities. In 2019 IEEE International Symposium on Information Theory (ISIT), pages 2224–2228. IEEE, 2019.
- [20] J. Li, A. Marsiglietti, and J. Melbourne. Further investigations of Rényi entropy power inequalities and an entropic characterization of s-concave densities. Geometric Aspects of Functional Analysis Israel Seminar (GAFA) 2017-2019, (Lecture Notes in Mathematics 2262), 2020.
- [21] J. Li and J. Melbourne. Further investigations of the maximum entropy of the sum of two dependent random variables. In 2018 IEEE International Symposium on Information Theory (ISIT), pages 1969–1972. IEEE, 2018.
- [22] M. Madiman and I. Kontoyiannis. Entropy bounds on Abelian groups and the Ruzsa divergence. *IEEE Transactions on Information Theory*, 64(1):77–92, 2016.
- [23] M. Madiman, J. Melbourne, and P. Xu. Forward and reverse entropy power inequalities in convex geometry. Convexity and Concentration, pages 427–485, 2017.
- [24] M. Madiman, J. Melbourne, and P. Xu. Rogozin's convolution inequality for locally compact groups. *Preprint*, arXiv:1705.00642, 2017.
- [25] A. W. Marshall, I. Olkin, and B. C. Arnold. *Inequalities: theory of majorization and its applications*. Springer Series in Statistics. Springer, New York, second edition, 2011.

- [26] A. Marsiglietti and V. Kostina. A lower bound on the differential entropy of log-concave random vectors with applications. *Entropy*, 20(3):185, 2018.
- [27] A. Marsiglietti and J. Melbourne. A Rényi entropy power inequality for log-concave vectors and parameters in [0, 1]. In 2018 IEEE International Symposium on Information Theory (ISIT), pages 1964–1968. IEEE, 2018.
- [28] A. Marsiglietti and J. Melbourne. On the entropy power inequality for the Rényi entropy of order [0, 1]. *IEEE Transactions on Information Theory*, 65(3):1387–1396, 2019.
- [29] A. Marsiglietti and J. Melbourne. Localization technique in the discrete setting with applications. arXiv preprint arXiv:2004.12005, 2020.
- [30] J. Melbourne and T. Tkocz. On the Rényi entropy of log-concave sequences. In *Proc. IEEE Int. Symp. Inform. Theory.* Los Angeles, 2020.
- [31] P.-M. Samson N. Gozlan, C. Roberto and P. Tetali. Transport proofs of some discrete variants of the Prékopa-Leindler inequality. arXiv preprint arXiv:1905.04038, 2019.
- [32] E. Ram and I. Sason. On Rényi entropy power inequalities. *IEEE Transactions on Information Theory*, 62(12):6800–6815, 2016.
- [33] A. Rényi. On measures of entropy and information. In Proc. 4th Berkeley Sympos. Math. Statist. and Prob., Vol. I, pages 547–561. Univ. California Press, Berkeley, Calif., 1961.
- [34] O. Rioul. Rényi entropy power inequalities via normal transport and rotation. Entropy, 20(9):641, 2018.
- [35] C. A. Rogers and G. C. Shephard. The difference body of a convex body. *Arch. Math.* (Basel), 8:220–233, 1957.
- [36] R. P. Stanley. Log-concave and unimodal sequences in algebra, combinatorics, and geometry. In *Graph theory and its applications: East and West (Jinan, 1986)*, volume 576 of *Ann. New York Acad. Sci.*, pages 500–535. New York Acad. Sci., New York, 1989.
- [37] P. Xu, J. Melbourne, and M. Madiman. Reverse entropy power inequalities for s-concave densities. In *Proceedings 2016 IEEE International Symposium on Information Theory*, pages 2284–2288, Barcelona, Spain, 2016.
- [38] Y. Yu. Letter to the editor: On an inequality of Karlin and Rinott concerning weighted sums of i.i.d. random variables. Adv. in Appl. Probab., 40(4):1223–1226, 2008.