NOTE ON THE p-DIVISIBILITY OF CLASS NUMBERS OF AN INFINITE FAMILY OF IMAGINARY QUADRATIC FIELDS.

SRILAKSHMI KRISHNAMOORTHY AND SUNIL KUMAR PASUPULATI

ABSTRACT. For any odd prime p, we construct an infinite family of imaginary quadratic fields whose class numbers are divisible by p. We give a corollary which settles Iizuka's conjecture for the case n=1 and p>2.

1. Introduction

Let K be a number field. The ideal class group Cl_K is defined to be the quotient group J_K/P_K , where J_K is the group of fractional ideals of K and P_K is the group of principal fractional ideals of K. It is well known that Cl_K is finite. The class number h_K of a number field K is the order of Cl_K . The ideal class group is one of the most basic and mysterious objects in algebraic number theory. The class group has drawn the attention of several authors. The divisibility properties of the class number of number fields play a significant role in understanding the structure of the ideal class groups of number fields. Cohen-Lenstra heuristics are a set of conjectures about this structure. In general, given a positive integer n, there is no characterization to find all quadratic fields whose class numbers are divisible by n. For n=3, these fields were characterized in [12]. For a given integer n>1, the Cohen-Lenstra heuristic [4] predicts that the proportion of imaginary quadratic fields with class number divisible by n should be positive. It has been proved by several authors that for every n>1, there exist infinitely many quadratic fields whose class number is divisible by n ([11], [10], [5], [2], [7], [14]).

B. H. Gross and D. E. Rohrlich proved that for any odd integer n > 3, there are infinitely many imaginary quadratic fields $(\mathbb{Q}\left(\sqrt{1-4U^n}\right), U > 1)$ whose class numbers are divisible by n. Furthermore Stéphane Louboutin [14] proved the same result by simplifying the Gross and Rohrlich's proof and proved the following result on divisibility of class number of $\mathbb{Q}(\sqrt{1-4U^k})$ for U > 2.

Theorem 1. If $k \in \mathbb{Z}^+$ be odd number, then for any integer $U \geq 2$ the ideal class groups of the imaginary quadratic fields $\mathbb{Q}(\sqrt{1-4U^k})$ contain an element of order k.

Murty [15] proved that the class number of $\mathbb{Q}\left(\sqrt{1-U^n}\right)$ is divisible by n if $1-U^n$ is square-free. We study the divisibility of class number of families $\mathbb{Q}(\sqrt{1-2m^p})$ by all odd primes p.

The following result on the 3-divisibility of the class number is proved by K. Chakraborty and A. Hoque (Theorem 3.2, [8]).

²⁰¹⁰ Mathematics Subject Classification. Primary: 11R29, Secondary: 11R11.

Key words and phrases. Class number, ideal class group, imaginary quadratic fields, Diophantine equation.

K. SRILAKSHMI AND P. SUNIL KUMAR

Theorem 2. The class number of $\mathbb{Q}(\sqrt{1-2m^3})$ is divisible by 3 for any odd integer m > 1.

We study the family $\mathbb{Q}(\sqrt{1-2m^p})$ for all odd primes p and prime power $m=q^r, r \in \mathbb{N}$. In the following theorem, we prove the p-divisibility of class numbers for this family by using the results of Yann Bugeaud and T. N. Shorey [1].

Theorem 3. For prime numbers $p, q \geq 3$ and $m = q^r, r \in \mathbb{N}$ such that $\mathbb{Q}(\sqrt{1-2m^p}) \neq \mathbb{Q}(\sqrt{-1})$, the class number of $\mathbb{Q}(\sqrt{1-2m^p})$ is divisible by p.

The condition $\mathbb{Q}(\sqrt{1-2m^p}) \neq \mathbb{Q}(\sqrt{-1})$, means that $2m^p-1$ is not a square. Siegel's theorem (Lemma 9) asserts that there are only finitely many $m \in \mathbb{Z}$ such that $2m^p-1$ is a square.

The Birch Swinnerton-Dyer conjecture is an elliptic curve analogue of the analytic class number formula. For any elliptic curve defined over \mathbb{Q} of rank zero and square-free conductor N, if $p \mid |E(\mathbb{Q})|$, under certain conditions on the Shafarevich-Tate group III_D , the first author [13] showed that $p \mid |\mathrm{III}_D|$ if and only if $p \mid h_K$, where $K = \mathbb{Q}(\sqrt{-D})$.

2. IIZUKA'S CONJECTURE

Y.Iizuka recently proves the following result on divisibility of the class numbers of imaginary quadratic fields in [9].

Theorem 4. There is an infinite family of pairs of imaginary quadratic fields $\mathbb{Q}(\sqrt{d})$ and $\mathbb{Q}(\sqrt{d+1})$ with $d \in \mathbb{Z}$ whose class numbers are both divisible by 3.

Based on the above theorem, Iizuka conjectured the following

Conjecture 5. (Iizuka) For any prime p and any positive integer n, there is an infinite family of n+1 successive real (or imaginary) quadratic fields

$$\mathbb{Q}(\sqrt{D}), \mathbb{Q}(\sqrt{D+1}), \cdots, \mathbb{Q}(\sqrt{D+n})$$

with $D \in \mathbb{Z}$ whose class numbers are divisible by p.

As a consequence of Theorem 1 and Theorem 3, we get a generalization of Theorem 4 for all odd prime numbers p and prove the following corollary.

Corollary 6. For every odd prime number there is an infinite family of pairs of imaginary quadratic fields $\mathbb{Q}(\sqrt{d})$ and $\mathbb{Q}(\sqrt{d+1})$ with $d \in \mathbb{Z}$ whose class numbers are both divisible by p.

Proof of the Corollary 6.

Fix an odd prime p. Consider the set

$$S_0 = \left\{ m \in \mathbb{Z}^+ | \text{ the class number of } \mathbb{Q}\left(\sqrt{1 - 2m^p}\right) \text{ is divisible by } p \right\}.$$

On the p-divisibility of class numbers of $\mathbb{Q}(\sqrt{d})$ and $\mathbb{Q}(\sqrt{d+1})$.

By Lemma 9, the equation $1 - 2x^p = -y^2$ has finitely many solutions $(x, y) \in \mathbb{Z} \times \mathbb{Z}$. Hence it follows from Theorem 3 that S_0 contains infinitely many odd prime powers, which implies that S_0 is an infinite set. For $m \in S_0$, the prime p divides the class number of

$$\mathbb{Q}(\sqrt{4(1-2m^p)^p}) = \mathbb{Q}(\sqrt{1-2m^p}).$$

Let $U = 2m^p - 1$. Then $U \ge 2$. Furthermore Theorem 1 implies that p divides the class number of $\mathbb{Q}(\sqrt{1-4U^p})$. Now look at

$$\mathbb{Q}(\sqrt{1-4U^p}) = \mathbb{Q}\left(\sqrt{1-4(2q^p-1)^p}\right) = \mathbb{Q}\left(\sqrt{4(1-2q^p)^p+1}\right).$$

Let $d = 4(1 - 2m^p)^p$. The prime p divides class numbers of $\mathbb{Q}(\sqrt{d}), \mathbb{Q}(\sqrt{d+1})$.

Now to conclude the corollary, we need to prove the set $\mathcal{A} = \left\{ \mathbb{Q}\left(\sqrt{1-2m^p}\right) | m \in S_0 \right\}$ is an infinite set. For every square-free integer $d_0 \neq 0$, let $f(x) = \frac{1-2x^p}{d_0}$. The polynomial f(x) has distinct roots in $\overline{\mathbb{Q}}$. Thus by Lemma 9, the equation $y^2 = f(x)$ has finitely many integral solutions. Hence the infiniteness of \mathcal{A} follows from that of S_0 .

Remark 7. The above corollary settles Iizuka's conjecture (5) for the case n = 1 and p > 2. We found a similar result for a different families of imaginary quadratic fields in [17]. J. Chattopadhyay and S. Muthukrishnan [3] answer a weaker version of Iizuka's conjecture for p = 3.

3. Preliminaries

We recall some known results and prove some lemmas that are necessary for proving our main theorem.

Definition 8. Let K be a number field and let S be a finite set of valuations on K, containing all the archimedean valuations. Then

$$R_S = \{ \alpha \in K \mid \nu(\alpha) > 0 \text{ for all } \nu \notin S \}$$

is called the set of S-integers.

Lemma 9. (Siegel's theorem, [16], Chapter IX, Theorem 4.3) Let K be a number field and S be a finite set of valuations on K, containing all the archimedean valuations. Let $f(X) \in K[X]$ be a polynomial of degree $d \geq 3$ with distinct roots in the algebraic closure \overline{K} of K. Then the equation $y^2 = f(x)$ has only finitely many solutions in S-integers $x, y \in R_S$.

We recall some results of Yann Bugeaud and T.N. Shorey [1] on solutions of Diophantine equation $D_1x^2 + D_2 = \lambda^2 k^y$, where D_1 and D_2 are coprime positive integers, $k \geq 2$ is an integer coprime with D_1D_2 and $\lambda = \sqrt{2}$, 2 such that $\lambda = 2$ if k is even.

Let us denote F_i to be the Fibonacci sequence defined by $F_0 = 0$, $F_1 = 1$ and $F_i = F_{i-1} + F_{i-2}$ for all $i \geq 2$. Let L_i be the Lucas sequence defined by $L_0 = 2$, $L_1 = 1$ and satisfying $L_i = L_{i-1} + L_{i-2}$ for all $i \geq 2$. Define the subsets $\mathcal{F}, \mathcal{G}, \mathcal{H}$ of $\mathbb{N} \times \mathbb{N} \times \mathbb{N}$ by

$$\mathcal{F} := \{ (F_{i-2\epsilon}, L_{i+\epsilon}, F_i) \mid i \ge 2, \epsilon \in \{\pm 1\} \},$$

$$\mathcal{G} := \{ (1, 4k^r - 1, k) \mid k \ge 2, r \ge 1 \},$$

 $\mathcal{H} := \{(D_1, D_2, k) \mid \text{there exist positive integers } r \text{ and } s \text{ such that }$

$$D_1 s^2 + D_2 = \lambda^2 k^r$$
 and $3D_1 s^2 - D_2 = \pm \lambda^2$.

Define $\mathcal{N}(\lambda, D_1, D_2, p)$ to be the number of $(x, y) \in \mathbb{Z}^+ \times \mathbb{Z}^+$ of the Diophantine equation $D_1 x^2 + D_2 = \lambda^2 p^y$.

Theorem 10. ([1], Theorem 1) Let p be a prime number. Then we have $\mathcal{N}(\lambda, D_1, D_2, p) \leq 1$ expect for $\mathcal{N}(2, 13, 3, 2) = \mathcal{N}(\sqrt{2}, 7, 11, 3) = \mathcal{N}(1, 2, 1, 3) = \mathcal{N}(2, 7, 1, 2) = \mathcal{N}(\sqrt{2}, 1, 1, 5) = \mathcal{N}(\sqrt{2}, 1, 1, 13) = \mathcal{N}(2, 1, 3, 7) = 2$ and when (D_1, D_2, p) belongs to one of the infinite families \mathcal{F}, \mathcal{G} and \mathcal{H} .

Lemma 11. For any odd prime q and any integer D > 3, the equation $Dx^2 + 1 = 2q^y$ has at most one solution $(x, y) \in \mathbb{Z}^+ \times \mathbb{Z}^+$.

Proof. Let $D_1 = D$ and $D_2 = 1$ and $\lambda = \sqrt{2}$. We first note that $(\lambda, D_1, D_2, q) \notin \{(2, 13, 3, 2), (\sqrt{2}, 7, 11, 3), (1, 2, 1, 3), (2, 7, 1, 2), (\sqrt{2}, 1, 1, 5), (\sqrt{2}, 1, 1, 13), (2, 1, 3, 7)\}$. By Theorem 10, it is enough to show that $(D_1, D_2, q) \notin \mathcal{F}$. If $(F_{i-\epsilon}, L_{i+\epsilon}, F_i) = (D_1, D_2, p)$, then $i = 2, \epsilon = -1$. Hence D = 2. This is not possible because D > 3. Therefore $(D_1, D_2, q) \notin \mathcal{F}$. If $(1, 4k^r - 1, k) = (D_1, D_2, q)$, then $D = D_1 = 1$. This is not possible. Therefore $(D_1, D_2, q) \notin \mathcal{G}$. If $D_1x^2 + 1 = 2q^y$, then $3D_1x^2 - 1 \ge D_1x^2 - 1 \ge 2q^y - 2 \ge 4$. Hence $(D_1, D_2, q) \notin \mathcal{H}$.

Proposition 12. For prime numbers $p, q \geq 3$ and $m = q^r, r \in \mathbb{N}$, let $\alpha = 1 + \sqrt{1 - 2m^p}$, then $\pm 2^{\frac{p-1}{2}}\alpha$ is not a p-th power of an algebraic integer in $\mathbb{Q}(\sqrt{1 - 2m^p})$.

Proof. Let d be the square-free part of $\sqrt{1-2m^p}$ with signature. Let $K = \mathbb{Q}(\sqrt{d})$ and \mathcal{O}_K be the ring of integers of K. Note that $-2^{\frac{p-1}{2}}\alpha$ is a p-th power in \mathcal{O}_K if and only if $2^{\frac{p-1}{2}}\alpha$ is a p-th power in \mathcal{O}_K . It is enough to show $2^{\frac{p-1}{2}}\alpha$ is not a p-th power. Suppose that,

(3.1)
$$2^{\frac{p-1}{2}}\alpha = \beta^p \text{ for some } \beta = a + b\sqrt{d} \in \mathcal{O}_K.$$

Then,

(3.2)
$$2^{\frac{p-1}{2}} \left(1 + \sqrt{1 - 2m^p} \right) = \sum_{j=0}^{\frac{p-1}{2}} \binom{p}{2j} a^{p-2j} b^{2j} d^j + \gamma \sqrt{d} \text{ for some } \gamma \in \mathbb{Z}.$$

By comparing constant terms on both sides, we have,

(3.3)
$$2^{\frac{p-1}{2}} = \sum_{j=0}^{\frac{p-1}{2}} \binom{p}{2j} a^{p-2j} b^{2j} d^j.$$

This implies that,

$$2^{\frac{p-1}{2}} = a \left(\sum_{j=0}^{\frac{p-1}{2}} \binom{p}{2j} a^{p-2j-1} b^{2j} d^j \right).$$

Hence a divides $2^{\frac{p-1}{2}}$.

On the p-divisibility of class numbers of $\mathbb{Q}(\sqrt{d})$ and $\mathbb{Q}(\sqrt{d+1})$.

Case 1:a is even.

We look at (3.1)

$$(3.4) 2^{\frac{p-1}{2}}\alpha = \left(a + b\sqrt{d}\right)^p,$$

applying the norm map on the both sides

$$(2m)^p = \left(a^2 - b^2 d\right)^p.$$

Hence we have

$$2m = a^2 - b^2 d$$

Since $2 \mid a$, we obtain $2 \mid b^2 d$. We deduce that $2 \mid b$ since d is odd. Taking divisibility of a^2, b^2 by 4 into consideration, we conclude that $4 \mid 2m$ but m is odd which contradicts the assumption that a is even. Case 2 : a is odd.

Suppose that a is odd. Since $a \mid 2^{\frac{p-1}{2}}$, this implies that $a = \pm 1$. Putting in Equation (3.1) we have

$$2^{\frac{p-1}{2}}(1+\sqrt{1-2m^p)} = \left(\pm 1 + b\sqrt{d}\right)^p,$$

applying the norm map on both sides we get

$$(2m)^p = (1 - b^2 d)^p$$
.

Rewriting the above equation using D = -d and $m = q^r$, we get

$$(3.5) 1 + Db^2 = 2q^r.$$

We observe that $1 - 2m^p = 1 - 2(q^r)^p = b'^2 d$ for some $b' \in \mathbb{Z}$. Rephrasing this equation we have,

$$(3.6) 1 + b'^2 D = 2q^{rp}.$$

Thus, from Equations (3.5) and (3.6), we get (b, r), (b', rp) are solutions of the equation $Dx^2 + 1 = q^y$, which is a contradiction to Lemma 11. Hence $\pm 2^{\frac{p-1}{2}}\alpha$ is not a p-th power in \mathcal{O}_K .

4. Proof of the theorem

We now prove the main theorem of this article.

Proof of Theorem 3. Let d be the square-free part of $1-2m^p$ with signature then $d\equiv 3\pmod 4$ and $K=\mathbb{Q}(\sqrt{d})$. Put $\alpha:=1+\sqrt{1-2m^p}$, then $N_{K/\mathbb{Q}}(\alpha)=2m^p$. Since $d\equiv 3\pmod 4$, the ideal (2) is ramified, there exists a prime ideal \mathcal{P} such that $(2)=\mathcal{P}^2$. Since the norm of α is $2m^p=2q^{rp}$, the prime decomposition of (α) is given by $(\alpha)=\mathcal{P}\mathcal{Q}^t$, for some positive integer t, where \mathcal{Q} is a prime which lies above q. Then $N_{K/\mathbb{Q}}(\alpha)=2q^t$, where $N(\mathcal{Q})=q$ (since q splits in $\mathbb{Q}(\sqrt{d})$ as $\left(\frac{d}{q}\right)=1$). Hence t=rp.

Consider the ideal $I := \mathcal{PQ}^{\frac{t}{p}}$, of K. Observe that

$$I^p = \mathcal{P}^p \mathcal{Q}^t = (2)^{\frac{p-1}{2}} \mathcal{P} \mathcal{Q}^t = (2)^{\frac{p-1}{2}} (\alpha) = (2^{\frac{p-1}{2}} \alpha).$$

We claim that the order of the ideal I in ideal class group is p. Suppose not, let $(\beta) = I$ for some β in \mathcal{O}_K . Then

$$(\beta^p) = (\beta)^p = I^p = (2^{\frac{p-1}{2}}\alpha).$$

K. SRILAKSHMI AND P. SUNIL KUMAR

Since the only units of \mathcal{O}_K are $\{1, -1\}$, this implies that I is a principal ideal if and only if $\pm 2^{\frac{p-1}{2}}\alpha$ is a power of p in \mathcal{O}_K . From Proposition 12, we know that $\pm 2^{\frac{p-1}{2}}\alpha$ is not a p-th power in \mathcal{O}_K . Hence the class group of $\mathbb{Q}(\sqrt{1-2m^p})$ has an element I of order p.

We prove a corollary of Theorem 3.

Corollary 13. For every odd prime p, there exist infinitely many imaginary biquadratic fields whose class number is divisible by p.

Proof. Fix an odd prime p. Consider the set

$$S_1 = \{ m \in \mathbb{Z}^+ | m \text{ is not a square, } m \equiv 1 \pmod{4} \text{ and}$$

the class number of $\mathbb{Q}(\sqrt{1-2m^p})$ is divisible by $p \}$.

By Lemma 9, the equation $1 - 2x^p = y^2$ has only finitely many solutions $(x, y) \in \mathbb{Z} \times \mathbb{Z}$. Hence it follows from Theorem 3 and Dirichlet's theorem on arithmetic progression that S_1 contains infinitely many primes q with $q \equiv 1 \pmod{4}$, which implies that S_1 is an infinite set.

For $m \in S_1$, consider the imaginary biquadratic field $K_m = \mathbb{Q}(\sqrt{1-2m^p},\sqrt{m})$. Denote $L_m^1 := \mathbb{Q}(\sqrt{1-2m^p}), L_m^2 := \mathbb{Q}(\sqrt{m})$ and $L_m^3 := \mathbb{Q}\left(\sqrt{1-2m^p}\sqrt{m}\right)$. Since m is not a square, L_m^2 is actually a quadratic field. We observe that $L_m^1 \neq L_m^2$ because $1-2m^p \equiv 3 \pmod 4$. Thus L_m^1, L_m^2 and L_m^3 are the three quadratic subfields of K_m . Let h_m, h_m^1, h_m^2 and h_m^3 be the class numbers of K_m, L_m^1, L_m^2 and L_m^3 respectively. Then by Lemma 2 in [6], we have $h_m = \frac{h_m^1 h_m^2 h_m^3}{2^i}$ where i = 0, 1. Since $m \in S_1$, the prime p divides h_m^1 . Since p is odd, p divides h_m . The infiniteness of the set $\{K_m | m \in S_1\}$ follows from that of the set S_1 .

ACKNOWLEDGEMENTS

We thank Prof. Kalyan Chakraborty for introducing this problem to us. We thank IISER Tiruvananthapuram for providing excellent working conditions. We thank Azizul Hoque for his comments and the careful reading of this article. We acknowledge SageMath for the numerical evidence of our theorems. We like to express our deep gratitude to the anonymous referee for carefully analyzing the article, refining several arguments and making suggestions for a better presentation.

References

- 1. Yann Bugeaud and T. N. Shorey, On the number of solutions of the generalized Ramanujan-Nagell equation, J. Reine Angew. Math. **539** (2001), 55–74. MR 1863854
- K. Chakraborty, A. Hoque, Y. Kishi, and P. P. Pandey, Divisibility of the class numbers of imaginary quadratic fields, J. Number Theory 185 (2018), 339–348. MR 3734353
- 3. Jaitra Chattopadhyay and Subramani Muthukrishnan, On the simultaneous 3-divisibility of class numbers of triples of imaginary quadratic fields, Acta Arith. 197 (2021), no. 1, 105–110. MR 4185918
- H. Cohen and H. W. Lenstra, Jr., Heuristics on class groups of number fields, Number theory, Noordwijkerhout 1983 (Noordwijkerhout, 1983), Lecture Notes in Math., vol. 1068, Springer, Berlin, 1984, pp. 33–62. MR 756082

- 5. J. H. E. Cohn, On the class number of certain imaginary quadratic fields, Proc. Amer. Math. Soc. 130 (2002), no. 5, 1275–1277. MR 1879947
- Gary Cornell, A note on the class group of composita, J. Number Theory 39 (1991), no. 1, 1–4.
 MR 1123163
- 7. Benedict H. Gross and David E. Rohrlich, Some results on the Mordell-Weil group of the Jacobian of the Fermat curve, Invent. Math. 44 (1978), no. 3, 201–224. MR 491708
- 8. Azizul Hoque and Kalyan Chakraborty, Divisibility of class numbers of certain families of quadratic fields, J. Ramanujan Math. Soc. **34** (2019), no. 3, 281–289. MR 4010381
- 9. Yoshichika Iizuka, On the class number divisibility of pairs of imaginary quadratic fields, J. Number Theory **184** (2018), 122–127. MR 3724158
- 10. Akiko Ito, A note on the divisibility of class numbers of imaginary quadratic fields $\mathbf{Q}(\sqrt{a^2-k^n})$, Proc. Japan Acad. Ser. A Math. Sci. 87 (2011), no. 9, 151–155. MR 2863357
- 11. Yasuhiro Kishi, Note on the divisibility of the class number of certain imaginary quadratic fields, Glasg. Math. J. **51** (2009), no. 1, 187–191. MR 2471686
- 12. Yasuhiro Kishi and Katsuya Miyake, Parametrization of the quadratic fields whose class numbers are divisible by three, J. Number Theory 80 (2000), no. 2, 209–217. MR 1740511
- 13. Srilakshmi Krishnamoorthy, A note on the Fourier coefficients of a Cohen-Eisenstein series, Int. J. Number Theory 12 (2016), no. 5, 1149–1161. MR 3529567
- 14. Stéphane R. Louboutin, On the divisibility of the class number of imaginary quadratic number fields, Proc. Amer. Math. Soc. 137 (2009), no. 12, 4025–4028. MR 2538563
- M. Ram Murty, The abc conjecture and exponents of class groups of quadratic fields, Number theory (Tiruchirapalli, 1996), Contemp. Math., vol. 210, Amer. Math. Soc., Providence, RI, 1998, pp. 85–95. MR 1478486
- Joseph H. Silverman, The arithmetic of elliptic curves, second ed., Graduate Texts in Mathematics, vol. 106, Springer, Dordrecht, 2009. MR 2514094
- 17. Jian-Feng Xie and Kuok Fai Chao, On the divisibility of class numbers of imaginary quadratic fields $\mathbb{Q}(\sqrt{D}), \mathbb{Q}(\sqrt{D+m})$, Ramanujan J. **53** (2020).

SRILAKSHMI KRISHNAMOORTHY

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH, THIRUVANANTHAPURAM, INDIA.

Email address: srilakshmi@iisertvm.ac.in

Sunil Kumar Pasupulati

INDIAN INSTITUTE OF SCIENCE EDUCATION AND RESEARCH, THIRUVANANTHAPURAM, INDIA.

Email address: sunil4960016@iisertvm.ac.in