

GENERATING FUNCTIONS OF PERMUTATIONS WITH RESPECT TO THEIR ALTERNATING RUNS

MIKLÓS BÓNA

ABSTRACT. We present a short, direct proof of the fact that the generating function of all permutations of a fixed length $n \geq 4$ is divisible by $(1+z)^m$, where $m = \lfloor (n-2)/2 \rfloor$.

1. INTRODUCTION

Let $p = p_1 p_2 \cdots p_n$ be a permutation. For an index $i \in [2, n-1]$, we say that p changes direction at i if $p_{i-1} < p_i > p_{i+1}$ or $p_{i-1} > p_i < p_{i+1}$. Furthermore, we say that p has k alternating runs if p changes directions a total of $k-1$ times.

Let $\text{run}(p)$ be the number of alternating runs of p , and let

$$R_n(z) = \sum_p z^{\text{run}(p)},$$

where the sum is taken over all permutations of length n . In this note, we prove that for $n \geq 4$, the polynomial $R_n(z)$ is divisible by a high power of $(1+z)$, namely by $(1+z)^m$, where $m = \lfloor (n-2)/2 \rfloor$.

This result was known before, by an analytic proof given by Herbert Wilf, that can be read also in [2], that was based on the relation between the Eulerian polynomials and $R_n(z)$, and also by an induction proof by Richard Ehrenborg and the present author [1] that also touched upon Eulerian polynomials. However, in this paper, we provide a direct, non-inductive proof.

2. A GROUP ACTION ON PERMUTATIONS

Let $p = p_1 p_2 \cdots p_n$ be a permutation. The *complement* of p is the permutation $\bar{p} = n+1-p_1 \ n+1-p_2 \ \cdots \ n+1-p_n$. For instance, the complement of 425613 is 352164. It is clear that p and \bar{p} have the same number of alternating runs, since the diagram of \bar{p} is just the diagram of p reflected through a horizontal line. In what follows, we will say *flipped* instead of *reflected through a horizontal line*. Note that this symmetry implies that all coefficients of $R_n(z)$ are even for $n \geq 2$.

Date: May 27, 2020.

Let s be a string of entries in p that are in consecutive positions. Let S be set of entries that occur in s , in other words, the *underlying set* of s . Then the *complement of s relative to S* is the string obtained from s so that for each j , the j th smallest entry of S is replaced by the j th largest entry of S . For example, if $s=24783$, then the complement of s relative to its underlying set S is 84327.

We will use a similar notion for sets. Let $T \subseteq U$ be finite sets. Let $T = \{t_1, t_2, \dots, t_j\}$, where the t_i are listed in increasing order. Let us say that t_i is the a_i th smallest element of U . Then the *vertical complement* of T with respect to U is the set consisting of the a_i th largest elements of U , for all i . For instance, if $T = \{1, 4, 6\}$, and $U = \{1, 2, 3, 4, 6, 8, 9\}$, then the vertical complement of T with respect to U is $\{3, 4, 9\}$. Indeed, T consists of the first, fourth, and fifth smallest elements of U , so its vertical complement with respect to U consists of the first, fourth, and fifth largest elements of U .

Definition 2.1. For $1 \leq i \leq n$, let c_i be the transformation on the set of all permutations $p = p_1 p_2 \cdots p_n$ that leaves the string $p_1 p_2 \cdots p_{i-1}$ unchanged, and replaces the string $p_i p_{i+1} \cdots p_n$ by its complement relative to its underlying set.

Note that $c_1(p) = \bar{p}$ and $c_n(p) = p$ for all p .

Example 2.2. Let $p = 315462$. Then $c_3(p) = 314526$, while $c_5(p) = 315426$.

Proposition 2.3. Let $n \geq 4$, and let $3 \leq i \leq n - 1$. Let p be any permutation of length n . Then one of p and $c_i(p)$ has exactly one more alternating run than the other.

Proof. As c_i does not change the number of runs of the string $p_1 p_2 \cdots p_{i-1}$ or the number of runs of the string $p_i p_{i+1} \cdots p_n$ and its image, all changes occur within the four-element string $p_{i-2} p_{i-1} p_i p_{i+1}$. There are only 24 possibilities for the pattern of these four entries, and it is routine to verify the statement for each of the possible 12 pairs. In fact, checking the six pairs in which $p_{i-2} < p_{i-1}$ is sufficient, for symmetry reasons. \square

In other words, $z^{\text{run}(p)} + z^{\text{run}(c_i(p))}$ is divisible by $1 + z$.

The following lemma is crucial for our purposes.

Lemma 2.4. Let $1 \leq i \leq j - 2 \leq n - 2$. Then for all permutations p of length n , the identity $c_i(c_j(p)) = c_j(c_i(p))$ holds.

Proof. Neither c_i nor c_j acts on any part of the initial segment $p_1 p_2 \cdots p_{i-1}$, so that segment, unchanged, will start both $c_i(c_j(p))$ and $c_j(c_i(p))$. The

ending segment $p_j p_{j+1} \cdots p_n$ gets flipped twice by both $c_i c_j$ and $c_j c_i$, so in the end, the pattern of the last $n - j + 1$ entries will be the same in $c_i(c_j(p))$ and $c_j(c_i(p))$, because in both permutations, it will be the same pattern as it was in p . The middle segment $p_i p_{i+1} \cdots p_j$ will get flipped once by both $c_i c_j$ and $c_j c_i$, so on both sides, the pattern of entries in positions $i, i + 1, \dots, j$ will be the complement of the pattern of $p_i p_{i+1} \cdots p_j$.

Finally, the *set* of entries in the last $n - j + 1$ positions is the same in both $c_i(c_j(p))$ and $c_j(c_i(p))$, since both sets are equal to the vertical complement of the set $\{p_j, p_{j+1}, \dots, p_n\}$ with respect to the set $\{p_i, p_{i+1}, \dots, p_n\}$ \square

Note that $c_i(c_j(p)) \neq p$, since the segment $p_i \cdots p_{j-1}$ is of length at least two, and gets flipped at least once.

Example 2.5. Continuing Example 2.2, we get that $c_3(c_5(315462)) = c_5(c_3(315462)) = 314562$.

Now let $n \geq 4$ be any integer. If n is even, let

$$\mathcal{C}_n = \{c_3, c_5, c_7, \dots, c_{n-1}\}.$$

If n is odd, let

$$\mathcal{C}_n = \{c_3, c_5, c_7, \dots, c_{n-2}\}.$$

In both cases, \mathcal{C}_n consists of $m = \lfloor (n-2)/2 \rfloor$ operations. Each of these operations are involutions, and by Lemma 2.4, they pairwise commute. Therefore, they define a group $G_m \cong \mathbb{Z}_2^m$ that acts on the set of all permutations of length n .

Note that with this set of generators, $c_i(c_j(p)) \neq p$, since the segment $p_i \cdots p_{j-1}$ is of length at least two, and gets flipped exactly once. Also note that the action of the individual c_i on the set of all permutations of length n is independent in the following sense. If p is a permutation, and $c_i(p)$ has one more alternating run than p , then $c_j(c_i(p))$ also has one more alternating run than $c_j(p)$.

The action of G_m on the set of all permutations of length n creates orbits of size 2^m .

Lemma 2.6. *Let A be any orbit of G_m on the set of all permutations of length n . Then the equality*

$$\sum_{p \in A} z^{\text{run}(p)} = z^a (1 + z)^m$$

holds, where a is a nonnegative integer.

Proof. Let $p \in A$, and going through p left to right, let us apply or not apply each element of \mathcal{C}_n so as to minimize the number of alternating runs of the obtained permutation. That is, if applying c_i increases the number of alternating run, then do not apply it, if it decreases the number of alternating runs, then apply it. Let q be the obtained permutation. Then we call q the *minimal* permutation in A , since among all permutations in A , it is q that has the smallest number of alternating runs. Now elements of A with i more alternating runs than q can be obtained from q by applying exactly i elements of \mathcal{C}_n to q . As there are $\binom{m}{i}$ ways to choose i such elements, our statement is proved by summing over i . \square

Theorem 2.7. *For $n \geq 4$, the equality*

$$R_n(z) = (1+z)^m \sum_q z^{\text{run}(q)}$$

holds, where the summation is over permutations q that are minimal in their orbit under the action of G_m . Here $m = \lfloor (n-2)/2 \rfloor$.

Proof. This follows from Lemma 2.6 by summing over all orbits. \square

REFERENCES

- [1] M. Bóna, R. Ehrenborg, A combinatorial proof of the log-concavity of the numbers of permutations with k runs. *J. Combin. Theory Ser. A* **90** (2000), no. 2, 293–303.
- [2] M. Bóna, *Combinatorics of Permutations*, Second Edition, CRC Press, 2012.
- [3] H. Wif Real Zeroes of polynomials that count runs and descending runs, unpublished manuscript, 1999.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF FLORIDA, 358 LITTLE HALL,
PO BOX 118105, GAINESVILLE, FL, 32611 – 8105 (USA)

E-mail address: bona@ufl.edu