

DEDICATED TO THE MEMORY OF PROFESSOR GUANG-YU SHEN

ON ENHANCED REDUCTIVE GROUPS (I):
PARABOLIC SCHUR ALGEBRAS AND THE DUALITIES RELATED TO
DEGENERATE DOUBLE HECKE ALGEBRAS

BIN SHU, YUNPENG XUE AND YUFENG YAO

ABSTRACT. An enhanced algebraic group \underline{G} of $G = \mathrm{GL}(V)$ over \mathbb{C} is a product variety $\mathrm{GL}(V) \times V$, endowed with an enhanced cross product. Associated with a natural tensor representation of \underline{G} , there are naturally Levi and parabolic Schur algebras \mathcal{L} and \mathcal{P} respectively. We precisely investigate their structures, and study the dualities on the enhanced tensor representations for variant groups and algebras. In this course, an algebraic model of so-called degenerate double Hecke algebras (DDHA) is produced, and becomes a powerful implement. The connection between \mathcal{L} and DDHA gives rise to two results for the classical representations of $\mathrm{GL}(V)$: (i) A duality between $\mathrm{GL}(V) \times \mathbf{G}_m$ and DDHA where \mathbf{G}_m is the one-dimensional multiplicative group; (ii) A branching duality formula. With aid of the above discussion, we further obtain a parabolic Schur-Weyl duality for $\underline{G} \rtimes \mathbf{G}_m$. What is more, the parabolic Schur subalgebra turns out to have only one block. The Cartan invariants for this algebra are precisely determined.

CONTENTS

Introduction	2
1. Semi-reductive groups and semi-reductive Lie algebras	4
2. Enhanced tensor representations and parabolic Schur algebras	5
3. Degenerate double Hecke algebras	14
4. Duality related to the degenerate double Hecke algebra and the branching duality formula	21
5. Enhanced tensor invariants and parabolic Schur-Weyl duality	26
6. Representations of the parabolic Schur algebra	30
Acknowledgement	34
References	34

2010 *Mathematics Subject Classification.* 20G05, 20G20, 20C30, 17B10.

Key words and phrases. enhanced-reductive algebraic groups, tensor representations, symmetric groups, Parabolic /Levi Schur algebras, degenerated double Hecke algebras, (Parabolic /Levi) Schur-Weyl duality, blocks, Cartan invariants.

This work is partially supported by the National Natural Science Foundation of China (12071136, 11671138 and 11771279), Shanghai Key Laboratory of PMMP (No. 13dz2260400).

INTRODUCTION

0.1. An algebraic group G is called a semi-reductive group if G is a semi-direct product of a reductive closed subgroup G_0 and the unipotent radical U . The study of semi-reductive algebraic groups and their Lie algebras becomes very important to lots of cases when the underground field is of characteristic $p > 0$ (see [15]). The examples what we are concerned arise from restricted simple Lie algebras of Cartan type (see [16], [18] and [19]). In the present paper, we have another natural example. Let $G = \mathrm{GL}(V)$ and ν be the natural representation of G on V . Then we have a typical enhanced reductive algebraic group $\underline{G} = G \times_{\nu} V$, which is a closed subgroup of $\mathrm{GL}(\underline{V})$ with \underline{V} being a one-dimensional extension of V . The enhanced reductive group \underline{G} is naturally a semi-reductive group. Some study of semi-reductive algebraic group will be done in some other place (to see [15]).

0.2. By the classical Schur-Weyl duality, the study of polynomial representations of general linear groups produces Schur algebras. Precisely, for a given infinite field \mathbb{F} , and $G = \mathrm{GL}(n, \mathbb{F})$ the general linear group over \mathbb{F} , the Schur algebra $S_{\mathbb{F}}(n, r)$ is exactly $\mathrm{End}_{\mathbb{F}}(E^{\otimes r})^{\mathfrak{S}_r}$ for $E = \mathbb{F}^n$. The clear structure makes Schur algebras become powerful and tractable in the study of polynomial representations of $\mathrm{GL}(n, \mathbb{F})$ (see [11], [14], *etc.*).

By analogy of this, the tensor representations of an enhanced group \underline{G} naturally produce the so-called enhanced Schur algebra $\mathcal{E}_{\mathbb{C}}(n, r)$ which are the algebras generated by the image of \underline{G} in the r th tensor representation on $\underline{V}^{\otimes r}$ for $V = \mathbb{C}^n$, respectively (see §2.5). Related with \underline{G} , there is canonically a short exact sequence of closed subgroups of GL_{n+1} : $\underline{G} \hookrightarrow \underline{G} \times \mathbf{G}_m \twoheadrightarrow \mathbf{G}_m$. And $\underline{G} \times \mathbf{G}_m$ is actually a parabolic subgroup of GL_{n+1} associated with the Levi subgroup $\mathrm{GL}_n \times \mathbf{G}_m$. Correspondingly, the so-called Levi and parabolic Schur algebras $\mathcal{L}(n, r)$ and $\mathcal{P}(n, r)$ naturally arise, which are by definition the image of $\mathrm{GL}_n \times \mathbf{G}_m$ and $\underline{G} \times \mathbf{G}_m$ in $\mathrm{End}_{\mathbb{F}}(\underline{V}^{\otimes r})$, respectively.

For simplicity, we work with $\mathbb{F} = \mathbb{C}$ in the main body of the text. Then we can realize $\mathcal{E}(n, r) := \mathcal{E}_{\mathbb{C}}(n, r)$ and $\mathcal{P}(n, r)$ as subalgebras of $S_{\mathbb{C}}(n+1, r)$. In the case $\mathbb{F} = \mathbb{C}$, $\mathrm{GL}(n, \mathbb{C})$ will be simply denoted by GL_n .

0.3. The main purposes are double. One is to develop the Levi, and parabolic Schur algebras and the related representations. The another one is to investigate dualities of variant groups and algebras in the enhanced tensor representations. In this course, an important algebraic model of so-called degenerate double Hecke algebras¹ (DDHAs for short) is introduced, in the same spirit of degenerate affine Hecke algebras and the related (see [5] and [2]). Roughly speaking, the DDHA \mathcal{H}_r of type A_{r-1} is a combination of varieties of subalgebras generated compatibly by the group algebras $\mathbb{C}\mathfrak{S}_r$ and $\mathbb{C}\mathfrak{S}_l$ for all positive integers l not bigger than r . The l th DDHA \mathcal{H}_r^l is an associative algebra defined by generators \mathbf{s}_i , and \mathbf{x}_{σ} , $i = 1, \dots, r-1$, and $\sigma \in \mathfrak{S}_l$, and by relations:

$$\mathbf{s}_i \mathbf{x}_{\sigma} = \mathbf{x}_{s_i \circ \sigma}, \quad \mathbf{x}_{\sigma} \mathbf{s}_i = \mathbf{x}_{\sigma \circ s_i} \text{ for } \sigma \in \mathfrak{S}_l, i < l;$$

$$\mathbf{s}_i \mathbf{x}_{\sigma} = \mathbf{x}_{\sigma} = \mathbf{x}_{\sigma} \mathbf{s}_i \text{ for } \sigma \in \mathfrak{S}_l, i > l.$$

together with the defining relations of \mathfrak{S}_r and of \mathfrak{S}_l (see (3.1)-(3.6) for the complete defining relations of a DDHA). This is an infinite-dimensional algebra. Nevertheless, the l th DDHA

¹In the literature, double affine Hecke algebras (DAHA) arising from the study of KZ equations are familiar to authors (*ref.* [6]). Our DDHA have no direct relation with the former.

\mathcal{H}_r^l naturally arises from the tensor representation $(\mathbb{C}^{n+1})^{\otimes r}$ over GL_n . The core operators in $\mathrm{End}_{\mathbb{C}}((\mathbb{C}^{n+1})^{\otimes r})$ for $l < r$ and $\sigma \in \mathfrak{S}_l$, are presented as below

$$x_\sigma = \Psi_l(\sigma) \otimes \mathrm{id}^{\otimes r-l}$$

where $\Psi_l(\sigma)$ is a position permutation by σ in the first l th tensor factors (see the paragraph around (3.7) for the precise meaning). From those x_σ and all usual operators s_i interchanging the i th and $(i+1)$ th factors in $(\mathbb{C}^{n+1})^{\otimes r}$, \mathcal{H}_r^l naturally arises. Correspondingly, $(\mathbb{C}^{n+1})^{\otimes r}$ becomes a natural module over \mathcal{H}_r . This representation is denoted by Ξ . Then $\Xi(\mathcal{H}_r)$ becomes finite-dimensional, denoted by $D(n, r)$. On the enhanced-tensor representation space $\underline{V}^{\otimes r}$, the role of pair $(\mathcal{H}_r, D(n, r))$ is somewhat a counterpart of the one of pair $(\mathrm{GL}(V), S_{\mathbb{C}}(n, r))$ when we consider the usual tensor representation space $V^{\otimes r}$.

The above DDHAs are powerful ingredients in the course of establishing dualities for the enhanced tensor representations.

0.4. Below are the main results.

Theorem 0.1. (1) (Theorem 2.10) The parabolic Schur algebra $\mathcal{P}(n, r)$ has a basis $\{\xi_{\tilde{\pi}_{\mathbf{s}, \mathbf{j}}} \mid (\tilde{\pi}_{\mathbf{s}, \mathbf{j}}) \in E\}$, and

$$\dim \mathcal{P}(n, r) = \sum_{k=0}^r \binom{n+k-1}{k} \binom{n+k}{k}.$$

There is a set of canonical generators $\{\theta_{\mathbf{s}, \mathbf{t}}, \xi_{\mathbf{i}, \mathbf{j}} \mid (\mathbf{s}, \mathbf{t}) \in \Lambda, (\mathbf{i}, \mathbf{j}) \in D/\mathfrak{S}_r\}$ for $\mathcal{P}(n, r)$.

(2) (Theorems 6.2 and 6.4) The parabolic Schur algebra $\mathcal{P}(n, r)$ has one block. The isomorphism classes of irreducible modules and the indecomposable projective modules (PIM for short) are parameterized by the set of dominant weights $\Lambda^+ := \bigcup_{l=1}^r \Lambda^+(n, l)$, respectively. For an irreducible module $D_{\gamma'}$ and a PIM P_{γ} with $\gamma, \gamma' \in \Lambda^+$, set $a_{\gamma, \gamma'} := (P_{\gamma} : D_{\gamma'})$, and $\ell(\gamma) = \#\mathfrak{S}_n \cdot \gamma$. Then the Cartan invariants are presented as

$$a_{\gamma, \gamma'} = \begin{cases} 1, & \text{if } \gamma' = \gamma; \\ \ell(\gamma'), & \text{if } |\gamma'| < |\gamma|; \\ 0, & \text{otherwise.} \end{cases}$$

With aid of the structural description of the Levi subalgebra $\mathcal{L}(n, r)$ of the parabolic Schur algebra $\mathcal{P}(n, r)$, we establish a duality between DDHA and $\mathrm{GL}_n \times \mathbf{G}_{\mathbf{m}}$ in $\mathrm{End}_{\mathbb{C}}((\mathbb{C}^{n+1})^{\otimes r})$ and its consequence a ‘‘duality branching formula’’ in dimensions. As its application, we finally obtain the Levi/Parabolic Schur-Weyl duality stated below.

Theorem 0.2. For $\underline{V} = \mathbb{C}^{n+1}$, denote by $(\underline{V}^{\otimes r}, \Phi)$ the tensor representations of $\mathrm{GL}_{n+1} = \mathrm{GL}(\underline{V}, \mathbb{C})$, and by $(\underline{V}^{\otimes r}, \Psi)$ the permutation representations of the symmetric group \mathfrak{S}_r .

(1) (Theorem 4.3) Let \mathcal{H}_r be the degenerate double Hecke algebra associated with \mathfrak{S}_r . Then the following Levi Schur-Weyl duality holds.

$$\begin{aligned} \mathrm{End}_{\mathbb{C}\Phi(\mathrm{GL}_n \times \mathbf{G}_{\mathbf{m}})}(\underline{V}^{\otimes r}) &= \Xi(\mathcal{H}_r); \\ \mathrm{End}_{\Xi(\mathcal{H}_r)}(\underline{V}^{\otimes r}) &= \mathbb{C}\Phi(\mathrm{GL}_n \times \mathbf{G}_{\mathbf{m}}). \end{aligned}$$

(2) (Corollary 4.5) For $\lambda \in \Lambda^+(n, l)$ (a partition of l with n parts), denote by S_l^λ the irreducible module of \mathfrak{S}_l corresponding to λ . The following dimension formula can be regarded as a duality to the classical branching rule. For $l \in \{1, 2, \dots, r\}$, and $\lambda \in \Lambda^+(n, l)$,

$$\sum_{\substack{\mu \in \Lambda^+(n+1, r) \\ \lambda \text{ interlaces } \mu}} \dim S_r^\mu = \binom{r}{l} \dim S_l^\lambda$$

(3) (Theorem 5.3) $\text{End}_{\mathbb{C}\Phi(\underline{G} \times \mathbf{G}_m)}(\underline{V}^{\otimes r}) = \Xi(\mathcal{H}_r)^V$ for $G = \text{GL}(V)$.

It is worth mentioning that the study of invariants beyond reductive groups is a challenge (see [7]-[9]). Consequently the invariant property of $\underline{G} \times \mathbf{G}_m$ as above has its own interest.

In the same spirit, one can consider more general case for Levi- and parabolic subgroups of GL_{n+1} .

0.5. Our paper is organized as follows. In the first section, we introduce some basic notions and notations for semi-reductive groups. Then we give some fundamental properties of semi-reductive groups. In the second section, we introduce the Levi/Parabolic Schur algebras, and investigate their structure. In the third section, we introduce the degenerate double Hecke algebras and demonstrate their representation meaning in the enhanced tensor products. The fourth section is devoted to the proof of Theorem 4.3 and Corollary 4.5. With the above, in the fifth section we first prove the Levi and parabolic Schur Weyl dualities and then give some tensor invariants. In the concluding section, we investigate the representations of the parabolic Schur algebras, obtaining the results on the blocks and Cartan invariants.

0.6. As to other aspects, there will be some investigations somewhere else (to see [13], [17], [20], etc.). The parabolic Schur algebras can be defined in prime characteristic. Their modular representations will be an interesting topic in the future. It is worth mentioning that the nilpotent cone of $\mathfrak{gl}(V) := \text{Lie}(\underline{G})$ for $G = \text{GL}(V)$ is the same as the enhanced nilpotent cone studied by Achar-Henderson in [1]. Adjoint nilpotent orbits of \underline{G} in $\mathfrak{gl}(V)$ are compatible with enhanced nilpotent orbits studied in [1]. With respect to those, there are some interesting phenomenon parallel to the $\text{GL}(V)$ -theory in prime characteristic, including Jantzen's realization of support varieties of Weyl modules by the closures of some nilpotent orbits (see [12]).

1. SEMI-REDUCTIVE GROUPS AND SEMI-REDUCTIVE LIE ALGEBEAS

In this section, all vector spaces and varieties are over a field \mathbb{F} which stands for either the complex number field \mathbb{C} , or an algebraically closed field \mathbf{k} of characteristic $p > 0$.

1.1. Notions and notations.

Definition 1.1. An algebraic group G over \mathbb{F} is called semi-reductive if $G = G_0 \ltimes U$ with G_0 being a reductive subgroup, and U the unipotent radical. Let $\mathfrak{g} = \text{Lie}(G)$, and $\mathfrak{g}_0 = \text{Lie}(G_0)$ and $\mathfrak{u} = \text{Lie}(U)$, then $\mathfrak{g} = \mathfrak{g}_0 \oplus \mathfrak{u}$.

Example 1.2. (Enhanced reductive algebraic groups) Let G_0 be a connected reductive algebraic group over \mathbb{F} , and (M, ρ) be a finite-dimensional rational representation of G_0 with representation space M over \mathbb{F} . Consider the product variety $G_0 \times M$. Regard M as an additive algebraic group. The variety $G_0 \times M$ is endowed with an enhanced cross product structure denoted by $G_0 \times_\rho M$, by defining for any $(g_1, v_1), (g_2, v_2) \in G_0 \times M$

$$(g_1, v_1) \cdot (g_2, v_2) := (g_1 g_2, \rho(g_1) v_2 + v_1). \quad (1.1)$$

Then by a straightforward computation it is easily known that $\underline{G}_0 := G_0 \times_\rho M$ becomes a group with identity $(e, 0)$ for the identity $e \in G_0$, and $(g, v)^{-1} = (g^{-1}, -\rho(g)^{-1}v)$. And $G_0 \times_\rho M$ has a subgroup G_0 identified with $(G_0, 0)$ and a subgroup M identified with (e, M) . Furthermore, \underline{G}_0 is connected since G_0 and M are irreducible varieties. We call \underline{G}_0 an *enhanced reductive algebraic group associated with the representation space M* . What is more, G_0 and M are closed subgroups of \underline{G}_0 , and M is a normal closed subgroup. Actually, we have $(g, w)(e, v)(g, w)^{-1} = (e, \rho(g)v)$ for any $(g, w) \in G_0$. From now on, we will write down \dot{g} for $(g, 0)$ and e^v for (e, v) unless other appointments. It is clear that $e^v \cdot e^w = e^{v+w}$ for $v, w \in V$.

Suppose $\underline{\mathfrak{g}}_0 = \text{Lie}(\underline{G}_0)$. Then $(M, \mathfrak{d}(\rho))$ becomes a representation of $\underline{\mathfrak{g}}_0$. Naturally, $\text{Lie}(\underline{G}_0) = \underline{\mathfrak{g}}_0 \oplus M$, with Lie bracket

$$[(X_1, v_1), (X_2, v_2)] := ([X_1, X_2], \mathfrak{d}(\rho)(X_1)v_2 - \mathfrak{d}(\rho)(X_2)v_1),$$

which is called an enhanced reductive Lie algebra.

Clearly, \underline{G}_0 is a semi-reductive group with M being the unipotent radical.

2. ENHANCED TENSOR REPRESENTATIONS AND PARABOLIC SCHUR ALGEBRAS

From now on, the ground field \mathbb{F} will be \mathbb{C} . We suppose $G = \text{GL}(V)$ over \mathbb{C} , and suppose that $\underline{G} := G \times_\nu V$ is an enhanced group of G associated with the natural representation ν on V . All representations for algebraic groups are always assumed to be rational ones.

2.1. Enhanced natural modules. An irreducible G -module becomes naturally an irreducible module of \underline{G} with trivial V -action. The isomorphism classes of irreducible rational representations of \underline{G} coincide with the ones of G .

Denote by \underline{V} the one-dimensional extension of $V \cong \mathbb{C}^n$ via the vector η , i.e. $\underline{V} = V \oplus \mathbb{C}\eta \cong \mathbb{C}^{n+1}$. We will call \underline{V} the enhanced space (of V). Naturally \underline{V} becomes a \underline{G} -module which is defined for any $\underline{g} := (g, v) \in \underline{G}$ and $\underline{u} = u + a\eta \in \underline{V}$ with $g \in G$, $v, u \in V$ and $a \in \mathbb{C}$, via

$$\underline{g} \cdot \underline{u} = \nu(g)u + av + a\eta. \quad (2.1)$$

It is not hard to see that this module is a rational module of \underline{G} . The corresponding rational representation of \underline{G} is denoted by $\underline{\nu}$, which gives rise to a short exact sequence of \underline{G} -modules

$$V \hookrightarrow \underline{V} \rightarrow \mathbb{C} \quad (2.2)$$

where \mathbb{C} means the one-dimensional trivial \underline{G} -module. The following fact is clear.

Lemma 2.1. *The enhanced reductive algebraic group $\underline{\text{GL}}(V)$ is a closed subgroup of $\text{GL}(\underline{V})$.*

2.2. Classical Schur-Weyl duality. Recall the classical Schur-Weyl duality. The natural representation of $\text{GL}(V)$ on V gives rise to a $G = \text{GL}(V)$ -module on the tensor product $V^{\otimes r}$ for any given positive integer r , with diagonal g -action for any $g \in G$. The corresponding representation is denoted by ϕ . This means

$$\phi(g)(v_1 \otimes v_2 \otimes \cdots \otimes v_r) = \nu(g)v_1 \otimes \nu(g)v_2 \otimes \cdots \otimes \nu(g)v_r$$

for any $g \in \text{GL}(V)$ and any monomial tensor product $v_1 \otimes \cdots \otimes v_r \in V^{\otimes r}$.

In the meanwhile, $V^{\otimes r}$ naturally becomes a \mathfrak{S}_r -module with permutation action. The corresponding \mathfrak{S}_r -representation on $V^{\otimes r}$ is denoted by ψ . This means that for any $\sigma \in \mathfrak{S}_r$,

$$\psi(\sigma)(v_1 \otimes v_2 \otimes \cdots \otimes v_r) = v_{\sigma^{-1}(1)} \otimes v_{\sigma^{-1}(2)} \otimes \cdots \otimes v_{\sigma^{-1}(r)}. \quad (2.3)$$

The classical Schur-Weyl duality shows that the images of ϕ and ψ are double centralizers in $\text{End}_{\mathbb{C}}(V^{\otimes r})$, i.e. for $G = \text{GL}(V)$

$$\begin{aligned}\text{End}_G(V^{\otimes r}) &= \mathbb{C}\psi(\mathfrak{S}_r) \\ \text{End}_{\mathfrak{S}_r}(V^{\otimes r}) &= \mathbb{C}\phi(G)\end{aligned}\tag{2.4}$$

Here $\mathbb{C}\psi(\mathfrak{S}_r)$ and $\mathbb{C}\phi(G)$ stand for the subalgebras of $\text{End}_{\mathbb{C}}(V^{\otimes r})$ generated by $\psi(\mathfrak{S}_r)$ and $\phi(G)$ respectively.

2.3. Enhanced tensor representations. Now consider the r th tensor product $\underline{V}^{\otimes r}$ for a fixed positive integer r , which becomes a \underline{G} -modules by diagonal action.

From the classical Schur-Weyl duality (see (2.4)), $V^{\otimes r}$ has the following decomposition as $\text{GL}(V) \times \mathfrak{S}_r$ -modules

$$V^{\otimes r} = \bigoplus_{\lambda} L^{\lambda} \otimes D^{\lambda}\tag{2.5}$$

where λ in the sum ranges over the set of partitions of r with n parts (zero parts are allowed), both L^{λ} and D^{λ} are the irreducible highest weight module of $\text{GL}(V)$ and the irreducible module of \mathfrak{S}_r respectively, associated with the partition λ (see [10]).

We now consider the composition factors of $\underline{V}^{\otimes r}$ as a \underline{G} -module. Keeping (2.2) and (2.5) in mind, one easily has the following.

Proposition 2.2. *As a \underline{G} -module, the following formula holds in the Grothendieck group of the rational \underline{G} -module category*

$$[\underline{V}^{\otimes r}] = \sum_{\mu} c_{\mu} [L^{\mu}]$$

where the sum is over partitions μ of $l \in \{0, 1, \dots, r\}$ of length $\leq n$, and

$$c_{\mu} = \binom{r}{l} \dim S_l^{\mu}.$$

2.4. A preliminary to enhanced Schur-Weyl dualities. Denote by Φ the representation of \underline{G} on the r -tensor product module $\underline{V}^{\otimes r}$. Then the following question is naturally raised.

Question 2.3. What is the centralizer of $\mathbb{C}\Phi(\underline{G})$ in $\text{End}_{\mathbb{C}}(\underline{V}^{\otimes r})$? What role does \mathfrak{S}_r play in the enhanced case as in the classical Schur-Weyl duality?

It is not trivial to give a complete answer which is left till §5 (see Theorem 5.3). Here we give some preliminary investigation. Note that $\text{GL}(\underline{V})$ contains a closed subgroup \underline{G} (Lemma 2.1). According to the classical Schur-Weyl duality, $\text{GL}(\underline{V})$ and \mathfrak{S}_r are dual in the sense of the double centralizers in $\text{End}_{\mathbb{C}}(\underline{V}^{\otimes r})$ (we will still denote by Φ the natural representation of $\text{GL}(\underline{V})$ on $\underline{V}^{\otimes r}$ if the context is clear). So the significant investigation of Question 2.3 is to decide the centralizer of $\mathbb{C}\Phi(\underline{G})$ in $\text{End}_{\mathbb{C}}(\underline{V}^{\otimes r})$. For this, we first have the following observation

Lemma 2.4. *For any $\xi \in \text{End}_{\underline{G}}(\underline{V}^{\otimes r})$, $\xi(V^{\otimes r}) \subset V^{\otimes r}$.*

Proof. For any nonzero $w \in V^{\otimes r}$, we want to show that $\xi(w) \in V^{\otimes r}$. We might as well suppose $\xi(w) \neq 0$. Then we write

$$\xi(w) = \sum_{i=1}^t \underline{u}_{i1} \otimes \underline{u}_{i2} \otimes \cdots \otimes \underline{u}_{ir} \in \underline{V}^{\otimes r} \quad (2.6)$$

with $\underline{u}_{ij} = u_{ij} + a_{ij}\eta$, where $u_{ij} \in V$, and $a_{ij} \in \mathbb{C}$, and the number t of summands in (2.6) is minimal among all possible expression of $\xi(w)$.

By the assumption, $\xi(\Phi(\underline{g})w) = \Phi(\underline{g})\xi(w)$ for any $\underline{g} := (g, v) \in \underline{G}$ with $g \in G$ and $v \in V$. In particular, we take some special element $(e, v) \in \underline{G}$, denoted by e^v . Then we have an equation

$$\xi(\Phi(e^v)w) = \Phi(e^v)\xi(w). \quad (2.7)$$

By (2.1), $\underline{\nu}(e^v)(\underline{u}_{ij}) = u_{ij} + a_{ij}v + a_{ij}\eta$. So we have

$$\Phi(e^v)\xi(w) = \sum_i (u_{i1} + a_{i1}v + a_{i1}\eta) \otimes \cdots \otimes (u_{ir} + a_{ir}v + a_{ir}\eta).$$

On the other hand, by (2.1) again we have $\underline{\nu}(e^v)u = u$ for any $u \in V$. Hence, $\Phi(e^v)w = w$. We have

$$\xi(\Phi(e^v)w) = \sum_i \underline{u}_{i1} \otimes \cdots \otimes \underline{u}_{ir}.$$

By comparing both sides of (2.7), the arbitrariness of v leads to all a_{ij} being equal to zero. The proof is completed. \square

Proposition 2.5. *Denote by Ψ the permutation representation of \mathfrak{S}_r on $\underline{V}^{\otimes r}$ defined as in (2.3). The following statements.*

- (1) $\mathbb{C}\Psi(\mathfrak{S}_r)$ is a subalgebra of $\text{End}_{\underline{G}}(\underline{V}^{\otimes r})$.
- (2) There is a surjective homomorphism of algebras

$$\text{End}_{\underline{G}}(\underline{V}^{\otimes r}) \twoheadrightarrow \mathbb{C}\Psi(\mathfrak{S}_r).$$

Proof. (1) Note that \underline{G} is a subgroup of $\text{GL}(\underline{V})$ and then $\text{End}_{\text{GL}(\underline{V})}(\underline{V}^{\otimes r}) \subset \text{End}_{\underline{G}}(\underline{V}^{\otimes r})$. This statement follows from the classical Schur-Weyl duality with respect to $\text{GL}(\underline{V})$ and \mathfrak{S}_r .

(2) By Lemma 2.4, we can define a map Res from $\text{End}_{\underline{G}}(\underline{V}^{\otimes r})$ to $\text{End}_G(V^{\otimes r})$ by sending ξ to $\xi|_{V^{\otimes r}}$. Then it is easily seen that Res is an algebra homomorphism. Furthermore, we assert that Res is surjective. Actually, by the classical Schur-Weyl duality we have $\text{End}_G(V^{\otimes r}) = \mathbb{C}\Psi(\mathfrak{S}_r)$. So any element of $\text{End}_G(V^{\otimes r})$ can be expressed as $\sum_i a_i \sigma_i$ which is a finite \mathbb{C} -linear combination of some $\sigma \in \mathfrak{S}_r$. Take ξ to be a morphism in $\text{End}_{\mathbb{C}}(\underline{V}^{\otimes r})$ sending $\underline{u}_1 \otimes \cdots \otimes \underline{u}_r \in \underline{V}^{\otimes r}$ to

$$\sum_i a_i \underline{u}_{\sigma_i^{-1}(1)} \otimes \cdots \otimes \underline{u}_{\sigma_i^{-1}(r)}.$$

Then ξ is \underline{G} -equivariant. And $\text{Res}(\xi)$ is exactly $\sum_i a_i \sigma_i \in \text{End}_G(V^{\otimes r})$. So the assertion is proved. \square

2.5. Levi and parabolic subalgebras of $S(n+1, r)$. From now on, we always assume $\dim V = n$. Set $\mathcal{A} := \text{End}_{\mathbb{C}}(\underline{V}^{\otimes r})$, and $\mathcal{S} = \mathbb{C}\Phi(\text{GL}(\underline{V}))$, $\mathcal{E} := \mathbb{C}\Phi(\underline{G})$ and $\mathcal{C} := \mathbb{C}\Psi(\mathfrak{S}_r)$ for $\underline{G} = \text{GL}(V) \times_{\nu} V$. Call \mathcal{E} an enhanced Schur algebra which reflects the polynomial representations of degree r for the enhanced reductive algebraic group \underline{G} . Obviously, \mathcal{E} is a subalgebra of the semi-simple algebra \mathcal{S} . Note that $\mathbb{C}\Phi(\text{GL}(\underline{V})) = \text{End}_{\mathbb{C}}(\underline{V}^{\otimes r})^{\mathfrak{S}_r}$ is actually the classical Schur algebra $S(n+1, r)$ (see [11]). Lemma 2.1 still holds in any case. Correspondingly, we denote \mathcal{E} by $\mathcal{E}(n, r)$ more precisely.

By the above arguments, we have a sequence of the following subalgebras in \mathcal{A} :

$$S(n, r) \subset \mathcal{E}(n, r) \subset S(n+1, r).$$

Furthermore, we will have more interesting and related subalgebras: Levi Schur algebras and parabolic Schur algebras.

2.5.1. Let us first recall some facts on the classical Schur algebras. The readers refer to [11] or [4] for the details.

For a given positive integer m , set $\underline{m} = \{1, 2, \dots, m\}$. Denote by $A(m, r)$ the space consisting of the elements expressible as polynomials which are homogeneous of degree r in the polynomial function $c_{i,j}$ ($i, j \in \underline{m}$) on $\text{GL}(m, \mathbb{C})$. Then $A(m, r)$ has a basis (modulo the order of factors in the monomials)

$$c_{\mathbf{i}, \mathbf{j}} = c_{i_1 j_1} c_{i_2 j_2} \cdots c_{i_r j_r}$$

for $\mathbf{i} = (i_1, \dots, i_r), \mathbf{j} = (j_1, \dots, j_r) \in \underline{m}^r$. The symmetric group \mathfrak{S}_r acts on the left on \underline{m}^r by $\sigma \cdot \mathbf{i} = (i_{\sigma^{-1}(1)}, \dots, i_{\sigma^{-1}(r)})$. Furthermore, \mathfrak{S}_r act also on the set $\underline{m}^r \times \underline{m}^r$ by $\sigma \cdot (\mathbf{i}, \mathbf{j}) = (\sigma \cdot \mathbf{i}, \sigma \cdot \mathbf{j})$. So we can define an equivalence relation \sim on $\underline{m}^r \times \underline{m}^r$, $(\mathbf{i}, \mathbf{j}) \sim (\mathbf{k}, \mathbf{l})$ if and only if $(\mathbf{k}, \mathbf{l}) = \sigma \cdot (\mathbf{i}, \mathbf{j})$ for some $\sigma \in \mathfrak{S}_r$. The number of the equivalence classes (or the orbits) in \underline{m}^r under such a \mathfrak{S}_r -action is just $\mathbf{d}_{m,r} := \binom{m^2+r-1}{r}$. Then $A(m, r)$ becomes a coalgebra of dimension $\mathbf{d}_{m,r}$ with the coproduct δ and the counit ε as below

$$\begin{aligned} \Delta : A(m, r) &\rightarrow A(m, r) \otimes A(m, r), \\ c_{\mathbf{i}, \mathbf{j}} &\mapsto \sum_{\mathbf{k}} c_{\mathbf{i}, \mathbf{k}} \otimes c_{\mathbf{k}, \mathbf{j}}; \end{aligned} \quad (2.8)$$

and $\varepsilon(c_{\mathbf{i}, \mathbf{j}}) = \delta_{\mathbf{i}, \mathbf{j}}$.

Alternatively, the classical Schur algebra $S(m, r)$ can be defined as the dual of the $A(m, r)$. So $S(m, r)$ has basis $\{\xi_{\mathbf{i}, \mathbf{j}} \mid (\mathbf{i}, \mathbf{j}) \in \underline{m}^r \times \underline{m}^r / \sim\}$ dual to the basis $\{c_{\mathbf{i}, \mathbf{j}} \mid (\mathbf{i}, \mathbf{j}) \in \underline{m}^r \times \underline{m}^r / \sim\}$ of $A(m, r)$. This means

$$\xi_{\mathbf{i}, \mathbf{j}}(c_{\mathbf{k}, \mathbf{l}}) = \begin{cases} 1, & \text{if } (\mathbf{i}, \mathbf{j}) \sim (\mathbf{k}, \mathbf{l}); \\ 0, & \text{otherwise.} \end{cases}$$

Furthermore, $S(m, r)$ is an associative algebra of dimension $\mathbf{d}_{m,r}$ with the following multiplication rule

- (S1) $\xi_{\mathbf{i}, \mathbf{j}} \xi_{\mathbf{k}, \mathbf{l}} = \sum_{\mathbf{p}, \mathbf{q}} a_{\mathbf{i}, \mathbf{j}, \mathbf{k}, \mathbf{l}, \mathbf{p}, \mathbf{q}} \xi_{\mathbf{p}, \mathbf{q}}$, where $a_{\mathbf{i}, \mathbf{j}, \mathbf{k}, \mathbf{l}, \mathbf{p}, \mathbf{q}}$ equals the number of the elements $\mathbf{s} \in \underline{m}^r$ satisfying $(\mathbf{i}, \mathbf{j}) \sim (\mathbf{p}, \mathbf{s})$ and $(\mathbf{k}, \mathbf{l}) \sim (\mathbf{s}, \mathbf{q})$.
- (S2) $\xi_{\mathbf{i}, \mathbf{j}} \xi_{\mathbf{k}, \mathbf{l}} = 0$ unless $\mathbf{j} \sim \mathbf{k}$.
- (S3) $\xi_{\mathbf{i}, \mathbf{i}} \xi_{\mathbf{j}, \mathbf{j}} = \xi_{\mathbf{i}, \mathbf{j}} = \xi_{\mathbf{i}, \mathbf{j}} \xi_{\mathbf{j}, \mathbf{j}}$.

2.5.2. *A key lemma for Schur algebras.* Let $\Phi_m : \mathrm{GL}(m, \mathbb{C}) \rightarrow \mathrm{End}_{\mathbb{C}}(V^{\otimes r})$ for $V = \mathbb{C}^m$. By the classical Schur algebra theory, we can identify the image of Φ_m with $S(m, r)$. So we can write the image precisely.

In general, take g from $\mathrm{GL}(m, \mathbb{C})$ with $g = (g_{pq})_{m \times m}$. Then

$$\Phi_m(g) = \sum_{(\mathbf{p}, \mathbf{q}) \in \underline{m}^r \times \underline{m}^r / \mathfrak{S}_r} a_{\mathbf{p}\mathbf{q}} \xi_{\mathbf{p}, \mathbf{q}} \text{ with } a_{\mathbf{p}\mathbf{q}} = \prod_{i=1}^r g_{p_i q_i} \text{ for } \mathbf{p} = (p_1, \dots, p_r), \mathbf{q} = (q_1, \dots, q_r). \quad (2.9)$$

Set $I_g := \{(\mathbf{p}, \mathbf{q}) \in \underline{m}^r \times \underline{m}^r / \mathfrak{S}_r \mid a_{\mathbf{p}\mathbf{q}} \neq 0\}$. Then $\Phi_m(g)$ can be expressed as

$$\Phi_m(g) = \sum_{(\mathbf{p}, \mathbf{q}) \in I_g} a_{\mathbf{p}\mathbf{q}} \xi_{\mathbf{p}, \mathbf{q}}. \quad (2.10)$$

Set $I^0(g) = \{(p, q) \in \underline{m} \times \underline{m} \mid g_{pq} \neq 0\}$, and set $\mathbb{C}^\times = \mathbb{C} \setminus \{0\}$. Denote $B(g) := \{h \in \mathrm{GL}(m, \mathbb{C}) \mid I^0(h) = I^0(g)\}$. Then $B(g)$ can be regarded as an intersection of a non-empty open subset in $\mathbb{C}^{\#I^0(g)}$ with $(\mathbb{C}^\times)^{\#I^0(g)}$. So it is still a nonzero open subset of $\mathbb{C}^{\#I^0(g)}$.

Fix an order for all elements of the set I_g . Set $\lambda(g) = \#I_g$. Then we can talk about the matrix for $\Phi_m(g)$ for $g \in \mathrm{GL}(m, \mathbb{C})$. The following observation is fundamental, which will be important to the sequel arguments.

Lemma 2.6. *Suppose $S(m, r)$ is the classical Schur algebra associated with $\mathrm{GL}(m, \mathbb{C})$ and degree r . Keep the notations as above. The following statements hold.*

- (1) *For any given $g \in \mathrm{GL}(m, \mathbb{C})$, there exist $\lambda(g)$ elements $h^{(i)} \in B(g)$, $i = 1, \dots, \lambda(g)$ such that the matrix $(a_{\mathbf{p}\mathbf{q}}^{(i)})_{\lambda(g) \times \lambda(g)}$ is invertible, where $a_{\mathbf{p}\mathbf{q}}^{(i)}$ is defined in the same sense as in (2.9) with respect to $h^{(i)}$.*
- (2) *Consequently, for any $(\mathbf{p}, \mathbf{q}) \in I_g$, $\xi_{\mathbf{p}, \mathbf{q}} = \sum_{i=1}^{\lambda(g)} c_i \Phi_m(h^{(i)})$ for some $c_i \in \mathbb{C}$.*
- (3) *Furthermore, for any $(\mathbf{p}, \mathbf{q}) \in \underline{m}^r \times \underline{m}^r$, there exists $g \in \mathrm{GL}(m, \mathbb{C})$ such that $(\mathbf{p}, \mathbf{q}) \in I_g$. Therefore, (2) is valid for any basis element $\xi_{\mathbf{p}, \mathbf{q}}$ of $S(m, r)$.*

Proof. (1) For $g = (g_{ij})_{m \times m} \in \mathrm{GL}(m, \mathbb{C})$, we set $l = \lambda(g)$ and $I_g = \{\tau_1, \dots, \tau_l\}$. Then

$$\Phi_m(g) = \sum_{k=1}^l a_{\tau_k} \xi_{\tau_k} \text{ with } a_{\tau_k} = \prod_{i=1}^r g_{p_i q_i} \text{ if } \tau_k = (\mathbf{p}, \mathbf{q}) \in \underline{m}^r \times \underline{m}^r / \mathfrak{S}_r.$$

For $s \in \mathbb{N}$, set

$$\mathfrak{P}_s = \{(c_{ij})_{m \times m} \in (\mathbb{C}^\times)^{m^2} \mid (c_{ij}^s g_{ij})_{m \times m} \in \mathrm{GL}(m, \mathbb{C})\}.$$

Each \mathfrak{P}_s is a nonempty open subset of \mathbb{A}^{m^2} . In particular, $\cap_{s=1}^{l-1} \mathfrak{P}_s \neq \emptyset$. For $\mathbf{c} = (c_{ij})_{m \times m} \in \cap_{s=1}^{l-1} \mathfrak{P}_s$, let $g_{\mathbf{c}, s} = (c_{ij}^s g_{ij})_{m \times m} \in \mathrm{GL}(m, \mathbb{C})$ for $s = 1, \dots, l-1$. Then

$$\Phi_m(g_{\mathbf{c}, s}) = \sum_{k=1}^l f_k(c_{ij})^s a_{\tau_k} \xi_{\tau_k}$$

where $f_k(x_{ij})$ is a monomial of degree r over the n^2 variables x_{ij} ($1 \leq i, j \leq n$) and $f_s(x_{ij}) \neq f_t(x_{ij})$ for any $s \neq t$. Therefore,

$$\left(\bigcap_{j=1}^{l-1} \mathfrak{P}_j \right) \cap \left(\bigcap_{1 \leq s \neq t < l} \mathfrak{X}_{st} \right) \neq \emptyset,$$

where

$$\mathfrak{X}_{st} = \{(c_{ij})_{m \times m} \in (\mathbb{C}^\times)^{m^2} \mid f_s(c_{ij}) \neq f_t(c_{ij})\} \text{ for } 1 \leq s \neq t \leq l.$$

Now take

$$\mathbf{c} = (c_{ij})_{m \times m} \in \left(\bigcap_{j=1}^{l-1} \mathfrak{P}_j \right) \cap \left(\bigcap_{1 \leq s \neq t < l} \mathfrak{X}_{st} \right).$$

Set $h^{(1)} = g$ and $h^{(i)} = g_{\mathbf{c}, i-1}$ ($2 \leq i \leq l$). Then the corresponding matrix $(a_{\mathbf{p}\mathbf{q}}^{(i)})_{\lambda(g) \times \lambda(g)}$ forms a Vandermonde one. It is desired.

(2) follows directly from (1).

(3) Take $c_1, \dots, c_m \in \mathbb{C} \setminus \{0\}$ such that $c_i \neq c_j$ for any $1 \leq i \neq j \leq m$, and $g = (g_{ij})$ with $g_{ij} = c_j^{i-1}$ for $1 \leq i, j \leq m$. Then $g \in \text{GL}(m, \mathbb{C})$ and $(\mathbf{p}, \mathbf{q}) \in I_g$ for any $(\mathbf{p}, \mathbf{q}) \in \underline{m}^r \times \underline{m}^r$. \square

2.5.3. Now we look for a set of generators of $\mathcal{E}(n, r)$. Generally, we denote by $\mathcal{F}(\text{GL}_m)$ the set of functions on $\text{GL}_m := \text{GL}(m, \mathbb{C})$ for a positive integer m . Then we can naturally regard $\mathcal{F}(\text{GL}_n)$ as a subset of $\mathcal{F}(\text{GL}_{n+1})$, $A(n, r)$ as a sub-coalgebra of $A(n+1, r)$ and $S(n, r)$ as a subalgebra of $S(n+1, r)$. Consider a map

$$\theta : \mathcal{E}(n, r) \rightarrow S(n+1, r)$$

sending $\Phi(\underline{g})$ to $\theta_{\underline{g}}$ for $\underline{g} = (g, v)$. This is an algebra homomorphism. Note that $(g, v) = (g, 0)(e, g^{-1}v)$. So $\theta_{\underline{g}} = \theta_{(g, 0)}\theta_{(e, g^{-1}v)}$. By the definition of enhanced groups (see (1.1)), under θ we can identify $\Phi(\text{GL}(V))$ with $S(n, r)$. Still set $e^v = (e, v)$ for $v \in V$. Denote $\Omega(n, r)$ the subalgebra of $\mathcal{E}(n, r)$ generated by θ_{e^v} with v ranging over V . Then the first question is to understand θ_{e^v} and the subalgebra $\Omega(n, r)$ generated by them.

In the standard basis elements $c_{\mathbf{i}\mathbf{j}}$ of $A(m, r)$, the factor $c_{i,j}$ stands for the function which associates to each $g \in \text{GL}_m$ its (i, j) -coefficient g_{ij} .

Let us turn to $\text{GL}(\underline{V})$. Firstly, from now on we will always fix a basis

$$\{\eta_i \mid i = 1, 2, \dots, n+1\} \text{ for } \underline{V} \text{ with } \eta_{n+1} = \eta; \eta_1, \dots, \eta_n \in V,$$

and then identify $\text{GL}(\underline{V})$ with GL_{n+1} . In particular, e^v becomes the following $(n+1) \times (n+1)$ matrix

$$\begin{pmatrix} 1 & 0 & 0 & \cdots & 0 & 0 & a_1 \\ 0 & 1 & 0 & \cdots & 0 & 0 & a_2 \\ 0 & 0 & 1 & \cdots & 0 & 0 & a_3 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & 0 & \cdots & 0 & 1 & a_n \\ 0 & 0 & 0 & \cdots & 0 & 0 & 1 \end{pmatrix}.$$

where $v = \sum_{i=1}^n a_i \eta_i \in V$.

In the sequel, we always set $\mathcal{N} := \{1, 2, \dots, n+1\}$ (note that, in general, we set $\underline{l} = \{1, 2, \dots, l\}$ for the positive integer l throughout the paper, only with an exception for $n+1$ because we intend to stress this special situation in our paper). For $\mathbf{s} = (s_1, \dots, s_n) \in \mathbb{N}^n$, $\mathbf{t} = (t_1, \dots, t_{n+1}) \in \mathbb{N}^{n+1}$ we set $|\mathbf{s}| = \sum s_i$, $|\mathbf{t}| = \sum_i t_i$. Furthermore, we denote by $(\dots i^d \dots)$

an r -tuple in \mathbb{N}^r , where i appears d times continuously. By a straightforward computation we have

$$\theta_{e^v} = \sum_{(\mathbf{s}, \mathbf{t})} \left(\prod_{k=1}^n a_k^{s_k} \right) \xi_{(1^{s_1} 2^{s_2} \dots n^{s_n} 1^{t_1} 2^{t_2} \dots n^{t_n} (n+1)^{t_{n+1}}), ((n+1)^{|\mathbf{s}|} 1^{t_1} 2^{t_2} \dots n^{t_n} (n+1)^{t_{n+1}})}. \quad (2.11)$$

Here (\mathbf{s}, \mathbf{t}) in the sum runs through the range $\Lambda := \{(\mathbf{s}, \mathbf{t}) \in \mathbb{N}^n \times \mathbb{N}^{n+1} \mid |\mathbf{s}| + |\mathbf{t}| = r\}$, and we appoint 0^0 to be 1 if it appears in the coefficient $a_{\mathbf{s}} := \prod_{k=1}^n a_k^{s_k}$. Denote by $\theta_{\mathbf{s}, \mathbf{t}}$ the summand in the expression in (2.11) corresponding to (\mathbf{s}, \mathbf{t}) .

Set $\mathbb{N}_r^n := \{\mathbf{s} \in \mathbb{N}^n \mid |\mathbf{s}| \leq r\}$. For $\mathbf{s} \in \mathbb{N}_r^n$, set $\Lambda_{\mathbf{s}} := \{\mathbf{t} \in \mathbb{N}^{n+1} \mid (\mathbf{s}, \mathbf{t}) \in \Lambda\}$, and set

$$\Theta_{\mathbf{s}} := \sum_{\mathbf{t} \in \Lambda_{\mathbf{s}}} \theta_{\mathbf{s}, \mathbf{t}}.$$

Then

$$\theta_{e^v} = \sum_{\mathbf{s} \in \mathbb{N}_r^n} a_{\mathbf{s}} \Theta_{\mathbf{s}}.$$

Lemma 2.7. *The subalgebra $\Omega(n, r)$ has a basis $\{\Theta_{\mathbf{s}} \mid \mathbf{s} \in \mathbb{N}_r^n\}$. Therefore, $\dim \Omega(n, r) = \sum_{k=0}^r \binom{n+k-1}{k}$.*

Proof. For $a \in \mathbb{C}$, take $v_a = \sum_{i=1}^n a^{r-i-1} \eta_i \in V$. Then

$$\theta_{e^{v_a}} = \sum_{\mathbf{s} \in \mathbb{N}_r^n} a^{|\mathbf{s}|} \Theta_{\mathbf{s}},$$

where $|\mathbf{s}| = \sum_{i=1}^n r^{i-1} s_i$. Then $|\mathbf{s}| \neq |\mathbf{s}'|$ for any distinct $\mathbf{s}, \mathbf{s}' \in \mathbb{N}_r^n$. Denote $\lambda(n, r) = \#\mathbb{N}_r^n$. Take $c \in \mathbb{C}^\times$ and c is not a root of unit, and $a_i = c^i$ for $0 \leq i \leq \lambda(n, r) - 1$. This implies that $\Theta_{\mathbf{s}}$ is in the subspace spanned by $\{\theta_{e^{v_{a_i}}} \mid 0 \leq i \leq \lambda(n, r) - 1\}$ for any $\mathbf{s} \in \mathbb{N}_r^n$. In particular, $\Theta_{\mathbf{s}}$ is in the subalgebra generated by θ_{e^v} for $v \in V$. Consequently, $\Omega(n, r)$ coincides with the subalgebra of $\mathcal{E}(n, r)$ generated by θ_{e^v} for $v \in V$. And we have $\Omega(n, r) = \sum_{\mathbf{s} \in \mathbb{N}_r^n} \mathbb{C} \Theta_{\mathbf{s}}$.

Next we need to show that all $\Theta_{\mathbf{s}}$ are linearly independent. For any $\mathbf{s} \in \mathbb{N}_r^n$, take $\mathbf{t} \in \mathbb{N}^{n+1}$ such that $(\mathbf{s}, \mathbf{t}) \in \Lambda$. Denote by $c_{\mathbf{s}, \mathbf{t}}$ the basis element in $A(n+1, r)$ corresponding to $\xi_{\mathbf{s}, \mathbf{t}}$. Then $\Theta_{\mathbf{s}}(c_{\mathbf{s}, \mathbf{t}}) = 1$, and $\Theta_{\mathbf{s}}(c_{\mathbf{s}', \mathbf{t}'}) = 0$ whenever $\mathbf{s} \neq \mathbf{s}'$. For any given combinator equation $\sum_{\mathbf{s}} k_{\mathbf{s}} \Theta_{\mathbf{s}} = 0$, taking the value at $c_{\mathbf{s}, \mathbf{t}}$ we have $k_{\mathbf{s}} = 0$. Hence, those $\Theta_{\mathbf{s}}$ are proved to be linearly independent, so that the dimension formula follows. \square

2.5.4. *Levi and Parabolic subalgebras of $S(n+1, r)$.* There are canonical imbeddings of algebraic groups $\mathrm{GL}_n \hookrightarrow \mathrm{GL}_{n+1}$ given by

$$g \mapsto \mathrm{Diag}(g, 1) := \begin{pmatrix} g & 0 \\ 0 & 1 \end{pmatrix} \quad \forall g \in \mathrm{GL}(n),$$

and $\mathbf{G}_{\mathbf{m}} \hookrightarrow \mathrm{GL}_{n+1}$ given by $c \mapsto \mathrm{Diag}(1, \dots, 1, c) \quad \forall c \in \mathbf{G}_{\mathbf{m}} = \mathbb{C}^\times$, respectively. In this sense, both GL_n and $\mathbf{G}_{\mathbf{m}}$ can be regarded as closed subgroups of GL_{n+1} . Clearly, the closed subgroup in GL_{n+1} generated by GL_n and $\mathbf{G}_{\mathbf{m}}$ is isomorphic to $\mathrm{GL}_n \times \mathbf{G}_{\mathbf{m}}$. We identify both throughout the paper. In the same sense, we identify the closed subgroup generated by \underline{G} and $\mathbf{G}_{\mathbf{m}}$ with $\mathbf{G}_{\mathbf{m}} \rtimes \underline{G}$ (note that $\mathbf{G}_{\mathbf{m}}$ normalizes \underline{G}). Denote by $\mathcal{L}(n, r)$ the image of $\mathrm{GL}_n \times \mathbf{G}_{\mathbf{m}}$ under Φ which we call the Levi subalgebra of $S(n+1, r)$. And denote by $\mathcal{P}(n, r)$

the image of $\mathbf{G}_m \ltimes \underline{G}$ under Φ , which is called the parabolic subalgebra of $S(n+1, r)$. Clearly, when $r \geq 2$, $\mathcal{P}(n, r)$ is an extension of $\mathcal{E}(n, r)$ with the following exact sequence

$$\mathcal{E}(n, r) \hookrightarrow \mathcal{P}(n, r) \twoheadrightarrow \mathcal{Z}. \quad (2.12)$$

where \mathcal{Z} denotes $\Phi(\mathbf{G}_m)$. Note that $\underline{V}^{\otimes r}$ can be decomposed into a direct sum of subspaces: $\underline{V}^{\otimes r} = \bigoplus_{l=0}^r \underline{V}_l^{\otimes r}$ with $\underline{V}_l^{\otimes r} = \sum_{\mathbf{i} \in \mathcal{N}_l^r} \mathbb{C}\eta_{\mathbf{i}}$, $l = 0, 1, \dots, r$ (see §3.3 for \mathcal{N}_l^r). So we can list a basis of the $(r+1)$ -dimensional subalgebra $\Phi(\mathbf{G}_m)$ of $S(n+1, r)$ which consists of ϵ_l , $l = 0, 1, \dots, r$. Here ϵ_l is defined via

$$\epsilon_l|_{\underline{V}_q^{\otimes r}} = \begin{cases} \text{id}_{\underline{V}_q^{\otimes r}}; & \text{if } q = l, \\ 0; & \text{otherwise.} \end{cases} \quad (2.13)$$

Let us further investigate the Levi subalgebra $\mathcal{L}(n, r)$. We consider the set

$$D := \mathfrak{S}_r \cdot \{(\mathbf{p}_l(n+1)^{r-l}), (\mathbf{q}_l(n+1)^{r-l}) \in \mathcal{N}^r \times \mathcal{N}^r \mid \mathbf{p}_l, \mathbf{q}_l \in \underline{n}^l, l = 0, 1, \dots, r\}.$$

By (S1)-(S3), the span of $\xi_{\mathbf{i}, \mathbf{j}}$ with (\mathbf{i}, \mathbf{j}) ranging over D forms a subalgebra of $S(n+1, r)$, denoted by $\mathcal{L}(n, r)'$.

For a given $(g, c) \in \text{GL}_n \times \mathbf{G}_m$ with $g = (g_{pq})_{n \times n}$, by the classical Schur algebra theory,

$$\Phi(\text{Diag}(g, c))|_{\underline{V}^{\otimes r}} = \sum_{(\mathbf{p}, \mathbf{q}) \in \underline{n}^r \times \underline{n}^r / \mathfrak{S}_r} a_{\mathbf{p}, \mathbf{q}} \xi_{\mathbf{p}, \mathbf{q}} \text{ with } a_{\mathbf{p}, \mathbf{q}} = \prod_{i=1}^r g_{p_i q_i}. \quad (2.14)$$

Recall the notation $I_g = \{(\mathbf{p}, \mathbf{q}) \in \underline{n}^r \times \underline{n}^r / \mathfrak{S}_r \mid a_{\mathbf{p}, \mathbf{q}} \neq 0\}$. Now let us return to the case of the enhanced tensor space. Then $\Phi(\text{Diag}(g, c))$ can be expressed as

$$\Phi(\text{Diag}(g, c)) = \sum_{(\mathbf{p}, \mathbf{q}) \in I_g} T_{\mathbf{p}, \mathbf{q}}^{g, c} \quad (2.15)$$

where $T_{\mathbf{p}, \mathbf{q}}^{g, c} = \sum_{\substack{(\mathbf{i}, \mathbf{j}) \in D / \mathfrak{S}_r \\ (\mathbf{i}, \mathbf{j}) \preceq (\mathbf{p}, \mathbf{q})}} a_{\mathbf{i}, \mathbf{j}} \xi_{\mathbf{i}, \mathbf{j}}$ with

$$a_{\mathbf{i}, \mathbf{j}} := c^{r-t} \prod_{k=1}^t g_{p_k q_k} \quad (2.16)$$

where $(\mathbf{i}, \mathbf{j}) \preceq (\mathbf{p}, \mathbf{q})$ means

$$(\mathbf{i}, \mathbf{j}) = ((p_1 \dots p_t (n+1)^{r-t}), (q_1 \dots q_t (n+1)^{r-t})) \quad (2.17)$$

with $(\mathbf{p}, \mathbf{q}) \sim ((p_1 \dots p_r), (q_1 \dots q_r))$ for some integer t between 0 and r . We set $\text{rank}_n(\mathbf{i}) := t$ in (2.17). By the above arguments, $\mathcal{L}(n, r) \subset \mathcal{L}(n, r)'$.

Lemma 2.8. *The following statements hold.*

- (1) *The Levi subalgebra $\mathcal{L}(n, r)$ coincides with $\mathcal{L}(n, r)'$, this is to say, it is spanned by $\xi_{\mathbf{i}, \mathbf{j}}$ with (\mathbf{i}, \mathbf{j}) ranging over D / \mathfrak{S}_r .*
- (2) *Let $\mathcal{L}(n, r)_t$ be the subspace of $\mathcal{L}(n, r)$ spanned by $\xi_{\mathbf{i}, \mathbf{j}}$ with $\text{rank}_n(\mathbf{i}) = t$. Then $\mathcal{L}(n, r) = \sum_{t=0}^r \mathcal{L}(n, r)_t$ is a graded algebra.*

Proof. The second statement directly follows from the first one and the multiplication formula of $S(n, r)$. As to the first statement, it suffices to prove that for any $(\mathbf{i}, \mathbf{j}) \in D$, $\xi_{\mathbf{i}, \mathbf{j}}$ lies in $\mathcal{L}(n, r)$. Indeed, it is obvious that there exists $g \in \text{GL}_n$ such that $(\mathbf{i}, \mathbf{j}) \in I_g$. By applying Lemma 2.6, $\xi_{\mathbf{i}, \mathbf{j}} = \sum \mathbb{C}\Phi(h^{(i)})$ for some $h^{(i)} \in B(g) \subset \text{GL}_n \hookrightarrow \text{GL}_{n+1}$. This is desired. \square

Thanks to Lemma 2.6, we have another result.

Lemma 2.9. *The space $T(n, r)$ spanned by $\theta_{\mathbf{s}, \mathbf{t}}$ for all $(\mathbf{s}, \mathbf{t}) \in \Lambda$ is just the subalgebra generated by $\Phi(\mathbf{G}_{\mathbf{m}})$ and all $\theta_{(\mathbf{t}, v)}$ with \mathbf{t} running through the subgroup $T(n) \subset \mathrm{GL}(n, \mathbb{C})$ consisting of diagonal matrices, and v ranging over V . Consequently, all $\theta_{\mathbf{s}, \mathbf{t}}$ are contained in $\mathcal{P}(n, r)$.*

2.6. The structure of the parabolic Schur algebra $\mathcal{P}(n, r)$. Now it is a position to investigate the structure of $\mathcal{P}(n, r)$. For any given element in \mathcal{N}^r , modulo the order we can write it in a standard form:

$$(1^{s_1} 2^{s_2} \dots n^{s_n} (n+1)^{r-|\mathbf{s}|})$$

where $\mathbf{s} := (s_1, \dots, s_n) \in \mathbb{N}_r^n$. So for any basis element $\xi_{\mathbf{i}, \mathbf{j}} \in S(n+1, r)$ we can write

$$\xi_{\mathbf{i}, \mathbf{j}} = \xi_{(1^{s_1} 2^{s_2} \dots n^{s_n} (n+1)^{r-|\mathbf{s}|}, \mathbf{j})} \quad (2.18)$$

with $\mathbf{j} \in \mathcal{N}^r$. Set

$$\begin{aligned} E := \{(\tilde{\pi}_{\mathbf{s}}, \mathbf{j}) := ((1^{s_1} 2^{s_2} \dots n^{s_n} (n+1)^{r-|\mathbf{s}|}), (j_1 \dots j_{|\mathbf{s}|} (n+1)^{r-|\mathbf{s}|})) \in \mathcal{N}^r \times \mathcal{N}^r \\ | \mathbf{s} \in \mathbb{N}_r^n, (j_1 \dots j_{|\mathbf{s}|}) \in \mathcal{N}^{|\mathbf{s}|}\}, \end{aligned} \quad (2.19)$$

and $\tilde{E} = \mathfrak{S}_r \cdot E$.

Theorem 2.10. *Keep the above notations. The following statements hold.*

- (1) *The parabolic Schur algebra $\mathcal{P}(n, r)$ is generated by $\Omega(n, r)$ and $\mathcal{L}(n, r)$ with product axioms as (S1)-(S3), this is to say, $\mathcal{P}(n, r)$ is generated by $\Theta_{\mathbf{s}}$ and $\xi_{\mathbf{i}, \mathbf{j}}$, with $\mathbf{s} \in \mathbb{N}_r^n$ and $(\mathbf{i}, \mathbf{j}) \in D/\mathfrak{S}_r$.*
- (2) *The parabolic Schur algebra $\mathcal{P}(n, r)$ has a basis $\{\xi_{\tilde{\pi}_{\mathbf{s}}, \mathbf{j}} \mid (\tilde{\pi}_{\mathbf{s}}, \mathbf{j}) \in E\}$, and*

$$\dim \mathcal{P}(n, r) = \sum_{k=0}^r \binom{n+k-1}{k} \binom{n+k}{k}.$$

- (3) *Set*

$$\mathcal{P}(n, r)_i = \mathbb{C}\text{-span}\{\xi_{\tilde{\pi}_{\mathbf{s}}, \mathbf{j}} \in E \mid |\mathbf{s}| \geq i\} \text{ for } i = 0, 1, \dots, r,$$

and

$$\mathcal{P}(n, r)_{r+1} = \mathbb{C}\text{-span}\{\xi_{\tilde{\pi}_{\mathbf{s}}, \mathbf{j}} \in E \mid |\mathbf{s}| = r, \mathbf{j} \in \mathcal{N}^r \setminus \underline{n}^r\}.$$

Then $\mathcal{P}(n, r)_{i+1}$ is an ideal of $\mathcal{P}(n, r)_i$, $i = 0, \dots, r$. Moreover, $\mathcal{P}(n, r)_i / \mathcal{P}(n, r)_{i+1} \cong S(n, i) \rtimes \mathfrak{a}_i$ as algebras, where \mathfrak{a}_i is a nonzero abelian ideal for $i = 0, \dots, r-1$, and $\mathcal{P}(n, r)_r / \mathcal{P}(n, r)_{r+1} \cong S(n, r)$.

- (4) *Furthermore, $\mathcal{P}(n, r)$ has another set of generators $\{\theta_{\mathbf{s}, \mathbf{t}}, \xi_{\mathbf{i}, \mathbf{j}} \mid (\mathbf{s}, \mathbf{t}) \in \Lambda, (\mathbf{i}, \mathbf{j}) \in D/\mathfrak{S}_r\}$.*

Proof. The statement (1) follows from Lemmas 2.7 and 2.8.

(2) Denote by $\mathcal{P}(n, r)'$ the subspace of $S(n+1, r)$ spanned by $\{\xi_{\tilde{\pi}_{\mathbf{s}}, \mathbf{j}} \mid (\tilde{\pi}_{\mathbf{s}}, \mathbf{j}) \in E\}$. By definition, $\mathcal{P}(n, r) \subset \mathcal{P}(n, r)'$. What remains is to show that for any $\mathbf{z} = \xi_{\tilde{\pi}_{\mathbf{s}}, \mathbf{j}}$ with $(\tilde{\pi}_{\mathbf{s}}, \mathbf{j}) \in E$, \mathbf{z} must fall in $\mathcal{P}(n, r)$. Modulo the order, we can write $(\tilde{\pi}_{\mathbf{s}}, \mathbf{j}) \in E$ as $((p_1 p_2 \dots p_s (n+1)^{r-s}), (q_1 \dots q_t (n+1)^{s-t} (n+1)^{r-s}))$ with $(p_1 p_2 \dots p_s, q_1 \dots q_t) \in \underline{n}^s \times \underline{n}^t$ for $t \leq s$. Then

$$\mathbf{z} = \xi_{p_1 p_2 \dots p_s (n+1)^{r-s}, q_1 \dots q_t (n+1)^{r-t}}.$$

Consider

$$\mathbf{x} := \xi_{p_1 p_2 \dots p_s (n+1)^{r-s}, q_1 \dots q_t p_{t+1} \dots p_s (n+1)^{r-s}} \in \mathcal{L}(n, r)$$

and

$$\mathbf{y} := \xi_{q_1 \dots q_t p_{t+1} \dots p_s (n+1)^{r-s}, q_1 \dots q_t (n+1)^{r-t}} \in \Omega(n, r).$$

Thanks to (S1), it follows that $\mathbf{xy} = (c+1)\mathbf{z}$ where

$$c = \#\{\sigma \cdot (q_1, \dots, q_t, p_{t+1}, \dots, p_s) \mid \sigma \in (\mathfrak{S}_s)_{\mathbf{q}}^{\mathbf{p}}\}$$

for

$$(\mathfrak{S}_s)_{\mathbf{q}}^{\mathbf{p}} := \{\sigma \in \mathfrak{S}_s \mid \sigma \cdot (p_1, \dots, p_s) = (p_1, \dots, p_s), \text{ while } \sigma \cdot (q_1, \dots, q_t, p_{t+1}, \dots, p_s) \neq (q_1, \dots, q_t, p_{t+1}, \dots, p_s)\}.$$

Consequently, by Lemmas 2.9 and 2.8 we have that \mathbf{z} really lies in $\mathcal{P}(n, r)$. The dimension formula is obvious.

(3) By (S1)-(S3), it is readily known that $\mathcal{P}(n, r)_{i+1}$ is an ideal of $\mathcal{P}(n, r)_i$, $i = 0, \dots, r-1$. And

$$\mathcal{P}(n, r)_i / \mathcal{P}(n, r)_{i+1} = \mathfrak{s}_i \oplus \mathfrak{a}_i$$

where

$$\mathfrak{s}_i = \mathbb{C}\text{-span}\{\overline{\xi_{\tilde{\pi}_s, \mathbf{j}}} \mid (\tilde{\pi}_s, \mathbf{j}) \in E_i^{(1)}\}, \mathfrak{a}_i = \mathbb{C}\text{-span}\{\overline{\xi_{\tilde{\pi}_s, \mathbf{k}}} \mid (\tilde{\pi}_s, \mathbf{k}) \in E_i^{(2)}\}$$

with

$$E_i^{(1)} = \{(\tilde{\pi}_s, \mathbf{j}) := ((1^{s_1} 2^{s_2} \dots n^{s_n} (n+1)^{r-i}), \mathbf{j} = (j_1 \dots j_i (n+1)^{r-i})) \mid \mathbf{s} \in \mathbb{N}^n, |\mathbf{s}| = i, (j_1 \dots j_i) \in \underline{n}^i\}.$$

and

$$E_i^{(2)} = \{(\tilde{\pi}_s, \mathbf{k}) := ((1^{s_1} 2^{s_2} \dots n^{s_n} (n+1)^{r-i}), (k_1 \dots k_i (n+1)^{r-i})) \mid \mathbf{s} \in \mathbb{N}^n, |\mathbf{s}| = i, \mathbf{k} = (k_1 \dots k_i) \in \mathcal{N}^i \setminus \underline{n}^i\}.$$

The following mapping

$$\begin{aligned} \Xi : \mathfrak{s}_i &\rightarrow S(n, i) \\ \overline{\xi_{\tilde{\pi}_s, \mathbf{j}}} &\mapsto \xi_{((1^{s_1} 2^{s_2} \dots n^{s_n}), (j_1 \dots j_i))} \end{aligned}$$

gives an algebra isomorphism between \mathfrak{s}_i and $S(n, i)$. Furthermore, \mathfrak{a}_i is an ideal of $\mathcal{P}(n, r)_i / \mathcal{P}(n, r)_{i+1}$.

(4) This statement follows from (1) and Lemma 2.9. \square

3. DEGENERATE DOUBLE HECKE ALGEBRAS

In this section, we introduce degenerate double Hecke algebras which will be important in the sequel arguments. For the symmetric group \mathfrak{S}_r , we denote by $s_l = (l, l+1)$ for $l = 1, \dots, r-1$, the transposition just interchanging l and $l+1$, and fixing the others.

3.1. Degenerate double Hecke algebras.

3.1.1. For given positive integers r and l with $r > l$, we consider the following algebra \mathcal{H}_r^l defined by generators $\{\mathbf{x}_\sigma \mid \sigma \in \mathfrak{S}_l\} \cup \{\mathbf{s}_i \mid i = 1, 2, \dots, r-1\}$ and relations as below.

$$\mathbf{s}_i^2 = 1, \quad \mathbf{s}_i \mathbf{s}_j = \mathbf{s}_j \mathbf{s}_i \text{ for } 0 < i \neq j \leq r-1, |j-i| > 1; \quad (3.1)$$

$$\mathbf{s}_i \mathbf{s}_j \mathbf{s}_i = \mathbf{s}_j \mathbf{s}_i \mathbf{s}_j \text{ for } 0 < i \neq j \leq r-1, |j-i| = 1; \quad (3.2)$$

$$\mathbf{x}_\sigma \mathbf{x}_\mu = \mathbf{x}_{\sigma \circ \mu} \text{ for } \sigma, \mu \in \mathfrak{S}_l; \quad (3.3)$$

$$\mathbf{s}_i \mathbf{x}_\sigma = \mathbf{x}_{s_i \circ \sigma}, \quad \mathbf{x}_\sigma \mathbf{s}_i = \mathbf{x}_{\sigma \circ s_i} \text{ for } \sigma \in \mathfrak{S}_l, i < l; \quad (3.4)$$

$$\mathbf{s}_i \mathbf{x}_\sigma = \mathbf{x}_\sigma = \mathbf{x}_\sigma \mathbf{s}_i \text{ for } \sigma \in \mathfrak{S}_l, i > l. \quad (3.5)$$

This is an infinite-dimensional associative algebra. We call \mathcal{H}_r^l the l th degenerate double Hecke algebra of \mathfrak{S}_r . By (3.3), the subalgebra X_l generated by \mathbf{x}_σ for $\sigma \in \mathfrak{S}_l$ is isomorphic to $\mathbb{C}\mathfrak{S}_l$. As well as being a subalgebra of \mathcal{H}_r^l , $\mathbb{C}\mathfrak{S}_r$ is also a quotient, via the homomorphism $\mathcal{H}_r^l \rightarrow \mathbb{C}\mathfrak{S}_r$ mapping $\mathbf{s}_i \mapsto s_i$ and $\mathbf{x}_\sigma \mapsto 0$ for each $i = 1, \dots, r-1$ and $\sigma \in \mathfrak{S}_l$.

Additionally, we make an appointment that $\mathcal{H}_r^0 := \langle \mathfrak{S}_r, X_0 \rangle$ with $X_0 = \mathbb{C}\mathbf{x}_\emptyset$ satisfying $f = f \cdot \mathbf{x}_\emptyset = \mathbf{x}_\emptyset \cdot f$ for $f \in \mathcal{H}_r^0$; and $\mathcal{H}_r^r := \langle \mathbb{C}\mathbf{s}_i, \mathbf{x}_\sigma \rangle$ with all $\mathbf{s}_i, \mathbf{x}_\sigma, i = 1, \dots, r-1$ and $\sigma \in \mathfrak{S}_r$ satisfying (3.1)-(3.4). Here and after, $\langle \square \rangle$ stands for a \mathbb{C} -algebra generated by \square . Then $\mathcal{H}_r^0 \cong \mathbb{C}\mathfrak{S}_r$, and $\mathcal{H}_r^r \cong \mathbb{C}\mathfrak{S}_r^{[2]}$ with $\mathfrak{S}_r^{[2]}$ being a group giving rise to a non-split extension $\mathfrak{S}_r \hookrightarrow \mathfrak{S}_r^{[2]} \rightarrow \mathfrak{S}_r$.

3.1.2. *Full degenerate double Hecke algebras.* Now we combine all \mathcal{H}_r^l ($l = 0, 1, \dots, r$) into a full degenerate double Hecke algebras.

Definition 3.1. The degenerate double Hecke algebra \mathcal{H}_r of \mathfrak{S}_r is an associative algebra with generators \mathbf{s}_i ($i = 1, \dots, r-1$), and $\mathbf{x}_\sigma^{(l)}$ for $\sigma \in \mathfrak{S}_l, l = 0, 1, \dots, r$, and with relations as (3.1)-(3.5) in which $\mathbf{x}_\sigma, \mathbf{x}_\mu$ are replaced by $\mathbf{x}_\sigma^{(l)}, \mathbf{x}_\mu^{(l)}$, and additional ones:

$$\mathbf{x}_\delta^{(l)} \mathbf{x}_\gamma^{(k)} = 0 \text{ for } \delta \in \mathfrak{S}_l, \gamma \in \mathfrak{S}_k \text{ with } k \neq l. \quad (3.6)$$

Naturally, as well as being a subalgebra of \mathcal{H}_r , $\mathbb{C}\mathfrak{S}_r$ is also a quotient of \mathcal{H}_r , via the homomorphism $\mathcal{H}_r \rightarrow \mathbb{C}\mathfrak{S}_r$ mapping $\mathbf{s}_i \mapsto s_i$ and $\mathbf{x}_\sigma^{(l)} \mapsto 0$ for each $i = 1, \dots, r-1$ and $\sigma \in \mathfrak{S}_l$, and $l = 0, 1, \dots, r$.

3.2. Degenerate double Hecke algebras arise from the following question:

Question 3.2. For the tensor representation $\underline{V}^{\otimes r}$ over the Levi-subgroup $\mathrm{GL}(V) \times \mathbf{G}_m$, $\mathrm{End}_{\mathbb{C}}(\underline{V}^{\otimes r})^{\mathrm{GL}(V) \times \mathbf{G}_m} = ?$

Let us begin the arguments with turning to $\underline{V}^{\otimes r}$ which is regarded as a $\mathrm{GL}(V)$ -module by fixing η and a \mathbf{G}_m -module by fixing V .

Keep the notations as before. In particular, we fix a basis $\{\eta_1, \dots, \eta_n; \eta_{n+1} := \eta\}$ for $\underline{V} = V \oplus \mathbb{C}\eta$ with $V = \sum_{i=1}^n \mathbb{C}\eta_i$. Associated with this basis, $\mathrm{GL}(V) = \mathrm{GL}(n, \mathbb{C})$, and $\mathrm{GL}(\underline{V}) = \mathrm{GL}(n+1, \mathbb{C})$. And $\mathrm{GL}(n, \mathbb{C})$ is canonically regarded as a subgroup of $\mathrm{GL}(n+1, \mathbb{C})$ by the established imbedding $\mathrm{GL}(V) \hookrightarrow \underline{\mathrm{GL}}(V) \hookrightarrow \mathrm{GL}(\underline{V})$ sending $g \in \mathrm{GL}(V)$ to $(g, 0) \in \underline{\mathrm{GL}}(V)$ (see Example 1.2 for the notations).

In this view, we already know that the image of $\mathrm{GL}(V) \times \mathbf{G}_m$ by Φ is $\mathcal{L}(n, r)$ (§2.5.4). In the following we will exactly determine $\mathrm{End}_{\mathcal{L}(n, r)}(\underline{V}^{\otimes r})$ and then deduce the Levi version of Schur-Weyl duality for $\mathrm{GL}(n, \mathbb{C}) \times \mathbf{G}_m$.

3.3. Recall that all $\eta_{\mathbf{i}} = \eta_{i_1} \otimes \eta_{i_2} \otimes \dots \otimes \eta_{i_r}$ for $\mathbf{i} = (i_1, \dots, i_r) \in \mathcal{N}^r$ form a basis of $\underline{V}^{\otimes r}$. For a given $\mathbf{j} = (j_1, \dots, j_r) \in \mathcal{N}^r$, there exists a unique $l \in \{0, 1, \dots, r\}$ and $\mathbf{j}' = (j'_1, \dots, j'_l) \in \underline{\mathcal{N}}^l$ such that $\mathbf{j} \sim (\mathbf{j}'(n+1)^{r-l})$, l is called the \underline{n} -rank of \mathbf{j} , denoted by $\mathrm{rk}_{\underline{n}}(\mathbf{j})$. All elements with \underline{n} -rank equal to l constitute a subset of \mathcal{N}^r , denoted by \mathcal{N}_l^r . Clearly, $\mathcal{N}^r = \bigcup_{l=0}^r \mathcal{N}_l^r$. And $\underline{V}^{\otimes r}$ is decomposed into a direct sum of subspaces: $\underline{V}^{\otimes r} = \bigoplus_{l=0}^r \underline{V}_l^{\otimes r}$ for $\underline{V}_l^{\otimes r} = \sum_{\mathbf{i} \in \mathcal{N}_l^r} \mathbb{C}\eta_{\mathbf{i}}$, $l = 0, 1, \dots, r$.

Clearly, each $\underline{V}_l^{\otimes r}$ is stabilized under \mathfrak{S}_r -action. Hence this action gives rise to a representation of \mathfrak{S}_r on $\underline{V}_l^{\otimes r}$, denoted by $\Psi|_l$. For a subset $I = \{i_1, \dots, i_l\} \subset \underline{r}$ whose elements are

assumed to be ordered increasingly, we can write a subspace $\underline{V}_I^{\otimes r}$ of $\underline{V}^{\otimes r}$ as

$$\underline{V}_I^{\otimes r} := \mathbb{C}\text{-span}\{\eta_{\mathbf{j}} = \eta_{j_1} \otimes \cdots \otimes \eta_{j_r} \mid j_{i_k} \in \underline{n}, k = 1, \dots, l; j_d = n+1 \text{ for } d \neq i_k\}.$$

This means, for any $\eta_{\mathbf{j}} \in \underline{V}_I^{\otimes r}$, there exists

$$\tau_I = \begin{pmatrix} 1 & 2 & \cdots & r \\ i_1 & i_2 & \cdots & i_r \end{pmatrix} \in \mathfrak{S}_r$$

such that $\Psi(\tau_I)\eta_{\mathbf{j}} = \eta_{\mathbf{j}_{i(n+1)^{r-l}}} \in \underline{V}_{\underline{l}}^{\otimes r}$ for some $\mathbf{j}_l \in \underline{n}^l$, where $\{i_{l+1}, \dots, i_n\} = \{j_d \mid d \in \underline{r} \setminus I\}$.

Conversely, for any $\eta_{\mathbf{j}} \in \underline{V}_{\underline{l}}^{\otimes r}$, there exists $\tau \in \mathfrak{S}_r$ and $\eta_{\mathbf{j}'}$ such that $\eta_{\mathbf{j}} = \Psi(\tau)(\eta_{\mathbf{j}'})$.

What is more, as a vector space

$$\underline{V}^{\otimes r} = \bigoplus_{I \subset \underline{r}} \underline{V}_I^{\otimes r}$$

where I in the sum ranges over all subsets of \underline{r} consisting of l elements. In the above, each $\underline{V}_I^{\otimes r}$ is a $\text{GL}(V)$ -module. The corresponding representation is a subrepresentation of $\text{GL}(V)$ on $\underline{V}_I^{\otimes r}$, the latter of which is also denoted by Φ for brevity.

For $I \subset \underline{r}$ with $\#I = l$, denote by $\text{Sym}(I)$ the symmetric group of I consisting all permutations of the I , which is isomorphic to \mathfrak{S}_l . Naturally, any $\sigma \in \text{Sym}(I)$ gives rise to a transformation on $\underline{V}_I^{\otimes r}$ which just permutes the set $\{\eta_{\mathbf{j}} \in \underline{V}_I^{\otimes r}\}$ via changing the position of factor η_{j_i} into the position of $\eta_{j_{\sigma^{-1}(i)}}$ for all $i \in I$ and fixing the other factors. This gives rise to an representation of $\text{Sym}(I)$ on $\underline{V}_I^{\otimes r}$. For $I = \underline{l}$, the corresponding symmetric group is directly denoted by \mathfrak{S}_l . The corresponding representation is denoted by $\Psi|_{\underline{l}}$.

3.4. Representations of \mathcal{H}_r on $\underline{V}^{\otimes r}$. On $V^{\otimes l}$, there is a permutation representation Ψ_l^V of \mathfrak{S}_l defined via $\Psi_l^V(\sigma)$ sending $v_1 \otimes v_2 \otimes \cdots \otimes v_l$ onto $v_{\sigma^{-1}(1)} \otimes v_{\sigma^{-1}(2)} \otimes \cdots \otimes v_{\sigma^{-1}(l)}$ for $\sigma \in \mathfrak{S}_l$. Keep in mind the notations $\underline{l} = \{1, 2, \dots, l\} \subset \underline{n}$, and $\underline{V}_{\underline{l}}^{\otimes r} = V^{\otimes l} \otimes \eta^{\otimes r-l}$. Extending Ψ_l^V , we define the following linear operator on $\underline{V}_{\underline{l}}^{\otimes r}$

$$x_\sigma = \Psi_l^V(\sigma) \otimes \text{id}^{\otimes r-l} \in \text{End}_{\mathbb{C}}(\underline{V}_{\underline{l}}^{\otimes r}) \quad (3.7)$$

for $\sigma \in \mathfrak{S}_l$. Next we extend x_σ to an element $x_\sigma^{\underline{l}}$ of $\text{End}_{\mathbb{C}}(\underline{V}_I^{\otimes r})$ by annihilating any other summand $\underline{V}_I^{\otimes r}$ with $I \neq \underline{l}$.

Now let us look at the representation meaning of $\mathbf{s}_i \in \mathcal{H}_r^l$ in $\text{End}_{\mathbb{C}}(\underline{V}_I^{\otimes r})$. This one keeps the role of $\Psi(s_i)$ such that the conjugation of $x_\sigma^{\underline{l}}$ by $\Psi(s_i)$ will be an operator translating $x_\sigma^{\underline{l}} \in \text{End}_{\mathbb{C}}(\underline{V}_{\underline{l}}^{\otimes r})$ to the forthcoming parallel one $x_\sigma^I \in \text{End}_{\mathbb{C}}(\underline{V}_I^{\otimes r})$ for $I = \{1, 2, \dots, l-1, l+1\}$.

In general, for $\eta_{\mathbf{j}} \in \underline{V}_I^{\otimes r}$ with $I \subset \underline{r}$ and $\#I = l$, we can write $\eta_{\mathbf{j}} = \Psi(\tau_I)\eta_{\mathbf{j}'_{i(n+1)^{r-l}}}$ for some $\mathbf{j}'_l \in \underline{n}^l$. Then $\Psi(\tau_I) \circ x_\sigma \circ \Psi(\tau_I^{-1})$ lies in $\text{End}_{\mathbb{C}}(\underline{V}_I^{\otimes r})$ for any $\sigma \in \mathfrak{S}_l$, which extends to an element of $\text{End}_{\mathbb{C}}(\underline{V}_I^{\otimes r})$ by annihilating any other summand $\underline{V}_J^{\otimes r}$ with $J \neq I$. This elements is denoted by x_σ^I . All x_σ^I ($\sigma \in \mathfrak{S}_l$) generate a subalgebra in $\text{End}_{\mathbb{C}}(\underline{V}_I^{\otimes r})$, denoted by E_I which is isomorphic to $\mathbb{C}\Psi_l^V(\mathfrak{S}_l)$. Set $x_e^{\underline{l}} := \sum_I x_e^I$, where I in the sum ranges over all subsets of \underline{r} containing l elements, and e represents the identity element in \mathfrak{S}_l . Then $x_e^{\underline{l}}$ is just the identity mapping on $\underline{V}_I^{\otimes r}$. Sometimes, e also indicates the identity element in \mathfrak{S}_r if the context is clear.

Now it is a position to demonstrate a representation of the degenerate double Hecke algebra \mathcal{H}_r on $\underline{V}^{\otimes r}$.

Lemma 3.3. *The following statements hold.*

- (1) For $1 \leq l \leq r$, there is an algebra homomorphism $\Xi_l : \mathcal{H}_r^l \rightarrow \text{End}_{\mathbb{C}}(\underline{V}_l^{\otimes r})$ defined by sending $\mathbf{s}_i \mapsto \Psi|_l(s_i)$ and $\mathbf{x}_\sigma \mapsto x_\sigma^l$.
- (2) For $l = 0$, there is an algebra homomorphism $\Xi_0 : \mathcal{H}_r^0 \rightarrow \text{End}_{\mathbb{C}}(\underline{V}_l^{\otimes r})$ defined by sending $\mathbf{s}_i \mapsto \Psi|_l(s_i) = \text{id}$.
- (3) For any $l \in \{0, 1, \dots, r\}$, $\Phi(\dot{g})$ for any $g \in \text{GL}(V)$ commutes with any elements from $\Xi_l(\mathcal{H}_r^l)$ in $\text{End}_{\mathbb{C}}(\underline{V}_l^{\otimes r})$.
- (4) Set $E_l := \bigoplus_{I \in \mathcal{S}_l} E_I$ with $\mathcal{S}_l := \{I \subset \underline{r} \mid \#I = l\}$, correspondingly $E_0 := E_\emptyset$. Then E_l is a left $\Psi|_l(\mathfrak{S}_r)$ -module under the conjugation, i.e. $\Psi|_l(\tau).x_\sigma^I = \Psi|_l(\tau)x_\sigma^I\Psi|_l(\tau)^{-1}$ for $\tau \in \mathfrak{S}_r$. Furthermore, $\Xi_l(\mathcal{H}_r^l) = \Psi|_l(\mathbb{C}\mathfrak{S}_r)E_l$.
- (5) On the enhanced tensor space $\underline{V}^{\otimes r}$, there is a representation Ξ of \mathcal{H}_r defined via:
 - (5.1) $\Xi|_{\mathbb{C}\mathfrak{S}_r} = \Psi$;
 - (5.2) For any $\mathbf{x}_\sigma \in \mathcal{H}_r^l$, $l = 0, 1, \dots, r$, $\Xi(\mathbf{x}_\sigma)|_{\underline{V}_l^{\otimes r}} = \Xi_l(\mathbf{x}_\sigma)$ and $\Xi(\mathbf{x}_\sigma)|_{\underline{V}_k^{\otimes r}} = 0$ for $k \neq l$.

Proof. (1) For $l \leq r$, we need to show that Ξ_l keeps the relations (3.1)-(3.5).

Recall that for any $1 \leq i \leq r-1$, $\mathbf{j} \in \mathcal{N}^r$, we have

$$\begin{aligned} \Xi_l(\mathbf{s}_i)(\eta_{\mathbf{j}}) &= \Psi|_l(s_i)(\eta_{\mathbf{j}}) \\ &= \eta_{\mathbf{s}_i(\mathbf{j})} \\ &= \eta_{j_1} \otimes \cdots \otimes \eta_{j_{i+1}} \otimes \eta_{j_i} \otimes \cdots \otimes \eta_{j_n}. \end{aligned}$$

Hence, it is readily known that

$$\Xi_l(\mathbf{s}_i)^2 = \text{id}, \quad \Xi_l(\mathbf{s}_i)\Xi_l(\mathbf{s}_j) = \Xi_l(\mathbf{s}_j)\Xi_l(\mathbf{s}_i) \text{ for } 0 \leq i \neq j \leq r-1, |j-i| > 1; \quad (3.8)$$

and

$$\Xi_l(\mathbf{s}_i)\Xi_l(\mathbf{s}_j)\Xi_l(\mathbf{s}_i) = \Xi_l(\mathbf{s}_j)\Xi_l(\mathbf{s}_i)\Xi_l(\mathbf{s}_j) \text{ for } 0 \leq i \neq j \leq r-1, |j-i| = 1. \quad (3.9)$$

For any $\sigma, \mu \in \mathfrak{S}_l$, and $\mathbf{j} = (\mathbf{j}_l(n+1)^{r-l})$ with $\mathbf{j}_l \in \underline{n}^l$, we have

$$\Xi_l(\mathbf{x}_\sigma) \circ \Xi_l(\mathbf{x}_\mu)(\eta_{\mathbf{j}}) = \eta_{\sigma\mu(\mathbf{j})} = \Xi_l(\mathbf{x}_{\sigma\mu})(\eta_{\mathbf{j}}).$$

And $\Xi_l(\mathbf{x}_\sigma) \circ \Xi_l(\mathbf{x}_\mu)(\eta_{\mathbf{k}}) = 0 = \Xi_l(\mathbf{x}_{\sigma\mu})(\eta_{\mathbf{k}})$ for any $\mathbf{k} \in \mathcal{N}^r$ with $\eta_{\mathbf{k}} \notin \underline{V}_l^r$. Hence,

$$\Xi_l(\mathbf{x}_\sigma) \circ \Xi_l(\mathbf{x}_\mu) = \Xi_l(\mathbf{x}_{\sigma\mu}) = \Xi_l(\mathbf{x}_\sigma\mathbf{x}_\mu), \quad \forall \sigma, \mu \in \mathfrak{S}_l. \quad (3.10)$$

For any $\sigma \in \mathfrak{S}_l$, $i < l$, and $\mathbf{j} = (\mathbf{j}_l(n+1)^{r-l})$ with $\mathbf{j}_l = (j_1, \dots, j_l) \in \underline{n}^l$, we have

$$\begin{aligned} \Xi_l(\mathbf{s}_i) \circ \Xi_l(\mathbf{x}_\sigma)(\eta_{\mathbf{j}}) &= \eta_{(j_{\sigma^{-1}(1)} \cdots j_{\sigma^{-1}(i+1)} j_{\sigma^{-1}(i)} \cdots j_{\sigma^{-1}(l)} (n+1)^{r-l})} \\ &= \Xi_l(\mathbf{x}_{\mathbf{s}_i \circ \sigma})(\eta_{\mathbf{j}}). \end{aligned}$$

And $\Xi_l(\mathbf{s}_i) \circ \Xi_l(\mathbf{x}_\sigma)(\eta_{\mathbf{k}}) = 0 = \Xi_l(\mathbf{x}_{\mathbf{s}_i \circ \sigma})(\eta_{\mathbf{k}})$ for any $\mathbf{k} \in \mathcal{N}^r$ with $\eta_{\mathbf{k}} \notin \underline{V}_l^r$. Hence,

$$\Xi_l(\mathbf{s}_i) \circ \Xi_l(\mathbf{x}_\sigma) = \Xi_l(\mathbf{s}_i\mathbf{x}_\sigma), \quad \sigma \in \mathfrak{S}_l, i < l. \quad (3.11)$$

Moreover, similar arguments yield that

$$\Xi_l(\mathbf{x}_\sigma) \circ \Xi_l(\mathbf{s}_i) = \Xi_l(\mathbf{x}_\sigma\mathbf{s}_i), \quad \sigma \in \mathfrak{S}_l, i < l. \quad (3.12)$$

and

$$\Xi_l(\mathbf{s}_i) \circ \Xi_l(\mathbf{x}_\sigma) = \Xi_l(\mathbf{x}_\sigma) = \Xi_l(\mathbf{x}_\sigma\mathbf{s}_i) \text{ for } \sigma \in \mathfrak{S}_l, i > l. \quad (3.13)$$

Now it follows from (3.8)-(3.13) that Ξ_l is an algebra homomorphism from \mathcal{H}_r^l to $\text{End}_{\mathbb{C}}(\underline{V}_l^{\otimes r})$.

(2) In this situation, Ξ_0 obviously keeps the relations (3.1)-(3.5). Hence, Ξ_0 is an algebra homomorphism from \mathcal{H}_r^0 to $\text{End}_{\mathbb{C}}(\underline{V}_l^{\otimes r})$.

(3) It suffices to show that for any $g \in \text{GL}(V)$, $1 \leq i \leq r-1$, $\sigma \in \mathfrak{S}_l$,

$$\Phi(\dot{g})\Xi_l(\mathbf{s}_i) = \Xi_l(\mathbf{s}_i)\Phi(\dot{g}) \text{ and } \Phi(\dot{g})\Xi_l(\mathbf{x}_\sigma) = \Xi_l(\mathbf{x}_\sigma)\Phi(\dot{g}).$$

Indeed,

$$\begin{aligned} \Phi(\dot{g})\Xi_l(\mathbf{s}_i)(\eta_{\mathbf{j}}) &= \Phi(\dot{g})(\eta_{j_1} \otimes \cdots \otimes \eta_{j_{i+1}} \otimes \eta_{j_i} \cdots \otimes \eta_{j_r}) \\ &= \Phi(\dot{g})(\eta_{j_1}) \otimes \cdots \otimes \Phi(\dot{g})(\eta_{j_{i+1}}) \otimes \Phi(\dot{g})(\eta_{j_i}) \cdots \otimes \Phi(\dot{g})(\eta_{j_r}) \\ &= \Xi_l(\mathbf{s}_i)\Phi(\dot{g})(\eta_{\mathbf{j}}). \end{aligned}$$

Hence, $\Phi(\dot{g})\Xi_l(\mathbf{s}_i) = \Xi_l(\mathbf{s}_i)\Phi(\dot{g})$.

Furthermore, for any $\mathbf{j} = (j_1, \dots, j_n) = (\mathbf{j}_l(n+1)^{r-l})$ with $\mathbf{j}_l = (j_1, \dots, j_l) \in \underline{n}^l$,

$$\begin{aligned} \Phi(\dot{g})\Xi_l(\mathbf{x}_\sigma)(\eta_{\mathbf{j}}) &= \Phi(\dot{g})(\eta_{j_{\sigma^{-1}(1)}} \otimes \cdots \otimes \eta_{j_{\sigma^{-1}(n)}}) \\ &= (\Phi(\dot{g})\eta_{j_{\sigma^{-1}(1)}}) \otimes \cdots \otimes (\Phi(\dot{g})\eta_{j_{\sigma^{-1}(n)}}) \\ &= \Xi_l(\mathbf{x}_\sigma)\Phi(\dot{g})(\eta_{\mathbf{j}}). \end{aligned}$$

And $\Phi(\dot{g})\Xi_l(\mathbf{x}_\sigma)(\eta_{\mathbf{k}}) = \Xi_l(\mathbf{x}_\sigma)\Phi(\dot{g})(\eta_{\mathbf{k}}) = 0$ for any $\mathbf{k} \in \mathcal{N}^r$ with $k_s = n+1$ for some $s \leq l$. This implies that $\Phi(\dot{g})\Xi_l(\mathbf{x}_\sigma) = \Xi_l(\mathbf{x}_\sigma)\Phi(\dot{g})$, as desired.

(4) The first part follows from the second one. We only need to prove the latter by different steps.

(i) First of all, by a direct check, $\Psi|_l(\mathbb{C}\mathfrak{S}_r)x_e^l$ is an associative algebra because $\Psi|_l(\tau).x_e^l = \Psi|_l(\tau)x_e^l\Psi|_l(\tau^{-1}) = x_e^l$ for $\tau \in \mathfrak{S}_r$. Correspondingly, it is a left module of $\Psi|_l(\mathbb{C}\mathfrak{S}_r)$.

(ii) By the above, it is not hard to see that $\Xi_l(\mathcal{H}_r^l)$ is spanned by $\Psi|_l(\tau)x_\sigma^l$ with $\tau \in \mathfrak{S}_r$, $\sigma \in \mathfrak{S}_l$ and $I \in \mathcal{I}$. Hence $\Xi_l(\mathcal{H}_r^l) = \Psi|_l(\mathbb{C}\mathfrak{S}_r)E_l$.

(iii) As to the algebra homomorphism, it can be directly verified.

(5) We need to show that Ξ keeps the relations (3.1)-(3.5). Since $\Psi|_{\mathfrak{S}_r}$ is a representation of \mathfrak{S}_r , Ξ keeps the relations (3.1)-(3.2). Moreover, note that $\Xi|_{\underline{V}_l^{\otimes r}} = \Xi_l$ for $0 \leq l \leq r$, we have

$$\Xi(\mathbf{x}_\sigma^l) \circ \Xi(\mathbf{x}_\mu^k) = \Xi_l(\mathbf{x}_\sigma^l)\Xi_k(\mathbf{x}_\mu^k) = \delta_{kl}\Xi_l(\mathbf{x}_{\sigma\circ\mu}^k) = \delta_{kl}\Xi(\mathbf{x}_\sigma^l\mathbf{x}_\mu^k) \text{ for } \sigma, \mu \in \mathfrak{S}_l, k, l \in \underline{r}. \quad (3.14)$$

$$\Xi(\mathbf{s}_i) \circ \Xi(\mathbf{x}_\sigma^l) = \Psi(s_i)\Xi_l(\mathbf{x}_\sigma^l) = \Xi_l(\mathbf{x}_{\mathbf{s}_i \circ \sigma}^l) = \Xi(\mathbf{s}_i\mathbf{x}_\sigma^l), \sigma \in \mathfrak{S}_l, i < l. \quad (3.15)$$

$$\Xi(\mathbf{x}_\sigma^l) \circ \Xi(\mathbf{s}_i) = \Xi_l(\mathbf{x}_\sigma^l)\Psi(s_i) = \Xi_l(\mathbf{x}_{\sigma \circ \mathbf{s}_i}^l) = \Xi(\mathbf{x}_\sigma^l\mathbf{s}_i), \sigma \in \mathfrak{S}_l, i < l. \quad (3.16)$$

$$\Xi(\mathbf{s}_i) \circ \Xi(\mathbf{x}_\sigma^l) = \Psi(s_i)\Xi_l(\mathbf{x}_\sigma^l) = \Xi_l(\mathbf{x}_\sigma^l) = \Xi(\mathbf{s}_i\mathbf{x}_\sigma^l) \text{ for } \sigma \in \mathfrak{S}_l, i > l. \quad (3.17)$$

$$\Xi(\mathbf{x}_\sigma^l) \circ \Xi(\mathbf{s}_i) = \Xi_l(\mathbf{x}_\sigma^l)\Psi(s_i) = \Xi_l(\mathbf{x}_\sigma^l) = \Xi(\mathbf{x}_\sigma^l\mathbf{s}_i) \text{ for } \sigma \in \mathfrak{S}_l, i > l. \quad (3.18)$$

So Ξ is an algebra homomorphism, thereby a representation of \mathcal{H}_r . \square

We will further investigate this representation in the next section.

3.5. Finite dimensional DDHAs. Keep the notations as above.

Definition 3.4. Set $D(n, r) := \Xi(\mathcal{H}_r)$, which we call the finite-dimensional degenerate double Hecke algebra of \mathfrak{S}_r (f.d. DHHA for short).

Set $D(n, r)_l := \Xi_l(\mathcal{H}_r^l)$ for $l = 0, 1, \dots, r$. We can extend the action of $D(n, r)_l$ on the whole of $\underline{V}^{\otimes r}$ as follows. Set

$$\Psi_l(\sigma) := \Psi|_{\underline{V}_l^{\otimes r}}(\sigma) \circ x_e^l.$$

Then Ψ_l defines a representation of \mathfrak{S}_r on $\underline{V}_l^{\otimes r}$. Now Ψ_l extends a representation of \mathfrak{S}_r on $\underline{V}^{\otimes r}$ by letting $\Psi_l(\mathfrak{S}_r)(\eta_{\mathbf{j}}) = 0$ for $\eta_{\mathbf{j}} \in \underline{V}_k^{\otimes r}$ with $k \neq l$. Similarly, each x_e^I extends to an element in $\text{End}_{\mathbb{C}}(\underline{V}^{\otimes r})$ with trivial action on $\underline{V}_k^{\otimes r}$ for $k \neq \#I$. So $D(n, r)_l$ annihilates $\underline{V}_k^{\otimes r}$ for $k \neq l$. Thus, each $D(n, r)_l$ becomes a subalgebra of \mathcal{A} , and two different such subalgebras are commutative.

Lemma 3.5. *Keep the notations as above. Then the following statements hold.*

(1) *The f.d. DDHA $D(n, r)$ is a direct sum of all $D(n, r)_l$ ($l = 0, \dots, r$), i.e.,*

$$D(n, r) = \bigoplus_{l=0}^r D(n, r)_l. \quad (3.19)$$

(2) *Set $d(n, l) := \sum_{\lambda \in P(l, n)} (\dim S_l^\lambda)^2$ with S_l^λ denoting the irreducible Specht module of \mathfrak{S}_l corresponding to $\lambda \in \text{Par}(l, n)$ (see §4.1 for the notation). Then $\dim D(n, r)_l = d(n, l) \binom{r}{l}^2$ and*

$$\dim D(n, r) = \sum_{l=0}^r d(n, l) \binom{r}{l}^2.$$

(3) *For $l \in \{0, 1, \dots, r\}$ there is a basis of E_l : $\{x_{\sigma_{l,i}}^I \mid I \in \mathcal{S}_l, \sigma_{l,i} \in \mathfrak{S}_l, i = 1, \dots, d(n, l)\}$ with $\sigma_{l,1} = \text{id}$, and $\binom{r}{l}^2$ elements $c_{J,I} \in \Psi|_l(\mathfrak{S}_r)$ such that $D(n, r)_l$ has a basis $\{c_{J,I} x_{\sigma_{l,i}}^I \mid (I, J) \in \mathcal{S}_l^2, i = 1, \dots, d(n, l)\}$. In particular, $x_e = \sum_{l=1}^r \sum_{I \in \mathcal{S}_l} x_e^I + x_\emptyset$ is just the identity of $D(n, r)$.*

(4) *The following decomposition of $D(n, r)$ into a direct sum of subspaces holds:*

$$D(n, r) = \mathbb{C}\Psi(\mathfrak{S}_r)x_e \oplus \bigoplus_{l=1}^r \bigoplus_{\substack{(J,I) \in \mathcal{S}_l^2 \\ i=1, \dots, d(n,l)}} \mathbb{C}c_{J,I}x_{\sigma_{l,i}}^I.$$

Proof. (1) By definition, $\bigoplus_{l=0}^r D(n, r)_l$ is really a direct sum in $\text{End}_{\mathbb{C}}(\underline{V}^{\otimes r})$. In the following, we prove (3.19). Let $\tau \in \mathfrak{S}_r$. Then for any $\mathbf{i} \in \mathcal{N}^r$, we have

$$\Psi(\tau)(\eta_{\mathbf{i}}) = \Psi|_s(\tau)(\eta_{\mathbf{i}}) = \sum_{l=0}^r \sum_{\substack{I \subset \underline{r} \\ \#I=l}} \Psi|_l(\tau)x_e^I(\eta_{\mathbf{i}}).$$

where $s = \text{rk}_{\underline{n}}(\mathbf{i})$. This implies that

$$\Psi(\tau) = \sum_{l=0}^r \sum_{\substack{I \subset \underline{r} \\ \#I=l}} \Psi|_l(\tau)x_e^I.$$

Hence,

$$D(n, r) \subset \bigoplus_{l=0}^r D(n, r)_l.$$

On the other hand, since

$$\Psi|_l(\sigma)(\eta_{\mathbf{i}}) = \delta_{l, \text{rk}_{\underline{n}}(\mathbf{i})} \Psi(\sigma)(\eta_{\mathbf{i}}) = \Psi(\sigma)x_e^l(\eta_{\mathbf{i}}), \forall \mathbf{i} \in \mathcal{N}^r,$$

it follows that

$$\Psi|_l(\sigma) = \Psi(\sigma)x_e^l.$$

This implies that

$$\bigoplus_{l=0}^r D(n, r)_l \subset D(n, r).$$

We complete the proof for (3.19).

(2) Due to (1), it suffices to prove the formula $\dim D(n, r)_l = d(n, l) \binom{r}{l}^2$. Firstly, for any given $I \in \mathcal{L}_l$, the subalgebra D_I generated by all x_σ^I ($\sigma \in \mathfrak{S}_l$) is isomorphic to $\mathbb{C}\Psi_l^V(\mathfrak{S}_l)$ which is of dimension equal to $d(n, l)$ by the representation theory of symmetric groups. So the dimensions of all those D_I are the same. Secondly, by Lemma 3.3(5) $D(n, r)_l$ is spanned by $\Psi(\tau)x_\sigma^I$ for all $\tau \in \mathfrak{S}_r$, $\sigma \in \mathfrak{S}_l$ and $I \subset \underline{r}$ with $\#I = l$. On the other hand, for any subsets $I = \{i_1 < i_2 < \cdots < i_l\}$, $J = \{j_1 < j_2 < \cdots < j_l\}$ of \underline{r} with $\#I = \#J = l$, take $\tau_{IJ} \in \mathfrak{S}_r$ such that $\tau_{IJ}(j_k) = i_k$ for $k = 1, \dots, l$. For any two τ_1 and $\tau_2 \in \mathfrak{S}_r$, if $\tau_1^{-1} \circ \tau_2$ stabilizes I , then $\Psi_l(\tau_1)x_\sigma^I = \Psi_l(\tau_2)x_{\sigma'}^I$ for some $\sigma' \in \mathfrak{S}_l$. Actually, for $\tau \in \mathfrak{S}_r$ satisfying $\tau|_I \in \text{Sym}(I)$ which is isomorphic to \mathfrak{S}_l , say by θ , we have $\Psi_l(\tau)x_\sigma^I = x_{\theta(\tau|_I) \circ \sigma}^I$ for any $\sigma \in \mathfrak{S}_l$, $I \subset \underline{r}$.

By the above arguments, we have that for any ordered pair (I, J) , the above τ_{IJ} changes a set of basis of D_J into the ones of D_I . There are $\binom{r}{l}^2$ such ordered pairs (I, J) . Summing up, along with the definition of D_I we have $\dim D(n, r)_l = d(n, l) \binom{r}{l}^2$, as desired.

(3) This statement follows from the arguments in (2).

(4) Note that the chosen basis elements for every E_I in the statement (3) include x_e^I . We make a proper adjustment of the basis such that the new basis contains x_e . Then the corresponding decomposition follows. \square

Remark 3.6. If $n > l$, then the following set

$$\{\Psi_l(\tau_{JI})x_\sigma^I \mid I \subset \underline{r}, J \subset \underline{r}, \#I = \#J = l, \sigma \in \mathfrak{S}_l\}$$

is a basis of $D(n, r)_l$. Moreover, the algebra structure of $D(n, r)_l$ is given by

$$(\Psi_l(\tau_{LK})x_\mu^K)(\Psi_l(\tau_{JI})x_\sigma^I) = \begin{cases} \Psi_l(\tau_{LI})x_{\mu \circ \sigma}^I, & \text{if } J = K; \\ 0, & \text{otherwise.} \end{cases} \quad (3.20)$$

Proof of Remark 3.6: Note that in this case, the elements x_σ^I for all $\sigma \in \mathfrak{S}_l$ and for any given I in the remark are linearly independent in D_I . The first statement follows from the arguments in the proof of Lemma 3.5(3). To show (3.20), we can assume that $I = \underline{l}$ without loss of generality. We note that the non-vanishing range of the operator $\Psi_l(\tau_{JI})x_\sigma^I$

is contained in $\underline{V}_J^{\otimes r}$ for $J \neq K$. Hence, $(\Psi_l(\tau_{LK})x_\mu^K)(\Psi_l(\tau_{JI})x_\sigma^I) = 0$. For the case $J = K$, we have

$$(\Psi_l(\tau_{LJ})x_\mu^J)(\Psi_l(\tau_{JI})x_\sigma^I)(\eta_{\mathbf{i}}) = \begin{cases} \Psi_l(\tau_{LI})\eta_{(\mu \circ \sigma)\mathbf{i}}, & \text{if } \mathbf{i} \in \mathcal{N}_l^r, i_k \in \underline{n}, \forall k \in I; \\ 0, & \text{otherwise.} \end{cases}$$

Hence, (3.20) follows.

4. DUALITY RELATED TO THE DEGENERATE DOUBLE HECKE ALGEBRA AND THE BRANCHING DUALITY FORMULA

In this section, with aid of some structural property of Levi and parabolic Schur algebras, we establish a duality between the degenerate double Hecke algebra \mathcal{H}_r and $\mathrm{GL}(n, \mathbb{C}) \times \mathbf{G}_m$ on the tensor space $(\mathbb{C}^{n+1})^{\otimes r}$. We will keep the notations as before.

4.1. Let us first recall the classical branching law for general linear groups. For positive integers m, r , set

$$\mathrm{Par}(r, m) = \{\mu = (\mu_1, \dots, \mu_m) \in \mathbb{N}^m \mid |\mu| = r, \mu_1 \geq \mu_2 \geq \dots \geq \mu_m\}.$$

An element from $\mathrm{Par}(r, m)$ is usually called a dominant weight, which is actually a partition of r into m parts (zero parts are allowed). For another weight $\lambda \in \mathrm{Par}(l, n)$ with $l \leq r$ and $n \leq m$, we call λ interlaces μ if $\mu_1 \geq \lambda_1 \geq \mu_2 \geq \lambda_2 \geq \dots \geq \mu_{m-1} \geq \lambda_{m-1} \geq \mu_m$ where $\lambda_{n+1} = \dots = \lambda_{m-1} = 0$. We denote $\lambda \lesssim \mu$ if λ interlaces μ .

For a given $\mu \in \mathrm{Par}(r, n+1)$, let L_{n+1}^μ be an irreducible $\mathrm{GL}(n+1, \mathbb{C})$ -module with highest weight μ . Then as a module over $\mathrm{GL}(n, \mathbb{C})$, there is a unique decomposition

$$L_{n+1}^\mu = \bigoplus_{l=0}^r \bigoplus_{\lambda \in \mathrm{Par}(l, n)} \delta_{\lambda \lesssim \mu} L_n^\lambda$$

with

$$\delta_{\lambda \lesssim \mu} = \begin{cases} 1, & \text{if } \lambda \lesssim \mu; \\ 0, & \text{otherwise.} \end{cases}$$

There is a similar branching law for symmetric groups (see (4.8)).

4.2. In the following, we give the decomposition of $\underline{V}_l^{\otimes r}$ as a \mathcal{H}_r^l -module into a direct sum of irreducible modules. For this, denote by L_n^λ the irreducible $\mathrm{GL}(n, \mathbb{C})$ -module with highest weight λ for $\lambda \in \mathrm{Par}(l, n)$, and denote by S_l^λ the irreducible Specht module over \mathfrak{S}_l corresponding to λ .

Recall that $\mathrm{GL}(n, \mathbb{C})$ has a standard Borel subgroup consisting of all upper triangular invertible matrices, denoted by \mathfrak{B} . Denote by \mathfrak{N} the unipotent radical of \mathfrak{B} which consists of all unipotent upper triangular matrices.

Lemma 4.1. *Keep the notations as above. As a \mathcal{H}_r^l -module, $\underline{V}_l^{\otimes r}$ decomposes into a direct sum of irreducible modules as follows*

$$\bigoplus_{\lambda \in \mathrm{Par}(l, n)} (D_l^\lambda)^{\oplus \dim L_n^\lambda} \tag{4.1}$$

where as a $\mathbb{C}\mathfrak{S}_l$ -module,

$$D_l^\lambda \cong \underbrace{S_l^\lambda \oplus S_l^\lambda \oplus \cdots \oplus S_l^\lambda}_{\binom{r}{l} \text{ times}}$$

Proof. For any given $J = \{j_1, j_2, \dots, j_l\} \subset \underline{r}$, we have a subspace $\underline{V}_J^{\otimes r}$ in $\underline{V}_l^{\otimes r}$, which is spanned by $\eta_{\mathbf{i}} = \eta_{i_1} \otimes \cdots \otimes \eta_{i_r}$ with $i_{j_1}, \dots, i_{j_l} \in \underline{n}$. Then

$$\underline{V}_l^{\otimes r} = \bigoplus_{\substack{J \subset \underline{r} \\ \#J=l}} \underline{V}_J^{\otimes r} \quad (4.2)$$

where in the sum, J ranges over all subsets of \underline{r} consisting of l -elements. The number of such J is exactly $\binom{r}{l}$. For a fixed J , $\underline{V}_J^{\otimes r}$ admits an \mathfrak{S}_l -action, which just permutes the position indicated by J , this is to say, for any $\sigma_l \in \mathfrak{S}_l$, $\sigma_l \cdot \eta_{\mathbf{i}} = \eta_{\sigma_l \cdot \mathbf{i}}$ if defining $\sigma_l \cdot \mathbf{i} = \delta \cdot ((\sigma_l \cdot (\mathbf{i}'_l))(n+1)^{r-l})$ for $\mathbf{i} = \delta \cdot (\mathbf{i}'_l(n+1)^{r-l})$. On the other hand, $\underline{V}_J^{\otimes r}$ becomes a $\text{GL}(V)$ -module with every factor η fixed.

Thus $\underline{V}_J^{\otimes r}$ can be regarded as an r -tensor space of V with $(\text{GL}_n \times \mathfrak{S}_l)$ -action. Thanks to the classical Schur-Weyl duality, as a $(\text{GL}_n \times \mathfrak{S}_l)$ -module we have the following decomposition

$$\underline{V}_J^{\otimes r} \cong \bigoplus_{\lambda \in \text{Par}(l, n)} L_n^\lambda \otimes S_l^\lambda. \quad (4.3)$$

Let us show the meaning of S_l^λ in the above decomposition by demonstrating the standard representatives in the isomorphism class of S_l^λ . For the simplicity of arguments, we might as well suppose $J = \underline{l}$ which is equal to $\{1, 2, \dots, l\}$ without any loss of generality. Take $\lambda = (\lambda_1, \lambda_2, \dots, \lambda_n) \in \text{Par}(l, n)$. Set

$$\eta(\lambda) = \underbrace{\eta_1 \otimes \cdots \otimes \eta_1}_{\lambda_1 \text{ factors}} \otimes \underbrace{\eta_2 \otimes \cdots \otimes \eta_2}_{\lambda_2 \text{ factors}} \otimes \cdots \otimes \underbrace{\eta_n \otimes \cdots \otimes \eta_n}_{\lambda_n \text{ factors}} \otimes \eta \otimes \cdots \otimes \eta.$$

Consider the orbit $\mathfrak{S}_l \cdot \eta(\lambda)$. The space spanned by this orbit forms an \mathfrak{S}_l -module, which is denoted by $I_{\underline{l}}^\lambda$. Then $I_{\underline{l}}^\lambda$ is actually the λ -weighted space of $\text{GL}(n, \mathbb{C})$ -module $\underline{V}_{\underline{l}}^{\otimes r}$. In particular, $I_{\underline{l}}^\lambda$ contains all maximal λ -weighted vectors in $\text{GL}(n, \mathbb{C})$ -module $\underline{V}_{\underline{l}}^{\otimes r}$, all of which by definition, are \mathfrak{N} -invariants in $I_{\underline{l}}^\lambda$. Furthermore, the invariant space in $I_{\underline{l}}^\lambda$ under the action of \mathfrak{N} is exactly the unique direct summand isomorphic to S_l^λ , in the complete reducible decomposition of $I_{\underline{l}}^\lambda$ as \mathfrak{S}_l -module (see [10, §9.1.1]). This summand is denoted by $S_{\underline{l}}^\lambda$. More generally, each $\underline{V}_I^{\otimes r}$ for $I \subset \underline{r}$ (with $\#I = l$) admits the corresponding I_I^λ which contains a unique direct summand S_I^λ paralleling to $S_{\underline{l}}^\lambda$. This means, there exists $\sigma \in \mathfrak{S}_r$ such that the conjugation by $\Psi|_l(\sigma)$ maps $S_{\underline{l}}^\lambda$ onto S_I^λ . Obviously,

$$D_l^\lambda := \bigoplus_{\substack{I \subset \underline{r} \\ \#I=l}} S_I^\lambda$$

is a $\Psi|_l(\mathfrak{S}_r)$ -module, which is as an \mathfrak{S}_l -module, isomorphic to the direct sum of $\binom{r}{l}$ copies of $S_{\underline{l}}^\lambda$.

Next we prove that D_l^λ is an irreducible module over \mathcal{H}_r^l . For this, we only need to show that for any given nonzero vector $w \in D_l^\lambda$, the cyclic \mathcal{H}_r^l -submodule W generated by w must coincide with D_l^λ itself. Now, we write $w = \sum_{\text{some } I} w_I$ with some nonzero $w_I \in \underline{V}_I^{\otimes r}$. Fix one

w_I . Recall \mathcal{H}_r^l has a subalgebra E_I by abuse of the notation (see §3.4), which is isomorphic to $\mathbb{C}\Psi_l^V(\mathfrak{S}_l)$. So the cyclic module $E_I w_I$ of E_I coincides with the irreducible $\mathbb{C}\mathfrak{S}_l$ -module S_l^λ . This implies that there is some x_σ^I for $\sigma \in \mathfrak{S}_l$ such that $x_\sigma^I w_I \neq 0$. On the other hand, by definition (see §3.4) $x_\sigma^I \sum_{J \neq I} w_J = 0$. Therefore, $x_\sigma^I w = x_\sigma^I w_I$ is a nonzero vector in S_l^λ . Hence the submodule W contains an irreducible E_I -module S_l^λ . By $\mathbb{C}\mathfrak{S}_r$ -action, we finally have that the \mathcal{H}_r^l -submodule W coincides with $\bigoplus_{\substack{J \subset [r] \\ \#J=l}} S_J^\lambda$, which is equal to D_l^λ .

According to the previous analysis, D_l^λ is really an irreducible \mathcal{H}_r^l -module. From the above arguments, along with (4.2) and (4.3), the lemma follows. \square

4.3. Let us first notice that $\underline{V}^{\otimes r}$ naturally becomes a representation space of $\mathcal{E}(n, r)$ (and of $S(n+1, r)$ more generally) by defining

$$\xi(\eta_{\mathbf{i}}) = \sum_{\mathbf{k} \in \mathcal{N}^r} \xi(c_{\mathbf{k}, \mathbf{i}}) \eta_{\mathbf{k}} \quad (4.4)$$

for any $\xi \in \mathcal{E}(n, r)$, and any basis elements $\eta_{\mathbf{i}} = \eta_{i_1} \otimes \cdots \otimes \eta_{i_r} \in$ of $\underline{V}^{\otimes r}$ ($\mathbf{i} \in \mathcal{N}^r$). This representation is denoted by Υ . We are actually considering $\Upsilon : \mathcal{E}(n, r) \rightarrow \text{End}_{\mathbb{C}}(\underline{V}^{\otimes r})$.

As before, let $\Phi : \text{GL}(n+1) \rightarrow \mathcal{A} := \text{End}_{\mathbb{C}}(\underline{V}^{\otimes r})$ be the natural representation. By the classical Schur algebra theory, we can identify the image of Φ with $S(n+1, r)$, in aid of Υ .

Consequently, we have the following corollary to Lemma 4.1.

Corollary 4.2. *Keep the notations as above. As a \mathcal{H}_r -module, $\underline{V}^{\otimes r}$ decomposes into a direct sum of irreducible modules as follows*

$$\bigoplus_{l=0}^r \bigoplus_{\lambda \in \text{Par}(l, n)} (D_l^\lambda)^{\oplus \dim L_l^\lambda}. \quad (4.5)$$

4.4. We have the following Levi version of Schur-Weyl duality from $\text{GL}(n+1, \mathbb{C})$ to $\text{GL}(n, \mathbb{C}) \times \mathbf{G}_{\mathbf{m}}$.

Theorem 4.3. *(Levi Schur-Weyl duality) Keep the above notations. The following statements hold.*

- (1) $\text{End}_{D(n, r)}(\underline{V}^{\otimes r}) = \mathbb{C}\Phi(\text{GL}(V) \times \mathbf{G}_{\mathbf{m}})$.
- (2) *The following duality holds:*

$$\begin{aligned} \text{End}_{\mathcal{L}(n, r)}(\underline{V}^{\otimes r}) &= D(n, r); \\ \text{End}_{D(n, r)}(\underline{V}^{\otimes r}) &= \mathcal{L}(n, r). \end{aligned} \quad (4.6)$$

Proof. (1) Recall that $\text{GL}(V) \times \mathbf{G}_{\mathbf{m}}$ can be regarded as a closed subgroup of $\text{GL}(\underline{V})$ (see §2.5.4). By Lemma 3.3(3) (along with some direct computation), $\mathbb{C}\Phi(\text{GL}(V) \times \mathbf{G}_{\mathbf{m}}) \subset \text{End}_{D(n, r)}(\underline{V}^{\otimes r})$. We need to show the opposite inclusion.

Note that $D(n, r) = \Xi(\mathcal{H}_r)$. So $\text{End}_{D(n, r)}(\underline{V}^{\otimes r}) \subset \text{End}_{\mathbb{C}\mathfrak{S}_r}(\underline{V}^{\otimes r}) = \mathbb{C}\Phi(\text{GL}(\underline{V}))$. The last equation is due to the classical Schur-Weyl duality for $\text{GL}(\underline{V})$ and \mathfrak{S}_r . Thus, for any $\phi \in \text{End}_{D(n, r)}(\underline{V}^{\otimes r})$, we can write $\phi = \sum_{h \in \text{GL}(\underline{V})} a_h \Phi(h)$ for $a_h \in \mathbb{C}$. In the following, we will show that ϕ must lie in $\mathbb{C}\Phi(\text{GL}(V) \times \mathbf{G}_{\mathbf{m}})$.

We identify $\text{GL}(\underline{V})$ with $\text{GL}(n+1, \mathbb{C})$, and $\Phi(\text{GL}(\underline{V}))$ with $S(n+1, r)$ by the classical theory of Schur algebras. Thus, we rewrite $\phi = \sum_{(\mathbf{i}, \mathbf{j}) \in \mathcal{N}^r \times \mathcal{N}^r / \mathfrak{S}_r} a_{\mathbf{ij}} \xi_{\mathbf{i}, \mathbf{j}}$ for $a_{\mathbf{ij}} \in \mathbb{C}$ satisfying

$a_{\mathbf{ij}} = a_{\tau, \mathbf{i}, \tau, \mathbf{j}}$ for any $\tau \in \mathfrak{S}_r$. Furthermore, $\phi = \sum_{l=0}^r \phi_l$ with

$$\phi_l = \sum_{\substack{(\mathbf{i}, \mathbf{j}) \in \mathcal{N}^r \times \mathcal{N}^r / \sim \\ \text{rk}_{\mathbb{N}} \mathbf{j} = l}} a_{\mathbf{ij}} \xi_{\mathbf{i}, \mathbf{j}}.$$

In this view, by Lemma 2.8 we only need to prove that all ϕ_l lie in the Levi subalgebra $\mathcal{L}(n, r)$. The arguments proceed in steps.

(1-1) By the assumption $\phi \in \text{End}_{D(n, r)}(\underline{V}^{\otimes r})$, $\phi \circ x_\sigma^l = x_\sigma^l \circ \phi$ for all $\sigma \in \mathfrak{S}_l$, and $l = 0, 1, \dots, r$. We first claim that ϕ stabilise $\underline{V}_l^{\otimes r}$ for every l .

Actually, if not so, then there must be a basis element $\eta_{\mathbf{j}} \in \underline{V}_l^{\otimes r}$ not obeying the stability claimed above for ϕ . From (4.4) it follows $\phi(\underline{V}_l^{\otimes r}) = \phi_l(\underline{V}_l^{\otimes r})$. Violation of the stability implies that $\phi_l(\eta_{\mathbf{j}}) = \eta_l + \eta_k$ with $\eta_l \in \underline{V}_l^{\otimes r}$ but $\eta_k \notin \underline{V}_l^{\otimes r}$. Suppose $\mathbf{j} = \tau(\mathbf{j}'(n+1)^{r-l})$ for $\tau \in \mathfrak{S}_r$, and $\mathbf{j}' \in \underline{n}^l$. Then $(\tau x_{\text{id}}^l \tau^{-1})(\eta_{\mathbf{j}}) = \eta_{\mathbf{j}}$. On the other side, $(\tau x_{\text{id}}^l \tau^{-1}) \circ \phi(\eta_{\mathbf{j}}) = \phi \circ (\tau x_{\text{id}}^l \tau^{-1})(\eta_{\mathbf{j}})$. This leads to a contradiction that $\tau x_{\text{id}}^l \tau^{-1}(\eta_l + \eta_k) = \eta_l + \eta_k$. So the claim is true.

(1-2) By (1-1), ϕ_l is commutative with x_σ^l for all $\sigma \in \mathfrak{S}_l$. From this, by the same arguments as in (1-1) it follows that ϕ_l stabilises $\underline{V}_l^{\otimes r}$. Hence ϕ_l is actually commutative with \mathfrak{S}_l in $\text{End}_{\mathbb{C}}(\underline{V}_l^{\otimes r})$.

Recall that $\underline{V}_l^{\otimes r} = V^{\otimes l} \times \eta^{\otimes r-l}$, and $x_\sigma^l = \Psi_l^V(\sigma) \otimes \text{id}^{\otimes l-r}$. By applying the classical Schur Weyl duality, we have that $\phi|_{\underline{V}_l^{\otimes r}} \in \Phi(\text{GL}(V)|_{\underline{V}_l^{\otimes r}} = \Phi(\text{GL}(V) \times \mathbf{G}_{\mathbf{m}})|_{\underline{V}_l^{\otimes r}})$. Note that $\phi|_{\underline{V}_l^{\otimes r}} = \phi_l|_{\underline{V}_l^{\otimes r}}$. From this, it is deduced that $\phi_l|_{\underline{V}_l^{\otimes r}} \in \Phi(\text{GL}(V) \times \mathbf{G}_{\mathbf{m}})|_{\underline{V}_l^{\otimes r}}$. By definition, we can further deduce that $\phi_l|_{\underline{V}_l^{\otimes r}} \in \Phi(\text{GL}(V) \times \mathbf{G}_{\mathbf{m}})|_{\underline{V}_l^{\otimes r}}$. Keep in mind $\phi_l|_{\underline{V}_k^{\otimes r}} = 0$ for $k \neq l$ in the same sense as in §4.3. By definition and Lemma 2.8(2), it follows that $\phi_l \in \mathcal{L}(n, r)_l \subset \mathcal{L}(n, r)$.

By the analysis in the beginning, we accomplish the proof of the first statement.

(2) The second equation in (4.6) follows from (1). Note that $\underline{V}^{\otimes r}$ is completely reducible over $D(n, r)$ (Corollary 4.2). The first equation in (4.6) follows from the classical double commutant theorem (see [10, §4.1.5]). \square

As a consequence of Theorem 4.3, we have the following result on classification of irreducible $D(n, r)$ -modules.

Corollary 4.4. *The set consisting of D_l^λ for $\lambda \in \text{Par}(l, n)$ and $l = 0, 1, \dots, r$ form a representative set of the isomorphism class of irreducible $D(n, r)$ -modules.*

Proof. Since $\mathcal{L}(n, r)$ is semisimple, so is $D(n, r)$ by Theorem 4.3 and the Double Commutant Theorem. Furthermore, each $D(n, r)_l$ is semisimple for $l = 0, 1, \dots, r$ by Lemma 3.5(1). Thanks to Lemma 4.1, D_l^λ is an irreducible $D(n, r)_l$ -module for any $\lambda \in \text{Par}(l, n)$, and

$D_l^\lambda \not\cong D_l^\mu$ for $\lambda \neq \mu$. Moreover,

$$\begin{aligned} \sum_{\lambda \in \text{Par}(l,n)} (\dim D_l^\lambda)^2 &= \sum_{\lambda \in \text{Par}(l,n)} \binom{r}{l}^2 (\dim S_l^\lambda)^2 \\ &= \binom{r}{l}^2 \sum_{\lambda \in \text{Par}(l,n)} (\dim S_l^\lambda)^2 \\ &= \binom{r}{l}^2 d(n,l) \\ &= \dim D(n,r)_l \quad (\text{Lemma 3.5(4)}). \end{aligned}$$

Hence, the set $\{D_l^\lambda \mid \lambda \in \text{Par}(l,n)\}$ exhausts all representatives of isomorphism classes of irreducible $D(n,r)_l$ -modules. Consequently, the desired assertion holds. \square

4.5. Branching duality formula. From Theorem 4.3 we derive a dimension relation between irreducible modules from $(\text{GL}_n, \mathfrak{S}_r)$ -duality, and irreducible modules from $(\text{GL}_{n+1}, \mathfrak{S}_r)$ -duality.

Corollary 4.5. *The following statements hold.*

(1) *Under the action of $\text{GL}(n, \mathbb{C}) \times \mathcal{H}_r$, the space of r -tensors over \mathbb{C}^{n+1} decomposes as*

$$(\mathbb{C}^{n+1})^{\otimes r} \cong \bigoplus_{l=0}^r \bigoplus_{\lambda \in \text{Par}(l,n)} L_n^\lambda \otimes D_l^\lambda.$$

(2) *(Branching duality formula) In the Schur-Weyl duality, the irreducible pairs (L_{n+1}^μ, S_r^μ) ($\mu \in \text{Par}(r, n+1)$) for $(\text{GL}_{n+1}, \mathfrak{S}_r)$ and the irreducible pairs $(L_n^\lambda, S_l^\lambda)$ ($\lambda \in \text{Par}(l, n)$) for $(\text{GL}_n, \mathfrak{S}_l)$ with $l = 1, \dots, r$ satisfy the following branching duality formula*

$$\sum_{\mu \in \text{Par}(r, n+1)} \delta_{\lambda \lesssim \mu} \dim S_r^\mu = \binom{r}{l} \dim S_l^\lambda \quad (4.7)$$

for any $\lambda \in \text{Par}(l, n)$ ($l \leq r$).

Proof. (1) Note that all $c \in \mathbf{G}_m$ act on $\underline{V}_l^{\otimes r}$ by a scalar for any l . Keep Lemma 4.1 in mind. So the decomposition follows from Theorem 4.3 and the classical double commutant theorem (see [10, §4.16]).

(2) It follows from (1) that as a $\text{GL}(n, \mathbb{C})$ -module

$$(\mathbb{C}^{n+1})^{\otimes r} \cong \bigoplus_{l=0}^r \bigoplus_{\lambda \in \text{Par}(l,n)} (L_n^\lambda)^{\oplus \binom{r}{l} \dim S_l^\lambda} = \sum_{l=0}^r \sum_{\lambda \in \text{Par}(l,n)} \left(\binom{r}{l} \dim S_l^\lambda \right) L_n^\lambda.$$

On the other hand, according to the classical Schur-Weyl duality, $(\mathbb{C}^{n+1})^{\otimes r}$ has the following decomposition, as a $\mathrm{GL}(n+1, \mathbb{C})$ -module:

$$\begin{aligned} (\mathbb{C}^{n+1})^{\otimes r} &\cong \bigoplus_{\mu \in \mathrm{Par}(r, n+1)} (L_{n+1}^\mu)^{\oplus \dim S_r^\mu} \\ &= \sum_{\mu \in \mathrm{Par}(r, n+1)} \sum_{\lambda \in \mathrm{Par}(l, n)} (\delta_{\lambda \lesssim \mu} \dim S_r^\mu) L_n^\lambda \\ &= \sum_{l=0}^r \sum_{\lambda \in \mathrm{Par}(l, n)} \sum_{\mu \in \mathrm{Par}(r, n+1)} (\delta_{\lambda \lesssim \mu} \dim S_r^\mu) L_n^\lambda \end{aligned}$$

Compare both decompositions. The equation (4.7) follows. \square

Remark 4.6. (1) When $l = r$, the formula (4.7) becomes trivial.

(2) When $l = r - 1$, the formula (4.7) becomes

$$\sum_{\mu \in \mathrm{Par}(r, n+1)} \delta_{\lambda \lesssim \mu} \dim S_r^\mu = r \dim S_{r-1}^\lambda$$

for $\lambda \in \mathrm{Par}(r-1, n)$. This formula can be regarded as a duality to the following classical branching rule (see [10, Corollary 9.2.7])

$$\mathrm{Res}_{\mathfrak{S}_{r-1}}^{\mathfrak{S}_r} S_r^\mu \cong \bigoplus_{\lambda \in \mathrm{Par}(r-1, n)} \delta_{\lambda \lesssim \mu} S_{r-1}^\lambda. \quad (4.8)$$

5. ENHANCED TENSOR INVARIANTS AND PARABOLIC SCHUR-WEYL DUALITY

Keep the notations as before. In this section, we give the answer to Question 2.3, then establish the parabolic Schur-Weyl dualities and give some applications.

5.1. Recall the notations in §2.5.3. In particular, \underline{V} has a basis $\{\eta_1, \dots, \eta_n; \eta_{n+1}\}$ with $\eta_i \in V$ for $i = 1, \dots, n$; $\eta_{n+1} = \eta$. Then $\underline{V}^{\otimes r}$ has a basis $\{\eta_{\mathbf{i}} = \eta_{i_1} \otimes \eta_{i_2} \otimes \dots \otimes \eta_{i_r} \mid \mathbf{i} = (i_1, i_2, \dots, i_r) \in \mathcal{N}^r\}$. Let us recall that $\underline{V}^{\otimes r}$ naturally becomes a representation space of $\mathcal{E}(n, r)$ (and of $S(n+1, r)$ more generally) by definition as in (4.4).

We first have the following basic observation.

Lemma 5.1. *Denote by $\mathcal{P}(n, r)_{l,0}$ the subspace of $\mathcal{P}(n, r)$ spanned by $\xi_{\mathbf{i}_{(n+1)^{r-l}, (n+1)^r}}$ with \mathbf{i}_l ranging over \underline{n}^l , and by $\mathcal{P}(n, r)_{\geq 0,0}$ the direct sum of all $\mathcal{P}(n, r)_{l,0}$ with $l = 0, 1, \dots, r$. Then the image of $\mathcal{P}(n, r)_{\geq 0,0}$ is exactly $(\underline{V}^{\otimes r})^{\mathfrak{S}_r}$.*

Proof. Recall that for any $l = 0, 1, \dots, r$, $\underline{V}_l^{\otimes r}$ has a basis consisting of $\eta_{\mathbf{i}} \in \underline{V}_l^{\otimes r}$ with \mathbf{i} ranging over \mathcal{N}_l^r . Note that $\mathbf{i} \sim \mathbf{i}_{l(n+1)^{r-l}}$ with $\mathbf{i}_l \in \underline{n}^l$. By definition, $\xi_{\mathbf{i}_{l(n+1)^{r-l}, (n+1)^r}}(\eta^{\otimes r})$ is a fundamental invariant in $(\underline{V}_l^{\otimes r})^{\mathfrak{S}_r}$ arising from the \mathfrak{S}_r -orbit of $\eta_{\mathbf{i}}$ (see (4.4)). Correspondingly, the image of $\mathcal{P}(n, r)_{l,0}$ coincides with $(\underline{V}_l^{\otimes r})^{\mathfrak{S}_r}$. When l ranges over $\{0, 1, \dots, r\}$, the lemma follows. \square

As a result, we have the following crucial lemma.

Lemma 5.2. *For any $\phi \in \mathrm{End}_{\mathcal{P}(n, r)}(\underline{V}^{\otimes r})$, ϕ must lie in $\mathbb{C}\Psi(\mathfrak{S}_r) + D(n, r)_{\mathrm{ann}}$ where $D(n, r)_{\mathrm{ann}} := \{\phi \in D(n, r) \mid \phi((\underline{V}^{\otimes r})^{\mathfrak{S}_r}) = 0\}$.*

Proof. Note that $\text{End}_{\mathcal{P}(n,r)}(\underline{V}^{\otimes r}) \subset \text{End}_{\mathcal{P}(n,r)}(\underline{V}^{\otimes r})$. From Theorem 4.3(2), $\text{End}_{\mathcal{P}(n,r)}(\underline{V}^{\otimes r}) \subset D(n,r)$. In the following, we will show that $\text{End}_{\mathcal{P}(n,r)}(\underline{V}^{\otimes r}) \cap D(n,r) \subset \mathbb{C}\Psi(\mathfrak{S}_r) + D(n,r)_{\text{ann}}$. By Lemma 3.5(4),

$$D(n,r) = \mathbb{C}\Psi(\mathfrak{S}_r)x_e \oplus \left(\bigoplus_{l=1}^r \bigoplus_{\substack{(J,I) \in \mathcal{S}_l^2 \\ i=1, \dots, d(n,l)}} \mathbb{C}c_{J,I}x_{\sigma_{l,i}}^I \right).$$

Thanks to Proposition 2.5, we only need to show that

$$\text{End}_{\mathcal{P}(n,r)}(\underline{V}^{\otimes r}) \cap \left(\bigoplus_{l=1}^r \bigoplus_{\substack{(J,I) \in \mathcal{S}_l^2 \\ i=1, \dots, d(n,l)}} \mathbb{C}c_{J,I}x_{\sigma_{l,i}}^I \right) \subset D(n,r)_{\text{ann}}. \quad (5.1)$$

Actually, for any given nonzero $\phi = \sum_{l=t}^s \phi_l$ with $0 < t \leq s \leq r$, and $\phi_l \in D(n,r)_l$, $l = t, t+1, \dots, s$, such that $\phi \in \text{End}_{\mathcal{P}(n,r)}(\underline{V}^{\otimes r})$ and both ϕ_s, ϕ_t are not zero. By the assumption, for any $\xi \in \mathcal{P}(n,r)_{\geq 0,0}$, we have $\xi \circ \phi = \phi \circ \xi$. Then on one side $\xi \circ \phi$ vanishes because $\phi = \sum_{l=t}^s \phi_l$ with $t > 0$. Hence, $\phi \circ \xi$ must be zero for all $\xi \in \mathcal{P}(n,r)_{\geq 0,0}$. Thanks to Lemma 5.1, the image of $\mathcal{P}(n,r)_{\geq 0,0}$ is exactly $(\underline{V}^{\otimes r})^{\mathfrak{S}_r}$. This means that ϕ annihilates the space $(\underline{V}^{\otimes r})^{\mathfrak{S}_r}$.

Summing up, (5.1) is proved. The lemma follows. \square

Generally, we have the following theorem.

Theorem 5.3. (*Parabolic Schur-Weyl duality*) *Keep the notations as above, in particular $\underline{G} = \underline{\text{GL}}(\underline{V})$. Set $D(n,r)^V = \{\phi \in D(n,r) \mid e^v \circ \phi \circ e^{-v} = \phi, \forall v \in V\}$. Then the following statements hold.*

- (1) $\text{End}_{\mathbb{C}\Phi(\underline{G} \rtimes \mathbf{G}_m)}(\underline{V}^{\otimes r}) = D(n,r)^V$.
- (2) *The above $D(n,r)^V$ can be described as*

$$\mathbb{C}\Psi(\mathfrak{S}_r) \subset D(n,r)^V \subset \mathbb{C}\Psi(\mathfrak{S}_r) + D(n,r)_{\text{ann}}.$$

Proof. (1) Keep in mind $\mathcal{P}(n,r) = \mathbb{C}\Phi(\underline{G} \rtimes \mathbf{G}_m)$, and $\underline{G} = G \times_{\nu} V$. Then the statement follows from Theorem 4.3.

(2) The first inclusion follows from Lemma 2.5(1). The second one is due to Lemma 5.2. \square

5.2. Example. Let us demonstrate the above theorem by an example. Keep the notations as before, and let $I = \{1, 2\}$, $I_i = \{i\}$ for $i = 1, 2$, and $\sigma \in \mathfrak{S}_2$ interchanging 1 and 2. Then we have the following demonstration of $D(n,r)^V$.

- (1) If $r = 1$ or 2, and $n \geq r$, then $D(n,r)^V = \mathbb{C}\Psi(\mathfrak{S}_r)$.
- (2) If $r = 2$ and $n = 1$, then $D(n,r)^V = \mathbb{C}\Psi(\mathfrak{S}_r) \oplus \mathbb{C}(x_e^I - x_{\sigma}^I)$.

Proof of the demonstration. Recall that \underline{V} has a basis $\eta_1, \dots, \eta_n; \eta_{n+1} := \eta$, where η_1, \dots, η_n forms a basis of V .

(1) In the case $r = 1$, it is obvious that $D(n,r)_{\text{ann}} = 0$. The assertion is obvious.

Now we suppose that $n \geq r = 2$. Then $D(n,r)$ has the following basis

$$\{x_e^I, x_{\sigma}^I, \Psi(\sigma)x_e^{I_1}, \Psi(\sigma)x_e^{I_2}, x_e^{I_1}, x_e^{I_2}, x_e^{\emptyset}\}.$$

Take any $\phi \in D(n, r)^V$, by the discussion as in Lemma 5.2, we can assume that

$$\phi = a_1 x_e^I + a_2 x_\sigma^I + a_3 \Psi(\sigma) x_e^{I_1} + a_4 \Psi(\sigma) x_e^{I_2} + a_5 x_e^{I_1} + a_6 x_e^{I_2} + a_7 x_e^\emptyset, \quad (5.2)$$

where $a_i \in \mathbb{C}$ for $1 \leq i \leq 7$. By the assumption, $\phi \circ \Phi(e^m) = \Phi(e^m) \circ \phi$. In particular, on one hand,

$$\begin{aligned} & \phi \circ \Phi(e^m)(\eta \otimes \eta) \\ &= \phi((\eta_1 + \eta) \otimes (\eta_1 + \eta)) \\ &= (a_1 + a_2)\eta_1 \otimes \eta_1 + (a_3 + a_6)\eta \otimes \eta_1 + (a_4 + a_5)\eta_1 \otimes \eta + a_7\eta \otimes \eta, \end{aligned} \quad (5.3)$$

and on the other hand,

$$\begin{aligned} & \Phi(e^m) \circ \phi(\eta \otimes \eta) \\ &= \Phi(e^m)(a_7\eta \otimes \eta) \\ &= a_7(\eta_1 + \eta) \otimes (\eta_1 + \eta) \\ &= a_7\eta_1 \otimes \eta_1 + a_7\eta \otimes \eta_1 + a_7\eta_1 \otimes \eta + a_7\eta \otimes \eta. \end{aligned} \quad (5.4)$$

By comparing both sides of (5.3) and (5.4), we have

$$a_1 + a_2 = a_3 + a_6 = a_4 + a_5 = a_7. \quad (5.5)$$

Moreover, on one hand,

$$\begin{aligned} & \phi \circ \Phi(e^m)(\eta_2 \otimes \eta) \\ &= \phi(\eta_2 \otimes (\eta_1 + \eta)) \\ &= a_1\eta_2 \otimes \eta_1 + a_2\eta_1 \otimes \eta_2 + a_3\eta \otimes \eta_2 + a_5\eta_2 \otimes \eta. \end{aligned} \quad (5.6)$$

On the other hand,

$$\begin{aligned} & \Phi(e^m) \circ \phi(\eta_2 \otimes \eta) \\ &= \Phi(e^m)(a_3\eta \otimes \eta_2 + a_5\eta_2 \otimes \eta) \\ &= a_3\eta_1 \otimes \eta_2 + a_3\eta \otimes \eta_2 + a_5\eta_2 \otimes \eta_1 + a_5\eta_2 \otimes \eta. \end{aligned} \quad (5.7)$$

By comparing both sides of (5.6) and (5.7), we have

$$a_1 = a_5, a_2 = a_3. \quad (5.8)$$

It follows from (5.5) and (5.8) that

$$a_1 = a_5 = a_6 := a, a_2 = a_3 = a_4 = b, a_7 = a + b.$$

Consequently,

$$\phi = a(x_e^I + x_e^{I_1} + x_e^{I_2} + x_e^\emptyset) + b(x_\sigma^I + \Psi(\sigma)x_e^{I_1} + \Psi(\sigma)x_e^{I_2} + x_e^\emptyset) = a\Psi(e) + b\Psi(\sigma) \in \mathbb{C}\Psi(\mathfrak{S}_r).$$

The assertion follows.

(2) Suppose $r = 2$ and $n = 1$. Take any $\phi \in D(n, r)^V$. As the arguments in (1), we can write ϕ as the form (5.2). By the assumption, $\phi \circ \Phi(e^m) = \Phi(e^m) \circ \phi$. In particular, on one

hand,

$$\begin{aligned}
& \phi \circ \Phi(e^n)(\eta_1 \otimes \eta) \\
&= \phi(\eta_1 \otimes (\eta_1 + \eta)) \\
&= (a_1 + a_2)\eta_1 \otimes \eta_1 + a_3\eta \otimes \eta_1 + a_5\eta_1 \otimes \eta,
\end{aligned} \tag{5.9}$$

and one the other hand,

$$\begin{aligned}
& \Phi(e^n) \circ \phi(\eta_1 \otimes \eta) \\
&= \Phi(e^n)(a_3\eta \otimes \eta_1 + a_5\eta_1 \otimes \eta) \\
&= (a_3 + a_5)\eta_1 \otimes \eta_1 + a_3\eta \otimes \eta_1 + a_5\eta_1 \otimes \eta.
\end{aligned} \tag{5.10}$$

By comparing both sides of (5.9) and (5.10), we have

$$a_1 + a_2 = a_3 + a_5. \tag{5.11}$$

Then it follow from (5.5) and (5.11) that

$$a_3 = a_4 := c, a_5 = a_6 := d, a_7 = a_1 + a_2 = c + d.$$

Consequently,

$$\begin{aligned}
\phi &= d(x_e^I + x_e^{I_1} + x_e^{I_2} + x_e^\emptyset) + c(x_\sigma^I + \Psi(\sigma)x_e^{I_1} + \Psi(\sigma)x_e^{I_2} + x_e^\emptyset) + (a_1 - c)x_e^I + (a_2 - d)x_\sigma^I \\
&= d\Psi(e) + c\Psi(\sigma) + (a_1 - c)(x_e^I - x_\sigma^I) \in \mathbb{C}\Psi(\mathfrak{S}_r) \oplus \mathbb{C}(x_e^I - x_\sigma^I).
\end{aligned}$$

Moreover, it is a routine to check that $x_e^I - x_\sigma^I \in D(n, r)^V$, and the assertion follows.

5.3. A conjecture. We propose the following conjecture.

Conjecture 5.4. When $n \geq r$, $D(n, r)^V$ coincides with $\mathbb{C}\Psi(\mathfrak{S}_r)$.

5.4. Enhanced tensor invariants. Identify $\underline{V}^{*\otimes r}$ with $(\underline{V}^{\otimes r})^*$. Then there is a natural $\underline{G} \rtimes \mathbf{G}_m$ -equivariant isomorphism of vector spaces

$$T : \underline{V}^{\otimes r} \otimes \underline{V}^{*\otimes r} \rightarrow \text{End}_{\mathbb{C}}(\underline{V}^{\otimes r}).$$

With the parabolic Schur-Weyl duality, we describe the enhanced tensor invariants.

Still let η_1, \dots, η_n be a basis of V , and let $\eta_{n+1} = \eta$. Then $\{\eta_i \mid i = 1, \dots, n+1\}$ constitute a basis of \underline{V} . Let $\eta_1^*, \dots, \eta_{n+1}^*$ be the dual basis of \underline{V}^* . For a multi-index $\mathbf{i} = (i_1, \dots, i_r) \in \mathbb{N}^r$. Still set $\eta_{\mathbf{i}} = \eta_{i_1} \otimes \dots \otimes \eta_{i_r}$, and $\eta_{\mathbf{i}}^* = \eta_{i_1}^* \otimes \dots \otimes \eta_{i_r}^*$. Then $\eta_{\mathbf{i}}$, and $\eta_{\mathbf{i}}^*$ form basis of $\underline{V}^{\otimes r}$ and $\underline{V}^{*\otimes r}$ respectively, when \mathbf{i} ranges over \mathbb{N}^r . Furthermore, the set $\{\eta_{\mathbf{i}} \otimes \eta_{\mathbf{j}}^* \mid (\mathbf{i}, \mathbf{j}) \in \mathbb{N}^r \times \mathbb{N}^r\}$ is a basis of $\underline{V}^{\otimes r} \otimes \underline{V}^{*\otimes r}$.

For $\sigma \in \mathfrak{S}_r$, define a mixed tensor C_σ by

$$C_\sigma = \sum_{\mathbf{i} \in \mathbb{N}^r} \eta_{\sigma, \mathbf{i}} \otimes \eta_{\mathbf{i}}^*.$$

Proposition 5.5. Let $\underline{G} = \underline{\text{GL}}(V)$. When $D(n, r)^V = \mathbb{C}\Psi(\mathfrak{S}_r)$, then the space of $\underline{G} \rtimes \mathbf{G}_m$ -invariants in the mixed-tensor product $\underline{V}^{\otimes r} \otimes \underline{V}^{*\otimes r}$ is generated by these C_σ with $\sigma \in \mathfrak{S}_r$.

Proof. By the definition of T mentioned before, for $\sigma \in \mathfrak{S}_r$ we have

$$T(C_\sigma)\eta_j = \sum_{i \in \mathbb{N}^r} \eta_i^*(\eta_j)\eta_{\sigma.i} = \eta_{\sigma.j}.$$

This means that $T(C_\sigma) = \Psi(\sigma)$. By the assumption, the $\underline{G} \rtimes \mathbf{G}_m$ -invariants in $\underline{V}^{\otimes r} \otimes \underline{V}^{*\otimes r}$ is isomorphic to $\sum_{\sigma \in \mathfrak{S}_r} \mathbb{C}\Psi(\sigma)$ via T . Hence, the proposition is proved. \square

6. REPRESENTATIONS OF THE PARABOLIC SCHUR ALGEBRA

Keep the notations as before. In this section, we proceed to study representations of the parabolic Schur algebra $\mathcal{P}(n, r)$. In the whole section, the term ‘‘a module of $\mathcal{P}(n, r)$ ’’ always means a right module of $\mathcal{P}(n, r)$.

6.1. Irreducible modules and PIMs. Recall $\mathcal{P}(n, r)$ has a sequence of right ideals of $\mathcal{P}(n, r)$:

$$\mathcal{P}(n, r) = \mathcal{P}(n, r)_0 \supset \mathcal{P}(n, r)_1 \supset \mathcal{P}(n, r)_2 \supset \cdots \supset \mathcal{P}(n, r)_r \supset \mathcal{P}(n, r)_{r+1}.$$

with quotients $\mathcal{P}(n, r)_i / \mathcal{P}(n, r)_{i+1} \cong S(n, i) \rtimes \mathfrak{a}_i$ (Lemma 2.10(3)). Those right ideals satisfy $\mathcal{P}(n, r)_i \mathcal{P}(n, r) = \mathcal{P}(n, r)_i$. Naturally, $S(n, r)$ naturally becomes a $\mathcal{P}(n, r)$ -module.

Consider the primitive idempotent decomposition of the identity element id in $\mathcal{P}(n, r)$

$$\text{id} = \sum_{\mathbf{s} \in \mathbb{N}_r^n} \xi_{\tilde{\pi}_{\mathbf{s}}, \tilde{\pi}_{\mathbf{s}}}. \quad (6.1)$$

We have

$$\mathcal{P}(n, r) = \sum_{\mathbf{s} \in \mathbb{N}_r^n} \xi_{\tilde{\pi}_{\mathbf{s}}, \tilde{\pi}_{\mathbf{s}}} \mathcal{P}(n, r). \quad (6.2)$$

Then $P_{\tilde{\pi}_{\mathbf{s}}} := \xi_{\tilde{\pi}_{\mathbf{s}}, \tilde{\pi}_{\mathbf{s}}} \mathcal{P}(n, r)$ is an indecomposable projective module (PIM² for short) of $\mathcal{P}(n, r)$. Furthermore, for $0 \leq l \leq r$ we set

$$\mathcal{P}(n, r)_{[l]} := \mathbb{C}\text{-span}\{\xi_{\tilde{\pi}_{\mathbf{s}}, \tilde{\pi}_{\mathbf{s}}} \in E \mid |\mathbf{s}| = l\}.$$

Then by (S1)-(S3) again,

$$\mathcal{P}(n, r)_{[l]} = \sum_{\substack{\mathbf{s} \in \mathbb{N}_r^n \\ |\mathbf{s}| = l}} P_{\tilde{\pi}_{\mathbf{s}}}, \quad (6.3)$$

6.1.1. Let us diverge by recalling some general results on irreducible modules for the classical Schur algebras. First of all, the Schur algebra $S(m, r)$ over \mathbb{C} is semisimple. For any $\pi \in \underline{m}^r$, π lies in some \mathfrak{S}_r -orbit from \underline{m}^r , say $\lambda = (\lambda_1, \dots, \lambda_m) \in \Lambda(m, r)$. The vector λ can also be regarded as a unordered partition of r into m parts, where zero parts being allowed. So the identity element $\text{id}_{S(m, r)}$ of $S(m, r)$ has the following primitive idempotent decomposition $\text{id}_{S(m, r)} = \sum_{\lambda \in \Lambda(m, r)} \xi_{\lambda, \lambda}$. Consequently, the right ideal generated by $\xi_{\lambda, \lambda}$ forms an irreducible module over $S(m, r)$, denoted by \tilde{D}_λ .

Consider the Weyl group \mathcal{W} of GL_m which is exactly \mathfrak{S}_m , and the action of \mathcal{W} on \underline{m}^r . The action is on the left, $w\pi = (w(\pi_1), \dots, w(\pi_r))$ for $w \in \mathcal{W}$ and $\pi = (\pi_1, \dots, \pi_r) \in \underline{m}^r$. This action commutes with that of \mathfrak{S}_r . Consequently, it is well-defined that \mathfrak{S}_m acts on $\Lambda(m, r)$

²short for Principal Indecomposable Modules.

as $w\lambda = (\lambda_{w^{-1}(1)}, \dots, \lambda_{w^{-1}(m)})$ for $w \in \mathscr{W}$ and $\lambda \in \Lambda(m, r)$. Each \mathscr{W} -orbit of $\Lambda(m, r)$ contains only one dominant weight $\gamma = (\gamma_1, \dots, \gamma_m)$ which satisfies $\gamma_1 \geq \gamma_2 \geq \dots \geq \gamma_m$. Denote by $\Lambda^+(m, r)$ the set of all dominant weights. Then $\Lambda^+(m, r)$ exactly consists of all partition of r admitting at most number m of nonzero parts, coinciding with $\text{Par}(r, m)$.

Lemma 6.1. *Suppose ν is a dominant weight, and shares the same \mathfrak{S}_m -orbit with λ . Then two $S(m, r)$ -irreducible modules \tilde{D}_λ and \tilde{D}_ν are isomorphic.*

Proof. Recall that $S(m, r) = \Phi(\text{GL}(m, \mathbb{C}))$ with $\text{GL}(m, \mathbb{C}) = \text{GL}(W)$ associated to a given basis $\eta_i, i = 1, \dots, m$ of $W \cong \mathbb{C}^m$. Suppose $\gamma = w(\lambda)$ for $w \in \mathscr{W}$. Note that w gives rise to an automorphism of $\text{GL}(W)$ by permutation via w on the basis $\eta_i, i = 1, \dots, m$. Naturally, w gives rise to an automorphism of $S(m, r)$ which interchanges two $S(m, r)$ -irreducible modules \tilde{D}_λ and \tilde{D}_γ . Hence both of them are isomorphic, as $S(m, r)$ -modules. \square

Thus the isomorphism classes of irreducible $S(m, r)$ -modules are parameterized by $\Lambda^+(m, r)$. For $\lambda \in \Lambda(m, r)$, the representative of the corresponding isomorphism classes of irreducibles is denoted by \tilde{D}_γ where $\gamma = \mathscr{W}(\lambda) \cap \Lambda^+(m, r)$. Denote by $\ell(\gamma)$ the number of different elements in the \mathscr{W} -orbit of γ . Then there is a direct sum decomposition of irreducible modules for $S(m, r)$:

$$S(m, r) \cong \sum_{\gamma \in \Lambda^+(m, r)} \tilde{D}_\gamma^{\oplus \ell(\gamma)}. \quad (6.4)$$

6.1.2. Let us turn back to the parabolic Schur algebra $\mathscr{P}(n, r)$. For a given $\mathbf{s} \in \mathbb{N}_r^n$ with $|\mathbf{s}| = l \leq r$, $\tilde{\pi}_{\mathbf{s}} = (1^{s_1} \dots n^{s_n} (n+1)^{r-l})$. By (S1)-(S3), $P_{\tilde{\pi}_{\mathbf{s}}} = \xi_{\tilde{\pi}_{\mathbf{s}}, \tilde{\pi}_{\mathbf{s}}} \mathscr{P}(n, r)_l$ and $\xi_{\tilde{\pi}_{\mathbf{s}}, \tilde{\pi}_{\mathbf{s}}} \mathscr{P}(n, r)_{l+1} = 0$. So $P_{\tilde{\pi}_{\mathbf{s}}}$ can be decomposed into a direct sum of two subspaces

$$P'_{\tilde{\pi}_{\mathbf{s}}} = \mathbb{C}\text{-span}\{\xi_{\tilde{\pi}_{\mathbf{s}}, \mathbf{p}_l} | \mathbf{p}_l \in \underline{n}^l\} \quad (6.5)$$

and

$$Q_{\tilde{\pi}_{\mathbf{s}}} = \mathbb{C}\text{-span}\{\xi_{\tilde{\pi}_{\mathbf{s}}, \mathbf{q}_l} | \mathbf{q}_l \in \mathcal{N}^l, \text{rk}_{\underline{n}}(\mathbf{q}_l) < l\}. \quad (6.6)$$

Furthermore $Q_{\tilde{\pi}_{\mathbf{s}}} = P_{\tilde{\pi}_{\mathbf{s}}} \mathscr{P}(n, r)_{l+1}$, and it becomes a $\mathscr{P}(n, r)$ -submodule of $P_{\tilde{\pi}_{\mathbf{s}}}$. Moreover, $Q_{\tilde{\pi}_{\mathbf{s}}}$ is the unique maximal submodule of $P_{\tilde{\pi}_{\mathbf{s}}}$, and the quotient $P_{\tilde{\pi}_{\mathbf{s}}}/Q_{\tilde{\pi}_{\mathbf{s}}} \cong \tilde{D}_{\tilde{\pi}_{\mathbf{s}}}$ as $S(n, l)$ -modules. By the arguments above, the irreducible $S(n, l)$ -module $\tilde{D}_{\tilde{\pi}_{\mathbf{s}}}$ is naturally an $\mathscr{P}(n, r)$ -module. Still denote by \mathscr{W} the Weyl group of GL_n . By the same arguments as in §6.1.1, $w \in \mathscr{W}$ gives rise to an automorphism of $\mathscr{P}(n, r)$. As $\mathscr{P}(n, r)$ -modules, $P_{\tilde{\pi}_{\mathbf{s}}} \cong P_\gamma$, and $\tilde{D}_{\tilde{\pi}_{\mathbf{s}}} \cong D_\gamma$ for some $\gamma \in \Lambda^+(n, l)$. Moreover,

$$\bigoplus_{\substack{l=0 \\ l \neq |\mathbf{s}|}}^r \mathscr{P}(n, r)_{[l]} \subseteq \text{Ann}_{\mathscr{P}(n, r)}(D_\gamma) \subsetneq \mathscr{P}(n, r).$$

This implies that $D_\gamma \not\cong D_{\gamma'}$ and $P_\gamma \not\cong P_{\gamma'}$ if $\gamma \in \Lambda^+(n, l)$ and $\gamma' \in \Lambda^+(n, l')$ with $l \neq l'$. Furthermore, by the classical theory of Schur algebras, $D_v \not\cong D_{v'}$ and $P_v \not\cong P_{v'}$ if $v, v' \in \Lambda^+(n, l)$ and $v \neq v'$. Summing up, we have the following classification results of irreducible $\mathscr{P}(n, r)$ -modules and of indecomposable projective $\mathscr{P}(n, r)$ -modules.

Theorem 6.2. *The set $\Lambda := \{\gamma \in \Lambda^+(n, l) \mid l = 0, \dots, r\}$ parameterize both of the isomorphism classes of irreducible $\mathcal{P}(n, r)$ -modules and of indecomposable projective $\mathcal{P}(n, r)$ -modules. The corresponding irreducible (resp. indecomposable projective) modules are D_γ (resp. P_γ) with $\gamma \in \Lambda$.*

6.2. Cartan invariants and block degeneracy for $\mathcal{P}(n, r)$. By the above arguments, the P_γ 's with $\gamma \in \bigcup_{l=0}^r \Lambda^+(n, l)$ form all PIM's for $\mathcal{P}(n, r)$, up to isomorphisms. In this subsection, we will precisely determine the Cartan invariants for the parabolic Schur algebra $\mathcal{P}(n, r)$. For that, we need the following preliminary result which describes the decomposition factors appearing in $Q_{\tilde{\pi}_{\mathbf{s}}}$ defined in (6.6) and their multiplicity.

Lemma 6.3. *Keep the notations as before. Then for any given $\mathbf{s} \in \mathbb{N}_r^n$ with $1 \leq |\mathbf{s}| = l \leq r$, the following equality holds in the Grothendieck group for the $\mathcal{P}(n, r)$ -module category.*

$$[Q_{\tilde{\pi}_{\mathbf{s}}}] = \sum_{i=0}^{l-1} \sum_{\gamma \in \Lambda^+(n, i)} \ell(\gamma)[D_\gamma] = \sum_{i=0}^{l-1} [S(n, i)], \quad (6.7)$$

where $\ell(\gamma)$ is defined as before, the number of different elements in the \mathcal{W} -orbit of γ .

Proof. The second equation of (6.7) is derived from (6.4). We only need to prove the first one.

Recall

$$Q_{\tilde{\pi}_{\mathbf{s}}} = \mathbb{C}\text{-span}\{\xi_{\tilde{\pi}_{\mathbf{s}}, \mathbf{q}_l(n+1)^{r-l}} \mid \mathbf{q}_l \in \mathcal{N}^l, \text{rk}_{\underline{n}}(\mathbf{q}_l) < l\}.$$

Set

$$Q_{\tilde{\pi}_{\mathbf{s}}}^{(i)} = \mathbb{C}\text{-span}\{\xi_{\tilde{\pi}_{\mathbf{s}}, \mathbf{q}_l(n+1)^{r-l}} \mid \mathbf{q}_l \in \mathcal{N}^l, \text{rk}_{\underline{n}}(\mathbf{q}_l) \leq l - i\} \text{ for } 1 \leq i \leq l.$$

By (S1)-(S3), each $Q_{\tilde{\pi}_{\mathbf{s}}}^{(i)}$ is a $\mathcal{P}(n, r)$ -submodule of $Q_{\tilde{\pi}_{\mathbf{s}}}$, and we have the following decreasing sequence of $\mathcal{P}(n, r)$ -modules:

$$Q_{\tilde{\pi}_{\mathbf{s}}} = Q_{\tilde{\pi}_{\mathbf{s}}}^{(1)} \supset Q_{\tilde{\pi}_{\mathbf{s}}}^{(2)} \supset \dots \supset Q_{\tilde{\pi}_{\mathbf{s}}}^{(l)} \supset 0.$$

Furthermore,

$$\bigoplus_{\substack{j=0 \\ j \neq l-i}}^r \mathcal{P}(n, r)_{[j]}$$

acts trivially on $Q_{\tilde{\pi}_{\mathbf{s}}}^{(i)}/Q_{\tilde{\pi}_{\mathbf{s}}}^{(i+1)}$ and $Q_{\tilde{\pi}_{\mathbf{s}}}^{(i)}/Q_{\tilde{\pi}_{\mathbf{s}}}^{(i+1)} \cong S(n, l-i)$ as $S(n, l-i)$ -modules. Consequently, $Q_{\tilde{\pi}_{\mathbf{s}}}^{(i)}/Q_{\tilde{\pi}_{\mathbf{s}}}^{(i+1)} \cong S(n, l-i)$ as $\mathcal{P}(n, l-i)$ -modules because $S(n, l-i)$ is a $\mathcal{P}(n, r)$ -module. Furthermore, by (6.4) we have

$$[Q_{\tilde{\pi}_{\mathbf{s}}}^{(i)}/Q_{\tilde{\pi}_{\mathbf{s}}}^{(i+1)}] = \sum_{\gamma \in \Lambda^+(n, l-i)} \ell(\gamma)[D_\gamma].$$

Consequently,

$$\begin{aligned}
[Q_{\tilde{\pi}_s}] &= \sum_{i=1}^l [Q_{\tilde{\pi}_s}^{(i)} / Q_{\tilde{\pi}_s}^{(i+1)}] \\
&= \sum_{i=1}^l \sum_{\gamma \in \Lambda^+(n, l-i)} \ell(\gamma) [D_\gamma] \\
&= \sum_{i=0}^{l-1} \sum_{\gamma \in \Lambda^+(n, i)} \ell(\gamma) [D_\gamma].
\end{aligned}$$

The proof is completed. \square

As usual, we denote by $(P_\gamma : D_{\gamma'})$ the multiplicity of the irreducible module $D_{\gamma'}$ appearing in composition series of the indecomposable projective module P_γ for $\gamma, \gamma' \in \bigcup_{l=0}^r \Lambda^+(n, l)$. We are now in the position to determine the Cartan invariants for the parabolic Schur algebra $\mathcal{P}(n, r)$.

Theorem 6.4. *Let $a_{\gamma, \gamma'} := (P_\gamma : D_{\gamma'})$ for any $\gamma, \gamma' \in \Lambda^+ := \bigcup_{l=0}^r \Lambda^+(n, l)$. Then*

$$a_{\gamma, \gamma'} = \begin{cases} 1, & \text{if } \gamma' = \gamma; \\ \ell(\gamma'), & \text{if } |\gamma'| < |\gamma|; \\ 0, & \text{otherwise.} \end{cases} \quad (6.8)$$

Consequently, the Cartan matrix $(a_{\gamma, \gamma'})_{\mathfrak{m} \times \mathfrak{m}}$ is an invertible and upper triangular one with diagonal entries being 1, where $\mathfrak{m} = \#\Lambda^+$.

Proof. If $\gamma \in \Lambda^+(n, 0)$, i.e., $\gamma = \mathbf{0} = (0, \dots, 0)$, then $P_{\mathbf{0}} = D_{\mathbf{0}}$ is irreducible, and the assertion (6.8) holds for $\gamma = \mathbf{0}$ and any $\gamma' \in \Lambda$. In the following, we assume $\gamma \in \Lambda^+(n, l)$ with $1 \leq l \leq r$. By the discussion in §6.1.2, P_γ has a unique maximal submodule Q_γ with $P_\gamma / Q_\gamma \cong D_\gamma$. Hence, it follows from Lemma 6.3 that

$$[P_\gamma] = [P_\gamma / Q_\gamma] + [Q_\gamma] = [D_\gamma] + \sum_{i=0}^{l-1} \sum_{v \in \Lambda^+(n, i)} \ell(\gamma) [D_v].$$

This implies the assertion (6.8). The other assertions are obvious. The proof is completed. \square

As a direct consequence of Theorem 6.4, we have the following result on block structure of the parabolic Schur algebra.

Corollary 6.5. *The parabolic Schur algebra $\mathcal{P}(n, r)$ has only one block.*

The block structure is closely related to the center for a finite-dimensional algebra (see [3, §1.8]). As to the latter, we have the following observation.

Proposition 6.6. *When $D(n, r)^V = \mathbb{C}\Psi(\mathfrak{S}_r)$, the center of $\mathcal{P}(n, r)$ is one-dimensional.*

Proof. Denote by \mathbb{C} the center of $\mathcal{P}(n, r)$. Note that $\mathcal{P}(n, r) \subset \text{End}_{\mathbb{C}}(\underline{V}^{\otimes r})$. We have

$$\mathbb{C} \subset \text{Cent}_{\mathcal{P}(n, r)}(\text{End}_{\mathbb{C}}(\underline{V}^{\otimes r})).$$

Hence $\mathbf{C} = \text{End}_{\mathcal{P}(n,r)}(\underline{V}^{\otimes r}) \cap \mathcal{P}(n,r)$. When $D(n,r)^V = \mathbb{C}\Psi(\mathfrak{S}_r)$, $\mathbf{C} = \mathbb{C}\Psi(\mathfrak{S}_r) \cap \mathcal{P}(n,r)$ by Theorem 5.3. Recall $\mathcal{P}(n,r)$ has a basis $\{\xi_{\tilde{\pi}_s, \mathbf{j}} \mid (\tilde{\pi}_s, \mathbf{j}) \in E\}$ (Theorem 2.10). For any given nonzero $\phi \in \mathbf{C}$, ϕ can be written in two forms

$$\phi = \sum_{(\tilde{\pi}_s, \mathbf{j}) \in E} a_{\tilde{\pi}_s, \mathbf{j}} \xi_{\tilde{\pi}_s, \mathbf{j}} \quad (6.9)$$

and

$$\phi = \sum_{\delta \in \mathfrak{S}_r} b_\delta \Psi(\delta). \quad (6.10)$$

We refine (6.10) as $\phi = \sum_{\delta \in S} b_\delta \Psi(\delta)$ where $S \subset \mathfrak{S}_r$ and all $b_\delta \neq 0$ for $\delta \in S$. Then we finally write (6.10) as

$$\begin{aligned} \phi &= \sum_{\delta \in S} b_\delta \sum_{\mathbf{s} \in \mathbb{N}_r^n} \xi_{\tilde{\pi}_s, \delta, \tilde{\pi}_s} \\ &= \sum_{\mathbf{s} \in \mathbb{N}_r^n} \sum_{\delta \in S} b_\delta \xi_{\tilde{\pi}_s, \delta, \tilde{\pi}_s}. \end{aligned}$$

If $S \neq \{\text{id}\}$, then we can take $\delta (\neq \text{id}) \in S$. In this case, there certainly exists $\mathbf{s} \in \mathbb{N}_r^n$ such that $(\tilde{\pi}_s, \delta(\tilde{\pi}_s)) \notin E$. This contradicts with (6.9).

So it is only possible that S coincides with $\{\text{id}\}$. Consequently, $\phi \in \mathbb{C}\text{id}$. Hence $\mathbf{C} = \mathbb{C}\text{id}$. The proof is completed. \square

ACKNOWLEDGEMENT

We are grateful to Chenliang Xue for his pointing out some wrong arguments and statements in an old version of the manuscript.

REFERENCES

- [1] P. Achar and A. Henderson, *Orbit closures in the enhanced nilpotent cone*, Adv. Math. 219 (2008), 27-62.
- [2] Arakawa and T. Suzuki, *Duality between $\mathfrak{sl}_n(\mathbb{C})$ and the degenerate affine Hecke algebra*, J. Algebra 209 (1998), 288-304.
- [3] D. J. Benson, *Representations and cohomology I: Basic representation theory of finite groups and associative algebras*, Cambridge University Press 1995.
- [4] B. Deng, J. Du, B. Parshall and J. Wang, *Finite dimensional algebras and quantum groups*, Mathematical Surveys and Monographs 150, American Mathematical Society, Providence, RI, 2008.
- [5] V. Drinfeld, *Degenerate affine Hecke algebras and Yangians*, Func. Anal. Appl. 20 (1986), 56-58.
- [6] I. Cherednik, *Double affine Hecke algebras*, LMS Lecture Notes Series 31, Cambridge University Press, 2005.
- [7] F. D. Grosshans, *The invariants of unipotent radicals of parabolic subgroups*, Invent. Math. 73 (1983), no. 1, 1-9.
- [8] F. D. Grosshans, *Hilbert's fourteenth problem for nonreductive groups*, Math. Z. 193 (1986), no. 1, 95-103.
- [9] F. D. Grosshans, *Algebraic homogeneous spaces and invariant theory*, Lecture Notes in Mathematics, 1673. Springer-Verlag, Berlin, 1997.
- [10] R. Goodman and N. R. Wallach, *Symmetry, representations, and invariants*, GTM 255, Springer, 2009.
- [11] J. A. Green, *Polynomial representations of GL_n* , LNM 830, Springer-Verlag, New York, 1980.
- [12] J. C. Jantzen, *Support varieties of Weyl modules*, Bull. London Math. Soc. 19 (1987), no. 3, 238-244.
- [13] B. Liu and D. Wang, *Enhanced Brauer algebras and related dualities*, preprint.

- [14] A. Mathas, *Iwahori-Hecke algebras and Schur algebras of the symmetric group*, ULS 15, American Math. Soc., Providence, R. I., 1999.
- [15] K. Ou, B. Shu and Y. Yao, *Chevalley restriction theorem for semi-reductive algebraic groups and its applications*, preprint (2019).
- [16] A. A. Premet and H. Strade, *Classification of finite dimensional simple Lie algebras in prime characteristics*, Representations of algebraic groups, quantum groups, and Lie algebras, 185-214, Contemp. Math., 413, Amer. Math. Soc., Providence, RI, 2006.
- [17] B. Shu, Y. Xue and Y. Yao, *Enhanced reductive groups (II): Nilpotent orbits*, preprint (2020).
- [18] H. Strade and R. Farnsteiner, *Modular Lie algebras and their representations*, Marcel Dekker, New York, 1988.
- [19] R. Wilson, *Automorphisms of graded Lie algebras of Cartan type*, Comm. Algebra 3(7), 591-613, 1975.
- [20] C. Xue, *Double Hecke algebras and related quantum Schur-Weyl dualities*, preprint.

SCHOOL OF MATHEMATICAL SCIENCES, EAST CHINA NORMAL UNIVERSITY, SHANGHAI, 200241, CHINA.
Email address: `bshu@math.ecnu.edu.cn`

SCHOOL OF MATHEMATICAL SCIENCES, EAST CHINA NORMAL UNIVERSITY, SHANGHAI, 200241, CHINA.
Email address: `1647227538@qq.com`

DEPARTMENT OF MATHEMATICS, SHANGHAI MARITIME UNIVERSITY, SHANGHAI, 201306, CHINA.
Email address: `yfyao@shmtu.edu.cn`