Optimal Anticodes, Diameter Perfect Codes, Chains and Weights

Luciano Panek and Navene Michele Paião Panek *†‡

Abstract

Let P be a partial order on $[n] = \{1, 2, ..., n\}$, \mathbb{F}_q^n be the linear space of n-tuples over a finite field \mathbb{F}_q and w be a weight on \mathbb{F}_q . In this paper, we consider metrics on \mathbb{F}_q^n induced by chain orders P over [n] and weights w over \mathbb{F}_q , and we determine the cardinality of all optimal anticodes and completely classify them. Moreover, we determine all diameter perfect codes for a set of relevant instances on the aforementioned metric spaces.

Key words: poset metric, pomset metric, NRT metric, perfect code, MDS code, anticode, diameter perfect code.

1 Introduction

Classically, coding theory takes place in linear spaces over finite fields or modules over rings, endowed with a metric, e.g., the linear space \mathbb{F}_q^n of all *n*-tuples over a finite field \mathbb{F}_q endowed with the Hamming metric or the module \mathbb{Z}_m^n of all *n*-tuples over a ring \mathbb{Z}_m of integers modulo m endowed with the Lee metric.

In a given metric space, codes that attain the sphere-packing bound are called perfect. A possible general setting for the existence problem of perfect codes is the class of distance regular graphs, introduced by Biggs (see [2]), including the nearly ubiquitous Hamming metric spaces (also called Hamming graphs).

The Johnson graphs and the Grassmann graphs are other examples of distance regular graphs (see [3]). For the Hamming graphs over \mathbb{F}_q , there are no nontrivial perfect codes except for the codes having the parameters of the Hamming codes and the two Golay codes. Martin and Zhu [9] showed that there are no nontrivial perfect codes in the Grassmann graphs. Determining all perfect codes is an open problem

^{*}Luciano Panek is with the Center of Exact Sciences and Engineering, State University of West Paraná, Foz do Iguaçu, Paraná, Brazil (e-mail: luciano.panek@unioeste.br).

[†]Nayene Michele Paião Panek is with the Center of Exact Sciences and Engineering, State University of West Paraná, Foz do Iguaçu, Paraná, Brazil (e-mail: nayene.panek@unioeste.br).

[‡]Copyright (c) 2021 IEEE. Personal use of this material is permitted. However, permission to use this material for any other purposes must be obtained from the IEEE by sending a request to pubs-permissions@ieee.org.

for Johnson graphs. It was conjectured by Delsarte in the 1970's that there are no nontrivial perfect codes in Johnson graphs (see [5]). See [7] and the references therein for progress towards proving the conjecture of Delsarte.

In his pioneer work [5], Delsarte also proved the following result:

Theorem 1 (Delsarte) Let $\Gamma = (V, E)$ be a distance regular graph. Let X and Y be subsets of V such that the nonzero distances occurring between vertices of X do not occur between vertices of Y. Then

$$|X| \cdot |Y| < |V|$$
.

Ahlswede, Aydinian, and Khachatrian in [1] gave the definition of diameter perfect code. They examined a variant of Theorem 1. Let $\Gamma = (V, E)$ be a distance regular graph. A subset A of V is called an *anticode* with diameter δ if δ is the maximum graph distance occurring between vertices of A. Anticodes with diameter δ having maximal size are called *optimal anticodes*. Let diam(A) be the diameter of an anticode A in Γ . Now let

$$A^*(D) \triangleq \max\{|A| : diam(A) \le D\}.$$

Theorem 2 Let $\Gamma = (V, E)$ be a distance regular graph. If C is a code in Γ with minimum distance D+1, then

$$A^*(D) \cdot |C| \le |V|. \tag{1}$$

Ahlswede $et\ al.$ continued with the following new definition. A code C with minimum distance D+1 is called $diameter\ perfect$ if equality holds in equation (1). This is a generalization of the usual definition of e-perfect code as balls of radius e are anticodes with diameter 2e.

In Hamming graphs, in addition to the Hamming and Golay codes, the extended Hamming and extended Golay codes are diameter perfect, as well as all MDS codes. In the Johnson graphs, no nontrivial e-perfect codes are known but all Steiner systems are diameter perfect codes. Nontrivial diameter perfect codes are also known in the Grassmann graphs. For more details, see [1].

Another possible general setting for the existence problem of perfect and diameter perfect codes is the class of the metric spaces introduced by Panek and Pinheiro in [11], the so-called weighted poset metric spaces, that include any additive metric space (e.g., the Hamming and Lee metric spaces), as well as the poset metric spaces [4] and the pomset metric spaces [14]. The class of these spaces is distinct from the class of the distance regular graph metric spaces and, thus, Delsarte's Theorem, does not apply. Fortunately, an alternative proof that was given in [6] can be slightly modified to work on this case (Theorem 5).

The poset metric spaces were introduced by Brualdi, Graves, and Lawrence in [4]. These metrics cast new light into many of the classical invariants of coding theory (such as minimum distance, packing and covering radius) and many of their basic results (concerning perfect and MDS codes, MacWilliams' identity, syndrome

decoding and so on) with several works published over the years, contributing to a better understanding of these invariants and their properties when considering the classical Hamming metric. As a comprehensive survey, we cite the book of Firer *et al.* [8].

For the distance regular graphs, the balls of radius e are anticodes with diameter larger than e and are optimal anticodes with diameter equal to 2e if C is a perfect code. For the weighted poset metric spaces, the diameter of the balls of radius e can be equal to e. The weighted poset metric is a mix of two extremal cases: the Hamming metric (determined by an anti-chain order and the Hamming weight on coordinates) with the Niederreiter-Rosenbloom-Tsfasman metric, introduced by Niederreiter in [10], and Rosenbloom and Tsfasman in [13] (determined by a chain order and the Hamming weight on coordinates; an ultrametric). For the Hamming metric, the diameter of a ball of radius e is greater than e. For the Niederreiter-Rosenbloom-Tsfasman metric the diameter of a ball of radius e is exactly equal to e.

In this work let us consider the particular setting of weighted poset metrics where the poset is a chain order (Section 3). We will show that the diameter of a ball with radius e is equal to e for all e if and only if the poset is a chain and the weight on coordinates is non-archimedean, the case where the weighted poset metric is an ultrametric (Section 4, Theorem 25). In addition, we will describe all optimal anticodes (Section 3.3, Theorem 19 and Theorem 20) and determine for some relevant instances all diameter perfect codes (Section 3.2, Corollary 14, Corollary 15 and Theorem 17; Section 3.4, Theorem 23). The Theorem 21 (Section 3.3) presents conditions on the weight on coordinates for a code to be diameter perfect. Section 2 is an introduction to weights, metrics, and related concepts used throughout this work.

2 Weights, Metrics and Related Concepts

2.1 Weights and Metrics

Let $[n] \triangleq \{1, 2, ..., n\}$ be a finite set with n elements and \leq_P be a partial order on [n]. We call the pair $P = ([n], \leq_P)$ a poset. If $k \leq_P j$ and $k \neq j$, we say that k is smaller than j and we write $k <_P j$. An ideal in $P = ([n], \leq_P)$ is a subset $I \subseteq [n]$ that contains every element in [n] smaller than or equal to some of its elements, i.e., if $j \in I$ and $k \leq_P j$ then $k \in I$. Given a subset $X \subset [n]$, we denote by $\langle X \rangle$ the smallest ideal containing X, called the ideal generated by X.

Let \mathbb{F}_q^n be the space of *n*-tuples over the finite field \mathbb{F}_q . Given a poset $P = ([n], \leq_P)$ and $u = (u_1, u_2, \ldots, u_n) \in \mathbb{F}_q^n$, the *support* of *u* is the set

$$supp(u) \triangleq \{i \in [n] : u_i \neq 0\}$$
.

The ideal $\langle supp(u) \rangle$ of P is denoted by I_u^P and its set of all maximal elements is denoted by M_u^P .

A map $w: \mathbb{F}_q^n \to \mathbb{N}$ is called a weight on \mathbb{F}_q^n if it satisfies the following properties:

- 1. $w(u) \ge 0$ for all $u \in \mathbb{F}_q^n$ and w(u) = 0 if and only if u = 0;
- 2. w(u) = w(-u) for all $u \in \mathbb{F}_q^n$;
- 3. $w(u+v) \leq w(u) + w(v)$ hold for all $u, v \in \mathbb{F}_q^n$ (triangle inequality).

It is clear that, given a weight w on \mathbb{F}_q^n , if we define the map $d: \mathbb{F}_q^n \times \mathbb{F}_q^n \to \mathbb{N}$ by

$$d(u, v) \triangleq w(u - v),$$

then d is a metric invariant by translations.

Definition 3 Given a poset $P = ([n], \leq_P)$ and a weight w on \mathbb{F}_q , the (P, w)-weight of $u \in \mathbb{F}_q^n$ is the non-negative integer

$$\varpi_{(P,w)}(u) \triangleq \sum_{i \in M_u^P} w(u_i) + M_w \cdot |I_u^P \setminus M_u^P|$$

where $M_w \triangleq \max\{w(\alpha) : \alpha \in \mathbb{F}_q\}$. If $u, v \in \mathbb{F}_q^n$, then their (P, w)-distance is defined by

$$d_{(P,w)}(u,v) \triangleq \varpi_{(P,w)}(u-v).$$

The (P, w)-weight $\varpi_{(P,w)}$ and the (P, w)-distance $d_{(P,w)}$ are also called weighted poset weight and weighted poset distance, respectively.

The (P, w)-weight is a weight on \mathbb{F}_q^n (see [11], Proposition 7), and therefore the (P, w)-distance is a metric invariant by translations, which combines and extends several classic weights of coding theory. When the weight w is the Hamming weight, the (P, w)-weight is the poset weight proposed by Brualdi $et\ al.$ in [4], and when the weight w is the Lee weight, the (P, w)-weight is the pomset weight proposed by Sudha and Selvaraj in [14] (see [11], Proposition 11). When P is the antichain order with n elements, i.e., $i \leq_P j$ in P if and only if i = j, the (P, w)-weight is an additive weight. The diagram in Figure 1 illustrates these facts.

Remark 4 We stress that the triangle inequality can fail if we consider supp(u) instead of M_u^P in the definition of $\varpi_{(P,w)}(u)$: for the usual order $1 \leq_P 2$ on [2] and the Lee weight w_L on \mathbb{F}_7 , if $u = (1,0), v = (-1,1) \in \mathbb{F}_7^2$, then $\varpi_{(P,w_L)}(u+v) \leq \varpi_{(P,w_L)}(u) + \varpi_{(P,w_L)}(v)$ if and only if $M_{w_L} \leq 2w_L(1)$, which is not true since $M_{w_L} = 3$ and $w_L(1) = 1$.

2.2 Anticodes and Diameter Perfect Codes

Let A be a subset of the weighted poset metric space $(\mathbb{F}_q^n, d_{(P,w)})$. The diameter of A is the maximum (P, w)-distance occurring between elements of A:

$$diam_{d_{(P,w)}}(A) \triangleq \max\{d_{(P,w)}(x,y) : x,y \in A\}.$$




Figure 1: A diagram of weights.

In this case we say that A is an anticode with diameter $diam_{d_{(P,w)}}(A)$. Let

$$A_{d_{(P,w)}}^*(D) \triangleq \max\{|A| : diam_{d_{(P,w)}}(A) \le D\}.$$

An anticode A is called D-optimal if $|A| = A_{d_{(P,w)}}^*(D)$.

The class of the weighted poset metric spaces $(\mathbb{F}_q^n, d_{(P,w)})$ is distinct from the class of the distance regular graph metric spaces and, thus, Delsarte's Theorem, does not apply. Fortunately, an alternative proof that was given in [6] can be slightly modified to work on this case (and for all metrics invariant by translations).

Theorem 5 If A and B are subsets of the weighted poset metric space $(\mathbb{F}_q^n, d_{(P,w)})$ such that nonzero (P, w)-distances occurring between elements in A do not occur between elements of B, then

$$|A| \cdot |B| \le q^n.$$

Proof. Suppose that A and B are subsets of \mathbb{F}_q^n such that nonzero (P, w)-distances occurring between elements in A do not occur between elements of B. Let

$$A+B\triangleq\{a+b:a\in A,\ b\in B\}.$$

Let us show that $|A + B| = |A| \cdot |B|$. Suppose $a_1 + b_1 = a_2 + b_2$ with $a_1, a_2 \in A$ and $b_1, b_2 \in B$. Since $d_{(P,w)}$ is invariant by translations,

$$d_{(P,w)}(a_1, a_2) = d_{(P,w)}(a_1 + b_1, a_2 + b_1)$$

= $d_{(P,w)}(a_2 + b_2, a_2 + b_1)$
= $d_{(P,w)}(b_2, b_1)$.

As nonzero (P, w)-distances occurring between elements in A do not occur between elements of B, it follows that $d_{(P,w)}(a_1, a_2) = d_{(P,w)}(b_2, b_1) = 0$. Therefore $a_1 = a_2$ and $b_1 = b_2$ and so $|A + B| = |A| \cdot |B|$. Since $A + B \subseteq \mathbb{F}_q^n$, we conclude that $|A| \cdot |B| = |A + B| \leq |\mathbb{F}_q^n| = q^n$.

A code is any subset $C \subseteq \mathbb{F}_q^n$ with minimum distance

$$d_{(P,w)}(C) \triangleq \min\{d_{(P,w)}(x,y): x,y \in C, \quad x \neq y\}.$$

Corollary 6 For all codes $C \subseteq \mathbb{F}_q^n$ with minimum distance $d_{(P,w)}(C) = D + 1$ we have

$$A_{d_{(P,w)}}^*(D) \cdot |C| \le q^n. \tag{2}$$

A code $C \subseteq \mathbb{F}_q^n$ is called diameter perfect if (2) holds with equality.

In this work, we will study the diameter perfect codes and optimal anticodes on $(\mathbb{F}_q^n, d_{(P,w)})$ when P is a chain order. We start by presenting the basic results of this weighted poset metric spaces.

3 Codes and Anticodes on Chain Orders

An order P on the finite set [n] is called a *chain* if every two elements are *comparable*, i.e., given $i, j \in [n]$ we have that either $i \leq_P j$ or $j \leq_P i$. In this case, the set can be labeled in such a way that $i_1 \leq_P i_2 \leq_P \ldots \leq_P i_n$. For the simplicity of the notation, in this situation, we will always assume that the chain order P is the usual chain order

$$1 \le 2 \le \ldots \le n$$
.

For the usual chain order $P = ([n], \leq)$ we have that $|\langle i \rangle| = i$ for each $i \in [n]$. So, given $0 \neq u = (u_1, \ldots, u_n) \in \mathbb{F}_q^n$,

$$\varpi_{(P,w)}(u) = w(u_i) + (i-1) \cdot M_w, \tag{3}$$

where $i = \max\{j : u_j \neq 0\}$. The metric space $(\mathbb{F}_q^n, d_{(P,w)})$ will be called weighted chain metric space. We remark that the weighted chain metric space is a particular case (a single chain) of the Niederreiter-Rosenbloom-Tsfasman spaces (several chains) if w is the Hamming weight. Originally, the Niederreiter-Rosenbloom-Tsfasman spaces were introduced by Niederreiter in [10] and Rosenbloom and Tsfasman in [13]. The Niederreiter-Rosenbloom-Tsfasman spaces are of special interest since they have several applications, as noted by Rosenbloom and Tsfasman (see [13]) and Park and Barg (see [12]).

In this section, we will away assume that $P = ([n], \leq)$ is the usual chain order and develop several results on codes and anticodes. Besides, we will omit the index P and write just d_w and ϖ_w for the weighted chain distance $d_{(P,w)}$ and weighted chain weight $\varpi_{(P,w)}$, respectively. In particular, if w is the Hamming weight, we will write just d and ϖ , the poset distance and poset weight, respectively.

We will denote by $B_w(u,r)$ the r-ball with center u and radius r relative to the weighted chain distance d_w , i.e.,

$$B_w(u,r) \triangleq \{v \in \mathbb{F}_q^n : d_w(u,v) \le r\}.$$

Remark 7 Writing $r = s + iM_w$ with $0 \le s < M_w$, if $s \ne w(a)$ for all $a \in \mathbb{F}_q$, then

$$B_w(u,r) = B_w(u,t+iM_w)$$

for all $s' \le t \le s$ where s' is the largest integer such that s' < s and s' = w(a) for some $a \in \mathbb{F}_q$.

From now on we will assume that $r = s + iM_w$ with s = w(a) for some $a \in \mathbb{F}_q$. Given $r \in \mathbb{N}$ such that $r \geq 1$ and w be a weight on \mathbb{F}_q , let

$$[r]_w \triangleq \{t \in [r] : t = w(a) \text{ for some } a \in \mathbb{F}_q\}$$

be the w-interval. We have that $[r]_w = [r] \cap Im(w)$, where Im(w) is the image of the map w.

Proposition 8 (Size of Ball) Let $D = S + R \cdot M_w$ be a non-negative integer and $x \in \mathbb{F}_q^n$. If S > 0, then

$$|B_w(x,D)| = q^R \cdot (1 + |w^{-1}([S]_w)|).$$

If S = 0, then

$$|B_w(x,D)| = q^R.$$

Proof. See Appendix 5. ■

If X is a subset of \mathbb{F}_q^n , the packing radius $R_w(X)$ is the largest positive integer number r such that any two r-balls centered at distinct elements of X are disjoint. In [11, Corollary 22] the authors show that

$$R_w(X) = M_w \cdot (d(C) - 1), \tag{4}$$

where d(C) is the minimum distance of C relative to the poset metric d.

We say that a code C is *perfect* if the union of the r-balls, $r = R_w(C)$, centered at the elements of C, covers \mathbb{F}_q^n .

3.1 MDS Codes

In this section, we describe all the MDS codes on (\mathbb{F}_q^n, d_w) . We shall see later (Section 3.2) that these will be the only diameter perfect codes for many instances of (\mathbb{F}_q^n, d_w) . We begin by showing that the minimum distance $d_w(C)$ is determined by d(C).

Given a weight w on \mathbb{F}_q , let $m_w \triangleq \min\{w(\alpha) : 0 \neq \alpha \in \mathbb{F}_q\}$.

Proposition 9 Let x_i be the i^{th} coordinate of $x \in \mathbb{F}_q^n$ and C be a code on \mathbb{F}_q^n . Then

$$d_w(C) = S_{w,C} + (d(C) - 1) \cdot M_w,$$

where $S_{w,C} \triangleq \min\{w(x_{d(C)} - y_{d(C)}) : x, y \in C, x_{d(C)} \neq y_{d(C)}\}$. Therefore, if C is a linear code, then

$$d_w(C) = m_w + (d(C) - 1) \cdot M_w.$$

Proof. Write $d_w(C) = S + R \cdot M_w$ with $m_w \leq S \leq M_w$. This implies that $d(c, c') \geq R+1$ for all $c \neq c' \in C$. Since $d(c, c') \geq d(C)$ for all $c \neq c' \in C$ and there are $c, c' \in C$ such that d(c, c') = d(C), we conclude that R + 1 = d(C), i.e., R = d(C) - 1. The

minimality of $d_w(C)$ implies that $S = \min\{w(x_{d(C)} - y_{d(C)}) : x, y \in C, x_{d(C)} \neq y_{d(C)}\}.$

A code $C \subseteq \mathbb{F}_q^n$ is said to be maximum distance separable (MDS) if its size |C| attains the Singleton bound

$$|C| \le q^{n - d(C) + 1}.$$

Proceeding with the same argument in the proof of Singleton bound (see [8], p. 63), we get: if C is an MDS code on \mathbb{F}_q^n , then

$$C = \{(x_y, y) : y \in \mathbb{F}_q^{n-d(C)+1}\},\$$

where $y \mapsto x_y$ is a map from $\mathbb{F}_q^{n-d(C)+1}$ into $\mathbb{F}_q^{d(C)-1}$. We now notice the following: since $|B_w(0, R_w(C))| = q^{d(C)-1}$ (see (4)),

$$|C| = q^{n-d(C)+1} \Leftrightarrow q^{d(C)-1} \cdot |C| = q^n$$

$$\Leftrightarrow |B_w(0, R_w(C))| \cdot |C| = q^n.$$

In short:

Theorem 10 In the weighted chain metric space (\mathbb{F}_q^n, d_w) , a code C is MDS if and only if C is perfect. Furthermore, if $C \subseteq \mathbb{F}_q^n$ is MDS (perfect), then

$$C = \{(x_y, y) : y \in \mathbb{F}_q^{n-d(C)+1}\},\$$

where $y \mapsto x_y$ is a map from $\mathbb{F}_q^{n-d(C)+1}$ into $\mathbb{F}_q^{d(C)-1}$.

3.2 Diameter Perfect Codes

In this section we will show that all MDS codes on (\mathbb{F}_q^n, d_w) are diameter perfect. We will also present some instances where (2) is equivalent to the Singleton bound. Consequently, in these instances, diameter perfect and MDS are equivalent concepts.

Let $A \subseteq \mathbb{F}_q^n$ such that $diam_{d_w}(A) \leq D$. Since $d_w(x,y) \leq D$ for all $x,y \in A$, we have that

$$A \subseteq B_w(x, D) \tag{5}$$

for each $x \in A$. Hence $|A| \leq |B_w(x,D)|$. Therefore,

$$A_{d_{w}}^{*}(D) \le |B_{w}(x,D)|,$$
 (6)

by Lemma 8 it follows that:

Theorem 11 Let $D = S + R \cdot M_w$ be a non-negative integer with $0 \le S < M_w$. If S > 0, then

$$A_{d_w}^*(D) \le q^R \cdot (1 + |w^{-1}([S]_w)|). \tag{7}$$

If S = 0, then

$$A_{d_w}^*(D) \le q^R.$$

Consequently, $A_{d_w}^*(D) < q^{R+1}$.

In Section 4 we will show that (7) holds with equality if and only if the weight w is non-archimedean.

Theorem 12 Let $D = \varpi_w(y)$ for some $y \in \mathbb{F}_q^n$. The following properties are equivalent:

- 1. $diam_{d_w}(B_w(x,D)) = D$ for all $x \in \mathbb{F}_q^n$;
- 2. $B_w(x, D)$ is D-optimal for all $x \in \mathbb{F}_q^n$;
- 3. If A is D-optimal, then $A = B_w(x, D)$ for some $x \in \mathbb{F}_q^n$;
- 4. $A_{dw}^*(D) = |B_w(0, D)|$.

Proof. (1) \Rightarrow (2): Let us suppose that $diam_{d_w}(B_w(x,D)) = D$ for all $x \in \mathbb{F}_q^n$. Since $A \subseteq B_w(x,D)$ for all $x \in A$ whenever $diam_{d_w}(A) \leq D$, we get $|B_w(x,D)| = A_{d_w}^*(D)$ for all $x \in \mathbb{F}_q^n$. Hence $B_w(x,D)$ is D-optimal for all $x \in \mathbb{F}_q^n$.

- $(2) \Rightarrow (3)$: Now suppose that $B_w(x, D)$ is D-optimal for all $x \in \mathbb{F}_q^n$ and let A be an anticode such that $|A| = A_{d_w}^*(D)$, i.e., A is D-optimal. Since $A \subseteq B_w(x, D)$ for all $x \in A$ and $|B_w(x, D)| = A_{d_w}^*(D)$, then $|A| = |B_w(x, D)|$ for all $x \in A$. Thus $A = B_w(x, D)$ for all $x \in A$.
- (3) \Rightarrow (4): Suppose that A is D-optimal and $A = B_w(x, D)$ for some $x \in \mathbb{F}_q^n$. Then $A_{d_w}^*(D) = |A| = |B_w(x, D)|$. Since d_w is invariant by translations, $|B_w(x, D)| = |B_w(0, D)|$. Thus $A_{d_w}^*(D) = |B_w(0, D)|$.
- $(4) \Rightarrow (1)$: Let us suppose that $A_{d_w}^*(D) = |B_w(0,D)|$. If A is D-optimal, then $|A| = |B_w(0,D)|$. Since $A \subseteq B_w(x,D)$ for all $x \in A$ and d_w is invariant by translation, $|A| = |B_w(0,D)| = |B_w(x,D)|$ for all $x \in A$, and hence $A = B_w(x,D)$ for all $x \in A$. This implies that $B_w(x,D)$ is D-optimal. Therefore $diam_{d_w}(B_w(x,D)) \leq D$. As $D = \varpi_w(y)$ for some $y \in \mathbb{F}_q^n$ and d_w is invariant by translation, $D = d_w(x,x+y)$, which implies that $diam_{d_w}(B_w(x,D)) = D$ for all $x \in A$. Thus $diam_{d_w}(B_w(x,D)) = D$ for all $x \in \mathbb{F}_q^n$.

Proposition 13 Let $D = R \cdot M_w$ and $x \in \mathbb{F}_q^n$. The following properties holds (and are equivalent):

- 1. $diam_{d_w}(B_w(x, D)) = D;$
- 2. $B_w(x, D)$ is D-optimal;
- 3. $B_w(0,D)$ is an R-dimensional subspace of \mathbb{F}_q^n .

Consequently, $A_{d_w}^*(D) = q^R$.

Proof. We have that

$$B_w(0,D) = \{(x_1,\ldots,x_R,0,\ldots,0) : x_1,\ldots,x_R \in \mathbb{F}_q\}.$$

Hence $B_w(0,D)$ is an R-dimensional subspace of \mathbb{F}_q^n . Therefore, $x-y \in B_w(0,D)$ for all $x,y \in B_w(0,D)$ and hence $diam_{d_w}(B_w(0,D)) = D$. Since d_w is invariant by

translations, $diam_{d_w}(B_w(x, D)) = D$ for all $x \in \mathbb{F}_q^n$. By Theorem 12 and Proposition 8 it follows that item 2 and $A_{d_w}^*(D) = q^R$ holds, and hence items 1, 2 and $A_{d_w}^*(D) = q^R$ are equivalent.

Now if $B_w(0, D)$ is D-optimal, $diam_{d_w}(B_w(0, D)) \leq D$, and since $D = R \cdot M_w$, we have that $x - \lambda y \in B_w(0, D)$ for all $x, y \in B_w(0, D)$ and $\lambda \in \mathbb{F}_q$, i.e., $B_w(0, D)$ is a subspace of \mathbb{F}_q^n .

Thus the items 1, 2, 3 and $A_{d_w}^*(D) = q^R$ holds and are all equivalent.

Let w be a weight on \mathbb{F}_q and D be a non-negative integer. We denote by $\lfloor D \rfloor_w$ the largest weight w(a) such that w(a) < D.

As $\lfloor D \rfloor_w = R \cdot M_w$ whenever $D = m_w + R \cdot M_w$, by Proposition 13 it follows that $A_{d_w}^*(\lfloor D \rfloor_w) = q^R$. Hence:

Corollary 14 Let C be a code with minimum distance $d_w(C) = m_w + (d(C) - 1) \cdot M_w$. Then

$$A_{d_m}^*(\lfloor d_w(C) \rfloor_w) \cdot |C| \le q^n \tag{8}$$

is equivalent to the Singleton bound. Therefore, if C is a code with minimum distance $d_w(C) = m_w + (d(C) - 1) \cdot M_w$, then C is diameter perfect if and only if C is MDS.

Let w_H be the Hamming weight on \mathbb{F}_q . Since $d_w(C) = d(C) \cdot M_w$ whenever $w = \lambda w_H$ for some integer $\lambda > 0$:

Corollary 15 If $w = \lambda w_H$ for some integer $\lambda > 0$, then

$$A_{dw}^*(|d_w(C)|_w) \cdot |C| \le q^n$$

is equivalent to the Singleton bound. Therefore, a code C is diameter perfect if and only if C is MDS.

In [1] Ahlswede *et al.* proved that MDS codes in Hamming spaces are diameter perfect. This is also our case:

Theorem 16 If C is an MDS code with minimum distance $d_w(C) = D$, then

$$A_{d_w}^*(\lfloor D\rfloor_w) \cdot |C| = q^n.$$

Proof. If C is an MDS code with minimum distance $d_w(C) = D$, then $D = m_w + (d(C) - 1) \cdot M_w$. This implies that $\lfloor D \rfloor_w = (d(C) - 1) \cdot M_w$, and hence $A_{d_w}^*(\lfloor D \rfloor_w) = q^{d(C)-1}$ (see Proposition 13). Since $|C| = q^{n-d(C)+1}$, the result follows.

The MDS codes (linear and non-linear) are described in Theorem 10.

Let C be a code on \mathbb{F}_q^n such that $|C| = q^k$ for some $0 \le k \le n$. Suppose $d_w(C) = S + (d(C) - 1) \cdot M_w$ with $m_w \le S < M_w$ be a distance (see Proposition 9).

If C is not an MDS code, then $|C| \leq q^{n-d(C)+1-i}$ for some integer $i \geq 1$, and since $A_{d_w}^*(\lfloor D \rfloor_w) < q^{d(C)}$ (see Theorem 11), we have

$$A_{d_w}^*(|D|_w) \cdot |C| < q^{n+1-i} \le q^n.$$

Now, if |C| is an MDS code, then $|C| = q^{n-d(C)+1}$ and $d_w(C) = m_w + (d(C)-1) \cdot M_w$. If $D = d_w(C)$, by Proposition 13, $A_{d_w}^*(\lfloor D \rfloor_w) = q^{d(C)-1}$. Thus

$$A_{d_w}^*(\lfloor D \rfloor_w) \cdot |C| = q^n.$$

In short:

Theorem 17 A code C of size power of q is diameter perfect if and only if C is MDS. In particular, a linear code C is diameter perfect if and only if C is MDS.

As we shall see in Theorem 19 and Theorem 20, not always the inequality (8) is equivalent to the Singleton bound.

3.3 Optimal Anticodes

In this section, we determine all the D-optimal anticodes. If $D = S + R \cdot M_w$, the idea is to partition the w-interval $[M_w]_w$ in "non-archimedean" and "not always non-archimedean" elements and analyze S (and consequently D) considering this partition. More precisely, if S is in the part "non-archimedean" then all D-optimal anticodes are balls of radius D (Theorem 19), and if S is in part "not always non-archimedean" then all D-optimal anticodes are proper parts of balls of radius D (Theorem 20).

We say that a weight w on \mathbb{F}_q^n is non-archimedean if

$$w(x+y) < \max\{w(x), w(y)\}\$$

for all $x, y \in \mathbb{F}_q^n$. Otherwise, we will say that the weight is archimedean.

Example 18 The Lee weight w_L on \mathbb{Z}_m is archimedean if $m \geq 4$: for x = y = 1,

$$w_L(x+y) > \max\{w(x), w(y)\}.$$

The Hamming weight w_H on \mathbb{Z}_m is non-archimedean.

Given an archimedean weight w on \mathbb{F}_q , let $m_w \leq S_w < M_w$ be the integer

$$S_w \triangleq \min\{\max\{w(a), w(b)\} : a, b \in \mathbb{F}_q \text{ and } w(a-b) > \max\{w(a), w(b)\}\}.$$

If w is non-archimedean, we define $S_w \triangleq M_w$. Notice that $S_w > 0$ for all weights w.

Theorem 19 Let (\mathbb{F}_q^n, d_w) be the weighted chain metric space and D be a non-negative integer. Write $D = S + R \cdot M_w$ with $0 \leq S < M_w$. If $0 \leq S < S_w$, then:

- 1. $diam_{d_w}(B_w(x,D)) = D;$
- 2. $B_w(x, D)$ is D-optimal for all $x \in \mathbb{F}_q^n$;
- 3. If A is D-optimal, then $A = B_w(x, D)$ for some $x \in \mathbb{F}_q^n$;
- 4. $B_w(0,D)$ is an R-dimensional subspace of \mathbb{F}_q^n if and only if S=0.

Consequently:

- 5. If S = 0, then $A_{dw}^*(D) = q^R$;
- 6. If $S \neq 0$, then $A_{d_w}^*(D) = q^R \cdot (1 + |w^{-1}([S]_w)|)$.

Proof. Since $w(a-b) \leq \max\{w(a), w(b)\}$ if $\max\{w(a), w(b)\} \leq S_w - 1$, $a, b \in \mathbb{F}_q$, and since $S < S_w$, it follows that $d_w(x,y) \leq D$ for all $x,y \in B_w(0,D)$. Hence $diam_{d_w}(B_w(0,D)) \leq D$. For $x \in \mathbb{F}_q^n$ such that $\varpi_P(x) = R + 1$ and $w(x_{R+1}) = S$ we have $d_w(0,x) = D$. Thus $diam_{d_w}(B_w(0,D)) = D$. As d_w is invariant by translations, we get

$$diam_{dw}(B_w(x,D)) = D$$

for all $x \in \mathbb{F}_q^n$. From this and Theorem 12, together with Proposition 8, we get the items 2, 3, 5 and 6.

Proposition 13 ensures that $B_w(0,D)$ is an R-dimensional subspace if S=0. Note that if $S_w=m_w$, then S=0. Assume now $m_w \leq S < S_w$. Put $x=(x_1,\ldots,x_{R-1},a,0,\ldots,0)$. If $B_w(0,D)$ is a linear subspace, then $w(\lambda a) \leq S$ for all $\lambda \in \mathbb{F}_q$ and for all $a \in \mathbb{F}_q$ such that $w(a) \leq S$. Since $S < M_w$ there is $b \in \mathbb{F}_q$ such that w(b) > S. For $\lambda = ba^{-1}$, we have $w(b) = w(\lambda a) \leq S$, a contradiction. Thus, if $B_w(0,D)$ is a linear subspace of \mathbb{F}_q^n , S=0.

If $A = B_w(x, D)$ and $S \ge S_w$ then $diam_{d_w}(A) > D$, hence A we will not be D-optimal. We will now describe the subsets of $B_w(x, D)$ that are D-optimal anticodes. For $S \ge S_w$, let $W_w(S)$ be a subset of \mathbb{F}_q of maximal size such that:

- 1. if $a \in W_w(S)$, then $S_w \leq w(a) \leq S$;
- 2. if $a \in W_w(S)$ and $0 \le w(b) \le S_w 1$, then $w(a b) \le S$;
- 3. if $a, b \in W_w(S)$, then $w(a b) \leq S$.

Let $\mathcal{W}_w(S)$ be the set of all subsets $W_w(S)$. Given integer non-negative R and $K \in \mathcal{W}_w(S)$, let

$$Y_{w,R}(K) \triangleq \{x \in \mathbb{F}_q^n : \varpi(x) = R+1 \text{ and } x_{R+1} \in K\}.$$

In other words, if $x \in Y_{w,R}(K)$, then:

- 1. $|I_x^P| = R + 1;$
- 2. $S_w + R \cdot M_w < d_w(0, x) < S + R \cdot M_w$;

- 3. $d_w(x,y) \leq S + R \cdot M_w$ for all $y \in \mathbb{F}_q^n$ such that $d_w(0,y) \leq (S_w 1) + R \cdot M_w$;
- 4. $d_w(x,y) \leq S + R \cdot M_w$ for all $y \in Y_{w,R}(K)$.

Theorem 20 Let (\mathbb{F}_q^n, d_w) be the weighted chain metric space and D be a non-negative integer. Write $D = S + R \cdot M_w$ with $0 \le S < M_w$ and let $D' = \lfloor S_w \rfloor_w + R \cdot M_w$. If $S \ge S_w$, then:

- 1. $D' \leq diam_{d_w}((x + Y_{w,R}(K)) \cup B_w(x,D')) \leq D$ for all $x \in \mathbb{F}_q^n$ and for all $K \in \mathcal{W}_w(S)$;
- 2. $(x+Y_{w,R}(K)) \cup B_w(x,D')$ is D-optimal for all $x \in \mathbb{F}_q^n$ and for all $K \in \mathcal{W}_w(S)$;
- 3. If A is D-optimal, then $A = (x + Y_{w,R}(K)) \cup B_w(x,D')$ for some $x \in \mathbb{F}_q^n$ and $K \in \mathcal{W}_w(S)$.

Consequently,

$$A_{d_w}^*(D) = q^R \cdot (1 + |w^{-1}([S_w - 1]_w)| + |W_w(S)|).$$

Proof. Let $K \in \mathcal{W}_w(S)$. If $x, y \in Y_{w,R}(K)$, then $\varpi_w(x) \leq D$, $\varpi_w(y) \leq D$ and $d_w(x,y) \leq D$. This implies that $Y_{w,R} \subseteq B_w(0,D)$ and $diam_{d_w}(Y_{w,R}) \leq D$. If

$$X_{w,R}(K) \triangleq Y_{w,R}(K) \cup B_w(0,D')$$

(see Figure 2), we can see that $diam_{d_w}(X_{w,R}(K)) \leq D$, and since $diam_{d_w}(B_w(0,D')) = D'$, it follows that $D' \leq diam_{d_w}(X_{w,R}(K)) \leq D$.




Figure 2: The set $X_{w,R}(K)$, where $\widetilde{D} = S_w + R \cdot M_w$.

We claim now that if $diam_{d_w}(A) \leq D$, then $A \subseteq x + X_{w,R}(K)$ for some $x \in \mathbb{F}_q^n$ and $K \in \mathcal{W}_w(S)$. Let $A_a = (-a) + A$ for some $a \in A$. We have $0 \in A_a$ and $diam_{d_w}(A_a) = diam_{d_w}(A)$. Since $A_a \subseteq B_w(0, D)$, $\varpi_w(x) \leq D$ for all $x \in A_a$:

• if $\varpi(x) < R+1$, then $x \in B_w(0, D')$;

- if $\varpi(x) = R + 1$ and $w(x_{R+1}) < S_w$, then $x \in B_w(0, D')$;
- if $\varpi(x) = R+1$ and $S_w \leq w(x_{R+1}) \leq S$, since $d_w(x,y) \leq D$ for all $y \in A_a$ such that $\varpi_P(y) = R+1$, then $w(x_{R+1}-b) \leq S$ for all $b \in \mathbb{F}_q$ such that $b = y_{R+1}$ for some $y \in A_a$.

Hence $A_a \subseteq X_{w,R}(K)$ for some $K \in \mathcal{W}_w(S)$, i.e., $A \subseteq x + X_{w,R}(K)$ with x = a. Since $diam_{d_w}(x + X_{w,R}(K)) \leq D$, it follows that $A_{d_w}^*(D) = |X_{w,R}(K)|$. As $Y_{w,R}(K) \cap B_w(0,D') = \emptyset$, we conclude that

$$A_{d_w}^*(D) = q^R \cdot (1 + |w^{-1}([S_w - 1]_w)| + |W_w(S)|),$$

and the desire result follows.

We conclude this section by showing that (2) depends only on w and d(C). Let $W^w(S)$ be a subset of \mathbb{F}_q of maximal size such that:

- 1. if $a, b \in W^w(S)$ and $a \neq b$, then $w(a b) \geq S$;
- 2. there are $a, b \in W^w(S)$ such that w(a b) = S.

Given an integer $m_w < S < M_w$ such that S = w(a) for some $a \in \mathbb{F}_q$, let

$$C = \{(0, \dots, 0, c_R, c_{R+1}, \dots, c_n) : c_{R+1}, \dots, c_n \in \mathbb{F}_q, c_R \in W^w(S)\}.$$

We have that C is a code in \mathbb{F}_q^n with minimum distance $d_w(C) = S + (R-1) \cdot M_w$ and size $q^{n-R} \cdot |W^w(S)|$. By Theorem 19, for $D = d_w(C)$ and suppose $S \leq S_w$, it follows that $A_{d_w}^*(\lfloor D \rfloor_w) = q^{R-1} \cdot (1 + |w^{-1}([S-1]_w)|)$. Hence,

$$A_{d_w}^*(\lfloor D \rfloor_w) \cdot |C| \le q^n$$

if and only if

$$(1 + |w^{-1}([S-1]_w)|) \cdot |W^w(S)| \le q.$$

Supposing now $S > S_w$, by Theorem 20,

$$A_{d_w}^*(\lfloor D\rfloor_w) = q^{R-1} \cdot (1 + |w^{-1}([S_w - 1]_w)| + |W_w(S - 1)|),$$

and hence,

$$A_{d_w}^*(\lfloor D \rfloor_w) \cdot |C| \le q^n$$

if and only if

$$(1 + |w^{-1}([S_w - 1]_w)| + |W_w(S - 1)|) \cdot |W^w(S)| \le q.$$

The case $S = m_w$ is equivalent to the Singleton bound (Corollary 14). In short:

Theorem 21 For the weighted chain metric space, we have that (2) is equivalent to

$$(1 + |w^{-1}([S-1]_w)|) \cdot |W^w(S)| \le q \tag{9}$$

for all $m_w < S \le S_w$ and

$$(1 + |w^{-1}([S_w - 1]_w)| + |W_w(S - 1)|) \cdot |W^w(S)| \le q \tag{10}$$

for all $S_w < S < M_w$. Therefore, a code C with minimum distance $d_w(C) = S + R \cdot M_w$ such that $m_w < S < M_w$ is diameter perfect if and only if (9) or (10) holds with equality.

If w is non-archimedean, then $S_w = M_w$. Therefore:

Corollary 22 For the weighted chain metric space (\mathbb{F}_q^n, d_w) with d_w ultrametric we have that (2) is equivalent to

$$(1 + |w^{-1}([S-1]_w)|) \cdot |W^w(S)| \le q \tag{11}$$

for all $S > m_w$. Therefore, a code C with minimum distance $d_w(C) = S + R \cdot M_w$ such that $S > m_w$ is diameter perfect if and only if (11) holds with equality.

3.4 Diameter Perfect Codes on \mathbb{F}_q^n with q Prime

Now let us suppose that q=p is a prime number. Let $C\subseteq \mathbb{F}_p^n$ be a code with $d_w(C)=S+(d(C)-1)\cdot M_w$. Since

$$1 + |w^{-1}([S-1]_w)| < p$$

for all $m_w < S < M_w$,

$$|W^w(S)| < p$$

for all $m_w < S < M_w$ and

$$1 + |w^{-1}([S_w - 1]_w)| + |W_w(S - 1)| < p$$

for all $S_w < S < M_w$, by Theorem 21 if C is diameter perfect, then $S = m_w$. By Corollary 14 and Theorem 16:

Theorem 23 Let p be a prime number. In (\mathbb{F}_p^n, d_w) the only diameter perfect codes are the MDS codes.

4 Non-Archimedean Weights and Ultrametrics

We say that a metric d on \mathbb{F}_q^n is an ultrametric if

$$d(x,y) \leq \max\{d(x,z),d(z,y)\}$$

for all $x, y, z \in \mathbb{F}_q^n$.

Proposition 24 The weighted chain distance d_w is an ultrametric if and only if w is a non-archimedean weight on \mathbb{F}_q .

Proof. See Appendix 6.

We know that if $d_{(P,w)}$ is the weighted chain distance with w non-archimedean, then $diam_{d_{(P,w)}}(B_{(P,w)}(x,D)) = D$ for all $x \in \mathbb{F}_q^n$ and for all D (Theorem 19 with $S_w = M_w$). Now, we will show that $diam_{d_{(P,w)}}(B_{(P,w)}(x,D)) = D$ for all $x \in \mathbb{F}_q^n$ and for all D only if $d_{(P,w)}$ is the weighted chain distance with w non-archimedean.

Theorem 25 Let $d_{(P,w)}$ be a weighted poset distance. Then

$$diam_{d_{(P,w)}}(B_{d_{(P,w)}}(x,D)) = D$$

for all $x \in \mathbb{F}_q^n$ and for all D if and only if $d_{(P,w)}$ is the weighted chain distance d_w and $d_{(P,w)}$ is an ultrametric.

Proof. Suppose that d_w is an ultrametric. By Proposition 24, w is non-archimedian. This implies that $S_w = M_w$. The "if" part follows from Theorem 19 with $S_w = M_w$.

Let us suppose now that $P=([n], \leq_P)$ is not a chain order. Then, there are $i, j \in [n]$ not comparable on P. Put $x, y \in \mathbb{F}_q^n$ with $supp(x)=\{i\}$ and $supp(y)=\{j\}$ such that $w(x_i)=w(y_j)$, assuming that $S=\varpi_w(x)\leq \varpi_w(y)=R$, we will have that $x,y\in B_w(0,R)$ and $d_w(x,y)=\varpi_w(x-y)>R$, which implies $diam_{d_w}(B_w(0,R))>R$. This shows that P is a chain order.

Suppose now that P is the chain order and d_w is not an ultrametric. By Proposition 24, w is archimedian, i.e., there are $a, b \in \mathbb{F}_q$ such that

$$w(a-b) > \max\{w(a), w(b)\}.$$

Let $x, y \in \mathbb{F}_q^n$ such that $supp(x) = supp(y) = \{R+1\}$ with $x_{R+1} = a$ and $y_{R+1} = b$. If $S = \max\{w(a), w(b)\}$, then

$$\varpi_w(x) = w(a) + R \cdot M_w < S + R \cdot M_w$$

and

$$\varpi_w(y) = w(b) + R \cdot M_w < S + R \cdot M_w.$$

Hence $x, y \in B_w(0, D)$ with $D = S + R \cdot M_w$. But

$$d_w(x,y) = \varpi_w(x-y)$$

$$= w(a-b) + R \cdot M_w$$

$$> S + R \cdot M_w$$

$$= D.$$

This show that $diam_{d_w}(B_w(0,D)) > D$, and the "only if" part follows.

From Theorem 25, together with Theorem 12 and Proposition 8, it follows that:

Corollary 26 Let (\mathbb{F}_q^n, d_w) be a weighted chain metric space and D be a non-negative integer. Write $D = S + R \cdot M_w$ with $0 \leq S < M_w$. Then d_w is an ultrametric if and only if any of the equivalent properties below holds:

- 1. $diam_{d_w}(B_w(x,D)) = D$ for all $x \in \mathbb{F}_q^n$;
- 2. $B_w(x,D)$ is D-optimal for all $x \in \mathbb{F}_q^n$;
- 3. If A is D-optimal, then $A = B_w(x, D)$ for some $x \in \mathbb{F}_q^n$;
- 4. $A_{d_w}^*(D) = q^R \text{ if } S = 0 \text{ and } A_{d_w}^*(D) = q^R \cdot (1 + |w^{-1}([S]_w)|) \text{ if } S \neq 0.$

5 Proof of Proposition 8

Proof. Since d_w is invariant by translations, we have that

$$|B_w(x,D)| = |B_w(0,D)|$$

for all $x \in \mathbb{F}_q^n$ and $D \ge 0$. Given $x \in \mathbb{F}_q^n$ such that either $\varpi(x) = R + 1$ and $w(x_{R+1}) > S$ or $\varpi(x) > R + 1$,

$$\varpi_w(x) = w(x_{\varpi(x)}) + (\varpi(x) - 1) \cdot M_w > D,$$

i.e., $x \notin B_w(0, D)$. Now for each $x \in \mathbb{F}_q^n$ such that either $\varpi(x) = R + 1$ and $w(x_{R+1}) \leq S$ or $\varpi(x) < R + 1$ we have that

$$\varpi_w(x) = w(x_{\varpi(x)}) + (\varpi(x) - 1) \cdot M_w \le D.$$

Hence $x \in B_w(0, D)$ if and only if either $\varpi(x) = R + 1$ and $w(x_{R+1}) \leq S$ or $\varpi(x) < R + 1$. Thus

$$|B_w(0,D)| = q^R \cdot |w^{-1}([S]_w)| + q^R,$$

and the desire result follows.

6 Proof of Proposition 24

Proof. If d_w is an ultrametric and there are $x, y \in \mathbb{F}_q$ such that $w(x + y) > \max\{w(x), w(y)\}$, taking $u, v \in \mathbb{F}_q^n$ with $supp(u) = supp(v) = \{i\}$ with $u_i = x$ and $v_i = -y$, we have that

$$\begin{split} d_w(u,v) &= \varpi_w(u-v) \\ &= w(u_i-v_i) + (i-1) \cdot M_w \\ &> \max\{w(x),w(y)\} + (i-1) \cdot M_w \\ &= \max\{w(x) + (i-1) \cdot M_w, \\ &\qquad \qquad w(y) + (i-1) \cdot M_w \} \\ &= \max\{\varpi_w(u),\varpi_w(v)\} \\ &= \max\{d_w(u,0),d_w(0,v)\}, \end{split}$$

which is a contradiction. Thus w is a non-archimedean weight.

Suppose now that w is a non-archimedean weight. We claim that ϖ_w is a non-archimedean weight on \mathbb{F}_q^n : if $x, y \in \mathbb{F}_q^n$ and $i = \max\{j : x_j + y_j \neq 0\}$, then

Hence

$$d_w(x,y) = \varpi_w(x-z-y+z)$$

$$\leq \max\{\varpi_w(x-z), \varpi_w(z-y)\}$$

$$= \max\{d_w(x,z), d_w(z,y)\}$$

for all $x, y \in \mathbb{F}_q^n$. Thus d_w is an ultrametric.

References

- [1] R. Ahlswede, H. K. Aydinian and L. H. Khachatrian, "On perfect codes and related concepts," *Designs, Codes and Cryptography*, vol. 22, pp. 221-237, 2001.
- [2] N. Biggs, "Perfect codes in graphs," Journal of Combinatorial Theory (B), vol. 15, pp. 289-296, 1973.
- [3] A. E. Brouwer, A. M. Cohen and A. Neumaier, *Distance-regular graphs*, Springer-Verlag, 1989.
- [4] R. Brualdi, J. S. Graves and M. Lawrence, "Codes with a poset metric," *Discrete Mathematics*, vol. 147, pp. 57-72, 1995.
- [5] P. Delsarte, "An algebraic approach to association schemes of coding theory," *Phillips J. Res.*, vol. 10, 1973.
- [6] M. Deza and P. Frankl, "On maximal numbers of permutations with given maximal or minimal distance," J. Combin. Theory Ser. A, vol. 22, 1977.
- [7] T. Etzion, "Configuration distribution and designs of codes in the Johnson scheme," J. Combin. Des., vol. 15, no. 1, pp. 15-34, 2007.
- [8] M. Firer, M. M. S. Alves, J. A. Pinheiro and L. Panek, *Poset codes: partial orders, metrics and coding theory*, Springer International Publishing, 2018.
- [9] J. Martin and X. J. Zhu, "Anticodes for the Grassmann and bilinear forms graphs," *Des. Codes Cryptogr.*, vol. 6, pp. 73-79, 1995.

- [10] H. Niederreiter, "A combinatorial problem for vector spaces over finite fields," *Discrete Mathematics*, vol. 96, pp. 221-228, 1991.
- [11] L. Panek and J. A. Pinheiro, "General approach to poset and additive metrics," in *IEEE Trans. Inf. Theory*, vol. 66, no. 11, pp. 6823-6834, Nov. 2020.
- [12] W. Park and A. Barg, "The ordered Hamming metric and ordered symmetric channels," in *IEEE International Symposium on Information Theory Proceeding*, pp. 2283-2287, IEEE, 2011.
- [13] M. Y. Rosenbloom and M. A. Tsfasman, "Codes for the *m*-metric," *Problems of Information Transmission*, vol. 33, no. 1, pp. 45-52, 1997.
- [14] I. G. Sudha and R. S. Selvaraj, "Codes with a pomset metric and constructions," *Designs, Codes and Cryptography*, vol. 86, pp. 875-892, 2018.