

# A Theory of Auto-Scaling for Resource Reservation in Cloud Services

Konstantinos Psychas, Javad Ghaderi  
Department of Electrical Engineering  
Columbia University

## ABSTRACT

We consider a distributed server system consisting of a large number of servers, each with limited capacity on multiple resources (CPU, memory, disk, etc.). Jobs with different rewards arrive over time and require certain amounts of resources for the duration of their service. When a job arrives, the system must decide whether to admit it or reject it, and if admitted, in which server to schedule the job. The objective is to maximize the expected total reward received by the system. This problem is motivated by control of cloud computing clusters, in which, jobs are requests for Virtual Machines or Containers that reserve resources for various services, and rewards represent service priority of requests or price paid per time unit of service by clients. We study this problem in an asymptotic regime where the number of servers and jobs' arrival rates scale by a factor  $L$ , as  $L$  becomes large. We propose a resource reservation policy that asymptotically achieves *at least*  $1/2$ , and under certain monotone property on jobs' rewards and resources, at least  $1 - 1/e$  of the optimal expected reward. The policy automatically scales the number of VM slots for each job type as the demand changes, and decides in which servers the slots should be created in advance, without the knowledge of traffic rates. It effectively tracks a low-complexity greedy packing of existing jobs in the system while maintaining only a small number,  $g(L) = \omega(\log L)$ , of reserved VM slots for high priority jobs that pack well.

## KEYWORDS

Scheduling, Loss Systems, Fluid Limits, Resource Allocation

## 1 INTRODUCTION

There has been a rapid migration of computing, storage, applications, and other services to cloud. By using cloud (e.g., Amazon AWS [2], Microsoft Azure [41], Google Cloud [20]), clients are no longer required to install and maintain their own infrastructure. Instead, clients use the cloud resources on demand, by procuring Virtual Machines (VMs) or Containers [5, 21] with specific configurations of CPU, memory, disk, and networking in the cloud data center, depending on their needs.

A key challenge for the cloud service providers is to efficiently support a wide range of services on their physical platform. They usually offer QoS guarantees (in SLAs) [6] for clients' applications and services, and allow the number of VM instances to scale up or down with demand to ensure QoS guarantees are met. For example, in Amazon EC2 auto-scaling [4], clients can define simple rules to launch or terminate VM instances as their application demand increases or decreases. Various predictive and reactive schemes have been proposed for dynamically allocating VMs to different services, e.g., [19, 25, 30, 38, 45, 47], however, they mostly assume a dedicated hosting model where VMs of each application run on

a dedicated set of servers. Such models do not consider potential consolidation of VMs in servers which is known to significantly improve efficiency and scalability [13, 49]. For instance, suppose a CPU-intensive VM, a disk-intensive VM, and a memory-intensive VM are located on three individual servers (for our purpose, we use the terms VM and Container interchangeably). The cloud operator can pack these VMs in a single server to fully utilize its resources along CPU, disk, and memory, then the two unused servers can be used to pack additional VMs and serve more requests. However, in the absence of an accurate estimate of the workload, or when the workload varies over time and space, it is not clear how many VM instances an application launches and which VMs must be packed in which servers to ensure efficiency.

In this paper, we consider a cloud data center consisting of a large number of servers. As an abstraction in our model, a VM is simply a multi-dimensional object (vector of resource requirements) that should be served by one server and *cannot be fragmented*. Each server has a limited fixed capacity on its available resources (CPU, memory, disk, networking). VM requests belong to a collection of VM types, each with a specific resource requirement vector, and a specific reward that represents its service priority or the price that will be paid per time unit of service by the client. When a VM request arrives, we must decide in an online manner whether to accept it, and, if so, in which server to schedule it. The objective is to *maximize the expected total reward* received by the system. Note that finding the right packing for a given workload is a hard combinatorial problem (related to multi-dimensional Knapsack [32]). The absence of accurate estimate of workload (VM traffic rates and service durations) makes the problem even more challenging. For instance, consider a simple scenario with three types of VMs with the following (CPU, memory) requirement and rewards: (0.6, 0.6) with reward 4, (0.7, 0.1) with reward 3, and (0.1, 0.7) with reward 3. Server's capacity is normalized to (1,1). Hence, a server can accommodate a single (0.6, 0.6) VM, or pack one (0.7, 0.1) VM and one (0.1, 0.7) VM together. Suppose there is one empty server, and a (0.6, 0.6) VM request arrives. Should we admit this request and receive a reward of 4, or reserve the server to pack one (0.7, 0.1) VM and one (0.1, 0.7) VM in future, which can potentially yield a maximum reward of 6?

This problem is related to the *Online Multiple Knapsack* problem, in which there is a set of bins of finite capacity, items with various sizes and profits arrive one by one, and the goal is to pack them in an *online* manner into the bins so as to maximize their total profit. In general, this problem does not have any competitive (constant approximation) algorithm [39], even when items are allowed to be removed from any bin at any time. Hence, proposed competitive algorithms focus on more restricted cases of the problem [14, 29].

In this paper, we study a stochastic version of the problem in an asymptotic regime, where the number of servers  $L$  is large and requests for VMs of type  $j$  arrive at rate  $\lambda_j L$ ,  $j = 1, \dots, J$ , and each requires service with mean duration  $1/\mu_j$ . The (normalized) load of the system is defined as  $\rho := (\lambda_j/\mu_j, j = 1, \dots, J)$ . This is the *heavy-traffic* regime, e.g. [26, 27, 31, 33, 42, 56, 58], and it has been shown that algorithms with good performance in such a regime also show good performance in other regimes. The interesting scenario occurs when not all VM requests can be scheduled (e.g.,  $\rho > \rho_c$  for a *critical load*  $\rho_c$  on the boundary of system capacity), in which case a fraction of the traffic has to be rejected even by the optimal policy. We propose an adaptive reservation policy that makes admission and packing decisions without the knowledge of  $\rho$ . Packing decisions include placement of admitted VM in one of the feasible servers, and migration of at most one VM across servers when a VM finishes its service.

## 1.1 Related Work

There is classical work on large loss networks, e.g. [7, 26, 27, 33], where calls with different bandwidth requirements and priorities arrive to a telecommunication network. Trunk reservation has been shown to be a robust and effective call admission policy in this setting, in which each call type is accepted if the residual link bandwidth is above a certain threshold for that type. The performance of trunk reservation policies has been analyzed in the asymptotic regime where the call arrival rates and link's capacity scale up by a factor  $N$ , as  $N \rightarrow \infty$ . This is different from our large-scale server model, where the server's capacity is "fixed" and only the number of servers scales (a.k.a. system scale-out as opposed to scale-up). This makes the problem significantly more difficult, because, due to resource fragmentation when packing VMs in servers, the resources of servers *cannot* be viewed as one giant pool; hence our policy not only needs to make admission decisions, but also decide in which server to place the admitted VM. Moreover, VMs have multi-dimensional resource as opposed to one-dimensional calls (bandwidth). If we restrict that every server can fit exactly one VM, our policy reduces to classical trunk reservation.

There has been past work on VM allocation [36, 37, 43, 44, 50, 59] and stochastic bin packing [18, 24, 51, 53, 54], however their models or objectives are different from ours. The works [36, 37, 43, 44] consider a queueing model where VM requests are placed in a queue and then served by the system. In this paper, we are considering a loss model without delay, i.e., each VM request upon arrival has to be served immediately, otherwise it is lost. The recent works [18, 51, 53, 54] study a system with an infinite number of servers and their objective is to minimize the number of occupied servers. The auto-scaling algorithm proposed in [23] also assumes such an infinite server model. These are different from our setting where we consider a finite number of servers and study the total reward of served VMs by the system, in the limit as the number of servers becomes large. In this regime, we have to address complex fluid limit behaviors, especially when the load is above the system capacity and VMs have different priorities.

The works [31, 42, 52, 58] study the blocking probability in a large-scale server system where *all VMs have the same reward*. The work [52] assumes a subcritical system load and only shows local

stability of fluid limits. The works [31, 42, 58] show that, under a power-of-d choices routing, the blocking probability drops much faster compared to the case of uniform random routing. However, there is no analysis of optimality, especially in a supercritical regime where even the optimal policy has a non-zero blocking probability. Moreover, such algorithms treat all VMs with the same priority (reward) when making decisions, thus a low priority VM can potentially block multiple high priority ones.

We remark that in real clouds, servers are monitored periodically [1, 17, 48], for resource management, security, recovery, billing, etc., hence scheduling decisions can be made based on available information about the global system state.

## 1.2 Contributions

We propose a dynamic resource reservation policy that makes admission and packing decisions based on the current system state, and prove that it asymptotically achieves *at least*  $1/2$ , and under certain monotone property on VMs' rewards and resources, at least  $1 - \frac{1}{e}$  of the optimal expected reward, as the number of servers  $L \rightarrow \infty$ . Further, simulations suggest that for real cloud VM instances, the achieved ratio is in fact very close to one.

The main features of our policy and analysis technique can be summarized as follows:

**Dynamic Reservation.** The policy *reserves slots* for VMs in advance. A slot for a VM type will reserve the VM's required resources on a specific server. An incoming VM request then will be admitted if there is enough reservation in the system, in which case it will fill an empty slot of that type. The policy effectively tracks a low-complexity greedy packing of existing VM requests in the system while maintaining only a small number  $g(L) = \omega(\log L)$  of empty slots (e.g.,  $(\log L)^{1+\epsilon}$ ), for VM types that have high priority at the current time. The reservation policy is *robust* and can automatically adapt to changes in the workload based on requests in the system and new arriving requests, without the knowledge of  $\rho$ .

**Analysis Technique.** Our proofs rely on analysis of fluid limits under the proposed policy, however, a major difficulty happens when the workload is above the critical load. In this regime the slot reservation process evolves at a much faster time-scale compared to the fluid-limit processes of the number of VMs and number of servers in different packing configurations in the system. To describe the behavior of fluid limits, we devise a careful analysis based on averaging the behavior of fluid-scale process over *small intervals of length*  $\omega(\log L/L)$ . We then introduce a *Lyapunov function* based on a Linear Program. It is designed to have a unique maximizer at a global greedy solution and determines the convergence properties of our policy in steady state.

## 1.3 Notations

For two positive-valued functions  $x(n)$  and  $y(n)$ , with  $n \in \mathbb{N}$ , we write  $x(n) = o(y(n))$  if  $\lim_{n \rightarrow \infty} x(n)/y(n) = 0$ , and  $x(n) = \omega(y(n))$  if  $y(n) = o(x(n))$ .  $\mathbf{1}(\cdot)$  is the indicator function.  $e_j$  denotes the  $j$ -th basis vector.  $t^-$  and  $t^+$  denote the times right before and after  $t$ .  $\mathbb{R}_+$  is the set of nonnegative real numbers.  $(\cdot)^+ = \max\{\cdot, 0\}$ .

## 2 MODEL AND DEFINITIONS

**Cloud Model.** We consider a collection of  $L$  servers denoted by the set  $\mathcal{L}$ . Each server  $\ell \in \mathcal{L}$  has a limited capacity on different resource types (CPU, memory, disk, networking, etc.). We assume there are  $n \geq 1$  types of resource.

**VM Model.** There is a collection of VM types denoted by the set  $\mathcal{J}$ . The VM types are indexed in arbitrary order from 1 to  $J$ . Each VM type  $j$  requires a vector of resources  $\mathbf{R}_j = (R_j^1, \dots, R_j^n)$ , where  $R_j^d$  is its requirement for the  $d$ -th resource,  $d = 1, \dots, n$ .

VMs are placed in servers and reserve the required resources. The sum of reserved resources by the VMs placed in a server should not exceed the server's capacity. A vector  $\mathbf{k} = (k_1, \dots, k_J) \in \mathbb{Z}_+^J$  is said to be a feasible configuration if the server can simultaneously accommodate  $k_1$  VMs of type 1,  $k_2$  VMs of type 2,  $\dots$ ,  $k_J$  VMs of type  $J$ . We use  $\mathcal{K}$  to denote the set of all feasible configurations (including the empty configuration  $\mathbf{0}_J$ ). The number of feasible configurations will be denoted by  $C := |\mathcal{K}|$ .

We define  $\mathcal{K}^{\mathcal{J}'}$  to be the set of feasible configurations that include only VMs from a subset of types  $\mathcal{J}' \subseteq \mathcal{J}$ , i.e.,

$$\mathcal{K}^{\mathcal{J}'} = \{\mathbf{k} \in \mathcal{K} : k_j = 0, \forall j \notin \mathcal{J}'\}. \quad (1)$$

We use  $K < \infty$  to denote the maximum number of VMs that can fit in a server. We use  $\mathbf{k}^\ell(t) = \mathbf{k}$  to denote that at time  $t$ , server  $\ell \in \mathcal{L}$  has configuration  $\mathbf{k}$ .

We do not necessarily need the resource requirements to be additive, only the monotonicity of the feasible configurations is sufficient, namely, if  $\mathbf{k} \in \mathcal{K}$ , and  $\mathbf{k}' \leq \mathbf{k}$  (component-wise), then  $\mathbf{k}' \in \mathcal{K}$ . This will allow sub-additive resources as well, when the cumulative resource used by the VMs in a configuration could be less than the sum of the resources used individually [46].

**Job and Reward Model.** Jobs for various VM types arrive to the system over time. We can consider two models for jobs:

(i) *Revenue interpretation:* a job of type  $j$  is a request to create a new VM of type  $j$ .

(ii) *Service interpretation:* a job of type  $j$  is a request that must be served by an existing VM of type  $j$  in the system.

To simplify the formulations and use one model to capture both interpretations, we assume that each VM can serve at most one job at any time. As we will see, our algorithm works based on creating "reserved VM slots" in advance. Hence, serving a newly arrived type- $j$  job can be interpreted as deploying a VM of type  $j$  in its reserved slot (revenue interpretation), or assigning it to an already deployed VM of type  $j$  in the slot (service interpretation).

Each job type  $j$  is associated with a reward  $u_j$  which represents its priority (service interpretation) or price paid per time unit of service (revenue interpretation).

We define the *feasible job placement*  $\hat{\mathbf{k}} = (\hat{k}_1, \dots, \hat{k}_J)$  to be the set of jobs that are simultaneously being served in a single server, where  $\hat{k}_j$  corresponds to the number of type- $j$  jobs. Note that by the definition of server configuration, it holds that  $\hat{\mathbf{k}} \leq \mathbf{k}$ , for some  $\mathbf{k} \in \mathcal{K}$ . Hence,  $\mathbf{k} - \hat{\mathbf{k}}$  can be viewed as the reserved VM slots, where  $k_j - \hat{k}_j$  is the number of reserved type- $j$  VM slots. We use  $\hat{\mathbf{k}}^\ell(t) = \hat{\mathbf{k}}$ , when at time  $t$ , the job placement in server  $\ell \in \mathcal{L}$  is  $\hat{\mathbf{k}}$ .

**Traffic Model.** Jobs of type  $j$  arrive according to a Poisson process of rate  $\lambda_j L$ , for a constant  $\lambda_j > 0$ . Once scheduled in a server (more accurately, in a reserved slot of type  $j$ ), a job of type  $j$  requires

an exponentially distributed service time with mean  $1/\mu_j$ , and generates *reward* at rate  $u_j$  during its service. We define the normalized workload of type- $j$  jobs as  $\rho_j := \lambda_j/\mu_j$  and the workload vector  $\boldsymbol{\rho} = (\rho_j, j \in \mathcal{J})$ .

*Definition 2.1 (Configuration Reward).* The reward  $U(\mathbf{k})$  of a configuration  $\mathbf{k} \in \mathcal{K}$  is defined as its total reward per unit time when its slots are full, i.e.,  $U(\mathbf{k}) := \sum_{j=1}^J u_j k_j$ .

*Definition 2.2 (Configuration Ordering).* For two vectors  $\mathbf{k}, \mathbf{k}' \in \mathcal{K}$ , we say  $\mathbf{k} > \mathbf{k}'$ , if either  $U(\mathbf{k}) > U(\mathbf{k}')$ , or  $U(\mathbf{k}) = U(\mathbf{k}')$  and considering the smallest  $j$  for which  $k_j \neq k'_j$ ,  $k_j > k'_j$ .

*Definition 2.3 (MaxReward).* Given a subset  $\mathcal{K}_s \subseteq \mathcal{K}$ , the maximum reward configuration of  $\mathcal{K}_s$  is defined as

$$\text{MAXREWARD}(\mathcal{K}_s) := \arg \max_{\mathbf{k} \in \mathcal{K}_s} U(\mathbf{k}),$$

where ties are broken based on the ordering in Definition 2.2.

*Definition 2.4 (State Variables).* Consider the system with  $L$  servers. We use  $X_{\mathbf{k}}^L(t)$  to denote the number of servers assigned to configuration  $\mathbf{k} \in \mathcal{K}$  at time  $t$ . To distinguish between servers assigned to the same configuration  $\mathbf{k}$ , we *index* them from 1 to  $X_{\mathbf{k}}^L(t)$ , starting from the most recent server assigned to  $\mathbf{k}$  (without loss of generality).

The system state at time  $t$  can then be described as

$$\mathbf{S}^L(t) := ((\mathbf{k}^\ell(t), \hat{\mathbf{k}}^\ell(t), c^\ell(t)), \ell \in \mathcal{L}), \quad (2)$$

where for each server  $\ell \in \mathcal{L}$ ,  $\mathbf{k}^\ell(t) \in \mathcal{K}$  is its configuration,  $\hat{\mathbf{k}}^\ell(t)$ , with  $\hat{\mathbf{k}}^\ell(t) \leq \mathbf{k}^\ell(t)$ , is its job placement, and  $c^\ell(t)$  is its index among the servers with configuration  $\mathbf{k}^\ell(t)$ .

The number of jobs of type  $j$  in the system at time  $t$  is given by

$$Y_j^L(t) = \sum_{\ell \in \mathcal{L}} \hat{k}_j^\ell(t). \quad (3)$$

We also define the vectors  $\mathbf{Y}^L(t) = (Y_j^L(t), j \in \mathcal{J})$ , and  $\mathbf{X}^L(t) = (X_{\mathbf{k}}^L(t), \mathbf{k} \in \mathcal{K})$ . Clearly  $\sum_{\mathbf{k} \in \mathcal{K}} X_{\mathbf{k}}^L(t) = L$  since there are  $L$  servers.

**Optimization Objective.** Given a Markov policy  $\pi$ , we define the expected reward of the policy per unit time as

$$F^\pi(L) = \lim_{t \rightarrow \infty} \mathbb{E} \left[ \sum_{j \in \mathcal{J}} Y_j^L(t) u_j \right]. \quad (4)$$

Our goal is to maximize the expected reward, i.e.,

$$\text{maximize}_\pi F^\pi(L), \quad (5)$$

where the maximization is over all Markov scheduling policies  $\pi$ . Hence, when jobs are requests for VMs, this optimization is a revenue maximization, whereas when jobs are requests to be served by existing VMs, it is a weighted QoS maximization where each service is weighted by its priority.

Note that under any Markov policy, the system state  $\mathbf{S}^L(t)$  is a continuous-time irreducible Markov chain over a finite state space, hence it is positive recurrent and (4) is well defined. Let  $\mathbf{X}^L(\infty)$  and  $\mathbf{Y}^L(\infty)$  be random vectors with the stationary distributions of  $\mathbf{X}^L(t)$  and  $\mathbf{Y}^L(t)$ , respectively, as  $t \rightarrow \infty$ . Note that if  $\mathbf{Y}^*(t)$  is the number of jobs in an  $M/M/\infty$  system in which every job is admitted, then  $\mathbf{Y}^L(\infty)$  is stochastically dominated by  $\mathbf{Y}^*(\infty)$  whose stationary distribution is Poisson with mean  $L\boldsymbol{\rho}$  [10].

We study the problem (5) in the asymptotic regime where the number of servers  $L \rightarrow \infty$ , while the job arrival rates are  $\lambda_j L, j \in \mathcal{J}$ . Note that we do not make any assumption on the values of  $\rho_j$ .

Notice that as  $t \rightarrow \infty$ , the scaled stationary random variables satisfy  $\frac{1}{L} \mathbf{X}^L(\infty) \leq \mathbf{1}$  and  $\frac{1}{L} \mathbf{Y}^L(\infty) \leq \frac{1}{L} \mathbf{Y}^*(\infty)$ . This implies that the sequence of scaled random variables is tight [8], therefore the (random) limits  $\mathbf{x}(\infty) := \lim_{L \rightarrow \infty} \frac{1}{L} \mathbf{X}^L(\infty)$ , and  $\mathbf{y}(\infty) := \lim_{L \rightarrow \infty} \frac{1}{L} \mathbf{Y}^L(\infty)$  exist along a subsequence of  $L$ . The limits satisfy  $x_k(\infty) \geq 0$ ,  $\sum_{k \in \mathcal{K}} x_k(\infty) = 1$ , and  $\mathbf{y}(\infty) \leq \boldsymbol{\rho}$ ,  $\mathbf{y}(\infty) \leq \sum_{k \in \mathcal{K}} x_k(\infty) \mathbf{k}$ .

To unify the algorithm descriptions for revenue maximization and QoS maximization, in the rest of the paper, we use the term “slot” of type  $j$  to refer to the resource (equal to a VM of type  $j$ ) reserved for one job of type  $j$  in a server. Filled slots have jobs already in them, while empty slots could accept jobs. Therefore, the term configuration applies to all the slots in a server, while placement applies to the filled slots in the server.

### 3 A STATIC OPTIMIZATION AND ITS GREEDY SOLUTION

Given a workload reference vector  $\hat{\mathbf{Y}}^L = (\hat{Y}_j^L, j \in \mathcal{J})$ , let  $F^*(L, \hat{\mathbf{Y}}^L)$  be the optimal value of the following linear program:

$$\max_{\mathbf{X}, \mathbf{Y}} \sum_j u_j Y_j \quad (6a)$$

$$\text{s.t. } Y_j \leq \hat{Y}_j^L, \forall j \in \mathcal{J} \quad (6b)$$

$$\sum_{k \in \mathcal{K}} X_k k_j \geq Y_j, \forall j \in \mathcal{J} \quad (6c)$$

$$\sum_{k \in \mathcal{K}} X_k = L, \quad X_k \geq 0, \forall k \in \mathcal{K} \quad (6d)$$

where  $\mathbf{Y}$  is the vector of jobs in the system, and  $\mathbf{X}$  is the vector of the number of servers assigned to each configuration. If we choose  $\hat{\mathbf{Y}}^L = \boldsymbol{\rho}L$ , this optimization will provide an upper bound on optimization (5), i.e.,  $F^\pi(L) \leq F^*(L, \boldsymbol{\rho}L)$ , for any Markov policy  $\pi$ . The interpretation of the result is as follows. The average number of type- $j$  jobs in the system cannot be more than its workload (Constraint (6b)), and further, it cannot be more than the average number of slots of type  $j$  in the servers (Constraint (6c)). The sum of number of servers in different configurations is  $L$ , so their average should also satisfy (6d).

As  $L \rightarrow \infty$ , the normalized objective value  $\frac{1}{L} F^*(L, \boldsymbol{\rho}L) \rightarrow U^*[\boldsymbol{\rho}]$ , which is the optimal value of the linear program below

$$\max_{\mathbf{x}, \mathbf{y}} \sum_j u_j y_j \quad (7a)$$

$$\text{s.t. } y_j \leq \rho_j, \forall j \in \mathcal{J} \quad (7b)$$

$$\sum_{k \in \mathcal{K}} k_j x_k \geq y_j, \forall j \in \mathcal{J} \quad (7c)$$

$$\sum_{k \in \mathcal{K}} x_k = 1, \quad x_k \geq 0, \forall k \in \mathcal{K} \quad (7d)$$

where  $x_k$  can be interpreted as the ideal fraction of servers which should be in configuration  $\mathbf{k}$  when  $L$  is large. Hence, one can consider a static reservation policy where the cloud cluster is partitioned and  $\lfloor x_k L \rfloor$  servers are assigned to each non-zero configuration  $\mathbf{k} \in \mathcal{K}$  (and the rest of servers can be empty to save resource

or used to serve more jobs). Then once a type- $j$  job arrives, it will be routed to an empty slot of type  $j$  in one of the servers, if any, otherwise it is rejected. This will provide an asymptotic optimal policy since it achieves the normalized reward  $U^*[\boldsymbol{\rho}]$ , as  $L \rightarrow \infty$ .

However, there are several issues with this approach: (i) solving optimization (6) or its relaxation (7) has a very high complexity, as the number of configurations is exponential in the number of job types  $J$ , and (ii) it requires knowing an accurate estimate of the workload  $\boldsymbol{\rho}$  which might not be available. Inaccurate estimates of workload can lead to poor performance for such policies, e.g., see [34] which illustrates that static reservation policies in classical loss networks can give very poor performance. Even if we have an estimate of the workload and approximate the solution to (7), to handle time-varying workloads, the new solution may require rearranging a large number of VMs and jobs to make their placements match the new solution. This is costly and also causes interruption of many jobs in service.

We first address the complexity issue, by presenting a greedy solution for the optimization, and analyze its asymptotic performance below.

#### 3.1 Greedy Solution

We describe a greedy algorithm, called *Greedy Placement Algorithm* (GPA), for solving optimization (6).

GPA takes as input the workload reference vector  $\hat{\mathbf{Y}}^L$ , and returns an assignment vector  $\hat{\mathbf{X}}^L$  which indicates which configurations should be used and in how many servers. The assignment consists of at most  $J$  configurations, which are found in  $J$  iterations. In each iteration  $i$ , GPA maintains a set of candidate job types  $\mathcal{J}[i]$ , and finds a configuration  $\mathbf{k}[i]$ . Initially  $\mathcal{J}[1] = \mathcal{J}$ . In iteration  $i$ :

- (1) It finds  $\mathbf{k}[i] = \text{MAXREWARD}(\mathcal{K}^{\mathcal{J}[i]})$ , which is the configuration of highest reward among the configurations that have jobs from the set  $\mathcal{J}[i]$ , according to Definition 2.3.
- (2) It computes the number of servers  $\hat{X}_{\mathbf{k}[i]}^L$  that should be assigned to  $\mathbf{k}[i]$ , until at least one of the job types  $j$ , for which  $k_j[\mathbf{k}[i]] > 0$ , has no more jobs left, or there are no more unused servers left. We refer to this job type as  $j^*$ .
- (3) It then creates  $\mathcal{J}[i+1]$  by removing job type  $j^*$  from  $\mathcal{J}[i]$ .

A pseudocode for GPA is given by Algorithm 1. We use the vector  $\hat{\mathbf{X}}^L = (\hat{X}_{\mathbf{k}}^L, \mathbf{k} \in \mathcal{K})$  to denote the output of GPA, which has at most  $J$  non-zero elements corresponding to  $\mathbf{k}[i], i = 1, \dots, J$ .

*Remark 1.* MAXREWARD finds the maximum reward configuration of a subset of job types, which is equivalent with *unbounded* Knapsack problem (unbounded number of items for each type). This problem is tractable with Pseudopolynomial algorithms to solve it exactly [3, 40] or fully polynomial approximation algorithms [28]. GPA needs to solve at most  $J$  instances of this problem. Note that the number of different instances of the problem is bounded and we can compute MAXREWARD for all of them offline as *they are not workload dependent*. This is in contrast to optimization (6), which is equivalent to multi Knapsack problem which is strongly NP-hard [32], and requires resolving when workload reference  $\hat{\mathbf{Y}}$  changes.

We next define the limit of  $\hat{\mathbf{X}}^L/L$  for input  $\hat{\mathbf{Y}}^L = L\boldsymbol{\rho}$ , as  $L \rightarrow \infty$ , which we refer to as *Global Greedy Assignment*. To describe

---

**Algorithm 1** Greedy Placement Algorithm (GPA)

---

```

1: function GPA( $\hat{Y}$ )
2:    $\mathbf{r} \leftarrow \hat{Y}$             $\triangleright$  tracks the vector of number of jobs left
3:    $N \leftarrow L$             $\triangleright$  tracks the number of servers left
4:    $i \leftarrow 1, \mathcal{J}[1] = \mathcal{J}$ 
5:   while  $\mathcal{J}[i] \neq \emptyset$  do
6:      $\mathbf{k}[i] \leftarrow \text{MAXREWARD}(\mathcal{K}^{\mathcal{J}[i]})$ 
7:      $j^* \leftarrow \arg \min_{j: k_j[i] > 0} \lceil \frac{r_j}{k_j[i]} \rceil$     $\triangleright$  break ties arbitrarily
8:      $\hat{X}_{\mathbf{k}[i]} \leftarrow \min \left( \lceil \frac{r_{j^*}}{k_{j^*}[i]} \rceil, N \right)$ 
9:      $\mathbf{r} \leftarrow \mathbf{r} - \hat{X}_{\mathbf{k}[i]} \mathbf{k}[i]$ 
10:     $N \leftarrow N - \hat{X}_{\mathbf{k}[i]}$ 
11:     $\mathcal{J}[i+1] \leftarrow \mathcal{J}[i] - \{j^*\}$ 
12:     $i \leftarrow i + 1$ 
13:  return  $\hat{X}_{\mathbf{k}[j]}, j = 1, \dots, J$ 

```

---

this assignment, we first define a unique ordering of the job types through the following proposition.

**PROPOSITION 3.1.** *For any permutation  $\sigma = (\sigma_1, \sigma_2, \dots, \sigma_J)$  of job types in  $\mathcal{J}$ , let  $\mathcal{J}_j^\sigma := \{\sigma_j, \dots, \sigma_J\}$ , and  $\mathbf{k}^{(j)} := \text{MAXREWARD}(\mathcal{K}^{\mathcal{J}_j^\sigma})$ . Given a workload  $\rho$ , there is a “unique” permutation  $\sigma = (\sigma_1, \sigma_2, \dots, \sigma_J)$  of job types, such that the following holds:*

1)  $\forall j \in \mathcal{J}, k_{\sigma_j}^{(j)} > 0$ , and there are constants  $z^{(j)}[\rho] \geq 0$ , such that

$$\rho_{\sigma_j} = \sum_{\ell=1}^j k_{\sigma_j}^{(\ell)} z^{(\ell)}[\rho], \quad (8)$$

2) for any two indexes  $j, j' \in \mathcal{J}$ , with  $j < j'$ , if

$$\rho_{\sigma_{j'}} = \sum_{\ell=1}^j k_{\sigma_{j'}}^{(\ell)} z^{(\ell)}[\rho], \quad (9)$$

then we should have  $\sigma_j < \sigma_{j'}$ .

**PROOF.** See Appendix A.  $\square$

The Global Greedy Assignment is defined as follows

**Definition 3.2 (Global Greedy Assignment).** Define the index  $I_\rho \leq J$  for which

$$\sum_{i=1}^{I_\rho-1} z^{(i)}[\rho] < 1, \quad \sum_{i=1}^{I_\rho} z^{(i)}[\rho] \geq 1,$$

with the convention that  $I_\rho = J + 1$  if  $\sum_{i=1}^J z^{(i)}[\rho] < 1$ . The *global greedy assignment*  $\mathbf{x}^{(g)}[\rho]$  is defined as

$$\mathbf{x}_{\mathbf{k}^{(i)}}^{(g)}[\rho] = \begin{cases} z^{(i)}[\rho], & \text{for } i < I_\rho \\ 0, & \text{for } i > I_\rho \\ 1 - \sum_{j=1}^{i-1} x_{\mathbf{k}^{(j)}}^{(g)}[\rho], & \text{for } i = I_\rho, \end{cases} \quad (10)$$

where  $\mathbf{k}^{(i)}$  and  $z^{(i)}[\rho]$ ,  $i = 1, \dots, J$ , were defined in Proposition 3.1, and  $\mathbf{k}^{(J+1)} := \mathbf{0}$  (empty configuration). We call the ordered configurations  $\mathbf{k}^{(i)}$ ,  $i = 1, \dots, J + 1$ , the “global greedy configurations” of workload  $\rho$ . For any configuration  $\mathbf{k} \in \mathcal{K}$  not in global greedy configurations,  $x_{\mathbf{k}}^{(g)}[\rho] = 0$ . When it is clear from the context, the dependency  $[\rho]$  will be omitted.

Since global greedy configurations  $\mathbf{k}^{(\ell)}$ ,  $\ell = 1, \dots, J + 1$ , depend on  $\rho$ , the following configurations will come in handy when the analysis needs to be agnostic to  $\rho$ .

**Definition 3.3 (Greedy Configurations).** The greedy configuration set  $\mathcal{K}^{(g)}$  includes all configurations that are output of  $\text{MAXREWARD}(\mathcal{K}^{\mathcal{J}'})$  for any  $\mathcal{J}' \subseteq \mathcal{J}$ . That is the set of all possible configurations which may be assigned by GPA, and the empty configuration. We define  $C^{(g)} := |\mathcal{K}^{(g)}|$ . We enumerate configurations of  $\mathcal{K}^{(g)}$  as  $\bar{\mathbf{k}}^{(i)}$ , for  $i = 1, \dots, C^{(g)}$ , such that  $\bar{\mathbf{k}}^{(i_1)} > \bar{\mathbf{k}}^{(i_2)}$  if  $i_1 < i_2$  (according to Definition 2.2), and  $\bar{\mathbf{k}}^{(C^{(g)})} = \mathbf{0}_J$ .

Notice that  $\{\mathbf{k}^{(j)}, j = 1, \dots, J + 1\} \subseteq \{\bar{\mathbf{k}}^{(i)}, i = 1, \dots, C^{(g)}\}$ , and their order is consistent with Definition 2.2, as defined below.

**Definition 3.4 (Mapping global greedy to greedy).** For any  $j, j' \in \{1, \dots, J + 1\}$ , with  $j < j'$ , there are indexes  $g_j, g_{j'} \in \{1, \dots, C^{(g)}\}$ , such that  $\mathbf{k}^{(j)} \equiv \bar{\mathbf{k}}^{(g_j)}$ ,  $\mathbf{k}^{(j')} \equiv \bar{\mathbf{k}}^{(g_{j'})}$ , and  $g_j < g_{j'}$ . We also define  $C_\rho^{(g)} := g_{I_\rho}$  to be the index for which  $\mathbf{k}^{(I_\rho)} \equiv \bar{\mathbf{k}}^{(C_\rho^{(g)})}$ .

The following proposition states the connection between GPA and Global Greedy Assignment  $\mathbf{x}_{\mathbf{k}}^{(g)}[\rho]$ .

**PROPOSITION 3.5.** *Let  $\hat{X}^L = \text{GPA}(L\rho)$ . Then*

$$\lim_{L \rightarrow \infty} \frac{\hat{X}_{\mathbf{k}}^L}{L} = x_{\mathbf{k}}^{(g)}[\rho], \quad \forall \mathbf{k} \in \mathcal{K}, \quad (11)$$

where  $x_{\mathbf{k}}^{(g)}[\rho]$  is the Global Greedy Assignment of Definition 3.2.

**PROOF.** See Appendix B.  $\square$

Note that clearly  $\mathbf{x}^{(g)}[\rho]$  is a feasible solution for optimization (7) and it is easy to see that its corresponding objective value is

$$U^{(g)}[\rho] := \sum_{j=1}^J u_j \sum_{\ell=1}^J k_j^{(\ell)} x_{\mathbf{k}^{(\ell)}}^{(g)}[\rho]. \quad (12)$$

It is also easy to see that in optimization (7) we can replace the inequality in (7c) with equality and the optimal value will not change. Let  $\mathbf{x}^*[\rho]$  be one such optimal solution to optimization (7) for workload  $\rho$ . Then the optimal objective value is

$$U^*[\rho] := \sum_{j \in \mathcal{J}} u_j \sum_{\mathbf{k} \in \mathcal{K}} k_j x_{\mathbf{k}}^*[\rho]. \quad (13)$$

The following corollary is immediate from Proposition 3.5.

**COROLLARY 3.6.** *Let  $F^{GPA}(L, \rho L)$  be the total reward of GPA in the system with  $L$  servers given reference workload  $\hat{Y}^L = \rho L$ . Then*

$$\lim_{L \rightarrow \infty} \frac{F^{GPA}(L, \rho L)}{F^*(L, \rho L)} = \frac{U^{(g)}[\rho]}{U^*[\rho]}.$$

The theorem below bounds the above ratio.

**THEOREM 3.7.** *The global greedy assignment  $\mathbf{x}^{(g)}[\rho]$  provides at least  $\frac{1}{2}$  of the optimal normalized reward, i.e.,  $\frac{U^{(g)}[\rho]}{U^*[\rho]} \geq \frac{1}{2}, \forall \rho \geq 0$ .*

PROOF. Consider the permutation of job types according to Proposition 3.1. By the global greedy definition and the feasibility of  $\mathbf{x}^*[\boldsymbol{\rho}]$ , for any job type  $\sigma_j$ ,  $j = 1, \dots, I_\rho - 1$ , we have

$$\sum_{\mathbf{k} \in \mathcal{K}} x_{\mathbf{k}}^* k_{\sigma_j} \leq \sum_{\ell=1}^j x_{\mathbf{k}^{(\ell)}}^{(g)} k_{\sigma_j}^{(\ell)} = \rho_{\sigma_j}, \quad (14)$$

from which it follows that

$$\sum_{j=1}^{I_\rho-1} \sum_{\mathbf{k} \in \mathcal{K}} x_{\mathbf{k}}^* k_{\sigma_j} u_{\sigma_j} \leq \sum_{j=1}^{I_\rho-1} \sum_{\ell=1}^j x_{\mathbf{k}^{(\ell)}}^{(g)} k_{\sigma_j}^{(\ell)} u_{\sigma_j} = \sum_{j=1}^{I_\rho-1} \rho_{\sigma_j} u_{\sigma_j} \leq U^{(g)}[\boldsymbol{\rho}]. \quad (15)$$

Also for the job types  $\sigma_j$ , for  $j = I_\rho, \dots, J$ , we have

$$\begin{aligned} \sum_{j=I_\rho}^J \sum_{\mathbf{k} \in \mathcal{K}} x_{\mathbf{k}}^* k_{\sigma_j} u_{\sigma_j} &= \sum_{\mathbf{k} \in \mathcal{K}} x_{\mathbf{k}}^* \sum_{j=I_\rho}^J k_{\sigma_j} u_{\sigma_j} \leq \\ \arg \max_{\mathbf{k} \in \mathcal{K}} \sum_{j=I_\rho}^J k_{\sigma_j} u_{\sigma_j} &\stackrel{(b)}{=} \sum_{j=I_\rho}^J k_{\sigma_j}^{(I_\rho)} u_{\sigma_j} \stackrel{(c)}{\leq} U^{(g)}[\boldsymbol{\rho}]. \end{aligned} \quad (16)$$

where (a) is due to the fact that  $\sum_{\mathbf{k} \in \mathcal{K}} x_{\mathbf{k}}^* = 1$ , (b) is by the definition of  $\mathbf{k}^{(I_\rho)}$ , and (c) is because  $U^{(g)}[\boldsymbol{\rho}]$  is a convex combination of rewards of  $\mathbf{k}^{(1)}, \dots, \mathbf{k}^{(I_\rho)}$ , which all have a reward no less than that of  $\mathbf{k}^{(I_\rho)}$ . Then adding (15) and (16), we get

$$U^*[\boldsymbol{\rho}] = \sum_{j=1}^J \sum_{\mathbf{k} \in \mathcal{K}} x_{\mathbf{k}}^* k_{\sigma_j} u_{\sigma_j} \leq 2U^{(g)}[\boldsymbol{\rho}]. \quad \square$$

Theorem 3.7 can be improved when job types and rewards satisfy a monotone greedy property described next.

*Definition 3.8.* We say the job types and the rewards have *monotone greedy* property if for any two instances of the optimization (7) with  $\boldsymbol{\rho}_1 \geq \boldsymbol{\rho}_2$ ,  $U^{(g)}[\boldsymbol{\rho}_1] \geq U^{(g)}[\boldsymbol{\rho}_2]$ .

It is easy to verify that any system with two job types always has the property in Definition 3.8. However, in general the property depends on the profile of jobs types and their rewards, and might not hold for adversarial profiles. The next theorem describes the improved bound when the monotone greedy property holds.

**THEOREM 3.9.** *If job types and rewards satisfy the monotone greedy property, then, for any  $\boldsymbol{\rho}$ ,  $\frac{U^{(g)}[\boldsymbol{\rho}]}{U^*[\boldsymbol{\rho}]} \geq 1 - 1/e$ .*

PROOF. Define a workload  $\boldsymbol{\rho}^* := \sum_{\mathbf{k} \in \mathcal{K}} \mathbf{k} x_{\mathbf{k}}^*[\boldsymbol{\rho}]$ . We notice that  $U^*[\boldsymbol{\rho}] = U^*[\boldsymbol{\rho}^*]$  in LP (7). Also by the monotone greedy property,  $U^{(g)}[\boldsymbol{\rho}] \geq U^{(g)}[\boldsymbol{\rho}^*]$ , since  $\boldsymbol{\rho} \geq \boldsymbol{\rho}^*$ . Hence, it suffices to prove the theorem for instances where  $\boldsymbol{\rho} = \boldsymbol{\rho}^*$  or in other words, instances for which, in the optimal solution, workload fits exactly in servers.

Consider now the projection of the workload  $\boldsymbol{\rho}^* = \boldsymbol{\rho}$  onto the global greedy configuration space  $\{\mathbf{k}^{(i)}[\boldsymbol{\rho}], i = 1, \dots, J\}$ . Since these configurations are independent, we can write

$$\boldsymbol{\rho}^* = \boldsymbol{\rho} = \sum_{i=1}^J z^{(i)}[\boldsymbol{\rho}] \mathbf{k}^{(i)}, \quad (17)$$

for  $z^{(i)}[\boldsymbol{\rho}]$  introduced in Proposition 3.1. For notational compactness, define  $q_i = z^{(i)}[\boldsymbol{\rho}]$ ,  $i = 1, \dots, J$ , and  $p_i = x_{\mathbf{k}^{(i)}}^{(g)}[\boldsymbol{\rho}]$ ,  $i = 1, \dots, I_\rho$ , and let  $W^{(i)} := U(\mathbf{k}^{(i)}) = \sum_{j=i}^J u_{\sigma_j} k_{\sigma_j}^{(i)}$ .

Then,

$$\begin{aligned} \sum_{j=i}^J q_j W^{(j)} &= \sum_{j=i}^J u_{\sigma_j} \sum_{\ell=i}^j q_\ell k_{\sigma_j}^{(\ell)} \stackrel{(a)}{\leq} \sum_{j=i}^J \rho_{\sigma_j} u_{\sigma_j} \\ &\stackrel{(b)}{\leq} \sum_{j=i}^J k_{\sigma_j}^{(i)} u_{\sigma_j} = W^{(i)}. \end{aligned} \quad (18)$$

Inequality (a) is because  $\sum_{\ell=i}^j q_\ell k_{\sigma_j}^{(\ell)} \leq \rho_{\sigma_j}$ , and Inequality (b) is because we assumed there is an assignment that can completely accommodate workload  $\boldsymbol{\rho}$ , and hence  $\rho_{\sigma_j}$  for  $j = i, \dots, J$ . If we remove all jobs with types  $1, \dots, i-1$  from assignment  $\mathbf{x}^*$ , the configurations used in the resulting assignment belong to the subset  $\mathcal{K}^{\{\sigma_i, \dots, \sigma_J\}}$  and  $\mathbf{k}^{(i)}$  is the configuration with the highest reward from this set.

An equivalent representation of (18) is that, for some constants  $b_i$ ,  $0 \leq b_i \leq 1$ ,  $i = 1, \dots, J$ ,

$$b_i W^{(i)} = \sum_{j=i}^J q_j W^{(j)}, \text{ and } (b_i - q_i) W^{(i)} = b_{i+1} W^{(i+1)}. \quad (19)$$

For completeness, we also define  $b_{J+1} = 1$ . Based on this representation, and using (17) and  $\boldsymbol{\rho} = \boldsymbol{\rho}^*$  by assumption, we get

$$\begin{aligned} \frac{U^{(g)}[\boldsymbol{\rho}]}{U^*[\boldsymbol{\rho}]} &= \frac{\sum_{i=1}^{I_\rho} p_i W^{(i)}}{\sum_{i=1}^J q_i W^{(i)}} = \frac{\sum_{i=1}^{I_\rho} p_i \prod_{j=1}^{i-1} \frac{b_j - q_j}{b_{j+1}} W^{(1)}}{b_1 W^{(1)}} = \\ &\sum_{i=1}^{I_\rho} \frac{p_i}{b_i} \prod_{j=1}^{i-1} \frac{b_j - q_j}{b_j} = 1 - \prod_{i=1}^{I_\rho} \left(1 - \frac{p_i}{b_i}\right). \end{aligned} \quad (20)$$

The right-hand side is minimized if  $b_i = 1$ ,  $i = 1, \dots, I_\rho$ , since  $p_i \geq 0$ . Then given  $\sum_{i=1}^{I_\rho} p_i = 1$ , the expression is minimized for  $p_i = 1/I_\rho$ ,  $i = 1, \dots, I_\rho$ , and its minimum value is  $1 - \left(1 - \frac{1}{I_\rho}\right)^{I_\rho} > 1 - e^{-1}$ .  $\square$

**PROPOSITION 3.10.** *The worst-case ratio of  $U^{(g)}[\boldsymbol{\rho}]/U^*[\boldsymbol{\rho}]$  is not greater than  $1 - 1/e$ .*

PROOF. We construct an adversarial example that achieves this bound. See Appendix C.  $\square$

Hence, the global greedy assignment achieves a factor within  $1/2$  to  $1 - 1/e$  of the optimal normalized reward in “all” the cases. Further, the bound  $1 - 1/e$  is tight when monotone greedy property holds. The assignment might actually achieve  $1 - 1/e$  in all the cases but requires a more careful analysis. In view of Corollary 3.6,  $\text{GPA}(\boldsymbol{\rho}L)$  asymptotically achieves the same factor of the optimal reward. In simulations in Section 7, based on cloud VM instances, we were not able to find any scenario where the ratio is below  $1 - 1/e$ , and in fact the ratio is much better ( $\approx 0.97$  on average).

However,  $\text{GPA}(\boldsymbol{\rho}L)$  requires the knowledge of  $\boldsymbol{\rho}$ . In the next section, we propose a dynamic reservation algorithm that is appropriate for use in online settings without the knowledge of  $\boldsymbol{\rho}$ . Its achievable normalized reward still converges to that of the global greedy assignment and it can also adapt to changes in the workload.

## 4 DYNAMIC RESERVATION ALGORITHM

We present a Dynamic Reservation Algorithm, called DRA, which makes admission decisions and configuration assignments, without the knowledge  $\rho$ . We first introduce the following notations:

- Recall the indexing of servers in the same configuration as in Definition 2.4. We use  $\ell_{k,i}$  to refer to the server with configuration  $k$  and index  $i$ .
- A key parameter of DRA is the *reservation factor*  $g(L)$ . It is the number of empty slots (safety margin) that the algorithm ideally wants to reserve for each job type if possible. For later analysis, we assume that  $g(L) = \omega(\log(L))$ , and is  $o(L)$ .

The configuration assignment occurs at *update times*. To simplify the analysis, we consider update times to be times when a job is admitted to or departs from the system. To avoid preemptions, only servers that are empty (have no jobs running) can be assigned to a new configuration.

At update time  $t$ , DRA updates the workload reference vector  $\hat{Y}^L(t)$  as

$$\hat{Y}^L(t) = Y^L(t) + g(L)\mathbf{1}, \quad (21)$$

where  $Y^L(t)$  is the vector of jobs in the system, *after* any job admission or job departure at time  $t$ .  $g(L)$  is the reservation factor as defined earlier.

Then DRA classifies the servers into two groups: *Accept Group* (AG) and *Reject Group* (RG). Servers in Accept Group keep their current configurations and DRA attempts to have all their slots filled by scheduling new jobs in them, while servers in Reject Group do not have desirable configurations and DRA attempts to make them empty, by not scheduling new jobs in them and possibly migrating their jobs to servers in Accept Group, so they can be reassigned to other configurations.

A pseudocode for DRA is given in Algorithm 2. It has three main components which we describe in detail below:


**Classification and Reassignment Algorithm (CRA).** This is the subroutine used by DRA to classify servers and possibly reassign some of them. It attempts to greedily reduce the disparity between the configuration assignment in the system  $X^L(t)$  and the output of GPA  $\hat{X}^L(t) = \text{GPA}(\hat{Y}^L(t))$ . To do so, it assigns *rank*s to servers in different configurations, which range from 1 to  $J + 1$ .

Initially, all servers are assigned rank  $J + 1$ . Any empty server of rank  $J + 1$  can be reassigned to reduce the disparity between  $X^L(t)$  and  $\hat{X}^L(t)$ . We use  $\ell_e$  to denote one of empty rank  $J + 1$  servers, and if no such server exists  $\ell_e = \emptyset$ .

Iterating over configurations  $k[i]$  found by GPA, for  $i = 1, \dots, J$ :

- If  $X_{k[i]}^L < \hat{X}_{k[i]}^L$ , it increases  $X_{k[i]}^L$  by reassigning any  $\ell_e$  to  $k[i]$ , until either (i) it matches  $\hat{X}_{k[i]}^L$ , or (ii)  $\ell_e = \emptyset$ . In either case, all servers of configuration  $k[i]$  get rank  $i$ .
- If  $X_{k[i]}^L(t) \geq \hat{X}_{k[i]}^L$ , it assigns rank  $i$  to all servers of configuration  $k[i]$  with indexes greater than  $X_{k[i]}^L(t) - \hat{X}_{k[i]}^L(t)$ .

We use  $I^*(t)$  to denote the first  $i$  for which  $X_{k[i]}^L$  cannot be matched to  $\hat{X}_{k[i]}^L$ , i.e. the first  $i$  at which  $\ell_e = \emptyset$ . If all configurations are matched, then  $I^*(t) = J$ . At the end of CRA, servers with rank greater than  $I^*(t)$  and index 1 in any configuration are classified as



**Figure 1: An example illustrating the state at the end of CRA.** Servers in each configuration are stacked from largest to smallest index.  $k[1], \dots, k[J]$  are the configurations returned by GPA. The dashed boxes indicate how many more servers need to be reassigned to a respective configuration to match the solution of GPA (horizontal line).  $I^*$  is the first  $i$  for which  $X_{k[i]}^L < \hat{X}_{k[i]}^L$  at the end of the procedure. Orange servers are the servers of Reject Group.

Reject Group, while the rest of the servers are classified as Accept Group.

See Figure 1 for an illustrative example for the state of CRA.

**Scheduling Arriving Job.** When DRA needs to schedule an arriving job of type  $j$ , it places the job in one of the servers of Accept Group with empty type- $j$  slot. If no such server exists, the job is rejected.

We use  $AG_j$  to denote one of the servers of Accept Group with empty type- $j$  slot. If no such server exists  $AG_j = \emptyset$ .

**Migrating Job after Departure.** Let  $RG_j$  denote the highest rank server among the Reject Group servers with type- $j$  jobs. If no such server exists,  $RG_j = \emptyset$ .

If a type- $j$  job departs from a server in Accept Group, DRA migrates one of the type- $j$  jobs from  $RG_j$  to the slot that emptied because of the departure, if  $RG_j \neq \emptyset$ .

**Initialization.** Initially servers have no indexes or classification (and might not even have configurations), so we need to specify how the system state is initialized (say at time 0) under DRA. If servers do not have configurations, but have jobs in them, we initialize  $k^\ell(0) = \hat{k}^\ell(0)$ , i.e., the configuration of each server  $\ell$  is set to its job placement. If servers have configurations, we keep their existing configuration. Indexing among the servers of a configuration can be arbitrary. We then run CRA that performs classification and reassigns any possibly empty servers.

*Remark 2.* Notice the duality of actions performed on arrivals and departures for any job type: jobs are admitted/migrated to empty slots in servers of Accept Group, and depart/migrate from filled slots in servers of Reject Group. The number of servers in Reject Group under our algorithm is at most one per configuration, i.e., at most  $C^{(g)}$  servers (constant independent of  $L$ ) which is negligible compared to the number of servers  $L$ , as  $L \rightarrow \infty$ . Further, job admissions and migrations are performed to slots which are already deployed in advance. The reservation factor  $g(L)$  is critical for maintaining enough deployed slots in the maximum reward configurations for future demand.

**Algorithm 2** DRA: Dynamic Reservation Algorithm

---

```

1: function CRA( $\hat{Y}^L, \mathbf{X}^L$ )
2:    $\hat{\mathbf{X}}^L \leftarrow \text{GPA}(\hat{Y}^L)$ .
3:   Set rank of all servers to  $J + 1$ .
4:    $I^* \leftarrow J$ 
5:   for  $i = 1$  to  $J$  do            $\triangleright J$  configurations found in GPA
6:      $Z \leftarrow 0, c \leftarrow X_{\mathbf{k}[i]}^L$     $\triangleright c$  is the index of server
7:     while  $Z < \hat{X}_{\mathbf{k}[i]}^L$  do
8:        $Z \leftarrow Z + 1, c \leftarrow c - 1$ 
9:       if  $c \leq 0$  then
10:        if  $\ell_e \neq \emptyset$  then
11:          Set rank of  $\ell_e$  to  $i$ .
12:          Reassign configuration of  $\ell_e$  to  $\mathbf{k}[i]$ .
13:        else
14:           $I^* \leftarrow \min(I^*, i)$ 
15:        else
16:          Set rank of  $\ell_{\mathbf{k}[i], c}$  to  $i$ .
17:   procedure ARRIVAL( $j, t$ )            $\triangleright$  Type- $j$  arrival at time  $t$ 
18:     if  $\text{AG}_j \neq \emptyset$  then
19:       Schedule job in  $\text{AG}_j$ .
20:       CRA( $Y^L(t) + g(L)\mathbf{1}, \mathbf{X}^L(t)$ )
21:     else
22:       Reject job.
23:   procedure DEPARTURE( $j, t$ )          $\triangleright$  Type- $j$  departure at time  $t$ 
24:     if  $\text{RG}_j \neq \emptyset$  and the slot emptied is in Accept Group then
25:       Migrate the job in  $\text{RG}_j$  to the slot that emptied.
26:     CRA( $Y^L(t) + g(L)\mathbf{1}, \mathbf{X}^L(t)$ )

```

---

In contrast, a naive static reservation algorithm, that solves (6) by replacing  $\hat{Y}$  with an estimate of workload, might require changing the configuration of a constant fraction of servers (the equivalent of Reject Group), as workload estimate changes. This would result in preemptions (or migrations) in  $O(L)$  interrupted servers.

Lastly, more accurate estimates of workload, if available, can be simply used in the input  $\hat{Y}$  to CRA, and CRA itself can be executed less regularly, depending on the complexity and convergence time tradeoff.

The following theorem states the main result regarding DRA.

**THEOREM 4.1.** *Let  $F^{\text{DRA}}(L)$  be the expected reward under DRA and  $F^*(L)$  be the optimal expected reward in optimization (5). Then*

$$\lim_{L \rightarrow \infty} \frac{F^{\text{DRA}}(L)}{F^*(L)} \geq \frac{1}{2}.$$

Further, under the monotone greedy property (Definition 3.8),

$$\lim_{L \rightarrow \infty} \frac{F^{\text{DRA}}(L)}{F^*(L)} \geq 1 - \frac{1}{e}.$$

**Remark 3.** Note that we did not make any assumption on the value of  $\rho$ , and Theorem 4.1 holds for any  $\rho$ . Define

$$\Lambda = \left\{ \mathbf{y} : \mathbf{y} \leq \sum_{\mathbf{k} \in \mathcal{K}} x_{\mathbf{k}} \mathbf{k}, \text{ for } x_{\mathbf{k}} \geq 0, \mathbf{k} \in \mathcal{K}, \sum_{\mathbf{k} \in \mathcal{K}} x_{\mathbf{k}} = 1 \right\}. \quad (22)$$

Theorem 4.1 holds even if  $\rho$  is outside  $\Lambda$ . In this scenario, a nonzero fraction of traffic has to be rejected even by the optimal policy.

The proof of Theorem 4.1 is based on analysis of fluid limits and a suitable Lyapunov function to show convergence, as we do next in Sections 5 and 6.

## 5 FLUID LIMITS UNDER DRA

We first define two useful variables, which are functions of the system state, and will be used in our convergence analysis.

*Definition 5.1 (Effective Number of Assigned Servers).* The effective number of servers in configuration  $\mathbf{k}$  is defined as

$$X_{\mathbf{k}}^{L(e)}(t) := \min(X_{\mathbf{k}}^L(t), \hat{X}_{\mathbf{k}}^L(t)). \quad (23)$$

Note that  $X_{\mathbf{k}}^{L(e)}(t) = \hat{X}_{\mathbf{k}}^L(t) = 0$  if  $\mathbf{k} \notin \{\bar{\mathbf{k}}^{(i)}, i = 1, \dots, C^{(g)}\}$ . With a minor abuse of terminology, we say the servers in configuration  $\mathbf{k}$  with indexes from  $X_{\mathbf{k}}^L(t) - X_{\mathbf{k}}^{L(e)}(t) + 1$  to  $X_{\mathbf{k}}^L(t)$ , have effective configuration  $\mathbf{k}$ .

*Remark 4.* Note that  $X_{\mathbf{k}}^{L(e)}(t)$  is *independent* of the indexing of servers in configuration  $\mathbf{k}$ . Also note if  $\mathbf{k} = \mathbf{k}[j]$ , where  $\mathbf{k}[j]$ ,  $j \leq J$ , is the  $j$ -th configuration returned by GPA at time  $t$ , then in DRA, servers with effective configuration  $\mathbf{k}[j]$  get rank  $j$ , and servers without effective configuration have rank  $J + 1$ .

*Definition 5.2.* Given an  $i \leq C^{(g)}$ , Reject Group servers can be divided as  $\text{RG} = \overline{\text{RG}}(i) \cup \text{RG}(i)$ . The servers with index 1 without effective configuration in  $\bar{\mathbf{k}}^{(\ell)}$ , for  $\ell = 1, \dots, i$ , belong to  $\overline{\text{RG}}(i)$ , while the rest of servers of Reject Group belong to  $\text{RG}(i)$ .

### 5.1 Effective Slot Deficit: $q$ Process

The job admission and configuration assignment under DRA crucially depends on the  $q$  process defined below.

*Definition 5.3.* For  $i \in \{1, \dots, C^{(g)}\}$ , and  $j \in \mathcal{J}$ , we define

$$q_{\bar{\mathbf{k}}^{(i)}, j}^L(t) := \sum_{\ell=1}^i X_{\bar{\mathbf{k}}^{(\ell)}}^{L(e)}(t) \bar{k}_j^{(\ell)} - Y_j^L(t) - g(L). \quad (24)$$

Note that,  $\forall j \in \mathcal{J}$ ,  $q_{\bar{\mathbf{k}}^{(i_2)}, j}^L(t) \geq q_{\bar{\mathbf{k}}^{(i_1)}, j}^L(t)$  if  $i_2 \geq i_1$ .

In words,  $q_{\bar{\mathbf{k}}^{(i)}, j}^L(t)$  measures the difference between the total number of type- $j$  slots (filled or empty) in servers that have effective configurations in the set  $\{\bar{\mathbf{k}}^{(\ell)} : \ell \leq i\}$  (see Definition 5.1), and the number of type- $j$  jobs in the system  $Y_j^L(t)$  and  $g(L)$  type- $j$  reservation slots.

Note that DRA (specifically GPA) will stop assigning configurations that have type- $j$  slots, once  $Y_j^L(t) + g(L)$  slots can be accommodated in servers with effective configuration in  $\{\bar{\mathbf{k}}^{(\ell)}, \ell \leq i\}$ . Since slots are created per server basis, by assigning configurations which each has at most  $K$  slots, we have  $q_{\bar{\mathbf{k}}^{(i)}, j}^L(t) < K$ .

To gain more insight, note that when  $q_{\bar{\mathbf{k}}^{(i)}, j}^L(t) \geq 0$  for an  $i \in \{1, \dots, C^{(g)}\}$ , it means type- $j$  jobs have enough reservation. When it is negative, it indicates the deficit of slots in servers with effective configuration  $\{\bar{\mathbf{k}}^{(\ell)}, \ell \leq i\}$ . When  $q_{\bar{\mathbf{k}}^{(i)}, j}^L(t) > -g(L) + JK$ , for an  $i \in \{1, \dots, C^{(g)}\}$ , a type- $j$  arrival at time  $t$  will certainly find a valid empty slot ( $\text{AG}_j \neq \emptyset$ ) and will be admitted. This is because the



number of empty slots of type  $j$  in Reject Group servers with any effective configuration is less than  $JK$ .

The  $\mathbf{q}$  process also determines the configuration assigned by CRA to an empty server  $\ell_e$  chosen for reassignment. The configuration would be  $\bar{\mathbf{k}}^{(i)}$ ,  $i < C^{(g)}$ , if:

$$\max_{j: \bar{k}_j^{(i)} > 0} q_{\bar{\mathbf{k}}^{(i)}, j}^L(t) \geq 0, \forall \ell \leq i - 1, \quad (25a)$$

$$\max_{j: \bar{k}_j^{(i)} > 0} q_{\bar{\mathbf{k}}^{(i)}, j}^L(t) < 0. \quad (25b)$$

This also implies that if only (25b) holds, the server would be assigned to one of the configurations  $\bar{\mathbf{k}}^{(\ell)}$ ,  $\ell = 1, \dots, i$ .

## 5.2 Existence of Fluid Limits

We define the scaled (normalized with  $L$ ) processes  $\mathbf{x}^{L(e)}(t)$ ,  $\mathbf{y}^L(t)$ , as follows. For  $i \in \{1, \dots, C^{(g)}\}$ , and  $j \in \mathcal{J}$ ,

$$\mathbf{x}_{\bar{\mathbf{k}}^{(i)}}^{L(e)}(t) = \frac{1}{L} X_{\bar{\mathbf{k}}^{(i)}}^{L(e)}(t), \quad \mathbf{y}_j^L = \frac{1}{L} Y_j^L(t),$$

and define  $\mathbf{z}^L(t) := (\mathbf{x}^{L(e)}(t), \mathbf{y}^L(t))$ . We also define the space

$$\mathcal{Z} = \left\{ (\mathbf{x}^{(e)}, \mathbf{y}) : \mathbf{y} \in \Lambda, x_{\bar{\mathbf{k}}^{(i)}}^{(e)} \geq 0, \sum_{i=1}^{C^{(g)}} x_{\bar{\mathbf{k}}^{(i)}}^{(e)} \leq 1, \sum_{i=1}^{C^{(g)}} x_{\bar{\mathbf{k}}^{(i)}}^{(e)} \bar{\mathbf{k}}^{(i)} \leq \mathbf{y} \right\}.$$

where  $\Lambda$  was defined in (22).

**PROPOSITION 5.4.** *Consider a sequence of systems with increasing  $L$ , and initializations  $\mathbf{z}^L(0) = (\mathbf{x}^{L(e)}(0), \mathbf{y}^L(0)) \in \mathcal{Z}$ , as  $L \rightarrow \infty$ . Then there is a subsequence of  $L$  such that  $\mathbf{x}^{L(e)}(t) \rightarrow \mathbf{x}^{(e)}(t)$ ,  $\mathbf{y}^L(t) \rightarrow \mathbf{y}(t)$ , along the subsequence. Any limit  $\mathbf{z}(t) := (\mathbf{x}^{(e)}(t), \mathbf{y}(t))$ ,  $t \geq 0$ , is called a fluid limit sample path. The convergence is almost surely u.o.c. (uniformly over compact time intervals) and the fluid limit sample paths are Lipschitz continuous.*

**PROOF.** Proof is standard, and can be found in Appendix D.  $\square$

## 5.3 Description of Fluid Limits

We provide an informal description of fluid limit equations here. The formal definitions and proofs can be found in Appendix G.

The properties of the fluid limit processes crucially depend on the  $\mathbf{q}$  process (Definition 5.3). First note that, from (24) and since  $q_{\bar{\mathbf{k}}^{(i)}, j}^L(t) < K$ , it follows that

$$\sum_{\ell=1}^{C^{(g)}} \bar{k}_j^{(\ell)} x_{\bar{\mathbf{k}}^{(\ell)}}^{(e)}(t) \leq y_j(t), \forall j \in \mathcal{J}. \quad (26)$$

Let  $x_{\emptyset}(t)$  be the fraction of servers which are empty and of rank  $J + 1$  at the fluid limit. When  $x_{\emptyset}(t) > 0$ , then CRA always finds empty rank  $J + 1$  servers available for reassignment. In this case, every job type will have enough empty slots, and all the arrivals will be admitted, i.e., we can find an  $\epsilon$  sufficiently small such that for every job type  $j$  and every time  $\tau \in [t, t + \epsilon]$ ,  $q_{\bar{\mathbf{k}}^{(C^{(g)})}, j}^L(\tau) \geq 0$ . Hence, noting that at the fluid limit type- $j$  jobs arrive at rate  $\lambda_j$  and

existing type- $j$  jobs depart at rate  $y_j(t)\mu_j$ ,

$$dy_j(t)/dt = \lambda_j - y_j(t)\mu_j, \forall j \in \mathcal{J}, \quad (27a)$$

$$\sum_{\ell=1}^{C^{(g)}} \bar{k}_j^{(\ell)} x_{\bar{\mathbf{k}}^{(\ell)}}^{(e)}(t) = y_j(t), \quad (27b)$$

where Equality (27b) is based on (24) and due to the fact that  $\lim_{L \rightarrow \infty} \frac{1}{L} q_{\bar{\mathbf{k}}^{(C^{(g)})}, j}^L(t) = 0$  in this case.

A major difficulty in describing fluid limits happens on the boundary  $x_{\emptyset}(t) = 0$ , i.e., when there are not always empty rank  $J + 1$  servers available for reassignment when CRA runs. In this case, let  $i^*(t)$  be the largest index in  $\{1, \dots, C^{(g)} - 1\}$  such that for every  $i \leq i^*(t)$ ,

$$\sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} x_{\bar{\mathbf{k}}^{(\ell)}}^{(e)}(t) = y_{j_i}(t), \text{ for some } j_i \in \mathcal{J}, \quad (28)$$

with the convention that  $i^*(t) = 0$  if (28) does not hold for  $i = 1$ . If  $i^*(t) < C^{(g)} - 1$ , then for  $L$  sufficiently large, and every time  $\tau \in [t, t + \epsilon]$  for  $\epsilon$  sufficiently small,

$$\max_{j: \bar{k}_j^{(i^*(t)+1)} > 0} q_{\bar{\mathbf{k}}^{(i^*(t)+1)}, j}^L(\tau) < 0. \quad (29)$$

Based on Definition 5.2, servers in  $\overline{\text{RG}}(i^*(t) + 1)$  have higher ranks compared to those in  $\text{RG}(i^*(t) + 1)$ , so any migrations by DRA will take place from  $\overline{\text{RG}}(i^*(t) + 1)$  first. We can then show that servers of  $\overline{\text{RG}}(i^*(t) + 1)$  empty at the fluid scale, at a rate of at least

$$\frac{\mu_{\min}}{JKC^2} \left( 1 - \sum_{\ell=1}^{i^*(t)+1} x_{\bar{\mathbf{k}}^{(\ell)}}^{(e)}(t) \right), \quad (30)$$

where  $\mu_{\min} := \min_{j \in \mathcal{J}} \mu_j$  (see Lemma F.3 in Appendix F).

The algorithm will reassign any such server that empties to one of configurations  $\bar{\mathbf{k}}^{(\ell)}$  for  $\ell = 1, \dots, i^*(t) + 1$ . If instead  $i^*(t) = C^{(g)} - 1$ , then it is uncertain whether servers that empty need to be reassigned to a new configuration or not, depending on whether  $\max_{j \in \mathcal{J}: \bar{k}_j^{(i)} > 0} q_{\bar{\mathbf{k}}^{(i)}, j}^L(\tau) < 0$ , for some  $i < C^{(g)}$  at time  $\tau \in [t, t + \epsilon]$ .

Hence, what we see is that, if  $x_{\emptyset}(t) = 0$ , when a server gets empty, it can be assigned to one of the configurations  $\bar{\mathbf{k}}^{(i)}$ ,  $i = 1, \dots, i^*(t) + 1$ . Exact characterization of these assignment rates, however, is not easy as they depend on values of processes  $q_{\bar{\mathbf{k}}^{(i)}, j}^L(\tau)$ ,  $i \in \{1, \dots, i^*(t)\}$ ,  $j \in \mathcal{J}$ , which evolve at a much faster time scale than the scaled processes  $\mathbf{x}^{L(e)}$  and  $\mathbf{y}^L$ . By the continuity of the fluid limit sample paths, at any regular time  $t$ , we can choose  $\epsilon$  small enough such that for all  $\tau \in [t, t + \epsilon]$ ,  $\mathbf{y}(\tau)$ , and  $\mathbf{x}^{(e)}(\tau)$  are approximately constant and equal to  $\mathbf{y}(t)$  and  $\mathbf{x}^{(e)}(t)$ , respectively (their actual change being of order  $\epsilon$ ). However, over the same interval, the  $q^L$  process makes  $O(L)$  transitions and its elements can change in the range  $[-LK, K]$ . This phenomenon is known as *separation of time scales* and has been also observed in other systems, e.g. [26, 27].

To further analyze fluid limits in our setting, we divide the interval  $[t, t + \epsilon]$  into smaller intervals of length  $\omega(\log L/L)$ , and infer properties for the fluid limits over  $[t, t + \epsilon]$  based on averaging the behavior of scaled processes over these smaller intervals, as  $L \rightarrow \infty$ , and  $\epsilon \rightarrow 0$ . To this end, we first make a few definitions.

Since the rate of change of any of the processes  $x_{\bar{k}^{(i)}}^{L(e)}(\tau)$  and  $y_j^L(\tau)$  over a subinterval is of interest, we give it a special name below.

*Definition 5.5 (Local Derivatives).* Given an interval  $[\tau_a, \tau_b]$ , we define the ‘‘local derivatives’’ of the scaled processes as

$$\nabla x_{\bar{k}^{(i)}}^{L(e)}[\tau_a, \tau_b] := \frac{x_{\bar{k}^{(i)}}^{L(e)}(\tau_b) - x_{\bar{k}^{(i)}}^{L(e)}(\tau_a)}{\tau_b - \tau_a} \quad i = 1, \dots, C^{(g)} \quad (31)$$

$$\nabla y_j^L[\tau_a, \tau_b] := \frac{y_j^L(\tau_b) - y_j^L(\tau_a)}{\tau_b - \tau_a} \quad j \in \mathcal{J}. \quad (32)$$

*Definition 5.6.* For any  $i \leq C_\rho^{(g)} - 1$ , we define a set

$$\mathcal{J}^{(i)} := \{j \in \mathcal{J} : \bar{k}_j^{(i)} > 0, \sum_{\ell=1}^i \bar{k}_j^{(\ell)} x_{\bar{k}^{(\ell)}}^{(g)} = \rho_j\}. \quad (33)$$

*Definition 5.7.* For given positive constants  $\alpha_i, i = 1, \dots, C_\rho^{(g)} - 1$ , we define  $C_\alpha(t)$  to be the largest index at time  $t$  such that  $C_\alpha(t) \leq \min(i^*(t), C_\rho^{(g)} - 1)$  and

$$\forall i \in [1, \dots, C_\alpha(t)] : x_{\bar{k}^{(i)}}^{(g)} - x_{\bar{k}^{(i)}}^{(e)}(t) < \alpha_i. \quad (34)$$

*5.3.1 Subinterval construction.* We first define a function  $f(L)$  below, which will control the length of subintervals.

*Definition 5.8.* The function  $f(L)$  is defined as

$$f(L) := \frac{\sqrt{g(L) \log(L)}}{L} \quad (35)$$

where  $g(L)$  is the reservation factor as defined in DRA.

We divide  $[t, t + \epsilon)$  into smaller intervals  $[\tau_n, \tau_{n+1})$ , such that

$$\tau_0 = t, \tau_n = \tau_{n-1} + D_{L,\epsilon}, n = 1, \dots, N_L, \quad (36)$$

where  $N_L = \lceil 1/f(L) \rceil$  is the number of such smaller intervals, and  $D_{L,\epsilon} = \frac{\epsilon}{N_L}$  is the length of each one. We then further divide each  $[\tau_n, \tau_{n+1})$  into a constant number  $M_n$  of subintervals  $[\tau_n^{(m-1)}, \tau_n^{(m)})$ ,  $m = 1, \dots, M_n$ ,  $\tau_n^{(0)} = \tau_n$ ,  $\tau_n^{(M_n)} = \tau_{n+1}$ . For every  $n$ , the sequence of stopping times  $\tau_n^{(m)}$  is recursively generated as follows:

Each time  $\tau_n^{(m)}$  is associated with a driving set of job indexes  $\tilde{\mathcal{J}}[m]$ , with the initialization  $\tilde{\mathcal{J}}[0] = \emptyset$  and  $\tau_n^{(0)} = \tau_n$ . Suppose  $\tilde{\mathcal{J}}[m-1] := \{j_i : i = 1, \dots, G_{m-1}\}$  at time  $\tau_n^{(m-1)}$ , where  $j_i \in \mathcal{J}^{(i)}$  (Definition 5.6). Define  $h^{\tilde{\mathcal{J}}[m-1],(\ell)}(t)$ ,  $\ell = 1, \dots, G_{m-1}$ , to be the (unique) solution to the following system of equations

$$\sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} h^{\tilde{\mathcal{J}}[m-1],(\ell)}(t) = \lambda_{j_i} - \mu_{j_i} y_{j_i}(t), \quad i = 1, \dots, G_{m-1}. \quad (37)$$

The next  $\tau_n^{(m)}$  is the earliest time  $\tau \in [\tau_n^{(m-1)}, \tau_{n+1})$  such that  $q_{\bar{k}^{(G_m)}, j}^L(\tau) \geq 0$  for some  $G_m \leq \min(G_{m-1} + 1, C_\alpha(t))$  and some  $j \in \mathcal{J}^{(G_m)}$ . Further, if  $G_m \leq G_{m-1}$ , we additionally require that

$$\sum_{\ell=1}^{G_m} \bar{k}_j^{(\ell)} h^{\tilde{\mathcal{J}}[m-1],(\ell)}(t) > \lambda_j - \mu_j y_j(t). \quad (38)$$

At such a time  $\tau$ , we set  $\tau_n^{(m)} = \tau$ , and the driving index set is set to

$$\tilde{\mathcal{J}}[m] := \{j'_i : i = 1, \dots, G_m\}, \quad (39)$$

where  $j'_i = j_i$  for  $i = 1, \dots, G_m - 1$ , and  $j'_m = j$ . Also,  $h^{\tilde{\mathcal{J}}[m],(\ell)}(t)$ ,  $\ell = 1, \dots, G_m$ , is set to the solution of the system of equations (37)

for the set  $\tilde{\mathcal{J}}[m]$ . If no time  $\tau \in [\tau_n^{(m-1)}, \tau_{n+1})$  satisfies the given conditions, then  $m = M_n$  and  $\tau_n^{(M_n)} = \tau_{n+1}$ .

The importance of quantities  $h^{\tilde{\mathcal{J}}[m],(i)}(t)$ ,  $i = 1, \dots, G_m$ , will become evident later where we will show (see Lemma G.2 in Appendix) that

$$\nabla x_{\bar{k}^{(i)}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] = h^{\tilde{\mathcal{J}},(i)}(t) + \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}}. \quad (40)$$

Hence, roughly, (37) gives the values of local derivatives, while when (38) occurs, the values of local derivatives change.

Note that the number of stopping times  $M_n$  in any interval  $[\tau_n, \tau_{n+1})$  is bounded. This is because the number of different driving sets  $\tilde{\mathcal{J}}[m]$  is finite and no set may appear twice in that sequence, since the comparison (38) induces a total ordering between the sets. Considering all possible driving set of indexes that may appear in the sequence, we have  $M_n \leq 1 + \sum_{i=1}^{C_\alpha(t)} \prod_{\ell=1}^i |\mathcal{J}^{(\ell)}| < \infty$ .

*5.3.2 Properties of fluid limits over subintervals.* Given an  $\epsilon_\rho > 0$ , we first define the set of fluid limit states

$$\Gamma[\epsilon_\rho] := \{(x^{(e)}, y) : y \leq \rho + \epsilon_\rho\} \cap \mathcal{Z}. \quad (41)$$

The following lemma states the invariant property of  $\Gamma[\epsilon_\rho]$ .

**LEMMA 5.9.** *If  $(x^{(e)}(0), y(0)) \in \mathcal{Z}$ , then for any  $\epsilon_\rho > 0$ , there is a time  $T_{\epsilon_\rho} > 0$  such that for all  $t \geq T_{\epsilon_\rho}$ ,  $(x^{(e)}(t), y(t)) \in \Gamma[\epsilon_\rho]$ . Further, convergence is uniform over all initial states in  $\mathcal{Z}$ .*

**PROOF.** See Appendix E.  $\square$

The following proposition states the behavior of scaled processes over the subintervals.

**PROPOSITION 5.10.** *For every  $m \in \{0, \dots, M_n - 1\}$ , let  $\tilde{\mathcal{J}}[m] = \{j_i : i = 1, \dots, G_m\}$  be the index set corresponding to time  $\tau_n^{(m)}$ , and  $\ell_m := G_m + 1$ . Then we can choose  $\alpha_i$ s in Definition 5.7, and  $\epsilon_\rho$  in (41) sufficiently small, such that, for any regular time  $t \geq T_{\epsilon_\rho}$ , with probability at least  $1 - o(L^{-2})$ , all the following properties hold:*

*P.1. For every  $i \in \{1, \dots, \ell_m - 1\}$ ,*

$$\sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} \nabla x_{\bar{k}^{(\ell)}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] = \lambda_{j_i} - \mu_{j_i} y_{j_i}(t) + \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}} \quad (42)$$

*P.2. If  $\ell_m < C_\rho^{(g)}$ ,*

$$\nabla x_{\bar{k}^{(\ell_m)}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] > \frac{\mu \min}{2} \alpha_{\ell_m} + \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}} \quad (43)$$

*P.3. If  $\ell_m = C_\rho^{(g)}$ ,*

$$\nabla x_{\bar{k}^{(C_\rho^{(g)})}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] > \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}} + \min \left\{ \frac{\mu \min}{JKC^2} \left( 1 - \sum_{i=1}^{C_\rho^{(g)}} x_{\bar{k}^{(i)}}^{(e)}(t) \right) - \sum_{i=1}^{C_\rho^{(g)}-1} \left( \nabla x_{\bar{k}^{(i)}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] \right)^+, \right. \\ \left. \min_{j: \bar{k}_j^{(C_\rho^{(g)})} > 0} \frac{\lambda_j - \mu_j y_j(t) - \sum_{i=1}^{C_\rho^{(g)}-1} \bar{k}_j^{(i)} \nabla x_{\bar{k}^{(i)}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}]}{\bar{k}_j^{(C_\rho^{(g)})}} \right\} \quad (44)$$

In words, (P.1.) states that, roughly, for any  $i < \ell_m$ , there is a job type  $j_i$  such that each of the effective number of servers with configurations  $\{\bar{\mathbf{k}}^{(\ell)}\}$  for  $\ell = 1, \dots, i$  changes at a rate that can accommodate exactly additional type- $j_i$  arrivals.

(P.2.) states that effective number of servers with configuration  $\bar{\mathbf{k}}^{(\ell_m)}$  increases by an amount proportional to  $\alpha_{\ell_m}$ . This implies that the rate at which  $x_{\bar{\mathbf{k}}^{(\ell_m)}}^{(e)}(t)$  converges to the global greedy solution is lower bounded by a constant independent of the system state.

(P.3.) describes the change in the effective number of servers in  $\bar{\mathbf{k}}^{(C_\rho^{(g)})}$ , the last configuration of the global greedy solution. The change either satisfies the same condition as (P.1.) or it is bounded by the difference of how fast Reject Group servers empty (based on (30) for  $i^*(t) = C_\rho^{(g)} - 1$ ) and at what rate they are assigned to configurations  $\bar{\mathbf{k}}^{(i)}$  for  $i < C_\rho^{(g)}$ .

PROOF OF PROPOSITION 5.10. The proof, including all supporting Lemmas, is provided in Appendix G.  $\square$

## 6 CONVERGENCE ANALYSIS

We show that the fluid limit of the effective configuration process  $\mathbf{x}^{(e)}(t)$  (which is a lower bound on the number of servers in each configuration) converges to the global greedy solution  $\mathbf{x}^{(g)}$ .

THEOREM 6.1. *Consider the fluid limits of the system under DRA, under any workload  $\rho$ , and any initial state  $\mathbf{z}(0) \in \mathcal{Z}$ . Then*

$$\lim_{t \rightarrow \infty} x_{\mathbf{k}}^{(e)}(t) = x_{\mathbf{k}}^{(g)}, \quad \mathbf{k} \in \mathcal{K}^{(g)}. \quad (45)$$

PROOF. Recall that  $\mathbf{z}(t) = (\mathbf{x}^{(e)}(t), \mathbf{y}(t))$ . We want to show that  $\mathbf{z}(t)$  converges to a point in the set  $\Gamma^*$  defined as

$$\Gamma^* := \{\mathbf{z} := (\mathbf{x}^{(e)}, \mathbf{y}) \in \Gamma[\epsilon_\rho] : x_{\mathbf{k}}^{(e)} = x_{\mathbf{k}}^{(g)}, \mathbf{k} \in \mathcal{K}^{(g)}\}. \quad (46)$$

where  $\Gamma[\epsilon_\rho]$  was defined in (41).

To show convergence, we use a Lyapunov function of the form

$$V(\mathbf{z}(t)) := \sum_{i=1}^{C_\rho^{(g)}} Z_i \left( x_{\bar{\mathbf{k}}^{(i)}}^{(g)} - x_{\bar{\mathbf{k}}^{(i)}}^{(e)}(t) \right) + Z \sum_{j=1}^J (y_j(t) - \rho_j)^+, \quad (47)$$

where  $Z$  and  $Z_i$ ,  $i \in \{1, \dots, C_\rho^{(g)}\}$ , are positive constants satisfying

$$Z > 4Z_1, \quad Z_i > \xi Z_{i+1}, \quad i = 1, \dots, C_\rho^{(g)} - 1, \quad (48)$$

for a  $Z_{C_\rho^{(g)}} > 0$ , and a sufficiently large constant  $\xi > 2K + 1$ .

The constants  $\epsilon_\rho$  and  $\xi$  will be chosen carefully to ensure the conditions of LaSalle's invariance principle [12, 35] hold for any  $\mathbf{z} \in \Gamma[\epsilon_\rho]$ , i.e.,

- (i) For any  $\mathbf{z} \in \Gamma[\epsilon_\rho]$ , we have  $V(\mathbf{z}) \geq 0$  and  $V(\mathbf{z}) = 0$  if and only if  $\mathbf{z} \in \Gamma^*$ ,
- (ii) For any  $\mathbf{z}(t) \in \Gamma[\epsilon_\rho] \setminus \Gamma^*$ ,  $dV(\mathbf{z}(t))/dt < 0$ , almost surely.

These conditions together with Lemma 5.9 will then imply that the limit points of trajectory  $\mathbf{z}(t)$  are in  $\Gamma^*$ .

We state each condition as a Proposition followed by its proof.

PROPOSITION 6.2. *Consider  $V(\mathbf{z})$  in (47), with coefficients in (48), for any  $\xi > (2K + 1)$ , and  $\epsilon_\rho > 0$ . Then we have  $V(\mathbf{z}) \geq 0$  for any  $\mathbf{z} \in \Gamma[\epsilon_\rho]$ , and  $V(\mathbf{z}) = 0$  if and only if  $\mathbf{z} \in \Gamma^*$ .*

PROOF OF PROPOSITION 6.2. Consider the following maximization problem over  $\boldsymbol{\eta} \in \mathbb{R}^{C_\rho^{(g)}}$ ,  $\boldsymbol{\theta} \in \mathbb{R}^J$ , where  $\eta_i$  corresponds to  $x_{\bar{\mathbf{k}}^{(i)}}^{(e)}(t)$  and  $\theta_j$  corresponds to  $(y_j(t) - \rho_j)^+$  in (47),

$$\max_{\boldsymbol{\eta}, \boldsymbol{\theta}} \sum_{i=1}^{C_\rho^{(g)}} Z_i \eta_i - \sum_{j=1}^J Z \theta_j \quad (49a)$$

$$\text{s.t.} \quad \sum_{i=1}^{C_\rho^{(g)}} \eta_i \leq 1, \quad (49b)$$

$$\sum_{i=1}^{C_\rho^{(g)}} \bar{k}_j^{(i)} \eta_i - \theta_j \leq \rho_j, \quad j = 1, \dots, J \quad (49c)$$

$$\theta_j \leq \epsilon_\rho, \quad j = 1, \dots, J \quad (49d)$$

$$\eta_i \geq 0, \quad i = 1, \dots, C_\rho^{(g)} \quad (49e)$$

$$\theta_j \geq 0, \quad j = 1, \dots, J. \quad (49f)$$

To prove the proposition, it is enough to show that the assignment  $(\boldsymbol{\eta}^{(g)}, \boldsymbol{\theta}^{(g)})$  that corresponds to the global greedy solution  $\mathbf{x}^{(g)}$  is the unique maximizer of the above LP. This assignment is

$$\begin{aligned} \eta_i^{(g)} &= x_{\bar{\mathbf{k}}^{(i)}}^{(g)}, \quad i = 1, \dots, C_\rho^{(g)}, \\ \theta_j^{(g)} &= 0, \quad j = 1, \dots, J. \end{aligned} \quad (50)$$

First note that (50) is a basic feasible solution for LP (49), i.e., it is a corner point of the LP's Polytope, since it is on the boundary of  $C_\rho^{(g)} + J$  independent inequalities (equal to the number of variables).

To show that (50) is the "unique maximizer", we need to verify that every neighboring corner point has lower objective value, and to do this, it suffices to verify that by moving along any valid direction within the Polytope, starting from assignment (50), the objective value is reduced. This proves that point (50) is locally optimal, which implies it is also global optimal, since the optimization is LP (and convex) [11]. In the rest of the proof, we use  $g_j$  to be the mapping in Definition 3.4 for  $j = 1, \dots, I_\rho$ , and  $\sigma_j$  to be the permutation of indexes  $\{1, \dots, J\}$  as defined in Proposition 3.1.

We define  $\Delta\eta_i := \eta'_i - \eta_i^{(g)}$  for  $i \in \{1, \dots, C_\rho^{(g)}\}$ , and  $\Delta\theta_j := \theta'_j - \theta_j^{(g)}$  for  $j \in \{1, \dots, J\}$ , where  $\eta'_i$  and  $\theta'_j$  are the values of a feasible point. We prove that the change in objective is negative considering only one positive  $\Delta\eta_i$  for some  $i \in \{1, \dots, C_\rho^{(g)}\} \setminus \{g_j : j = 1, \dots, I_\rho\}$ , while the other  $\Delta\eta_i$ s in this set are 0, and constraints (49b)–(49f) are not violated. This suffices because any feasible point can be constructed as a convex summation of the changes  $\Delta\eta_i$  and if individual changes reduce objective, their convex sum will reduce the objective too.

Suppose  $i^* \in \{1, \dots, C_\rho^{(g)}\} \setminus \{g_j : j = 1, \dots, I_\rho\}$  is the index for which  $\Delta\eta_{i^*} > 0$ . A feasible point will necessarily satisfy the following set of equations, which correspond to  $C_\rho^{(g)} + J$  constraints (specifically, (49b), (49c) for  $j \in \{\sigma_{j'} : j' = 1, \dots, I_\rho\}$ , and (49f) for  $j = 1, \dots, J$ ) which held as equalities at point (50),

$$-\Delta\theta_j \leq 0, \quad j = 1, \dots, J,$$

$$-\Delta\eta_{i^*} < 0; \Delta\eta_i = 0, i \neq i^*, i \in \{1, \dots, C_\rho^{(g)}\} \setminus \{g_j : j = 1, \dots, I_\rho\},$$

$$\bar{k}_{\sigma_j}^{(i^*)} \Delta\eta_{i^*} + \sum_{\ell=1}^j \bar{k}_{\sigma_j}^{(\ell)} \Delta\eta_{g_\ell} - \Delta\theta_{\sigma_j} \leq 0, \quad j = 1, \dots, I_\rho - 1,$$

$$\Delta\eta_{i^*} + \sum_{j=1}^{I_\rho} \Delta\eta_{g_j} \leq 0.$$

(51)

Notice that the conditions (51) are not necessarily sufficient so even if all of them are satisfied the resulting point may be infeasible. Nevertheless, we prove that in any case the objective function will be reduced. The change in value of objective function is given by

$$\Delta F := \sum_{\ell=1}^{I_\rho} Z_{g_\ell} \Delta \eta_{g_\ell} + Z_{i^*} \Delta \eta_{i^*} - Z \sum_{j=1}^J \Delta \theta_j. \quad (52)$$

Given the conditions (51), we show (52) will be negative by finding constants  $\beta > 0$ ,  $\beta_j > 0$ ,  $j = 1, \dots, I_\rho$ , and  $\gamma_j > 0$ ,  $j = 1, \dots, J$ , such that

$$\begin{aligned} \Delta F &= \beta(-\Delta \eta_{i^*}) + \sum_{j=1}^{I_\rho-1} \beta_j \left( \bar{k}_{\sigma_j}^{(i^*)} \Delta \eta_{i^*} + \sum_{\ell=1}^j \bar{k}_{\sigma_j}^{(g_\ell)} \Delta \eta_{g_\ell} - \Delta \theta_{\sigma_j} \right) \\ &\quad + \beta_{I_\rho} \left( \Delta \eta_{i^*} + \sum_{\ell=1}^{I_\rho} \Delta \eta_{g_\ell} \right) + \sum_{j=1}^J \gamma_j (-\Delta \theta_j). \end{aligned} \quad (53)$$

It is not difficult to show by matching the coefficients of (52) and (53) that the values of  $\beta$ ,  $\beta_j$ , for  $j = 1, \dots, I_\rho$  and  $\gamma_j$  for  $j = 1, \dots, J$ , are strictly positive for the choice of  $Z$  and  $Z_i$ 's in the proposition's statement. The details can be found in Appendix H. ■

**PROPOSITION 6.3.** *For function  $V(\mathbf{z})$ , as defined in (47) and (48), there is a constant  $\xi > 2K + 1$ , such that if  $\mathbf{z}(t) \in \Gamma[\epsilon_\rho] \setminus \Gamma^*$ , then  $\frac{d}{dt} V(\mathbf{z}(t)) < 0$ .*

To prove Proposition 6.3, we first prove the following lemma for the local derivatives over subintervals  $[\tau_n, \tau_{n+1}]$  defined in Section 5.3.1.

**LEMMA 6.4.** *Consider the Lyapunov function  $V(\mathbf{z})$  defined in (47). We can choose the constant  $\xi > 2K + 1$  sufficiently large such that the following holds. If at a regular time  $t$ ,  $V(\mathbf{z}(t)) > \epsilon_V$ , for some  $\epsilon_V > 0$ , then there is a  $\delta(\epsilon_V) > 0$  such that for any  $n \in \{0, \dots, N_L - 1\}$ ,*

$$\sum_{i=1}^{C_\rho^{(g)}} Z_i \nabla x_{\mathbf{k}^{(i)}}^{L(e)} [\tau_n, \tau_{n+1}] > \delta(\epsilon_V) + \sum_{j=1}^J Z \frac{d}{dt} (y_j(t) - \rho_j)^+ + o(1), \quad (54)$$

with probability greater than  $1 - o(L^{-2})$

**PROOF OF LEMMA 6.4.** The proof of Lemma 6.4 is based on using (i) properties of fluid limits in Proposition 5.10, and (ii) the boundedness of local derivatives (Lemma F.2 in Appendix F), and (iii) the fact that  $\frac{d}{dt} (y_j(t) - \rho_j)^+ \leq -\mu_j (y_j(t) - \rho_j)^+$ .

The detailed proof can be found in Appendix I. ■

Finally, by using Lemma 6.4, we can show that change of  $V(\mathbf{z}(t))$  is negative, almost surely, by averaging the change of  $V(\mathbf{z}(t))$  over all the subintervals  $[\tau_n, \tau_{n+1}]$  of  $[t, t + \epsilon)$ , as we do below.

**PROOF OF PROPOSITION 6.3.** Note that at any regular time  $t$ ,

$$\frac{d}{dt} V(\mathbf{z}(t)) = - \sum_{i=1}^{C_\rho^{(g)}} Z_i \frac{d}{dt} x_{\mathbf{k}^{(i)}}^{L(e)}(t) + \sum_{j=1}^J Z \frac{d}{dt} (y_j(t) - \rho_j)^+, \quad (55)$$

and  $\frac{d}{dt} x_{\mathbf{k}^{(i)}}^{L(e)}(t) = \lim_{\epsilon \rightarrow 0} \lim_{L \rightarrow \infty} \frac{x_{\mathbf{k}^{(i)}}^{L(e)}(t+\epsilon) - x_{\mathbf{k}^{(i)}}^{L(e)}(t)}{\epsilon}$ . Hence, using the division of  $[t, t + \epsilon)$  into  $N_L$  subintervals  $[\tau_n, \tau_{n+1}]$  of equal size,

as defined in Section 5.3.1, we can write

$$\begin{aligned} \frac{d}{dt} V(\mathbf{z}(t)) &= - \lim_{\epsilon \rightarrow 0} \lim_{L \rightarrow \infty} \frac{1}{N_L} \sum_{n=1}^{N_L} \sum_{i=1}^{C_\rho^{(g)}} Z_i \nabla x_{\mathbf{k}^{(i)}}^{L(e)} [\tau_n, \tau_{n+1}] \\ &\quad + \sum_{j=1}^J Z \frac{d}{dt} (y_j(t) - \rho_j)^+ \\ &\stackrel{(a)}{<} - \delta(\epsilon_V) - \lim_{\epsilon \rightarrow 0} \lim_{L \rightarrow \infty} \frac{1}{N_L} \sum_{n=1}^{N_L} o(1) \stackrel{(b)}{=} -\delta(\epsilon_V) < 0, \end{aligned}$$

where in (a) we used (54) of Lemma 6.4 in every subinterval  $[\tau_n, \tau_{n+1}]$  and in (b) we used the property that  $\sum_{n=1}^{N_L} o(1)/N_L = o(1)$ .

Let  $E_L$  be the event that

$$- \frac{1}{N_L} \sum_{n=1}^{N_L} \sum_{i=1}^{C_\rho^{(g)}} Z_i \nabla x_{\mathbf{k}^{(i)}}^{L(e)} [\tau_n, \tau_{n+1}] + \sum_{j=1}^J Z \frac{d}{dt} (y_j(t) - \rho_j)^+ > 0.$$

The probability that (54) holds for all  $N_L$  subintervals, is at least  $1 - N_L o(L^{-2}) = 1 - o(L^{-1})$ , which follows from  $N_L = \Theta(1/f(L))$  based on Definition 5.8. Hence,  $\mathbb{P}(E_L) < o(L^{-1})$ , and  $\frac{d}{dt} V(\mathbf{z}(t)) < 0$  holds in probability. We can further show that convergence is almost sure. This is because  $\sum_{L=1}^{\infty} \mathbb{P}(E_L) < \sum_{L=1}^{\infty} o(L^{-1}) < \infty$ , and by the Borel-Cantelli Lemma [8],  $\frac{d}{dt} V(\mathbf{z}(t)) < 0$ , almost surely. ■

Propositions 6.2 and 6.3 complete the proof of Theorem 6.1. □

**PROOF OF THEOREM 4.1.** The proof follows from Theorem 6.1 and Theorems 3.7 and 3.9. The details are standard and can be found in Appendix J. □

## 7 SIMULATION RESULTS


### 7.1 Evaluation using synthetic traffic

In this section, we evaluate the approximation ratio and convergence properties of DRA. We start by choosing the VM types considering the VM instances offered by major cloud providers like Google Cloud, are mainly optimized for either memory, CPU, or regular usage. Further, instances are priced proportional to the resources they request, with each resource having a base pricing rate. To simplify simulations, we considered instances that only have memory and CPU requirements. In particular, we used representative VM

vCPU		Memory: GB per vCPU		
Small	Large	High	Low	Regular
2,4, or 8	32 or 64	8 or 16	1 or 2	4

**Table 1: The representative VM instances from Google Cloud based on combination of vCPU and Memory.**

instances, based on combination of vCPU and memory in Table 1. Lastly, each vCPU usage generates 8 reward per unit time, while each GB of memory generates 1. This choice was made based on the relative pricing of CPU and memory of VMs offered by Google Cloud, according to which 8 GB memory is approximately priced as much as 1 vCPU [22]. We generated random collections of VM types, each with three small and three large VMs, with vCPU and memory chosen randomly from Table 1. Servers always have capacity of 80 vCPUs and 640 GB of memory. The normalized workload



**Figure 2: Global greedy vs. optimal, as workload  $\alpha\rho$  increases. The rewards coincide outside the marked points.**

$\rho_j$  for each VM type  $j$  is selected uniformly at random between 0.2 to 2. The statistics we obtained based on 50 randomly generated VM collections and workloads was that, in 23 of them reward of global greedy was identical to the optimal, on average its ratio compared to optimal was 0.972 and in the worst case it was no less than 0.86. Recall that optimal can be found by solving optimization (7). For the rest of simulations, we considered a subset of the worst-case VM collection and its corresponding workload, namely, VM types are: (1, 1), (4, 16), (2, 32), (32, 256), and  $\rho$  rounded to (2, 1/2, 4/3, 1).


To better understand how workload may affect the approximation ratio, we study this worst-case example and scale its workload  $\rho$  by a factor  $\alpha$  that ranges from 0 to 10. Figure 2 shows the reward for the global greedy  $U^{(g)}[\alpha\rho]$  and the optimal reward  $U^*[\alpha\rho]$ . We notice there are two critical  $\alpha$  points. Before the first point, the workload is low enough such that the global greedy assignment can fully accommodate it, hence its reward is the same as the optimal which should also be able to accommodate the full workload. The second point is a point above which the workload is high such that it is possible to assign the configuration of maximum reward to all servers without leaving any slots empty. In this case, both the rewards will coincide again, and take the maximum possible value.

In Figure 2, the two critical points are  $\alpha = 6/7$  and  $\alpha = 6$ . The worst ratio between the reward of global greedy and the optimal occurs at  $\alpha = 1$ , which is  $\approx 0.862$ . Note that in general  $U^{(g)}[\alpha\rho]$  and  $U^*[\alpha\rho]$  might coincide even between the critical points although this is not the case for this example.


To study the impact of the number of servers  $L$ , we run DRA in systems with various number of servers, and compare the obtained average normalized reward (normalized with  $L$ ) with the global greedy reward  $U^{(g)}[\rho]$ , and the optimal reward  $U^*[\rho]$ . The arrivals are generated at rate  $\rho_j L$ , and service times are exponentially distributed with mean 1. The result is depicted in Figure 3, which clearly shows that as the number of servers  $L$  becomes large, DRA approaches the global greedy reward and 86% of the optimal reward. Further, Figure 4 shows how the reward of DRA evolves over time and converges to the global greedy reward when  $L = 180$ .

## 7.2 Evaluation using real traffic trace

We evaluate our algorithm using a more realistic setting with arrival and service times extracted from a Google cluster dataset [57]. In particular, we extracted tasks which were completed within the time window of the trace and used the first 1 million in all simulations.



**Figure 3: The reward of DRA as a fraction of the optimal reward (left y-axis), and that of the global greedy (right y-axis).**



**Figure 4: Convergence of the reward of DRA to that of the global greedy assignment over time when  $L = 180$  servers.**

Tasks were mapped to types by setting their resource requirements to be the largest of the requested resources and rounding it up to the closest power of 1/2. Their reward was set to be equal to their rounded size multiplied by a factor that depends on their priority. Factor is 1, 3, 9 for priorities 0, 1, 2 respectively. Tasks have the same type if both their priority and normalized size are equal. The size of servers is normalized to 1.

We compare the performance of DRA and three other algorithms:

**Upper bound:** It solves optimization (6) with  $\hat{Y}(t)$  being the number of jobs in an infinite server system that rejects no jobs. This gives an upper bound on the performance of any algorithm.

**Power-of- $d$ -choices:** Upon an arrival, it picks  $d$  servers and attempts to schedule the job arrived in the least loaded server if it fits [58]. We picked  $d = 5$ , but behavior of the algorithm is not expected to change significantly for larger  $d$ .

**DRA+Preemption:** This is simply an extension to our algorithm that preempts some of the jobs of priority 0, when a job of type  $j$  with priority 1 or 2 gets rejected. Notice that preemptions of low priority jobs is already considered in similar scenarios that in Google cluster setting [55]. Specifically, our algorithm attempts to preempt jobs of priority 0 starting from those of smallest size. Considering reservation factor is  $g(L)$  and size of type- $j$  job is  $s_j$ , preemptions will stop if the total size of preempted jobs is  $g(L)s_j$  or no more priority 0 jobs are available. The algorithm finds which jobs to preempt, if any, the same way it finds jobs to migrate so this addition needs minimal changes in implementation.




Figure 5: The comparison of rewards for different number of servers based on the Google trace.




Figure 6: Comparison of the reward over time of different algorithms for a part of the Google trace.

Figure 5 shows the performance results (the time-average of rewards) with varying number of servers. Especially, considering preemptions in DRA makes a great difference. Note that the upper bound may be impossible to achieve by any algorithm.

To give more insight, in Figure 6 we plot the total reward over time for all algorithms for a part of the simulation of 1000 servers, including the corresponding total size of arrivals of all job types of each priority. We notice that power-of- $d$ -choices algorithm can be better than DRA in parts of trace in which a spike in demand of priority 0 jobs is followed by a spike in demand of priority 1 jobs. This is because reservation of DRA is not sufficient to account for spikes in demand, while power-of- $d$ -choices does not efficiently use the resources of all servers and may have more free capacity when a spike occurs. DRA with preemptions is particularly effective in such scenarios as it does not need to reserve resources in advance. In addition, it makes efficient use of the resources of all the servers the same way DRA does and thus is strictly better than both of the other algorithms in almost all parts of the trace.

## 8 CONCLUSIONS

In this paper, we proposed a VM reservation and admission policy that operates in an online manner and can guarantee at least  $1/2$  (and under certain monotone property,  $1 - 1/e$ ) of the optimal

expected reward. Assumptions such as Poisson arrivals and exponential service times are made to simplify the analysis, and the policy itself does not rely on this assumption. The policy strikes a balance between good VM packing and serving high priority VM requests, by maintaining only a small number  $g(L) = \omega(\log L)$  of reserved VM slots at any time. Our techniques for analysis of fluid-scale processes on the boundary in our problem, and the design of LP-based Lyapunov functions with a unique maximizer at the given desired equilibrium, can be of interest on their own.

Although we considered that the policy classifies and reassigns servers at arrival and departure events, this was only to simplify the analysis, and in practice CRA can make such updates periodically, by factoring all arrival or departures in the past period in its input for the current period. Further, if a more accurate estimate of the workload is available, we can incorporate that estimate in the vector  $\hat{Y}$  used by DRA, to improve the convergence time.

Moreover, the policy can be extended to a multi-pool server system, where constant fractions of servers belong to different server types. We postpone the details to a future work.

## REFERENCES

- [1] Giuseppe Aceto, Alessio Botta, Walter De Donato, and Antonio Pescapè. 2013. Cloud monitoring: A survey. *Computer Networks* 57, 9 (2013), 2093–2115.
- [2] Amazon AWS 2019. *Amazon Web Services (AWS)*. <https://aws.amazon.com/>
- [3] Rumen Andonov, Vincent Poirriez, and Sanjay Rajopadhye. 2000. Unbounded knapsack problem: Dynamic programming revisited. *European Journal of Operational Research* 123, 2 (2000), 394–407.
- [4] AWS auto-scaler 2019. *Amazon EC2 Auto-Scaler*. <https://docs.aws.amazon.com/autoscaling/>
- [5] AWS container 2019. *Amazon AWS Containers*. <https://aws.amazon.com/containers/>
- [6] AWS SLA 2019. *Amazon AWS Service Level Agreements (SLAs)*. <https://aws.amazon.com/legal/service-level-agreements/>
- [7] N. G. Bean, R. J. Gibbens, and S. Zachary. 1995. Asymptotic analysis of single resource loss systems in heavy traffic, with applications to integrated networks. *Advances in Applied Probability* 27, 1 (1995), 273–292. <https://doi.org/10.2307/1428107>
- [8] Patrick Billingsley. 2008. *Probability and measure*. John Wiley & Sons.
- [9] Patrick Billingsley. 2013. *Convergence of probability measures*. John Wiley & Sons, 2nd edition.
- [10] Gunter Bolch, Stefan Greiner, Hermann Meer, and Kishor Trivedi. 2006. *Queueing Networks and Markov Chains: Modeling and Performance Evaluation With Computer Science Applications, Second Edition*. Vol. 95.
- [11] Stephen Boyd and Lieven Vandenbergh. 2004. *Convex Optimization*. Cambridge University Press, New York, NY, USA.
- [12] Cyril Cohen and Damien Rouhling. 2017. A Formal Proof in Coq of LaSalle’s Invariance Principle. In *Interactive Theorem Proving*, Mauricio Ayala-Rincón and César A. Muñoz (Eds.). Springer International Publishing, Cham, 148–163.
- [13] Antonio Corradi, Mario Fanelli, and Luca Foschini. 2014. VM consolidation: A real case based on OpenStack Cloud. *Future Generation Computer Systems* 32 (2014), 118–127.
- [14] Marek Cygan, Lukasz Jeż, and Jiří Sgall. 2016. Online knapsack revisited. *Theory of Computing Systems* 58, 1 (2016), 153–190.
- [15] David Pollard 2015. *MiniEmpirical*. <http://www.stat.yale.edu/~pollard/Books/Mini/>
- [16] S.N. Ethier and T.G. Kurtz. 2009. *Markov Processes: Characterization and Convergence*. Wiley.
- [17] Apache Software Foundation. 2019. Apache Hadoop Yarn. <http://hadoop.apache.org/docs/current/hadoop-yarn/hadoop-yarn-site/YARN.html>
- [18] Javad Ghaderi, Yuan Zhong, and Rayadurgam Srikant. 2014. Asymptotic optimality of BestFit for stochastic bin packing. *ACM SIGMETRICS Performance Evaluation Review* 42, 2 (2014), 64–66.
- [19] Mostafa Ghobaei-Arani, Sam Jabbehdari, and Mohammad Ali Pourmina. 2018. An autonomic resource provisioning approach for service-based cloud applications: A hybrid approach. *Future Generation Computer Systems* 78 (2018), 191–210.
- [20] Google Cloud 2019. *Google cloud computing services*. <https://cloud.google.com/>
- [21] Google Kubernetes 2019. *Kubernetes at Google Cloud*. <https://cloud.google.com/kubernetes/>
- [22] Google Pricing 2020. *Google Compute Engine All Pricing*. <https://cloud.google.com/compute/all-pricing>

- [23] Yang Guo, Alexander Stolyar, and Anwar Walid. 2018. Online vm auto-scaling algorithms for application hosting in a cloud. *IEEE Transactions on Cloud Computing* (2018).
- [24] Varun Gupta and Ana Radovanovic. 2012. Online stochastic bin packing. *arXiv preprint arXiv:1211.2687* (2012).
- [25] Rui Han, Li Guo, Moustafa M Ghanem, and Yike Guo. 2012. Lightweight resource scaling for cloud applications. In *EEE/ACM International Symposium on Cluster, Cloud and Grid Computing (ccgrid 2012)*. 644–651.
- [26] PJ Hunt and TG Kurtz. 1994. Large loss networks. *Stochastic Processes and their Applications* 53, 2 (1994), 363–378.
- [27] PJ Hunt, CN Laws, et al. 1997. Optimization via trunk reservation in single resource loss systems under heavy traffic. *The Annals of Applied Probability* 7, 4 (1997), 1058–1079.
- [28] Oscar H Ibarra and Chul E Kim. 1975. Fast approximation algorithms for the knapsack and sum of subset problems. *Journal of the ACM (JACM)* 22, 4 (1975), 463–468.
- [29] Kazuo Iwama and Shiro Taketomi. 2002. Removable online knapsack problems. In *International Colloquium on Automata, Languages, and Programming*. Springer, 293–305.
- [30] Jing Jiang, Jie Lu, Guangquan Zhang, and Guodong Long. 2013. Optimal cloud resource auto-scaling for web applications. In *EEE/ACM International Symposium on Cluster, Cloud, and Grid Computing*. 58–65.
- [31] A Karthik, Arpan Mukhopadhyay, and Ravi R Mazumdar. 2017. Choosing among heterogeneous server clouds. *Queueing Systems* 85, 1-2 (2017), 1–29.
- [32] Hans Kellerer, Ulrich Pferschy, and David Pisinger. 2004. Multidimensional knapsack problems. In *Knapsack problems*. Springer, 235–283.
- [33] Frank P Kelly. 1991. Loss networks. *The annals of applied probability* (1991), 319–378.
- [34] Peter B. Key. 1990. *Optimal control and trunk reservation in loss networks*. Vol. 4. Teachers College Library - Columbia University. 203–242 pages. <https://doi.org/10.1017/S0269964800001558>
- [35] Joseph LaSalle. 1960. Some Extensions of Liapunov’s Second Method. *IRE Transactions on Circuit Theory* 7, 4 (December 1960), 520–527. <https://doi.org/10.1109/TCT.1960.1086720>
- [36] Siva Theja Maguluri, Rayadurgam Srikant, and Lei Ying. 2012. Stochastic models of load balancing and scheduling in cloud computing clusters. In *2012 Proceedings IEEE Infocom*. IEEE, 702–710.
- [37] Siva Theja Maguluri, Rayadurgam Srikant, and Lei Ying. 2014. Heavy traffic optimal resource allocation algorithms for cloud computing clusters. *Performance Evaluation* 81 (2014), 20–39.
- [38] Ming Mao, Jie Li, and Marty Humphrey. 2010. Cloud auto-scaling with deadline and budget constraints. In *IEEE/ACM International Conference on Grid Computing*. 41–48.
- [39] Alberto Marchetti-Spaccamela and Carlo Vercellis. 1995. Stochastic on-line knapsack problems. *Mathematical Programming* 68, 1-3 (1995), 73–104.
- [40] Silvano Martello and Paolo Toth. 1990. An exact algorithm for large unbounded knapsack problems. *Operations research letters* 9, 1 (1990), 15–20.
- [41] Microsoft Azure. 2019. *Microsoft cloud computing service*. <https://azure.microsoft.com/>
- [42] Arpan Mukhopadhyay, A Karthik, Ravi R Mazumdar, and Fabrice Guillemin. 2015. Mean field and propagation of chaos in multi-class heterogeneous loss models. *Performance Evaluation* 91 (2015), 117–131.
- [43] Konstantinos Psychas and Javad Ghaderi. 2017. On non-preemptive VM scheduling in the cloud. *Proceedings of the ACM on Measurement and Analysis of Computing Systems* 1, 2 (2017), 35.
- [44] Konstantinos Psychas and Javad Ghaderi. 2018. Randomized Algorithms for Scheduling Multi-Resource Jobs in the Cloud. *IEEE/ACM Transactions on Networking* 26, 5 (2018), 2202–2215.
- [45] Chenhao Qu, Rodrigo N Calheiros, and Rajkumar Buyya. 2018. Auto-scaling web applications in clouds: A taxonomy and survey. *ACM Computing Surveys (CSUR)* 51, 4 (2018), 1–33.
- [46] Saffraz Rampersaud and Daniel Grosu. 2014. A sharing-aware greedy algorithm for virtual machine maximization. In *IEEE 13th International Symposium on Network Computing and Applications*. 113–120.
- [47] Nilabja Roy, Abhishek Dubey, and Aniruddha Gokhale. 2011. Efficient autoscaling in the cloud using predictive models for workload forecasting. In *IEEE 4th International Conference on Cloud Computing*. 500–507.
- [48] Jin Shao, Hao Wei, Qianxiang Wang, and Hong Mei. 2010. A runtime model based monitoring approach for cloud. In *2010 IEEE 3rd International Conference on Cloud Computing*. IEEE, 313–320.
- [49] Weijia Song, Zhen Xiao, Qi Chen, and Haipeng Luo. 2013. Adaptive resource provisioning for the cloud using online bin packing. *IEEE Trans. Comput.* 63, 11 (2013), 2647–2660.
- [50] Mark Stillwell, Frederic Vivien, and Henri Casanova. 2012. Virtual machine resource allocation for service hosting on heterogeneous distributed platforms.
- [51] Alexander L Stolyar. 2013. An infinite server system with general packing constraints. *Operations Research* 61, 5 (2013), 1200–1217.
- [52] Alexander L Stolyar. 2017. Large-scale heterogeneous service systems with general packing constraints. *Advances in Applied Probability* 49, 1 (2017), 61–83.
- [53] Alexander L Stolyar and Yuan Zhong. 2013. A large-scale service system with packing constraints: Minimizing the number of occupied servers. In *ACM SIGMETRICS Performance Evaluation Review*, Vol. 41. ACM, 41–52.
- [54] Alexander L Stolyar and Yuan Zhong. 2015. Asymptotic optimality of a greedy randomized algorithm in a large-scale service system with general packing constraints. *Queueing Systems* 79, 2 (2015), 117–143.
- [55] Abhishek Verma, Luis Pedrosa, Madhukar Korupolu, David Oppenheimer, Eric Tune, and John Wilkes. 2015. Large-scale cluster management at Google with Borg. *Proceedings of the Tenth European Conference on Computer Systems - EuroSys ’15* (2015), 1–17. <https://doi.org/10.1145/2741948.2741964>
- [56] Ward Whitt. 1985. Blocking when service is required from several facilities simultaneously. *AT&T technical journal* 64, 8 (1985), 1807–1856.
- [57] John Wilkes. 2011. *Google Cluster Data*. <https://github.com/google/cluster-data>
- [58] Qiaomin Xie, Xiaobo Dong, Yi Lu, and Rayadurgam Srikant. 2015. Power of d choices for large-scale bin packing: A loss model. *ACM SIGMETRICS Performance Evaluation Review* 43, 1 (2015), 321–334.
- [59] Yangming Zhao, Yifan Huang, Kai Chen, Minlan Yu, Sheng Wang, and Dongsheng Li. 2015. Joint VM placement and topology optimization for traffic scalability in dynamic datacenter networks. *Computer Networks* 80 (2015), 109–123.

## APPENDIX

### A PROOF OF PROPOSITION 3.1

We omit the notation  $[\rho]$  for compactness. Also we use the following notations for shorthand purposes

$$\text{amin}(j) := \arg \min_{i: k_i^{(j)} > 0} \frac{\rho_i}{k_i^{(j)}} - \sum_{\ell=1}^{j-1} z^{(\ell)} \frac{k_i^{(\ell)}}{k_i^{(j)}}, \quad (56)$$

$$\text{assgn}(j) := \min_{i: k_i^{(j)} > 0} \frac{\rho_i}{k_i^{(j)}} - \sum_{\ell=1}^{j-1} z^{(\ell)} \frac{k_i^{(\ell)}}{k_i^{(j)}}.$$

As a convention, if minimum is attained by more than one indexes, the lowest one is chosen. We define

$$\sigma := (\text{amin}(1), \text{amin}(2), \dots, \text{amin}(J)), \quad (57)$$

$$z^{(j)} := \text{assgn}(j), \quad j = 1, \dots, J.$$

We can verify that  $\sigma$  and its corresponding values  $z^{(j)}$  satisfy all the conditions of Proposition 3.1. It remains to prove that this permutation  $\sigma$  is unique.

Suppose there is another permutation  $\sigma' := \{\sigma'_1, \sigma'_2, \dots, \sigma'_J\}$  that satisfies the properties of Proposition 3.1 and  $j$  is the lowest index for which  $\sigma_j \neq \sigma'_j$ . We define  $D_i^j := \frac{\rho_i}{k_i^{(j)}} - \sum_{\ell=1}^{j-1} z^{(\ell)} \frac{k_i^{(\ell)}}{k_i^{(j)}}$  and compare  $D_{\sigma_j}^j$  to  $D_{\sigma'_j}^j$ . We will reach a contradiction in all possible cases, which proves that permutation  $\sigma$  is unique.

- (1) If  $D_{\sigma_j}^j > D_{\sigma'_j}^j$ , then  $\sigma_j := \text{amin}(j)$  is not the minimizer of (56) and this contradicts the definition of  $\text{amin}(j)$ .
- (2) If  $D_{\sigma_j}^j = D_{\sigma'_j}^j$ , then we consider the index  $j_a$  for which  $\sigma_j = \sigma'_{j_a}$  and the index  $j_b$  for which  $\sigma'_j = \sigma_{j_b}$ . This implies

$$D_{\sigma_{j_b}}^j = D_{\sigma'_j}^j = D_{\sigma_j}^j = D_{\sigma'_{j_a}}^j = z^{(j)}, \quad (58)$$

or equivalently

$$\rho_{\sigma'_{j_a}} = \sum_{\ell=1}^j k_{\sigma'_{j_a}}^{(\ell)} z^{(\ell)} = \rho_{\sigma'_j}, \quad (59)$$

$$\rho_{\sigma_{j_b}} = \sum_{\ell=1}^j k_{\sigma_{j_b}}^{(\ell)} z^{(\ell)} = \rho_{\sigma_j}.$$



We also notice that  $j < j_a$ , since  $\sigma_j \neq \sigma'_i$  for  $i = 1, \dots, j$  and similarly  $j < j_b$ . Then, considering (59), assumption (9),  $j < j_a$  and  $j < j_b$ , we get  $\sigma'_j < \sigma'_{j_a}$  and  $\sigma_j < \sigma_{j_b}$  which are contradictory as they imply

$$\sigma_j < \sigma_{j_b} = \sigma'_j < \sigma'_{j_a} = \sigma_j.$$

- (3) If  $D_{\sigma_j}^j < D_{\sigma'_j}^j$ , then we consider the index  $j_a$  for which  $\sigma_j = \sigma'_{j_a}$ . Then for permutation  $\sigma'$  to be valid, there should be constants  $z^{(\ell)} \geq 0$  for  $\ell = 1, \dots, j_a$  such that,

$$\rho_{\sigma'_{j_a}} = \sum_{\ell=1}^{j_a} k_{\sigma'_{j_a}}^{(\ell)} z^{(\ell)}. \quad (60)$$

On the other hand,

$$\sum_{\ell=1}^{j_a} k_{\sigma'_{j_a}}^{(\ell)} z^{(\ell)} = \rho_{\sigma'_{j_a}} \equiv \rho_{\sigma_j} < \sum_{\ell=1}^j k_{\sigma'_{j_a}}^{(\ell)} z^{(\ell)}. \quad (61)$$

where (a) is a consequence of  $D_{\sigma_j}^j < D_{\sigma'_j}^j$  when  $z^{(j)} = D_{\sigma'_j}^j$ .

From (61), we also get

$$\sum_{\ell=j+1}^{j_a} k_{\sigma'_{j_a}}^{(\ell)} z^{(\ell)} < 0, \quad (62)$$

which contradicts the assumption  $z^{(\ell)} \geq 0$  for  $\ell = j+1, \dots, j_a$ , if we consider  $k_{\sigma'_{j_a}}^{(\ell)} \geq 0$  for  $\ell = j+1, \dots, j_a$ .

## B PROOF OF PROPOSITION 3.5

For  $\mathbf{k} \notin \mathcal{K}^{(g)}$ , it is obvious that  $\hat{X}_{\mathbf{k}}^L = 0$ , since  $\mathbf{k}$  is never assigned by GPA for any input. Thus  $\lim_{L \rightarrow \infty} \frac{\hat{X}_{\mathbf{k}}^L}{L} = 0 = x_{\mathbf{k}}^{(g)}$ . Hence, it remains to prove the proposition for  $\mathbf{k} \in \mathcal{K}^{(g)}$ , i.e., for  $\bar{\mathbf{k}}^{(i)}$ ,  $i = 1, \dots, C^{(g)}$ . For this we will use the following Lemma.

LEMMA B.1. For  $i = 1, \dots, C^{(g)}$ ,

$$|\hat{X}_{\bar{\mathbf{k}}^{(i)}}^L - Lx_{\bar{\mathbf{k}}^{(i)}}^{(g)}| \leq (K+1)^{i-1}, \quad (63)$$

where  $\hat{X}_{\bar{\mathbf{k}}^{(i)}}^L = \text{GPA}(L\rho)$  when the number of servers is  $L$  and  $K$  is an upper bound on the maximum number of jobs in any configuration.

PROOF. For shorthand purposes, define  $\hat{X}_i^a := \hat{X}_{\bar{\mathbf{k}}^{(i)}}^L(t)$  and  $\hat{X}_i^b := Lx_{\bar{\mathbf{k}}^{(i)}}^{(g)}$ . By definition of global greedy assignment, for every  $i \in \{1, \dots, C^{(g)}\}$ , one of the following holds:

- (1) There is some  $j \in \mathcal{J}$  such that  $L\rho_j$  fits exactly in  $\hat{X}_i^b$  servers assigned to  $\bar{\mathbf{k}}^{(\ell)}$  for  $\ell = 1, \dots, i$ , i.e.,

$$L\rho_j = \sum_{\ell=1}^i \hat{X}_i^b \bar{k}_j^{(\ell)}. \quad (64)$$

- (2) All servers are assigned to one of the configurations  $\bar{\mathbf{k}}^{(\ell)}$  for  $\ell = 1, \dots, i$ , i.e.,

$$\hat{X}_i^b = L - \sum_{\ell=1}^{i-1} \hat{X}_i^b. \quad (65)$$

Similarly, for GPA, there is an index  $I_L$  such that one of the following holds:

- (1) For  $i \in \{1, \dots, I_L - 1\}$ , there is  $j \in \mathcal{J}$  such that  $L\rho_j$  jobs fit in  $\hat{X}_i^a$  servers assigned to  $\bar{\mathbf{k}}^{(\ell)}$  for  $\ell = 1, \dots, i$ , but not in  $\hat{X}_i^a$  servers assigned to  $\bar{\mathbf{k}}^{(\ell)}$  for  $\ell = 1, \dots, i-1$  and  $\hat{X}_i^a - 1$  servers assigned to  $\bar{\mathbf{k}}^{(i)}$ . This implies that

$$\hat{X}_i^a \bar{k}_j^{(i)} \geq L\rho_j - \sum_{\ell=1}^{i-1} \hat{X}_i^a \bar{k}_j^{(\ell)}, \quad (66)$$

$$(\hat{X}_i^a - 1) \bar{k}_j^{(i)} < L\rho_j - \sum_{\ell=1}^{i-1} \hat{X}_i^a \bar{k}_j^{(\ell)}.$$

- (2) For  $i \in \{I_L, \dots, C^{(g)}\}$ , all servers are assigned to one of the configurations  $\bar{\mathbf{k}}^{(\ell)}$  for  $\ell = 1, \dots, i$ , i.e.,

$$\hat{X}_i^a = L - \sum_{\ell=1}^{i-1} \hat{X}_i^a. \quad (67)$$

We can show inductively that for any  $i \in \{1, \dots, I_L - 1\}$  and for large enough  $L$ , if (66) holds then (64) holds for the same job type  $j$ . By assuming otherwise we can easily reach a contradiction (details are omitted). This means we can replace  $L\rho_j$  in (66) with the right hand side of (64). Also with similar arguments we can prove that if (67) holds then (65) holds as well. Therefore, for  $i = 1$ , we either get  $\hat{X}_1^a = \hat{X}_1^b = L$  or  $\hat{X}_1^a \bar{k}_j^{(1)} \geq \hat{X}_1^b \bar{k}_j^{(1)} > (\hat{X}_1^a - 1) \bar{k}_j^{(1)}$ . Hence, in either case, we have  $|\hat{X}_1^a - \hat{X}_1^b| < 1$ , which proves (63) for  $i = 1$ . Now suppose the statement is true for indexes  $1, \dots, i-1$ . We show that it is also true for  $i$ .

If (66) holds, then by replacing  $L\rho_j$  in (66) with the right-hand-side of (64), we get

$$|\hat{X}_i^a - \hat{X}_i^b| < 1 + \sum_{\ell=1}^{i-1} \frac{\bar{k}_j^{(\ell)}}{\bar{k}_j^{(i)}} |X_\ell^a - X_\ell^b|.$$

Hence, noting that  $\frac{\bar{k}_j^{(\ell)}}{\bar{k}_j^{(i)}} \leq K$ ,  $\ell = 1, \dots, i-1$ , we get

$$|\hat{X}_i^a - \hat{X}_i^b| \leq 1 + \sum_{\ell=1}^{i-1} K(1+K)^{\ell-1} = (1+K)^{i-1}.$$

If instead (67) holds, then since (65) also holds, and we get

$$\begin{aligned} |\hat{X}_i^a - \hat{X}_i^b| &\leq \sum_{\ell=1}^{i-1} |X_\ell^a - X_\ell^b| \\ &\leq \sum_{\ell=1}^{i-1} (1+K)^{\ell-1} < (1+K)^{i-1}. \end{aligned}$$

This completes the proof of (63) for arbitrary  $i$ .  $\square$

The proposition then follows since for any  $i \in \{1, \dots, C^{(g)}\}$ ,

$$\lim_{L \rightarrow \infty} \left| \frac{\hat{X}_{\bar{\mathbf{k}}^{(i)}}^L}{L} - x_{\bar{\mathbf{k}}^{(i)}}^{(g)} \right| \leq \lim_{L \rightarrow \infty} \frac{(K+1)^{i-1}}{L} = 0.$$

## C PROOF OF PROPOSITION 3.10

Consider a system with  $J$  job types. Suppose type- $i$  jobs, for each  $i = 1, \dots, J-1$ , can fit  $J$  times in an empty server, and type- $J$  jobs can fit  $N+1$  times. Suppose the configuration that uses 1 job of each type  $i$  and  $N$  jobs of type  $J$  is feasible as well. The aforementioned



configurations will be maximal if we assume we have  $J + 1$  resources and

- each type- $i$  job,  $i = 1, \dots, J - 1$ , occupies  $1/J$  of resource  $i$  and  $1/J$  of resource  $J + 1$ .
- each type- $J$  job occupies  $1/(N + 1)$  of resource  $J$  and  $1/(JN)$  of resource  $J + 1$ .

Assume that each type- $i$  job,  $i = 1, \dots, J - 1$ , gives reward  $u_i = \frac{1}{J} \left( \frac{J-1}{J} \right)^{i-1} u$ , and each type- $J$  job gives a reward  $u_J = \frac{1}{N+1} \left( \frac{J-1}{J} \right)^{J-1} u$ . Let the workload  $\rho$  be such that  $\rho_i = 1$  for  $i = 1, \dots, J - 1$  and  $\rho_J = N$ .

In this example, the global greedy assignment assigns only the  $J$  configurations that consist of a single job type and each one is assigned to  $\frac{1}{J}$  fraction of servers. The normalized reward of  $\mathbf{x}^{(g)}$  is

$$U^{(g)}(J, N) = \frac{1}{J} \left( \sum_{i=1}^{J-1} J \frac{1}{J} \left( \frac{J-1}{J} \right)^{i-1} + (N+1) \frac{1}{N+1} \left( \frac{J-1}{J} \right)^{J-1} \right) u = \left( 1 - (1 - 1/J)^J \right) u. \quad (68)$$

The optimal assignment assigns the configuration that uses 1 job of each type  $i$  and  $N$  jobs of type  $J$  to all servers. The normalized reward of  $\mathbf{x}^*$  is therefore

$$U^*(J, N) = \left( \sum_{i=1}^{J-1} \frac{1}{J} \left( \frac{J-1}{J} \right)^{i-1} + N \frac{1}{N+1} \left( \frac{J-1}{J} \right)^{J-1} \right) u = \left( 1 - (1 - 1/J)^{J-1} + \frac{N}{N+1} (1 - 1/J)^{J-1} \right) u. \quad (69)$$

From these, the result is obvious, as

$$\lim_{N \rightarrow \infty} \frac{U^{(g)}(J, N)}{U^*(J, N)} = \frac{\left( 1 - (1 - 1/J)^J \right)}{\left( 1 - (1 - 1/J)^{J-1} + (1 - 1/J)^{J-1} \right)} = 1 - (1 - 1/J)^J, \quad (70)$$

and

$$\lim_{J \rightarrow \infty} \lim_{N \rightarrow \infty} \frac{U^{(g)}(J, N)}{U^*(J, N)} = 1 - \lim_{J \rightarrow \infty} (1 - 1/J)^J = (1 - 1/e). \quad (71)$$

## D PROOF OF PROPOSITION 5.4

For the proof of this proposition we will need the following Lemma

LEMMA D.1. For  $i = 1, \dots, C^{(g)}$ , the absolute jump in  $\hat{X}_{\mathbf{k}^{(i)}}^L(t)$ ,  $X_{\mathbf{k}^{(i)}}^L(t)$ ,  $X_{\mathbf{k}^{(i)}}^{L(e)}(t)$ , after a job arrival or departure event, is at most  $(K + 1)^{i-1}$ , where  $K$  is an upper bound on the maximum number of jobs in any configuration.

PROOF. We first prove the result for  $\hat{X}_{\mathbf{k}^{(i)}}^L(t)$ . We consider  $\sum_{i=1}^{C^{(g)}-1} \hat{X}_{\mathbf{k}^{(i)}}^L(t) < L$  before and after an event, as otherwise the range of change of any  $\hat{X}_{\mathbf{k}^{(i)}}^L(t)$  will be even smaller, because of the extra constraint. Consider an arrival or departure event takes place. We denote the values  $\hat{X}_{\mathbf{k}^{(i)}}^L(t)$  for  $i \in \{1, \dots, C^{(g)}\}$ , as given by Algorithm 1, before and after the event by  $\hat{X}_i^a$  and  $\hat{X}_i^b$  respectively.

We define  $i^*$  to be the first index in  $\{1, \dots, C^{(g)}\}$  for which  $\hat{X}_{i^*}^a \neq \hat{X}_{i^*}^b$  so for  $\ell \in \{1, \dots, i^* - 1\}$  we have  $\hat{X}_\ell := \hat{X}_\ell^a = \hat{X}_\ell^b$ . We

also define for  $j \in \{1, \dots, J\}$ ,  $Y_j := Y_j^L(t) + g(L)$ , where  $Y_j^L(t)$  is the number of jobs in the system before the event. Finally we define  $\zeta$  to be 1 if the event is arrival and  $-1$  if the event is departure.

We prove by induction that for  $i \geq i^*$ ,  $|X_i^a - X_i^b| \leq (1 + K)^{i-i^*}$ . We start with the base case  $i = i^*$ . Before any event, we know there is some  $j \in \mathcal{J}$  such that  $Y_j$  jobs fit in  $\hat{X}_\ell^a$  servers assigned to  $\mathbf{k}^{(\ell)}$  for  $\ell = 1, \dots, i$ , but not in  $\hat{X}_\ell^a$  servers assigned to  $\mathbf{k}^{(\ell)}$  for  $\ell = 1, \dots, i - 1$  and  $\hat{X}_i^a - 1$  servers assigned to  $\mathbf{k}^{(i)}$ . This implies

$$\hat{X}_i^a \bar{k}_j^{(i)} \geq Y_j - \sum_{\ell=1}^{i-1} \hat{X}_\ell \bar{k}_j^{(\ell)}, \text{ and } (\hat{X}_i^a - 1) \bar{k}_j^{(i)} < Y_j - \sum_{\ell=1}^{i-1} \hat{X}_\ell \bar{k}_j^{(\ell)}.$$

We can use similar argument after the event when  $Y_j$  changes to  $Y_j + \zeta$ , i.e.,

$$\hat{X}_i^b \bar{k}_j^{(i)} \geq Y_j + \zeta - \sum_{\ell=1}^{i-1} \hat{X}_\ell \bar{k}_j^{(\ell)}, \text{ and } (\hat{X}_i^b - 1) \bar{k}_j^{(i)} < Y_j + \zeta - \sum_{\ell=1}^{i-1} \hat{X}_\ell \bar{k}_j^{(\ell)}.$$

Algebraic manipulations based on this set of equations shows

$$|\hat{X}_i^a - \hat{X}_i^b| < \frac{|\zeta|+1}{\bar{k}_j^{(i)}} \leq 2,$$

or equivalently  $|\hat{X}_i^a - \hat{X}_i^b| \leq 1$ .

Now consider  $i > i^*$  and suppose for  $\ell = i^*, \dots, i - 1$ ,  $|X_\ell^a - X_\ell^b| \leq (1 + K)^{\ell-i}$ . Similar to the arguments for the base case, the following equations have to hold for a job type  $j' \in \mathcal{J}$ ,

$$\begin{aligned} \sum_{\ell=i^*}^i \hat{X}_\ell^a \bar{k}_{j'}^{(\ell)} &\geq Y_{j'} - \sum_{\ell=1}^{i^*-1} \hat{X}_\ell \bar{k}_{j'}^{(\ell)}, \\ \sum_{\ell=i^*}^{i-1} \hat{X}_\ell^a \bar{k}_{j'}^{(\ell)} + (\hat{X}_i^a - 1) \bar{k}_{j'}^{(i)} &< Y_{j'} - \sum_{\ell=1}^{i^*-1} \hat{X}_\ell \bar{k}_{j'}^{(\ell)}, \\ \sum_{\ell=i^*}^i \hat{X}_\ell^b \bar{k}_{j'}^{(\ell)} &\geq Y_{j'} - \sum_{\ell=1}^{i^*-1} \hat{X}_\ell \bar{k}_{j'}^{(\ell)}, \\ \sum_{\ell=i^*}^{i-1} \hat{X}_\ell^b \bar{k}_{j'}^{(\ell)} + (\hat{X}_i^b - 1) \bar{k}_{j'}^{(i)} &< Y_{j'} - \sum_{\ell=1}^{i^*-1} \hat{X}_\ell \bar{k}_{j'}^{(\ell)}. \end{aligned}$$

With algebraic manipulations, we get

$$|\hat{X}_i^a - \hat{X}_i^b| < 1 + \sum_{\ell=i^*}^{i-1} \frac{\bar{k}_{j'}^{(\ell)}}{\bar{k}_{j'}^{(i)}} |X_\ell^a - X_\ell^b|.$$

Hence, considering  $\frac{\bar{k}_{j'}^{(\ell)}}{\bar{k}_{j'}^{(i)}} \leq K$  for  $\ell = i^*, \dots, i - 1$ , we get

$$|\hat{X}_i^a - \hat{X}_i^b| \leq 1 + \sum_{\ell=i^*}^{i-1} K(1 + K)^{\ell-i^*} = (1 + K)^{i-i^*}.$$

The result for  $X_{\mathbf{k}^{(i)}}^L(t)$  then follows by noticing:

- (1) If  $X_{\mathbf{k}^{(i)}}^L(t) \geq \hat{X}_{\mathbf{k}^{(i)}}^L(t)$  then after an event  $X_{\mathbf{k}^{(i)}}^L(t)$  may not increase more than what  $\hat{X}_{\mathbf{k}^{(i)}}^L(t)$  does, which is at most  $(K + 1)^{i-1}$ . Similarly, it may not decrease more than the increase of  $X_{\mathbf{k}^{(i)}}^L(t)$  for  $\ell = 1, \dots, i - 1$  which is at most

$$\sum_{\ell=1}^{i-1} (K + 1)^{\ell-1} < (K + 1)^{i-1},$$

or more than the decrease of  $\hat{X}_{\mathbf{k}^{(i)}}^L(t)$  which is again  $(K + 1)^{i-1}$ .

Notice that the last claim assumes  $X_{\mathbf{k}^{(i)}}^L(t) = \hat{X}_{\mathbf{k}^{(i)}}^L(t)$ , because in case  $X_{\mathbf{k}^{(i)}}^L(t) > \hat{X}_{\mathbf{k}^{(i)}}^L(t)$  it means server of Reject Group assigned to  $\mathbf{k}^{(i)}$  is not empty so maximum decrease of  $X_{\mathbf{k}^{(i)}}^L(t)$  is 1 when that server empties.

- (2) If  $X_{\mathbf{k}^{(\ell)}}^L(t) < \hat{X}_{\mathbf{k}^{(\ell)}}^L(t)$  then no server not assigned to a configuration  $\mathbf{k}^{(\ell)}$  for  $\ell = 1, \dots, i-1$  will be empty. Then after an event  $X_{\mathbf{k}^{(\ell)}}^L(t)$  may not increase more than the decrease of  $X_{\mathbf{k}^{(\ell)}}^L(t)$  for  $\ell = 1, \dots, i-1$  which is at most

$$\sum_{\ell=1}^{i-1} (K+1)^{\ell-1} \leq (K+1)^{i-1} - 1.$$

and decrease of  $X_{\mathbf{k}}^L(t)$  with  $\mathbf{k} \notin \{\mathbf{k}^{(\ell)} : \ell = 1, \dots, i\}$ , which is at most 1 since none of them was empty and at most one may empty after each event. Thus, the total decrease of all servers that may be reassigned to  $\mathbf{k}^{(i)}$  is no more than  $(K+1)^{i-1}$ . Also decrease is at most  $(K+1)^{i-1}$  following the same argument as in the case  $X_{\mathbf{k}^{(i)}}^L(t) \geq \hat{X}_{\mathbf{k}^{(i)}}^L(t)$ .

- (3) For any  $\mathbf{k} \notin \{\mathbf{k}^{(i)} : i = 1, \dots, C^{(g)} - 1\}$ ,  $X_{\mathbf{k}}^L$  may only decrease and the decrease will be at most

$$\sum_{\ell=1}^{C^{(g)}-1} (K+1)^{\ell-1} < (K+1)^{C^{(g)}-1}.$$

Finally, it trivially follows that the maximum change of  $X_{\mathbf{k}^{(i)}}^{L(e)}(t)$  is  $(K+1)^{i-1}$  as well, for  $i = 1, \dots, C^{(g)} - 1$ , since

$$X_{\mathbf{k}^{(i)}}^{L(e)}(t) = \min(\hat{X}_{\mathbf{k}^{(i)}}^L(t), X_{\mathbf{k}^{(i)}}^L(t)).$$

□

We can now prove the existence of fluid limits of the process  $X_{\mathbf{k}}^{L(e)}(t)$ , for  $\mathbf{k} = \mathbf{k}^{(i)}$ ,  $i = 1, \dots, C^{(g)}$ . For each job type  $j$ , we define two independent unit-rate Poisson processes  $\Pi_j^a(\cdot)$  and  $\Pi_j^d(\cdot)$ . By the Functional Strong Law of Large Numbers, almost surely,

$$\frac{\Pi_j^a(Lt)}{L} \rightarrow t, \quad \text{u.o.c.} \quad \frac{\Pi_j^d(Lt)}{L} \rightarrow t, \quad \text{u.o.c.} \quad (72)$$

where u.o.c means uniformly over compact time intervals.

Define  $h_{j,\mathbf{k}}^a(S^L(t))$  and  $h_{j,\mathbf{k}}^d(S^L(t))$  to be the amount of change in  $X_{\mathbf{k}}^{L(e)}(t)$  due to an arrival and departure of a type- $j$  job, respectively, at state  $S^L(t)$ . Then the process  $X_{\mathbf{k}}^{L(e)}(t)$  can be described as

$$X_{\mathbf{k}}^{L(e)}(t) = X_{\mathbf{k}}^{L(e)}(0) + A_{\mathbf{k}}^L(0, t) - D_{\mathbf{k}}^L(0, t) \quad (73)$$

where, for any  $0 \leq t_1 < t_2$ , without loss of generality, we construct the arrival and departure processes for the  $L$ -th system, and the corresponding jumps, as

$$A_{\mathbf{k}}^L(t_1, t_2) = \sum_{j=1}^J \frac{\Pi_j^a(\int_{t_1}^{t_2} \lambda_j L ds)}{\sum_{n=1}^{\infty} \lambda_j} h_{j,\mathbf{k}}^a(S^L(T_n)),$$

$$D_{\mathbf{k}}^L(t_1, t_2) = \sum_{j=1}^J \frac{\Pi_j^d(\int_{t_1}^{t_2} \mu_j Y_j^L(s) ds)}{\sum_{n=1}^{\infty} \mu_j} h_{j,\mathbf{k}}^d(S^L(T_n)),$$

where  $T_n$  is the time of the  $n$ -th jump in corresponding Poisson processes. By Lemma D.1,  $|h_{j,\mathbf{k}}^a(S^L(t))|, |h_{j,\mathbf{k}}^d(S^L(t))| \leq (1+K)^{C^{(g)}-1} := M$ . Then the scaled processes  $\frac{1}{L}A_{\mathbf{k}}^L(t_1, t_2)$  and  $\frac{1}{L}D_{\mathbf{k}}^L(t_1, t_2)$  in (73) are asymptotically Lipschitz continuous, which implies that they have

a convergent subsequence [16]. This is because for any  $t_1 < t_2$ ,

$$\limsup_L \frac{1}{L} \left| A_{\mathbf{k}}^L(t_1, t_2) \right| \leq \limsup_L \frac{1}{L} \Pi_j^a \left( \int_{t_1}^{t_2} \lambda_j L ds \right) M$$

$$= \limsup_L \frac{1}{L} \Pi_j^a (\lambda_j L (t_2 - t_1)) M = \lambda_j M (t_2 - t_1), \quad (74)$$

where we used (72) to get almost sure convergence. We can similarly bound  $\frac{1}{L}D_{\mathbf{k}}^L(t_1, t_2)$  by noting that  $Y_j^L(s) \leq LK$ . Hence, with the stated initialization, the scaled process  $X_{\mathbf{k}}^{L(e)}(t)/L$  converges to a Lipschitz continuous sample path  $x_{\mathbf{k}}^{(e)}(t)$  along the subsequence [16]. Similarly, it can be shown that the fluid limits of processes  $x_{\mathbf{k}}^L(t)$  and  $\hat{x}_{\mathbf{k}}^L(t)$  exist and they are Lipschitz continuous.

Similarly,  $Y_j^L(t)$  increases by at most 1 every time a type- $j$  job arrives and decreases by 1 every time a type- $j$  job in the system departs. Hence, the limit of  $y_j^{(L)}(t)$  also exists by asymptotic Lipschitz continuity.

## E PROOF OF LEMMA 5.9

For each job type  $j$ , the number of type- $j$  jobs in the system is bounded by the number of type- $j$  jobs in an  $M/M/\infty$  system where all arrivals are accepted. This implies that  $y_j(t)$  is also bounded by the fluid limit of type- $j$  jobs in the  $M/M/\infty$  system, i.e.,

$$y_j(t) \leq y_j(0) + \lambda_j t - \int_0^t y_j(s) \mu_j ds. \quad (75)$$

This implies  $y_j(t) \leq \rho_j + (y_j(0) - \rho_j)e^{-\mu_j t}$ . Considering that for any initial state  $\mathbf{z}(0)$ ,  $y_j(0) \leq K$ , we can get that  $y_j(t) < \rho_j + \epsilon_\rho$  if  $t > T_{\epsilon_\rho, j}$  where  $T_{\epsilon_\rho, j} = \frac{-\log \epsilon_\rho + \log K - \rho_j}{\mu_j}$ . Finally, we can choose  $T_{\epsilon_\rho} := \max_{j \in \mathcal{J}} T_{\epsilon_\rho, j}$ .

## F BOUNDS ON THE CHANGE OF SCALED PROCESSES

In this section, we provide a few lemmas which will be used in the proofs later. Their proofs are straightforward and based on concentration inequalities for Poisson distribution.

**LEMMA F.1.** *Consider a time interval  $[\tau_a, \tau_b)$ , and a Poisson process  $N$ , with  $N[\tau_a, \tau_b]$  being the number of events of the process in  $[\tau_a, \tau_b)$ , and function  $f(L)$  as given in Definition 5.8. Then we have:*

*If rate of  $N$  is at least  $L\lambda$  and length of  $[\tau_a, \tau_b)$  is at least  $cf(L)$ ,*

$$\mathbb{P}\left(N[\tau_a, \tau_b] > L\lambda cf(L) + o(Lf(L))\right) \geq 1 - o(L^{-2}). \quad (76)$$

*If  $N$  has rate exactly  $L\lambda$  and length of  $[\tau_a, \tau_b)$  is at least  $cf(L)$ ,*

$$\mathbb{P}\left(N[\tau_a, \tau_b] = L\lambda cf(L) + o(Lf(L))\right) \geq 1 - o(L^{-2}). \quad (77)$$

*Lastly if  $N$  has rate at most  $L\lambda$ , and length of  $[\tau_a, \tau_b)$  is at most  $cf(L)$ ,*

$$\mathbb{P}\left(N[\tau_a, \tau_b] < L\lambda cf(L) + o(Lf(L))\right) \geq 1 - o(L^{-2}). \quad (78)$$

**PROOF.** The proofs of all the cases are based on the tail bounds of Poisson distribution. Specifically, we use the following bounds [15].

For a Poisson random variable  $X$  with mean  $\lambda$  we have

$$\begin{aligned}\mathbb{P}(X - \lambda > x) &\leq \exp\left(-\frac{x^2}{2(\lambda+x)}\right), \\ \mathbb{P}(X - \lambda < -x) &\leq \exp\left(-\frac{x^2}{2(\lambda+x)}\right), \\ \mathbb{P}(|X - \lambda| > x) &\leq 2 \exp\left(-\frac{x^2}{2(\lambda+x)}\right).\end{aligned}$$

Then, in the case that the rate is at least  $L\lambda$  and length of interval is at least  $cf(L)$ , we have that for any  $\epsilon' > 0$

$$\mathbb{P}(N[\tau_a, \tau_b] - cf(L)\lambda L > \epsilon' f(L)L) < \exp\left(-\frac{(\epsilon')^2 f(L)L}{2c\lambda + 2\epsilon'}\right). \quad (79)$$

Last expression is  $o(L^{-2})$  which can be shown by taking its logarithm and using the fact that  $\log L = o(f(L)L)$  by Definition 5.8.

Since  $\epsilon'$  was arbitrary, we eventually get

$$\mathbb{P}(N[\tau_a, \tau_b] > cf(L)\lambda L + o(f(L)L)) < o(L^{-2}). \quad (80)$$

Other cases can be shown in a similar way.  $\square$

LEMMA F.2. Consider a time interval  $[\tau_a, \tau_b] \subset [t_n, t_{n+1}]$ , with  $t_n$  defined in Section 5.3.1. Assume that the interval is of length at most  $\epsilon f(L)$ , for function  $f(L)$  as in Definition 5.8, and constant  $\epsilon > 0$  sufficiently small. Then, with probability at least  $1 - o(L^{-2})$ ,

$$x_{\bar{\mathbf{k}}(i)}^{L(e)}(\tau_b) - x_{\bar{\mathbf{k}}(i)}^{L(e)}(\tau_a) > -B_i \epsilon f(L) + o(f(L)), \quad (81)$$

and

$$q_{\bar{\mathbf{k}}(i),j}^L(\tau_b) - q_{\bar{\mathbf{k}}(i),j}^L(\tau_a) > -B_{i+1} \epsilon L f(L) + o(L f(L)) \quad (82)$$

where  $B_i := (K+1)^{i-1} 2(K\mu_{max} + \sum_{j=1}^J \lambda_j)$ .

PROOF OF LEMMA F.2. The state changes only at arrivals and departures. By definition of  $\tau_n$ ,  $\tau_0 = t$ , and by Lipschitz continuity of  $\mathbf{y}(t)$ , for any time  $\tau \in [\tau_n, \tau_{n+1}]$ , and  $j \in \mathcal{J}$ ,  $y_j^L(\tau) = y_j(t) + O(\epsilon)$ , almost surely, for  $L$  large enough along the subsequence.

Let  $N_{ad}[\tau_a, \tau_b]$  be the number of arrival or departure events of any job type in the interval  $[\tau_a, \tau_b]$ . This process is Poisson with rate at most  $L(\mu_j y_j(t) + O(\epsilon) + \lambda_j) < L\hat{R}$ , where  $\hat{R} := 2(K\mu_{max} + \sum_{j=1}^J \lambda_j)$ . Also  $\epsilon f(L)$  is an upper bound on length of interval  $[\tau_a, \tau_b]$ , so by applying Lemma F.1, we have

$$\mathbb{P}(N_{ad}[\tau_a, \tau_b] - \epsilon \hat{R} f(L)L > o(f(L)L)) < o(L^{-2}). \quad (83)$$

Now suppose the event  $N_{ad}[\tau_a, \tau_b] < \epsilon \hat{R} f(L)L + o(f(L)L)$  holds. The absolute change that occurs to variables  $X_{\bar{\mathbf{k}}(i)}^{L(e)}(\tau)$  after each event is at most  $(K+1)^{i-1}$  according to Lemma D.1, hence

$$\begin{aligned}X_{\bar{\mathbf{k}}(i)}^{L(e)}(\tau_b) - X_{\bar{\mathbf{k}}(i)}^{L(e)}(\tau_a) &\geq -N_{ad}[\tau_a, \tau_b](K+1)^{i-1} \geq \\ &-\hat{R}(K+1)^{i-1} \epsilon L f(L) + o(L f(L)) = -B_i \epsilon L f(L) + o(L f(L)).\end{aligned}$$

Dividing both sides by  $L$  we get (81). Based on (24), we can write

$$\begin{aligned}q_{\bar{\mathbf{k}}(i),j}^L(\tau_b) - q_{\bar{\mathbf{k}}(i),j}^L(\tau_a) &= \\ \sum_{\ell=1}^i \bar{k}_j^{(\ell)} \left( X_{\bar{\mathbf{k}}^{(\ell)}}^{L(e)}(\tau_b) - X_{\bar{\mathbf{k}}^{(\ell)}}^{L(e)}(\tau_a) \right) - Y_j^L(\tau_b) + Y_j^L(\tau_a).\end{aligned} \quad (84)$$

Then (82) follows by considering

$$\sum_{\ell=1}^i \bar{k}_j^{(\ell)} \left( X_{\bar{\mathbf{k}}^{(\ell)}}^{L(e)}(\tau_b) - X_{\bar{\mathbf{k}}^{(\ell)}}^{L(e)}(\tau_a) \right) \geq -K \sum_{\ell=1}^i B_\ell \epsilon L f(L) + o(L f(L)).$$

and

$$-Y_j^L(\tau_b) + Y_j^L(\tau_a) \geq -\hat{R} \epsilon L f(L) + o(L f(L)). \quad (85)$$

$\square$

LEMMA F.3. Consider function  $f(L)$  as in Definition 5.8 and, a time interval  $[\tau_a, \tau_b]$  with length at most  $f(L)$ . Suppose for some  $i \in \{1, \dots, C^{(g)} - 1\}$ , we have that at any time  $\tau \in [\tau_a, \tau_b]$

$$\max_{j \in \mathcal{J}: \bar{k}_j^{(i)} > 0} q_{\bar{\mathbf{k}}^{(i)},j}^L(\tau) < 0. \quad (86)$$

Let  $N_e[\tau_a, \tau_b]$  be the number of times that servers in  $\overline{\text{RG}}(i)$  empty during  $[\tau_a, \tau_b]$ . Then with probability  $1 - o(L^{-2})$ ,

$$\frac{N_e[\tau_a, \tau_b]}{L(\tau_b - \tau_a)} > \frac{\mu_{min}}{JKC^2} \left( 1 - \sum_{\ell=1}^i x_{\bar{\mathbf{k}}^{(\ell)}}^{(e)}(t) \right) + \frac{o(f(L))}{\tau_b - \tau_a}. \quad (87)$$

PROOF. Let  $p := 1 - \sum_{\ell=1}^i x_{\bar{\mathbf{k}}^{(\ell)}}^{(e)}(t)$ , which is strictly positive and is the number of servers without effective configuration in  $\bar{\mathbf{k}}^{(\ell)}$  for  $\ell = 1, \dots, i$ . Notice that due to (86), rank 1 servers in this set will always belong to Reject Group servers  $\overline{\text{RG}}(i)$  (Definition 5.2).

Since there are at most  $C$  different configurations, one of the configurations, say  $\mathbf{k}$ , is assigned to at least  $\frac{p}{C}$  servers without effective configuration in  $\bar{\mathbf{k}}^{(\ell)}$  for  $\ell = 1, \dots, i$  at the fluid limit at time  $t$ , i.e.,  $x_{\mathbf{k}}(t) \geq p' := \frac{p}{C} > 0$ . We also define set  $\mathcal{J}^* := \{j : k_j > 0\}$  with cardinality  $J^*$ .

Due to migrations performed on departure instances, at least one server in  $\overline{\text{RG}}(i)$  gets empty, when, for every  $j \in \mathcal{J}^*$ , the number of type- $j$  jobs that departs is at least the number of type- $j$  jobs that are in the servers of  $\overline{\text{RG}}(i)$ . This is because any type- $j$  departure will create a new empty type- $j$  slot in the servers of  $\overline{\text{RG}}(i)$ , if there is a type- $j$  job in any of them. Hence one of the servers in this set will empty after at most  $CK$  jobs of each job type in  $\mathcal{J}^*$  depart, where  $CK$  is an upper bound on the total number of jobs of any type that can be in Reject Group servers. Then we need to bound

$$\mathbb{P}\left(N_e[\tau_a, \tau_b] < \frac{p' \mu_{min} L}{JKC} (\tau_b - \tau_a) - \epsilon L f(L)\right). \quad (88)$$

If  $\tau_b - \tau_a < \frac{\epsilon f(L) J K C}{p' \mu_{min}}$ , then

$$\begin{aligned}\mathbb{P}\left(N_e[\tau_a, \tau_b] - \frac{p' \mu_{min} L}{J} (\tau_b - \tau_a) < -\epsilon K C L f(L)\right) &\leq \\ \mathbb{P}(N_e[\tau_a, \tau_b] < 0) &= 0.\end{aligned}$$

If instead  $\tau_b - \tau_a \geq \frac{\epsilon f(L) J K C}{p' \mu_{min}}$ , then we consider the counting process  $D_{\mathcal{J}^*}(\tau)$  defined as follows.  $D_{\mathcal{J}^*}(\tau_a) = 0$  and it is incremented at times  $\tau^{(i)}$  for  $i \in \mathbb{Z}^+$  if  $\tau^{(i)}$  is the first time since  $\tau^{(i-1)}$  at which at least one departure of type  $j$  occurred for all  $j \in \mathcal{J}^*$  and as a convention  $\tau^{(0)} = \tau_a$ . Based on arguments so far, the process  $N_e[\tau_a, \tau_b]$  will increment by at least 1 between times  $\tau^{(i)}$  and  $\tau^{(i+CK)}$ , for  $i \geq 0$ , i.e.,  $N_e[\tau_a, \tau_b] \geq \left\lfloor \frac{D_{\mathcal{J}^*}(\tau_b)}{CK} \right\rfloor$ . Then applying Lemma F.1 to process  $D_{\mathcal{J}^*}$ , which has rate at least  $\frac{p' \mu_{min} L}{J}$ , we get

$$\begin{aligned}\mathbb{P}\left(N_e[\tau_a, \tau_b] - \frac{p' \mu_{min} L}{JCK} (\tau_b - \tau_a) < -\epsilon K C L f(L)\right) &\leq \\ \mathbb{P}\left(\left\lfloor \frac{D_{\mathcal{J}^*}(\tau_b)}{CK} \right\rfloor - \frac{p' \mu_{min} L}{JCK} (\tau_b - \tau_a) < -\epsilon K C L f(L)\right) &= o(L^{-2}).\end{aligned}$$

We have thus proven, since  $\epsilon$  can be arbitrarily close to 0, that with probability  $1 - o(L^{-2})$

$$N_e[\tau_a, \tau_b] > \frac{\rho' \mu_{\min} L}{JCK} (\tau_b - \tau_a) + o(Lf(L)), \quad (89)$$

which implies (87).  $\blacksquare$

## G PROOF OF PROPOSITION 5.10

We prove Proposition 5.10 for the following values of parameters:

- $\alpha_i, i \in \{1, \dots, C_\rho^{(g)} - 1\}$ , is given by

$$\alpha_i := \delta \frac{\mu_{\min} v^{i - C_\rho^{(g)}} v_i}{\mu_{\max} 8K^2 C}, \quad (90)$$

where  $K$  is the maximum number of jobs in a server,  $\mu_{\min} := \min_{j \in \mathcal{J}} \mu_j$ ,  $\mu_{\max} := \max_{j \in \mathcal{J}} \mu_j$ ,  $v := 12K \frac{\mu_{\max}}{\mu_{\min}}$ ,  $v_i := 1$  if  $i \leq C_\alpha(t)$  else  $v_i := K$  and  $\delta$  is a positive constant sufficiently small such that

$$\delta < \frac{1}{2JC} x_{\bar{k}^{(C_\rho^{(g)})}}^{(g)} \quad (91)$$

and for any two indexes  $j_a, j_b \in \{1, \dots, C^{(g)}\}$ ,

$$\rho_{j_a} > \sum_{i=1}^{j_b} x_{\bar{k}^{(i)}}^{(g)} \bar{k}_{j_a}^{(i)} + \delta, \text{ or } \rho_{j_a} = \sum_{i=1}^{j_b} x_{\bar{k}^{(i)}}^{(g)} \bar{k}_{j_a}^{(i)}. \quad (92)$$

- $\epsilon_\rho$  is chosen as

$$\epsilon_\rho = \frac{\alpha_1}{2}. \quad (93)$$

Before presenting the main proof, we state a few lemmas.

LEMMA G.1. *The fraction of servers without effective configuration in the set  $\{\bar{k}^{(i)} : i = 1, \dots, C_\alpha(t)\}$ , with  $C_\alpha(t)$  as in Definition 5.7, is at least  $\frac{1}{2} x_{\bar{k}^{(C_\rho^{(g)})}}^{(g)}$ .*

PROOF. The bound can be inferred as follows

$$1 - \sum_{\ell=1}^i x_{\bar{k}^{(\ell)}}^{(e)}(t) \geq x_{\bar{k}^{(C_\rho^{(g)})}}^{(g)} + \sum_{\ell=1}^i \left( x_{\bar{k}^{(\ell)}}^{(g)} - x_{\bar{k}^{(\ell)}}^{(e)}(t) \right) > x_{\bar{k}^{(C_\rho^{(g)})}}^{(g)} - \sum_{\ell=1}^i \alpha_\ell > x_{\bar{k}^{(C_\rho^{(g)})}}^{(g)} - \sum_{\ell=1}^i \frac{\delta}{2C_\rho^{(g)}} > x_{\bar{k}^{(C_\rho^{(g)})}}^{(g)} / 2.$$

To get this result we used that  $i \leq C_\alpha(t) < C_\rho^{(g)}$ ,  $\sum_{\ell=1}^{C_\rho^{(g)}} x_{\bar{k}^{(\ell)}}^{(g)} = 1$ ,  $\alpha_i < \frac{\delta}{2C_\rho^{(g)}}$  from (90) and (91).  $\blacksquare$

LEMMA G.2. *Consider an interval  $[\tau_a, \tau_b] \subseteq [\tau_n^{(m)}, \tau_n^{(m+1)})$ , with  $\tau_n^{(m)}$  being defined in Section 5.3.1 and the corresponding driving set of indexes  $\bar{\mathcal{J}}[m] = \{j_i : i = 1, \dots, G\}$ . If the following holds for  $i = 1, \dots, G$*

$$\sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} \nabla x_{\bar{k}^{(\ell)}}^{L(e)}[\tau_a, \tau_b] = \lambda_{j_i} - \mu_{j_i} y_{j_i}(t) + \frac{o(f(L))}{\tau_b - \tau_a}, \quad (94)$$

then

$$\nabla x_{\bar{k}^{(i)}}^{L(e)}[\tau_a, \tau_b] = h^{\bar{\mathcal{J}},(i)}(t) + \frac{o(f(L))}{\tau_b - \tau_a}, \quad (95)$$

PROOF. The proof is by induction. For  $i = 1$  we have

$$\nabla x_{\bar{k}^{(1)}}^{L(e)}[\tau_a, \tau_b] = \frac{\lambda_{j_1} - \mu_{j_1} y_{j_1}(t)}{\bar{k}_{j_1}^{(1)}} + \frac{o(f(L))}{\tau_b - \tau_a} = h^{\bar{\mathcal{J}},(1)}(t) + \frac{o(f(L))}{\tau_b - \tau_a}.$$

Now assume that for every  $\ell \in \{1, \dots, i-1\}$ ,

$$\nabla x_{\bar{k}^{(\ell)}}^{L(e)}[\tau_a, \tau_b] = h^{\bar{\mathcal{J}},(\ell)}(t) + \frac{o(f(L))}{\tau_b - \tau_a}.$$

Then (94) implies

$$\begin{aligned} \nabla x_{\bar{k}^{(i)}}^{L(e)}[\tau_a, \tau_b] &= \frac{\lambda_{j_i} - \mu_{j_i} y_{j_i}(t) - \sum_{\ell=1}^{i-1} \bar{k}_{j_i}^{(\ell)} \nabla x_{\bar{k}^{(\ell)}}^{L(e)}[\tau_a, \tau_b]}{\bar{k}_{j_i}^{(i)}} + \frac{o(f(L))}{\tau_b - \tau_a} \stackrel{(a)}{=} \\ &= \frac{\lambda_{j_i} - \mu_{j_i} y_{j_i}(t) - \sum_{\ell=1}^{i-1} \bar{k}_{j_i}^{(\ell)} h^{\bar{\mathcal{J}},(\ell)}(t)}{\bar{k}_{j_i}^{(i)}} + \frac{o(f(L))}{\tau_b - \tau_a} = \\ &= h^{\bar{\mathcal{J}},(i)}(t) + \frac{o(f(L))}{\tau_b - \tau_a}. \end{aligned}$$

In (a), we used the fact that sum of a finite number of  $\frac{o(f(L))}{\tau_b - \tau_a}$  terms is still  $\frac{o(f(L))}{\tau_b - \tau_a}$ .  $\blacksquare$

LEMMA G.3. *Suppose  $\mathbf{z}(t) \in \Gamma[\epsilon_\rho]$ , and for every  $i \in \{1, \dots, G\}$ ,  $h^{\bar{\mathcal{J}}[m],(i)}(t)$  is defined as in (37), and  $G \leq C_\alpha(t)$  with  $C_\alpha(t)$  as in Definition 5.7. Then*

$$|h^{\bar{\mathcal{J}}[m],(i)}(t)| < \mu_{\max} \alpha_i + \mu_{\max} \sum_{\ell=1}^{i-1} 2(K+1)^{i-\ell} \alpha_\ell. \quad (96)$$

PROOF. Let  $\bar{\mathcal{J}} = \bar{\mathcal{J}}[m]$ . The proof is by induction.

**Base Case:** It suffices to show that  $h^{\bar{\mathcal{J}},(1)}(t) < \mu_{\max} \alpha_1$  and  $h^{\bar{\mathcal{J}},(1)}(t) > -\mu_{\max} \alpha_1$ .

To show  $h^{\bar{\mathcal{J}},(1)}(t) < \mu_{\max} \alpha_1$ , consider  $j_1$  which is the first index of  $\bar{\mathcal{J}}$ , then

$$h^{\bar{\mathcal{J}},(1)}(t) = \frac{\lambda_{j_1} - \mu_{j_1} y_{j_1}(t)}{\bar{k}_{j_1}^{(1)}} \stackrel{(a)}{=} \mu_{j_1} \left( x_{\bar{k}^{(1)}}^{(g)} - x_{\bar{k}^{(1)}}^{(e)}(t) \right) < \mu_{\max} \alpha_1.$$

In (a), we used (33) for  $i = 1$ , according to which  $\rho_{j_1} = \bar{k}_{j_1}^{(1)} x_{\bar{k}^{(1)}}^{(g)}$  and  $y_{j_1}(t) = \bar{k}_{j_1}^{(1)} x_{\bar{k}^{(1)}}^{(e)}(t)$ .

To show  $h^{\bar{\mathcal{J}},(1)}(t) > -\mu_{\max} \alpha_1$ , we use the fact that  $y_{j_1}(t) - \rho_{j_1} < \epsilon_\rho$  (since  $\mathbf{z}(t) \in \Gamma[\epsilon_\rho]$ ) which can be applied as

$$h^{\bar{\mathcal{J}},(1)}(t) = \frac{\lambda_{j_1} - \mu_{j_1} y_{j_1}(t)}{\bar{k}_{j_1}^{(1)}} > -\frac{\mu_{j_1} \epsilon_\rho}{\bar{k}_{j_1}^{(1)}} \stackrel{(a)}{\geq} -\mu_{\max} \alpha_1. \quad (97)$$

In (a), we used that  $\epsilon_\rho < \alpha_1$ , which is due to (93).

**Inductive Case:** We assume (96) is true for all indexes up to  $i-1$ . Then we can upper and lower bound  $h^{\bar{\mathcal{J}},(i)}(t)$  as follows.

To show  $h^{\bar{\mathcal{J}},(i)}(t) < \mu_{\max}\alpha_i + \mu_{\max} \sum_{\ell=1}^{i-1} 2(K+1)^{i-\ell}\alpha_\ell$ , let  $j_i$  be the  $i$ th index in  $\bar{\mathcal{J}}$ , then

$$\begin{aligned} h^{\bar{\mathcal{J}},(i)}(t) &= \frac{\lambda_{j_i} - \mu_{j_i}y_{j_i}(t) - \sum_{\ell=1}^{i-1} \bar{k}_{j_i}^{(\ell)} h^{\bar{\mathcal{J}},(\ell)}(t)}{\bar{k}_{j_i}^{(i)}} \stackrel{(a)}{=} \\ &= \frac{\mu_{j_i} \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} (x_{\bar{k}^{(\ell)}}^{(g)} - x_{\bar{k}^{(\ell)}}^{(e)}(t)) - \sum_{\ell=1}^{i-1} \bar{k}_{j_i}^{(\ell)} h^{\bar{\mathcal{J}},(\ell)}(t)}{\bar{k}_{j_i}^{(i)}} < \\ &= \mu_{\max}\alpha_i + \mu_{\max} \sum_{\ell=1}^{i-1} K\alpha_\ell + \sum_{\ell=1}^{i-1} K\mu_{\max}\alpha_\ell + \\ &= \sum_{\ell=1}^{i-1} K\mu_{\max} \sum_{\ell'=1}^{\ell-1} 2(1+K)^{\ell-\ell'}\alpha_{\ell'} \leq \mu_{\max}\alpha_i + 2\mu_{\max} \sum_{\ell=1}^{i-1} (1+K)^{i-\ell}\alpha_\ell \end{aligned}$$

In (a), we used (33), according to which  $\rho_{j_i} = \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} x_{\bar{k}^{(\ell)}}^{(g)}$  and  $y_{j_i}(t) = \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} x_{\bar{k}^{(\ell)}}^{(e)}(t)$ . The rest of the inequalities come from recursive application of (96) and algebraic manipulations.

To show  $h^{\bar{\mathcal{J}},(i)}(t) > -\mu_{\max}\alpha_i - \mu_{\max} \sum_{\ell=1}^{i-1} 2(K+1)^{i-\ell}\alpha_\ell$ , we use  $y_{j_i}(t) - \rho_{j_i} < \epsilon_\rho$  as follows,

$$\begin{aligned} h^{\bar{\mathcal{J}},(i)}(t) &= \frac{\lambda_{j_i} - \mu_{j_i}y_{j_i}(t) - \sum_{\ell=1}^{i-1} \bar{k}_{j_i}^{(\ell)} h^{\bar{\mathcal{J}},(\ell)}(t)}{\bar{k}_{j_i}^{(i)}} = \\ &= \frac{\lambda_{j_i} - \mu_{j_i}y_{j_i}(t) - \sum_{\ell=1}^{i-1} \bar{k}_{j_i}^{(\ell)} h^{\bar{\mathcal{J}},(\ell)}(t)}{\bar{k}_{j_i}^{(i)}} > \\ &= -\mu_{\max}\alpha_i - \sum_{\ell=1}^{i-1} K\mu_{\max}\alpha_\ell - \sum_{\ell=1}^{i-1} K\mu_{\max} \sum_{\ell'=1}^{\ell-1} 2(1+K)^{\ell-\ell'}\alpha_{\ell'} > \\ &= -\mu_{\max}\alpha_i - 2\mu_{\max} \sum_{\ell=1}^{i-1} (1+K)^{i-\ell}\alpha_\ell. \end{aligned}$$

In particular for (a) we used that  $\epsilon_\rho < \alpha_1 < \alpha_i$ , which is due to (93) and (90). We also made recursive use of (96). ■

LEMMA G.4. *If  $\mathbf{z}(t) \in \Gamma[\epsilon_\rho]$ , then for any  $i \in \{1, \dots, C_\alpha(t) + 1\}$ :*

$$x_{\bar{k}^{(i)}}^{(g)} - x_{\bar{k}^{(i)}}^{(e)}(t) > -\epsilon_\rho - K \sum_{\ell=1}^{i-1} \alpha_\ell. \quad (98)$$

PROOF. Since  $i-1 \leq C_\alpha(t) < C_\rho^{(g)}$  then for  $j = \sigma_{i-1}$ ,

$$\rho_j = \sum_{\ell=1}^{i-1} \bar{k}_j^{(\ell)} x_{\bar{k}^{(\ell)}}^{(g)}. \quad (99)$$

Considering that  $j$  next, we can prove (98) as follows.

$$\begin{aligned} x_{\bar{k}^{(i)}}^{(g)} - x_{\bar{k}^{(i)}}^{(e)}(t) &\stackrel{(a)}{\geq} \frac{\rho_j - y_j(t) - \sum_{\ell=1}^{i-1} \bar{k}_j^{(\ell)} (x_{\bar{k}^{(\ell)}}^{(g)} - x_{\bar{k}^{(\ell)}}^{(e)}(t))}{\bar{k}_j^{(i)}} \stackrel{(b)}{\geq} \\ &= \frac{-\epsilon_\rho - \sum_{\ell=1}^{i-1} \bar{k}_j^{(\ell)} \alpha_\ell}{\bar{k}_j^{(i)}} \geq -\epsilon_\rho - K \sum_{\ell=1}^{i-1} \alpha_\ell. \end{aligned}$$

In (a) we used (99) and the fact that  $\sum_{\ell=1}^{i-1} \bar{k}_j^{(\ell)} x_{\bar{k}^{(\ell)}}^{(e)}(t) \leq y_j(t)$ . In (b) we used  $\rho_j - y_j(t) \geq -\epsilon_\rho$  as implied by (41), and  $-x_{\bar{k}^{(\ell)}}^{(g)} + x_{\bar{k}^{(\ell)}}^{(e)}(t) > -\alpha_\ell$  as implied by (34) for  $\ell \leq i-1 \leq C_\alpha(t)$ . □

#### MAIN PROOF OF PROPOSITION 5.10:

As a reminder in what follows we will use the notations  $\bar{\mathcal{J}}[m] = \{j_i : i = 1, \dots, G_m\}$  and  $\ell_m := G_m + 1$ , given in description of Proposition. If not clear from context, we will make the association of  $j_i$  with  $m$  explicit using also the notation  $j_i[m]$ . Notice that for any  $m \in \{0, \dots, M_n - 1\}$ ,  $\ell_m \leq C_\alpha(t)$ .

**Proof of Property P.1:** This property follows from two claims.

*Claim 1.* Consider  $m \in \{0, \dots, M_n - 1\}$ . If for every  $i \in \{1, \dots, \ell_m - 1\}$ ,  $q_{\bar{k}^{(i)}, j_i[m]}^L(\tau_n^{(m)}) = o(Lf(L))$  with probability  $1 - o(L^{-2})$ , then (42) holds with probability  $1 - o(L^{-2})$ .

*Claim 2.* For every  $m \in \{0, \dots, M_n - 1\}$  and for every  $i \in \{1, \dots, \ell_m - 1\}$ , we have that  $q_{\bar{k}^{(i)}, j_i[m]}^L(\tau_n^{(m)}) = o(Lf(L))$  with probability  $1 - o(L^{-2})$ .

#### Proof of Claim 1:

**Base Case  $i = 1$ :**

Let  $j_1 = j_1[m]$ . For (42) to be true, it suffices to prove, that for any time  $\tau \in (\tau_n^{(m)}, \tau_n^{(m+1)})$ ,

$$\bar{k}_{j_1}^{(1)} \nabla x_{\bar{k}^{(1)}}^{L(e)}[\tau_n^{(m)}, \tau] = \lambda_{j_1} - \mu_{j_1}y_{j_1}(t) + \frac{o(f(L))}{\tau - \tau_n^{(m)}}, \quad (100)$$

with probability  $1 - o(L^{-2})$  or equivalently, for any  $\epsilon > 0$ ,

$$\begin{aligned} \mathbb{P} \left( \left| \bar{k}_{j_1}^{(1)} (x_{\bar{k}^{(1)}}^{L(e)}(\tau) - x_{\bar{k}^{(1)}}^{L(e)}(\tau_n^{(m)})) \right. \right. \\ \left. \left. - (\lambda_{j_1} - \mu_{j_1}y_{j_1}(t))(\tau - \tau_n^{(m)}) \right| \leq \epsilon f(L) \right) > 1 - o(L^{-2}). \end{aligned} \quad (101)$$

For a given  $\tau$ , consider  $\tau'$  to be the latest time in  $[\tau_n^{(m)}, \tau]$  such that

$$\max_{j \in \mathcal{J}: \bar{k}_j^{(1)} > 0} q_{\bar{k}^{(1)}, j}^L(\tau') \geq 0. \quad (102)$$

This time always exists since (102) holds for  $\tau' = \tau_n^{(m)}$ . To prove (101) then, it is sufficient to prove

$$\begin{aligned} \mathbb{P} \left( \left| \bar{k}_{j_1}^{(1)} (x_{\bar{k}^{(1)}}^{L(e)}(\tau') - x_{\bar{k}^{(1)}}^{L(e)}(\tau_n^{(m)})) \right. \right. \\ \left. \left. - (\lambda_{j_1} - \mu_{j_1}y_{j_1}(t))(\tau' - \tau_n^{(m)}) \right| \leq \frac{\epsilon f(L)}{2} \right) = 1 - o(L^{-2}), \end{aligned} \quad (103)$$

and

$$\begin{aligned} \mathbb{P} \left( \left| \bar{k}_{j_1}^{(1)} (x_{\bar{k}^{(1)}}^{L(e)}(\tau) - x_{\bar{k}^{(1)}}^{L(e)}(\tau')) \right. \right. \\ \left. \left. - (\lambda_{j_1} - \mu_{j_1}y_{j_1}(t))(\tau - \tau') \right| \leq \frac{\epsilon f(L)}{2} \right) = 1 - o(L^{-2}). \end{aligned} \quad (104)$$

**Proof of (103):** We will now prove (103) by considering two cases depending on length of  $\tau' - \tau_n^{(m)}$ .

We consider

$$\tau' - \tau_n^{(m)} \leq \frac{\epsilon f(L)}{4KR} \quad (105)$$

or

$$\tau' - \tau_n^{(m)} > \frac{\epsilon f(L)}{4KR}, \quad (106)$$

where  $R := \sum_{j \in \mathcal{J}} \lambda_j + K\mu_{\max}$

**Case (105):** We notice  $x_{\bar{k}^{(1)}}^{L(e)}(\tau)$  will change by at most  $1/L$  at each arrival or departure according to Lemma D.1 and thus  $\bar{k}_{j_1}^{(1)} x_{\bar{k}^{(1)}}^{L(e)}(\tau)$  will change by at most  $K/L$ .

The number of arrivals and departures in  $[\tau_n^{(m)}, \tau']$  is stochastically bounded by a Poisson Process of rate  $\left(\sum_{j \in \mathcal{J}} \lambda_j + K\mu_{max}\right)L$  on an interval of length at most  $\frac{\epsilon f(L)}{4KR}$ , which, according to Lemma F.1, with probability  $1 - o(L^{-2})$  is at most

$$\left(\sum_{j \in \mathcal{J}} \lambda_j + K\mu_{max}\right)L \frac{\epsilon f(L)}{4KR} + o(Lf(L)) = \frac{\epsilon Lf(L)}{4K} + o(Lf(L)),$$

therefore,

$$\mathbb{P}\left(\bar{k}_{j_1}^{(1)} \left(x_{\bar{k}^{(1)}}^{L(e)}(\tau') - x_{\bar{k}^{(1)}}^{L(e)}(\tau_n^{(m)})\right) \leq \frac{\epsilon f(L)}{4}\right) \geq 1 - o(L^{-2}). \quad (107)$$

Considering (105) holds, we also have

$$\mathbb{P}\left((\lambda_{j_1} - \mu_{j_1} y_{j_1}(t)) (\tau' - \tau_n^{(m)}) \leq \frac{\epsilon f(L)}{4}\right) = 1 \quad (108)$$

It is now easy to verify that equations (107) and (108) imply (103).

**Case (106):** In this case we notice using Lemma F.1 for the process of jobs of type  $j_1$  in the system which is Poisson with rate  $L(\lambda_{j_1} - \mu_{j_1} y_{j_1}(t))$ , that with probability  $1 - o(L^{-2})$

$$\begin{aligned} o(Lf(L)) &= q_{\bar{k}^{(1)}, j_1}^L(\tau_n^{(m)}) \\ &= q_{\bar{k}^{(1)}, j_1}^L(\tau') - \bar{k}_{j_1}^{(1)} L(x_{\bar{k}^{(1)}}^{L(e)}(\tau') - x_{\bar{k}^{(1)}}^{L(e)}(\tau_n^{(m)})) \\ &\quad + Ly_{j_1}^L(\tau') - Ly_{j_1}^L(\tau_n^{(m)}) \\ &\leq K - \bar{k}_{j_1}^{(1)} L(x_{\bar{k}^{(1)}}^{L(e)}(\tau') - x_{\bar{k}^{(1)}}^{L(e)}(\tau_n^{(m)})) \\ &\quad + L(\lambda_{j_1} - \mu_{j_1} y_{j_1}(t))(\tau' - \tau_n^{(m)}) + o(Lf(L)), \end{aligned}$$

or equivalently, since trivially  $K = o(Lf(L))$ ,

$$\begin{aligned} \frac{x_{\bar{k}^{(1)}}^{L(e)}(\tau') - x_{\bar{k}^{(1)}}^{L(e)}(\tau_n^{(m)})}{\tau' - \tau_n^{(m)}} &\leq \\ \frac{1}{\bar{k}_{j_1}^{(1)}} (\lambda_{j_1} - \mu_{j_1} y_{j_1}(t)) + \frac{o(f(L))}{\tau' - \tau_n^{(m)}} &\stackrel{(a)}{=} \\ \frac{1}{\bar{k}_{j_1}^{(1)}} (\lambda_{j_1} - \mu_{j_1} y_{j_1}(t)) + o(1), & \end{aligned} \quad (109)$$

where in (a) we just used (106). Let

$$j' := \arg \max_{j \in \mathcal{J}: \bar{k}_j^{(1)} > 0} q_{\bar{k}^{(1)}, j}^L(\tau'). \quad (110)$$

Then we also have, using Lemma F.1 for the process of jobs of type  $j'$  in the system which is Poisson with rate  $L(\lambda_{j'} - \mu_{j'} y_{j'}(t))$ , that

with probability  $1 - o(L^{-2})$

$$\begin{aligned} 0 &\leq q_{\bar{k}^{(1)}, j'}^L(\tau') = q_{\bar{k}^{(1)}, j'}^L(\tau_n^{(m)}) \\ &\quad + \bar{k}_{j'}^{(1)} L(x_{\bar{k}^{(1)}}^{L(e)}(\tau') - x_{\bar{k}^{(1)}}^{L(e)}(\tau_n^{(m)})) - Ly_{j'}^L(\tau') + Ly_{j'}^L(\tau_n^{(m)}) \\ &\leq K + \bar{k}_{j'}^{(1)} L(x_{\bar{k}^{(1)}}^{L(e)}(\tau') - x_{\bar{k}^{(1)}}^{L(e)}(\tau_n^{(m)})) \\ &\quad - L(\lambda_{j'} - \mu_{j'} y_{j'}(t))(\tau' - \tau_n^{(m)}) + o(Lf(L)), \end{aligned}$$

from which, since trivially  $K = o(Lf(L))$ , it follows

$$\frac{x_{\bar{k}^{(1)}}^{L(e)}(\tau') - x_{\bar{k}^{(1)}}^{L(e)}(\tau_n^{(m)})}{\tau' - \tau_n^{(m)}} \geq \frac{\lambda_{j'} - \mu_{j'} y_{j'}(t)}{\bar{k}_{j'}^{(1)}} + o(1). \quad (111)$$

Considering that (38) does not hold for index  $j = j'$  and  $G_m = 1$ , we get

$$\frac{\lambda_{j'} - \mu_{j'} y_{j'}(t)}{\bar{k}_{j'}^{(1)}} \geq h\bar{\mathcal{T}}[m, (1)](t) = \frac{\lambda_{j_1} - \mu_{j_1} y_{j_1}(t)}{\bar{k}_{j_1}^{(1)}}. \quad (112)$$

From (109), (111) and (112) we get

$$\frac{x_{\bar{k}^{(1)}}^{L(e)}(\tau') - x_{\bar{k}^{(1)}}^{L(e)}(\tau_n^{(m)})}{\tau' - \tau_n^{(m)}} = \frac{\lambda_{j_1} - \mu_{j_1} y_{j_1}(t)}{\bar{k}_{j_1}^{(1)}} + o(1), \quad (113)$$

which holds with probability  $1 - o(L^{-2})$  and therefore it implies (103).

**Proof of (104):** We will now prove (104) by considering two cases depending on length of  $\tau - \tau'$ .

We consider

$$\tau - \tau' \leq \frac{\epsilon f(L)}{4KR} \quad (114)$$

or

$$\tau - \tau' > \frac{\epsilon f(L)}{4KR}, \quad (115)$$

where  $R := \sum_{j \in \mathcal{J}} \lambda_j + K\mu_{max}$ .

We further assume that

$$\epsilon < \frac{\mu_{j_1} \alpha_1}{4KR}. \quad (116)$$

**Case (114):** Following the same arguments as in the case of (105) we can infer the equivalent of (107) and (108) for interval  $(\tau', \tau)$ , i.e.

$$\mathbb{P}\left(\bar{k}_{j_1}^{(1)} \left(x_{\bar{k}^{(1)}}^{L(e)}(\tau) - x_{\bar{k}^{(1)}}^{L(e)}(\tau')\right) \leq \frac{\epsilon f(L)}{4}\right) \geq 1 - o(L^{-2}) \quad (117)$$

and

$$\mathbb{P}\left((\lambda_{j_1} - \mu_{j_1} y_{j_1}(t)) (\tau - \tau') \leq \frac{\epsilon f(L)}{4}\right) = 1, \quad (118)$$

which imply (104).

**Case (115):** First we will prove that

$$q_{\bar{k}^{(1)}, j_1}^L(\tau') \geq -\epsilon^2 Lf(L) + o(Lf(L)), \quad (119)$$

or equivalently

$$1/Lq_{\bar{k}^{(1)}, j_1}^L(\tau') - 1/Lq_{\bar{k}^{(1)}, j_1}^L(\tau_n^{(m)}) \geq -\epsilon^2 f(L) + o(f(L)), \quad (120)$$

which both will hold with probability  $1 - o(L^{-2})$ . The analysis for this is same as with the proof of (103) so we highlight only the parts that are different.

- Instead of considering cases  $\tau' - \tau_n^{(m)} \leq \frac{\epsilon f(L)}{4KR}$  and  $\tau' - \tau_n^{(m)} > \frac{\epsilon f(L)}{4KR}$ , we should consider  $\tau' - \tau_n^{(m)} \leq \frac{\epsilon^2 f(L)}{2(K+1)R}$  and  $\tau' - \tau_n^{(m)} > \frac{\epsilon^2 f(L)}{2(K+1)R}$ .
- If interval is short, we can bound the absolute change of variable  $1/Lq_{\bar{k}^{(1)}, j_1}^L(\tau)$  which changes by at most  $(K+1)/L$  after each arrival or departure.
- If interval is long, we can still prove that the equivalent of (109) is satisfied as equality and considering  $y_{j_1}^L(\tau') - y_{j_1}^L(\tau_n^{(m)}) = \lambda_{j_1} - \mu_{j_1} y_{j_1}(t) + o(f(L))$ , we can get through (109) that  $1/Lq_{\bar{k}^{(1)}, j_1}^L(\tau') = o(f(L))$  with probability  $1 - o(L^{-2})$ .

In this case, because of (119) we get

$$\begin{aligned} o(Lf(L)) - \epsilon^2 Lf(L) &\leq q_{\bar{k}^{(1)}, j_1}^L(\tau') = q_{\bar{k}^{(1)}, j_1}^L(\tau) \\ &- \bar{k}_{j_1}^{(1)} L \left( x_{\bar{k}^{(1)}}^{L(e)}(\tau) - x_{\bar{k}^{(1)}}^{L(e)}(\tau') \right) + Ly_{j_1}^L(\tau) - Ly_{j_1}^L(\tau') \leq \\ K - \bar{k}_{j_1}^{(1)} L \left( x_{\bar{k}^{(1)}}^{L(e)}(\tau) - x_{\bar{k}^{(1)}}^{L(e)}(\tau') \right) \\ &+ L(\lambda_{j_1} - \mu_{j_1} y_{j_1}(t))(\tau - \tau') + o(Lf(L)), \end{aligned} \quad (121)$$

from which, after considering  $K = o(Lf(L))$ , it follows that

$$\begin{aligned} \frac{x_{\bar{k}^{(1)}}^{L(e)}(\tau) - x_{\bar{k}^{(1)}}^{L(e)}(\tau')}{\tau - \tau'} &\leq \frac{\lambda_{j_1} - \mu_{j_1} y_{j_1}(t)}{\bar{k}_{j_1}^{(1)}} \\ &+ \frac{\epsilon^2 f(L)}{\bar{k}_{j_1}^{(1)}(\tau - \tau')} + \frac{o(f(L))}{\tau - \tau'} \stackrel{(a)}{\leq} \\ \mu_{j_1} (x_{\bar{k}^{(1)}}^{(g)} - x_{\bar{k}^{(1)}}^{(e)}(t)) + \frac{4K\epsilon R}{\bar{k}_{j_1}^{(1)}} + o(1) &\stackrel{(b)}{<} \\ \mu_{j_1} \alpha_1 + \mu_{j_1} \alpha_1 + o(1) &< 2\mu_{max} \alpha_1 + o(1). \end{aligned} \quad (122)$$

In (a) we applied (115) and properties of  $j_1$  which come from (28) and (33) for  $i = 1$ , while in (b) we used (34) and (116).

Also from Lemma F.3 and given that all the servers that empty during  $[\tau', \tau]$  will be assigned to configuration  $\bar{k}^{(1)}$ , we will have that with probability  $1 - o(L^{-2})$ ,

$$x_{\bar{k}^{(1)}}^{L(e)}(\tau) - x_{\bar{k}^{(1)}}^{L(e)}(\tau') \geq \frac{\mu_{min}}{JKC^2} \left( 1 - x_{\bar{k}^{(1)}}^{(e)}(t) \right) (\tau - \tau') + o(f(L)) \quad (123)$$

or equivalently,

$$\begin{aligned} \frac{x_{\bar{k}^{(1)}}^{L(e)}(\tau) - x_{\bar{k}^{(1)}}^{L(e)}(\tau')}{\tau - \tau'} &\geq \frac{\mu_{min}}{JKC^2} \left( 1 - x_{\bar{k}^{(1)}}^{(e)}(t) \right) + o(1) > \\ \frac{\mu_{min} x_{\bar{k}^{(c_p)}}^{(g)}}{2JKC^2} + o(1), \end{aligned} \quad (124)$$

where (a) is due to Lemma G.1. So far we proved that if (115) holds, (122) and (124) also hold.

However, considering (90) we have

$$2\mu_{max} \alpha_1 < \frac{\mu_{min} x_{\bar{k}^{(c_p)}}^{(g)}}{2JKC^2} \quad (125)$$

and because of that, the probability that (122) and (124) are both true is  $o(L^{-2})$ . This means that (114) holds with probability at least

$1 - o(L^{-2})$ , and thus the analysis of (114) is sufficient for (104) to hold.

**Inductive Case  $i > 1$ :** For  $i \in \{1, \dots, \ell_m\}$  we have according to the assumptions of Claim 2 that  $q_{\bar{k}^{(i)}, j_i[m]}^L(\tau_n^{(m)}) = o(Lf(L))$  with probability  $1 - o(L^{-2})$ .

What we need to prove is that under this assumption, for any time  $\tau \in (\tau_n^{(m)}, \tau_n^{(m+1)})$ ,  $i \in \{1, \dots, \ell_m\}$  and  $j_i := j_i[m]$ ,

$$\sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} \nabla x_{\bar{k}^{(\ell)}}^{L(e)}[\tau_n^{(m)}, \tau] = \lambda_{j_i} - \mu_{j_i} y_{j_i}(t) + \frac{o(f(L))}{\tau - \tau_n^{(m)}}. \quad (126)$$

with probability  $1 - o(L^{-2})$ , so it is sufficient to only consider the case  $q_{\bar{k}^{(i)}, j_i[m]}^L(\tau_n^{(m)}) = o(Lf(L))$ . Equivalently, it suffices to show that for any  $\epsilon > 0$ ,

$$\begin{aligned} \mathbb{P} \left( \left| \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} (x_{\bar{k}^{(\ell)}}^{L(e)}(\tau) - x_{\bar{k}^{(\ell)}}^{L(e)}(\tau_n^{(m)})) \right. \right. \\ \left. \left. - (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t))(\tau - \tau_n^{(m)}) \right| \leq \epsilon f(L) \right) \geq 1 - o(L^{-2}). \end{aligned} \quad (127)$$

For a given  $\tau$ , let  $\tau'$  to be the latest time in  $(\tau_n^{(m)}, \tau)$  such that

$$\max_{j \in \mathcal{J}: \bar{k}_j^{(i)} > 0} q_{\bar{k}^{(i)}, j}^L \geq 0. \quad (128)$$

Using the same argument as in the base case, this time always exists. To prove (127) then, it is sufficient to prove

$$\begin{aligned} \mathbb{P} \left( \left| \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} (x_{\bar{k}^{(\ell)}}^{L(e)}(\tau') - x_{\bar{k}^{(\ell)}}^{L(e)}(\tau_n^{(m)})) \right. \right. \\ \left. \left. - (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t))(\tau' - \tau_n^{(m)}) \right| \leq \frac{\epsilon f(L)}{2} \right) \geq 1 - o(L^{-2}), \end{aligned} \quad (129)$$

and

$$\begin{aligned} \mathbb{P} \left( \left| \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} (x_{\bar{k}^{(\ell)}}^{L(e)}(\tau) - x_{\bar{k}^{(\ell)}}^{L(e)}(\tau')) \right. \right. \\ \left. \left. - (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t))(\tau - \tau') \right| \leq \frac{\epsilon f(L)}{2} \right) \geq 1 - o(L^{-2}). \end{aligned} \quad (130)$$

**Proof of (129):** We will now prove (129) by considering two cases depending on length of  $\tau' - \tau_n^{(m)}$ , i.e., we have either

$$\tau' - \tau_n^{(m)} \leq \frac{\epsilon f(L)}{4(1+K)^i R} \quad (131)$$

or

$$\tau' - \tau_n^{(m)} > \frac{\epsilon f(L)}{4(1+K)^i R}, \quad (132)$$

where  $R := \sum_{j \in \mathcal{J}} \lambda_j + K\mu_{max}$

**Case (131):** We notice  $x_{\bar{k}^{(\ell)}}^{L(e)}(\tau)$  will change by at most  $(1+K)^{\ell-1}/L$  at each arrival or departure according to Lemma D.1 and thus  $\sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} x_{\bar{k}^{(\ell)}}^{L(e)}(\tau)$  will change by at most  $\frac{(1+K)^i - 1}{L}$ .

For the number of arrivals and departures in  $[\tau_n^{(m)}, \tau']$  which are Poisson processes with means at most  $\left( \sum_{j \in \mathcal{J}} \lambda_j + K\mu_{max} \right) L$  on an interval of length at most  $\frac{\epsilon f(L)}{4(1+K)^i R}$  we have, according to

Lemma F.1, that with probability  $1 - o(L^{-2})$  they are at most

$$\left( \sum_{j \in \mathcal{J}} \lambda_j + K\mu_{\max} \right) L \frac{\epsilon f(L)}{4(1+K)^i R} + o(Lf(L)) = \frac{\epsilon Lf(L)}{4(1+K)^i} + o(Lf(L)),$$

therefore,

$$\mathbb{P} \left( \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} (x_{\bar{k}(\ell)}^{L(e)}(\tau') - x_{\bar{k}(\ell)}^{L(e)}(\tau_n^{(m)})) \leq \frac{\epsilon f(L)}{4} \right) = 1 - o(L^{-2}). \quad (133)$$

Considering (131) holds, in this case we clearly have

$$\mathbb{P} \left( (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t)) (\tau' - \tau_n^{(m)}) \leq \frac{\epsilon f(L)}{4} \right) = 1 \quad (134)$$

It is now easy to verify that equations (133) and (134) imply (129).

**Case (132):** In this case we notice,

$$\begin{aligned} o(Lf(L)) &= q_{\bar{k}^{(i)}, j_i}^L(\tau_n^{(m)}) = q_{\bar{k}^{(i)}, j_i}^L(\tau') \\ &\quad - \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} L(x_{\bar{k}(\ell)}^{L(e)}(\tau') - x_{\bar{k}(\ell)}^{L(e)}(\tau_n^{(m)})) + Ly_{j_i}^L(\tau') - Ly_{j_i}^L(\tau_n^{(m)}) \\ &\leq K - \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} L(x_{\bar{k}(\ell)}^{L(e)}(\tau') - x_{\bar{k}(\ell)}^{L(e)}(\tau_n^{(m)})) \\ &\quad + L(\lambda_{j_i} - \mu_{j_i} y_{j_i}(t))(\tau' - \tau_n^{(m)}) + o(Lf(L)), \end{aligned}$$

which holds with probability  $1 - o(L^{-2})$  by applying Lemma F.1 in the last step. It therefore follows that, with the same probability,

$$\begin{aligned} \frac{x_{\bar{k}^{(i)}}^{L(e)}(\tau') - x_{\bar{k}^{(i)}}^{L(e)}(\tau_n^{(m)})}{\tau' - \tau_n^{(m)}} &\leq \\ \frac{\lambda_{j_i} - \mu_{j_i} y_{j_i}(t)}{\bar{k}_{j_i}^{(i)}} - \sum_{\ell=1}^{i-1} \frac{\bar{k}_{j_i}^{(\ell)} (x_{\bar{k}(\ell)}^{L(e)}(\tau') - x_{\bar{k}(\ell)}^{L(e)}(\tau_n^{(m)}))}{\bar{k}_{j_i}^{(i)} (\tau' - \tau_n^{(m)})} + \frac{o(f(L))}{\tau' - \tau_n^{(m)}} &\stackrel{(a)}{=} \\ \frac{\lambda_{j_i} - \mu_{j_i} y_{j_i}(t) - \sum_{\ell=1}^{i-1} \bar{k}_{j_i}^{(\ell)} h^{\bar{\mathcal{J}}[m](\ell)}(t)}{\bar{k}_{j_i}^{(i)}} + o(1), & \end{aligned} \quad (135)$$

where in (a) we applied (95) of Lemma G.2 for indexes  $1, \dots, i-1$ , and used (132).

Next, let

$$j' := \arg \max_{j \in \mathcal{J}: \bar{k}_j^{(i)} > 0} q_{\bar{k}^{(i)}, j}^L(\tau').$$

Then again we have that, with probability  $1 - o(L^{-2})$ ,

$$\begin{aligned} 0 &\leq q_{\bar{k}^{(i)}, j'}^L(\tau') = q_{\bar{k}^{(i)}, j'}^L(\tau_n^{(m)}) + \\ &\quad \sum_{\ell=1}^i \bar{k}_{j'}^{(\ell)} L(x_{\bar{k}(\ell)}^{L(e)}(\tau') - x_{\bar{k}(\ell)}^{L(e)}(\tau_n^{(m)})) - L(y_{j'}^L(\tau') - y_{j'}^L(\tau_n^{(m)})) \\ &\leq K + \sum_{\ell=1}^i \bar{k}_{j'}^{(\ell)} L(x_{\bar{k}(\ell)}^{L(e)}(\tau') - x_{\bar{k}(\ell)}^{L(e)}(\tau_n^{(m)})) \\ &\quad - (\lambda_{j'} - \mu_{j'} y_{j'}(t))(\tau' - \tau_n^{(m)}) + o(Lf(L)), \end{aligned}$$

from which it follows that

$$\begin{aligned} \frac{x_{\bar{k}^{(i)}}^{L(e)}(\tau') - x_{\bar{k}^{(i)}}^{L(e)}(\tau_n^{(m)})}{\tau' - \tau_n^{(m)}} &\geq \\ \frac{\lambda_{j'} - \mu_{j'} y_{j'}(t)}{\bar{k}_{j'}^{(i)}} - \sum_{\ell=1}^{i-1} \frac{\bar{k}_{j'}^{(\ell)} (x_{\bar{k}(\ell)}^{L(e)}(\tau') - x_{\bar{k}(\ell)}^{L(e)}(\tau_n^{(m)}))}{\bar{k}_{j'}^{(i)} (\tau' - \tau_n^{(m)})} + \frac{o(f(L))}{\tau' - \tau_n^{(m)}} &\stackrel{(a)}{=} \\ \frac{\lambda_{j'} - \mu_{j'} y_{j'}(t) - \sum_{\ell=1}^{i-1} \bar{k}_{j'}^{(\ell)} h^{\bar{\mathcal{J}}[m](\ell)}(t)}{\bar{k}_{j'}^{(i)}} + o(1). & \end{aligned} \quad (136)$$

where in (a) we applied (95) of Lemma G.2 for indexes  $1, \dots, i-1$ , and used (132).

Considering (37) holds, that (38) does not hold for  $j = j'$  and  $G_m = i$ , we get

$$\begin{aligned} \frac{\lambda_{j'} - \mu_{j'} y_{j'}(t) - \sum_{\ell=1}^{i-1} \bar{k}_{j'}^{(\ell)} h^{\bar{\mathcal{J}}',(\ell)}(t)}{\bar{k}_{j'}^{(i)}} &\geq \\ h^{\bar{\mathcal{J}}[m],(i)}(t) = \frac{\lambda_{j_i} - \mu_{j_i} y_{j_i}(t) - \sum_{\ell=1}^{i-1} \bar{k}_{j_i}^{(\ell)} h^{\bar{\mathcal{J}}[m],(\ell)}(t)}{\bar{k}_{j_i}^{(i)}}. & \end{aligned} \quad (137)$$

From (135), (136) and (137) we get

$$\begin{aligned} \frac{x_{\bar{k}^{(i)}}^{L(e)}(\tau') - x_{\bar{k}^{(i)}}^{L(e)}(\tau_n^{(m)})}{\tau' - \tau_n^{(m)}} &= \\ \frac{\lambda_{j_i} - \mu_{j_i} y_{j_i}(t) - \sum_{\ell=1}^{i-1} \bar{k}_{j_i}^{(\ell)} h^{\bar{\mathcal{J}}[m],(\ell)}(t)}{\bar{k}_{j_i}^{(i)}} + o(1), & \end{aligned}$$

which holds with probability  $1 - o(L^{-2})$  and therefore it implies (129).

**Proof of (130):** We will now prove (130) by considering two cases depending on length of  $\tau - \tau'$ .

We consider

$$\tau - \tau' \leq \frac{\epsilon f(L)}{4(1+K)^i R} \quad (138)$$

or

$$\tau - \tau' > \frac{\epsilon f(L)}{4(1+K)^i R}, \quad (139)$$

where  $R := \sum_{j \in \mathcal{J}} \lambda_j + K\mu_{\max}$ , and

$$\epsilon < \frac{\mu_{j_i} \alpha_i}{4(1+K)^i R}. \quad (140)$$

**Case (138):** Following the same arguments as in the Case of (131) we can infer the equivalent of (133) and (134) for interval  $(\tau', \tau)$ , i.e.

$$\lim_{L \rightarrow \infty} \mathbb{P} \left( \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} (x_{\bar{k}(\ell)}^{L(e)}(\tau) - x_{\bar{k}(\ell)}^{L(e)}(\tau')) \leq \frac{\epsilon f(L)}{4} \right) = 1 - o(L^{-2}) \quad (141)$$

and

$$\mathbb{P} \left( (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t))(\tau - \tau') \leq \frac{\epsilon f(L)}{4} \right) = 1 - o(L^{-2}). \quad (142)$$

which imply (130).



**Case (139):** First we will prove that

$$q_{\bar{k}^{(i)}, j_i}^L(\tau') \geq o(Lf(L)) - \epsilon^2 Lf(L), \quad (143)$$

or equivalently

$$1/Lq_{\bar{k}^{(i)}, j_i}^L(\tau') - 1/Lq_{\bar{k}^{(i)}, j_i}^L(\tau_n^{(m)}) \geq o(f(L)) - \epsilon^2 f(L). \quad (144)$$

Both will hold with probability  $1 - o(L^{-2})$ . The analysis is same as with the proof of (129) with the following changes.

- Instead of considering cases  $\tau' - \tau_n^{(m)} \leq \frac{\epsilon f(L)}{4(1+K)^i R}$  and  $\tau' - \tau_n^{(m)} > \frac{\epsilon f(L)}{4(1+K)^i R}$ , we should consider  $\tau' - \tau_n^{(m)} \leq \frac{\epsilon^2 f(L)}{2(1+K)^i R}$  and  $\tau' - \tau_n^{(m)} > \frac{\epsilon^2 f(L)}{2(1+K)^i R}$ .
- If interval is short, we can bound the absolute change of variable  $1/Lq_{\bar{k}^{(i)}, j_i}^L(\tau)$  which changes by at most  $(1+K)^i/L$ .
- If interval is long, we can still prove that the equivalent of (135) is satisfied as equality and considering  $y_{j_i}^L(\tau') - y_{j_i}^L(\tau_n^{(m)}) = \lambda_{j_i} - \mu_{j_i} y_{j_i}(t) + o(f(L))$ , we can get through (135) that  $1/Lq_{\bar{k}^{(i)}, j_i}^L(\tau') = o(f(L))$  with probability  $1 - o(L^{-2})$ .

In this case, because of (143) we get

$$\begin{aligned} o(Lf(L)) - \epsilon^2 Lf(L) &\leq q_{\bar{k}^{(i)}, j_i}^L(\tau') = q_{\bar{k}^{(i)}, j_i}^L(\tau) \\ &- \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} L(x_{\bar{k}^{(\ell)}}^{L(e)}(\tau) - x_{\bar{k}^{(\ell)}}^{L(e)}(\tau')) + Ly_{j_i}^L(\tau) - Ly_{j_i}^L(\tau') \leq \\ K - \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} L(x_{\bar{k}^{(\ell)}}^{L(e)}(\tau) - x_{\bar{k}^{(\ell)}}^{L(e)}(\tau')) + L(\lambda_{j_i} - \mu_{j_i} y_{j_i}(t))(\tau - \tau') \\ &+ o(Lf(L)), \end{aligned}$$

from which it follows

$$\begin{aligned} \frac{x_{\bar{k}^{(i)}}^{L(e)}(\tau) - x_{\bar{k}^{(i)}}^{L(e)}(\tau')}{\tau - \tau'} &\leq \frac{\lambda_{j_i} - \mu_{j_i} y_{j_i}(t)}{\bar{k}_{j_i}^{(i)}} \\ &- \sum_{\ell=1}^{i-1} \frac{\bar{k}_{j_i}^{(\ell)} (x_{\bar{k}^{(\ell)}}^{L(e)}(\tau) - x_{\bar{k}^{(\ell)}}^{L(e)}(\tau')) - \epsilon^2 f(L)}{\bar{k}_{j_i}^{(i)}(\tau - \tau')} + \frac{o(f(L))}{\tau - \tau'} \stackrel{(a)}{<} \\ \mu_{j_i} \sum_{\ell=1}^i \frac{\bar{k}_{j_i}^{(\ell)}}{\bar{k}_{j_i}^{(i)}} (x_{\bar{k}^{(\ell)}}^{(g)} - x_{\bar{k}^{(\ell)}}^{(e)}(t)) - \sum_{\ell=1}^{i-1} \frac{\bar{k}_{j_i}^{(\ell)} h^{\bar{J}[m], (\ell)}(t)}{\bar{k}_{j_i}^{(i)}} \\ &+ \frac{\epsilon 4(1+K)^i R}{\bar{k}_{j_i}^{(i)}} + o(1) \stackrel{(b)}{<} \mu_{\max}(\alpha_i + \sum_{\ell=1}^{i-1} K\alpha_{\ell'}) \\ &+ \sum_{\ell=1}^{i-1} K\mu_{\max} \left( \alpha_{\ell} + \sum_{\ell'=1}^{\ell-1} 2(1+K)^{\ell-\ell'} \alpha_{\ell'} \right) + \\ 4\epsilon(1+K)^i R + o(1) &\stackrel{(c)}{<} 2\mu_{\max}\alpha_i + 2\mu_{\max} \sum_{\ell=1}^{i-1} (1+K)^{i-\ell} \alpha_{\ell} + o(1). \end{aligned} \quad (145)$$

In (a) we applied the properties of  $j_i$ , from (28), (33), and (95) of Lemma G.2, for indexes  $1, \dots, i-1$ , and then replaced  $\tau - \tau'$  with its bound from (139). In (b), we used property (34), and the fact that  $\frac{\bar{k}_{j_i}^{(\ell)}}{\bar{k}_{j_i}^{(i)}} \leq K$  for  $\ell = 1, \dots, i-1$ . In (c), we used (140) and simplified.

Also from Lemma F.3 and given that all the servers that empty during  $[\tau', \tau]$  will be assigned to configuration  $\bar{k}^{(\ell)}$  for  $\ell = 1, \dots, i$ , we will have that with probability  $1 - o(L^{-2})$ ,

$$\begin{aligned} \frac{x_{\bar{k}^{(i)}}^{L(e)}(\tau) - x_{\bar{k}^{(i)}}^{L(e)}(\tau')}{\tau - \tau'} &\geq \frac{\mu_{\min}}{JKC^2} \left( 1 - \sum_{\ell=1}^i x_{\bar{k}^{(\ell)}}^{(e)}(t) \right) \\ &- \sum_{\ell=1}^{i-1} \frac{(x_{\bar{k}^{(\ell)}}^{L(e)}(\tau) - x_{\bar{k}^{(\ell)}}^{L(e)}(\tau'))^+}{\tau - \tau'} + \frac{o(f(L))}{\tau - \tau'} \stackrel{(a)}{=} \\ \frac{\mu_{\min}}{JKC^2} \left( 1 - \sum_{\ell=1}^i x_{\bar{k}^{(\ell)}}^{(e)}(t) \right) &- \sum_{\ell=1}^{i-1} h^{\bar{J}[m], (\ell)}(t)^+ \\ &+ o(1) \stackrel{(b)}{>} \frac{\mu_{\min}}{JKC^2} \frac{x_{\bar{k}^{(c\rho)}}^{(c\rho)}}{2} - 2\mu_{\max} \sum_{\ell=1}^{i-1} \frac{(1+K)^{i-\ell}}{K} \alpha_{\ell} + o(1). \end{aligned} \quad (146)$$

In (a) we applied (95) of Lemma G.2 for indexes  $1, \dots, i-1$  and used (139). In (b) we applied Lemma G.1 and equation (96) for indexes  $1, \dots, i-1$  and simplified.

So far we proved that if (139) holds, (145) and (146) also hold.

However, considering (90) we have

$$\begin{aligned} 2\mu_{\max}\alpha_i + 2\mu_{\max} \sum_{\ell=1}^{i-1} (1+K)^{i-\ell} \alpha_{\ell} &\stackrel{(a)}{<} \\ \frac{\mu_{\min}}{JKC^2} \frac{x_{\bar{k}^{(c\rho)}}^{(c\rho)}}{2} - 2\mu_{\max} \sum_{\ell=1}^{i-1} \frac{(1+K)^{i-\ell}}{K} \alpha_{\ell} &\end{aligned} \quad (147)$$

and because of that, the probability that (145) and (146) are both true is  $o(L^{-2})$ . This means that (138) holds with probability at least  $1 - o(L^{-2})$ , and thus the analysis of (138) is sufficient for (104) to hold.

### Proof of Claim 2:

We will prove the result inductively on  $m$ . For  $m = 0$ ,  $\ell_m = 0$  so there is nothing to prove, so we will start with the base case  $m = 1$  and then move on to the inductive step.

#### Base case $m = 1$ :

In this case  $\ell_m = 1$  and  $0 \leq q_{\bar{k}^{(1)}, j_i[1]}^L(\tau_n^{(1)}) < K$  from which it trivially follows that  $q_{\bar{k}^{(1)}, j_i[1]}^L(\tau_n^{(1)}) = o(Lf(L))$ .

#### Inductive case $m > 1$ , Base case $i = \ell_m$ :

$$0 \leq q_{\bar{k}^{(i)}, j_i[m]}^L(\tau_n^{(m)}) < K$$

from which it trivially follows that

$$q_{\bar{k}^{(i)}, j_i[m]}^L(\tau_n^{(m)}) = o(Lf(L)).$$

**Inductive case  $m > 1$ , Base case  $i < \ell_m$ :** In this case, we know by the induction hypothesis that  $q_{\bar{k}^{(i)}, j_i[m-1]}^L(\tau_n^{(m-1)}) = o(Lf(L))$ . Further, notice that  $j_i[m-1] = j_i[m]$  for  $i = 1, \dots, \ell_m - 1$  so for simplicity we will refer to both as  $j_i$ . Then we have that with

probability  $1 - o(L^{-2})$ ,

$$\begin{aligned} q_{\bar{k}^{(i)}, j_i}^L(\tau_n^{(m)}) &\geq q_{\bar{k}^{(i)}, j_i}^L(\tau_n^{(m-1)}) + \\ &L \left( \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} \left( x_{\bar{k}^{(\ell)}}^{(e)}(\tau_n^{(m)}) - x_{\bar{k}^{(\ell)}}^{(e)}(\tau_n^{(m-1)}) \right) - (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t)) \right) \times \\ &(\tau_n^{(m)} - \tau_n^{(m-1)}) + o(Lf(L)) \stackrel{(a)}{=} o(Lf(L)). \end{aligned} \quad (148)$$

In (a) we have used that (42) holds for index  $m-1$  in place of  $m$ .

**Proof of Property P.2.:**

*Definition G.5.* For an index  $i$  for which  $1 \leq i \leq \ell_m$ , the time  $\tau_n^{(m,i)}$  is defined as the latest time in  $[\tau_n^{(m)}, \tau_n^{(m+1)})$ , such that

$$\max_{j \in \mathcal{J}; \bar{k}_j^{(i)} > 0} q_{\bar{k}^{(i)}, j}^L(\tau_n^{(m,i)}) \geq 0. \quad (149)$$

Also let  $j(i) := \arg \max_{j \in \mathcal{J}; \bar{k}_j^{(i)} > 0} q_{\bar{k}^{(i)}, j}^L(\tau_n^{(m,i)})$ .

For (P.2.) to be true, it suffices that for any  $\epsilon > 0$

$$\begin{aligned} &\mathbb{P} \left( x_{\bar{k}^{(\ell_m)}}^{(e)}(\tau_n^{(m+1)}) - x_{\bar{k}^{(\ell_m)}}^{(e)}(\tau_n^{(m)}) \right. \\ &\quad \left. - \mu_{\min} \frac{\alpha \ell_m}{2} (\tau_n^{(m+1)} - \tau_n^{(m)}) > -\epsilon f(L) \right) = 1 - o(L^{-2}). \end{aligned} \quad (150)$$

To prove (150), it is sufficient to prove

$$\begin{aligned} &\mathbb{P} \left( x_{\bar{k}^{(\ell_m)}}^{(e)}(\tau_n^{(m, \ell_m)}) - x_{\bar{k}^{(\ell_m)}}^{(e)}(\tau_n^{(m)}) \right. \\ &\quad \left. - \mu_{\min} \frac{\alpha \ell_m}{2} (\tau_n^{(m, \ell_m)} - \tau_n^{(m)}) > -\frac{\epsilon f(L)}{2} \right) = 1 - o(L^{-2}), \end{aligned} \quad (151)$$

and

$$\begin{aligned} &\mathbb{P} \left( x_{\bar{k}^{(\ell_m)}}^{(e)}(\tau_n^{(m+1)}) - x_{\bar{k}^{(\ell_m)}}^{(e)}(\tau_n^{(m, \ell_m)}) \right. \\ &\quad \left. - \mu_{\min} \frac{\alpha \ell_m}{2} (\tau_n^{(m+1)} - \tau_n^{(m, \ell_m)}) > -\frac{\epsilon f(L)}{2} \right) = 1 - o(L^{-2}). \end{aligned} \quad (152)$$

**Proof of (151):** We will now prove (151) by considering two cases depending on length of  $\tau_n^{(m, \ell_m)} - \tau_n^{(m)}$ .

We consider

$$\tau_n^{(m, \ell_m)} - \tau_n^{(m)} \leq \frac{\epsilon f(L)}{4(1+K)\ell_m^{-1}R} \quad (153)$$

or

$$\tau_n^{(m, \ell_m)} - \tau_n^{(m)} > \frac{\epsilon f(L)}{4(1+K)\ell_m^{-1}R} \quad (154)$$

where  $R := \sum_{j \in \mathcal{J}} \lambda_j + K\mu_{\max}$

**Case (153):** We notice  $x_{\bar{k}^{(\ell_m)}}^{(e)}(\tau)$  will change by at most  $(1+K)\ell_m^{-1}/L$  at each arrival or departure according to Lemma D.1.

Using Lemma F.1, with probability  $1 - o(L^{-2})$ , The number of arrivals and departures in  $[\tau_n^{(m)}, \tau_n^{(m, \ell_m)}]$  can be bounded by

$$\begin{aligned} &\left( \sum_{j \in \mathcal{J}} \lambda_j + K\mu_{\max} \right) L \frac{\epsilon f(L)}{4(1+K)\ell_m^{-1}R} + o(Lf(L)) = \\ &\frac{\epsilon Lf(L)}{4(1+K)\ell_m^{-1}} + o(Lf(L)), \end{aligned} \quad (155)$$

therefore

$$\mathbb{P} \left( x_{\bar{k}^{(\ell_m)}}^{(e)}(\tau_n^{(m, \ell_m)}) - x_{\bar{k}^{(\ell_m)}}^{(e)}(\tau_n^{(m)}) \leq \frac{\epsilon f(L)}{4} \right) = 1 - o(L^{-2}). \quad (156)$$

Considering (153) holds, we trivially have

$$\mathbb{P} \left( \mu_{\min} \frac{\alpha \ell_m}{2} (\tau_n^{(m, \ell_m)} - \tau_n^{(m)}) \leq \frac{\epsilon f(L)}{4} \right) = 1 \quad (157)$$

It is now easy to verify that (156) and (157) imply (151).

**Case (154):**

Let  $j := j(\ell_m)$ . In this case, we have

$$\begin{aligned} 0 &\leq q_{\bar{k}^{(\ell_m)}, j}^L(\tau_n^{(m, \ell_m)}) = q_{\bar{k}^{(\ell_m)}, j}^L(\tau_n^{(m)}) \\ &+ \sum_{\ell=1}^{\ell_m} \bar{k}_j^{(\ell)} L \left( x_{\bar{k}^{(\ell)}}^{L(e)}(\tau_n^{(m, \ell_m)}) - x_{\bar{k}^{(\ell)}}^{L(e)}(\tau_n^{(m)}) \right) \\ &- L \left( y_j^L(\tau_n^{(m, \ell_m)}) - y_j^L(\tau_n^{(m)}) \right) \\ &\leq K + \sum_{\ell=1}^{\ell_m} \bar{k}_j^{(\ell)} L \left( x_{\bar{k}^{(\ell)}}^{L(e)}(\tau_n^{(m, \ell_m)}) - x_{\bar{k}^{(\ell)}}^{L(e)}(\tau_n^{(m)}) \right) \\ &- L(\lambda_j - \mu_j y_j(t)) \left( \tau_n^{(m, \ell_m)} - \tau_n^{(m)} \right) + o(Lf(L)), \end{aligned} \quad (158)$$

which holds with probability  $1 - o(L^{-2})$  by application of Lemma F.1 in the last step. If  $\ell_m > C_\alpha(t)$  then, since (34) does not hold for  $i = \ell_m$ , we have

$$\frac{\lambda_j - \mu_j y_j(t)}{\bar{k}_j^{(\ell_m)}} > \frac{\mu_j \alpha \ell_m + \mu_j \sum_{\ell=1}^{\ell_m-1} \bar{k}_j^{(\ell)} (x_{\bar{k}^{(\ell)}}^{(g)} - x_{\bar{k}^{(\ell)}}^{(e)}(t))}{\bar{k}_j^{(\ell_m)}}, \quad (159)$$

otherwise, by property (92) for  $j_a = j$  and  $j_b = \ell_m$  we have

$$\frac{\lambda_j - \mu_j y_j(t)}{\bar{k}_j^{(\ell_m)}} > \frac{\mu_j \delta + \mu_j \sum_{\ell=1}^{\ell_m} \bar{k}_j^{(\ell)} (x_{\bar{k}^{(\ell)}}^{(g)} - x_{\bar{k}^{(\ell)}}^{(e)}(t))}{\bar{k}_j^{(\ell_m)}}. \quad (160)$$

Based on the above inequality, we can write

$$\begin{aligned} &\frac{x_{\bar{k}^{(\ell_m)}}^{L(e)}(\tau_n^{(m, \ell_m)}) - x_{\bar{k}^{(\ell_m)}}^{L(e)}(\tau_n^{(m)})}{\tau_n^{(m, \ell_m)} - \tau_n^{(m)}} \geq \\ &\frac{\lambda_j - \mu_j y_j(t)}{\bar{k}_j^{(\ell_m)}} - \frac{\sum_{\ell=1}^{\ell_m-1} \bar{k}_j^{(\ell)} \left( x_{\bar{k}^{(\ell)}}^{(e)}(\tau_n^{(m, \ell_m)}) - x_{\bar{k}^{(\ell)}}^{(e)}(\tau_n^{(m)}) \right)}{\sum_{\ell=1}^{\ell_m-1} \bar{k}_j^{(\ell)} (\tau_n^{(m, \ell_m)} - \tau_n^{(m)})} \\ &+ \frac{o(f(L))}{\tau_n^{(m, \ell_m)} - \tau_n^{(m)}} \stackrel{(a)}{\geq} \mu_j \min \left( \frac{\delta}{\bar{k}_j^{(\ell_m)}} - (x_{\bar{k}^{(\ell_m)}}^{(g)} - x_{\bar{k}^{(\ell_m)}}^{(e)}(t)), \alpha \ell_m \right) \\ &+ \frac{\mu_j \sum_{\ell=1}^{\ell_m-1} \bar{k}_j^{(\ell)} (x_{\bar{k}^{(\ell)}}^{(g)} - x_{\bar{k}^{(\ell)}}^{(e)}(t)) - \sum_{\ell=1}^{\ell_m-1} \bar{k}_j^{(\ell)} h^{\mathcal{J}[m], (\ell)}(t)}{\bar{k}_j^{(\ell_m)}} \\ &+ o(1) \stackrel{(b)}{\geq} \left( \mu_{\min} \min \left( \delta - (\epsilon_\rho + K \sum_{\ell=1}^{\ell_m-1} \alpha_\ell), \alpha \ell_m \right) \right. \\ &\quad \left. - \mu_{\max} \sum_{\ell=1}^{\ell_m-1} K(\epsilon_\rho + K \sum_{\ell'=1}^{\ell-1} \alpha_{\ell'}) \right. \\ &\quad \left. - \sum_{\ell=1}^{\ell_m-1} K\mu_{\max} \left( \alpha_\ell + \sum_{\ell'=1}^{\ell-1} 2(1+K)\ell^{-\ell'} \alpha_{\ell'} \right) \right) + o(1) \stackrel{(c)}{>} \\ &\frac{\mu_{\min} \alpha \ell_m}{2} + o(1). \end{aligned} \quad (161)$$

In (a) we used: 1) (159) and (160), 2) equation (96) for indexes  $1, \dots, \ell_m - 1$ , and 3) replaced  $\tau_n^{(m, \ell_m)} - \tau_n^{(m)}$  with its bound from (154). In (b) we used Lemma G.4 for indexes  $1, \dots, \ell_m$ . Finally, in (c) we used (90) and (93).

We have thus shown by (161) that

$$\frac{x_{\bar{k}^{(\ell_m)}}^{L(e)}(\tau_n^{(m, \ell_m)}) - x_{\bar{k}^{(\ell_m)}}^{L(e)}(\tau_n^{(m)})}{\tau_n^{(m, \ell_m)} - \tau_n^{(m)}} > \frac{\mu \min \alpha \ell_m}{2} + o(1)$$

with probability  $1 - o(L^{-2})$  which is equivalent to (151).

**Proof of (152):** We will now prove (152) by considering two cases depending on length of  $\tau_n^{(m+1)} - \tau_n^{(m, \ell_m)}$  and reach a contradiction for each of them.

We consider

$$\tau_n^{(m+1)} - \tau_n^{(m, \ell_m)} \leq \frac{\epsilon f(L)}{4(1+K)\ell_m^{-1}(\sum_{j \in \mathcal{J}} \lambda_j + K\mu_{max})} \quad (162)$$

or

$$\tau_n^{(m+1)} - \tau_n^{(m, \ell_m)} > \frac{\epsilon f(L)}{4(1+K)\ell_m^{-1}(\sum_{j \in \mathcal{J}} \lambda_j + K\mu_{max})} \quad (163)$$

**Case (162):** Following the same arguments as in the Case of (153) we can infer the equivalent of (156) and (157) for interval  $(\tau_n^{(m, \ell_m)}, \tau_n^{(m+1)})$ , i.e.

$$\mathbb{P}\left(x_{\bar{k}^{(\ell_m)}}^{(e)}(\tau_n^{(m+1)}) - x_{\bar{k}^{(\ell_m)}}^{(e)}(\tau_n^{(m, \ell_m)}) \leq \frac{\epsilon f(L)}{4}\right) = 1 - o(L^{-2}), \quad (164)$$

and

$$\mathbb{P}\left(\frac{\mu \min \alpha \ell_m}{2}(\tau_n^{(m+1)} - \tau_n^{(m, \ell_m)}) \leq \frac{\epsilon f(L)}{4}\right) = 1 - o(L^{-2}). \quad (165)$$

which imply (152).

**Case (163):** In this case we notice, using Lemma F.3 that with probability  $1 - o(L^{-2})$

$$\begin{aligned} & (x_{\bar{k}^{(\ell_m)}}^{L(e)}(\tau_n^{(m+1)}) - x_{\bar{k}^{(\ell_m)}}^{L(e)}(\tau_n^{(m, \ell_m)})) \\ & + \sum_{\ell=1}^{\ell_m-1} \left(x_{\bar{k}^{(\ell)}}^{L(e)}(\tau_n^{(m+1)}) - x_{\bar{k}^{(\ell)}}^{L(e)}(\tau_n^{(m, \ell_m)})\right)^+ > \end{aligned} \quad (166)$$

$$\frac{\mu \min}{JKC^2} \left(1 - \sum_{\ell=1}^{\ell_m} x_{\bar{k}^{(\ell)}}^{(e)}(t)\right) (\tau_n^{(m+1)} - \tau_n^{(m, \ell_m)}) + o(f(L)),$$

or equivalently

$$\begin{aligned} & \frac{x_{\bar{k}^{(\ell_m)}}^{L(e)}(\tau_n^{(m+1)}) - x_{\bar{k}^{(\ell_m)}}^{L(e)}(\tau_n^{(m, \ell_m)})}{\tau_n^{(m+1)} - \tau_n^{(m, \ell_m)}} \stackrel{(a)}{>} \\ & \frac{\mu \min}{JKC^2} \left(1 - \sum_{\ell=1}^{\ell_m} x_{\bar{k}^{(\ell)}}^{(e)}(t)\right) - \sum_{\ell=1}^{\ell_m-1} h^{(\ell)}(t)^+ + \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m, \ell_m)}} \stackrel{(b)}{>} \\ & \frac{\mu \min}{JKC^2} \frac{x^{(g)}}{\bar{k}^{(C_\rho^{(g)})}} - \sum_{\ell=1}^{\ell_m-1} \mu_{max} \left(\alpha_\ell + 2 \sum_{\ell'=1}^{\ell-1} (1+K)^{\ell-\ell'} \alpha_{\ell'}\right) + o(1) \stackrel{(c)}{>} \\ & \frac{\mu \min \alpha \ell_m}{2} + o(1). \end{aligned} \quad (167)$$

Inequality (a) comes from applying to (166), the equation (95) of Lemma G.2 for indexes  $1, \dots, \ell_m - 1$ . In (b) we used Lemma G.1

and equation (96) for indexes  $1, \dots, \ell_m - 1$ . Finally, in (c) we used (90).

We have thus shown through (167) that

$$\frac{x_{\bar{k}^{(\ell_m)}}^{L(e)}(\tau_n^{(m+1)}) - x_{\bar{k}^{(\ell_m)}}^{L(e)}(\tau_n^{(m, \ell_m)})}{\tau_n^{(m+1)} - \tau_n^{(m, \ell_m)}} > \frac{\mu \min \alpha \ell_m}{2} + o(1)$$

with probability  $1 - o(L^{-2})$ , which is equivalent to (152).

**Proof of Property P.3.:** Consider  $j = j(C_\rho^{(g)})$  given from Definition G.5.

For (P.3.) to be true, it suffices to prove that for any  $\epsilon > 0$ ,

$$\begin{aligned} P_1 := & \mathbb{P}\left(\sum_{\ell=1}^{C_\rho^{(g)}} \bar{k}_j^{(\ell)} \left(x_{\bar{k}^{(\ell)}}^{(e)}(\tau_n^{(m, C_\rho^{(g)})}) - x_{\bar{k}^{(\ell)}}^{(e)}(\tau_n^{(m)})\right) \right. \\ & \left. - (\lambda_j - \mu_j y_j(t))(\tau_n^{(m, C_\rho^{(g)})} - \tau_n^{(m)}) > -\frac{\epsilon f(L)}{2}\right) = 1 - o(L^{-2}), \end{aligned} \quad (168)$$

and

$$\begin{aligned} P_2 := & \mathbb{P}\left(x_{\bar{k}^{(C_\rho^{(g)})}}^{(e)}(\tau_n^{(m+1)}) - x_{\bar{k}^{(C_\rho^{(g)})}}^{(e)}(\tau_n^{(m, C_\rho^{(g)})}) \right. \\ & + \sum_{\ell=1}^{C_\rho^{(g)}-1} \left(x_{\bar{k}^{(\ell)}}^{(e)}(\tau_n^{(m+1)}) - x_{\bar{k}^{(\ell)}}^{(e)}(\tau_n^{(m, C_\rho^{(g)})})\right)^+ \\ & \left. - \mu \min \frac{1 - \sum_{i=1}^{C_\rho^{(g)}} x_{\bar{k}^{(i)}}^{(e)}(t)}{JKC^2} (\tau_n^{(m+1)} - \tau_n^{(m, C_\rho^{(g)})}) > -\frac{\epsilon f(L)}{2}\right) \\ & = 1 - o(L^{-2}). \end{aligned} \quad (169)$$

To show why this is sufficient we first introduce the following notations

$$\begin{aligned} f_1[\tau_a, \tau_b] := & \mu \min \frac{1 - \sum_{i=1}^{C_\rho^{(g)}} x_{\bar{k}^{(i)}}^{(e)}(t)}{JKC^2} - \sum_{i=1}^{C_\rho^{(g)}-1} \nabla x_{\bar{k}^{(i)}}^{(e)}[\tau_a, \tau_b]^+ \\ f_2[\tau_a, \tau_b] := & \min_{j \in \mathcal{J}} \left( \frac{\lambda_j - \mu_j y_j(t) - \sum_{i=1}^{C_\rho^{(g)}-1} \bar{k}_j^{(i)} \nabla x_{\bar{k}^{(i)}}^{(e)}[\tau_a, \tau_b]}{\bar{k}_j^{(C_\rho^{(g)})}} \right) \end{aligned} \quad (170)$$

Then if (168) and (169) indeed hold, we can get that

$$\begin{aligned} & \mathbb{P} \left( x_{\bar{k}^{(C_\rho^{(g)})}}^{(e)}(\tau_n^{(m+1)}) - x_{\bar{k}^{(C_\rho^{(g)})}}^{(e)}(\tau_n^{(m)}) > \right. \\ & \left. \min(f_1[\tau_n^{(m)}, \tau_n^{(m+1)}], f_2[\tau_n^{(m)}, \tau_n^{(m+1)}]) - \epsilon f(L) \right) \geq \\ & \mathbb{P} \left( x_{\bar{k}^{(C_\rho^{(g)})}}^{(e)}(\tau_n^{(m, C_\rho^{(g)})}) - x_{\bar{k}^{(C_\rho^{(g)})}}^{(e)}(\tau_n^{(m)}) > \right. \\ & \left. \min(f_1[\tau_n^{(m)}, \tau_n^{(m, C_\rho^{(g)})}], f_2[\tau_n^{(m)}, \tau_n^{(m, C_\rho^{(g)})}]) - \frac{\epsilon f(L)}{2} \right) \end{aligned} \quad (171)$$

$$\begin{aligned} & \mathbb{P} \left( x_{\bar{k}^{(C_\rho^{(g)})}}^{(e)}(\tau_n^{(m+1)}) - x_{\bar{k}^{(C_\rho^{(g)})}}^{(e)}(\tau_n^{(m, C_\rho^{(g)})}) > \right. \\ & \left. \min(f_1[\tau_n^{(m, C_\rho^{(g)})}, \tau_n^{(m+1)}], f_2[\tau_n^{(m, C_\rho^{(g)})}, \tau_n^{(m+1)}]) - \frac{\epsilon f(L)}{2} \right) \\ & \geq P_1 P_2 = 1 - o(L^{-2}). \end{aligned}$$

**Proof of (168):** We will now prove (168) by considering two cases depending on length of  $\tau_n^{(m, C_\rho^{(g)})} - \tau_n^{(m)}$ .

We consider

$$\tau_n^{(m, C_\rho^{(g)})} - \tau_n^{(m)} \leq \frac{\epsilon f(L)}{4(1+K)^{C_\rho^{(g)}} R} \quad (172)$$

or

$$\tau_n^{(m, C_\rho^{(g)})} - \tau_n^{(m)} > \frac{\epsilon f(L)}{4(1+K)^{C_\rho^{(g)}} R}, \quad (173)$$

where  $R := \sum_{j \in \mathcal{J}} \lambda_j + K\mu_{\max}$ .

**Case (172):** We notice  $x_{\bar{k}^{(\ell)}}^{L(e)}(\tau)$  will change by at most  $(1+K)^{\ell-1}/L$  at each arrival or departure according to Lemma D.1 and thus  $\sum_{\ell=1}^{C_\rho^{(g)}} \bar{k}_j^{(\ell)} x_{\bar{k}^{(\ell)}}^{L(e)}(\tau)$  will change by at most  $(1+K)^{C_\rho^{(g)}}/L - 1/L$ .

Using Lemma F.1, with probability  $1 - o(L^{-2})$ , the number of arrivals and departures in interval  $[\tau_n^{(m)}, \tau_n^{(m, C_\rho^{(g)})}]$  of length at most  $\frac{\epsilon f(L)}{4(1+K)^{C_\rho^{(g)}} R}$  is at most

$$\begin{aligned} & \left( \sum_{j \in \mathcal{J}} \lambda_j + K\mu_{\max} \right) L \frac{\epsilon f(L)}{4(1+K)^{C_\rho^{(g)}} R} + o(Lf(L)) = \\ & \frac{\epsilon Lf(L)}{4(1+K)^{C_\rho^{(g)}}} + o(Lf(L)), \end{aligned}$$

therefore

$$\mathbb{P} \left( \sum_{\ell=1}^{C_\rho^{(g)}} \bar{k}_j^{(\ell)} (x_{\bar{k}^{(\ell)}}^{(e)}(\tau_n^{(m, C_\rho^{(g)})}) - x_{\bar{k}^{(\ell)}}^{(e)}(\tau_n^{(m)})) \leq \frac{\epsilon f(L)}{4} \right) = 1 - o(L^{-2}). \quad (174)$$

Considering (172) holds, we will also have

$$\mathbb{P} \left( (\lambda_j - \mu_j y_j(t)) (\tau_n^{(m, C_\rho^{(g)})} - \tau_n^{(m)}) \leq \frac{\epsilon f(L)}{4} \right) = 1. \quad (175)$$

It is now easy to verify that equations (174) and (175) imply (168).

**Case (173):** In this case we notice, using Lemma F.1 for the process of jobs of type  $j$  in the system which is Poisson with rate

$L(\lambda_j - \mu_j y_j(t))$ , that with probability  $1 - o(L^{-2})$

$$\begin{aligned} 0 & \leq q_{\bar{k}^{(C_\rho^{(g)})}, j}^{L}(\tau_n^{(m, \ell m)}) = q_{\bar{k}^{(C_\rho^{(g)})}, j}^{L}(\tau_n^{(m)}) + \\ & \sum_{\ell=1}^{C_\rho^{(g)}} \bar{k}_j^{(\ell)} L (x_{\bar{k}^{(\ell)}}^{L(e)}(\tau_n^{(m, \ell m)}) - x_{\bar{k}^{(\ell)}}^{L(e)}(\tau_n^{(m)})) \\ & - Ly_j^L(\tau_n^{(m, C_\rho^{(g)})}) - Ly_j^L(\tau_n^{(m)}) \\ & \leq K + \sum_{\ell=1}^{C_\rho^{(g)}} \bar{k}_j^{(\ell)} L \left( x_{\bar{k}^{(\ell)}}^{L(e)}(\tau_n^{(m, C_\rho^{(g)})}) - x_{\bar{k}^{(\ell)}}^{L(e)}(\tau_n^{(m)}) \right) \\ & - L(\lambda_j - \mu_j y_j(t)) (\tau_n^{(m, C_\rho^{(g)})} - \tau_n^{(m)}) + o(Lf(L)), \end{aligned} \quad (176)$$

from which it follows

$$\begin{aligned} & \sum_{\ell=1}^{C_\rho^{(g)}} \bar{k}_j^{(\ell)} \left( x_{\bar{k}^{(C_\rho^{(g)})}}^{L(e)}(\tau_n^{(m, C_\rho^{(g)})}) - x_{\bar{k}^{(C_\rho^{(g)})}}^{L(e)}(\tau_n^{(m)}) \right) \geq \\ & (\lambda_j - \mu_j y_j(t)) (\tau_n^{(m, C_\rho^{(g)})} - \tau_n^{(m)}) + o(Lf(L)), \end{aligned} \quad (177)$$

which implies (168).

**Proof of (169):** We will now prove (169) by considering two cases depending on length of  $\tau_n^{(m+1)} - \tau_n^{(m, C_\rho^{(g)})}$  and reach a contradiction for each of them.

We consider

$$\tau_n^{(m+1)} - \tau_n^{(m, C_\rho^{(g)})} \leq \frac{\epsilon f(L)}{4(1+K)^{C_\rho^{(g)}-1} R} \quad (178)$$

or

$$\tau_n^{(m+1)} - \tau_n^{(m, C_\rho^{(g)})} > \frac{\epsilon f(L)}{4(1+K)^{C_\rho^{(g)}-1} R} \quad (179)$$

where  $R := \sum_{j \in \mathcal{J}} \lambda_j + K\mu_{\max}$ .

**Case (178):** We notice  $x_{\bar{k}^{(\ell)}}^{(e)}(\tau)$  will change by at most  $(1+K)^{\ell-1}/L$  at each arrival or departure according to Lemma D.1 and thus  $(x_{\bar{k}^{(e)}}^{(e)}(\tau) + \sum_{\ell=1}^{C_\rho^{(g)}-1} x_{\bar{k}^{(\ell)}}^{(e)}(\tau)^+)$  will change by at most  $\frac{(1+K)^{C_\rho^{(g)}}}{KL}$ .

Using Lemma F.1, with probability  $1 - o(L^{-2})$ , the number of arrivals and departures in interval  $[\tau_n^{(m, C_\rho^{(g)})}, \tau_n^{(m+1)}]$  of length at most  $\frac{\epsilon f(L)}{4(1+K)^{C_\rho^{(g)}-1} R}$  is at most

$$\begin{aligned} & \left( \sum_{j \in \mathcal{J}} \lambda_j + K\mu_{\max} \right) L \frac{\epsilon f(L)}{4(1+K)^{C_\rho^{(g)}-1} R} + o(Lf(L)) = \\ & \frac{\epsilon Lf(L)}{4(1+K)^{C_\rho^{(g)}-1}} + o(Lf(L)), \end{aligned}$$

therefore

$$\begin{aligned} & \mathbb{P} \left( \sum_{\ell=1}^{C_\rho^{(g)}} \bar{k}_j^{(\ell)} \left( x_{\bar{k}^{(\ell)}}^{(e)}(\tau_n^{(m+1)}) - x_{\bar{k}^{(\ell)}}^{(e)}(\tau_n^{(m, C_\rho^{(g)})}) \right) \leq \frac{\epsilon f(L)}{4} \right) = \\ & 1 - o(L^{-2}). \end{aligned} \quad (180)$$

Considering (178) holds, we will also have

$$\mathbb{P} \left( \mu_{\min} \frac{1 - \sum_{i=1}^{C_\rho^{(g)}} x_{\mathbf{k}^{(i)}}^{(e)}(t)}{JKC^2} \left( \tau_n^{(m+1)} - \tau_n^{(m, C_\rho^{(g)})} \right) \leq \frac{\epsilon f(L)}{4} \right) = 1. \quad (181)$$

It is now easy to verify that equations (180) and (181) imply (169).

**Case (179):** In this case we notice that whenever a server without effective configuration in set  $\mathcal{K} := \{\mathbf{k}^{(\ell)} : \ell = 1, \dots, C_\rho^{(g)}\}$  empties, it will be assigned to one of the configurations of  $\mathcal{K}$ . This statement is equivalent to the following, considering the bound of Lemma F.3,

$$\begin{aligned} & \left( x_{\bar{\mathbf{k}}^{(C_\rho^{(g)})}}^{L(e)}(\tau_n^{(m+1)}) - x_{\bar{\mathbf{k}}^{(C_\rho^{(g)})}}^{L(e)}(\tau_n^{(m, C_\rho^{(g)})}) \right) \\ & + \sum_{\ell=1}^{C_\rho^{(g)}-1} \left( x_{\bar{\mathbf{k}}^{(\ell)}}^{L(e)}(\tau_n^{(m+1)}) - x_{\bar{\mathbf{k}}^{(\ell)}}^{L(e)}(\tau_n^{(m, C_\rho^{(g)})}) \right)^+ \\ & - \frac{\mu_{\min}}{JKC^2} \left( 1 - \sum_{\ell=1}^{C_\rho^{(g)}} x_{\bar{\mathbf{k}}^{(\ell)}}^{(e)}(t) \right) \left( \tau_n^{(m+1)} - \tau_n^{(m, C_\rho^{(g)})} \right) > o(f(L)). \end{aligned}$$

Since this holds based on Lemma F.3 with probability  $1 - o(L^{-2})$ , it implies (169).

## H DETAILS OF PROOF OF PROPOSITION 6.2

We notice that the last equation of the system (51) is the same as the previous ones, if  $\Delta\eta_{i^*}$  has a coefficient  $\bar{k}_{\sigma_{I\rho}}^{(i^*)} := 1$ ,  $\Delta\eta_{g_j}$  has a coefficient  $\bar{k}_{\sigma_{I\rho}}^{(g_\ell)} := 1$  for  $\ell = 1, \dots, I_\rho$  and  $\Delta\theta_{\sigma_{I\rho}} = 0$ . Thus, in what follows we analyze the system in its most general form where

$$\Delta\eta_{i^*} + \sum_{j=1}^{I_\rho} \Delta\eta_{g_j} \leq 0$$

is replaced with

$$\bar{k}_{\sigma_{I\rho}}^{(i^*)} \Delta\eta_{i^*} + \sum_{\ell=1}^{I_\rho} \bar{k}_{\sigma_{I\rho}}^{(g_\ell)} \Delta\eta_{g_\ell} - \Delta\theta_{\sigma_{I\rho}} \leq 0.$$

As we showed in the main proof of Proposition 6.2, the values of  $\beta$  and  $\beta_j$ ,  $j = 1, \dots, I_\rho$ ,  $\gamma_j$ ,  $j = 1, \dots, J$ , which we want to prove they are positive, are given by the following system of equations

$$\begin{aligned} Z_{g_\ell} &= \sum_{j=\ell}^{I_\rho} \beta_j \bar{k}_{\sigma_j}^{(g_\ell)} \\ Z_{i^*} &= -\beta + \sum_{j=1}^{I_\rho} \beta_j \bar{k}_{\sigma_j}^{(i^*)} \\ Z &= \gamma_j + \beta_j \quad j \in \{\sigma_\ell : \ell = 1, \dots, I_\rho\} \\ Z &= \gamma_j \quad j \in \{\sigma_\ell : \ell = I_\rho + 1, \dots, J\}. \end{aligned} \quad (182)$$

It is straightforward from (182) that  $\beta_{I_\rho} = \frac{Z_{\sigma_{I\rho}}}{\bar{k}_{\sigma_{I\rho}}^{(g_{I\rho})}} > 0$ . We will

now show  $\bar{k}_{\sigma_\ell}^{(g_\ell)} \beta_\ell > Z_{g_\ell}/2 > 0$  for  $\ell = 1, \dots, I_\rho - 1$ , when  $Z_i > (2K+1)Z_{i+1}$ ,  $i = 1, \dots, C_\rho^{(g)} - 1$  based on assumptions. For shorthand

purposes we also define  $C := 2K + 1$ . The proof is as follows

$$\begin{aligned} \bar{k}_{\sigma_\ell}^{(g_\ell)} \beta_\ell &= Z_{g_\ell} - \sum_{j=\ell+1}^{I_\rho} \bar{k}_{\sigma_j}^{(g_\ell)} \beta_j \stackrel{(a)}{\geq} Z_{g_\ell} - K \sum_{j=\ell+1}^{I_\rho} Z_{g_j} \stackrel{(b)}{>} \\ & Z_{g_\ell} - K \sum_{j=\ell+1}^{I_\rho} Z_{g_\ell} C^{\ell-j} > Z_{g_\ell} \left( 1 - \frac{K}{C-1} \right) = Z_{g_\ell}/2. \end{aligned} \quad (183)$$

In (a) we used (182), according to which, considering  $\beta_{\ell'} > 0$  for  $\ell' = \ell + 1, \dots, I_\rho$ , we have  $\bar{k}_{\sigma_j}^{(g_j)} \beta_j < Z_{g_j}$  or  $\bar{k}_{\sigma_j}^{(g_\ell)} \beta_j < \frac{\bar{k}_{\sigma_j}^{(g_\ell)}}{\bar{k}_{\sigma_j}^{(g_j)}} Z_{g_j} \leq K Z_{g_j}$  and in (b) we used that for  $\ell < j$ ,  $Z_{g_j} < C^{\ell-j} Z_{g_{j+\ell-j}} \leq C^{\ell-j} Z_{g_\ell}$ .

To prove  $\beta > 0$ , suppose  $m$  is the lowest index for which  $\bar{k}_{\sigma_m}^{(i^*)} > 0$ . Then we will have

$$\beta = -Z_{i^*} + \sum_{j=1}^{I_\rho} \beta_j \bar{k}_{\sigma_j}^{(i^*)} > -Z_{i^*} + \beta_m \bar{k}_{\sigma_m}^{(i^*)} \stackrel{(a)}{>} -Z_{i^*} + \frac{\bar{k}_{\sigma_m}^{(i^*)}}{\bar{k}_{\sigma_m}^{(g_m)}} Z_{\sigma_m}/2 \stackrel{(b)}{>} 0.$$

Inequality (a) uses just that  $\bar{k}_{\sigma_\ell}^{(g_\ell)} \beta_\ell > Z_{\sigma_\ell}/2$  for  $\ell = m$ , which we have already proved. Inequality (b) follows considering that  $g_m < i^*$  and that  $Z_{i^*} < Z_{i^*+1}/C < \frac{\bar{k}_{\sigma_m}^{(i^*)}}{\bar{k}_{\sigma_m}^{(g_m)}} Z_{g_m}/2$ .

To show that  $g_m < i^*$  we notice that  $\bar{\mathbf{k}}^{(i^*)}$  and  $\bar{\mathbf{k}}^{(g_m)}$  are two different configurations whose job types belong to  $\{\sigma_m, \dots, \sigma_J\}$  and  $\bar{\mathbf{k}}^{(g_m)}$  is the configuration of maximum reward that has this property, so its index as given from Definition 3.3 should be lower than  $i^*$ .

Lastly we need to show that  $\gamma_j > 0$  for  $j = 1, \dots, J$ . If  $j \in \{I_\rho + 1, \dots, J\}$  then  $\gamma_j = Z > 0$ . If  $j \in \{1, \dots, I_\rho\}$  then by using that for  $\ell = 1, \dots, C_\rho^{(g)}$  we have  $Z > Z_\ell$  because of (48) and  $Z_{g_\ell} \geq \beta_\ell$  because of (182), we get

$$\gamma_j = Z - \beta_j > Z_{g_j} - \beta_j \geq 0. \quad (184)$$

## I PROOF OF LEMMA 6.4

Consider the function  $f(L)$  as in Definition 5.8. Recall that  $Z_i > \xi Z_{i+1}$  for  $i = 1, \dots, C_\rho^{(g)} - 1$  and  $Z_{C_\rho^{(g)}} > 0$ . We choose  $\xi$  such that:

$$\xi > \frac{\mu_{\max}}{\mu_{\min}} \left( 12K^2 + 16CK^2 \frac{2(\mu_{\max}K + \sum_{j=1}^J \lambda_j)}{\delta} \left( 12K \frac{\mu_{\max}}{\mu_{\min}} \right)^{C_\rho^{(g)}} \right), \quad (185)$$

and  $Z$  is chosen such that  $Z > 4Z_1$ . We first show the following lemma.

**LEMMA I.1.** For any  $m \in \{1, \dots, M_n\}$ , with probability greater than  $1 - o(L^{-2})$ ,

$$\begin{aligned} & \sum_{i=1}^{C_\rho^{(g)}} Z_i \left( x_{\bar{\mathbf{k}}^{(i)}}^{L(e)}(\tau_n^{(m+1)}) - x_{\bar{\mathbf{k}}^{(i)}}^{L(e)}(\tau_n^{(m)}) \right) + \\ & \sum_{j=1}^J Z(\tau_n^{(m+1)} - \tau_n^{(m)}) (\mu_j y_j(t) - \lambda_j)^+ \geq \\ & \delta(\epsilon_V)(\tau_n^{(m+1)} - \tau_n^{(m)}) + o(f(L)). \end{aligned} \quad (186)$$

PROOF. We will use P.1., P.2. and P.3. to refer to the properties in Proposition 5.10 and whenever we apply such a property any resulting relation holds with probability  $1 - o(L^{-2})$ .

For compactness we also define

$$W^{(m)}(t) := \sum_{i=1}^{C_\rho^{(g)}} Z_i \nabla x_{\bar{k}^{(i)}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] + \sum_{j=1}^J Z(\mu_j y_j(t) - \lambda_j)^+.$$

Our objective is thus to find  $\delta(\epsilon_V)$  such that

$$W^{(m)}(t) \geq \delta(\epsilon_V) + \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}}, \quad (187)$$

with probability  $1 - o(L^{-2})$ .

We will analyze two separate cases depending on whether  $\ell_m < C_\rho^{(g)}$  or  $\ell_m = C_\rho^{(g)}$ .

**Case  $\ell_m < C_\rho^{(g)}$ :** Considering Lemma F.2 and Property P.2., we first can get the following bound

$$\begin{aligned} & \sum_{i=\ell_m}^{C_\rho^{(g)}} Z_i \left( x_{\bar{k}^{(i)}}^{L(e)}(\tau_n^{(m+1)}) - x_{\bar{k}^{(i)}}^{L(e)}(\tau_n^{(m)}) \right) > \\ & \left( Z_{\ell_m} \alpha_{\ell_m} - \sum_{i=\ell_m+1}^{C_\rho^{(g)}} Z_i B_i \right) \left( \tau_n^{(m+1)} - \tau_n^{(m)} \right) + o(f(L)) \stackrel{(a)}{>} \\ & Z_{\ell_m} \alpha_{\ell_m} / 2 (\tau_n^{(m+1)} - \tau_n^{(m)}) + o(f(L)), \end{aligned} \quad (188)$$

where (a) follows from definitions of  $\xi$ , for which  $Z_i > \xi Z_{i+1}$ , and  $\alpha_{\ell_m}$  given in (185) and (90) respectively. If we further show that

$$\sum_{i=1}^{\ell_m-1} Z_i \nabla x_{\bar{k}^{(i)}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] + \sum_{j=1}^J Z(\mu_j y_j(t) - \lambda_j)^+ \geq \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}}, \quad (189)$$

then it follows from (189) and (188) that

$$W^{(m)}(t) \geq Z_{\ell_m} \alpha_{\ell_m} / 2 + \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}}. \quad (190)$$

One way to show (189) is to find constants  $Z_i^*$  for  $i = 1, \dots, \ell_m - 1$  such that

$$\begin{aligned} & \sum_{i=1}^{\ell_m-1} Z_i \nabla x_{\bar{k}^{(i)}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] + \sum_{j=1}^J Z \mu_j (y_j(t) - \rho_j)^+ \geq \\ & \sum_{i=1}^{\ell_m-1} Z_i^* (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t)) + \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}} \end{aligned} \quad (191)$$

and

$$Z_i^* (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t)) \geq 0 \quad i = 1, \dots, \ell_m - 1. \quad (192)$$

A choice of constants  $Z_i^*$  that satisfies (191) and (192) for  $i = 1, \dots, \ell_m - 1$  is

$$Z_i^* = Z'_i - \mathbb{1}(\lambda_{j_i} - \mu_{j_i} y_{j_i}(t) < 0) Z 2^{-i}$$

where constants  $Z'_i$  are given by the following system of equations

$$\sum_{\ell=i}^{\ell_m-1} \bar{k}_{j_i}^{(\ell)} Z'_\ell = Z_i \quad i = 1, \dots, \ell_m - 1. \quad (193)$$

We will now justify why this choice of  $Z_i^*$  satisfies (191) and (192). The requirement (191) can be inferred by adding the next two relationships.

The first relationship is

$$\begin{aligned} & \sum_{i=1}^{\ell_m-1} Z'_i (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t)) \stackrel{(a)}{=} \\ & \sum_{i=1}^{\ell_m-1} Z'_i \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} \nabla x_{\bar{k}^{(\ell)}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] + \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}} \stackrel{(b)}{=} \\ & \sum_{i=1}^{\ell_m-1} Z_i \nabla x_{\bar{k}^{(i)}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] + \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}}, \end{aligned} \quad (194)$$

where (a) is due to P.1. for  $i = 1, \dots, \ell_m - 1$  and (b) is due to (193).

The second relationship is

$$\begin{aligned} & - \sum_{i=1}^{\ell_m-1} Z 2^{-i} (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t)) \mathbb{1}(\mu_{j_i} y_{j_i}(t) > \lambda_{j_i}) = \\ & \sum_{j=1}^J Z(\mu_j y_j(t) - \lambda_j)^+ \sum_{i=1}^{\ell_m-1} 2^{-i} \mathbb{1}(j_i = j) \leq \sum_{j=1}^J Z(\mu_j y_j(t) - \lambda_j)^+. \end{aligned} \quad (195)$$

Finally, we should prove that (192) also holds. If  $\lambda_{j_i} - \mu_{j_i} y_{j_i}(t) \geq 0$  it suffices that  $Z_i^* > 0$  or  $Z'_i > 0$ . For this, we will show recursively that

$$2Z_i > Z'_i > 0, \quad i = 1, \dots, \ell_m - 1, \quad (196)$$

For  $i = \ell_m - 1$ , using (193), we have

$$2Z_{\ell_m-1} > Z_{\ell_m-1} \frac{1}{\bar{k}_{j_{\ell_m-1}}^{(\ell_m-1)}} = Z'_{\ell_m-1} > 0, \quad (197)$$

while for  $i < \ell_m - 1$ ,

$$\begin{aligned} & 2Z_i > Z_i \frac{1}{\bar{k}_{j_i}^{(i)}} - \sum_{\ell=i+1}^{\ell_m-1} \frac{\bar{k}_{j_i}^{(\ell)}}{\bar{k}_{j_i}^{(i)}} Z'_\ell \\ & \stackrel{(a)}{=} Z'_i > \frac{1}{\bar{k}_{j_i}^{(i)}} \left( Z_i - \sum_{\ell=i+1}^{\ell_m-1} K Z'_\ell \right) \\ & > \frac{1}{\bar{k}_{j_i}^{(i)}} \left( Z_i - \sum_{\ell=i+1}^{\ell_m-1} K 2Z_\ell \right) \stackrel{(b)}{>} \frac{1}{\bar{k}_{j_i}^{(i)}} Z_{\ell_m-1} > 0, \end{aligned} \quad (198)$$

where in (a) we used (193), and in (b) we used  $Z_\ell > (2K+1)Z_{\ell+1}$  for any  $\ell < \ell_m - 1$ . Notice that this claim is consistent with assumption  $Z_\ell > \xi Z_{\ell+1}$ , since  $\xi > 2K+1$ .

If  $\lambda_{j_i} - \mu_{j_i} y_{j_i}(t) < 0$ , it suffices that  $Z_i^* < 0$  or  $Z'_i < 2^{-i} Z$ . We can get this result from (196), which we proved earlier, as follows

$$Z'_i < 2Z_i \leq 2(2K+1)^{-i+1} Z_1 < 2^{-i} Z. \quad (199)$$

where the last inequality is because  $4Z_1 < Z$ .

**Case  $\ell_m = C_\rho^{(g)}$ :** For notation compactness, for  $j \in \mathcal{J}$  such that  $\bar{k}_j^{(C_\rho^{(g)})} > 0$ , we define:

$$f_j(t) := \frac{1}{\bar{k}_j^{(C_\rho^{(g)})}} \left( \lambda_j - \mu_j y_j(t) - \sum_{i=1}^{C_\rho^{(g)}-1} \bar{k}_j^{(i)} h^{\mathcal{J}[m],(i)}(t) \right),$$

$$f^*(t) := \frac{\mu \min}{JKC^2} \left( 1 - \sum_{i=1}^{C_\rho^{(g)}} x_{\mathbf{k}^{(i)}}^{(e)}(t) \right) - \sum_{i=1}^{C_\rho^{(g)}-1} \left( h^{\mathcal{J}[m],(i)}(t) \right)^+,$$

$$j' := \arg \min_{j \in \mathcal{J}: \bar{k}_j^{(C_\rho^{(g)})} > 0} f_j(t).$$

If  $C_\rho^{(g)} = C^{(g)}$  the set  $\{j \in \mathcal{J} : \bar{k}_j^{(C_\rho^{(g)})} > 0\}$  is empty in which case we consider  $j' = \emptyset$ . We distinguish two sub-cases.

**Subcase  $j' = \emptyset$  or  $f_{j'}(t) \geq f^*(t)$ :** In this case it suffices to find constants  $Z_i^*$  for  $i = 1, \dots, \ell_m$ , such that

$$\sum_{i=1}^{C_\rho^{(g)}} Z_i \nabla x_{\mathbf{k}^{(i)}}^{(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] + \sum_{j=1}^J Z \mu_j (y_j(t) - \rho_j)^+ \geq$$

$$\sum_{i=1}^{C_\rho^{(g)}-1} Z_i^* (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t)) + Z_{C_\rho^{(g)}}^* \frac{\mu \min}{JKC^2} \left( 1 - \sum_{i=1}^{C_\rho^{(g)}} x_{\mathbf{k}^{(i)}}^{(e)}(t) \right) \quad (200)$$

$$+ \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}},$$

and if we further define

$$W_{C_\rho^{(g)}}^a(\mathbf{z}(t)) := \sum_{i=1}^{C_\rho^{(g)}-1} Z_i^* (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t)) +$$

$$Z_{C_\rho^{(g)}}^* \frac{\mu \min}{JKC^2} \left( 1 - \sum_{i=1}^{C_\rho^{(g)}} x_{\mathbf{k}^{(i)}}^{(e)}(t) \right). \quad (201)$$

then

$$Z_i^* (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t)) \geq 0 \quad i = 1, \dots, C_\rho^{(g)} - 1, \quad (202)$$

$$Z_{C_\rho^{(g)}}^* \frac{\mu \min}{JKC^2} \left( 1 - \sum_{i=1}^{C_\rho^{(g)}} x_{\mathbf{k}^{(i)}}^{(e)}(t) \right) \geq 0, \quad (203)$$

and

$$W_{C_\rho^{(g)}}^a(\mathbf{z}(t)) > 0 \Leftrightarrow \mathbf{z}(t) \notin \Gamma^*. \quad (204)$$

A choice of  $Z_i^*$  that satisfies those requirements is

$$Z_{C_\rho^{(g)}}^* = Z_{C_\rho^{(g)}}, \quad (205)$$

and

$$Z_i^* = Z'_i - \mathbb{1}(\lambda_{j_i} - \mu_{j_i} y_{j_i}(t) < 0) Z 2^{-i} \quad (206)$$

where constants  $Z'_i$  are given by the following system of equations

$$\mathbb{1}(h^{\mathcal{J}[m],(i)}(t) > 0) Z_{C_\rho^{(g)}}^* + \sum_{\ell=i}^{C_\rho^{(g)}-1} \bar{k}_{j_i}^{(\ell)} Z'_\ell = Z_i \quad i = 1, \dots, C_\rho^{(g)} - 1. \quad (207)$$

We will now justify why this choice of  $Z_i^*$  satisfies (200) and (202) and (203). To prove (200) we can add three relationships (208),

(209) and (210) that we prove below. The first relationship is

$$\sum_{i=1}^{C_\rho^{(g)}-1} Z'_i (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t)) \stackrel{(a)}{=} \sum_{i=1}^{C_\rho^{(g)}-1} Z'_i \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} \nabla x_{\mathbf{k}^{(\ell)}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] + \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}} = \quad (208)$$

$$\sum_{i=1}^{C_\rho^{(g)}-1} Z_i \nabla x_{\mathbf{k}^{(i)}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] + \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}},$$

where (a) is due to P.1. for  $i = 1, \dots, C_\rho^{(g)} - 1$ . The second relationship is

$$Z_{C_\rho^{(g)}}^* \frac{\mu \min}{JKC^2} \left( 1 - \sum_{i=1}^{C_\rho^{(g)}} x_{\mathbf{k}^{(i)}}^{(e)}(t) \right) \stackrel{(a)}{\leq} Z_{C_\rho^{(g)}}^* \nabla x_{\bar{\mathbf{k}}^{(C_\rho^{(g)})}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}]$$

$$+ Z_{C_\rho^{(g)}}^* \sum_{i=1}^{C_\rho^{(g)}-1} \left( \nabla x_{\mathbf{k}^{(i)}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] \right)^+ + \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}} \stackrel{(b)}{=} Z_{C_\rho^{(g)}}^* \nabla x_{\bar{\mathbf{k}}^{(C_\rho^{(g)})}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}]$$

$$+ Z_{C_\rho^{(g)}}^* \sum_{i=1}^{C_\rho^{(g)}-1} \mathbb{1}(h^{\mathcal{J}[m],(i)}(t) > 0) \nabla x_{\mathbf{k}^{(i)}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}]$$

$$+ \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}}, \quad (209)$$

where (a) is due to P.3. and assumption  $j' = \emptyset$  or  $f_{j'}(t) \geq f(t)$ , while (b) is due to Lemma G.2. The third relationship is

$$\sum_{i=1}^{C_\rho^{(g)}-1} Z 2^{-i} (\mu_{j_i} y_{j_i}(t) - \lambda_{j_i}) \mathbb{1}(y_{j_i}(t) > \rho_{j_i}) =$$

$$\sum_{j=1}^J Z (\mu_j y_j(t) - \lambda_j)^+ \sum_{i=1}^{C_\rho^{(g)}-1} 2^{-i} \mathbb{1}(j_i = j) < \sum_{j=1}^J Z \mu_j (y_j(t) - \rho_j)^+. \quad (210)$$

Then to prove (202) it suffices to show, just like in the case  $\ell_m < C_\rho^{(g)}$ , that for  $i = 1, \dots, C_\rho^{(g)} - 1$ ,  $Z_i^* > 0$  if  $\lambda_{j_i} - \mu_{j_i} y_{j_i}(t) \geq 0$ , and  $Z_i^* < 0$  otherwise, while for (203) to be true we need  $Z_{C_\rho^{(g)}}^* > 0$ ,

since  $(1 - \sum_{i=1}^{C_\rho^{(g)}} x_{\mathbf{k}^{(i)}}^{(e)}(t)) \geq 0$ .

If we define  $Z'_{C_\rho^{(g)}} := Z_{C_\rho^{(g)}}^*$  then it suffices to show recursively

$$2Z_i > Z'_i > 0 \quad i = 1, \dots, C_\rho^{(g)}. \quad (211)$$

The process is very similar to the case  $\ell_m < C_\rho^{(g)}$ . We can now prove (204) as follows.

Considering

$$1 - \sum_{i=1}^{C_\rho^{(g)}} x_{\bar{k}^{(i)}}^{(e)}(t) = \sum_{i=1}^{C_\rho^{(g)}} \left( x_{\bar{k}^{(i)}}^{(g)} - x_{\bar{k}^{(i)}}^{(e)}(t) \right)$$

and for  $i = 1, \dots, C_\rho^{(g)} - 1$ ,

$$\mu_{j_i} y_{j_i}(t) - \lambda_{j_i} = \sum_{\ell=1}^i \bar{k}_{j_i}^{(\ell)} (x_{\bar{k}^{(\ell)}}^{(e)}(t) - x_{\bar{k}^{(\ell)}}^{(g)}).$$

then  $W_{C_\rho^{(g)}}^a(\mathbf{z}(t)) = 0$ , if and only if  $x_{\bar{k}^{(i)}}^{(g)} = x_{\bar{k}^{(i)}}^{(e)}(t)$  for  $i = 1, \dots, C_\rho^{(g)}$  or equivalently if and only if  $\mathbf{z}(t) \in \Gamma^*$ . If we define the vectors  $\mathbf{x}(t) := (x_{\bar{k}^{(i)}}^{(e)}(t))_{i=1, \dots, C_\rho^{(g)}}$ ,  $\mathbf{x} := (x_{\bar{k}^{(i)}}^{(g)})_{i=1, \dots, C_\rho^{(g)}}$ ,  $\Delta \mathbf{x}(t) := \mathbf{x}(t) - \mathbf{x}$  and consider  $x$  the zero vector in space  $\Gamma[\epsilon_\rho]$ , then we can also verify  $W_{C_\rho^{(g)}}^a(\mathbf{z}(t))$  satisfies the subadditive and absolutely scalable properties, i.e

$$\begin{aligned} W_{C_\rho^{(g)}}^a(\Delta \mathbf{x}_1(t)) + W_{C_\rho^{(g)}}^a(\Delta \mathbf{x}_2(t)) &\leq W_{C_\rho^{(g)}}^a(\Delta \mathbf{x}_1(t) + \Delta \mathbf{x}_2(t)) \\ W_{C_\rho^{(g)}}^a(\alpha \Delta \mathbf{x}(t)) &= |\alpha| W_{C_\rho^{(g)}}^a(\Delta \mathbf{x}(t)). \end{aligned} \quad (212)$$

Thus  $W_{C_\rho^{(g)}}^a(\mathbf{z}(t))$  has the properties of norm in space  $\Gamma[\epsilon_\rho]$  and since this space has finite dimensions all of its norms are equivalent which means there is  $c^a$  such that

$$W_{C_\rho^{(g)}}^a(\mathbf{z}(t)) \geq c^a V(\mathbf{z}(t)) > c^a \epsilon_V \quad (213)$$

**Subcase**  $f_{j'}(t) < f(t)$ : In this case it suffices to prove

$$\begin{aligned} \sum_{i=1}^{C_\rho^{(g)}} Z_i \nabla x_{\bar{k}^{(i)}}^{(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] + \sum_{j=1}^J Z \mu_j (y_j(t) - \rho_j)^+ &\geq \\ \sum_{i=1}^{C_\rho^{(g)}} Z_i^* (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t)) + \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}}, \end{aligned} \quad (214)$$

and

$$Z_i^* (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t)) \geq 0 \quad i = 1, \dots, C_\rho^{(g)}. \quad (215)$$

A choice of values of  $Z_i^*$  that satisfies those requirements for  $i = 1, \dots, C_\rho^{(g)} - 1$  is

$$Z_i^* = Z'_i - \mathbb{1}(\lambda_{j_i} - \mu_{j_i} y_{j_i}(t) < 0) Z 2^{-i} \quad (216)$$

and if we further define

$$W_{C_\rho^{(g)}}^b(\mathbf{z}(t)) := \sum_{i=1}^{C_\rho^{(g)}} Z_i^* (\lambda_{j_i} - \mu_{j_i} y_{j_i}(t)) = 0. \quad (217)$$

then

$$Z_{C_\rho^{(g)}}^* = Z'_{C_\rho^{(g)}} - \mathbb{1}(\lambda_{j'} - \mu_{j'} y_{j'}(t) < 0) Z 2^{-C_\rho^{(g)}}. \quad (218)$$

and

$$W_{C_\rho^{(g)}}^b(\mathbf{z}(t)) > 0 \Leftrightarrow \mathbf{z}(t) \notin \Gamma^*. \quad (219)$$

Constants  $Z'_i$  are given by the following system of equations

$$\begin{aligned} \bar{k}_{j'}^{(i)} Z'_{C_\rho^{(g)}} + \sum_{\ell=i}^{C_\rho^{(g)}-1} \bar{k}_{j'}^{(\ell)} Z'_\ell &= Z_i \quad i = 1, \dots, C_\rho^{(g)} - 1, \\ \bar{k}_{j'}^{(C_\rho^{(g)})} Z'_{C_\rho^{(g)}} &= Z_{C_\rho^{(g)}}. \end{aligned} \quad (220)$$

We will now justify why this choice of  $Z_i^*$  satisfies (214) and (215). To prove (214) we can add relationships (208) and (210) from sub-case  $f_{j'}(t) \geq f(t)$  and (221) proven next.

$$\begin{aligned} Z'_{C_\rho^{(g)}} (\lambda_{j'} - \mu_{j'} y_{j'}(t)) &\stackrel{(a)}{\leq} \\ Z'_{C_\rho^{(g)}} \sum_{i=1}^{C_\rho^{(g)}-1} \bar{k}_{j'}^{(i)} \nabla x_{\bar{k}^{(i)}}^{L(e)}[\tau_n^{(m)}, \tau_n^{(m+1)}] + \frac{o(f(L))}{\tau_n^{(m+1)} - \tau_n^{(m)}}, \end{aligned} \quad (221)$$

where (a) is due to P.3. and assumption  $f_{j'}(t) < f(t)$ .

By using systems of equations (216) and (218), we can show, similarly to the case  $\ell_m < C_\rho^{(g)}$ , that for  $i = 1, \dots, C_\rho^{(g)}$ ,  $Z_i^* > 0$  if  $\lambda_{j_i} - \mu_{j_i} y_{j_i}(t) \geq 0$  and  $Z_i^* < 0$  otherwise, by proving recursively

$$2Z_i > Z'_i > 0 \quad i = 1, \dots, C_\rho^{(g)}. \quad (222)$$

Similarly with  $W_{C_\rho^{(g)}}^a(\mathbf{z}(t))$ , we can show (219) as  $W_{C_\rho^{(g)}}^b(\mathbf{z}(t))$  satisfies all the properties of a norm in space  $\Gamma[\epsilon_\rho]$ .

Also, since this space has finite dimensions all of its norms are equivalent which means there is  $c^b$  such that

$$W_{C_\rho^{(g)}}^b(\mathbf{z}(t)) \geq c^b V(\mathbf{z}(t)) > c^b \epsilon_V. \quad (223)$$

**Determining**  $\delta(\epsilon_V)$ : Expressions (190), (213) and (223) give a lower bound on  $W^{(m)}(t)$  in three different cases, thus for any  $m \in \{1, \dots, M_n\}$  a value of  $\delta(\epsilon_V)$  is the minimum of these expressions i.e.

$$\delta(\epsilon_V) := \min \left( \min_{i=1, \dots, C_\rho^{(g)}-1} Z_i \alpha_i / 2, c^a \epsilon_V, c^b \epsilon_V \right). \quad (224)$$

This completes the proof.  $\square$

Summing (186) in Lemma I.1, over all  $m \in \{1, \dots, M_n\}$ , we get

$$\begin{aligned} \sum_{i=1}^{C_\rho^{(g)}} Z_i \left( X_{\bar{k}^{(i)}}^{L(e)}(\tau_{n+1}) - X_{\bar{k}^{(i)}}^{L(e)}(\tau_n) \right) &\geq \\ \delta(\epsilon_V) (\tau_{n+1} - \tau_n) - \sum_{j=1}^J Z (\tau_{n+1} - \tau_n) \mu_j (y_j(t) - \rho_j)^+ + o(f(L)). \end{aligned} \quad (225)$$

Since each of (186) holds with probability  $1 - o(L^{-2})$  and (225) is a finite sum of them, it will also be satisfied with probability  $1 - o(L^{-2})$ .

To show that (225) implies (54), it remains to prove that

$$\frac{d}{dt} (y_j(t) - \rho_j)^+ \leq -\mu_j (y_j(t) - \rho_j)^+. \quad (226)$$

If  $y_j(t) < \rho_j$ , then  $(y_j(t) - \rho_j)^+ = 0$ , and obviously

$$\frac{d}{dt} (y_j(t) - \rho_j)^+ = \frac{d}{dt} 0 = 0. \quad (227)$$



If  $y_j(t) > \rho_j$ , then  $(y_j(t) - \rho_j)^+ = y_j(t) - \rho_j$  and hence

$$\begin{aligned} \frac{d}{dt}(y_j(t) - \rho_j)^+ &= \frac{d}{dt}y_j(t) \\ &\leq \lambda_j - y_j(t)\mu_j = -\mu_j(y_j(t) - \rho_j), \end{aligned} \quad (228)$$

where in the last inequality we used the fact that rate that type- $j$  jobs at fluid limit are admitted cannot be more than  $\lambda_j$  (not all type- $j$  jobs that arrive are admitted), and existing type- $j$  jobs in the system depart at rate  $\mu_j y_j(t)$  in fluid limit.

## J PROOF OF THEOREM 4.1

Using Theorem 6.1, we first show that, as  $L \rightarrow \infty$ , the sequence of stationary random variables  $\mathbf{x}^{L(e)(\infty)}$  converges in distribution to  $\mathbf{x}^{(g)}$  (the unique global greedy assignment), i.e.,

$$\mathbf{x}^{L(e)(\infty)} \implies \mathbf{x}^{(g)}.$$

By Theorem 6.1, given an  $\epsilon_1 > 0$ , we can choose  $t_{\epsilon_1}$  large enough such that  $\|\mathbf{x}^{(e)}(t) - \mathbf{x}^{(g)}\| \leq \epsilon_1$ , for  $t \geq t_{\epsilon_1}$ . Also by Theorem 5.4, we can choose a subsequence of  $L_n$  of  $L$  such that  $x_k^{L_n(e)}(t) \rightarrow x_k^{(e)}(t)$  (u.o.c). Now for an  $\epsilon_2 > 0$ , and  $L_n$  large enough, we can choose an  $\epsilon_3$  such that uniformly over all initial states  $\mathbf{z}^{L_n}(0)$  we have  $\|\mathbf{z}^{L_n}(0) - \mathbf{z}(0)\| \leq \epsilon_3$  and

$$\mathbb{P}\left(\|\mathbf{x}^{L_n(e)}(t_{\epsilon_1}) - \mathbf{x}^{(e)}(t_{\epsilon_1})\| < \epsilon_1\right) > 1 - \epsilon_2.$$

This is true because otherwise for a sequence of initial states  $\mathbf{z}^{L_n}(0) \rightarrow \mathbf{z}(0)$ ,  $\mathbb{P}\left(\|\mathbf{x}^{L_n(e)}(t_{\epsilon_1}) - \mathbf{x}^{(e)}(t_{\epsilon_1})\| < \epsilon_1\right) \leq 1 - \epsilon_2$ , which is impossible because, almost surely,  $\mathbf{x}^{L_n(e)}(t_{\epsilon_1}) \rightarrow \mathbf{x}^{(e)}(t_{\epsilon_1})$ . Hence,

$$\begin{aligned} &\mathbb{P}\left(\|\mathbf{x}^{L_n(e)}(t_{\epsilon_1}) - \mathbf{x}^{(g)}\| < 2\epsilon_1\right) > \\ &\mathbb{P}\left(\|\mathbf{x}^{L_n(e)}(t_{\epsilon_1}) - \mathbf{x}^{(e)}(t_{\epsilon_1})\| + \|\mathbf{x}^{(e)}(t_{\epsilon_1}) - \mathbf{x}^{(g)}\| < 2\epsilon_1\right) \\ &> \mathbb{P}\left(\|\mathbf{x}^{L_n(e)}(t_{\epsilon_1}) - \mathbf{x}^{(e)}(t_{\epsilon_1})\| < \epsilon_1\right) \\ &> 1 - \epsilon_2 \end{aligned}$$

which implies  $\mathbf{x}_k^{L_n(e)(\infty)} \implies \mathbf{x}_k^{(g)}$ , since  $\epsilon_1, \epsilon_2$  were chosen arbitrarily. Since this holds for every subsequence  $L_n$  of  $L$ , and all converge to the same limit  $\mathbf{x}^{(g)}$ , we can conclude  $\mathbf{x}^{L(e)(\infty)} \implies \mathbf{x}^{(g)}$  (e.g., see Theorem 2.6 of [9]).

Next, denote the normalized reward of the system at time  $t$  under DRA as  $U^L(t) = \frac{1}{L}F^{DRA}(t)$ , and the normalized reward at time  $t$  under the optimal policy as  $U^{\star L}(t) = \frac{1}{L}F^{\star}(t)$ . Then,

$$U^L(\infty) \geq \sum_{i=1}^{C^{(g)}} \sum_{j=1}^J x_{\bar{k}^{(i)}}^{L(e)(\infty)} \bar{k}_j^{(i)} - \sum_{j=1}^J \frac{g(L) + K}{L} u_j, \quad (229)$$

since there are at most  $g(L) + K$  empty slots for each job type in all the servers with effective configuration. Further note that random variables  $\{U^L(\infty)\}$ , and  $\{x_{\bar{k}^{(i)}}^{L(e)(\infty)}\}$  are uniformly integrable, so they also converge in expectation (e.g., see Theorem 3.5 of [9]), hence taking the expectations from both sides and letting  $L \rightarrow \infty$ ,

$$\lim_{L \rightarrow \infty} \mathbb{E}[U^L(\infty)] \geq \sum_{i=1}^{C^{(g)}} \sum_{j=1}^J x_{\bar{k}^{(i)}}^{(g)} \bar{k}_j^{(i)} = U^{(g)}[\boldsymbol{\rho}] \stackrel{(a)}{\geq} \frac{U^{\star}[\boldsymbol{\rho}]}{2}, \quad (230)$$

where (a) is a direct consequence of Theorem 3.7 (and  $\frac{1}{2}$  can be replaced with  $(1 - 1/e)$  if Theorem 3.9 is used). Also note that for the optimal algorithm  $\mathbb{E}[U^{\star L}(\infty)] \leq U^{\star}[\boldsymbol{\rho}]$ , since  $U^{\star}[\boldsymbol{\rho}]$  is the maximum possible normalized reward. Using this, and dividing both sides of (230) by  $\mathbb{E}[U^{\star L}(\infty)]$ , completes the proof.