

# SUITABLE SETS FOR STRONGLY TOPOLOGICAL GYROGROUPS

FUCAI LIN\*, TINGTING SHI, AND MENG BAO

ABSTRACT. A discrete subset  $S$  of a topological gyrogroup  $G$  with the identity  $0$  is said to be a *suitable set* for  $G$  if it generates a dense subgyrogroup of  $G$  and  $S \cup \{0\}$  is closed in  $G$ . In this paper, it was proved that each countable Hausdorff topological gyrogroup has a suitable set; moreover, it is shown that each separable metrizable strongly topological gyrogroup has a suitable set.

## 1. INTRODUCTION

In 1990, K.H. Hofmann and S.A. Morris in [15] introduced the concept of a suitable set for a topological group as an example of a 'thin' closed generating set. It was shown that each locally compact group has a suitable set. Fundamental results on suitable sets for topological groups were obtained by Comfort et al. in [8] and Dikranjan et al. in [9] and [10]. I. Guran in [14], F. Lin, A. Ravsky, and T. Shi in [18] considered suitable sets for paratopological groups. In 2003, T. Banakh and I. Protasov generalized Guran's results to left topological groups in [3].

A generalization of a group, gyrogroup (see Definition 2.2 below) was introduced by A.A. Ungar [27] in 2002, while studying a  $c$ -ball  $\mathbb{R}_c^3 = \{\mathbf{v} \in \mathbb{R}^3 : \|\mathbf{v}\| < c\}$  of relativistically admissible velocities endowed with Einstein velocity addition  $\oplus_E$ . Recall that for vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{R}_c^3$

$$\mathbf{u} \oplus_E \mathbf{v} = \frac{1}{1 + \frac{\langle \mathbf{u}, \mathbf{v} \rangle}{c^2}} \left( \mathbf{u} + \frac{1}{\gamma_{\mathbf{u}}} \mathbf{v} + \frac{1}{c^2} \frac{\gamma_{\mathbf{u}}}{1 + \gamma_{\mathbf{u}}} \langle \mathbf{u}, \mathbf{v} \rangle \mathbf{u} \right),$$

where

$$\gamma_{\mathbf{u}} = \frac{1}{\sqrt{1 - \frac{\|\mathbf{u}\|^2}{c^2}}}$$

is the Lorentz factor. It turned out that  $(\mathbb{R}_c^3, \oplus_E)$  is a gyrogroup, which fails to be a group, because the operation  $\oplus_E$  is not associative. Recently, the topic of gyrogroups was investigated by many scholars, see [12, 13, 16, 17, 19, 20, 21, 22, 23, 24, 25, 27].

In 2017, W. Atiponrat [2] introduced the concept of topological gyrogroups, which is a generalization of a topological group. Namely, a topological gyrogroup  $G$  is a gyrogroup  $(G, \oplus)$  endowed with a topology such that the multiplication map  $\oplus$  from  $G \times G$  to  $G$  is jointly continuous and the inverse map  $\ominus : G \rightarrow G$  is continuous. In turned out

---

1991 *Mathematics Subject Classification*. Primary 54H11; 22A05, Secondary 20B30; 20N99; 54A25; 54C35; 54D65; 54E35.

*Key words and phrases*. suitable sets; gyrogroups; (strongly) topological gyrogroups; left precompact; pseudocompact; separable space; metrizable space.

The first author is supported by the NSFC (No.11571158), the Natural Science Foundation of Fujian Province (No. 2017J01405) of China, the Program for New Century Excellent Talents in Fujian Province University, and the Institute of Meteorological Big Data-Digital Fujian and Fujian Key Laboratory of Data Science and Statistics. The second author is supported by the Young and middle-aged project in Fujian Province (No. JAT190397).

\* Corresponding author.

that topological gyrogroups possess nice properties. In particular, Z. Cai, S. Lin, and W. He in [7] proved that every topological gyrogroup is a rectifiable space, so every first-countable topological gyrogroup is metrizable. Then R. Shen in [22] proved that every weakly first-countable paratopological left-loop is first-countable. M. Bao and F. Lin introduced the concept of strongly topological gyrogroups, and proved that every feathered strongly topological gyrogroup is paracompact, every  $T_0$  strongly topological gyrogroup is completely regular and every  $T_0$  strongly topological gyrogroup with a countable pseudocharacter is submetrizable, see [4, 5, 6].

In this paper, we mainly consider suitable sets for (strongly) topological gyrogroups. A subset  $S$  of a topological gyrogroup  $G$  is said to be a *suitable set* for  $G$  if (1)  $\overline{\langle S \rangle} = G$ , (2)  $S$  has the discrete topology, and (3)  $S \cup \{0\}$  is closed in  $G$ . We show that each countable Hausdorff topological gyrogroup has a suitable set, and each separable metrizable strongly topological gyrogroup has a suitable set, which generalizes some results for topological groups in [8, 9].

All spaces throughout this paper are supposed to be Hausdorff, unless the opposite is not stated. Let  $\mathbb{N}$  be the set of all positive integers and  $\omega$  the first infinite ordinal. Let  $X$  be a topological space, and let  $A$  be a subset of  $X$ . The *closure* of  $A$  in  $X$  is denoted by  $\overline{A}$ . For undefined notation and terminology, the reader may refer to [1, 11].

## 2. MOTIVATION AND PRELIMINARIES

In this section, we provide a motivation to study suitable sets in topological gyrogroups. Also we recall and introduce notions and notation used in the paper.

**Definition 2.1.** [2] A *groupoid* is a pair  $(G, \oplus)$ , where  $G$  is a nonempty set and  $\oplus$  is a binary operation on  $G$ . A function  $f$  from a groupoid  $(G_1, \oplus_1)$  to a groupoid  $(G_2, \oplus_2)$  is called a *groupoid homomorphism*, if  $f(x \oplus_1 y) = f(x) \oplus_2 f(y)$  for any elements  $x, y \in G_1$ . Furthermore, a bijective groupoid homomorphism from a groupoid  $(G, \oplus)$  to itself will be called a *groupoid automorphism*. We denote for a set of all automorphisms of a groupoid  $(G, \oplus)$  by  $\text{Aut}(G, \oplus)$ .

**Definition 2.2.** [26] A groupoid  $(G, \oplus)$  is called a *gyrogroup*, if its binary operation satisfies the following conditions.

(G1) There exists a unique identity element  $0 \in G$  such that  $0 \oplus a = a = a \oplus 0$  for all  $a \in G$ .

(G2) For each  $x \in G$ , there exists a unique inverse element  $\ominus x \in G$  such that  $\ominus x \oplus x = 0 = x \oplus (\ominus x)$ .

(G3) There exists a map  $\text{gyr} : G \times G \rightarrow \text{Aut}(G, \oplus)$ , such that  $x \oplus (y \oplus z) = (x \oplus y) \oplus \text{gyr}[x, y](z)$  for all  $z \in G$ .

(G4) For any  $x, y \in G$ ,  $\text{gyr}[x \oplus y, y] = \text{gyr}[x, y]$ .

**Definition 2.3.** [23] A nonempty subset  $H$  of a gyrogroup  $(G, \oplus)$  is called a *subgyrogroup* of  $G$  (denoted by  $H \leq G$ ), provided the following conditions hold.

(i) The restriction  $\oplus|_{H \times H}$  is a binary operation on  $H$ , i.e.  $(H, \oplus|_{H \times H})$  is a groupoid.

(ii) For any  $x, y \in H$ , the restriction of  $\text{gyr}[x, y]$  to  $H$ ,  $\text{gyr}[x, y]|_H : H \rightarrow \text{gyr}[x, y](H)$ , is a bijective homomorphism.

(iii)  $(H, \oplus|_{H \times H})$  is a gyrogroup.

A subgyrogroup  $H$  of  $G$  is said to be an  *$L$ -subgyrogroup* [23], denoted by  $H \leq_L G$ , if  $\text{gyr}[a, h](H) = H$  for all  $a \in G$  and  $h \in H$ .

**Definition 2.4.** [2] A triple  $(G, \tau, \oplus)$  is called a *topological gyrogroup*, provided the following conditions hold.

- (1)  $(G, \tau)$  is a topological space.
- (2)  $(G, \oplus)$  is a gyrogroup.
- (3) The binary operation  $\oplus : G \times G \rightarrow G$  is jointly continuous, where  $G \times G$  is endowed with the product topology and the inversion  $\ominus : G \rightarrow G, x \mapsto \ominus x$ , is continuous.

It is easy to see that each topological group is a topological gyrogroup  $(G, \tau, \oplus)$  provided we put  $\text{gyr}[x, y](z) = z$  all  $x, y, z \in G$ . A well-known example of a topological gyrogroup, which is not a topological group, is the following *Möbius topological gyrogroup*.

**Example 2.5.** [2] Let  $D$  be a open unit disk  $\{z \in \mathbb{C} : |z| < 1\}$  in the complex plane. Define a Möbius addition  $\oplus_M : D \times D \rightarrow D$  putting

$$a \oplus_M b = \frac{a + b}{1 + \bar{a}b} \text{ for all } a, b \in D.$$

Then  $(D, \oplus_M)$  is a gyrogroup with

$$\text{gyr}[a, b](c) = \frac{1 + \bar{a}b}{1 + \bar{a}b} c \text{ for any } a, b, c \in D.$$

But  $(D, \oplus_M)$  is not a group, because the operation  $\oplus_M$  is not associative. Indeed, it is easy to check that  $(1/2 \oplus_M i/2) \oplus_M (-1/2) \neq 1/2 \oplus_M (i/2 \oplus_M (-1/2))$ . If  $\tau$  the usual topology on  $D$  then  $(D, \tau, \oplus_M)$  is a topological gyrogroup.

**Definition 2.6.** [4] A topological gyrogroup  $G$  is a *strongly topological gyrogroup* if there exists a neighborhood base  $\mathcal{U}$  of 0 such that  $\text{gyr}[x, y](U) = U$  for each  $x, y \in G$  and  $U \in \mathcal{U}$ . In this case we shall say that  $G$  is a strongly topological gyrogroup with a neighborhood base  $\mathcal{U}$  of 0. Clearly, we may assume that  $U$  is symmetric for each  $U \in \mathcal{U}$ .

We claim that  $(D, \tau, \oplus_M)$  in Example 2.5 is a strongly topological gyrogroup [4]. Indeed, for any  $n \in \omega$ , let  $U_n = \{x \in D : |x| \leq \frac{1}{n}\}$ . Then,  $\mathcal{U} = \{U_n : n \in \omega\}$  is a neighborhood base of 0. Moreover, since  $1 + \bar{a}b = 1 + \bar{a}b$  for each  $a, b \in D$ , we have  $|\frac{1 + \bar{a}b}{1 + \bar{a}b}| = 1$ . Therefore, we see that  $\text{gyr}[x, y](U) \subset U$ , for any  $x, y \in D$  and each  $U \in \mathcal{U}$ . By [23, Proposition 2.6] it follows that  $\text{gyr}[x, y](U) = U$ .

Moreover, Möbius gyrogroups, Einstein gyrogroups, and Proper velocity gyrogroups, that were studied in [12, 13, 26], are all strongly topological gyrogroups, see [4].

**Definition 2.7.** [15] Let  $G$  be a topological gyrogroup and  $S$  a subset of  $G$ . Then  $S$  is said to be a *suitable set* for  $G$  if  $S$  is discrete in itself, generates a dense subgroup of  $G$ , and  $S \cup \{0\}$  is closed in  $G$ .

By the same notations of [9], let  $\mathcal{S}$  (resp.,  $\mathcal{S}_c$ ) be the class of topological gyrogroups having a suitable (resp., closed suitable) set. It turns out that very often the subset  $S$  of the group  $G$  has the stronger property to generate  $G$ , instead of generating just a dense subgroup of  $G$ . We denote by  $\mathcal{S}_g$  and  $\mathcal{S}_{cg}$  the corresponding subclasses of  $\mathcal{S}$  and  $\mathcal{S}_c$ , respectively.

The following proposition generalizes [8, Proposition 1.4].

**Proposition 2.8.** If a topological gyrogroup  $(G, \oplus)$  has a suitable set, then  $G$  is Hausdorff or  $|G| \leq 2$ .

*Proof.* Assume that  $G$  is not Hausdorff and  $|G| \geq 3$ . Let  $S$  be the suitable set for  $G$ . Since  $G$  is not Hausdorff and  $T_0$  and  $T_3$  are equivalent in topological gyrogroups by [2], for every  $g \in G$  a set  $\overline{\{g\}}$  contains a point  $h \neq g$ . By the assumption of  $|G| \geq 3$ , it follows that  $S \cup \{0\}$  has at least two points. Take an arbitrary point  $s \in S \setminus \{0\}$ . Since  $S$  is discrete in itself, we have  $S \cap \overline{\{s\}} = \{s\}$ . Further,  $S \cup \{0\}$  is closed in  $G$ , thus  $\overline{\{s\}} = \{s, 0\}$ . Therefore,  $\overline{\{0\}} = \{s, 0\}$ . It follows that  $S$  has at most two points,  $s$  and  $0$ . Moreover, since  $\overline{\{0\}}$  is a gyrogroup, it is clear that  $s \oplus s = 0$ . Then,  $s = \ominus s = 0$ , that is,  $G = S$ , this is a contradiction.  $\square$

Recall that given a space  $X$ , a *pseudocharacter*  $\psi(X)$  of  $x$  is the smallest infinite cardinal  $\kappa$  such that any point of  $X$  is an intersection of at most  $\kappa$  open subsets of  $X$  and *extent*  $e(X)$  is the supremum of cardinalities of closed discrete subspaces of  $X$ . Similarly to the proof of [9, Lemma 2.3], we can show the following

**Proposition 2.9.** A Hausdorff topological gyrogroup  $G$  which has a suitable set satisfies  $d(G) \leq e(G) \cdot \psi(G)$ .

*Proof.* We assume that  $A$  is a suitable set for  $G$ . If  $U$  is an open neighborhood of  $0$  in  $G$ , then  $A \setminus U$  is discrete and closed in  $G$ , which implies  $|A \setminus U| \leq e(G)$ . Pick a family  $\gamma$  of open sets in  $G$  such that  $\bigcap \gamma = \{0\}$  and  $|\gamma| = \psi(G)$ . Since  $A \setminus \{0\} \subset \bigcup \{A \setminus U : U \in \gamma\}$ , it follows that  $|A| \leq e(G) \cdot \psi(G)$ . The subgroup  $H = \langle A \rangle$  of  $G$  satisfies  $|H| \leq |A| \cdot \aleph_0$ . Since  $A$  is a suitable set and  $H$  is dense in  $G$ , we can conclude that

$$d(G) \leq |H| \leq |A| \cdot \aleph_0 \leq e(G) \cdot \psi(G).$$

$\square$

Therefore, it is natural to have the following result.

**Corollary 2.10.** A non-separable Lindelöf Hausdorff topological gyrogroup of countable pseudocharacter does not have a suitable set.

**Example 2.11.** There exists a non-separable Lindelöf Hausdorff topological gyrogroup  $G$  of countable pseudocharacter such that  $G$  does not have a suitable set and  $G$  is not a topological group.

*Proof.* Let  $D$  be the topological gyrogroup in Example 2.5, and let  $H$  be the Lindelöf non-separable topological group with countable pseudocharacter in (a) of [9, Theorem 2.4]. Then  $D$  has a suitable set by in the following Corollary 4.15 and  $H$  does not have any suitable set. Moreover, the product  $G = D \times H$  is a Lindelöf non-separable topological gyrogroup with countable pseudocharacter, hence it does not have any suitable set by Corollary 2.10. Clearly,  $G$  is not a topological group.  $\square$

In this paper we mainly consider the following question.

**Question 2.12.** If  $G$  belongs to some class  $\mathcal{C}$  of Hausdorff topological gyrogroups, does  $G$  have a suitable set?

### 3. COUNTABLE TOPOLOGICAL GYROGROUP WITH A SUITABLE SET

In this section, we study the suitable sets in the class  $\mathcal{C}$  of Hausdorff countable topological gyrogroups. We prove that every Hausdorff countable topological gyrogroup  $G$  has a closed discrete subset  $S$  such that  $\langle S \rangle = G$ . First, we need some lemmas.

Let  $G$  be a gyrogroup. Fix an  $n \in \mathbb{N}$ . For any  $x_1, \dots, x_n \in G$  and  $\varepsilon_1, \dots, \varepsilon_n \in \{-1, 1\}$ , denote by  $R[\varepsilon_1 x_1, \dots, \varepsilon_n x_n]$  the set of all elements which is added some brackets in the summand  $\varepsilon_1 x_1 \oplus \dots \oplus \varepsilon_n x_n$  such that the summand belongs to  $G$ , where

$$\varepsilon_i x_i = \begin{cases} x_i, & \varepsilon_i = 1; \\ \ominus x_i, & \varepsilon_i = -1. \end{cases}$$

Clearly,  $R[\varepsilon_1 x_1, \dots, \varepsilon_n x_n]$  is a countable set, so enumerate  $R[\varepsilon_1 x_1, \dots, \varepsilon_n x_n]$  as

$$\{f_m(\varepsilon_1 x_1, \dots, \varepsilon_n x_n) : m \in \mathbb{N}\}.$$

If  $A_1, \dots, A_n \subset G$ , then we denote  $R[\varepsilon_1 A_1, \dots, \varepsilon_n A_n]$  and  $f_m(\varepsilon_1 A_1, \dots, \varepsilon_n A_n)$  as the sets

$$\bigcup_{x_1 \in A_1, \dots, x_n \in A_n} R[\varepsilon_1 x_1, \dots, \varepsilon_n x_n] \text{ and } \bigcup_{x_1 \in A_1, \dots, x_n \in A_n} f_m(\varepsilon_1 x_1, \dots, \varepsilon_n x_n),$$

respectively.

In the class of topological gyrogroups, since the multiplication is jointly continuous and the inverse is continuous, it is easy to prove the following lemma.

**Lemma 3.1.** *Let  $a_1, a_2, \dots, a_n$  be points of a topological gyrogroup  $G$ , and let  $V$  be a neighborhood of the point  $f_m(\varepsilon_1 a_1, \dots, \varepsilon_n a_n)$ . Then there exists neighborhoods  $U_1, \dots, U_n$  of  $a_1, \dots, a_n$  in  $G$  respectively such that  $f_m(\varepsilon_1 U_1, \dots, \varepsilon_n U_n) \subset V$ .*

A topological space  $X$  is *zero-dimensional* if it has a base consisting of clopen subsets.

**Lemma 3.2.** *Let  $G$  be a nondiscrete Hausdorff topological gyrogroup and  $U$  a nonempty open subset which generates  $G$ . Then every point  $x \in U$  has an open neighborhood  $V_x$  of  $x$  such that  $V_x \subset U$  and  $\langle U \setminus \overline{V_x} \rangle = G$ . In particular, if  $G$  is zero-dimensional, then  $V_x$  can be chosen to be clopen in  $G$ .*

*Proof.* Let  $U$  be a nonempty open subset which generates  $G$ . Take an arbitrary point  $x \in U$ . Since  $G$  is not discrete, it is obvious that  $U \setminus \{x\}$  is dense in  $U$ , then it follows that  $\langle U \setminus \{x\} \rangle$  is dense in  $\langle U \rangle = G$ . Moreover, since  $U \setminus \{x\}$  is open in  $G$  and every open subgyrogroup is closed in  $G$  by [2, Proposition 7], we can conclude that  $\langle U \setminus \{x\} \rangle$  is open and closed in  $G$ . Therefore,  $\langle U \setminus \{x\} \rangle = G$ .

Since  $x \in \langle U \setminus \{x\} \rangle$ , there exist  $y_1, y_2, \dots, y_n \in U \setminus \{x\}$ ,  $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_n \in \{1, -1\}$  and  $m \in \mathbb{N}$  such that  $x = f_m(\varepsilon_1 y_1, \dots, \varepsilon_n y_n)$ , where

$$\varepsilon_i y_i = \begin{cases} y_i, & \varepsilon_i = 1; \\ \ominus y_i, & \varepsilon_i = -1. \end{cases}$$

Because each  $y_i \neq x$ , we can find an open neighborhood  $O$  of  $x$  such that  $y_i \notin \overline{O} \subset U$ , for  $i = 1, \dots, n$ . Then for each  $i \in \{1, \dots, n\}$ , there is an open neighborhood  $O_i$  of  $y_i$  such that  $O_i \subset U$ ,  $O \cap O_i = \emptyset$ , and  $f_m(\varepsilon_1 O_1, \dots, \varepsilon_n O_n) \subset O$  by Lemma 3.1. Put  $W = f_m(\varepsilon_1 O_1, \dots, \varepsilon_n O_n)$ . Then  $W$  is an open neighborhood of  $x$ . By the regularity of  $G$ , there exists an open neighborhood  $\overline{V_x}$  of  $x$  such that  $\overline{V_x} \subset W \subset O$ . Therefore,  $O_i \subset U \setminus O \subset U \setminus \overline{V_x}$ , for  $i = 1, 2, \dots, n$ . So,  $\overline{V_x} \subset W \subset \langle U \setminus \overline{V_x} \rangle$ . Thus  $\langle U \setminus \overline{V_x} \rangle = \langle U \rangle = G$ .

It is obvious that the last statement of this lemma.  $\square$

Now we can prove our main theorem in this section.

**Theorem 3.3.** *Every countable Hausdorff topological gyrogroup  $G$  belongs to  $\mathcal{S}_{cg}$ .*

*Proof.* If  $G$  is finitely generated or discrete, then the theorem is clear. Therefore, we may suppose that  $G$  is neither finitely generated nor discrete. Enumerate  $G$  as  $\{g_n : n < \omega\}$ .

It suffice to find a subset  $S$  in  $G$  and an open neighborhood  $U_n$  of  $g_n$ , for each  $n < \omega$  satisfying  $\langle S \rangle = G$  and  $U_n \cap S$  is finite.

Next we will by induction to find a clopen set  $V_n$  in  $G$  and a finite set  $S_n \subset G$  for each  $n < \omega$  so that the following conditions hold:

- (i)  $g_n \in \bigcup_{i=0}^n V_i$  for each  $n \in \omega$ ;
- (ii)  $G = \langle G \setminus (\bigcup_{i=0}^n V_i) \rangle$  for each  $n \in \omega$ ;
- (iii) for each  $n > 0$ ,  $V_n \subset G \setminus (\bigcup_{i=0}^{n-1} V_i)$ ;
- (iv)  $V_i \cap S_n = \emptyset$ , for  $i < n$ ; and
- (v)  $g_n \in \langle \bigcup_{i=0}^n S_i \rangle$  for each  $n \in \omega$ .

Then set  $U_n = \bigcup_{i=0}^n V_i$  for each  $n \in \omega$ , and put  $S = \bigcup_{n < \omega} S_n$ . Clearly,  $\langle S \rangle = G$  and  $U_n \cap S$  is finite for each  $n \in \omega$ .

Therefore, it suffices to construct  $S_n$  and  $V_n$  inductively as follows.

Set  $S_0 = \{g_0\}$ . Since every Hausdorff topological gyrogroup is regular and every countable non-empty regular space is zero-dimensional [11, Corollary 6.2.8], it follows that the countable topological gyrogroup  $G$  is zero-dimensional. By Lemma 3.2, there exists a clopen neighborhood  $V_0$  of  $g_0$  such that  $G = \langle G \setminus V_0 \rangle$ .

Assume that the finite sets  $S_0, S_1, \dots, S_k$  and clopen sets  $V_0, V_1, \dots, V_k$  have been defined satisfying the above properties (i)-(v). Clearly, if  $g_{k+1} \in \langle \bigcup_{i=0}^k S_i \rangle$ , then set  $S_{k+1} = \emptyset$ . If  $g_{k+1} \notin \langle \bigcup_{i=0}^k S_i \rangle$ , then it follows from (ii) that there exist

$$y_1, y_2, \dots, y_m \in G \setminus \left( \bigcup_{i=0}^k V_i \right), \varepsilon_1, \varepsilon_2, \dots, \varepsilon_m \in \{1, -1\}$$

and  $n \in \mathbb{N}$  such that

$$g_{k+1} = f_n(\varepsilon_1 y_1, \dots, \varepsilon_m y_m).$$

Set  $S_{k+1} = \{y_1, y_2, \dots, y_m\}$ . Thus both (iv) and (v) are satisfied.

Obviously, if  $g_{k+1} \in \bigcup_{i=0}^k V_i$ , then put  $V_{k+1} = \emptyset$ . If  $g_{k+1} \notin \bigcup_{i=0}^k V_i$ , then it follows from Lemma 3.2 that there exists a clopen neighborhood  $V_{k+1}$  of  $g_{k+1}$  such that  $V_{k+1} \subset G \setminus (\bigcup_{i=0}^k V_i)$  and  $G = \langle G \setminus (\bigcup_{i=0}^{k+1} S_i) \rangle$ . Then (i)-(iii) are all satisfied.

Therefore, the sets  $S_n$  and  $V_n$  are defined for all  $n$  with the required properties.  $\square$

**Corollary 3.4.** [8] *Every countable Hausdorff topological group  $G$  belongs to  $\mathcal{S}_{cg}$ .*

#### 4. A STRONGLY TOPOLOGICAL GYROGROUP WITH A SUITABLE SET

In this section, we mainly prove that every separable metrizable strongly topological gyrogroup has a suitable set. First, we need some lemmas.

**Lemma 4.1.** *Suppose that  $(G, \tau, \oplus)$  is a strongly topological gyrogroup with a symmetric neighborhood base  $\mathcal{U}$  at 0. Suppose further that  $U, V, W$  are all open neighborhoods of 0 such that  $V \oplus V \subset W$ ,  $W \oplus W \subset U$  and  $V, W \in \mathcal{U}$ . If a subset  $A$  of  $G$  is  $U$ -disjoint (that is, if  $b \notin a \oplus U$ , for any distinct  $a, b \in A$ ), then for each  $x \in G$  the set  $x \oplus V$  intersects at most one of the element of the family  $\{a \oplus V : a \in A\}$ . In particular, the family of open sets  $\{a \oplus V : a \in A\}$  is discrete in  $G$ .*

*Proof.* We need to show that, for every  $x \in G$ , the open neighborhood  $x \oplus V$  of  $x$  intersects at most one element of the family  $\{a \oplus V : a \in A\}$ . We assume the contrary that, for some  $x \in G$ , there exist distinct elements  $a, b \in A$  such that  $(x \oplus V) \cap (a \oplus V) \neq \emptyset$  and  $(x \oplus V) \cap (b \oplus V) \neq \emptyset$ . We show that  $b \in a \oplus U$  as follows.

Since  $(x \oplus V) \cap (a \oplus V) \neq \emptyset$ , we have that there exist  $v_1, v_2 \in V$  such that  $x \oplus v_1 = a \oplus v_2$ . Then,  $a = (a \oplus v_2) \oplus \text{gyr}[a, v_2](\ominus v_2) = (x \oplus v_1) \oplus \text{gyr}[a, v_2](\ominus v_2)$ . Therefore,

$$\begin{aligned} a &\in (x \oplus v_1) \oplus \text{gyr}[a, v_2](V) \\ &= (x \oplus v_1) \oplus V \\ &= x \oplus (v_1 \oplus \text{gyr}[v_1, x](V)) \\ &= x \oplus (v_1 \oplus V) \\ &\subset x \oplus (V \oplus V) \\ &\subset x \oplus W. \end{aligned}$$

Thus,  $a \in x \oplus W$ . By the same method, we also have  $b \in x \oplus W$ .

Therefore, there exists  $w_1 \in W$  such that  $a = x \oplus w_1$ . Then,

$$x = a \oplus \text{gyr}[x, w_1](\ominus w_1) \in a \oplus \text{gyr}[x, w_1](W) = a \oplus W.$$

Hence,

$$\begin{aligned} b &\in (a \oplus W) \oplus W \\ &= a \oplus (W \oplus \text{gyr}[W, a](W)) \\ &= a \oplus (W \oplus W) \\ &\subset a \oplus U. \end{aligned}$$

□

Let  $G$  be a topological gyrogroup. For  $\kappa$  an infinite cardinal, the topological gyrogroup  $G$  is said to be *left  $\kappa$ -totally bounded* if for every nonempty open subset  $U$  of  $G$  there is  $F \subset G$  such that  $|F| < \kappa$  and  $G = F \oplus U$ . We denote  $lb(G)$  by the least cardinal  $\kappa \geq \omega$  such that  $G$  is left  $\kappa$ -totally bounded. Each left  $\omega$ -totally bounded topological gyrogroup is also called *left precompact*.

**Lemma 4.2.** *Let  $G$  be a strongly topological gyrogroup with  $lb(G) = \kappa$ . If  $\tau < \kappa$ , then there exist an open neighborhood  $V$  of 0 and a subset  $\{p_\alpha : \alpha < \tau\}$  such that for each  $p \in G$  the set  $p \oplus V$  intersects at most one of the elements of the family  $\{p_\alpha \oplus V : \alpha < \tau\}$ .*

*Proof.* Since  $lb(G) = \kappa$  and  $\tau < \kappa$ , it follows that there exists a nonempty open neighborhood  $U$  of 0 in  $G$  such that no  $F \subset G$  with  $|F| \leq \tau$  satisfies  $G = F \oplus U$ . By induction, it is easy to find a set  $\{p_\alpha : \alpha < \tau\}$  such that each  $p_\alpha$  satisfies  $p_\alpha \notin \bigcup_{\beta < \alpha} (p_\beta \oplus U)$ . Then, from Lemma 4.1 we can find a nonempty open neighborhood  $V$  of 0 in  $G$  such that for each  $p \in G$  the set  $p \oplus V$  intersects at most one of the elements of the family  $\{p_\alpha \oplus V : \alpha < \tau\}$ . □

The strongly topological gyrogroup  $G$  in Example 2.5 is left precompact and non-pseudocompact. However, the following result shows that each pseudocompact strongly topological gyrogroup is left precompact.

**Theorem 4.3.** *Suppose that  $(G, \tau, \oplus)$  is a strongly topological gyrogroup with a symmetric open neighborhood base  $\mathcal{U}$  at 0. If  $G$  is pseudocompact, then it will be left precompact.*

*Proof.* Let  $U$  be an arbitrary symmetric open neighborhood of 0 in  $G$  and  $V, W \in \mathcal{U}$  such that  $V \oplus V \subset W$  and  $W \oplus W \subset U$ . Let

$$\mathcal{F} = \{A \subset G : (b \oplus V) \cap (a \oplus V) = \emptyset, \text{ for any distinct } a, b \in A\}.$$

Define  $\leq$  in  $G$  such that  $A_1 \leq A_2$  if and only if  $A_1 \subset A_2$ , for any  $A_1, A_2 \in \mathcal{F}$ . Then,  $(\mathcal{F}, \leq)$  is a poset and the union of any chain of  $V$ -disjoint sets is again a  $V$ -disjoint

set. Therefore, it follows from Zorn's Lemma that there exists a maximal element  $A$  in  $\mathcal{F}$  so that  $\{a \oplus V : a \in A\}$  is a disjoint family of non-empty open sets in  $G$ . By Lemma 4.1, the family of open sets  $\{a \oplus V : a \in A\}$  is discrete in  $G$ . What's more,  $G$  is pseudocompact, we have that  $A$  is finite. Finally, we show that  $A \oplus U = G$  as follows.

Take an arbitrary  $x \in G$ . If  $x \notin A$ , then it follows from the maximality of  $A$  that there exists  $a \in A$  such that  $(x \oplus V) \cap (a \oplus V) \neq \emptyset$ . Then, there exist  $v_1, v_2 \in V$  such that  $x \oplus v_1 = a \oplus v_2$ . By the right cancellation law, we have that

$$\begin{aligned}
x &= (x \oplus v_1) \oplus \text{gyr}[x, v_1](\ominus v_1) \\
&= (a \oplus v_2) \oplus \text{gyr}[x, v_1](\ominus v_1) \\
&\in (a \oplus v_2) \oplus \text{gyr}[x, v_1](V) \\
&= (a \oplus v_2) \oplus V \\
&= a \oplus (v_2 \oplus \text{gyr}[v_2, a](V)) \\
&= a \oplus (v_2 \oplus V) \\
&\subset a \oplus (V \oplus V) \\
&\subset a \oplus U.
\end{aligned}$$

Therefore,  $A \oplus U = G$ . □

**Theorem 4.4.** *Suppose that  $(G, \tau, \oplus)$  is a strongly topological gyrogroup with a symmetric open neighborhood base  $\mathcal{U}$  at 0, and that  $H$  is an open  $L$ -subgyrogroup of  $G$ . If  $H$  has a suitable set, then  $G$  has a suitable set. If  $H$  has a closed suitable set, then  $G$  has a closed suitable set.*

*Proof.* Let  $S$  be a suitable set for  $H$ . Since  $H$  is a  $L$ -subgyrogroup, two distinct cosets of  $H$  are disjoint. Then let  $A$  select one point from each coset of  $H$  in  $G$  such that  $0 \notin A$  and  $|A \cap (g \oplus H)| = 1$  for each  $x \in G$ . We claim that  $S \cup A$  is suitable for  $G$ .

Indeed,  $S \cup \{0\}$  and  $H$  are all closed in  $G$ , thus  $S \cup A$  is discrete in  $G \setminus \{0\}$ , then there is at most an accumulation point 0 since  $S \cup A \cup \{0\}$  is closed in  $G$ . Now it suffices to prove that  $\langle S \cup A \rangle$  is dense in  $G$ . Since  $\langle S \rangle$  is dense in  $H$ , the subgyrogroup  $\langle S \cup A \rangle$  is dense in  $G$ . If  $S$  is closed in  $H$ , then  $S \cup A$  is closed in  $G$ . □

However, the following question is open.

**Question 4.5.** *Suppose that  $(G, \tau, \oplus)$  is a strongly topological gyrogroup, and that  $H$  is an open subgyrogroup of  $G$  with a suitable set. Does  $H$  have a suitable set?*

The following lemma gives a partial answer to Question 4.5.

**Lemma 4.6.** *Suppose that  $(G, \tau, \oplus)$  is a separable strongly topological gyrogroup with a symmetric open neighborhood base  $\mathcal{U}$  at 0,  $H$  is an open subgyrogroup of  $G$ . If  $H$  has a suitable set, then  $G$  has a suitable set. If  $H$  has a closed suitable set, then  $G$  has a closed suitable set.*

*Proof.* Let  $S$  be a suitable set for  $H$ . Since  $G$  is separable, there exists a countable subset  $A = \{g_n : n \in \omega\}$  of  $G$  such that  $g_0 = 0$  and  $\overline{A} = G$ , then  $A \oplus H = G$  since  $H$  is open in  $G$ . Then, by induction on  $n$ , we can choose a subset  $B$  of  $A$  satisfies the following conditions:

- (i)  $B$  is closed discrete;
- (ii)  $\overline{\langle B \cup S \rangle} = G$ ;
- (iii)  $B \cap (g \oplus H) = \{g\}$ .

Indeed, take  $g_0 = \{0\}$ . If  $H = G$ , then let  $B = \{0\}$ ; otherwise,  $G \setminus H \neq \emptyset$ , since  $G \setminus H$  is open, there exists a minimum  $n_1 \in \mathbb{N}$  such that  $g_{n_1} \in (g_{n_1} \oplus H) \setminus H$  and  $g_i \in H$  for any  $i < n_1$ . Assume we have defined the points  $g_0, g_{n_1}, \dots, g_{n_k}$  such that  $g_{n_i} \in (g_{n_i} \oplus H) \setminus \bigcup_{j < i} (g_{n_j} \oplus H)$  for each  $i \leq k$  and  $g_j \in \bigcup_{i=0}^{k-1} (g_{n_i} \oplus H)$  for any  $n_m \leq j < n_{m+1}$  and  $m \leq k-1$ . If  $\bigcup_{i=0}^k (g_{n_i} \oplus H) = G$ , let  $B = \{g_{n_i} : i \leq k\}$ ; otherwise, the set  $G \setminus \bigcup_{i \leq k} (g_{n_i} \oplus H)$  is a nonempty open subset of  $G$ , then there exists a minimum  $n_{k+1} \in \mathbb{N}$  such that  $g_{n_{k+1}} \in (g_{n_{k+1}} \oplus H) \setminus \bigcup_{i \leq k} (g_{n_i} \oplus H)$  and  $g_j \in \bigcup_{i \leq k} (g_{n_i} \oplus H)$  for each  $j \leq n_{k+1}$ . If there exists  $N \in \mathbb{N}$  such that  $\bigcup_{i \leq N} (g_{n_i} \oplus H) = G$ , then  $B = \{g_{n_i} : i \leq N\}$  is a finite set; otherwise, put  $B = \{g_{n_i} : i \in \omega\}$ . By our construction of  $B$ , it is easy to see that  $B$  satisfies the conditions (i)-(iii).

By (ii),  $\langle S \cup B \rangle$  is dense in  $G$ . Moreover,  $S \cup \{0\}$  and  $H$  are all closed in  $G$ , thus  $(S \cup A) \setminus \{0\}$  is discrete in  $G \setminus \{0\}$ , then there is at most an accumulation point 0 since  $S \cup A \cup \{0\}$  is closed in  $G$ . If  $S$  is closed in  $H$ , then  $S \cup A$  is closed in  $G$ .  $\square$

**Lemma 4.7.** *Suppose that  $(G, \tau, \oplus)$  is a strongly topological gyrogroup with a symmetric open neighborhood base  $\mathcal{U}$  at 0,  $B$  is a left precompact subset of  $G$  and  $S$  is dense in  $B$ . Then, for every neighborhood  $U$  of 0 in  $G$ , there is a finite set  $K \subset S$  such that  $B \subset K \oplus U$ .*

*Proof.* We assume that  $U$  is an arbitrary neighborhood of 0 in  $G$  and  $V \in \mathcal{U}$  such that  $V \oplus V \subset U$ . Since  $B$  is left precompact in  $G$ , there exists a finite set  $F$  in  $G$  such that  $B \subset F \oplus V$ . Take an arbitrary  $x \in F$  such that  $B \cap (x \oplus V) \neq \emptyset$ . Then  $S \cap (x \oplus V) \neq \emptyset$  and we pick a point  $y_x \in S \cap (x \oplus V)$ . Then the finite set

$$K_1 = \{y_x : x \in F \text{ and } B \cap (x \oplus V) \neq \emptyset\}$$

is contained in  $S$ . We claim  $B \subset K_1 \oplus U$ .

Indeed, if  $b \in B$ , then there exists  $x \in F$  such that  $b \in x \oplus V$ , so  $b \in B \cap (x \oplus V) \neq \emptyset$ . Therefore,  $y_x \in x \oplus V$ . We can find  $v_1 \in V$  such that  $y_x = x \oplus v_1$ . Then

$$\begin{aligned} x &= (x \oplus v_1) \oplus \text{gyr}[x, v_1](\ominus v_1) \\ &= y_x \oplus \text{gyr}[x, v_1](\ominus v_1) \\ &\in y_x \oplus \text{gyr}[x, v_1](V) \\ &= y_x \oplus V. \end{aligned}$$

Thus,

$$\begin{aligned} b &\in x \oplus V \\ &\subset (y_x \oplus V) \oplus V \\ &= y_x \oplus (V \oplus \text{gyr}[V, y_x](V)) \\ &= y_x \oplus (V \oplus V) \\ &\subset y_x \oplus U \\ &\subset K_1 \oplus U. \end{aligned}$$

$\square$

**Lemma 4.8.** *Every subgyrogroup  $H$  of a left precompact strongly topological gyrogroup  $G$  is left precompact.*

*Proof.* Take an arbitrary open neighborhood  $U$  of 0 in  $H$ , then there is an open neighborhood  $V$  of 0 in  $G$  such that  $V \cap H = U$ . Since  $H$  is a left precompact subset of  $G$ , by Lemma 4.7, we can find a finite set  $F \subset H$  such that  $H \subset F \oplus V$ . Therefore, for every  $h \in H$ , there exist  $f \in F$  and  $v \in V$  such that  $h = f \oplus v$ . Thus,

$v = (\ominus f) \oplus h \in H \oplus H \subset H$ . Then,  $v \in V \cap H = U$ . It follows that  $H \subset F \oplus U$ , that is,  $H = F \oplus U$ .  $\square$

By Theorem 4.3 and Lemma 4.8, we have the following corollary.

**Corollary 4.9.** *Every subgyrogroup  $H$  of a pseudocompact strongly topological gyrogroup  $G$  is left precompact.*

**Lemma 4.10.** *Suppose that  $(G, \tau, \oplus)$  is a strongly topological gyrogroup with a symmetric open neighborhood base  $\mathcal{U}$  at 0, and suppose that  $G$  is non-pseudocompact left precompact with a countable dense subgyrogroup  $P$ . Then there exists a subset  $L \subset P$  such that  $L$  is closed discrete in  $G$  and  $\langle L \rangle = P$ . In particular,  $L$  is suitable for  $G$ .*

*Proof.* Since  $G$  is not pseudocompact, we can fix a sequence  $\{U_n : n \in \omega\}$  of non-empty open subsets of  $G$  such that  $U_n \in \mathcal{U}$ ,  $\overline{U_{n+1}} \subset U_n$  for each  $n \in \omega$  and  $\bigcap \{U_n : n \in \omega\} = \emptyset$ . Let  $\{x_n : n \in \omega\}$  be an enumeration of elements of  $P$ .

We construct an increasing sequence  $\{L_k : k \in \omega\}$  of finite subsets of  $P$  by induction which satisfies the following conditions:

- (1)  $x_k \in \langle L_k \rangle$ ;
- (2)  $L_{k+1} \setminus L_k \subset U_k$ ;
- (3)  $G = \langle L_k \rangle \oplus U_k$ .

Since the subgyrogroup  $P$  is dense in  $G$ , it follows from [2, Lemma 9] that  $G = \overline{P} \subset P \oplus U_0$ . So  $G = P \oplus U_0$ . Since  $G$  is left precompact, it follows from Lemma 4.7 that we can find a finite subset  $K_0$  of  $P$  such that  $K_0 \oplus U_0 = G$ . Therefore, for any  $x_0 \in G$ , there exist  $a_0 \in K_0, u_0 \in U_0$  such that  $x_0 = a_0 \oplus u_0$ . Then  $u_0 = (\ominus a_0) \oplus x_0 \in P$  and let  $L_0 = K_0 \cup \{u_0\}$ .

We assume that for some  $n \in \omega$  we have defined an increasing sequence  $L_0, \dots, L_n$  of finite subsets of  $P$  which satisfies (1) and (3) for each  $k \leq n$  and (2) for every  $k < n$ . Since  $P$  is dense in  $G$ , it is clear that  $\langle U_n \cap P \rangle$  is dense in the gyrogroup  $G_n = \langle U_n \rangle$ . Thus,  $\langle U_n \cap P \rangle \oplus U_{n+1} = G_n$ . It follows from Lemma 4.8 that  $G_n$  is left precompact, hence we can find a finite subset  $F_{n+1}$  of  $\langle U_n \cap P \rangle$  such that  $F_{n+1} \oplus U_{n+1} = G_n$ . Clearly, we can find a finite subset  $K_{n+1}$  of  $U_n \cap P$  with  $F_{n+1} \subset \langle K_{n+1} \rangle$ , so  $\langle K_{n+1} \rangle \oplus U_{n+1} = G_n$ . Let  $L'_{n+1} = L_n \cup K_{n+1}$ . By (3), we have

$$\begin{aligned}
G &= \langle L_n \rangle \oplus U_n \\
&\subset \langle L'_{n+1} \rangle \oplus G_n \\
&= \langle L'_{n+1} \rangle \oplus (\langle K_{n+1} \rangle \oplus U_{n+1}) \\
&= (\langle L'_{n+1} \rangle \oplus \langle K_{n+1} \rangle) \oplus \text{gyr}[\langle L'_{n+1} \rangle, \langle K_{n+1} \rangle](U_{n+1}) \\
&= (\langle L'_{n+1} \rangle \oplus \langle K_{n+1} \rangle) \oplus U_{n+1} \\
&= \langle L'_{n+1} \rangle \oplus U_{n+1}.
\end{aligned}$$

Therefore, there exist  $a_{n+1} \in \langle L'_{n+1} \rangle, u_{n+1} \in U_{n+1}$  such that  $x_{n+1} = a_{n+1} \oplus u_{n+1}$ . Since  $a_{n+1} \in \langle L'_{n+1} \rangle \subset P$  and  $x_{n+1} \in P$ , it follows that  $u_{n+1} = (\ominus a_{n+1}) \oplus x_{n+1} \in P$ . Then let  $L_{n+1} = L'_{n+1} \cup \{u_{n+1}\}$ . It is clear that  $L_{n+1}$  is a finite subset of  $P$  and  $L_n \subset L_{n+1}$ . At the same time,  $\langle L_{n+1} \rangle \oplus U_{n+1} = G$ . Moreover,  $L_{n+1} \setminus L_n \subset K_{n+1} \cup \{u_{n+1}\} \subset U_n$ . Therefore, we complete the construction.

Finally, set  $L = \bigcup \{L_n : n \in \omega\}$ . It follows from (2) that  $L \setminus \overline{U_k} \subset L_k$  is a finite set for each  $k \in \omega$ . Then,  $\bigcap \{\overline{U_n} : n \in \omega\}$  implies that  $L$  is a closed discrete subset of  $G$ . Moreover, (1) guarantees that  $\langle L \rangle = P$ .  $\square$

Now we can prove one of main results in this section.

**Theorem 4.11.** *Suppose that  $(G, \tau, \oplus)$  is a non-pseudocompact strongly topological gyrogroup with a symmetric open neighborhood base  $\mathcal{U}$  at 0. If  $G$  is separable, then  $G \in \mathcal{S}_c$ .*

*Proof.* It follows from [5] that  $G$  is Tychonoff. We divide the proof into two cases:

**Case 1:**  $G$  is not left precompact.

Then  $G$  has a neighborhood  $U$  of 0 such that  $F \oplus U \neq G$  for any finite  $F \subset G$ . By Lemma 4.1, we need to take  $V, W \in \mathcal{U}$  such that  $V \oplus V \subset W$  and  $W \oplus W \subset U$ , then there exists a subset  $A = \{a_n : n \in \omega\} \subset G$  with  $a_i \neq a_j$  if  $i \neq j$  and the family  $\gamma = \{a_n \oplus V : n \in \omega\}$  is discrete in  $G$ . Let  $B$  a countable dense subset of  $G$ , and set  $B_V = B \cap V = \{b_n : n \in \omega\}$ . We prove that  $S = A \cup (\bigcup_{n \in \mathbb{N}} a_n \oplus b_n)$  is a suitable set of open subgyrogroup  $G_1 = \langle V \cup A \rangle$ .

Since  $\langle B_V \rangle$  is dense in  $\langle V \rangle$  and  $B_V \subset \langle S \rangle$ , we have that  $\langle S \rangle$  is dense in  $G_1$ . For every  $g \in G$ , there exists a neighborhood  $O$  of 0 in  $G$  such that  $(g \oplus O) \cap S \subset \{a_n, a_n \oplus b_n\}$ . Therefore,  $S$  is closed and discrete and hence it is a suitable set of  $G_1$ . Then since  $G_1$  is an open subgyrogroup of  $G$ , it follows from Lemma 4.6 that  $G$  has a closed suitable set.

**Case 2:**  $G$  is left precompact.

Since  $G$  is non-pseudocompact, we can choose a discrete family  $\gamma = \{U_n : n \in \omega\}$  of non-empty open subsets of  $G$ . Let  $B = \{d_n : n \in \mathbb{N}\}$  be a countable dense subset of  $G$ .

Since  $G$  is precompact, for every  $n \in \omega$ , there exists a finite subset  $A_n = \{a(n, i) : 1 \leq i \leq m_n\}$  of  $G$  such that  $A_n \oplus U_n = G$ . Fix an  $n \in \omega$  and define  $H_n^i = \{d_n\} \cap (a(n, i) \oplus U_n)$  for each  $i \leq m_n$ . Then  $H_n = \bigcup \{H_n^i : 1 \leq i \leq m_n\}$ .

The set  $T_n = \bigcup \{(\ominus a(n, i)) \oplus H_n^i : 1 \leq i \leq m_n\}$  is closed and discrete in  $G$  and lies in  $U_n$ . Since the family  $\gamma$  is discrete, the set  $T = \bigcup \{T_n : n \in \omega\}$  is closed and discrete in  $G$ . Let  $A = \bigcup \{A_n : n \in \omega\}$ . For every  $n \in \omega$ , choose a point  $y_n \in U_n$  such that  $d_n \in A_n \oplus y_n$  and denote by  $G_2$  the closure of  $P = \langle A \cup \{y_n : n \in \omega\} \rangle$  in  $G$ . The gyrogroup  $G_2$  is closed and left precompact by Lemma 4.8. Moreover, for each  $n \in \omega$ ,  $G_2 \cap U_n \neq \emptyset$ , so  $G_2$  is not pseudocompact.

It follows from Lemma 4.10 that there is a closed discrete subset  $L$  of  $G_2$  such that  $\langle L \rangle = P$ . We find that  $T \cup L$  is closed and discrete in  $G$  and  $P \subset \langle T \cup L \rangle \supset \langle T \cup A \rangle \supset B$ . Hence  $\langle T \cup L \rangle$  is dense in  $G$ . Therefore,  $T \cup L$  is a closed suitable set for  $G$ .  $\square$

**Lemma 4.12.** *Let  $G$  be a compact metrizable strongly topological gyrogroup. Then  $G$  has a closed suitable set.*

*Proof.* Since  $G$  is compact metrizable, it is separable, hence there exists a countable dense subgroup  $P$ . Let  $P = \{x_n : n \in \omega\}$  be an enumeration of  $P$ . Moreover, we can choose a decreasing sequence  $\{U_n : n \in \omega\}$  of open neighborhoods of the identity 0 in  $G$  satisfying the following conditions:

- (1)  $U_{n+1} \oplus U_{n+1} \subset U_n$  for each  $n \in \omega$ ;
- (2)  $\bigcap_{n \in \omega} U_n = \{0\}$ .

By the same construction of Lemma 4.10, we can find an increasing sequence  $\{L_k : k \in \omega\}$  of finite subsets of  $P$  by induction which satisfies the following conditions:

- (1)  $x_k \in \langle L_k \rangle$ ;
- (2)  $L_{k+1} \setminus L_k \subset U_k$ ;
- (3)  $G = \langle L_k \rangle \oplus U_k$ .

By a similar proof of Lemma 4.10, we can find a closed discrete subset  $L$  for  $P$ . Then  $L$  is a closed suitable set for  $G$  since  $P$  is dense in  $G$ .  $\square$

A space  $X$  is *paracompact*, if each its open cover has a locally finite open refinement. A space  $X$  is *submetrizable*, if there exists a continuous injective map of  $X$  to a metrizable space.

**Corollary 4.13.** *Suppose that  $(G, \tau, \oplus)$  is a separable left precompact Hausdorff strongly topological gyrogroup of countable pseudocharacter with a symmetric open neighborhood base  $\mathcal{U}$  at 0. If  $P$  is a countable dense subgyrogroup of  $G$ , then there exists a discrete subset  $L$  of  $P$  such that  $L$  is closed in  $G \setminus \{0\}$  and  $P = \langle L \rangle$ . So  $L$  is a suitable set for both  $P$  and  $G$ .*

*Proof.* If  $G$  is not pseudocompact, then it follows from Lemma 4.10 that the conclusion holds. Assume that  $G$  is pseudocompact, then from [4, 5] that each strongly topological gyrogroup of countable pseudocharacter is paracompact and submetrizable, hence it is compact and metrizable, thus  $G$  has a closed suitable set by Lemma 4.12.  $\square$

A space  $X$  is a  $\sigma$ -space if it has a  $\sigma$ -locally finite network.

**Corollary 4.14.** *Suppose that  $(G, \tau, \oplus)$  is a strongly topological gyrogroup. If  $G$  is a separable  $\sigma$ -space then  $G$  has a suitable set.*

*Proof.* Since each  $\sigma$ -space has a countable pseudocharacter,  $G$  has countable pseudocharacter. If  $G$  is not pseudocompact, the conclusion holds from Theorem 4.11. From [4, 5], each strongly topological gyrogroup of countable pseudocharacter is paracompact and submetrizable, hence it is compact and metrizable, thus separable precompact, so we can apply Lemma 4.13 to conclude that  $G$  has a suitable set.  $\square$

**Corollary 4.15.** *Suppose that  $(G, \tau, \oplus)$  is a strongly topological gyrogroup. If  $G$  is a separable metrizable space, then  $G$  has a suitable set.*

We now close our paper with the following three questions.

**Question 4.16.** *Suppose that  $(G, \tau, \oplus)$  is a metrizable strongly topological gyrogroup, does  $G$  have a suitable set?*

**Question 4.17.** *Does each locally compact (strongly) topological gyrogroup have a suitable set? What if the space is compact?*

**Question 4.18.** *Suppose that  $(G, \tau, \oplus)$  is a Hausdorff strongly topological gyrogroup with a symmetric open neighborhood base  $\mathcal{U}$  at 0 which satisfies  $d(G) < b(G)$ , does  $G$  have a closed suitable set?*

**Acknowledgments.** The authors thank to Alex Ravsky for valuable remarks and corrections and all other sort of help related to the content of this paper.

## REFERENCES

- [1] A.V. Arhangel'skiĭ, M. Tkachenko, *Topological Groups and Related Structures*, Atlantis Press and World Sci., 2008.
- [2] W. Atiponrat, *Topological gyrogroups: generalization of topological groups*, Topol. Appl., **224** (2017) 73–82.
- [3] T. Banakh, I. Protasov, *Ball Structures and Colorings of Graphs and Groups*, VNTL Publ., Lviv, 2003.
- [4] M. Bao, F. Lin, *Feathered gyrogroups and gyrogroups with countable pseudocharacter*, Filomat, **33**(16) (2019) 5113–5124.
- [5] M. Bao, F. Lin, *Submetrizable strongly topological gyrogroups*, arxiv.org/abs/2003.06132.
- [6] M. Bao, F. Lin, *Quotient with respect to admissible  $L$ -subgyrogroups*, arxiv.org/abs/2003.08843v2.
- [7] Z. Cai, S. Lin, W. He, *A note on Paratopological Loops*, Bulletin of the Malaysian Math. Sci. Soc., **42**(5) (2019) 2535–2547.

- [8] W.W. Comfort, S.A. Morris, D. Robbie, S. Svetlichny, M. Tkachenko, *Suitable sets for topological groups*, Topol. Appl., **86** (1998) 25–46.
- [9] D. Dikranjan, M. Tkachenko, V. Tkachuk, *Some topological groups with and some without suitable sets*, Topol. Appl., **98** (1999) 131–148.
- [10] D. Dikranjan, M. Tkačenko, V. Tkachuk, *Topological groups with thin generating sets*, J. Pure Appl. Algebra, **145**:2 (2000), 123–148.
- [11] R. Engelking, General Topology (revised and completed edition), Heldermann Verlag, Berlin, 1989.
- [12] M. Ferreira, *Factorizations of Möbius gyrogroups*, Adv. Appl. Clifford Algebras, **19** (2009) 303–323.
- [13] M. Ferreira, G. Ren, *Möbius gyrogroups: A Clifford algebra approach*, J. Algebra, **328** (2011) 230–253.
- [14] I. Guran, *Suitable sets for paratopological groups*, Abstracts of 4-th International Algebraic Conference in Ukraine (Lviv, 2003), 87–88.
- [15] K.H. Hoffmann, S.A. Morris, *Weight and  $c$* , J. Pure Appl. Algebra, **68** (1990) 181–194.
- [16] F. Lin, R. Shen, *On rectifiable spaces and paratopological groups*, Topol. Appl., **158** (2011) 597–610.
- [17] F. Lin, C. Liu, S. Lin, *A note on rectifiable spaces*, Topol. Appl., **159** (2012) 2090–2101.
- [18] F. Lin, A. Ravsky, *Suitable sets for paratopological groups*, [arXiv:2005.08233v1](https://arxiv.org/abs/2005.08233v1)
- [19] F. Lin, *Compactly generated rectifiable spaces or paratopological groups*, Math. Commun., **18** (2013) 417–427.
- [20] F. Lin, J. Zhang, K. Zhang, *Locally  $\sigma$ -compact rectifiable spaces*, Topol. Appl., **193** (2015) 182–191.
- [21] L.V. Sabinin, L.L. Sabinin, L.V. Sbitneva, *On the notion of gyrogroup*, Aequ. Math., **56** (1998) 11–17.
- [22] R. Shen, *The first-countability of paratopological left-loops*, Topol. Appl., Article 107190.
- [23] T. Suksumran, K. Wiboonton, *Isomorphism theorems for gyrogroups and  $L$ -subgyrogroups*, J. Geom. Symmetry Phys., **37** (2015) 67–83.
- [24] T. Suksumran, *Essays in mathematics and its applications: in honor of Vladimir Arnold*, in: P.M. Pardalos, T.M. Rassias (Eds.), *The Algebra of Gyrogroups: Cayley’s Theorem, Lagrange’s Theorem, and Isomorphism Theorems*, Springer, 2016, 369–437.
- [25] T. Suksumran, *Special subgroups of gyrogroups: commutators, nuclei and radical*, Math. Interdiscip. Res, **1** (2016) 53–68.
- [26] A.A. Ungar, *Analytic Hyperbolic Geometry and Albert Einstein’s Special Theory of Relativity*, World Scientific, Hackensack, New Jersey, 2008.
- [27] A.A. Ungar, *Beyond the Einstein addition law and its gyroscopic Thomas precession: The theory of gyrogroups and gyrovectors spaces*, Fundamental Theories of Physics, vol. 117, Springer, Netherlands, 2002.

FUCAI LIN: SCHOOL OF MATHEMATICS AND STATISTICS, MINNAN NORMAL UNIVERSITY, ZHANGZHOU 363000, P. R. CHINA

*E-mail address:* [linfucai2008@aliyun.com](mailto:linfucai2008@aliyun.com); [linfucai@mnnu.edu.cn](mailto:linfucai@mnnu.edu.cn)

TINGTING SHI: SCHOOL OF MATHEMATICS AND STATISTICS, MINNAN NORMAL UNIVERSITY, ZHANGZHOU 363000, P. R. CHINA

*E-mail address:* [277653220@qq.com](mailto:277653220@qq.com)

MENG BAO: SCHOOL OF MATHEMATICS AND STATISTICS, MINNAN NORMAL UNIVERSITY, ZHANGZHOU 363000, P. R. CHINA

*E-mail address:* [mengbao95213@163.com](mailto:mengbao95213@163.com)