# ASYMPTOTICALLY MEAN VALUE HARMONIC FUNCTIONS IN DOUBLING METRIC MEASURE SPACES

TOMASZ ADAMOWICZ<sup>1</sup>, ANTONI KIJOWSKI, AND ELEFTERIOS SOULTANIS<sup>2</sup>

ABSTRACT. We consider weakly and strongly asymptotically mean value harmonic (amv-harmonic) functions on metric measure spaces which, in the classical setting, are known to coincide with harmonic functions.

We demonstrate that, in non-collapsed RCD-spaces with vanishing metric measure boundary, Cheeger harmonic functions are weakly amv-harmonic and that, in Carnot groups, weak amv-harmonicity equivalently characterizes harmonicity in the sense of the sub-Laplacian. Moreover, in the first Heisenberg group, we prove a Blaschke–Privaloff–Zaremba type theorem which yields the equivalence of both weak and strong amv-harmonicity with harmonicity in the sense of the sub-Laplacian.

In doubling metric measure spaces we show that strongly amv-harmonic functions are Hölder continuous for any exponent below one. More generally, we define the class of functions with finite amv-norm and show that functions in this class belong to a fractional Hajłasz–Sobolev space and their blow-ups satisfy the mean-value property.

In the toy model case of weighted Euclidean domains, we identify the elliptic PDE characterizing amv-harmonic functions, and use this to point out that Cheeger harmonic functions in RCD-spaces need not be weakly amv-harmonic without the non-collapsing condition.

*Keywords*: Asymptotic mean value property, Synthetic curvature bounds, Carnot groups, harmonic functions, Dirichlet forms, Gromov–Hausdorff convergence, Korevaar–Schoen energy, mean value property, metric measure boundary, Sobolev spaces.

Mathematics Subject Classification (2010): Primary: 31E05; Secondary: 53C23, 35R03.

#### **CONTENTS**

1. Introduction	2
1.1. Harmonic and amv-harmonic functions in singular spaces	2
1.2. Amv-harmonicity in general doubling measure spaces	5
1.3. Weighted Euclidean domains	7
2. Preliminaries	8
2.1. Notation and conventions	8
2.2. Doubling measures and averaging operators	9
2.3. Hajłasz–Sobolev spaces	10
2.4. Cheeger and Korevaar–Schoen energy, and Dirichlet forms	11
3. Refined averaging and mean value harmonic functions	14
3.1. Local Lipschitz continuity of mean value-harmonic functions	14
3.2. Mean value-harmonic functions of polynomial growth	17
4. Hölder continuity of strongly amv-harmonic functions	19
5. Blow-ups of Hajłasz–Sobolev functions with finite AMV-norm	22
5.1. Pointed measured Gromov–Hausdorff convergence	22
5.2. Gromov–Hausdorff convergence and averaging operators	25
5.3. Blow-ups of Sobolev functions with finite amv-norm	26

<sup>1</sup>T. Adamowicz was supported by a grant of National Science Center, Poland (NCN), UMO-2017/25/B/ST1/01955.

<sup>&</sup>lt;sup>2</sup>E. Soultanis was supported by Swiss National Foundation grant no. 182423.

6. Weakly amv-harmonic functions	29
6.1. Symmetrized <i>r</i> -laplacian and the Korevaar–Schoen energy	29
6.2. Carnot groups	30
6.3. RCD-spaces	32
7. Improving regularity: the BPZ-theorem for amv-harmonic functions	35
8. Weighted Euclidean spaces. Elliptic PDEs and amv-harmonic functions	38
8.1. $\operatorname{AMV}_{loc}^{p}(\Omega_{w})$ as a second order Sobolev space	39
8.2. PDE characterization of amv-harmonicity	43
References	44

### 1. INTRODUCTION

1.1. Harmonic and amv-harmonic functions in singular spaces. A central feature of harmonic functions on Euclidean domains is the mean value property. Although the Euclidean structure is crucial for this its validity, the mean value property can be expressed using only the metric and the measure, in terms of having vanishing *r*-laplacian. We define the *r*-laplacian of a function  $u \in L^1_{loc}(X)$  on a metric measure space  $X = (X, d, \mu)$  as follows

$$\Delta_r^{\mu} u(x) = \Delta_r u(x) = \frac{u_{B_r(x)} - u(x)}{r^2}, \quad x \in X,$$
(1)

where  $u_{B_r(x)}$  stands for the mean-value of u over a ball  $B_r(x)$ . The classical mean value property states that, in a Euclidean domain  $\Omega$ , a harmonic function u satisfies  $\Delta_r u(x) = 0$ , for all  $0 < r < \text{dist}(x, \partial \Omega)$ . Functions on a domain of a metric measure space with this property are called *mean value harmonic* (mv-harmonic), see Definition 3.1.

It turns out that harmonic functions rarely enjoy the mean value property outside the Euclidean setting and instead satisfy an *asymptotic mean value property*, where the pointwise limit  $r \rightarrow 0$  in (1) vanishes. For example, this is the case of smooth Riemannian manifolds, whereas the mean value property for harmonic functions on manifolds is known to hold only on the so-called harmonic manifolds. The Lichnerowicz conjecture, proven for manifolds of dimension 2-5, characterizes harmonic manifolds as either flat or rank-one symmetric, see Example 4 in [1] and references therein. The converse statement, namely when the asymptotic mean value property implies that a function satisfies the appropriate Laplace equation is known as the Blaschke–Privaloff–Zaremba (BPZ) theorem, and will be discussed in more detail below, see Theorem 1.4.

Apart from the classical setting, the *r*-laplacian also arises in approximation problems of Riemannian manifolds by graphs [14], and the mean value property plays a role in geometric group theory in Kleiner's proof of Gromov's polynomial growth theorem [47] (see also [57] for a notion of amv-laplacian on metric measure spaces). Furthermore, [21, Theorem 1] indirectly relates amvharmonicity to functions with bounded variation. Namely, the result characterizes  $C^1$ -minimal surfaces S by observing that a certain piecewise constant function  $f_S$  is pointwise amv-harmonic in the sense that  $\Delta_r f_S \to 0$  pointwise, as  $r \to 0$  on S. Moreover, the proof of [21, Theorem 1] uses the relation between the amv-harmonic operator and a nondegenerate 1-Laplacian:  $\operatorname{div}(\nabla/\sqrt{1+|\nabla|^2})$ . In the setting of Carnot groups the *r*-Laplacian and its relations to the subelliptic harmonic functions have been studied, for instance, in [2, 24, 25].

In this paper we consider two notions of *asymptotically mean value harmonic* (amv-harmonic) functions, arising from different ways to interpret the limit in the equation

$$\lim_{r \to 0} \Delta_r u = 0$$

More precisely, we define strong and weak amv-harmonicity as follows.

**Definition 1.1.** Let  $X = (X, d, \mu)$  be a metric measure space.

(1) A function  $u \in L^1_{loc}(X)$  is strongly anv-harmonic, if

$$\lim_{r \to 0} \|\Delta_r u\|_{L^{\infty}(K)} = 0$$

for any compact set  $K \subset X$ ;

(2) A function  $u \in L^2_{loc}(X)$  on a metric measure space X is *weakly anv-harmonic* if

$$\lim_{r \to 0} \int_X \varphi \Delta_r u \mathrm{d}\mu = 0$$

for every compactly supported Lipschitz function  $\varphi$ .

We also say that a weakly (resp. strongly) amv-harmonic function *has the weak (resp. strong) amv-property.* 

Our first two results establish the existence of many weakly amv-harmonic functions in metric measure spaces with strong geometric control. (In what follows, by *harmonic* functions we mean continuous local minimizers of the Cheeger energy, cf. Definition 2.3.) In both theorems, two key features play a central role: that the Korevaar–Schoen energy defines a Dirichlet form, and that the spaces do not have boundary in a measure theoretic sense. We refer to Definitions 2.4 and 1.2 for the precise formulations.

First we consider spaces satisfying synthetic Riemannian curvature conditions (RCD-spaces). We refer, for instance, to [5, 6, 66] for more details on synthetic curvature bounds (see also Section 6.3). An RCD(K, N)-space  $X = (X, d, \mu)$  is called *non-collapsed* if  $\mu = \mathcal{H}^N$ , where  $\mathcal{H}^N$  denotes the *N*-dimensional Hausdorff measure.

**Theorem 1.1.** Let  $\Omega \subset X$  be a domain in a non-collapsed RCD(K, N)-space  $X = (X, d, \mathcal{H}^N)$  with vanishing *metric measure boundary. Then harmonic functions in*  $\Omega$  *are weakly anv-harmonic in*  $\Omega$ .

In particular harmonic functions on Alexandrov spaces with vanishing mm-boundary are weakly amv-harmonic. At the end of the introduction we observe that the non-collapsing condition in Theorem 1.1 is necessary. We remark that, for locally Lipschitz continuous functions, the converse implication in Theorem 1.1 also holds, see Remark 6.4.

Our second result demonstrates that rectifiability is not essential for the connection between harmonic and weakly amv-harmonic functions. Namely, we identify a class of homogeneous norms on Carnot groups under which harmonic functions have the weak amv-property. We remark that, since the Cheeger energy arises from the horizontal gradient, harmonic functions in Carnot groups are equivalently characterized as solutions to the sub-Laplacian equation, which are known to satisfy certain *weighted* mean value properties, see [11, Chapter 5.5] and [2].

Recall that an *s*-step Carnot group  $\mathbb{G}$  with Lie algebra  $\mathfrak{g} = V_1 \oplus \ldots \oplus V_s$  can be identified with  $\mathbb{R}^m$ , where  $m = v_1 + \cdots + v_s$  and  $v_i := \dim V_i$ , for  $i = 1, \ldots, s$  (see Section 6.2). For the statement of our next result, we denote by  $z = (z^{(1)}, z^{(2)}, \ldots, z^{(s)})$  the coordinates in  $\mathbb{G}$ , where  $z^{(i)} \in \mathbb{R}^{v_i}$ . We furthermore denote by  $z' = (z^{(2)}, \ldots, z^{(s)})$  the coordinates corresponding to the non-horizontal space.

**Theorem 1.2.** Let  $\Omega \subset \mathbb{G}$  be an open domain in a Carnot group  $\mathbb{G}$  and suppose  $\rho$  is a homogeneous pseudonorm on  $\mathbb{G}$ , given by

$$\rho(z) := F(|z^{(1)}|, z'), \quad z = (z^{(1)}, z') \in \mathbb{G},$$
(2)

for some function  $F : \mathbb{R} \times \mathbb{R}^{m-v_1} \to \mathbb{R}$  that is  $C^1$  outside the origin and  $\partial_1 F > 0$ .

If  $\Omega$  denotes the metric measure space  $(\Omega, \rho, \mathcal{H}^Q)$ , where  $\mathcal{H}^Q$  is the Haar measure on  $\mathbb{G}$ , then a function  $u \in HW^{1,2}_{loc}(\Omega)$  is harmonic if and only if it is weakly amv-harmonic.

In the claim above  $HW_{loc}^{1,2}(\Omega)$  denotes the horizontal Sobolev space. Pseudonorms satisfying (2) include e.g. the Korányi–Reimann metric, see also Remark 6.2. In the special case that  $\mathbb{G} = \mathbb{H}^1$  is

the first Heisenberg group, the results in [24] imply that harmonicity is also characterized by strong amv-harmonicity.

**Corollary 1.3.** Let  $\Omega \subset \mathbb{H}^1$  be a domain in the first Heisenberg group equipped with the Korányi–Reimann metric. Then a Sobolev function  $u \in HW_{loc}^{1,2}(\Omega)$  is harmonic if and only if it is strongly anv-harmonic.

Indeed, if *u* is harmonic, then by [24, Lemma 3.3] it is strongly amv-harmonic. The opposite implication is a consequence of the following Blaschke–Privaloff–Zaremba type theorem, see also 7.1 in Section 7.

**Theorem 1.4.** Let  $\Omega \subset \mathbb{H}_1$  be a domain in the first Heisenberg group  $\mathbb{H}_1$  equipped with the Korányi–Reimann metric  $d_{\mathbb{H}^1}$  and let  $f : \Omega \to \mathbb{R}$  be a continuous function satisfying

$$\lim_{r \to 0^+} \Delta_r f(x) = 0$$

for every  $x \in \Omega$ . Then f is a sub-elliptic harmonic function, i.e.  $f \in C^{\infty}(\Omega)$  and  $\Delta_{\mathbb{H}_1} f = 0$  in  $\Omega$ .

Furthermore, similar results hold on Riemannian manifolds with Ricci curvature bounded below, see Proposition 7.2, and in Alexandrov spaces, see Proposition 7.3.

The equivalence between harmonic and strongly amv-harmonic functions in non-collapsed RCD-spaces is not known. By existing regularity theory, weakly amv-harmonic functions in RCD-spaces and Carnot groups are a posteriori known to be locally Lipschitz regular, and exist in abundance, see e.g. [39, Theorem 1.2] and [11].

Next we briefly discuss the two key features (KS-Dirichlet form and vanishing mm-boundary) mentioned after Definition 1.1.

*Spaces with a Korevaar–Schoen Dirichlet form.* Theorems 1.1 and 1.2 follow from a general principle relating the approximate Korevaar–Schoen energy to a *symmetrized r*-laplacian. In contrast to the *r*-laplacian, the symmetrized *r*-laplacian is a self-adjoint operator, defined for a locally integrable function *u* on a metric measure space  $X = (X, d, \mu)$  by

$$\widetilde{\Delta}_{r}u(x) := \frac{1}{2} \int_{B_{r}(x)} \left[ \frac{1}{\mu(B_{r}(x))} + \frac{1}{\mu(B_{r}(y))} \right] \frac{u(y) - u(x)}{r^{2}} \mathrm{d}\mu(y), \quad x \in X.$$

Suppose the Korevaar–Schoen energy is a Dirichlet form on X in the sense of Definition 2.4. Then the symmetrized r-laplacians  $\tilde{\Delta}_r u$  converge in a suitable weak sense, as  $r \to 0$ , to the Korevaar– Schoen laplacian  $\Delta_{KS} u$  of u for any  $u \in D(\Delta_{KS})$ , see Section 2.4 and Proposition 6.2. It follows that local minimizers of the Korevaar–Schoen energy are characterized as functions whose symmetrized r-laplacian vanishes in the weak sense, as  $r \to 0$ .

Both RCD-spaces and Carnot groups have the property that the Korevaar–Schoen energy is a Dirichlet form. For RCD-spaces, this follows from [31] (in the non-collapsed case also from work of Sturm [63, 64]), while for Carnot groups it is implied by [65]. See Propositions 6.4 and 6.3 for precise statements.

Vanishing metric measure boundary. To connect harmonic functions (continuous local energy minimizers) to weak amv-harmonicity, we need to prove that the difference  $\Delta_r - \tilde{\Delta}_r$  of the *r*-laplacian and the symmetrized *r*-laplacian vanishes in the limit  $r \to 0$  in a suitable sense. The assumption of vanishing metric measure boundary enters into play here. While in Carnot groups translation invariance of the measure implies that  $\Delta_r - \tilde{\Delta}_r$  vanishes identically for all r > 0, the situation is more delicate in RCD-spaces.

In this paper we will work with the notion of *vanishing metric measure boundary* (mm-boundary), introduced in [41]. It can be defined in terms of the Bishop–Gromov densities at scale r. Here  $\theta^N(\mu, x)$ 

denotes the *Bishop–Gromov density* at point  $x \in X$  defined for a measure  $\mu$  as follows

$$\theta^N(\mu, x) := \limsup_{r \to 0} \theta^N_r(\mu, x), \quad \theta^N_r(\mu, x) := \frac{\mu(B_r(x))}{\omega_N r^N}.$$
(3)

In the case  $\mu = \mathcal{H}^N$ , we denote by  $\theta^N(\cdot) := \theta^N(\mathcal{H}^N, \cdot)$ .

**Definition 1.2** (Definition 1.5 in [41]). Let  $X = (X, d, \mu)$  be an *N*-rectifiable space. We say that *X* has vanishing mm-boundary, if the signed Radon measures

$$\frac{1 - \theta_r^N(\mu, \cdot)}{r} \mathrm{d}\mu, \quad 0 < r \le 1$$

are uniformly bounded in the total variation norm, and converge weakly to zero as  $r \to 0$ .

Examples of spaces with vanishing mm-boundary include Riemannian manifolds without boundary with Lipschitz continuous metrics, convex hypersurfaces in Euclidean spaces and two-dimensional Alexandrov spaces without boundaries, see Example 1.12, Theorem 1.8 and Theorem 5.3 in [41].

By the Bishop–Gromov theorem (see Theorem 6.5), the limit in  $\theta^N(x)$  exists and satisfies  $\theta^N(x) \le 1$ for every  $x \in X$ , and  $\theta^N(x) = 1$  for  $\mathcal{H}^N$ -almost every  $x \in X$ , for a non-collapsed RCD(K, N)-space X. In Section 6 we prove that, in non-collapsed RCD-spaces with vanishing mm-boundary, the difference  $\Delta_r - \widetilde{\Delta}_r$  indeed converges to zero as  $r \to 0$  in a suitable sense, cf. Lemma 6.7 and Corollary 6.8.

We remark that in the non-collapsed RCD-spaces one studies also other notions of boundary, defined by using the stratification of its singular set, see [22, 42]. For a non-collapsed RCD(K, N)-space, the singular set S can be characterized as

$$\mathcal{S} = \{ x \in X : \ \theta^N(x) < 1 \},\$$

cf. [22, Corollary 1.7] and [42, (2.13)]. Having vanishing mm-boundary is a rather strong condition that is not yet well understood. The relationship between vanishing topological and mm-boundary is defined and investigated in [41] for Alexandrov spaces where it is conjectured that Alexandrov-spaces with empty (topological) boundary have vanishing mm-boundary, see [41, Section 1.3]. In RCD-spaces, no relationship is known to the authors of the present manuscript between vanishing mm-boundary and the boundaries defined in [22] and [42], though Ricci-limit spaces with both-sided curvature bounds are expected to have vanishing mm-boundary, cf. [41, Problem 8.13].

1.2. **Amv-harmonicity in general doubling measure spaces.** In doubling metric measure spaces, on the other hand, neither of the essential features mentioned above is available, and no existence or regularity results on weakly amv-harmonic functions are known to us. We instead consider strongly amv-harmonic functions. While their existence in this generality is not clear (see the discussion preceding Theorem 1.6), the next theorem provides Hölder regularity, for all exponents below one, of strongly amv-harmonic functions.

**Theorem 1.5.** Let  $\Omega \subset X$  be a domain in a complete doubling metric measure space. Then any strongly *amv-harmonic function on*  $\Omega$  *is locally*  $\alpha$ -Hölder continuous for any  $\alpha \in (0, 1)$ .

The proof of Theorem 1.5 is not quantitative, and we do not know whether strongly amv-harmonic functions are locally Lipschitz continuous, even under some additional Sobolev regularity (see how-ever Theorem 1.8 for local Lipschitz continuity of mv-harmonic functions).

Since the *r*-laplacians  $\Delta_r$  do not asymptotically agree with the self-adjoint operators  $\overline{\Delta}_r$ , and there is no meaningful limit operator defined on a large enough space of test functions, to achieve Theorem 1.5 we instead define the space of functions with finite *amv-norm*; given a metric measure space  $(X, d, \mu)$ , and a number  $p \in [1, \infty]$ , set

$$\operatorname{AMV}^{p}(X) := \left\{ u \in L^{p}(X) : \|u\|_{\operatorname{AMV}^{p}} < \infty \right\},$$
(4)

where

$$\|u\|_{\mathrm{AMV}^p} := \limsup \|\Delta_r u\|_{L^p(X)}$$

is the amv-norm of u. The class  $AMV_{loc}^p(X)$  is defined as the space of function  $u \in L_{loc}^p(X)$  for which  $\limsup \|\Delta_r u\|_{L^p(K)} < \infty$  for every compact  $K \subset X$ .

For functions with (locally) finite amv-norm, we prove that the blow-ups at almost every point satisfy the global mean value property (see Definition 3.1), which in general is very rare. While on Finsler manifolds Theorem 1.6 below is trivial – in the sense that the blow-ups are linear functions, which always satisy the mean value property – it can be seen as an infinitesimal connection between amv- and mean value harmonicity, and may serve as an obstruction to having many amv-harmonic functions on metric spaces that are too irregular.

**Theorem 1.6.** Let  $\Omega \subset X$  be an open subset of a proper locally doubling length space  $X = (X, d, \mu)$ , and let  $1 . Suppose <math>u \in M^{1,p}_{loc}(\Omega) \cap AMV^p_{loc}(\Omega)$  and  $(r_k)$  is a positive sequence converging to zero. Then for  $\mu$ -almost every  $x \in \Omega$ , any approximate tangent map  $u_{\infty} : X_{\infty} \to \mathbb{R}$  at x, subordinate to a subsequence of  $(r_k)$ , is mv-harmonic.

Here  $M_{loc}^{1,p}(\Omega)$  is the Hajłasz–Sobolev space, cf. Section 2.3. Hajłasz–Sobolev functions have approximate tangents at almost every point (cf. Proposition 5.8), while having finite amv-norm guarantees that any approximate tangent has the strong mean value property. See Section 5.3 for the proof of Theorem 1.6.

We do not know whether the requirement  $u \in M^{1,p}_{loc}(\Omega)$  in Theorem 1.6 is redundant. That is, we do not know whether the amv-norm bounds the Hajłasz–Sobolev norm of a function. The following theorem is a weaker statement in this direction, yielding fractional Hajłasz–Sobolev regularity of functions with finite amv-norm.

**Theorem 1.7.** Let  $\Omega \subset X$  be an open subset of a proper locally doubling metric measure space  $X = (X, d, \mu)$ , and let  $u \in AMV_{loc}^{p}(\Omega)$ . Then  $u \in M_{loc}^{\alpha,p}(\Omega)$  for every  $0 < \alpha < 1$ .

Theorem 1.7 is not quantitative, and gives no explicit bounds on the fractional Hajłasz–Sobolev gradient in terms of the amv-norm. We refer to Section 4 and Proposition 4.1 which directly implies Theorem 1.7. Combined with the Morrey embedding theorem for fractional Hajłasz–Sobolev spaces (see Proposition 2.1 and [69, Corollary 1.4]), we obtain Theorem 1.5 and the stronger Corollary 4.2 from Theorem 1.7.

The proof of Proposition 4.1 relies on an iterated argument involving a refined averaging operator, introduced in Section 3. The action of the refined averaging operator leaves strongly mv-harmonic functions invariant (in a local sense). This observation can be used to prove local Lipschitz regularity of mv-harmonic functions, see Theorem 1.8 below. This improves Corollaries 5.1 and 5.2 in [1], where local Lipschitz regularity is obtained, respectively, for uniform measures and measures satisfying the annular decay condition and under the assumption that the underlying space supports the (1, p)-Poincaré inequality.

**Theorem 1.8.** Let  $\Omega \subset X$  be an open subset of a complete locally doubling metric measure space  $X = (X, d, \mu)$ , and  $u \in L^1_{loc}(\Omega)$  an mv-harmonic function on  $\Omega$ . Then u is locally Lipschitz and satisfies the bound

$$\operatorname{LIP}(u|_{B_r(x_0)}) \le \frac{C}{r} \inf_{c \in \mathbb{R}} \oint_{B_{3r}(x_0)} |u - c| \mathrm{d}\mu$$
(5)

whenever  $\bar{B}_{3r}(x_0) \subset \Omega$ .

*Mean value harmonic functions with polynomial growth.* In Section 3.2 we study dimension upper bounds for the linear space of mean-value harmonic functions on metric measure spaces with polynomial growth rate at most m, denoted  $\mathcal{H}^m$ . These results go back to [19, 20] for Riemannian manifolds with non-negative Ricci curvature, see also [40, 39] for Alexandrov and RCD-spaces. We prove that in a

complete doubling metric space *X* equipped with the measure satisfying the  $\alpha$ -annular decay (cf. Definition 2.1) it holds that

$$\dim \mathcal{H}^m(X) \le Cm^{Q-\alpha},$$

where the constant  $C = C(Q, \alpha)$  depends only on the doubling exponent Q and  $\alpha$ , see Proposition 3.4 for the detailed formulation of our result. The proof is a straightforward adaptation of an argument of Li [52].

1.3. Weighted Euclidean domains. As a toy model, we consider weighted Euclidean domains. Let  $\|\cdot\|$  be a norm on  $\mathbb{R}^n$  and  $\Omega \subset \mathbb{R}^n$  a domain. We denote by  $\mathcal{H}^n$  the Hausdorff *n*-measure, taken with respect to the metric induced by  $\|\cdot\|$ , restricted to  $\Omega$ . By a *weight* on  $\Omega$  we mean a strictly positive locally Lipschitz function  $w : \Omega \to (0, \infty)$ . In what follows, we consider the corresponding weighted Euclidean domain  $\Omega_w := (\Omega, \|\cdot\|, wd\mathcal{H}^n)$ .

Observe that  $\Omega_w$  has vanishing mm-boundary if and only if  $w \equiv 1/J(\|\cdot\|)$ , where  $J(\|\cdot\|)$  denotes the Jacobian of the norm  $\|\cdot\|$  (see Definition 5 and Lemma 6(ii) in [46]). Indeed, for any compact  $K \subset \Omega$  and r > 0 such that  $B_r(x) \subset K$ , the estimate

$$|1 - \theta_r^n(w d\mathcal{H}^n, x)| \ge |1 - J(\|\cdot\|)w(x)| - J(\|\cdot\|) \operatorname{LIP}(w|_K)r, \quad x \in K$$

implies that, if  $\frac{1-\theta_r^n(wd\mathcal{H}^n,)}{r}wd\mathcal{H}^n$  is uniformly bounded in total variation norm, then  $w \equiv 1/J(\|\cdot\|)$ .

Thus the *r*-laplacian  $\Delta_r^w$  on  $\Omega_w$  may fail to be asymptotically self-adjoint. Nevertheless, weak any-harmonicity in  $\Omega_w$  can characterized by an elliptic PDE. Consider the unbounded operator

$$L_w u := \frac{1}{2} \operatorname{div}(M \nabla u) + \frac{1}{w} \langle \nabla w, M \nabla u \rangle, \tag{6}$$

where  $\langle \cdot, \cdot \rangle$  is the standard Euclidean inner product, and  $M \in \mathbb{R}^{n \times n}$  denotes the positive definite *matrix of second moments* of the unit ball  $B^n \subset \mathbb{R}^n$  of  $\|\cdot\|$  given by

$$m_{ij} := \int_{B^n} y_i y_j \mathrm{d}\mathcal{H}^n(y), \quad 1 \le i, j \le n,$$

see [45, Sections 4.1-4.3].

**Theorem 1.9.** Let  $u \in W^{1,2}_{loc}(\Omega)$  for a domain  $\Omega \subset \mathbb{R}^n$ . Then the following are equivalent.

- (1)  $L_w u = 0$  in  $\Omega$  in the weak sense,
- (2) *u* is weakly amv-harmonic in  $\Omega_w$ , and
- (3)  $u \in AMV_{loc}^2(\Omega_w)$  and  $\Delta_r^w u \to 0$  in  $L^2_{loc}(\Omega_w)$  as  $r \to 0$ .

Moreover, if in addition  $w \in C^{\infty}(\Omega)$ , then (1), (2) and (3) are equivalent to

(4) *u* is strongly amv-harmonic in  $\Omega_w$ .

We refer to [57] and [13] for some related results. The non-trivial implications in Theorem 1.9 boil down to regularity theory of elliptic PDE's, and a characterization of  $AMV_{loc}^2(\Omega_w)$  as a second order Sobolev space, see Theorem 8.2 in Section 8.

In view of classical existence theory, weakly and strongly amv-harmonic functions are abundant under the hypotheses of Theorem 1.9 and their further studies in the weighted setting could be interesting per se. In contrast, mean value harmonic functions (with  $\Omega = \mathbb{R}^n$ ) satisfy an overdetermined system of PDE's – in particular, if  $\|\cdot\|$  is an  $l^p$ -norm ( $p \neq 2$ ), mv-harmonic functions on (unweighted)  $\mathbb{R}^2$  form a finite dimensional space; see [45].

We finish the introduction by observing a consequence of Theorem 1.9: harmonic functions in RCDspaces need not be amv-harmonic without the non-collapsing condition. Indeed, consider  $\Omega = \mathbb{R}^n$  with the standard Euclidean norm, and  $w = e^{-f}$ , where  $f \in C^{\infty}(\mathbb{R}^n)$ . Then  $X_w := (\mathbb{R}^n, w d\mathcal{H}^n)$  is an RCD(K, N)-space, for  $K \in \mathbb{R}$  and N > n, if

$$\operatorname{Hess} f \ge K \operatorname{id} \otimes \operatorname{id} + \frac{1}{N - n} \nabla f \otimes \nabla f,$$

cf. [7] and [67]. In this case the Laplacian  $\Delta_w$  associated to the Cheeger energy

$$Ch_w(\varphi) := \frac{1}{2} \int_{\mathbb{R}^n} |\nabla \varphi|^2 w \mathrm{d}\mathcal{H}^n$$

is given by

$$\Delta_w \varphi = \Delta \varphi - \langle \nabla f, \nabla \varphi \rangle, \quad \varphi \in D(\Delta_w),$$

whereas the "amv-operator"  $L_w$  in (6) satisfies

$$L_w \varphi := \frac{1}{n+2} \left[ \frac{1}{2} \Delta \varphi - \langle \nabla f, \nabla \varphi \rangle \right], \quad \varphi \in AMV^2(X_w).$$

It follows that harmonic functions, i.e. weak solutions of  $\Delta_w u = 0$ , are distinct from weakly amvharmonic functions on  $X_w$ . It is also noteworthy that the factor 1/2 makes  $L_w$  a non-self-adjoint operator unless w is constant.

#### 2. Preliminaries

2.1. Notation and conventions. Throughout this article, a metric measure space  $X = (X, d, \mu)$  is a separable metric space (X, d) equipped with a Radon measure  $\mu$  that is finite and non-trivial on balls, i.e.  $0 < \mu(B) < \infty$  for all balls  $B \subset X$ .

Given a subset  $F \subset X$  of a metric space and r > 0, we denote

$$N_r(F) = \{x \in X : \operatorname{dist}(x, F) < r\}$$
 and  $\overline{N}_r(F) = \{x \in X : \operatorname{dist}(x, F) \le r\}$ 

the open and closed *r*-neighbourhood of *F* (note that  $\overline{N}_r(F)$  need not be the closure of  $N_r(F)$  unless *X* is a length space). For  $x \in X$ , we denote by  $B_r(x) := N_r(\{x\})$  and  $\overline{B}_r(x) := \overline{N}_r(\{x\})$ , respectively, an open and closed ball centered at *x* with radius *r*. Given  $s \ge 0$ , the Hausdorff *s*-measure  $\mathcal{H}^s$  is defined by

$$\mathcal{H}^{s}(A) = \lim_{\delta \to 0} \inf \left\{ \sum_{i} \omega_{s} (\operatorname{diam} A_{i}/2)^{s} : A \subset \bigcup_{i} A_{i}, \operatorname{diam}(A_{i}) \leq \delta \right\}, \quad A \subset X,$$

where  $\omega_s = \frac{\pi^{s/2}}{\Gamma(s/2+1)}$ . The normalizing constant is chosen so that  $\mathcal{H}^n$  coincides with the Lebesgue measure on  $\mathbb{R}^n$ .

The Lipschitz constant of a map  $f: (X, d_x) \to (Y, d_Y)$  between metric spaces is

$$\operatorname{LIP}(f) := \sup_{x \neq y} \frac{d_Y(f(x), f(x))}{d_X(x, y)},$$

and the pointwise Lipschitz slope is

$$\operatorname{Lip} f(x) := \limsup_{r \to 0} \sup_{0 < d(y,x) < r} \frac{d_Y(f(x), f(y))}{r}, \quad x \in X.$$

Given  $u \in L^1_{loc}(X)$  and a  $\mu$ -measurable set  $A \subset X$  with  $\mu(A) > 0$ ,

$$u_A = \oint_A u \mathrm{d}\mu = \frac{1}{\mu(A)} \int_A u \mathrm{d}\mu$$

denotes the average of u over A.

2.2. Doubling measures and averaging operators. A measure  $\mu$  on a separable metric space X is called *locally doubling* if, for every compact  $K \subset X$ , there exists  $r_K > 0$  and a constant  $C_K > 0$ , such that

$$\mu(B_{2r}(x)) \le C_K \mu(B_r(x)) \tag{7}$$

for every  $x \in K$  and  $0 < r \le r_K$ . If  $\mu$  is locally doubling, then for every compact  $K \subset X$  there exists a constant C > 0 for which

$$\frac{\mu(B_r(y))}{\mu(B_R(x))} \ge C\left(\frac{r}{R}\right)^Q, \quad x \in K, \ y \in B_R(x), \ 0 < r \le R \le r_K,$$
(8)

where  $Q = \log_2 C_K$ . If the constant  $C_K$  can be chosen independently of the set  $K \subset X$ , and  $r_K = \infty$ , then we say that  $\mu$  is *doubling*, denote  $C_K = C_{\mu}$  and the number  $Q = \log_2 C_{\mu}$  is called the *doubling* exponent of  $\mu$ .

The following definition is due to Buckley, see [15, Section 1], and is stronger than the doubling condition.

**Definition 2.1.** Let  $(X, d, \mu)$  be a metric measure space with a doubling measure  $\mu$ . We say that X satisfies the  $\alpha$ -annular decay property with some  $\alpha \in (0, 1]$  if there exists  $A \ge 1$  such that for all  $x \in X$ , r > 0 and  $\epsilon \in (0, 1)$  it holds that

$$\mu\left(B(x,r)\setminus B(x,r(1-\epsilon))\right) \le A\epsilon^{\alpha}\mu(B(x,r)).$$
(9)

If  $\alpha = 1$ , then we say that *X* satisfies the *strong annular decay property*.

Examples of spaces with strong annular decay property include geodesic metric spaces with uniform measures and Heisenberg groups  $\mathbb{H}^n$  equipped with a left-invariant Haar measures. By [15, Corollary 2.2], a length space with a doubling measure has the  $\alpha$ -annular decay property for some  $\alpha \in (0,1]$  with  $\alpha$  depending only on a doubling constant of the measure. (In fact, it is enough for a metric measure space to be the so-called  $(\alpha, \beta)$ -chain space to conclude that the space has the  $\delta$ annular decay property, see Theorem 2.1 in [15]).

Let  $(X, d, \mu)$  be a metric measure space and r > 0. Given a locally integrable function  $u \in L^1_{loc}(X)$ , we denote by

$$A^{\mu}_{r}u(x) = A_{r}u(x) = \oint_{B_{r}(x)} u \mathrm{d}\mu, \quad x \in X,$$

the *r*-average function of *u*. Note that  $A_r u(x) = u_{B_r(x)}$ .

We will use the two notations interchangeably, depending on whether we want to view the average as a number, or an operator on a function space. Indeed, the function  $A_r u : X \to \mathbb{R}$  is measurable, and  $A_r$  defines a bounded linear operator  $A_r : L^1(X) \to L^1(X)$  if and only if  $a_r \in L^{\infty}(X)$ , where

$$a_r(x) = \int_{B_r(x)} \frac{\mathrm{d}\mu(y)}{\mu(B_r(y))}, \quad x \in X.$$

Moreover, in this case the operator norm satisfies  $||A_r||_{L^1 \to L^1} = ||a_r||_{L^{\infty}}$ , see [3, Theorem 3.3]. This is true in particular when  $\mu$  is a doubling measure. On the other hand, it is true that by the Lebesgue differentiation theorem

$$u(x) = \lim_{r \to 0} A_r u(x)$$
 for almost every  $x \in X$ ,

if  $\mu$  is infinitesimally doubling, cf. [38, Remark 3.4.29].

If X is doubling as a metric space, then there exists C > 0 so that  $||A_r||_{L^p \to L^p} \leq C$  for every r > 0and every  $1 \leq p < \infty$ , cf. [3, Theorem 3.5]. However,  $A_r$  is not a self-adjoint operator; the formal adjoint  $A_r^*$  of  $A_r$  is given by

$$(A_r^{\mu})^* u(x) = A_r^* u(x) := \int_{B_r(x)} \frac{u(y) \mathrm{d}\mu(y)}{\mu(B_r(y))}, \quad x \in X,$$

for  $u \in L^1_{loc}(X)$ . Indeed, a direct computation using the Fubini theorem yields that

$$\int_{X} vA_{r}u \mathrm{d}\mu = \int_{X} uA_{r}^{*}v \mathrm{d}\mu, \quad u \in L^{p}(X), \ v \in L^{q}(X),$$
(10)

where 1/p + 1/q = 1.

We may express the *r*-laplacian using the averaging operator as

$$\Delta_r u = \frac{A_r u - u}{r^2}, \quad u \in L^1_{loc}(X)$$

Next, we denote by

$$\Delta_r^* u := \frac{A_r^* u - u}{r^2}, \quad u \in L^1_{loc}(X),$$
(11)

the formal adjoint of the *r*-laplacian. Note that if  $A_r : L^p(X) \to L^p(X)$  is bounded, then  $\Delta_r : L^p(X) \to L^p(X)$  and  $\Delta_r^* : L^q(X) \to L^q(X)$  are both bounded, where 1/p + 1/q = 1.

**Remark 2.1.** Identity (10) together with (11) imply the following Green-type formula for  $u \in L^p(X)$ ,  $v \in L^q(X)$  where 1/p + 1/q = 1:

$$\int_X v\Delta_r u \mathrm{d}\mu = \int_X u\Delta_r^* v \mathrm{d}\mu.$$

The asymptotic behaviour of  $\Delta_r - \Delta_r^*$  as  $r \to 0$  plays an important role in Section 6.

**Remark 2.2.** While most of our results will be formulated for metric measure spaces, they encompass the case of an open set  $\Omega \subset X$  in the introduction. Indeed, an open subset  $\Omega \subset X$  of a metric measure space can be regarded as a metric measure space  $\Omega = (\Omega, d|_{\Omega}, \mu|_{\Omega})$ . In particular, *if X is locally doubling*, *then*  $\Omega$  *is locally doubling*.

2.3. Hajłasz–Sobolev spaces. We briefly recall fractional Hajłasz–Sobolev spaces and their local variants.

**Definition 2.2.** Let  $(X, d, \mu)$  be a metric measure space, and  $1 \le p \le \infty$ ,  $0 < \alpha \le 1$ . The *local fractional Hajtasz-Sobolev space*  $M_{loc}^{\alpha,p}(X)$  consists of all Borel functions  $u \in L_{loc}^p(X)$  with the following property: there exists a null set  $N \subset X$  and, for every compact  $K \subset X$ , a non-negative function  $g_K \in L_{loc}^p(X)$  and  $r_K > 0$  such that

$$|u(x) - u(y)| \le d(x, y)^{\alpha} [g_K(x) + g_K(y)], \quad x, y \in K \setminus N, \ d(x, y) < r_K.$$
(12)

If we can choose  $g_K = g \in L^p(X)$  independently of K, and  $r_K = \infty$ , we say that  $u \in M^{\alpha,p}(X)$ , and call g a Hajłasz gradient of u.

Fractional Sobolev–Hajłasz spaces on connected doubling spaces (and more generally RD-spaces, cf. [50]) coincide with a Triebel–Lizorkin space for appropriate parameters. For more details on the connection between Sobolev–Hajłasz, Triebel–Lizorkin and Besov spaces we refer to [50, 51]. Here we record the following Morrey embedding theorem for fraction! Sobolev–Hajłasz spaces, see [69, Corollary 1.4].

**Proposition 2.1** (Fractional Morrey embedding). Let  $B \subset X$  be a compact ball in a doubling metric measure space so that (8) holds in B with exponent Q. Let  $0 < \alpha \le 1$ ,  $0 and suppose that <math>\alpha p > Q$ . Then there exists a constant C > 0 so that every  $u \in M^{\alpha,p}(B)$  satisfies

$$|u(x) - u(y)| \le Cd(x,y)^{\alpha - Q/p} \left( \oint_B g^p \mathrm{d}\mu \right)^{1/p}, \quad x, y \in B,$$

whenever  $g \in L^p(B)$  is a Hajłasz gradient of u.

Proposition 2.1 also follows from [34, Theorem 8.7] by considering the metric space  $(X, d^{\alpha}, \mu)$ . We omit the details.

2.4. **Cheeger and Korevaar–Schoen energy, and Dirichlet forms.** In this section we recall some basic notions in calculus in metric measure spaces, needed in the discussion of the harmonic functions according to Cheeger, see Theorems 1.1 and 1.2, in Section 6 and in the proof of Proposition 5.8. Moreover, we discuss the Korevar–Schoen energy and its relation to the Dirichlet forms.

We say that a nonnegative Borel function g on metric space X is an *upper gradient* of a function  $u : X \to [-\infty, \infty]$ , if for all nonconstant rectifiable curves  $\gamma : [0, \ell(\gamma)] \to X$ , parametrized by arc length, we have

$$|u(x) - u(y)| \le \int_{\gamma} g \, ds$$

where x and y are the two endpoints of curve  $\gamma$ . If at least one of the values |u(x)|, |u(y)| is infinite, then we apply convention that  $\int_{\gamma} g \, ds = \infty$ . Recall that a family of rectifiable curves in X is of zero p-modulus if there is a non-negative Borel measurable function  $g \in L^p(X)$  such that  $\int_{\gamma} g \, ds = \infty$  for each  $\gamma \in \Gamma$ . We refer, for instance, to Chapter 5 in [38] for the definition of the p-modulus of a family of curves and its basic properties. We say that g is a p-weak upper gradient of u if the above inequality holds outside the collection of curves of p-modulus zero. However, it turns out that the more handy unique gradient to work with, is the so called *minimal p-weak upper gradient*  $g_u \in L^p(X)$ , which is the p-weak upper gradient with smallest  $L^p$ -norm, see e.g. the discussion in [38].

Let us recall the following version of Sobolev spaces on the metric measure space  $(X, d, \mu)$ ; see [38] for more on this space. Let  $u : X \to [-\infty, \infty]$  be a measurable function and set

$$||u||_{N^{1,p}(X)}^p := \int_X |u|^p \, d\mu + \inf_g \int_X g^p \, d\mu,$$

where the infimum is taken over all *p*-weak upper gradients *g* of *u*. With this notation we define the *Newtonian space* on *X* as follows:  $N^{1,p}(X) = \{u : ||u||_{N^{1,p}(X)} < \infty\} / \sim$ , where functions *u* and *v* are equivalent, denoted  $u \sim v$ , if  $||u - v||_{N^{1,p}(X)} = 0$ . Similarly one defines the local Newtonian space  $N^{1,p}_{loc}(\Omega)$ , for a domain  $\Omega \subset X$ , see [38, Chapter 7] and [8] for precise definitions of these spaces and their further properties.

If  $X = \mathbb{R}^n$  with the Lebesgue measure and the Euclidean distance, then  $N^{1,p}(\Omega)$  (resp.  $N^{1,p}_{loc}(\Omega)$ ) coincides with the Sobolev space  $W^{1,p}(\Omega)$  (resp.  $W^{1,p}_{loc}(\Omega)$ ) and  $g_u = |\nabla u|$ .

*Cheeger energy.* Let  $X = (X, d, \mu)$  be a metric measure space and  $1 \le p < \infty$ . Following Cheeger's original work [18], the Cheeger energy  $Ch_p : L^p(X) \to [0, \infty]$  is defined as the  $L^p$ -relaxation of the slope

$$u \mapsto \frac{1}{p} \int_X (\operatorname{Lip} u)^p \mathrm{d}\mu.$$

It follows from the work of Shanmugalingam and Ambrosio–Colombo—Di Marino, cf. [62, 4], that for Newton–Sobolev functions the Cheeger energy coincides with the *p*-integral of the minimal *p*-weak upper gradient, i.e.

$$Ch_p(u) = \frac{1}{p} \int_X g_u^p \mathrm{d}\mu, \tag{13}$$

where  $g_u$  is the minimal *p*-weak upper gradient of *u*. We refer to [38, 8] for an account of upper gradients and Newton–Sobolev spaces. Cheeger's definition guarantees that  $Ch_p$  is lower semicontinuous with respect to  $L^p$ -convergence.

**Definition 2.3.** Let  $\Omega \subset X$  be a domain in a metric measure space. A continuous function  $u \in N^{1,2}_{loc}(\Omega)$  is called harmonic, if

$$Ch_2(u) \le Ch_2(u+\varphi) \tag{14}$$

for any  $\varphi \in \text{LIP}_c(\Omega)$ .

The continuity of Cheeger harmonic functions is imposed as a part of our definition. However, in path-connected doubling metric measure spaces supporting the appropriate Poincaré inequality, local Hölder continuity of Cheeger harmonic functions can be proven, also the Lipschitz regularity under stronger assumptions, see [49, Theorem 1.1] for details.

*Korevaar–Schoen energy.* In [48], the authors defined a Sobolev space on a metric measure space X by considering what is now called as the *Korevaar–Schoen energy* for  $u \in L^2_{loc}(X)$ :

$$E_{KS}^2(u) := \sup_{\varphi \in C_c(X), 0 < \varphi \le 1} \limsup_{r \to 0} \int_X \varphi(x) e_r^2(u) \mathrm{d}\mu, \tag{15}$$

where the approximate energy densities  $e_r^2(u)d\mu$  are given by

$$e_r^2(u)(x) = \frac{1}{2} \oint_{B_r(x)} \left| \frac{u(y) - u(x)}{r} \right|^2 \mathrm{d}\mu(y), \quad x \in X.$$

For each r > 0, the approximate energy densities arise from a symmetric bi-linear form  $e_r : L^2_{loc}(X) \times L^2_{loc}(X) \to L^1_{loc}(X)$  given by

$$e_r(u,v)(x) = \frac{1}{2} \int_{B_r(x)} \frac{u(y) - u(x)}{r} \frac{v(y) - v(x)}{r} d\mu(y), \quad x \in X,$$

which satisfies

$$e_r(u,u) = e_r^2(u).$$

Furthermore, we define

$$E_r(u,v) := \int_X e_r(u,v) \mathrm{d}\mu.$$

The Sobolev space  $KS^{1,2}(X)$  consisting of functions  $u \in L^2(X)$  for which  $E^2_{KS}(u) < \infty$ , equipped with the norm

$$||u||_{KS^{1,2}} = (||u||_{L^2}^2 + E_{KS}^2(u))^{1/2}$$

is known as the *Korevaar–Schoen space*. In doubling metric measure spaces it is known that there is a continuous embedding  $M^{1,2}(X) \hookrightarrow KS^{1,2}(X) \hookrightarrow N^{1,2}(X)$ . Moreover, in complete doubling spaces supporting a 2-Poincaré inequality the two spaces are equal with equivalent norms, cf. [38, Theorem 10.5.3].

*Dirichlet forms.* The Korevaar–Schoen energy does not give rise to a Dirichlet form in general metric measure spaces, since the limit in (15) need not exist.

Here, we follow [12, Part I] and say that a Dirichlet form on  $L^2(X)$  is a lower semicontinuous quadratic functional  $E : \mathcal{F} \to [0, \infty]$  on a dense domain  $\mathcal{F} \subset L^2(X)$ , which itself is a Hilbert space when equipped with the norm

$$||f||_{\mathcal{F}} := (||f||_{L^2}^2 + E(f))^{1/2}, \quad f \in \mathcal{F},$$

satisfying

$$E(\min\{f,1\}) \le E(f), \quad f \in \mathcal{F}.$$

We denote by  $\mathcal{E} : \mathcal{F} \times \mathcal{F} \to \mathbb{R}$  the bi-linear form associated to *E* and assume, throughout this paper, the following strong locality property: *if*  $f, g \in \mathcal{F}$  and *g* is constant on a neighbourhood of spt *f*, then  $\mathcal{E}(f,g) = 0$ .

The form  $\mathcal{E}$  admits a generator  $\Delta_{\mathcal{E}}$  – which we refer to as the laplacian associated to the Dirichlet form. That is,  $\Delta_{\mathcal{E}} : D(\Delta_{\mathcal{E}}) \to L^2(X)$  is a self-adjoint operator with domain  $D(\Delta_{\mathcal{E}}) \subset \mathcal{F}$  satisfying the following:

$$\int f \Delta_{\mathcal{E}} g \mathrm{d}\mu = -\mathcal{E}(f,g), \quad f \in \mathcal{F}, \ g \in D(\Delta_{\mathcal{E}}).$$

Moreover,  $\mathcal{E}$  admits a *Carré du champ*, i.e. a positive, symmetric and continuous bi-linear operator  $\Gamma : \mathcal{F} \times \mathcal{F} \to L^1(X)$  with the property that

$$\begin{split} \mathcal{E}(f,g) &= \int_X \Gamma(f,g) \mathrm{d}\mu, \quad f,g \in \mathcal{F}, \text{ and} \\ \mathcal{E}(hf,g) + \mathcal{E}(f,hg) &= \mathcal{E}(h,fg) + 2 \int_X h \Gamma(f,g) \mathrm{d}\mu, \quad f,g,h \in \mathcal{F} \cap L^\infty(X). \end{split}$$

We may localize a given Dirichlet form E on  $L^2(X)$  to a domain  $\Omega \subset X$ . We say that  $f \in \mathcal{F}_{loc}(\Omega)$  if there is a sequence  $(f_k) \subset \mathcal{F}$  and open sets  $\Omega_k \subset \Omega$  such that  $\Omega_k \subset \operatorname{spt}(f_k)$ ,  $\Omega = \bigcup_k \Omega_k$ , and  $f_k = f$  on  $\Omega_k$ . By the strong locality, we may define the Carré du Champ

$$\Gamma: \mathcal{F}_{loc}(\Omega) \times \mathcal{F}_{loc}(\Omega) \to L^1_{loc}(\Omega), \quad \Gamma(f,g) = \Gamma(f_k,g_k) \text{ on } \Omega_k,$$

where  $\bigcup_k \Omega_k = \Omega$  and  $f = f_k$  and  $g = g_k$  on  $\Omega_k$ . We can correspondingly localize  $\Delta_{\mathcal{E}}$ , and we denote by  $D(\Delta_{\mathcal{E}}; \Omega)$  the domain of  $\Delta_{\mathcal{E}}$  localized to  $\Omega$ , cf. [12, Section 7].

The Cheeger and Korevaar–Schoen energies do not define Dirichlet forms in the sense defined above in general. Indeed, while the Cheeger energy is lower semicontinuous, it need not be a quadratic form, and the Korevaar–Schoen energy is neither quadratic nor lower semicontinuous in general.

However, in RCD spaces and Carnot groups the Cheeger energy is quadratic and therefore a Dirichlet form. In the former, this gives rise to  $\Gamma$ -calculus and the formulation of the Bochner inequality, and in the latter the Dirichlet structure arising from the Cheeger energy coincides with horizontal gradients. We refer to Section 6 for a more precise discussion. In particular in these classes the Korevaar–Schoen energy is known to be a constant multiple of the Cheeger energy and thus a posteriori defines a Dirichlet form, see [31].

*Spaces with a Korevaar–Schoen Dirichlet form.* The Korevaar–Schoen energy has a distinguished role in this paper due to its close connection with the symmetrized *r*-laplacian, see Section 6.1. For the sake of a unified treatment of weakly anv-harmonic functions in Section 6, we introduce the following terminology for spaces where the Korevaar–Schoen energy defines a Dirichlet form.

**Definition 2.4.** Let  $X = (X, d, \mu)$  be a locally compact metric measure space. We say that X has a Korevaar–Schoen Dirichlet form (abbreviated KS-Dirichlet form), if  $E_{KS}$  is lower semicontinuous with respect to  $L^2$ -convergence and, for all  $u \in KS^{1,2}(X)$ , there exists an integrable function  $e^2(u)$  satisfying

$$\lim_{r\to 0} \int_X \varphi e_r^2(u) \mathrm{d}\mu = \int_X \varphi e^2(u) \mathrm{d}\mu$$

for all  $\varphi \in C_c(X)$ . The function  $e^2(u)$  is called the energy density of u.

By the previous discussion, elaborated in Section 6, RCD spaces and Carnot groups equipped with certain homogeneous norms have a KS-Dirichlet form. Sturm [63, 64] introduced the class of MCP(K, N)-spaces, which includes RCD-spaces and showed that an energy closely related to  $E_{KS}^2$  is a Dirichlet form on MCP(K, N). Note that here we do not consider the existence and lower semicontinuity of the limit in the Korevaar–Schoen energy for maps with metric space targets. This fact is known for RCD and MCP-spaces while for Carnot groups it is – to our knowledge – an open question.

If X has a KS-Dirichlet form, then the domain of  $E_{KS}^2$  is  $KS^{1,2}(X)$ , and the Carré du Champ is given by

$$e(u,v) := \frac{1}{4} [e^2(u+v) - e^2(u-v)] = \lim_{r \to 0} e_r(u,v),$$

where the convergence is in the weak sense of measures. We denote by  $\mathcal{E}_{KS}$  and  $\Delta_{KS}$  the bi-linear form, and its generator, associated to  $E_{KS}^2$ .

# 3. Refined averaging and mean value harmonic functions

In this section we consider mean value harmonic functions. In particular, we show their local Lipschitz regularity assuming merely the doubling property of the underlying measure, see Theorem 1.8. We also prove a dimension bound on the space of mv-harmonic functions with polynomial growth in the spirit of the celebrated results of Colding–Minicozzi [19, 20] confirming Yau's conjecture, see Proposition 3.4. Our approach emphasizes the role of the averaging operators.

**Definition 3.1.** Let  $X = (X, d, \mu)$  be a metric measure space. We say that a function  $u \in L^1_{loc}(X)$  is *mean value harmonic* (or has the *mean value property*) if

$$\Delta_r u = 0 \text{ on } K$$

for any compact set  $K \subset X$  and  $r < r_K := \sup\{\rho > 0 : \overline{N}_r(K) \text{ is compact}\}.$ 

If *X* is a complete doubling metric measure space with doubling exponent *Q*, and  $\Omega \subset X$  a domain, we denote by  $\mathcal{H}(\Omega)$  the space of mv-harmonic functions on  $\Omega$ . Note that, if  $u \in \mathcal{H}(\Omega)$ , then

$$u(x) = u_{B_r(x)}, \quad x \in \Omega, \ r < \operatorname{dist}(x, X \setminus \Omega).$$
(16)

The mean value property (16) has strong implications. In particular, a Harnack inequality holds on balls: if  $u \in \mathcal{H}(\Omega)$  is non-negative and  $\overline{B}_{4r}(x_0) \subset \Omega$ , then

$$\sup_{B_r(x_0)} u \le C \inf_{B_r(x_0)} u,\tag{17}$$

where the constant depends only on the doubling constant of  $\mu$ , see [1, Section 4]. A standard argument then gives the Harnack inequality for non-negative mv-harmonic functions, cf. [1, Proposition 4.2]. We refer to [1] for further properties of mv-harmonic functions in metric measure spaces. In the next section we see that the mean value property (16) also yields more regularity than one might initially expect.

3.1. Local Lipschitz continuity of mean value-harmonic functions. If a measure  $\mu$  has the  $\alpha$ -annular decay property (see Definition 2.1), then [1, Theorem 4.2] shows that mv-harmonic functions are  $\alpha$ -Hölder continuous. In this section we will prove that in fact, mv-harmonic functions are *Lipschitz* continuous even when the measure does not satisfy the annular decay condition. The idea of the proof is to consider a refined averaging process, wherein we average over the radius as well as the space variable. Throughout the subsection we use the abbreviation

$$u_r := A_r u$$

for  $u \in L^1_{loc}(X)$  and r > 0. Given such a function we define

$$A^{r}u(x) = u^{r}(x) := \frac{2}{r} \int_{r/2}^{r} u_{t}(x) dt, \quad x \in X.$$
(18)

The following elementary lemma will play a crucial role in the estimates proving that  $A^{r}u$  is locally Lipschitz. We use the notation

$$A_{r,R}(x) = B_R(x) \setminus B_r(x)$$

for  $x \in X$  and  $r \leq R$ , with the convention that  $B_r(x) = \emptyset$  for  $r \leq 0$  and  $A_{r,R}(x) = \emptyset$  if r > R.

**Lemma 3.1.** Let  $f \in L^1_{loc}(X)$  be a nonnegative function and  $x \in X$ . Let  $0 \le r \le R < \infty$ , and  $-\infty < d_1 \le d_2 < \infty$  Then

$$\int_{r}^{R} \int_{A_{t+d_{1},t+d_{2}}(x)} f \mathrm{d}\mu \, \mathrm{d}t \le (d_{2} - d_{1}) \int_{A_{r+d_{1},R+d_{2}}(x)} f \mathrm{d}\mu.$$

*Proof.* Let  $g : \mathbb{R} \to \mathbb{R}$  be the function defined as follows: g(t) = 0 for  $t \leq 0$  and

$$g(t) = \int_{B_t(x)} f \mathrm{d}\mu, \quad t \ge 0$$

Then *g* is a nondecreasing function. If  $R + d_1 \le r + d_2$ , then  $R - r \le d_2 - d_1$  and we have the trivial estimate

$$\int_{r}^{R} \int_{A_{t+d_{1},t+d_{2}}(x)} f d\mu \, dt \leq \int_{r}^{R} \int_{A_{r+d_{1},R+d_{2}}(x)} f d\mu dt \leq (d_{2}-d_{1}) \int_{A_{r+d_{1},R+d_{2}}(x)} f d\mu.$$

Assume  $R + d_1 > r + d_2$ . Then

$$\int_{r}^{R} \int_{A_{t+d_{1},t+d_{2}}(x)} f d\mu dt = \int_{r}^{R} [g(t+d_{2}) - g(t+d_{1})] dt = \int_{r+d_{2}}^{R+d_{2}} g dt - \int_{r+d_{1}}^{R+d_{1}} g dt$$
$$= \int_{R+d_{1}}^{R+d_{2}} g dt - \int_{r+d_{1}}^{r+d_{2}} g dt \le (d_{2} - d_{1})g(R+d_{2}) - (d_{2} - d_{1})g(r+d_{1})$$
$$= (d_{2} - d_{1}) \int_{A_{r+d_{1},R+d_{2}}(x)} f d\mu.$$

**Proposition 3.2.** Let  $X = (X, d, \mu)$  be a locally doubling metric measure space,  $u \in L^1_{loc}(X)$ . Let  $K \subset X$  be compact and  $r_K > 0$  such that (7) holds for  $r < r_K$ . Then, for any  $r < r_K$ , the function  $u^r$  is Lipschitz on K and, for any  $c \in \mathbb{R}$ , satisfies the Hajłasz type estimate

$$|u^{r}(x) - u^{r}(y)| \leq \frac{Cd(x,y)}{r} \left( \oint_{B_{2r}(x)} |u - c| \mathrm{d}\mu + \oint_{B_{2r}(y)} |u - c| \mathrm{d}\mu \right), \quad x, y \in K, \ d(x,y) < r,$$
(19)

where the constant C depends only on the doubling constant of  $\mu$  on K. In particular

$$\operatorname{LIP}(u^r|_{B_r(x_0)}) \le \frac{C}{r} \oint_{B_{3r}(x_0)} |u - c| \mathrm{d}\mu$$

whenever  $\bar{B}_{3r}(x_0) \subset X$  is compact.

*Proof of Proposition 3.2.* Let us first prove the Lipschitz estimate assuming (19). Let  $K = B_r(x_0)$ ,  $c \in R$  and assume that  $x, y \in B_r(x_0)$  with d(x, y) < r. Then (19) directly yields

$$|u(x) - u(y)| \le \frac{Cd(x,y)}{r} \left( \oint_{B_{2r}(x)} |u - c| \mathrm{d}\mu + \oint_{B_{2r}(y)} |u - c| \mathrm{d}\mu \right) \le \frac{Cd(x,y)}{r} \oint_{B_{3r}(x_0)} |u - c| \mathrm{d}\mu.$$

If  $d(x,y) \ge r$ , then  $d(x,x_0) + d(y,x_0) \le 2d(x,y)$ , and therefore

$$\begin{aligned} |u^{r}(x) - u^{r}(y)| &\leq |u^{r}(x) - u^{r}(x_{0})| + |u^{r}(y) - u^{r}(x_{0})| \leq \frac{C(d(x, x_{0}) + d(y, x_{0}))}{r} \int_{B_{3r}(x_{0})} |u - c| \mathrm{d}\mu \\ &\leq \frac{2Cd(x, y)}{r} \int_{B_{3r}(x_{0})} |u - c| \mathrm{d}\mu. \end{aligned}$$

Thus it suffices to prove (19). Let K and r be as in the claim and denote by  $C_K$  the doubling constant of  $\mu$  in K. Given  $x, y \in K$  with  $d := d(x, y) \leq r$  and  $r/2 \leq t < r$ , we have that for the symmetric difference of two balls it holds

$$B_t(x) \triangle B_t(y) \subset A_{t-d,t+d}(x) \subset N_{2r}(K).$$

Recall that  $N_{2r}(K)$  stands for the 2*r*-neighbourhood of *K*. This and the doubling property of  $\mu$  on *K* yield, for any  $c \in \mathbb{R}$ , the estimate

$$\begin{aligned} |u_t(x) - u_t(y)| &= \left| \int_{B_t(x)} (u - c) d\mu - \int_{B_t(y)} (u - c) d\mu \right| \\ &\leq \frac{\mu(B_t(x) \triangle B_t(y))}{\mu(B_t(x))} \int_{B_t(y)} |u - c| d\mu + \frac{1}{\mu(B_t(x))} \int_{B_t(x) \triangle B_t(y)} |u - c| d\mu \\ &\leq \frac{C_K^2 \mu(A_{t-d,t+d}(x))}{\mu(B_r(x))} \int_{B_r(y)} |u - c| d\mu + \frac{C_K}{\mu(B_r(x))} \int_{A_{t-d,t+d}(x)} |u - c| d\mu \end{aligned}$$

By Lemma 3.1 – applied twice ( $f \equiv 1$  and f = |u - c|) with  $d_1 = -d$  and  $d_2 = d$  – the above inequality implies

$$\begin{split} \int_{r/2}^{r} |u_t(x) - u_t(y)| \mathrm{d}t &\leq \frac{2C_K^2 d\mu (A_{r/2-d,r+d}(x))}{\mu (B_r(x))} \oint_{B_r(y)} |u - c| \mathrm{d}\mu + \frac{2C_K d}{\mu (B_r(x))} \int_{A_{r/2-d,r+d}(x)} |u - c| \mathrm{d}\mu \\ &\leq 2C_K^3 d\left( \oint_{B_{2r}(y)} |u - c| \mathrm{d}\mu + \oint_{B_{2r}(x)} |u - c| \mathrm{d}\mu \right), \end{split}$$

where in the last inequality we also appeal to the doubling property (8) of  $\mu$ . Using this estimate we obtain

$$|u^{r}(x) - u^{r}(y)| \leq \frac{2}{r} \int_{r/2}^{r} |u_{t}(x) - u_{t}(y)| \mathrm{d}t \leq \frac{4C_{K}^{3}d(x,y)}{r} \left( \oint_{B_{2r}(y)} |u - c| \mathrm{d}\mu + \oint_{B_{2r}(x)} |u - c| \mathrm{d}\mu \right),$$
(19).

pro ıg (19)

*Proof of Theorem 1.8.* By considering the metric measure space  $(\Omega, d, \mu|_{\Omega})$  we may assume that  $\Omega = X$ , cf. Remark 2.2. Since u is mv-harmonic, for every compact  $K \subset X$  there exists  $r_K > 0$  so that  $u = u_r$ on K for all  $r < r_K$ . In particular,  $u = u^r$  on K whenever  $r < r_K$ . The Lipschitz continuity of u and the estimate (5) then follows by Proposition 3.2. 

**Remark 3.1.** When X is complete and  $\mu$  is globally doubling, mv-harmonic functions satisfy  $u = u_R$ on X, for any R > 0. Consequently (5) yields

$$\operatorname{LIP}(u|_{B_R(p)}) \le \frac{C}{R} \int_{B_{3R}(p)} |u - u(p)| \mathrm{d}\mu, \quad p \in X.$$

Remark 3.1 together with Harnack's inequality (17) imply, in particular, that there are no nonconstant mv-harmonic functions of sublinear growth. A function *u* is said to have sublinear growth, if

$$\limsup_{R \to \infty} \frac{1}{R} \sup_{B_R(p)} |u| = 0$$

for some, and hence any  $p \in X$ .

The following observation can be considered as a counterpart of Cheng's result for harmonic functions with sublinear growth on complete manifolds with nonnegative Ricci curvature, see Corollary 1.5 in [53], and it is also related to the celebrated Phragmen-Lindelöf theorem.

**Corollary 3.3.** Let  $X = (X, d, \mu)$  be a doubling metric measure space. If u is mv-harmonic and has sublinear growth, then it is constant.

In the next section we discuss mean value harmonic functions with prescribed growth in more detail.

3.2. Mean value-harmonic functions of polynomial growth. In [19, 20] Colding and Minicozzi proved a conjecture of Yau on the finite dimensionality of the space of harmonic functions (in the sense of solutions to the Beltrami–Laplace equation) with polynomial growth of degree m. The showed that, in a Riemannian n-manifold M of non-negative Ricci curvature the following bound holds:

$$\dim \mathcal{H}^m(M) \le C(n)m^{n-1}.$$

This result has been extended to Alexandrov and RCD-spaces, cf. [40, 39]. An argument of Li [52] uses the doubling property and the mean value inequality of subharmonic functions to obtain the estimate

$$\dim \mathcal{H}^m(M) \le C(n)m^Q \tag{20}$$

for manifolds with a measure satisfying (8) and a global Poincaré inequality. Thus estimate (20) remains valid in the context of mv-harmonic functions on doubling spaces.

In fact a modification of the same argument improves the bound (20) if  $\mu$  satisfies an annular decay property. We follow the strategy in [52], see Lemmas 3.6 and 3.7, and present the modifications needed for our result.

For m > 0, let  $\mathcal{H}^m(X)$  denote the space of  $u \in \mathcal{H}(X)$  with growth rate at most m. A function  $u \in \mathcal{H}(X)$  is said to have *growth rate at most* m if, for some (and hence all)  $p \in X$ , there exists C > 0 so that

$$|u(x)| \le C(1+d_p(x))^m, \quad x \in X,$$

where  $d_p: X \to \mathbb{R}$  is the distance function  $x \mapsto d(p, x)$ .

**Proposition 3.4.** Let  $X = (X, d, \mu)$  be a complete doubling metric measure space with doubling exponent  $Q := \log_2 C_{\mu} > 1$ , and suppose  $\mu$  has  $\alpha$ -annular decay, cf. (9). Then, for any m > 0, we have that

$$\dim \mathcal{H}^m(X) \le Cm^{Q-\alpha},$$

where the constant  $C = C(Q, \alpha)$  depends only on Q and  $\alpha$ .

Doubling measures on length spaces always satisfy an annular decay property for some  $\alpha$ , see the discussion following Definition 2.1. Thus, Proposition 3.4 implies the following corollary.

**Corollary 3.5.** Let  $(X, d, \mu)$  be a complete geodesic doubling metric measure space with Q > 1. Then there exists  $\delta > 0$ , depending only on the doubling constant of  $\mu$ , so that

$$\dim \mathcal{H}^m(X) \le C(Q)m^{Q-\delta}.$$

Let  $X = (X, d, \mu)$  be a complete doubling metric measure space, where  $\mu$  has  $\alpha$ -annular decay. Given R > 0, we define a bi-linear form

$$A_R(u,v) := \int_{B_R(p)} uv d\mu, \quad u, v \in \mathcal{H}(X),$$

and note that  $A_R$  is symmetric and positive semidefinite. It follows from the proof of [40, Lemma 3.4] that, for any finite dimensional vector subspace  $V \subset \mathcal{H}(X)$ , there exists a radius  $R_0 > 0$  so that  $A_R$  is an inner product on V for every  $R > R_0$ .

In order to prove Proposition 3.4 we need the following auxiliary results.

**Lemma 3.6.** Let V be a k-dimensional linear subspace of  $\mathcal{H}^m(X)$ . For any  $p \in X$ ,  $\beta > 1$ ,  $\delta > 0$ ,  $R_0 > 0$  there exists  $R > R_0$  such that if  $u_1, \ldots, u_k$  is an orthonormal basis for V with respect to the inner product  $A_{\beta R}$ , then

$$\int_{B_R(p)} (u_1^2 + \ldots + u_k^2) \, d\mu \ge \frac{k}{\beta^{2m+Q+\delta}}$$

*Proof.* The proof of [52, Lemma 2] for manifolds carries over to the setting of metric measure spaces under our assumptions. See also [40, Lemma 3.7], where the lemma is proven in the setting of Alexandrov spaces, and [39, Lemma 5.2] for the formulation of the lemma in the  $RCD^*(0, N)$  spaces.

**Lemma 3.7.** Let V be a k-dimensional linear subspace of  $\mathcal{H}^m(X)$ . Then, there exists a constant C = C(Q) such that for any base  $u_1, \ldots, u_k$  of V, any  $p \in X$ , R > 0 and any  $\epsilon \in (0, \frac{1}{2})$  it holds that

$$\int_{B_R(p)} (u_1^2 + \ldots + u_k^2) \, d\mu \le \frac{C}{\epsilon^{Q-\alpha}} \sup_{\sum_{i=1}^k a_i^2 = 1} \int_{B_{(1+\epsilon)R}(p)} |a_1 u_1 + \ldots + a_k u_k|^2 \mathrm{d}\mu$$

*Proof.* The proof of Lemma 3.7 is a modification of the corresponding one of [52, Lemma 1]. We follow closely the proof of Lemma 5.3 in [39]. Fix  $q \in B_r(p)$  and define  $V_q := \{u \in V : u(q) = 0\}$ . The subspace  $V_q \subset V$  is of  $\operatorname{codim} V_q \leq 1$  (since, if  $u, v \notin V_q$ , then  $u - \frac{u(q)}{v(q)}v \in V_q$ ).

There exists an orthogonal change of variables *A* on *V* such that  $A(u_i) := v_i$  for all i = 1, ..., k with  $v_i \in V_q$  for i = 2, ..., k.

We recall the relevant part of Proposition 3.1 in [1] which states that, if  $f \in \mathcal{H}(X)$ , then  $F \circ f$  is subharmonic provided  $F : f(X) \to \mathbb{R}$  is convex. We apply this result for  $f = v_1$  and  $F(s) = s^2$  to obtain that

$$\sum_{i=1}^{k} u_i^2(q) = \sum_{i=1}^{k} v_i^2(q) = v_1^2(q) \le \int_{B_{(1+\epsilon)R-d_p(q)}(q)} v_1^2(z) \, d\mu(z)$$
$$\le \sup_{\substack{a_i \in \mathbb{R}, i=1,\dots,k\\ \sum_i a_i^2 = 1}} \frac{1}{\mu(B_{(1+\epsilon)R-d_p(q)}(q))} \int_{B_{(1+\epsilon)R}(p)} \left| \sum_{i=1}^{k} a_i u_i(z) \right|^2 \, d\mu(z).$$
(21)

We apply (8) to obtain that

$$\frac{1}{\mu(B_{(1+\epsilon)R-d_p(q)}(q))} \le \frac{C_{\mu}^2}{\mu(B_{(1+\epsilon)R}(p))} \left(\frac{(1+\epsilon)R}{(1+\epsilon)R-d_p(q)}\right)^Q.$$

Hence, upon integrating (21), we arrive at

$$\sum_{j=1}^{k} A_{R}(u_{j}, u_{j})$$

$$\leq \frac{C}{\mu(B_{R}(p))} \left( \int_{B_{R}(p)} \left( 1 + \epsilon - \frac{d_{p}(q)}{R} \right)^{-Q} \mathrm{d}\mu(q) \right) \sup_{\sum_{i=1}^{k} a_{i}^{2} = 1} \int_{B_{(1+\epsilon)R}(p)} |a_{1}u_{1} + \ldots + a_{k}u_{k}|^{2} \mathrm{d}\mu.$$
(22)

Denote

$$f: [0,1] \to \mathbb{R}, \quad f(t) = (1 + \epsilon - t)^{-Q},$$

and note that the claim follows directly from (22) and the estimate

$$\oint_{B_R(p)} f(d_p/R) \mathrm{d}\mu \le \frac{C}{\epsilon^{Q-\alpha}}.$$
(23)

To obtain this, note that f is smooth and increasing, and thus

$$\int_{B_R(p)} f(d_p/R) \mathrm{d}\mu = \int_0^\infty \mu(B_R(p) \cap \{f \circ (d_p/R) \ge \lambda\}) \mathrm{d}\lambda = \int_{-\infty}^1 f'(s)\mu(B_R(p) \cap \{d_p \ge sR\}) \mathrm{d}s$$

after a change of variables  $f(s) = \lambda$ . The  $\alpha$ -annular decay implies

$$\mu(B_R(p) \cap \{d_p \ge sR\}) = \mu(A_{sR,R}(p)) \le C(1-s)^{\alpha} \mu(B_R(p))$$

and therefore

$$\int_{B_R(p)} f(d_p/t) \mathrm{d}\mu \leq C \int_{-\infty}^1 f'(s)(1-s)^\alpha \mathrm{d}s \leq CQ \int_{-\infty}^1 (1+\epsilon-s)^{-Q-1+\alpha} \mathrm{d}s$$

$$= \frac{CQ}{Q-\alpha} \frac{1}{\epsilon^{Q-\alpha}},$$

establishing (23).

*Proof of Proposition 3.4.* Let *R* be large enough,  $\beta = 1 + \epsilon$  and  $\epsilon = 1/(2m)$ . Let  $\{u_1, \ldots, u_k\}$  be an orthonormal basis with respect to  $A_{\beta R}$ . Combining the estimates in Lemmas 3.6 and 3.7 we obtain

$$\frac{k}{(1+\epsilon)^{2m+Q+\delta}} \le \sum_{j=1}^{k} A_R(u_j, u_j) \le \frac{C(Q)}{\epsilon^{Q-\alpha}},$$

since

$$\sup_{\sum_{i=1}^{k} a_{i}^{2}=1} \int_{B_{(1+\epsilon)R}(p)} |a_{1}u_{1} + \ldots + a_{k}u_{k}|^{2} \mathrm{d}\mu = 1.$$

Thus

$$k \le C(Q)(1 + 1/(2m))^{2m + Q + \delta}(2m)^{Q - \alpha} \le Cm^{Q - \alpha}$$

after letting  $\delta \rightarrow 0$ .

# 4. HÖLDER CONTINUITY OF STRONGLY AMV-HARMONIC FUNCTIONS

In this section we prove Theorems 1.5 and 1.7. Throughout the section  $X = (X, d, \mu)$  denotes a locally compact and doubling metric measure space with doubling exponent Q. We define the sharp maximal function  $\mathcal{M}^{\#}u$  of a locally integrable function u on X as

$$\mathcal{M}_R^{\#}u(x) := \sup_{0 < r < R} \oint_{B_r(x)} |u - u_{B_r(x)}| \mathrm{d}\mu, \quad x \in X.$$

By standard maximal function estimates we have that, for any compact  $K \subset X$ ,

$$\int_{K} (\mathcal{M}_{R}^{\#} u)^{p} \mathrm{d}\mu \leq C \int_{N_{R}(K)} |u|^{p} \mathrm{d}\mu$$

with a constant *C* depending only on *p* an the local doubling constant of  $\mu$  on *K*, as long as  $R < r_K$  (cf. (7)). We use this observation in the proof of the next proposition without explicit mention.

**Proposition 4.1.** Let  $\Omega \subset X$  be a domain, and  $u \in AMV^p(\Omega)$ . Then  $u \in M^{1/2,p}_{loc}(\Omega)$ . Moreover, if  $u \in M^{\alpha,p}_{loc}(\Omega)$  for some  $\alpha \in (0,1)$ , then  $u \in M^{\alpha',p}_{loc}(\Omega)$ , where

$$\alpha' = \frac{2 - 1/p}{3 - \alpha - 1/p} > \alpha.$$

*Proof.* Let  $K \subset \Omega$  be compact, and define

$$R_K = \frac{1}{6} \min\{\operatorname{dist}(K, X \setminus \Omega)^2, r_K^2, 1\},\$$

where  $r_K$  is given by the locally doubling condition (7). Let  $x, y \in K$  satisfy  $d(x, y) < R_K$ . For any  $r \in (d(x, y), r_K)$  we have that

$$|u(x) - u(y)| \le |u(x) - A^{r}u(x)| + |u(y) - A^{r}u(y)| + |A^{r}u(x) - A^{r}u(y)| \le 2r \int_{0}^{r} (|\Delta_{t}u(x)| + |\Delta_{t}u(y)|) dt + \frac{Cd(x,y)}{r} \int_{B_{3r}(x)} |u - u_{B_{3r}(x)}| d\mu,$$
(24)

where the third term is estimated using Proposition 3.2 with  $c = u_{B_{3r}(x)}$  and the first two terms follow from the estimate

$$\begin{aligned} |u(z) - A^{r}u(z)| &\leq \frac{2}{r} \int_{\frac{r}{2}}^{r} \left| u(z) - \oint_{B_{t}(z)} u(w) \mathrm{d}\mu(w) \right| \mathrm{d}t = \frac{2}{r} \int_{\frac{r}{2}}^{r} t^{2} \left| \Delta_{t}u(z) \right| \mathrm{d}t \\ &\leq 2r \int_{\frac{r}{2}}^{r} \left| \Delta_{t}u(z) \right| \mathrm{d}t \leq 2r \int_{0}^{r} \left| \Delta_{t}u(z) \right| \mathrm{d}t, \quad z = x, y. \end{aligned}$$

By choosing  $r = d(x, y)^{1/2}$  we have that  $d(x, y) < r < r_K$  and thus, by (24), we obtain

$$|u(x) - u(y)| \le d(x, y)^{1/2} [g(x) + g(y)],$$

where

$$g(x) = 2 \int_0^{r_K} |\Delta_t u(x)| \mathrm{d}t + C \mathcal{M}_{3r_K}^{\#} u(x).$$

This proves the first claim.

Suppose  $u \in M_{loc}^{\alpha,p}(\Omega)$ , and let  $g_K$  be the Hajłasz gradient and  $\tilde{r}_K$  the radius in Definition 2.2. Set

$$R_K = \frac{1}{6} \min\{\tilde{r}_K, \operatorname{dist}(K, X \setminus \Omega)^{3-\alpha-1/p}, r_K^2\}.$$

From (24) we obtain that as long as  $6r < \tilde{r}_K$  and  $d(x, y) < R_K$ , then

$$|u(x) - u(y)| \le Cr^{2-1/p} \left( \int_0^r (|\Delta_t u(x)|^p + |\Delta_t u(y)|^p) dt \right)^{1/p} + C \frac{d(x,y)}{r^{1-\alpha}} [A_{6r}g_K(x) + A_{6r}g_K(y)].$$
(25)

Indeed, applying the Hölder inequality to the first term in the right-hand side of (24) we obtain, up to a multiplicative constant, that

$$r \int_0^r |\Delta_t u(x)| \, \mathrm{d}t \le r \left( \int_0^r |\Delta_t u(x)|^p \, \mathrm{d}t \right)^{\frac{1}{p}} r^{\frac{p-1}{p}} = r^{2-\frac{1}{p}} \left( \int_0^r |\Delta_t u(x)|^p \, \mathrm{d}t \right)^{\frac{1}{p}}.$$

The second term in (24) is estimated using the Hajłasz inequality, up to a multiplicative constant, in the following way

$$\begin{aligned} \frac{d(x,y)}{r} \oint_{B_{3r}} |u - u_{B_{3r}(x)}| \mathrm{d}\mu &= \frac{d(x,y)}{r} \oint_{B_{3r}(x)} \int_{B_{3r}(x)} u(x) - \int_{B_{3r}(x)} u(z) \mathrm{d}\mu(x) \left| \mathrm{d}\mu(w) \right| \\ &\leq \frac{d(x,y)}{r} \oint_{B_{3r}(x)} \int_{B_{3r}(x)} \int_{B_{3r}(x)} u(w) - u(z) |\mathrm{d}\mu(z) \mathrm{d}\mu(w) \\ &\leq \frac{d(x,y)}{r} \int_{B_{3r}(x)} \int_{B_{3r}(x)} d(w,z)^{\alpha} (g_K(w) + g_K(z)) \mathrm{d}\mu(z) \mathrm{d}\mu(w) \\ &\leq \frac{d(x,y)}{r} (6r)^{\alpha} \int_{B_{3r}(x)} \int_{B_{3r}(x)} (g_K(w) + g_K(z)) \mathrm{d}\mu(z) \mathrm{d}\mu(w) \\ &\leq C \frac{d(x,y)}{r^{1-\alpha}} [A_{6r}g_K(x) + A_{6r}g_K(y)]. \end{aligned}$$

We choose r in (25) such that

$$r^{2-1/p} = \frac{d(x,y)}{r^{1-\alpha}},$$

i.e.

$$r = d(x, y)^{1/(3-\alpha-1/p)} < r_K^{1/(3-\alpha-1/p)} < \tilde{r}_K,$$

to obtain

$$|u(x) - u(y)| \le d(x, y)^{\alpha'} [g(x) + g(y)]$$

where

$$\alpha' = \frac{2 - 1/p}{3 - \alpha - 1/p}$$

and

$$g(x) = C\left(\int_0^{\tilde{r}_K} |\Delta_t u(x)|^p \mathrm{d}t\right)^{1/p} + C\mathcal{M}_{\tilde{r}_K}g_K(x)$$

г	
∟	_

Theorem 1.7 follows immediately from Proposition 4.1.

*Proof of Theorem* 1.7. Define  $\alpha_0 = 1/2$  and

$$\alpha_{n+1} = \frac{2 - 1/p}{3 - \alpha_n - 1/p}, \quad n \ge 0.$$

We see that  $\alpha_n$  forms an increasing sequence converging to 1. By Proposition 4.1 we have that  $u \in M_{loc}^{\alpha_k,p}(\Omega)$  for every k. The claim follows.

In light of the fractional Morrey embedding (Proposition 2.1), Theorem 1.7 immediately yields the following corollary, which also contains Theorem 1.5.

**Corollary 4.2.** Let  $\Omega \subset X$  be a domain in a doubling metric measure space with doubling exponent Q. If p > Q and  $u \in AMV_{loc}^{p}(\Omega)$ , then u is locally  $\beta$ -Hölder continuous for every  $\beta < 1 - Q/p$ . In particular a strongly anv-harmonic function u on  $\Omega$  is  $\beta$ -Hölder continuous for any  $\beta < 1$ .

*Proof.* Given  $\beta < 1 - Q/p$ , choose  $\alpha < 1$  which satisfies  $\alpha p > Q$  and  $\beta < \alpha - Q/p$ . By Proposition 2.1 u is locally  $(\alpha - Q/p)$ -Hölder continuous, and thus locally  $\beta$ -Hölder continuous.

If u is strongly any-harmonic on  $\Omega$ , then in particular  $u \in AMV_{loc}^{p}(\Omega)$  for every  $p < \infty$ . This finishes the proof.

### 5. BLOW-UPS OF HAJŁASZ–SOBOLEV FUNCTIONS WITH FINITE AMV-NORM

To study blow-ups of functions with finite amv-norm, we first review pointed measured Gromov– Hausdorff convergence (pmGH-convergence) of spaces and functions. In the literature there are several variants of pmGH-convergence. Here we follow the presentation of [43, Section 2], and refer the interested reader to [43, 38, 23] for more discussion and the relationship between the various notions.

5.1. Pointed measured Gromov-Hausdorff convergence. Let  $F_m \subset Z$  for m = 1, 2, ... and  $F \subset Z$  be closed sets in a metric space Z. We say that a sequence  $(F_m)$  Hausdorff-converges to F, denoted  $F_m \to F$ , if

$$\lim_{m \to \infty} \sup_{z \in F_m \cap B_R(q)} \operatorname{dist}_Z(z, F) = 0 \text{ and } \lim_{m \to \infty} \sup_{z \in F \cap B_R(q)} \operatorname{dist}_Z(z, F_m) = 0$$

for every  $q \in Z$  and R > 0. If  $\nu_m$ , for m = 1, 2, ... and  $\nu$  are Radon measures on Z, we say that a sequence  $(\nu_m)$  converges to  $\nu$  weakly, denoted  $\nu_m \rightharpoonup \nu$ , if

$$\lim_{m \to \infty} \int_Z \varphi \mathrm{d}\nu_m = \int_Z \varphi \mathrm{d}\nu$$

for every continuous function  $\varphi : Z \to \mathbb{R}$  with bounded support.

**Lemma 5.1.** Let  $(\nu_m)$  be a sequence of measures on Z converging weakly to  $\nu$ , and suppose  $(F_m)$  is a sequence of compact sets Hausdorff-converging to a compact set F. Then

$$\limsup_{m \to \infty} \nu_m(F_m) \le \nu(F) \le \liminf_{m \to \infty} \nu_m(N_{\varepsilon}(F_m))$$

for any  $\varepsilon > 0$ .

*Proof.* Given  $\varepsilon > 0$  there exists  $m_0$  such that  $F \subset N_{\varepsilon}(F_m)$  and  $F_m \subset N_{\varepsilon}(F)$ , whenever  $m \ge m_0$ . By [38, Remark 11.4.1] we obtain

$$\limsup_{m \to \infty} \nu_m(F_m) \le \limsup_{m \to \infty} \nu(\bar{N}_{\varepsilon}(F)) \le \nu(\bar{N}_{\varepsilon}(F)).$$

Taking infimum over  $\varepsilon > 0$  we obtain the left-hand side inequality in the assertion. Similarly, to see the second inequality we use [38, Remark 11.4.1] and estimate

$$\nu(F) \le \nu(N_{\varepsilon}(F)) \le \liminf_{m \to \infty} \nu_m(N_{\varepsilon}(F)) \le \liminf_{m \to \infty} \nu_m(N_{2\varepsilon}(F_m)).$$

For the next definition, we recall that a pointed metric measure space  $X = (X, d, \mu, p)$  consists of a metric measure space  $(X, d, \mu)$  and a distinguished point  $p \in X$ . We consider only proper spaces here.

**Definition 5.1.** A sequence  $(X_m) = (X_m, d_m, \mu_m, p_m)$  of pointed proper metric measure spaces *pmGH*converges to a pointed proper metric measure space  $X = (X, d, \mu, p)$ , denoted  $X_m \xrightarrow{pmGH} X$ , if there exists a pointed proper metric space (Z, q) and isometric embeddings  $\iota_m : X_m \to Z$  for m = 1, 2, ...and  $\iota : X \to Z$  so that

- (1)  $\iota_m(p_m) = \iota(p) = q$  for all *m*, and  $\iota_m(X_m)$  Hausdorff-converges to  $\iota(X)$ ;
- (2) the measures  $\iota_{m*}\mu_m$  weakly converge to  $\iota_*\mu$ , as  $m \to \infty$ .

We also define Gromov–Hausdorff convergence for sequences of functions. Note that in Definition 5.2, the measures play no role. Since we consider pointed measured spaces, we nevertheless include them (see also Definition 5.3).

**Definition 5.2.** Let  $u_m : X_m \to \mathbb{R}$  for m = 1, 2, ... and  $u : X \to \mathbb{R}$  be functions on pointed proper metric measure spaces. We say that the sequence  $(u_m)$  *Gromov–Hausdorff converges to u*, denoted  $u_m \xrightarrow{GH} u$ , if there are isometric embeddings  $\iota_m : X_m \to Z \ m = 1, 2, ...$  and  $\iota : X \to Z$  satisfying (1) and (2) in Definition 5.1 and

(3) 
$$u_m(z_m) \to u(z)$$
 whenever  $z_m \in X_m$ ,  $z \in X$ , and  $\iota_m(z_m) \to \iota(z)$ , as  $m \to \infty$ .

The embeddings  $\iota_m : X_m \to Z$  and  $\iota : X \to Z$  satisfying (1) and (2) (resp. (3)) in Definition 5.1 are said to *realize* the convergence  $X_m \xrightarrow{pmGH} X$  (resp.  $u_m \xrightarrow{GH} u$ ).

Two central properties of Gromov–Hausdorff convergence are its compactness properties (see Proposition 5.2 below) and the stability of many important properties in metric geometry and analysis under the Gromov–Hausdorff convergence. For our purposes, the stability of length spaces and the doubling property of the measure is important; for a detailed discussion see [38, Section 11] and [43].

The following compactness property is proved in [43, Proposition 3], see also [38]. For a pointed metric space (X, d, p) and  $\varepsilon, R > 0$ , let  $N_X(\varepsilon, R)$  denote the maximal number of disjoint closed balls of radius  $\varepsilon$  inside a ball  $B_R(p)$ . We say that a sequence  $(X_m, d_m, p_m)$  of pointed metric spaces is *totally bounded* if

$$\sup_{m} N_{X_m}(\varepsilon, R) < \infty$$

for every  $\varepsilon, R > 0$ .

**Proposition 5.2** (cf. Proposition 3, [44]). Let  $(X_m) = (X_m, d_m, \mu_m, p_m)$  be a totally bounded sequence of proper metric measure spaces, satisfying

$$\sup_{m} \mu_m(B_R(p_m)) < \infty \quad \text{for every } R > 0.$$
(26)

Then there exists a subsequence and a pointed proper metric measure space  $X = (X, d, \mu, p)$  so that  $X_m \xrightarrow{pmGH} X$ .

There is also a compactness result for sequences of functions on pointed spaces, which can be proved using Proposition 5.2 and a diagonal argument, as in the proof of the Arzela–Ascoli theorem. Under a different notion of convergence (which is equivalent to ours under the hypotheses there), Proposition 5.3 appears in [44]. For the additional statement (2), see [38, Section 11] and [43].

To state the result, let  $(X_m) = (X_m, d_m, \mu_m, p_m)$  be a sequence of pointed proper metric measure spaces. We say that a sequence  $(f_m)$  of functions  $f_m : X_m \to \mathbb{R}$  for m = 1, 2, ... is *equicontinuous* if, for every  $\varepsilon, R > 0$ , there exists  $\delta > 0$  such that if  $x_m, y_m \in B_R(p_m)$  satisfy  $d(x_m, y_m) < \delta$ , then  $|f_m(x_m) - f_m(y_m)| < \varepsilon$  uniformly in m.

**Proposition 5.3.** Let  $(X_m) = (X_m, d_m, \mu_m, p_m)$  be a totally bounded sequence of pointed proper metric measure spaces satisfying (26). If a sequence of functions  $(f_m)$ , where  $f_m : X_m \to \mathbb{R}$  for m = 1, 2, ..., is an equicontinuous sequence of functions, for which

$$\sup|f_m(p_m)| < \infty,$$

then there exists a subsequence of  $(f_m)$  and a continuous function  $f : X \to \mathbb{R}$ , defined on a proper pointed metric measure space X, for which  $f_m \xrightarrow{GH} f$ .

Moreover,

- (1) if each  $f_m$  is L-Lipschitz, then f is L-Lipschitz;
- (2) if each  $X_m$  is a doubling length space with doubling constants not exceeding a positive constant C, then X is a length space with doubling constant not exceeding  $C^2$ .

To study tangents of Hajłasz–Sobolev functions, we also consider a notion of weak convergence for functions. The following definition is a slight modification of the weak convergence in [23].

**Definition 5.3.** Let  $(X_m) = (X_m, d_m, \mu_m, p_m)$  and  $X = (X, d, \mu, p)$  be pointed proper metric measure spaces. A sequence  $(u_m)$  of functions  $u_m \in L^1_{loc}(X_m)$  converges *weakly* to  $u \in L^1_{loc}(X)$ , denoted  $u_m \stackrel{GH}{\rightharpoonup} u$ , if there exist isometric embeddings  $\iota_m : X_m \to Z$ ,  $\iota : X \to Z$  satisfying (1) and (2) in Definition 5.1, and for which

(3')  $\iota_{m*}((u_m)+d\mu_m) \rightharpoonup \iota_*(u_+d\mu)$  and  $\iota_{m*}((u_m)-d\mu_m) \rightharpoonup \iota_*(u_-d\mu)$ , as  $m \to \infty$ . Here, for a function  $f: Z \to \mathbb{R}$  we denote

$$f_+ = \max\{f, 0\}$$
 and  $f_- = -\min\{f, 0\}$ .

The Gromov-Hausdorff convergence of functions is analogous to the uniform convergence on compact sets and indeed coincides with this notion if  $X_m = X = Z$  for all m. The weak convergence of functions as in Definition 5.3 corresponds to weak convergence of signed measures. In keeping with these analogies, we indeed have the natural implication between the two notions.

**Lemma 5.4.** Let metric measure spaces  $(X_m)$  and X be as in Definition 5.3. Suppose further, that a sequence  $(u_m)$  of functions  $u_m \in L^1_{loc}(X_m)$  Gromov–Hausdorff converges to a continuous function  $u : X \to \mathbb{R}$ . Then  $(u_m)$  converges to u weakly.

More precisely, if  $\iota_m : X_m \to Z$ ,  $\iota : X \to Z$  realize the convergence  $u_m \xrightarrow{GH} u$ , then

$$\lim_{m \to \infty} \int_{X_m} (\varphi \circ \iota_m)(u_m)_{\pm} \mathrm{d}\mu_m = \int_X (\varphi \circ \iota) u_{\pm} \mathrm{d}\mu$$

for any boundedly supported continuous  $\varphi: Z \to \mathbb{R}$ .

*Proof.* It is easy to see that if  $u_m \xrightarrow{GH} u$ , then  $(u_m)_{\pm} \xrightarrow{GH} u_{\pm}$ , and the embeddings realizing the first convergence also realize the latter convergence. Thus we may assume that  $u_m$  is non-negative for all m, and the embeddings  $\iota_m : X_m \to Z$ ,  $\iota : X \to Z$  realize the convergence  $u_m \xrightarrow{GH} u$ . It suffices to show that  $\iota_{m*}(u_m d\mu) \to \iota_*(u d\mu)$ .

Let  $\tilde{u}: Z \to \mathbb{R}$  be a continuous extension of  $u \circ \iota^{-1}: \iota(X) \to \mathbb{R}$ , and set

$$\tilde{u}_m = u_m \circ \iota_m^{-1}|_{\iota_m(X_m)}$$

Given any continuous  $\varphi : Z \to \mathbb{R}$  with bounded (thus compact) support, we have

$$\int_{X_m} (\varphi \circ \iota_m) u_m \mathrm{d}\mu_m - \int_X (\varphi \circ \iota) u \, \mathrm{d}\mu = \int_Z \varphi \left( \tilde{u}_m - \tilde{u} \right) \iota_{m*}(\mathrm{d}\mu_m) + \int_Z \varphi \, \tilde{u} \, \iota_{m*}(\mathrm{d}\mu_m) - \int_Z \varphi \, \tilde{u} \, \iota_*(\mathrm{d}\mu).$$

Since  $\iota_{m*}(d\mu_m) \rightharpoonup \iota_*(d\mu)$ , it suffices to prove that

$$\int_{Z} \varphi \left( \tilde{u}_m - \tilde{u} \right) \iota_{m*}(\mathrm{d}\mu_m) \to 0$$

If  $B \subset Z$  is a closed ball containing the support of  $\varphi$ , we obtain

$$\begin{split} \limsup_{m \to \infty} \left| \int_{Z} \varphi \left( \tilde{u}_{m} - \tilde{u} \right) \iota_{m*}(\mathrm{d}\mu_{m}) \right| &\leq \limsup_{m \to \infty} \|\varphi\|_{L^{\infty}(B)} \mu_{m}(\iota_{m}^{-1}B) \|\tilde{u}_{m} - \tilde{u}\|_{L^{\infty}(\iota_{m}(X_{m}) \cap B)} \\ &\leq \|\varphi\|_{L^{\infty}(B)} \mu(\iota^{-1}B) \limsup_{m \to \infty} \|\tilde{u}_{m} - \tilde{u}\|_{L^{\infty}(\iota_{m}(X_{m}) \cap B)}. \end{split}$$

Suppose  $\limsup_{m\to\infty} \|\tilde{u}_m - \tilde{u}\|_{L^{\infty}(\iota_m(X_m)\cap B)} > \varepsilon_0$  for some  $\varepsilon_0 > 0$ . Then, there is a sequence  $(x_m)$  such that  $x_m \in X_m$  with the corresponding sequence  $(z_m)$  defined  $z_m := \iota_m(x_m) \in B$  and

$$|\tilde{u}_m(z_m) - \tilde{u}(z_m)| = |u_m(x_m) - \tilde{u}(\iota_m(x_m))| > \varepsilon_0$$

for *m* large enough. Since *B* is compact, a subsequence of  $(z_m)$ , denoted the same for simplicity, satisfies  $z_m \to z \in \iota(X)$  for some *z*, as  $m \to \infty$ . By the Gromov–Haudorff convergence  $u_m \xrightarrow{GH} u$  and the continuity of  $\tilde{u}$ , we obtain

$$\lim_{m \to \infty} |u_m(x_m) - \tilde{u}(\iota_m(x_m))| = |u(\iota^{-1}(z)) - \tilde{u}(z)| = 0,$$

which is a contradiction. This completes the proof.

5.2. Gromov–Hausdorff convergence and averaging operators. In this section we show the stability of operators  $A_r$  and  $A_r^*$  under the Gromov-Hausdorff convergence.

**Proposition 5.5.** Let  $(X_m) = (X_m, d_m, \mu_m, p_m)$  and  $X = (X, d, \mu, p)$  be proper locally doubling length spaces. Suppose the sequence  $(u_m)$  of functions  $u_m \in L^1_{loc}(X_m)$  converges weakly to  $u \in L^1_{loc}(X)$ , and let  $\iota_m : X_m \to Z$  for m = 1, 2, ... and  $\iota : X \to Z$  realize this convergence. If  $z_m \in X_m$  for m = 1, 2, ... and  $z \in X$  are such that  $\iota_m(z_m) \to \iota(z)$ , as  $m \to \infty$ , then

$$\lim_{m \to \infty} \int_{B_r(z_m)} u_m \mathrm{d}\mu_m = \int_{B_r(z)} u \mathrm{d}\mu$$

for any r > 0.

*Proof.* By considering functions  $u_{m\pm}$  for  $m \ge 1$ , we may assume without the loss of generality that all the functions  $u_m$  and u are non-negative. We note that

$$\iota_m(B_r(z_m)) \to \iota(B_r(z)), \quad m \to \infty$$
(27)

for every r > 0 in the sense of Hausdorff-convergence. Indeed, given  $q \in Z$ , R > 0 and arbitrary  $\varepsilon > 0$  the convergence  $\iota_m(X_m) \to \iota(X)$  and  $\iota_m(z_m) \to \iota(z)$  for  $m \to \infty$  imply that, for large enough m

$$B_R(q) \cap \iota_m(B_r(z_m)) = B_R(p) \cap \iota_m(X_m) \cap B_r(\iota_m(z_m)) \subset N_\varepsilon(\iota(X)) \cap B_{r+\varepsilon}(\iota(z)) = N_\varepsilon(\iota(B_r(z)))$$

and

$$B_R(q) \cap \iota(B_r(z)) = B_R(p) \cap \iota(X) \cap B_r(\iota(z))) \subset N_{\varepsilon}(\iota_m(X_m)) \cap B_{r+\varepsilon}(\iota_m(z_m)) = N_{\varepsilon}(\iota(B_r(z_m))).$$

The convergence (27) follows. Note that the property of being a length space was used in the last equality, cf. [38, Lemma 11.3.10]. By Lemma 5.1 we have, for any  $\varepsilon > 0$ , that

$$\limsup_{m \to \infty} \nu_m(\bar{B}_r(z_m)) \le \nu(\bar{B}_r(z)) \le \liminf_{m \to \infty} \nu_m(B_{\varepsilon+r}(z_m))$$

where  $d\nu = ud\mu$  and  $\nu_m = u_m d\mu_m$ . Recall from Section 3.1 notation  $A_{r,R}(x) = \overline{B}_R(x) \setminus B_r(x)$  for  $x \in \mathbb{Z}$  and  $r \leq R$ . It holds that

$$\liminf_{m \to \infty} \nu_m(B_{\varepsilon+r}(z_m)) = \liminf_{m \to \infty} [\nu_m(\bar{B}_r(z_m)) + \nu_m(A_{r,r+\varepsilon}(z_m))]$$
  
$$\leq \liminf_{m \to \infty} \nu_m(B_r(z_m)) + \limsup_{m \to \infty} \nu_m(A_{r,r+\varepsilon}(z_m)).$$

The argument used to establish (27) also yields that

$$A_{r,r+\varepsilon}(z_m) \to A_{r,r+\varepsilon}(z)$$

in the sense of Hausdorff-convergence. Applying Lemma 5.1 once more we obtain

$$\limsup_{m \to \infty} \nu_m(\bar{B}_r(z_m)) \le \nu(\bar{B}_r(z)) \le \liminf_{m \to \infty} \nu_m(B_r(z_m)) + \nu(A_{r,r+\varepsilon}(z)).$$

Since  $\mu$  is a locally doubling measure on a length space, it has an annular decay property (see the discussion after Definition 2.1). Therefore,  $\lim_{\varepsilon \to 0} \mu(A_{r,r+\varepsilon}(z)) = 0$  which, by the absolute continuity of  $\nu$  with respect to  $\mu$ , implies  $\lim_{\varepsilon \to 0} \nu(A_{r,r+\varepsilon}(z)) = 0$ . We have obtained

$$\limsup_{m \to \infty} \nu_m(B_r(z_m)) \le \nu(B_r(z)) \le \liminf_{m \to \infty} \nu_m(B_r(z_m))$$

which completes the proof.

25

Proposition 5.5 has the following immediate corollary.

**Corollary 5.6.** Let  $(X_m) = (X_m, d_m, \mu_m, p_m)$  and  $X = (X, d, \mu, p)$  be proper locally doubling length spaces, and  $(u_m)$  be a sequence of functions  $u_m \in L^1_{loc}(X_m)$  converging weakly to  $u \in L^1_{loc}(X)$ . Then

(a)  $A_r^{\mu_m} u_m \xrightarrow{GH} A_r^{\mu} u$ , and (b)  $(A_r^{\mu_m})^* u_m \xrightarrow{GH} (A_r^{\mu})^* u$ 

for each r > 0. In particular, if u is continuous, then

 $\Delta^{\mu_m}_r u_m \stackrel{GH}{\rightharpoonup} \Delta^{\mu}_r u \quad and \quad (\Delta^{\mu_m}_r)^* u_m \stackrel{GH}{\rightharpoonup} (\Delta^{\mu}_r)^* u.$ 

*Proof.* The first claim follows directly from Proposition 5.5. Note that, if  $\iota_m : X_m \to Z$  and  $\iota : X \to Z$  realize the convergence then  $X_m \xrightarrow{pmGH} X$ , Proposition 5.5 implies, in particular, that

$$\mu_m(B_r(z_m)) \to \mu(B_r(z))$$
 whenever  $\iota_m(z_m) \to \iota(z)$ 

It follows that, for any r > 0, the sequence  $(f_m)$  of functions  $f_m \in L^1_{loc}(X_m)$ ,

$$f_m(z) = \frac{u_m(z)}{\mu_m(B_r(z))}$$

weakly converges to

$$f(z) := \frac{u(z)}{\mu(B_r(z))}.$$

This yields part (b) of the assertion.

5.3. Blow-ups of Sobolev functions with finite amv-norm. Consider a proper locally doubling length space  $X = (X, d, \mu)$ . Given a point  $x \in X$  and r > 0, the pointed metric measure space

$$X_r = (X, d_r, \mu_r, x), \quad d_r := \frac{d}{r}, \quad \mu_r := \frac{1}{\mu(B(x, r))}\mu$$

is called a *rescaling* of X (at x by r).

Let  $(r_m)$  be a sequence of positive numbers converging to zero, and denote  $X_m := X_{r_m}$ . A pointed measured Gromov–Hausdorff limit  $X_{\infty}$  of  $X_m$  is called a *tangent space of* X *at* x *subordinate to*  $(r_m)$ .

Similarly, if  $f : X \to \mathbb{R}$  is a function,  $x \in X$  and r > 0, then a function

$$f_r := \frac{f - f(x)}{r} : X_r \to \mathbb{R}$$

is a rescaling of f at x by r. A Gromov–Hausdorff limit  $f_{\infty} : X_{\infty} \to \mathbb{R}$  of a sequence  $(f_m)$ , where  $f_m := f_{r_m}$  for m = 1, 2, ... is called a *tangent of* f at x subordinate to  $(r_m)$ . If the convergence  $f_m \to f_{\infty}$  as  $m \to \infty$  is weak (cf. Definition 5.3), then we say that  $f_{\infty}$  is an *approximate tangent of* f at x, subordinate to  $(r_m)$ .

It is worth remarking that, in general, tangents are highly non-unique – different sequences can produce different limits. However, any sequence of rescalings is totally bounded and satifies (26). Moreover,  $f_m(x) = 0$  for all m, and thus Proposition 5.3 implies the existence of tangents of Lipschitz functions at any point.

**Proposition 5.7.** Let X be a proper locally doubling metric measure space and  $f : X \to \mathbb{R}$  an L-Lipschitz function. Fix a sequence  $(r_m)$  of positive numbers converging to zero. Then, for any  $x \in X$ , there exists a subsequence of the rescalings  $f_m : X_m \to \mathbb{R}$  at x, and a tangent function  $f_\infty : X_\infty \to \mathbb{R}$  such that  $f_m \xrightarrow{GH} f_\infty$ .

In particular, any tangent space  $X_{\infty}$  is doubling, and any tangent  $f_{\infty} : X_{\infty} \to \mathbb{R}$  is *L*-Lipschitz. Next, we present a variant for Hajłasz–Sobolev functions.

26

**Proposition 5.8.** Let X be a proper locally doubling metric measure space, p > 1, and  $u \in M^{1,p}(X)$ . Given a sequence  $r_m$ , for  $\mu$ -almost every point  $x \in X$ , there is a subsequence of the rescalings  $u_m : X_m \to \mathbb{R}$  and a Lipschitz function  $u_\infty : X_\infty \to \mathbb{R}$ , so that  $u_m \stackrel{GH}{=} u_\infty$ .

It can be shown that the Lipschitz constant of  $u_{\infty}$  satisfies the following estimates for  $g_u$  the minimal upper gradient of u

$$\frac{1}{C}g_u(x) \le \operatorname{LIP}(u_\infty) \le Cg_u(x)$$

for a constant depending only on the local doubling constant of  $\mu$  near x.

*Proof of Proposition 5.8.* Let  $g \in L^p(X)$  be a Hajłasz upper gradient of u, and set

$$E_n = \{ x \in X : g(x) > n \}, \quad n \in \mathbb{N}.$$

Then

$$\lim_{n \to \infty} \mu(E_n) = 0 \quad \text{and} \quad \lim_{n \to \infty} \int_{E_n} g d\mu = 0$$

Hence, there exists a null set  $N \subset X$  for which every  $x \in X \setminus N$  has the property that

$$\lim_{r \to 0} \frac{\mu(B_r(x) \cap E_n)}{\mu(B_r(x))} = 0 \quad \text{and} \quad \lim_{r \to 0} \frac{1}{\mu(B_r(x))} \int_{B_r(x) \cap E_n} g(y) d\mu(y) = 0$$
(28)

for some  $n \in \mathbb{N}$ . We fix  $x \in X \setminus N$  and  $n \in \mathbb{N}$  satisfying (28).

Note that  $u|_{X \setminus E_n}$  is 2*n*-Lipschitz and let  $\tilde{u}_n : X \to \mathbb{R}$  be a 2*n*-Lipschitz extension of  $u|_{X \setminus E_n}$ . By Proposition 5.7 there is a subsequence of the rescalings  $(\tilde{u}_n)_m : X_m \to \mathbb{R}$  and a Lipschitz function  $\tilde{u} : X_\infty \to \mathbb{R}$  so that  $(\tilde{u}_n)_m \xrightarrow{GH} \tilde{u}$ .

 $\tilde{u}: X_{\infty} \to \mathbb{R}$  so that  $(\tilde{u}_n)_m \xrightarrow{GH} \tilde{u}$ . We show that, for this subsequence, the rescalings  $u_m: X_m \to \mathbb{R}$  converge weakly to  $\tilde{u}$ . (This is different from claiming that  $(\tilde{u}_n)_m \xrightarrow{GH} \tilde{u}$ , which follows from Lemma 5.4.)

Let *Z* be a proper metric space and  $\iota_m : X_m \to Z$  isometric embeddings realizing the convergence  $(\tilde{u}_n)_m \to \tilde{u}$ . Given  $\varphi \in C_b(Z)$ , fix a large number R > 0 so that spt  $\varphi \subset B_R(\iota_m(x))$  for all  $m \in \mathbb{N}$ . Then  $\operatorname{spt}(\varphi \circ \iota_m) \subset B_R^{X_m}(x) = B_{r_mR}(x)$ . We have

$$\begin{aligned} \left| \int_{X_m} \varphi \circ \iota_m \, u_m \mathrm{d}\mu_m - \int_{X_\infty} \varphi \circ \iota_\infty \, \tilde{u} \mathrm{d}\mu_\infty \right| \\ &= \left| \int_X \varphi \circ \iota_m \, (\tilde{u}_n)_m \mathrm{d}\mu_m - \int_{X_\infty} \varphi \circ \iota_\infty \, \tilde{u} \mathrm{d}\mu_\infty + \int_{E_n} \varphi \circ \iota_m \, [u_m - (\tilde{u}_n)_m] \mathrm{d}\mu_m \right| \\ &\leq \left| \int_X \varphi \circ \iota_m \, (\tilde{u}_n)_m \mathrm{d}\mu_m - \int_{X_\infty} \varphi \circ \iota_\infty \, \tilde{u} \mathrm{d}\mu_\infty \right| + \int_{E_n} |\varphi \circ \iota_m \, u_m| \mathrm{d}\mu_m + \int_{E_n} |\varphi \circ \iota_m \, (\tilde{u}_n)_m| \mathrm{d}\mu_m \end{aligned}$$

The first term converges to zero since  $(\tilde{u}_n)_m \xrightarrow{GH} \tilde{u}$ . We may estimate the second term by

$$\int_{E_n} |\varphi \circ \iota_m| \, |u_m| \mathrm{d}\mu_m \le \frac{\|\varphi\|_{\infty}}{\mu(B_{r_m}(x))} \int_{E_n \cap B_{r_mR}(x)} (g(x) + g(y)) \mathrm{d}\mu(y)$$

which, by (28) converges to zero as  $m \to \infty$ . Similarly,

$$\int_{E_n} |\varphi \circ \iota_m| \, |(\tilde{u}_n)_m| \mathrm{d}\mu_m \le 2n \|\varphi\|_{\infty} \frac{\mu(E_n \cap B_{r_m R}(x))}{\mu(B_{r_m}(x))}$$

converges to zero as  $m \to \infty$ . This completes the proof.

By a suitable cut-off argument, we obtain the following corollary whose proof we omit.

**Corollary 5.9.** Let  $\Omega \subset X$  be a domain in a proper locally doubling metric measure space, and let  $u \in M_{loc}^{1,p}(\Omega)$ . Given a sequence  $r_m \downarrow 0$ , for  $\mu$ -almost every  $x \in \Omega$ , there is a subsequence of the rescalings  $u_m : X_m \to \mathbb{R}$  at x, and Lipschitz function  $u_\infty : X_\infty \to \mathbb{R}$ , so that  $u_m \stackrel{GH}{\rightharpoonup} u_\infty$ .

We are now ready to prove that having finite amv-norm forces tangent maps to be mean value harmonic.

*Proof of Theorem 1.6.* By Corollary 5.9, u has approximate tangent maps, subordinate to  $(r_m)$ , for  $\mu$ -almost every  $x \in X$ . For any  $k \in \mathbb{N}$ ,  $r \in \mathbb{Q}_+$ , and compact  $K \subset \Omega$ , we have

$$\int_{K} \liminf_{m \to \infty} \oint_{B(x,kr_m)} |\Delta_{r_m r} u| d\mu \, d\mu(x) \le \liminf_{m \to \infty} \int_{K} \oint_{B(x,kr_m)} |\Delta_{r_m r} u| d\mu \, d\mu(x) \le C \limsup_{\rho \to 0} \int_{K} |\Delta_{\rho} u| d\mu,$$

cf. [3, Theorem 3.3]. Thus, almost every  $x_0 \in X$  has the following property: for every subsequence of  $(r_m)$  there exists a further subsequence (not relabeled), for which

$$\lim_{m \to \infty} \int_{B(x_0, kr_m)} |\Delta_{r_m r} u| d\mu < \infty \quad \text{for every } k \in \mathbb{N} \text{ and } r \in \mathbb{Q}_+.$$
(29)

Let  $x_0 \in \Omega$  be a point where (29) and the claim of Corollary 5.9 holds for  $(r_m)$ . Furthermore, let  $(X_{\infty}, d_{\infty}, \mu_{\infty}, x_{\infty})$  be a pointed measured Gromov–Hausdorff limit of a sequence

$$X_m = (X, d_m, \mu_m, x_0) = \left(X, \frac{d}{r_n}, \frac{\mu}{\mu(B_{r_m}(x_0))}, x_0\right),$$

and  $u_{\infty}: X_{\infty} \to \mathbb{R}$  a weak limit of the sequence

$$u_m := \frac{u - u(x_0)}{r_m} : X_m \to \mathbb{R}.$$

for a subsequence of  $(r_m)$ . We pass to a further subsequence (again not relabeled) for which (29) holds. Note that

$$A_r^{\mu_m} u_m(z) - u_m(z) = \oint_{B_{r_m r}(z)} \frac{u(y) - u(z)}{r_m} d\mu(y) = r_m r^2 \Delta_{r_m r} u(z)$$

for any  $z \in X_m$  and r > 0. Let  $\iota_m : X_m \to Z$  and  $\iota : X_\infty \to Z$  realize the convergence  $u_m \stackrel{GH}{\rightharpoonup} u_\infty$ . Corollary 5.6 implies that

$$\begin{split} \int_{X_{\infty}} \varphi \circ \iota_{\infty}(z) \left[ A_{r}^{\mu_{\infty}} u_{\infty}(z) - u_{\infty}(z) \right] \mathrm{d}\mu_{\infty}(z) &= \lim_{m \to \infty} \int_{X_{m}} \varphi \circ \iota_{m}(z) \left[ A_{r}^{\mu_{m}} u_{\infty}(z) - u_{m}(z) \right] \mathrm{d}\mu_{m}(z) \\ &= \lim_{m \to \infty} r_{m} r^{2} \int_{X_{m}} \varphi \circ \iota_{m}(z) \Delta_{r_{m}r} u(z) \mathrm{d}\mu_{m}(z) \end{split}$$

for every compactly supported  $\varphi \in C(Z)$ . Fix such  $\varphi \in C_c(Z)$ , and not that for large enough k, the support of  $\varphi \circ \iota_m$  satisfies  $\operatorname{spt}(\varphi \circ \iota_m) \subset B_{kr_m}(x_0)$  for all m.

Since

$$\left| \int_{X_m} \varphi \circ \iota_m(z) \Delta_{r_m r} u(z) d\mu_m(z) \right| \leq \frac{\|\varphi \circ \iota_m\|_{\infty}}{\mu(B_{r_m}(x_0))} \int_{B_{kr_m}(x_0)} |\Delta_{r_m r} u(z)| d\mu(z)$$
$$\leq C_{k,\varphi} \int_{B_{kr_m}(x_0)} |\Delta_{r_m r} u| d\mu < \infty$$

it follows that

$$\int_{X_{\infty}} \varphi \circ \iota_{\infty}(z) \left[ (u_{\infty})_{B_{\infty}(z,r)} - u_{\infty}(z) \right] d\mu_{\infty}(z) = 0.$$

Since  $\varphi$  is arbitrary, this establishes the claim for all rational r > 0. The claim follows for all r > 0 by continuity, since  $X_{\infty}$  is a length space and  $\mu_{\infty}$  a doubling measure.

## 6. WEAKLY AMV-HARMONIC FUNCTIONS

We now turn to weakly amv-harmonic functions. In this section we prove Theorems 1.1 and 1.2. We begin by relating the Korevaar–Schoen energy to the symmetrized *r*-laplacians, thus establishing the general principle that local minimizers correspond to functions with vanishing symmetrized *r*-laplacian, cf. Proposition 6.2.

In the case of Carnot groups the symmetrized *r*-laplacian agrees with the *r*-laplacian. Thus Theorem 1.2 follows directly from Proposition 6.2, once we establish that Carnot groups have a Korevaar–Schoen Dirichlet form, see Proposition 6.3.

Finally we consider non-collapsed RCD-spaces. After reviewing their relevant basic properties we show that having vanishing mm-boundary implies that  $\Delta_r$  and  $\widetilde{\Delta}_r$  agree asymptotically (Lemma 6.7). Since by [31] the Korevaar–Schoen energy is a Dirichlet form, Theorem 1.1 follows.

6.1. Symmetrized *r*-laplacian and the Korevaar–Schoen energy. Throughout this subsection,  $X = (X, d, \mu)$  denotes a generic metric measure space. To define the symmetrized *r*-laplacian, we denote by  $k_r : X \times X \to \mathbb{R}$  the symmetric mean value kernel:

$$k_r(x,y) = \frac{1}{2}\chi_{[0,r]}(d(x,y))\left(\frac{1}{\mu(B_r(x))} + \frac{1}{\mu(B_r(y))}\right), \quad x,y \in X, r > 0.$$
(30)

**Definition 6.1.** Let  $X = (X, d, \mu)$  be a metric measure space, and r > 0. The *symmetrized r-laplacian* of a function  $u \in L^1_{loc}(X)$  is given by

$$\widetilde{\Delta}_r u(x) = \int_X k_r(x, y) \frac{u(y) - u(x)}{r^2} \mathrm{d}\mu(y), \quad x \in X.$$

Let  $u \in L^1_{loc}(X)$  and r > 0. A direct calculation yields the following identity (cf. formula (11)).

$$\widetilde{\Delta}_r u = \frac{1}{2} \left( \Delta_r u + \Delta_r^* u - u \Delta_r^* 1 \right), \quad x \in X.$$
(31)

Moreover, we have the following elementary identities (recall definitions of  $e_r$  and  $E_r$  in Section 2.4). Lemma 6.1. Let  $u, v \in L^2_{loc}(X)$  and r > 0. Then

$$\Delta_r(uv) = u\Delta_r v + 2e_r(u,v) + v\Delta_r u.$$

In particular, it holds that

(a) 
$$\int_X v \widetilde{\Delta}_r u \mathrm{d}\mu = -E_r(u, v)$$

and

(b) 
$$\int_X v(\Delta_r u - \widetilde{\Delta}_r u) d\mu = \frac{1}{2} \int_X v(x) \oint_{B_r(x)} \frac{\delta_r(x,y)}{r} \frac{u(y) - u(x)}{r} d\mu(y) d\mu(x)$$

and whenever  $u, v \in L^2(X)$ . Here

$$\delta_r(x,y) = 1 - \frac{\mu(B_r(x))}{\mu(B_r(y))}.$$

*Proof.* For any  $x \in X$ , we have

$$\begin{aligned} u(x)\Delta_r v(x) &+ 2e_r(u,v)(x) + v(x)\Delta_r u(x) \\ &= \int_{B_r(x)} \frac{u(x)(v(y) - v(x)) + (u(y) - u(x))(v(y) - v(x)) + v(x)(u(y) - u(x))}{r^2} d\mu(y) \\ &= \int_{B_r(x)} \frac{u(y)v(y) - u(x)v(x)}{r^2} d\mu(y) = \Delta_r(uv)(x), \end{aligned}$$

proving the first identity in the claim. Together with Remark 2.1, the identity (31) implies (a): indeed, we have that

$$2\int_{X} v\widetilde{\Delta}_{r} u d\mu = \int_{X} v(\Delta_{r} u + \Delta_{r}^{*} u - u\Delta_{r}^{*} 1) d\mu = \int_{X} (v\Delta_{r} u + v\Delta_{r}^{*} u - uv\Delta_{r}^{*} 1) d\mu$$
$$= \int_{X} (v\Delta_{r} u + u\Delta_{r} v - \Delta_{r}(uv)) d\mu = -2\int_{X} e_{r}(u, v) d\mu$$

The identity (b) follows directly from the observation that

$$\Delta_r u(x) - \widetilde{\Delta}_r u(x) = \frac{1}{2} \int_{B_r(x)} \left[ \frac{1}{\mu(B_r(x))} - \frac{1}{\mu(B_r(y))} \right] \frac{u(y) - u(x)}{r^2} \mathrm{d}\mu(y)$$

The characterization of local minimizers of the Korevaar–Schoen energy in terms of the symmetrized r-laplacians – in spaces with a KS-Dirichlet form – is a straightforward consequence of the preceding observations.

**Proposition 6.2.** Suppose X is a metric measure space where the Korevaar–Schoen energy  $E_{KS}^2$  exists in the sense of Definition 2.4. Then the following are equivalent for a function  $u \in KS_{loc}^{1,2}(\Omega)$  in a domain  $\Omega \subset X$ :

- (1) *u* is a local minimizer of  $E_{KS}^2$  in  $\Omega$ ;
- (2) for every  $\varphi \in KS^{1,2}(\Omega)$  with compact support in  $\Omega$  we have

$$\lim_{r \to 0} \int_X \varphi \widetilde{\Delta}_r u \mathrm{d}\mu = 0.$$

*Proof.* Recall that local minimizers of  $E_{KS}^2$  in  $\Omega$  are characterized by the property that  $\mathcal{E}_{KS}(\varphi, u) = 0$  for every  $\varphi \in KS^{1,2}(\Omega)$  with compact support in  $\Omega$ . Recall also, that the Carré du Champ for  $\mathcal{E}_{KS}$  is e(u, v). By Lemma 6.1 and the existence of the Korevaar–Schoen energy, this is equivalent with

$$\lim_{r \to 0} \int_X \varphi \widetilde{\Delta}_r u \mathrm{d}\mu = 0.$$

**Remark 6.1.** More generally, we have the following: if  $u \in D(\Delta_{KS}; \Omega)$ , then

$$\int_{X} \varphi \Delta_{KS} u \mathrm{d}\mu = -\mathcal{E}_{KS}(\varphi, u) = \lim_{r \to 0} \int_{X} \varphi \widetilde{\Delta}_{r} u \mathrm{d}\mu$$

for each  $\varphi \in KS^{1,2}(\Omega)$  with compact support in  $\Omega$ .

6.2. **Carnot groups.** A Carnot group  $\mathbb{G}$  is a connected and simply connected nilpotent Lie group whose Lie algebra  $\mathfrak{g}$  is finite dimensional and admits a stratification of step  $s \in \mathbb{N}$ 

$$\mathfrak{g} = V_1 \oplus \cdots \oplus V_s$$

with the property that  $[V_1, V_j] = V_{j+1}$  for j = 1, ..., s - 1 and  $[V_1, V_s] = \{0\}$ .  $v_i := \dim V_i$  and set  $h_0 = 0$ ,  $h_i = v_1 + \cdots + v_i$  and  $m = h_s$ . By using the exponential map, every Carnot group  $\mathbb{G}$  of step s is isomorphic as a Lie group to  $(\mathbb{R}^m, \cdot)$ , with the group operation given by the Baker–Campbell–Hausdorff formula.

The stratification gives rise to a family of dilations  $\delta_t : \mathbb{G} \to \mathbb{G}$  for t > 0, given by

$$\delta_t(x_1,\ldots,x_m) = (t^{\sigma_1}x_1,\ldots,t^{\sigma_s}x_m),$$

where the homogeneities  $\sigma_i$  are defined by  $\sigma_j := i$  for  $h_{i-1} < j \le h_i$ .

A continuous function  $\rho : \mathbb{G} \to [0,\infty)$  is said to be a *pseudonorm* on *G*, if  $\rho$  satisfies the following conditions:

(1) 
$$\rho(\delta_r(x)) = r\rho(x)$$
 for every  $r > 0$  and  $x \in G$ ,

(2)  $\rho(x) > 0$  if and only if  $x \neq 0$ ,

(3)  $\rho$  is symmetric, i.e.  $\rho(x^{-1}) = \rho(x)$  for every  $x \in G$ .

A pseudonorm defines a pseudodistance on a group. Examples of such pseudodistances include the Korányi–Reimann distance and the following :

$$|x|_{\mathbb{G}} := |(x^{(1)}, \dots, x^{(s)})|_{\mathbb{G}} := \left(\sum_{j=1}^{s} ||x^{(j)}||^{\frac{2s!}{j}}\right)^{\frac{1}{2s!}} \qquad \mathbf{d}(x, y) := |y^{-1} \cdot x|_{\mathbb{G}},\tag{32}$$

where  $x^{(i)} = (x_{h_{i-1}+1}, \dots, x_{h_i})$  for  $i = 1, \dots, s$ , and  $||x^{(j)}||$  stands for Euclidean norm in  $\mathbb{R}^{v_j}$ . We observe that with the above notation, the coordinates in  $\mathbb{G}$  can be given a form  $x = (x^{(1)}, x^{(2)}, \dots, x^{(s)})$ .

The number  $Q := \sum_{i=1}^{k} i \dim(V_i)$  is called the *homogeneous dimension* of  $\mathbb{G}$ . The Lebesgue measure  $\mathcal{L}^m := \mu$  is the Haar measure on  $\mathbb{G}$ . Moreover, for all t > 0 it holds  $\mu(\delta_t(E)) = t^Q \mu(E)$ , for a measurable  $E \subset \mathbb{G}$ . In particular,  $\mu(B(x,r)) = cr^Q$  for any left-invariant metric (*c* depends on the choice of metric) and the Hausdorff-dimension of  $\mathbb{G}$  equals Q.

Next we come to Sobolev spaces. Let  $\Omega \subset \mathbb{G}$  be a domain. We define the horizontal Sobolev space  $HW^{1,p}(\Omega)$  as space of measurable functions  $u \in L^p(\Omega)$  whose distributional horizontal derivatives  $X_j u$  are  $L^p$ -functions. We denote the horizontal gradient of u by

$$\nabla_H u = \sum_{j=1}^k (X_j u) X_j.$$

The norm  $||u||_{1,p}^p = ||u||_{L^p(\Omega)}^p + ||\nabla_H u||_{L^p(\Omega)}^p$  makes  $HW^{1,p}(\Omega)$  into a Banach space.

The various different notions on Sobolev spaces on  $\Omega \subset \mathbb{G}$  agree, since  $\mathbb{G}$  supports a 1-Poincaré inequality, cf. [36]. We mention here that, for  $u \in HW^{1,2}(\Omega)$ , the function  $|\nabla_H u|$  is a minimal weak upper gradient of u.

Below we will also use the approximate Pansu differential  $Du_x : \mathbb{G} \to \mathbb{R}$  of a Sobolev function  $u \in W^{1,2}_{loc}(\Omega)$  at almost every  $x \in \Omega$ , which can be seen as a natural derivative in Carnot groups and plays an important role in function and mapping theory (e.g. in quasiconformal analysis). The following property relating the Pansu differential and the horizontal gradient of a function can be found in [58, Remark 3.3], see also [17, Corollary 3.5].

$$Du_x(z) = \langle \nabla_H u(x), z^{(1)} \rangle, \quad z = (z^{(1)}, \dots, z^{(s)}) \in \mathbb{G}.$$

**Proposition 6.3.** Let  $\Omega \subset \mathbb{G}$  be a domain a Carnot group and  $\rho$  be a pseudonorm on  $\mathbb{G}$  of the form

$$\rho(z) = F(|z^{(1)}|, z'), \quad z = (z^{(1)}, z') \in \mathbb{G},$$
(33)

where  $z' = (z^{(2)}, \ldots, z^{(s)})$  and  $F : \mathbb{R} \times \mathbb{R}^{m-v_1} \to \mathbb{R}$  is  $C^1$  outside the origin and  $\partial_1 F > 0$ . Then for every  $u \in HW^{1,2}(\Omega)$  we have

$$e^2(u) = c|\nabla_H u|^2$$

almost everywhere, where *c* is a constant depending only on  $\rho$ , the data of  $\mathbb{G}$ , and e(u) is the KS-energy density of *u*, *cf*. Definition 2.4.

**Remark 6.2.** A particular class of homogeneous norms satisfying (33) is given as follows. Consider a norm N on  $\mathbb{R}^s$  with  $\partial_1 N > 0$ , and norms  $\|\cdot\|_i$  on  $V_i$  for i = 2, ..., s. The function

$$\rho(z) := N(|z^{(1)}|, ||z^{(2)}||_2^{1/2}, \dots, ||z^{(s)}||_s^{1/s}), \quad z = (z^{(1)}, \dots, z^{(s)}) \in \mathbb{G}$$
(34)

is a homogeneous norm. This class includes the Korányi–Reimann norm on  $\mathbb{H}^1$  and more generally norms of the form (33).

Proof. By the proof of [65, Theorem 4.1], we have that the KS-energy density exists and satisfies

$$e(u)(x) = \oint_{B_{\rho}} |Du_x(z)|^2 \mathrm{d}z = \oint_{B_{\rho}} |\langle \nabla_H u, z^{(1)} \rangle|^2 \mathrm{d}z$$

for a.e  $x \in \Omega$ . The last equality follows from the remark above. Here  $B_{\rho}$  is the unit ball with respect to the pseudonorm  $\rho$ .

Since every two pseudonorms are equivalent (see [11, Proposition 5.1.4] and [2, Remark 2]) we have, in particular, that by (32) it holds

$$\{z \in \mathbb{G} : |z|_{\mathbb{G}} \le 1/C\} \subset \{z \in \mathbb{G} : F(|z^{(1)}|, z') \le 1\} \subset \{z \in \mathbb{G} : |z|_{\mathbb{G}} \le C\}$$

This observation together with the assumptions on  $\rho$  imply that, for each  $z' \in M$ , where  $M \subset \mathbb{R}^{m-v_1}$  is a bounded (and closed) set, the function  $F(\cdot, z')$  is invertible and thus there exists a continuous function  $G: M \to \mathbb{R}$  such that for each fixed z'

$$\{(z^{(1)}, z') \in \mathbb{G} : \rho(z^{(1)}, z') \le 1\} = \{(z^{(1)}, z') \in \mathbb{G} : |z^{(1)}| \le G(z')\}.$$

Using this and the change of variables, we compute

$$\begin{split} \int_{B_{\rho}} |\langle \nabla_{H}u, z^{(1)} \rangle|^{2} \mathrm{d}z^{(1)} &= \int_{M} \left( \int_{\{|z^{(1)}| \leq G(z')\}} |\langle \nabla_{H}u(x), z^{(1)} \rangle|^{2} \mathrm{d}z^{(1)} \right) \mathrm{d}z' \\ &= \int_{\{z' \in M: G(z') \geq 0\}} G(z')^{2+k} \left( \int_{\{|z^{(1)}| \leq 1\}} |\langle \nabla_{H}u(x), z^{(1)} \rangle|^{2} \mathrm{d}z^{(1)} \right) \mathrm{d}z' \\ &= c \int_{\{|z^{(1)}| \leq 1\}} |\langle \nabla_{H}u(x), z^{(1)} \rangle|^{2} \mathrm{d}z^{(1)}, \end{split}$$

where *c* is a constant. Indeed, since *G* is continuous and *M* is bounded, the integral of  $G^{2+k}$  is finite. Then, by [31, Lemma 4.18] we have that

$$\int_{\{|z^{(1)}|\leq 1\}} |\langle \nabla_H u(x), z^{(1)} \rangle|^2 \mathrm{d} z^{(1)} = c(v_1) |\nabla_H u(x)|^2.$$

This completes the proof.

We finish the subsection by providing the proof of Theorem 1.2.

*Proof of Theorem 1.2.* From Proposition 6.3 it follows that the KS-energy is a constant multiple of the Cheeger energy, and thus a Dirichlet form. Moreover, it follows that the KS-laplacian is a constant multiple of the sub-Laplacian. Proposition 6.2 now directly implies Theorem 1.2.

6.3. **RCD-spaces.** RCD-spaces arise from the study of the synthetic curvature bounds, initially stated in terms of convexity properties of entropy functionals in Wasserstein spaces for Riemannian manifolds, cf. [66, 63]. A *Riemannian* theory of the resulting *curvature dimension conditions* (CD(K, N)conditions) is obtained by requiring in addition the quadratic property for the Cheeger energy, see Section 2.4. The notion of *RCD*-spaces was introduced and substantially developed by Ambrosio– Gigli–Savare, cf. [5]. In terms of the  $\Gamma$ -calculus introduced in Section 2.4, the RCD(K, N)-condition with parameters  $K \in \mathbb{R}$  and  $N \ge 1$ , can be characterized as:

$$\frac{1}{2}\int |\nabla f|^2 \Delta g \mathrm{d}\mu \ge \int g\left(K|\nabla f|^2 + \langle \nabla f, \nabla \Delta f \rangle + \frac{1}{N}(\Delta f)^2\right) \mathrm{d}\mu$$

for any  $f \in D(\Delta)$  with  $\Delta f \in W^{1,2}(X)$  and any  $g \in L^{\infty}(X) \cap D(\Delta)$ . We refer, for instance, to [66] for a survey of synthetic curvature conditions.

By a result of Gigli–Tyulenev [31] the Korevaar–Schoen energy is a constant multiple of the Cheeger energy, and thus a Dirichlet form, on RCD-spaces.

**Proposition 6.4** (cf. Theorem 2.18 and Proposition 4.19 in [31]). Let X be an RCD(K, N)-space, for  $K \in \mathbb{R}$  and  $N \geq 1$ . Then there is a constant c = c(K, N) so that, if  $\Omega \subset X$  is a domain, then for any  $u \in KS^{1,2}(\Omega)$  we have that

$$\int_{\Omega} |\nabla u|^2 \mathrm{d}\mu = c(N) E_{KS}^2(u;\Omega),$$

where  $|\nabla u|$  is the minimal weak upper gradient of u.

We remark that by Proposition 6.4 local minimizers of  $E_{KS}^2$  are exactly the local minimizers of  $Ch_2$ . It is known, cf. [49], that local minimizers of  $Ch_2$  on a domain  $\Omega$  are locally Lipschitz continuous in  $\Omega$  and are characterized by the equation  $\Delta u = 0$ .

RCD-spaces also enjoy geometric properties akin to Riemannian manifolds. Of interest to us is the Bishop–Gromov inequality, due to Sturm [64]. In order to state it, we recall the following auxiliary functions. For numbers  $K \in \mathbb{R}$  and  $N \ge 1$ , define  $s_{K,N} : [0, \infty) \to \mathbb{R}$  by

$$s_{K,N}(t) = \begin{cases} \sqrt{\frac{N-1}{K}} \sin\left(t\sqrt{\frac{K}{N-1}}\right), & K > 0\\ t, & K = 0\\ \sqrt{\frac{N-1}{-K}} \sinh\left(t\sqrt{-\frac{K}{N-1}}\right), & K < 0. \end{cases}$$

Furthermore, let

$$v_{K,N}(r) := N\omega_N \int_0^r s_{K,N}(t)^{N-1} \mathrm{d}t, \quad r \ge 0.$$

**Theorem 6.5** (cf. Theorem 2.3, Remark 5.3 in [64]). Let X be an RCD(K, N)-space, for  $K \in \mathbb{R}$  and  $N \ge 1$ . Then for any  $x \in X$ , the function

$$r \mapsto \frac{\mu(B_r(x))}{v_{K,N}(r)}$$

*is non-increasing on the interval*  $\left(0, \pi \sqrt{\frac{N-1}{\max\{K,0\}}}\right)$ .

Remark 6.3. Note that

$$\lim_{r \to 0} \frac{v_{K,N}(r)}{\omega_N r^N} = 1 \text{ and } v_{K,N}(r) \le \omega_N r^N + C_{K,N} r^{N+2}, \quad r < 1/C_{K,N}$$

for a suitable constant  $C_{K,N}$ , cf. [41]. In particular, in non-collapsed RCD-spaces, the Bishop–Gromov density  $\theta^N$  exists and satisfies  $\theta^N \leq 1$  at every point.

We record the following consequence of the Bishop–Gromov inequality as a corollary.

**Corollary 6.6.** Let X be a non-collapsed RCD(K, N)-space. Then  $\mathcal{H}^N$  is locally Ahlfors N-regular.

In general, RCD(K, N)-spaces are rectifiable but the dimension may be strictly smaller than N (see, for example weighted Euclidean spaces in the Introduction). Non-collapsed RCD(K, N)-spaces on the other hand are N-rectifiable.

We next employ the vanishing metric measure boundary to obtain an asymptotic connection between the *r*-laplacian and the symmetrized *r*-laplacian.

**Lemma 6.7.** Suppose X is a non-collapsed RCD(K, N)-space and has vanishing mm-boundary. Then

$$\lim_{r \to 0} \int_X \varphi(x) \oint_{B_r(x)} \left| \frac{\delta_r(x, y)}{r} \right| d\mathcal{H}^N(y) d\mathcal{H}^N u(x) = 0$$

for every  $\varphi \in C_c(X)$ .

*Proof.* It suffices to prove the claim for non-negative  $\varphi$ . Recall the definition of  $\delta_r(x,y) := 1 - \mathcal{H}^N(B_r(x))/\mathcal{H}^N(B_r(y))$ . If K is a compact set and  $x, y \in K$ , then

$$|\delta_r(x,y)| = \left|\frac{\theta_r^N(y) - \theta_r^N(x)}{\theta_r^N(y)}\right| \le C_K(|\theta_r^N(y) - 1| + |\theta_r^N(x) - 1|),$$
(35)

where  $C_K$  is the Ahlfors regularity constant on K. By Remark 6.3 we have that

$$\left(\frac{\theta_r^N - 1}{r}\right)_{-} \le C_{K,N}r$$

for small r > 0. This and the fact that X has vanishing mm-boundary imply that

$$\lim_{r \to 0} \int_X \varphi \left| \frac{\theta_r^N - 1}{r} \right| d\mathcal{H}^N = 0$$
(36)

for every  $\varphi \in C_c(X)$ . From (35) we obtain

$$\int_{X} \varphi \int_{B_{r}(x)} \left| \frac{\delta_{r}(x,y)}{r} \right| d\mu(y) d\mathcal{H}^{N}(x)$$
  
$$\leq C_{K'} \int_{X} \varphi \left| \frac{\theta_{r}^{N}(x) - 1}{r} \right| d\mathcal{H}^{N}(x) + C_{K'} \int_{X} \varphi(x) \int_{B_{r}(x)} \left| \frac{\theta_{r}^{N}(y) - 1}{r} \right| d\mathcal{H}^{N}(y) d\mathcal{H}^{N}(x)$$

where  $\varphi \in C_c(X)$  is non-negative,  $K' = \overline{N}_{r_0}(\operatorname{spt} \varphi)$  is a compact neighbourhood (for small enough  $r_0 > 0$ ), and  $r < r_0$ . The first term converges to zero by (36). The second term equals

$$\int_{X} A_{r}^{*} \varphi(x) \left| \frac{\theta_{r}^{N} - 1}{r} \right| d\mathcal{H}^{N}(x)$$
(37)

and also converges to zero. Indeed, let us observe that

$$\begin{split} |A_r^*\varphi(x) - \varphi(x)| &= \left| \int_{B_r(x)} \frac{\varphi(y)}{\mathcal{H}^N(B_r(y))} \mathrm{d}\mathcal{H}^N(y) - \varphi(x) \right| = \left| \int_{B_r(x)} \frac{\varphi(y)}{\mathcal{H}^N(B_r(y))} - \frac{\varphi(x)}{\mathcal{H}^N(B_r(x))} \mathrm{d}\mathcal{H}^N(y) \right| \\ &\leq \left| \int_{B_r(x)} \frac{\varphi(y) - \varphi(x)}{\mathcal{H}^N(B_r(y))} \mathrm{d}\mathcal{H}^N(y) \right| + \left| \int_{B_r(x)} \varphi(x) \left( \frac{1}{\mathcal{H}^N(B_r(y))} - \frac{1}{\mathcal{H}^N(B_r(x))} \right) \mathrm{d}\mathcal{H}^N(y) \right| \\ &\leq 2 \left\| \varphi \right\|_{\infty} \int_{B_r(x)} \frac{1}{\mathcal{H}^N(B_r(y))} \mathrm{d}\mathcal{H}^N(y) + \left\| \varphi \right\|_{\infty} \left( 1 + \int_{B_r(x)} \frac{1}{\mathcal{H}^N(B_r(y))} \mathrm{d}\mathcal{H}^N(y) \right), \end{split}$$

which is bounded (by a constant M) since  $a_r \in L^{\infty}$ . The above estimate holds for all  $x \in N_{2r}(\operatorname{spt} \varphi)$ , for  $x \notin N_{2r}(\operatorname{spt} \varphi)$  both  $A_r^*\varphi$  and  $\varphi$  vanish. Now we focus on proving convergence to zero in (37) as  $r \to 0$ .

$$\left| \int_X A_r^* \varphi(x) \left| \frac{\theta_r^N - 1}{r} \right| \mathrm{d}\mathcal{H}^N(x) \right| \le \int_X |A_r^* \varphi(x) - \varphi(x)| \left| \frac{\theta_r^N - 1}{r} \right| \mathrm{d}\mathcal{H}^N(x) + \int_X |\varphi(x)| \left| \frac{\theta_r^N - 1}{r} \right| \mathrm{d}\mathcal{H}^N(x).$$

The second term goes to zero by our assumptions on vanishing metric boundary, hence we take care of the first term. Fix  $\varepsilon > 0$  and consider  $r < \frac{\varepsilon}{2}$ :

$$\int_{X} |A_{r}^{*}\varphi(x) - \varphi(x)| \left| \frac{\theta_{r}^{N} - 1}{r} \right| d\mathcal{H}^{N}(x) = \int_{N_{\epsilon}(\operatorname{spt}\varphi)} |A_{r}^{*}\varphi(x) - \varphi(x)| \left| \frac{\theta_{r}^{N} - 1}{r} \right| d\mathcal{H}^{N}(x)$$
$$\leq M \int_{N_{\epsilon}(\operatorname{spt}\varphi)} \left| \frac{\theta_{r}^{N} - 1}{r} \right| d\mathcal{H}^{N}(x)$$

In order to show that the above expression goes to 0 we choose a continuous function  $\psi$  with spt  $\psi \subset N_{2\epsilon}(\operatorname{spt} \varphi)$  and constantly equal to 1 on  $N_{\epsilon}(\operatorname{spt} \varphi)$ . Then, we estimate in the following way

$$\int_{N_{\epsilon}(\operatorname{spt}\varphi)} \left| \frac{\theta_r^N - 1}{r} \right| d\mathcal{H}^N(x) \le \int_X \psi(x) \left| \frac{\theta_r^N - 1}{r} \right| d\mathcal{H}^N(x).$$

where the right-hand side goes to 0 as  $r \rightarrow 0$ .

**Corollary 6.8.** Let X be a non-collapsed RCD(K, N)-space with vanishing mm-boundary. Then

$$\lim_{r \to 0} \left| \int_X \varphi(\widetilde{\Delta}_r u - \Delta_r u) \mathrm{d}\mathcal{H}^N \right| = 0$$

whenever  $u, \varphi \in \text{LIP}(\Omega)$  and u or  $\phi$  have compact support in  $\Omega$ .

*Proof.* By a cut-off argument we may assume that both  $\varphi$  and u have compact support in  $\Omega$ . By Lemma 6.1 we have that

$$\left| \int_{X} \varphi(\widetilde{\Delta}_{r} u - \Delta_{r} u) \mathrm{d}\mathcal{H}^{N} \right| \leq \int_{X} |\varphi(x)| \operatorname{LIP}(u) \oint_{B_{r}(x)} \left| \frac{\delta_{r}(x, y)}{r} \right| \mathrm{d}\mathcal{H}^{N}(y) \, \mathrm{d}\mathcal{H}^{N}(x).$$

The claim now follows from Lemma 6.7.

*Proof of Theorem 1.1.* Let  $u : \Omega \to \mathbb{R}$  be harmonic. In particular, u is locally Lipschitz. By Lemma 6.4 u is a local minimizer of  $E_{KS}^2$  in  $\Omega$ . Theorem 6.2 implies that

$$\lim_{r\to 0} \int_{\Omega} \varphi \widetilde{\Delta}_r u \mathrm{d}\mu = 0$$

for every  $u \in KS^{1,2}(\Omega)$  with compact support in  $\Omega$ . This and Corollary 6.8 imply that

$$\lim_{r \to 0} \int_X \varphi \Delta_r u \mathrm{d}\mu = 0$$

for every  $\varphi \in \text{LIP}_c(\Omega)$ .

**Remark 6.4.** It follows from Corollary 6.8 and the argument above that, if  $u : \Omega \to \mathbb{R}$  is *a locally Lipschitz continuous* weakly amv-harmonic function, then *u* is Cheeger harmonic.

## 7. IMPROVING REGULARITY: THE BPZ-THEOREM FOR AMV-HARMONIC FUNCTIONS

In this section we discuss Blaschke-Privaloff-Zaremba type theorems (BPZ theorems, for short). In its classical version in the setting of Euclidean spaces, see e.g. [55, Theorem 2.1.5], the BPZ theorem asserts that given an open set in  $\mathbb{R}^n$  a continuous pointwise amv-harmonic function solves locally the Laplace equation. Thus, the pointwise nullity of the amv-harmonic operator  $\lim_{r\to 0^+} \Delta_r$  improves the regularity of amv-harmonic functions. Below we show that this is also the case of amv-harmonic functions in Heisenberg group  $\mathbb{H}_1$  and on Riemannian manifolds. Moreover, we show counterpart of the BPZ theorem in the Alexandrov spaces and discuss the relation between the amv-harmonic functions defined in the weak sense and the Laplacian in Alexandrov spaces.

The setting of the Heisenberg group  $\mathbb{H}_1$ . Solutions to the subelliptic Laplace equation are  $C^2$  due to results e.g. in [16] and [56]. The Dirichlet problem on Korányi–Reimann balls for the continuous boundary data has the classical  $C^2$ -solution for harmonic sub-elliptic equation in  $\mathbb{H}_1$ , see [27]. However, it is shown in [37], that for  $\mathbb{H}_n$  for  $n \ge 2$  balls in the Carnot-Carathéodory distance are not regular at the characteristic points. Therefore, due to the approach we take in the proof of Theorem 7.1 below, we will restrict our discussion to the case of  $\mathbb{H}_1$  and balls with respect to the *Korányi–Reimann* distance.

For the readers convenience we recall that our model for  $\mathbb{H}^1$  is the group  $(\mathbb{R}^3, \cdot)$  where the group law is given by

$$(x, y, t) \cdot (x', y', t') = (x + x', y + y', t + t' - 2xy' + 2x'y).$$

By using this group law, one introduces a frame of left-invariant vector fields which agree with the standard basis at the origin:

$$X := \partial_x + 2y\partial_t, \quad Y := \partial_y - 2x\partial_t, \quad T := \partial_t.$$

The Korányi–Reimann distance is a metric defined by

$$d_{\mathbb{H}^1}(p,q) := \|q^{-1}p\|_{\mathbb{H}^1}, \text{ where } \|(x,y,t)\|_{\mathbb{H}^1} = \sqrt[4]{(x^2+y^2)^2+t^2}.$$

Let  $p_0 \in \mathbb{H}_1$  and R > 0. An open ball in  $\mathbb{H}_1$  centered at  $p_0$  with radius R with respect to metric  $d_{\mathbb{H}^1}$  is defined as follows:  $B(p_0, R) := \{p \in \mathbb{H}_1 : \|p^{-1}p_0\|_{\mathbb{H}^1} < R\}$ . For further discussion on Carnot groups we refer to Section 6.2.

**Theorem 7.1** (The Blaschke-Privaloff-Zaremba theorem for antv-harmonic functions in  $\mathbb{H}_1$ ). Let  $\Omega \subset \mathbb{H}_1$  be a domain in the first Heisenberg group  $\mathbb{H}_1$  equipped with metric  $d_{\mathbb{H}^1}$  and let  $f : \Omega \to \mathbb{R}$  be a continuous satisfying

$$\lim_{r \to 0} \Delta_r u(x) = 0, \quad x \in \Omega.$$

*Then f is a sub-elliptic harmonic function, i.e. for all*  $x \in \Omega$  *it holds that* 

$$\lim_{r \to 0^+} \Delta_r f(x) = \frac{1}{3\pi} \Delta_{\mathbb{H}_1} f(x) = 0.$$

The constant  $-\frac{1}{3\pi}$  is computed in [24, Lemma 3.3]. We further remark that the same lemma shows the Blaschke-Privaloff-Zaremba theorem for  $C^2$ -functions.

*Proof.* In the proof we follow the original idea of Privaloff developed for the setting of  $\mathbb{R}^n$ , see [61, Theorem II]. Let  $p_0 \in \Omega$  and  $B = B(p_0, R)$  be a ball centered at  $p_0$  with radius R > 0 such that  $B \subset \Omega$ . Theorem in [27] allows us to infer that the sub-elliptic Dirichlet problem on B with the boundary data f has the unique solution, denoted by u, such that  $u \in C(\overline{B})$  and  $u = f|_{\partial B}$ . Set  $\phi = f - u$ . Then  $\phi \in C(\overline{B})$  and  $\phi|_{\partial B} \equiv 0$ . The assertion will be proven if we show that  $\phi \equiv 0$  in B. We argue by contradiction. Namely, suppose that there exists  $q \in B$  such that  $\phi(q) \neq 0$  and without the loss of generality we assume that  $\phi(q) < 0$ . Let us define the following function on B

$$F(p) = \phi(p) + \frac{\phi(q)}{2} \left( \frac{\|(p^{-1}p_0)_H\|_{\mathbb{R}^2}^2 - R^2}{R^2} \right)$$

where  $||(p^{-1}p_0)_H||_{\mathbb{R}^2}$  stands for the Euclidean length in  $\mathbb{R}^2$  of the horizontal part of point  $p^{-1}p_0 \in \Omega$ . It follows that  $F \in C(\overline{B})$ ,  $F|_{\partial B} \ge 0$  and that F(q) < 0. Hence, there is  $q_m \in \overline{B}$  such that  $F(q_m) = \min_{\overline{B}} F$  (in fact,  $q_m \in B$ ). Moreover,  $\Delta_r F(q_m) \ge 0$  for all  $r \le R$  and by direct computations we verify that

$$\Delta_r F(q_m) = \Delta_r \phi(q_m) + \frac{\phi(q)}{2} \Delta_r \left( \frac{\|(p^{-1}p_0)_H\|_{\mathbb{R}^2}^2 - R^2}{R^2} \right) (q_m)$$

Therefore, upon applying the definition of  $\Delta_r$ , we arrive at the following estimate

$$0 \le \Delta_r F(q_m) = \Delta_r f(q_m) - \Delta_r u(q_m) + \frac{\phi(q)}{2} \Delta_r \left( \frac{\|(p^{-1}p_0)_H\|_{\mathbb{R}^2}^2 - R^2}{R^2} \right) (q_m).$$
(38)

Let us denote the coordinates of p and  $p_0$  as follows: p = (x, y, t) and  $p_0 = (x_0, y_0, t_0)$ . Then

$$\frac{\phi(q)}{2}\Delta_{\mathbb{H}_1}\left(\frac{\|(p^{-1}p_0)_H\|_{\mathbb{R}^2}^2 - R^2}{R^2}\right)(q_m) = \frac{\phi(q)}{2R^2}\Delta_{\mathbb{H}_1}\left((x - x_0)^2 + (y - y_0)^2\right)(q_m)$$

Recall, that

$$\Delta_{\mathbb{H}_1} = \partial_x^2 + \partial_y^2 + (4x^2 + 4y^2)\partial_t^2 + 4y\partial_x\partial_t - 4x\partial_y\partial_t$$

Therefore, in our case the sub-Laplacian reduces to the Laplacian in  $\mathbb{R}^2$ 

$$\Delta_{\mathbb{H}_1} \left( (x - x_0)^2 + (y - y_0)^2 \right) (q_m) = \Delta \left( (x - x_0)^2 + (y - y_0)^2 \right) (q_m) = 4$$

Since f is assumed to be pointwise amv-harmonic and  $u \in C^2(B)$ , the definition of amv-harmonic functions together with [24, Lemma 3.3] imply that upon  $r \to 0^+$  it holds that  $\Delta_r f(q_m) \to 0$  and  $\Delta_r u(q_m) \to 0$ . Moreover, by applying [24, Lemma 3.3] again we obtain that

$$\lim_{r \to 0^+} \frac{\phi(q)}{2} \Delta_r \left( \frac{\|(p^{-1}p_0)_H\|_{\mathbb{R}^2}^2 - R^2}{R^2} \right) (q_m) = \frac{1}{3\pi} \frac{\phi(q)}{2} \Delta_{\mathbb{H}_1} \left( \frac{\|(p^{-1}p_0)_H\|_{\mathbb{R}^2}^2 - R^2}{R^2} \right) (q_m) = \frac{2\phi(q)}{3\pi R^2} < 0,$$

since above we assume that  $\phi(q) < 0$ . In a consequence, we get that  $0 \le \frac{2\phi(q)}{3\pi R^2} < 0$  contradicting our assumption. The proof of the theorem is completed.

*The setting of Riemannian manifolds.* For the readers convenience, we sketch the proof of the BPZ-theorem in the Riemannian setting. The argument is analogous to that in the proof of Theorem 7.1

**Proposition 7.2** (The Blaschke-Privaloff-Zaremba theorem on manifolds). Let (M, g) be an *n*-dimensional Riemannian manifold with Ricci curvature bounded below by some  $K \in \mathbb{R}$ . Let further  $\Omega \subset M$  be a domain in M and let  $f : \Omega \to \mathbb{R}$  be a continuous function in  $\Omega$  satisfying

$$\lim_{n \to \infty} \Delta_r u(x) = 0, \quad x \in \Omega.$$

Then f solves the Beltrami-Laplace equation in  $\Omega$ .

*Proof.* As in the proof for Theorem 7.1 we argue by contradiction. Fix  $z_0 \in \Omega$  and a ball  $B = B(z_0, R) \subset \Omega$  for a small enough radius R > 0, which will be determined below. As previously, we find function u as the unique solution to the Beltrami–Laplace Dirichlet problem  $\Delta_M u = 0$  in B, such that  $u \in C(\overline{B})$  and  $u = f|_{\partial B}$ . The existence of such u follows, for instance from the general theory for metric measure spaces, since the Riemannian manifold M with the Ricci curvature bounded from below satisfies the doubling condition and the 1-Poincaré inequality (see [8, Chapter A.7] and [36] for details).

As in the proof of Theorem 7.1 we set  $\phi = f - u$  and notice that  $\phi \in C(\overline{B})$  and  $\phi|_{\partial B} \equiv 0$ . In order to prove the assertion we show that  $\phi \equiv 0$  in B. We argue by contradiction and suppose that there exists  $y \in B$  such that  $\phi(y) \neq 0$  and without the loss of generality we assume that  $\phi(y) < 0$ .

Similarly as in the proof of Theorem 7.1 we define function F as follows

$$F(z) = \phi(z) + \frac{\phi(y)}{2} \left( \frac{\text{dist}_M(z, z_0)^2 - R^2}{R^2} \right),$$

where  $dist_M(\cdot, z_0)$  is a distance function to  $z_0 \in \Omega$  defined in metric g on M. The direct computations for the Beltrami-Laplace operator, see the proof of [54, Theorem 4.1], allow us to conclude that

$$\Delta_M(\operatorname{dist}_M(z, z_0)^2) = 2 + 2H_{z_0}(z)\operatorname{dist}_M(z, z_0),$$

where  $H_{z_0}(z)$  stands for the mean curvature of  $\partial B(z_0, r)$  at z for  $r = \text{dist}_M(z, z_0)$ . According to the discussion in [59], it holds that  $H_{z_0}(z) = \frac{n-1}{r} - \frac{1}{3}\text{Ric}_{z_0}(v, v)r + O(r^2)$ , asymptotically as  $r \to 0$ , where  $v \in S^{n-1} \subset T_{z_0}M$  (notice that this formula arises as a perturbation of the mean-curvature formula for a sphere in the zero-curvature case). Therefore, by taking R small enough, and hence small enough r < R, we obtain that  $\Delta_M(\text{dist}_M(z, z_0)^2) > c > 0$  on B.

As previously, in order to get the contradiction we need to conclude that  $\Delta_r f(y_m) \to 0$  and  $\Delta_r u(y_m) \to 0$  as  $r \to 0^+$ . The first convergence follows, as f is amv-harmonic, while the second follows from Claim (1), Chapter 6.32 in [68]. Indeed, by [68] solutions of the Beltrami–Laplace equation are smooth (assumptions of [68, Chapter 6.32] require M to be compact, but we need this result locally). Similarly to the Euclidean argument, based on expanding u in the Taylor expansion up to the second order, one can show that  $\lim_{r\to 0^+} \Delta_r u(y_m) \leq \frac{1}{2(n+2)} \Delta_M u(y_m) = 0$ , see [70, Corollary 3.2.].

Thus, by analogous reasoning as in the previous theorem we obtain that

$$0 \leq \lim_{r \to 0^+} \frac{\phi(y)}{2} \Delta_r \left( \frac{\operatorname{dist}_M(z, z_0)^2 - R^2}{R^2} \right) (y_m) \leq \frac{\phi(y)}{4(n+2)} \Delta_M \left( \frac{\operatorname{dist}_M(z, z_0)^2 - R^2}{R^2} \right) (y_m) \\ \leq \frac{c\phi(y)}{4(n+2)R^2} < 0.$$

This contradicts our assumption that  $\phi(y) < 0$ . Hence, the proof of the theorem is completed.

The setting of the Alexandrov spaces. Another setting where a BPZ type theorem can be proven is the setting of Alexandrov spaces. Namely, let M be Alexandrov space without boundary and let  $\Omega \subset M$  be a bounded domain. For a semiconcave function  $u : \Omega \to \mathbb{R}$  (see Preliminaries in [71]) a natural Laplacian  $\mathcal{L}_u$  defined via the Dirichlet form is a signed Radon measure and its nonsingular part in the Lebesgue decomposition of  $\mathcal{L}_u$  satisfies

$$\Delta u(x) = n \oint_{\Sigma_x} H_x u(\xi, \xi) d\xi, \tag{39}$$

for almost all points  $x \in M$ , where  $H_x$  denotes the Perelman Hessian while  $\Sigma_x$  stands for the space of directions at x, see [60] and [71, Preliminaries]. We combine this observation with (2.8) in [71, Lemma 2.3] to state the following variant of the BPZ theorem in Alexandrov spaces.

**Proposition 7.3** (The Blaschke-Privaloff-Zaremba theorem in Alexandrov spaces). Let  $u : \Omega \to \mathbb{R}$  be a semiconcave function defined on a domain  $\Omega$  of Alexandrov space M without boundary. Then for all x in the set of regular points of u (i.e. for almost all points in M) it holds that

$$\lim_{r \to 0^+} \Delta_r u(x) = \frac{1}{2(n+2)} \Delta u(x).$$

In particular, a semiconcave function u on  $\Omega$  is harmonic (in the sense of the Laplacian  $\Delta$ ) if and only if  $\lim_{r\to 0^+} \Delta_r u(x) = 0$  for every regular point x of u.

The proof of this observation is the immediate consequence of the proof of [71, Lemma 2.3]. We remark that the theorem shows that for a large class of functions the amv-Laplace harmonic functions coincide with those which are harmonic in Alexandrov spaces with respect to Laplacian  $\Delta$ . However, notice that semiconcave functions are locally Lipschitz and posses a Taylor expansion up to the second order (with respect to Perelman's Hessian). Therefore, in the setting of Alexandrov spaces a characterization of the amv-harmonic functions stated in Proposition 7.3 holds under stronger assumptions than just a continuity as it is in  $\mathbb{R}^n$ ,  $\mathbb{H}_1$  and on manifolds (cf., respectively, [55, Theorem 2.1.5] and Theorem 7.1 and Proposition 7.2). Finally let us comment that, since Alexandrov spaces without boundary are conjectured to have vanishing mm-boundary (see [41, Section 1.3]) and this is known to be the case for two-dimensional Alexandrov spaces (Theorem 1.8 in [41]), the characterization part of Proposition 7.3 is related to our Theorem 1.1.

# 8. WEIGHTED EUCLIDEAN SPACES. ELLIPTIC PDES AND AMV-HARMONIC FUNCTIONS

In this final section we consider amv-harmonicity in weighted Euclidean domains and prove Theorem 1.9 which characterizes weak (and in some cases strong) amv-harmonic functions as weak solutions of the elliptic PDE, cf. (6):

$$L_w u = 0.$$

To set up notation, fix a norm  $\|\cdot\|$  on  $\mathbb{R}^n$  and a domain  $\Omega \subset \mathbb{R}^n$ . Denote  $\mathcal{H}^n_{\Omega} := \mathcal{H}^n|_{\Omega}$ , where the Hausdorff measure is taken with respect to the metric induced by  $\|\cdot\|$ . Given a positive, locally Lipschitz function  $w : \Omega \to (0, \infty)$ , we consider the metric measure spaces

$$\Omega := (\Omega, \|\cdot\|, \mathcal{H}^n_{\Omega}) \text{ and } \Omega_w = (\Omega, \|\cdot\|, w\mathcal{H}^n_{\Omega}).$$

Throughout the section,  $B^n$  denotes the (open) unit ball  $B_1(0)$  with respect to  $\|\cdot\|$ . We will use the shorthand notation

$$dy = d\mathcal{H}^n_{\Omega}(y), \quad A_r := A_r^{\mathcal{H}^n_{\Omega}}, \text{ and } \Delta_r := \Delta_r^{\mathcal{H}^n_{\Omega}},$$

as well as

$$A_r^w := A_r^{w \mathcal{H}_\Omega^n}$$
 and  $\Delta_r^w := \Delta_r^{w \mathcal{H}_\Omega^n}$ 

The following elementary facts will be used throughout the sequel.

**Remark 8.1.** Since w is continuous and strictly positive on  $\Omega$ , the measure  $w\mathcal{H}^n_{\Omega}$  is locally doubling. Moreover,

- (1)  $A_r^w f \to f$  locally uniformly in  $\Omega$  as  $r \to 0$ , whenever  $f : \Omega \to \mathbb{R}$  is continuous;
- (2) For each  $p \in [1, \infty]$ ,  $L^p_{loc}(\Omega) = L^p_{loc}(\Omega_w)$  as sets, and  $L^p$ -convergence on compact subsets of  $\Omega$  with respect to  $\mathcal{H}^n_{\Omega}$  and  $w\mathcal{H}^n_{\Omega}$  agree.

The next lemma provides two different representations for  $\Delta_r^w$  in terms of  $\Delta_r$  and will prove useful in the next subsection.

**Lemma 8.1.** Suppose  $f \in L^1_{loc}(\Omega)$ . Then

$$\Delta_r^w f = \frac{1}{A_r w} (\Delta_r (fw) - f \Delta_r w), \tag{40}$$

and

$$\Delta_r^w f = \Delta_r f + \frac{1}{A_r w} \langle f, w \rangle_r, \tag{41}$$

where

$$\langle f,g \rangle_r(x) := \int_{B_r(x)} \frac{f(y) - A_r f(x)}{r} \frac{g(y) - A_r g(x)}{r} \mathrm{d}y,$$

for  $f, g \in L^1_{loc}(\Omega)$  such that  $fg \in L^1_{loc}(\Omega)$ , and all balls  $B_r(x) \subset \Omega$ .

Proof. The first identity is a direct consequence of the pointwise identity

$$(f(y) - f(x))w(y) = f(y)w(y) - f(x)w(x) + f(x)(w(x) - w(y)), \quad x, y \in \Omega.$$

To see the second assertion (41), fix r > 0 and denote for brevity  $\mu := w \mathcal{H}_{\Omega}^{n}$ . Then

$$\frac{\langle f, w \rangle_r(x)}{A_r w(x)} = \frac{1}{A_r w(x)} \int_{B_r(x)} \frac{(f(y) - A_r f(x))(w(y) - A_r w(x))}{r^2} dy$$
  
=  $\frac{1}{A_r w(x)} \int_{B_r(x)} \frac{f(y)w(y) - A_r w(x)f(y) - A_r f(x)w(y) + A_r f(x)A_r w(x)}{r^2} dy$   
=  $\frac{1}{r^2 A_r w(x)} \left[ \int_{B_r(x)} f w dy - A_r f(x)A_r w(x) \right] = \frac{A_r^w f(x) - A_r f(x)}{r^2}$   
=  $\Delta_r^w f(x) - \Delta_r f(x), \quad x \in \Omega.$ 

8.1. AMV<sup>*p*</sup><sub>*loc*</sub>( $\Omega_w$ ) as a second order Sobolev space. To prove Theorem 1.9, we start by analyzing the operator (6). Note that, for  $u \in L^1_{loc}(\Omega)$ ,

$$L_w u := \frac{1}{2} \operatorname{div}(M \nabla u) + \frac{1}{w} \langle \nabla w, M \nabla u \rangle$$

is a distribution, and is given by a locally integrable function when  $u \in W^{2,1}_{loc}(\Omega)$ . Recall that the matrix of second moments of  $B^n$  is defined as

$$M = (m_{ij}), \quad m_{ij} = \int_{B^n} y_i y_j \mathrm{d}y, \quad i, j = 1, \dots, n.$$

 $\square$ 

**Remark 8.2.** The matrix M is symmetric and positive definite, cf. [45, Sections 4.1-4.3]. Since  $B^n$  is symmetric we have that

$$\oint_{B_r(x)} (y-x)_i \mathrm{d}y = r \oint_{B^n} z_i \mathrm{d}z = 0, \quad i = 1, \dots, n$$

for  $x \in \mathbb{R}^n$  and r > 0.

In this subsection we show that, when  $u \in W^{2,p}_{loc}(\Omega)$ , we have  $\Delta^w_r u \to L_w u$  in  $L^p_{loc}(\Omega)$ . More specifically, we have the following theorem.

**Theorem 8.2.** Let  $p \in (1, \infty)$ . Then  $W^{1,p}_{loc}(\Omega) \cap AMV^p_{loc}(\Omega_w) = W^{2,p}_{loc}(\Omega)$  and, for every  $u \in W^{2,p}_{loc}(\Omega)$ , we have that  $\Delta^w_r u \xrightarrow{r \to 0} L_w u$  in  $L^p_{loc}(\Omega)$ .

We will prove Theorem 8.2 by reducing it to the unweighted case (using Lemma 8.1), where the result follows by standard theory in a straightforward way.

The following observation is well known, but we briefly recall the proof.

**Lemma 8.3.** If  $u \in C^2(\Omega)$ , then

$$\Delta_r u \to \frac{1}{2} \operatorname{div}(M \nabla u)$$

locally uniformly in  $\Omega$ , as  $r \to 0$ .

*Proof.* Let  $x \in \Omega$  and  $r < \operatorname{dist}(x, \partial \Omega)$ . The Taylor expansion of u yields

$$u(x+rz) - u(x) = r \sum_{i=1}^{n} \partial_i u(x) z_i + \frac{1}{2} r^2 \sum_{i,j=1}^{n} \partial_{ij} u(x) z_i z_j + E_r(x,z),$$

where  $E_r(x,z)/r^2 \to 0$  locally uniformly in  $\Omega \times \overline{B}^n$ , as  $r \to 0$ . Thus we have

$$\Delta_r u(x) = \int_{B^n} \frac{u(x+rz) - u(x)}{r^2} dz = \frac{1}{2} \sum_{i,j=1}^n \partial_{ij} u(x) \int_{B^n} z_i z_j dz + \int_{B^n} \frac{E_r(x,z)}{r^2} dz,$$

from which the claim follows.

**Proposition 8.4.** Let p > 1. Then  $\operatorname{AMV}_{loc}^{p}(\Omega) = W_{loc}^{2,p}(\Omega)$  and, for each  $u \in W_{loc}^{2,p}(\Omega)$ , we have that  $\Delta_{r}u \to \frac{1}{2}\operatorname{div}(M\nabla u)$ (42)

in  $L^p_{loc}(\Omega)$ , as  $r \to 0$ .

*Proof.* Assume  $u \in W^{2,p}_{loc}(\Omega)$  and set

$$R(x,y) := u(y) - u(x) - \sum_{i=1}^{n} \partial_i u(x)(y-x)_i, \quad x, y \in \Omega.$$

Let  $\Omega' \subset \Omega$  be a compactly contained  $W^{2,p}$ -extension domain. Then, for a ball (respectively, a unit sphere)  $B_r^E$  ( $\partial B_1^E$ ) in the Euclidean norm, by [14, Thm 2.5, Thm 4.1 and (21)] we have that

$$\lim_{r \to 0} \int_{\Omega'} \oint_{B_r^E(x)} \left| \frac{R(x,y)}{r^2} \right|^p \mathrm{d}y \mathrm{d}x = c(n,p) \int_{\Omega'} \int_{\partial B_1^E(0)} \left| 2 \sum_{i \le j} \partial_{ij} u(x) e_i e_j \right|^p \mathrm{d}e \,\mathrm{d}x. \tag{43}$$

By Remark 8.2 we see that

$$\int_{B_r(x)} \frac{R(x,y)}{r^2} dy = \Delta_r u(x), \quad \bar{B}_r(x) \subset \Omega.$$
(44)

Together with (43) and the fact that  $\|\cdot\|$  is comparable to the Euclidean norm, (44) implies that

$$\limsup_{r \to 0} \int_{\Omega'} |\Delta_r u|^p \mathrm{d}x \le \limsup_{r \to 0} \int_{\Omega'} \oint_{B_r(x)} \left| \frac{R(x,y)}{r^2} \right|^p \mathrm{d}y \mathrm{d}x \le c \lim_{r \to 0} \int_{\Omega'} \oint_{B_r^E(x)} \left| \frac{R(x,y)}{r^2} \right|^p \mathrm{d}y \mathrm{d}x < \infty.$$

Since any point  $x \in \Omega$  has a neighbourhood  $\Omega'$  which is a  $W^{2,p}$ -extension domain, it follows that  $u \in AMV_{loc}^{p}(\Omega)$ .

Conversely, suppose  $u \in AMV_{loc}^{p}(\Omega)$ . Then, for any positive sequence  $(r_m)$  converging to zero there is a further subsequence and a function  $g \in L_{loc}^{p}(\Omega)$  such that  $\Delta_{r_m} u \rightharpoonup g$  weakly in  $L_{loc}^{p}(\Omega)$  as  $m \rightarrow \infty$ . In particular, for any  $\varphi \in C_c^2(\Omega)$ , we have

$$\int_{\Omega} \varphi g \mathrm{d}x = \lim_{m \to \infty} \int_{\Omega} \varphi \Delta_{r_m} u \mathrm{d}x = \lim_{m \to \infty} \int_{\Omega} \Delta_{r_m} \varphi u \mathrm{d}x = \frac{1}{2} \int_{\Omega} \mathrm{div}(M \nabla \varphi) u \mathrm{d}x,$$

since  $\Delta_r \varphi \to \frac{1}{2} \operatorname{div}(M \nabla \varphi)$  locally uniformly in  $\Omega$ , as  $r \to 0$ . This shows that g is unique and agrees with the distribution  $\frac{1}{2} \operatorname{div}(M \nabla u)$  on  $\Omega$ . Thus  $\operatorname{div}(M \nabla u) \in L^p_{loc}(\Omega)$  and we have that  $u \in W^{2,p}_{loc}(\Omega)$ , see e.g. [33, Theorem 6.29], applied for the differential operator  $P = \Delta$ .

It remains to prove the convergence in (42). For this assume  $u \in W^{2,p}_{loc}(\Omega)$ . Denote by  $\nabla^2 u$  the matrix of second weak partial derivatives of u and by  $Q_s(x)$  a concentric cube centered at x with side length s. Then, by [9, Thm 3.3. lemma 3.4] we observe that there exist constants  $C, \sigma \ge 1$ , such that C = C(n), for which

$$|\Delta_r u(x)| \le \frac{C(n)}{r^2} \int_{Q_{\sigma r}(x)} \frac{|\nabla^2 u|(y)}{|x-y|^{n-2}} \mathrm{d}y \le \mathcal{M}_{\sqrt{n}\sigma r} |\nabla^2 u|(x), \quad B_{\sigma r}(x) \subset \Omega.$$
(45)

Let  $\Omega' \subset \Omega$  be compactly contained and let  $(u_m)$  be a sequence of smooth functions converging to u in  $W^{2,p}(\Omega')$  as  $m \to \infty$ . In particular,  $|\nabla^2(u-u_m)| \to 0$  and  $\operatorname{div}(M\nabla(u-u_m)) \to 0$  in  $L^p(\Omega')$  as  $m \to \infty$ . By Lemma 8.3 and (45) we obtain

$$\begin{split} &\lim_{r \to \infty} \|\Delta_{r} u - \frac{1}{2} \operatorname{div}(M \nabla u)\|_{L^{p}(\Omega')} \\ &\leq \lim_{r \to 0} \sup_{r \to 0} \left( \|\Delta_{r}(u - u_{m})\|_{L^{p}(\Omega')} + \|\Delta_{r} u_{m} + \frac{1}{2} \operatorname{div}(M \nabla u_{m})\|_{L^{p}(\Omega')} + \frac{1}{2} \|\operatorname{div}(M \nabla (u - u_{m}))\|_{L^{p}(\Omega')} \right) \\ &\leq C \lim_{r \to 0} \sup_{r \to 0} \|\mathcal{M}_{\sqrt{n}\sigma r}|\nabla^{2}(u - u_{m})|\|_{L^{p}(\Omega')} + \frac{1}{2} \|\operatorname{div}(M \nabla (u - u_{m}))\|_{L^{p}(\Omega')} \\ &\leq C' \|\nabla^{2}(u - u_{m})\|_{L^{p}(\Omega')} + \frac{1}{2} \|\operatorname{div}(M \nabla (u - u_{m}))\|_{L^{p}(\Omega')}. \end{split}$$

Upon letting  $m \to \infty$  the claim follows.

In order to deal with the "inner product" term in (41) we have the following lemma.

**Lemma 8.5.** If  $f, g \in W^{1,1}_{loc}(\Omega)$  and in addition  $g \in L^n_{loc}(\Omega)$ , then

$$\langle f,g\rangle_r \xrightarrow{r \to 0} \langle M\nabla f,\nabla g\rangle$$

pointwise almost everywhere in  $\Omega$ .

*Proof.* The Sobolev embedding implies that  $f \in L_{loc}^{n/(n-1)}(\Omega)$ . Thus  $fg \in L_{loc}^{1}(\Omega)$  and  $\langle f, g \rangle_r$  is finite. Sobolev functions in  $W_{loc}^{1,1}$  satisfy the following approximation by tangent planes for almost every  $x \in \Omega$ , cf. [26, Chapter 6.2.1 Theorem 2]:

$$\lim_{r \to 0} \left( \int_{B_1^n(x)} \left| \frac{f(x+rz) - f(x) - \nabla f(x) \cdot (rz)}{r} \right|^{n/(n-1)} \mathrm{d}z \right)^{(n-1)/n} = 0.$$
(46)

It follows that

$$\lim_{r \to 0} \left( \oint_{B_1^n(x)} \left| \frac{f(x+rz) - A_r f(x) - \nabla f(x) \cdot (rz)}{r} \right|^{n/(n-1)} \mathrm{d}z \right)^{(n-1)/n} = 0, \tag{47}$$

since

$$\begin{split} & \left( \int_{B_1^n(x)} \left| \frac{f(x+rz) - A_r f(x) - \nabla f(x) \cdot (rz)}{r} \right|^{n/(n-1)} \mathrm{d}z \right)^{(n-1)/n} \\ & \leq \left( \int_{B_1^n(x)} \left| \frac{f(x+rz) - f(x) - \nabla f(x) \cdot (rz)}{r} \right|^{n/(n-1)} \mathrm{d}z \right)^{(n-1)/n} + \left| \frac{A_r f(x) - f(x)}{r} \right| \\ & \leq 2 \left( \int_{B_1^n(x)} \left| \frac{f(x+rz) - f(x) - \nabla f(x) \cdot (rz)}{r} \right|^{n/(n-1)} \mathrm{d}z \right)^{(n-1)/n} . \end{split}$$

For the next calculations we use the shorthand

$$R_r^f(x,z) := \frac{f(x+rz) - A_r f(x) - \nabla f(x) \cdot (rz)}{r}, \quad \bar{B}_r(x) \subset \Omega$$

The definition of the matrix M yields

$$\langle M\nabla f(x), \nabla g(x) \rangle = \int_{B_1^n(x)} \sum_{i,j=1}^n \partial_i f(x) \partial g_j(x) z_i z_j \mathrm{d}z = \int_{B_1^n(x)} (\nabla f(x) \cdot z) (\nabla g \cdot z) \mathrm{d}z$$

which may be expanded to

$$\begin{split} \langle M\nabla f(x), \nabla g(x) \rangle &= \int_{B_1^n(x)} \frac{(f(x+rz) - A_r f(x))}{r} (\nabla g(x) \cdot z) \mathrm{d}z - \int_{B_1^n(x)} R_r^f(x, z) (\nabla g(x) \cdot z) \mathrm{d}z \\ &= \int_{B_1^n(x)} \frac{f(x+rz) - A_r f(x)}{r} \frac{g(x+rz) - A_r g(x)}{r} \mathrm{d}z \\ &- \int_{B_1^n(x)} [R_r^g(x, z) (\nabla f(x) \cdot z) + R_r^f(x, z) (\nabla g(x) \cdot z)] \mathrm{d}z. \end{split}$$

Thus

$$\langle f,g\rangle_r(x) - \langle M\nabla f(x),\nabla g(x)\rangle = \int_{B_1^n(x)} [R_r^g(x,z)(\nabla f(x)\cdot z) + R_r^f(x,z)(\nabla g(x)\cdot z)] \mathrm{d}z$$

tends to zero as  $r \to 0$  for almost every  $x \in \Omega$ , by (47).

The proof of Theorem 8.2 can now be reduced to the unweighted case using the results above. *Proof of Theorem 8.2.* We first observe that, if  $u \in W_{loc}^{1,p}(\Omega)$ , then

$$\begin{aligned} |\langle u, w \rangle_r(x)| &\leq \int_{B_r(x)} \frac{|u - A_r u(x)|}{r} \frac{|w - A_r w(x)|}{r} \mathrm{d}y \\ &\leq \|\nabla w\|_{L^{\infty}(B_r(x))} \int_{B_r(x)} |u - A_r u(x)| / r \mathrm{d}y \leq C \|\nabla w\|_{L^{\infty}(B_r(x))} \mathcal{M}_r |\nabla u|(x). \end{aligned}$$

It follows that, for each compactly contained  $\Omega' \subset \Omega$  there exists a function  $g \in L^p(\Omega')$  such that

$$|\langle u, w \rangle_r| \le g \tag{48}$$

almost everywhere in  $\Omega'$ . Together with Lemma 8.5, (48) implies that

$$\langle u, w \rangle_r \to \langle M \nabla u, \nabla w \rangle \quad \text{in } L^p_{loc}(\Omega)$$

$$\tag{49}$$

Suppose  $u \in W^{2,p}_{loc}(\Omega)$ . Then  $\langle u, w \rangle_r \to \langle M \nabla u, \nabla w \rangle$  and  $\Delta_r u \to \frac{1}{2} \operatorname{div}(M \nabla u)$  in  $L^p_{loc}(\Omega)$ . Thus Lemma 8.1 (41) implies that

$$\Delta_r^w u \xrightarrow{r \to 0} L_w u$$

in  $L^p_{loc}(\Omega)$ , which also implies that  $u \in AMV^p_{loc}(\Omega_w)$ .

Conversely, if  $u \in W^{1,p}_{loc}(\Omega) \cap AMV^p_{loc}(\Omega_w)$ , Lemma 8.1(41) and (48) imply that, for every compactly contained  $\Omega' \subset \Omega$  there exists a function  $g \in L^p(\Omega')$  such that

$$|\Delta_r u| \le |\Delta_r^w u| + \frac{1}{A_r w}g$$
 on  $\Omega'$ .

Thus  $u \in AMV_{loc}^{p}(\Omega)$  which, by Proposition 8.4, implies that  $u \in W_{loc}^{2,p}(\Omega)$ .

Using Lemma 8.1 (40) it is possible to prove that, for  $w \in W^{2,\infty}_{loc}(\Omega)$ , the classes  $AMV^p_{loc}(\Omega_w)$  agree with  $W_{loc}^{2,p}(\Omega)$ .

**Theorem 8.6.** Suppose  $w \in W^{2,\infty}_{loc}(\Omega_w)$ , and  $p \in (1,\infty)$ . Then  $\operatorname{AMV}^p_{loc}(\Omega_w) = W^{2,p}_{loc}(\Omega)$ . For every  $u \in \operatorname{AMV}^p_{loc}(\Omega_w)$  we have that  $\Delta^w_r u \to L_w u$  in  $L^p_{loc}(\Omega)$ , as  $r \to 0$ .

*Proof.* We sketch the proof and omit the details.

Identity (40) yields  $|\Delta_r^w(uw)| \leq A_r w |\Delta_r^w u| + |u| |\Delta_r w|$  for  $u \in AMV_{loc}^p(\Omega_w)$ . By Proposition 8.4 we have that  $uw \in W_{loc}^{2,p}(\Omega)$  which implies  $u \in W_{loc}^{2,p}(\Omega)$  since  $w^{-1} \in W_{loc}^{2,\infty}(\Omega)$ . Using (40) and Proposition 8.4 we have the convergence

$$\Delta_r^w u \xrightarrow{r \to 0} \frac{1}{2w} (\operatorname{div}(M\nabla(uw))) - u \operatorname{div}(M\nabla w)) = \frac{1}{2w} (u \operatorname{div}(M\nabla w) + 2\langle \nabla w, M\nabla u \rangle + w \operatorname{div}(M\nabla u) - u \operatorname{div}(M\nabla w)) = L_w u \text{whenever } u \in W^{2,p}_{L_v}(\Omega).$$

in  $L^p_{loc}(\Omega)$ , whenever  $u \in W^{2,p}_{loc}(\Omega)$ .

8.2. PDE characterization of amv-harmonicity. We are now in a position to prove Theorem 1.9. For clarity, we formulate the following weak counterpart of Proposition 8.4 as a separate lemma. We omit the proof, which follows e.g. by an approximation argument from Proposition 8.4, or via a direct argument in the spirit of Proposition 8.4.

**Lemma 8.7.** Let  $u \in W^{1,2}_{loc}(\Omega)$ . Then  $\Delta_r u \to \frac{1}{2} \operatorname{div}(M \nabla u)$  weakly, i.e.

$$\lim_{r \to 0} \int_{\Omega} \varphi \Delta_r u \mathrm{d}x = -\frac{1}{2} \int \langle \nabla \varphi, M \nabla u \rangle \mathrm{d}x$$

for every  $\varphi \in C_c^{\infty}(\Omega)$ .

*Proof of Theorem 1.9.* We first prove the equivalences (1)–(3). The implication (3)  $\implies$  (2) is clear, and  $(2) \implies (1)$  follows from Lemmas 8.7 and 8.1(41). Indeed, together with (49) in the proof of Theorem 8.2, we have that

$$0 = \lim_{r \to 0} \int_{\Omega} \varphi \Delta_r^w u dx = \lim_{r \to 0} \left( \int_{\Omega} \varphi \Delta_r u \, dx + \int_{\Omega} \frac{\varphi}{A_r w} \langle u, w \rangle_r \, dx \right)$$
$$= -\frac{1}{2} \int_{\Omega} \langle \nabla \varphi, M \nabla u \rangle dx + \int_{\Omega} \frac{\varphi}{w} \langle \nabla w, M \nabla u \rangle dx$$

for every  $\varphi \in C_c^{\infty}(\Omega)$ , which is the weak formulation of  $L_w u = 0$ .

Thus it suffices to prove the implication (1)  $\implies$  (3). By elementary regularity theory, see e.g. [32, Thm 8.8], a weak solution  $u \in W_{loc}^{1,2}(\Omega)$  to  $L_w u = 0$  satisfies  $u \in W_{loc}^{2,2}(\Omega)$ . Theorem 8.2 now directly implies that (3) holds.

43

It remains to prove the equivalence of (1)–(3) with (4) assuming smoothness of w. Since (4)  $\implies$  (2), it suffices to prove that (1)  $\implies$  (4). Since weak solutions of  $L_w u = 0$  with  $w \in C^{\infty}(\Omega)$  are smooth (see e.g. [32, Cor 8.11]), Lemma 8.3 and 8.1 (40) imply that

$$\Delta_r^w u = \frac{1}{A_r w} (\Delta_r(uw) - u\Delta_r w) \to \frac{1}{2w} (\operatorname{div}(M\nabla(uw)) - u\operatorname{div}(M\nabla w)) = L_w u = 0$$

locally uniformly in  $\Omega$ , as  $r \to 0$ .

**Remark 8.3.** The assumption  $w \in C^{\infty}(\Omega)$  in the second part of Theorem 1.9 is by no means optimal, but is sufficient for the purposes of the discussion at the end of the introduction, where it was pointed out that the (weak or strong) amv-property need not hold for harmonic functions on RCD-spaces without the non-collapsing condition.

#### REFERENCES

- T. ADAMOWICZ, M. GACZKOWSKI, P. GÓRKA, Harmonic functions on metric measure spaces, Rev. Mat. Complut. 32(1) (2019), 141–186.
- [2] T. ADAMOWICZ, B. WARHURST, Mean value property and harmonicity on Carnot–Carathéodory groups, Potential Anal. 52 (3) (2020), 497–525. doi.org/10.1007/s11118-018-9740-4.
- [3] J. M. ALDAZ, Boundedness of averaging operators on geometrically doubling metric spaces Ann. Acad. Sci. Fenn. Math. 44(1) (2019), 497–503.
- [4] L. AMBROSIO, M. COLOMBO, S. DI MARINO, Sobolev spaces in metric measure spaces: reflexivity and lower semicontinuity of slope, Adv. Stud. Pure Math., 67 (2015), 1–58.
- [5] L. AMBROSIO, N. GIGLI, G. SAVARÉ, Metric measure spaces with Riemannian Ricci curvature bounded from below, Duke Math. J., 163(7) (2014), 1405–1490.
- [6] L. AMBROSIO, N. GIGLI, G. SAVARÉ, Calculus and heat flow in metric measure spaces and applications to spaces with Ricci bounds from below, Invent. Math., 195(2) (2014), 289–391.
- [7] L. AMBROSIO, N. GIGLI, G. SAVARÉ, Bakry-Émery curvature-dimension condition and Riemannian Ricci curvature bounds, Ann. Probab. 43(1) (2015), 339–404.
- [8] A. BJÖRN, J. BJÖRN, *Nonlinear Potential Theory on Metric Spaces*, EMS Tracts in Mathematics, 17, European Math. Soc., Zurich, 2011.
- [9] B. BOJARSKI, P. HAJŁASZ, P. STRZELECKI, Improved C<sup>k,λ</sup> approximation of higher order Sobolev functions in norm and capacity, Indiana Univ. Math. J. 51(3) (2002), 507–540.
- [10] B. BOJARSKI, L. IHNATSYEVA, J. KINNUNEN, How to recognize polynomials in higher order Sobolev spaces Math. Scand. 112(2) (2013), 161–181.
- [11] A. BONFIGLIOLI, E. LANCONELLI, F. UGUZZONI, Stratified Lie Groups and Potential Theory for Their Sub-Laplacians, Springer Monographs in Mathematics, Springer, 2007.
- [12] N. BOULEAU, F. HIRSCH, Dirichlet forms and analysis on Wiener space, De Gruyter Studies in Mathematics, 14, Berlin, 1991. x+325 pp.
- [13] A. BOSE, Functions satisfying a weighted average property, Trans. Amer. Math. Soc. 118 (1965), 472–487.
- [14] D. BURAGO, S. IVANOV, Y. KURYLEV, Spectral stability of metric-measure Laplacians Israel J. Math. 232(1) (2019), 125– 158.
- [15] S. BUCKLEY, Is the maximal function of a Lipschitz function continuous?, Ann. Acad. Sci. Fenn. Math. 24(2) (1999), 519– 528.
- [16] L. CAPOGNA, Regularity of quasi-linear equations in the Heisenberg group, Comm. Pure Appl. Math. 50(9) (1997), 867– 889.
- [17] L. CAPOGNA, M. COWLING, Conformality and Q-harmonicity in Carnot groups, Duke Math. J. 135(3) (2006), 455–479.
- [18] J. CHEEGER, Differentiability of Lipschitz functions on metric measure spaces, Geom. Funct. Anal. 9(3) (1999), 428–517.
- [19] T. H. COLDING, W. P. MINICOZZI II, Harmonic Functions on Manifolds, Annals of Math. (2) 146(3), 725–747 (1997).
- [20] T. H. COLDING, W. P. MINICOZZI II, *Harmonic functions with polynomial growth*, J. Differential Geom. 46(1) (1997), 1–77.
- [21] A. CÓRDOBA, J. OCÁRIZ, A note on generalized laplacians and minimal surfaces, arXiv:1812.07897 to appear in Bull. Lond. Math. Soc., doi: 10.1112/blms.12314.
- [22] G. DE PHILIPPIS, N. GIGLI, Non-collapsed spaces with Ricci curvature bounded from below, J. Éc. polytech. Math. 5 (2018), 613–650.
- [23] S. ERIKSSON-BIQUE, J. GILL, P. LAHTI, N. SHANMUGALINGAM, Asymptotic behavior of BV functions and sets of finite perimeter in metric measure spaces, arXiv:1810.05310.

- [24] F. FERRARI, Q. LIU, J. MANFREDI, On the characterization of p-harmonic functions on the Heisenberg group by mean value properties, Discrete Contin. Dyn. Syst. 34 (7) (2014), 2779–2793.
- [25] F. FERRARI, A. PINAMONTI, Characterization by asymptotic mean formulas of q-harmonic functions in Carnot groups, Potential Anal. 42(1) (2015), 203–227.
- [26] L. EVANS, R. GARIEPY, *Measure theory and fine properties of functions* (Revised edition) ,Textbooks in Mathematics, CRC Press, Boca Raton, FL, 2015.
- [27] B. GAVEAU, J. VAUTHIER, The Dirichlet problem for the subelliptic Laplacian on the Heisenberg group. II, Canad. J. Math. 37(4) (1985), 760–766.
- [28] N. GIGLI, On the differential structure of metric measure spaces and applications, Mem. Amer. Math. Soc. 236 (2015), no. 1113, vi+91 pp.
- [29] N. GIGLI, Lecture notes on differential calculus on RCD spaces, Publ. Res. Inst. Math. Sci. 54(4) (2018), 855–918.
- [30] N. GIGLI, Nonsmooth differential geometry An approach tailored for spaces with Ricci curvature bounded from below, Mem. Amer. Math. Soc. 251 (2018), no. 1196, v+161 pp.
- [31] N. GIGLI, A. TYULENEV, Korevaar–Schoen's energy on strongly rectifiable spaces, arXiv:2002.07440v1.
- [32] D. GILBARG, N. S. TRUDINGER, *Elliptic Partial Differential Equations of Second Order*. Reprint of the 1998 edition. Classics in Mathematics. Springer-Verlag, Berlin, 2001. xiv+517 pp.
- [33] G. GRUBB, Distributions and operators, Graduate Texts in Mathematics, 252. Springer, New York, 2009. xii+461 pp.
- [34] P. HAJŁASZ, Sobolev spaces on metric-measure spaces. Heat kernels and analysis on manifolds, graphs, and metric spaces (Paris, 2002), 173–218, Contemp. Math., 338, Amer. Math. Soc., Providence, RI, 2003.
- [35] P. HAJŁASZ, J. KINNUNEN, Holder quasicontinuity of Sobolev functions on metric spaces, Rev. Mat. Iberoamericana 14(3) (1998), 601–622.
- [36] P. HAJŁASZ, P. KOSKELA, Sobolev Met Poincaré, Mem. Amer. Math. Soc., 145 (2000).
- [37] W. HANSEN, H. HUEBER, The Dirichlet problem for sub-Laplacians on nilpotent Lie groups–geometric criteria for regularity, Math. Ann. 276(4) (1987), 537–547.
- [38] J. HEINONEN, P. KOSKELA, N. SHANMUGALINGAM, J. TYSON, *Sobolev spaces on metric measure spaces. An approach based on upper gradients*, New Mathematical Monographs, 27. Cambridge University Press, Cambridge, 2015.
- [39] B. HUA, M. KELL, C. XIA, Harmonic functions on metric measure spaces, arXiv:1308.3607.
- [40] B. HUA, Harmonic functions of polynomial growth on singular spaces with nonnegative Ricci curvature, Proc. Amer. Math. Soc. 139(6) (2011), 2191–2205.
- [41] V.KAPOVITCH, A.LYTCHAK, A.PETRUNIN, *Metric-Measure Boundary and Geodesic Flow on Alexandrov spaces*, to appear in J. Eur. Math. Soc.
- [42] V. KAPOVITCH, A. MONDINO, On the topology and the boundary of N-dimensional RCD(K,N) spaces, arXiv:1907.02614.
- [43] S. KEITH, Modulus and the Poincaré inequality on metric measure spaces, Math. Z. 245(2) (2003), 255–292.
- [44] S. KEITH, A differentiable structure for metric measure spaces, Adv. Math. 183(2) (2004), 271–315.
- [45] A. KIJOWSKI, Characterization of mean value harmonic functions on norm induced metric measure spaces with weighted Lebesgue measure, Electron. J. Differential Equations 2020, Paper No. 8, 26 pp.
- [46] B. KIRCHHEIM, Rectifiable metric spaces: local structure and regularity of the Hausdorff measure, Proc. Amer. Math. Soc. 121(1) (1994), 113–123.
- [47] B. KLEINER, A new proof of Gromov's theorem on groups of polynomial growth, J. Amer. Math. Soc. 23(3) (2010), 815–829.
- [48] N. KOREVAAR, R. SCHOEN, Sobolev spaces and harmonic maps for metric space targets, Comm. Anal. Geom. 1(3-4) (1993), 561–659.
- [49] P. KOSKELA, K. RAJALA, N. SHANMUGALINGAM, Lipschitz continuity of Cheeger-harmonic functions in metric measure spaces, J. Funct. Anal. 202(1) (2003), 147–173.
- [50] P. KOSKELA, D. YANG, Y. ZHOU, A characterization of Hajłasz–Sobolev and Triebel–Lizorkin spaces via grand Littlewood– Paley functions, J. Funct. Anal. 258(8) (2010), 2637–2661.
- [51] P. KOSKELA, D. YANG, Y. ZHOU, Pointwise characterizations of Besov and Triebel–Lizorkin spaces and quasiconformal mappings, Adv. Math. 226(4) (2011), 3579–3621.
- [52] P. LI, Harmonic sections of polynomial growth, Math. Res. Lett. 4(1) (1997), 35–44.
- [53] P. LI, *Harmonic functions and applications to complete manifolds,* XIV Escola de Geometria Diferencial. [XIV School of Differential Geometry] Instituto de Matemática Pura e Aplicada (IMPA), Rio de Janeiro, 2006.
- [54] P. LI, Geometric analysis. Cambridge Studies in Advanced Mathematics, 134. Cambridge University Press, Cambridge, 2012. x+406.
- [55] J. LLORENTE, Mean value properties and unique continuation, Commun. Pure Appl. Anal. 14(1) (2015), 185–199.
- [56] J. MANFREDI, G. MINGIONE, Regularity results for quasilinear elliptic equations in the Heisenberg group, Math. Ann. 339(3) (2007), 485–544.
- [57] A. MINNE, D. TEWODROSE, Asymptotic Mean Value Laplacian in Metric Measure Spaces, arXiv:1912.00259.
- [58] R. MONTI, F. SERRA CASSANO, Surface measures in Carnot-Carathéodory spaces, Calc. Var. Partial Differential Equations 13(3) (2001), 339–376.

## TOMASZ ADAMOWICZ, ANTONI KIJOWSKI, AND ELEFTERIOS SOULTANIS $^{\rm 2}$

- [59] F. PACARD, X. XU, Constant mean curvature spheres in Riemannian manifolds, Manuscripta Math. 128(3) (2009), 275–295.
- [60] A. PETRUNIN, *Alexandrov meets Lott–Villani–Sturm*, Münster J. of Math. 4 (2011), 53–64, http://cn.arxiv.org/abs/1003.5948v1.
- [61] I. PRIVALOFF, Sur les fonctions harmoniques, Rec. Math. Moscou (Mat. Sbornik), 32 (1925), 464-471.
- [62] N. SHANMUGALINGAM, Newtonian spaces: an extension of Sobolev spaces to metric measure spaces, Rev. Mat. Iberoamericana 16(2) (2000), 243–279.
- [63] K.-T. STURM, On the geometry of metric measure spaces. I, Acta Math. 196(1) (2006), 65–131.
- [64] K.-T. STURM, On the geometry of metric measure spaces. II. Acta Math. 196(1) (2006), 133–177.
- [65] K.-H. TAN, Sobolev classes and horizontal energy minimizers between Carnot-Carathéodory spaces, Commun. Contemp. Math. 6(2) (2004), 315–347.
- [66] C. VILLANI, Synthetic theory of Ricci curvature bounds, Jpn. J. Math. 11(2) (2016), 219–263.
- [67] F.-Y. WANG, Equivalent semigroup properties for the curvature-dimension condition, Bull. Sci. Math. 135(6-7) (2011), 803–815.
- [68] F. W. WARNER, Foundations of differentiable manifolds and Lie groups. Corrected reprint of the 1971 edition, Graduate Texts in Mathematics, 94. Springer-Verlag, New York-Berlin, 1983.
- [69] D. YANG, New characterizations of Hajłasz-Sobolev spaces on metric spaces, Sci. China Ser. A 46(5) (2003), 675–689.
- [70] H.-C. ZHANG, X. ZHONG, X.-P. ZHU, Quantitative gradient estimates for harmonic maps into singular spaces, Sci. China Math. 62(11) (2019), 2371–2400.
- [71] H.-C. ZHANG, X.-P. ZHU, Yau's gradient estimates on Alexandrov spaces, J. Differential Geom. 91(3) (2012), 445–522.

T.A.: INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES, ŚNIADECKICH 8, WARSAW, 00-656, POLAND *E-mail address*: tadamowi@impan.pl

A.K.: INSTITUTE OF MATHEMATICS, POLISH ACADEMY OF SCIENCES, ŚNIADECKICH 8, WARSAW, 00-656, POLAND *E-mail address*: akijowski@impan.pl

E.S.: UNIVERSITY OF FRIBOURG, DEPARTMENT OF MATHEMATICS, CHEMIN DU MUSÉE 23, CH-1700 FRIBOURG *E-mail address*: elefterios.soultanis@gmail.com