

A CLASS OF HIGHER INDUCTIVE TYPES IN ZERMELO-FRAENKEL SET THEORY

ANDREW W SWAN

ABSTRACT. We define a class of higher inductive types that can be constructed in the category of sets under the assumptions of Zermelo-Fraenkel set theory without the axiom of choice or the existence of uncountable regular cardinals. This class includes the example of unordered trees of any arity.

1. INTRODUCTION

Higher inductive types are one of the key ideas in homotopy type theory [Uni13]. We think of an (ordinary) inductive type as the smallest type closed under certain algebraic operations or *point constructors*. For instance, we define the type of countably branching trees T to be the smallest type closed under the following operations.

$$\begin{aligned} \text{leaf} &: T \\ \text{node} &: (\omega \rightarrow T) \rightarrow T \end{aligned}$$

Within type theory we formalise the idea that T is the smallest type with the above point constructors using recursion or induction terms. However, semantically, it is often more convenient to think in terms of initial algebras. We say an *algebra* for the above constructors is a type X together with a map $1 + X^\omega \rightarrow X$. T is then the initial object in the category of algebras. This is a classic example of a *W-type*, as defined by Moerdijk and Palmgren [MP00].

For higher inductive types, one not only has point constructors, but also *path constructors*, which add proofs of identities of terms. Higher inductive types are usually considered within HoTT and have well understood semantics within models of HoTT [LS20], [CHM18], [CH19]. However, since they are defined within the language of type theory, one might also consider whether they hold in interpretations of extensional type theory in locally cartesian closed categories, and in particular the category of sets, **Set**.

One of the simplest examples of higher inductive type is pushouts. In **Set** these can be implemented as pushouts in the usual categorical sense. It follows that **Set** contains all of the n -dimensional spheres, although there is not much you can say about them without the univalence axiom, and indeed they turn out to be trivial in **Set**. Quotients and image factorisations are examples of simple colimits that play a useful role even within models of extensional type theory [Mai05], [AB04].

There are also more complicated examples of higher inductive types that are non trivial in extensional type theory, and even **Set**, within the framework of *quotient inductive types* [AK16]. In fact our examples of interest fall within a smaller class

Date: July 22, 2021.

with a simpler definition and clearer semantics. This class was studied by Blass [Bla83] under the name *free algebras subject to identities* and by Fiore, Pitts and Steenkamp in [FPS20] under the name *QW-types* (we will refer to them by the latter name). A well known example of such a type is that of “unordered countably branching trees.” We modify the definition of T above to get the higher inductive type T_{Sym} by adding equations as follows, where we write $\text{Sym}(\omega)$ for the type of permutations $\omega \rightarrow \omega$.

$$\begin{aligned} \text{leaf} &: T_{\text{Sym}} \\ \text{node} &: (\omega \rightarrow T_{\text{Sym}}) \rightarrow T_{\text{Sym}} \\ \text{perm} &: \prod_{f: \omega \rightarrow T_{\text{Sym}}} \prod_{\pi: \text{Sym}(\omega)} \text{node}(f) = \text{node}(f \circ \pi) \end{aligned}$$

Altenkirch, Capriotti, Dijkstra, Kraus and Forsberg include this in [ACD⁺18], as a non trivial example of a quotient inductive(-inductive) type. As they remark, the obvious construction of T_{Sym} as a quotient of T requires the axiom of choice.¹ Fiore, Pitts and Steenkamp showed that in fact it is an example of a *QW-type* [FPS20, Example 2].

Blass showed that all *QW-types* can be constructed in **Set** under the assumption that regular cardinals are unbounded in the class of all ordinals. More generally free algebras can be constructed in cocomplete categories from the existence of regular cardinals of sufficiently high cardinality via the general techniques of Kelly [Kel80]. For example this plays an important role in the construction of higher inductive types by Lumsdaine and Shulman [LS20]. The existence of an unbounded class of regular ordinals is usually a reasonable one. It follows from very weak versions of choice, such as **WISC** [vdB12] and a variant is often assumed in constructive set theory [AR01, Section 10]. It is also the case that every inaccessible cardinal is in particular regular.²

However, Gitik [Git80] has constructed a model of **ZF** in which ω is the only regular cardinal.³ Moreover, Blass showed that the assumption is strictly necessary, by constructing a *QW-type* which is isomorphic to the collection of ordinals hereditarily of countable cofinality, if it exists. He deduced by Gitik’s result that this gives an example of a *QW-type* that does not provably exist in **Set** under the assumptions of **ZF**.⁴

Fiore, Pitts and Steenkamp in loc. cit. gave an electronically verified proof that *QW-types* can be constructed in type theory using Agda sized types and universes closed under inductive-inductive types. We can see from Blass’ counterexample that some combination of these assumptions for **Set** must lead to the existence of uncountable regular cardinals. In a second paper [FPS21] the same authors showed

¹For the special case above, countable choice would be enough.

²Note, however that even in the presence of inaccessible cardinals it can be useful to have proofs that are valid in **ZF** without further assumption: if κ is inaccessible, then V_κ is a transitive model of **ZF**, and so any proof valid in **ZF** can be carried out inside it (e.g. to construct HITs that belong to V_κ), but V_κ itself does not contain inaccessible cardinals without further assumptions on κ .

³under certain large cardinal assumptions

⁴Using Blass’ argument and a later paper by Gitik [Git85] one can also show the following. For every successor ordinal α there is a transitive model of **ZF** where Blass’ example of a *QW-type* exists and is precisely the set of all ordinals less than ω_α .

that the weak choice principle **WISC** can be used in place of Agda sized types, again verifying the result electronically.

On the other hand, some higher inductive types can be constructed in **Set** without choice or unbounded regular cardinals. In addition to colimits, as mentioned above, the author showed in [Swa18] that W -types with reductions exist in any boolean topos, including **Set**. A similar argument shows that Sojakova's notion of W -suspensions [Soj15] also exist in all boolean toposes.⁵

In this paper we will see a new class of QW -types that can be constructed in **Set** under **ZF**, without any assumptions of choice or existence of regular cardinals, that we call *image preserving* QW -types. This will include the example of unordered countably branching trees above, and more generally unordered trees of any arity. The proof is based on a construction of a set of all hereditarily countable sets due to Jech [Jec82]. We will be able to recover hereditarily countable sets as a special case, and moreover a later generalisation due to Holmes [Hol14].

Acknowledgements. I am grateful to Jonas Frey, Simon Henry and Thomas Streicher for helpful discussion and suggestions.

I gratefully acknowledge the support of the Air Force Office of Scientific Research through MURI grant FA9550-15-1-0053. Any opinions, findings and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the AFOSR.

2. IMAGE PRESERVING QW -TYPES

We now define our class of higher inductive types that we will construct in **Set**. It will be clear by the definition that this is a special case of QW -types [FPS20].

Definition 2.1. A *polynomial* is a set A together with a family of sets $(B_a)_{a \in A}$. We will refer to elements of A as *constructors* and say B_a is the *arity* of the constructor $a \in A$.

Definition 2.2. Given a polynomial $(B_a)_{a \in A}$, an *algebra* is a set X together with a function $s: \sum_{a \in A} X^{B_a} \rightarrow X$. We refer to such s as an *algebra structure* on X .

If (X, s) and (Y, t) are algebras, we say a *homomorphism* is a function $h: X \rightarrow Y$ such that for all $a \in A$ and $f: B_a \rightarrow X$ we have $h(s(a, f)) = t(a, h \circ f)$.

Remark 2.3. Although we assumed X and Y are sets in the definition above, we can also define algebra structures and homomorphisms for classes by replacing functions with class functions.

Definition 2.4. Given a polynomial $(B_a)_{a \in A}$, an *image preserving equation* over $(B_a)_{a \in A}$ consists of a set V , and $a, b \in A$ together with $l: B_a \rightarrow V$ and $r: B_b \rightarrow V$ such that the image of l is equal to the image of r .

A *family of image preserving equations* consists of a set E together with a family of image preserving equations $(V_e, a_e, b_e, l_e, r_e)_{e \in E}$.

Definition 2.5. Given a polynomial $(B_a)_{a \in A}$ and a family of image preserving equations, $(V_e, a_e, b_e, l_e, r_e)_{e \in E}$, an *algebra* is an algebra (X, s) for the polynomial $(B_a)_{a \in A}$ such that for every $e \in E$ and every function $h: V_e \rightarrow X$ we have $s(a_e, h \circ l_e) = s(b_e, h \circ r_e)$.

⁵A proof is left as an exercise for the reader.

Example 2.6. Suppose we are given a set B . We consider the polynomial with two constructors of arity 0 and B . We consider the set of image preserving equations with set of variables B , and $l, r: B \rightarrow B$ defined by $l = 1_B$ and $r = \pi$ for each permutation $\pi \in \text{Sym}(B)$. The initial algebra is then the set of *unordered trees of arity B* . In particular, we can take $B = \omega$ to get unordered countably branching trees.

Example 2.7. Given any polynomial $(B_a)_{a \in A}$ and set V we can consider the set of *all* image preserving equations. We will see in the next section how this allows us to recover hereditarily countable sets and more generally hereditarily small sets.

Example 2.8. We will be able to deduce from the main theorem that Blass' example of a QW -type that cannot be constructed in \mathbf{ZF} cannot be viewed as an image preserving QW -type. However, for illumination we will give a more intuitive direct reason why it does not satisfy the definition. In Blass' example, the initial algebra is expected to behave like the collection of all ordinals of countable cofinality. In particular there is an operation sup which takes a sequence $(\alpha_n)_{n < \omega}$ and is expected to behave like the supremum of the countable sequence of ordinals $(\alpha_n)_{n < \omega}$. In particular we should identify $\text{sup}((\alpha_n)_{n < \omega})$ and $\text{sup}((\beta_n)_{n < \omega})$ whenever α_n is cofinal in β_n and vice versa. However, this is much weaker than α_n and β_n containing exactly the same elements (possibly in a different order).

3. HEREDITARILY SMALL SETS

As part of [Jec82] Jech showed how to construct the set of all hereditarily countable sets in \mathbf{ZF} . This was later generalised by Holmes [Hol14] who showed that for any set B we can construct a set containing all sets hereditarily with cardinality less than or equal to B . We in fact give a very slight further generalisation of Holmes' result by instead considering families of sets $(B_a)_{a \in A}$. In order to derive this from the main theorem 5.12, we first need to clarify the relation between image preserving equations and the class of hereditarily small sets, which is the topic of this section.

Definition 3.1. Given a family of sets $(B_a)_{a \in A}$ we say a set X is *small relative to $(B_a)_{a \in A}$* if for some $a \in A$ there exists a surjection $B_a \rightarrow X$. We say X is *hereditarily small relative to $(B_a)_{a \in A}$* if X is small relative to $(B_a)_{a \in A}$ and all of its elements are hereditarily small relative to $(B_a)_{a \in A}$. We write $H((B_a)_{a \in A})$ for the class of hereditarily small sets.

We view $H((B_a)_{a \in A})$ as an algebra on the polynomial $(B_a)_{a \in A}$ as follows. Given $a \in A$ and $f: B_a \rightarrow H((B_a)_{a \in A})$, we define $s(a, f)$ to be $\{f(b) \mid b \in B_a\}$.

Lemma 3.2. $H((B_a)_{a \in A})$ satisfies all image preserving equations relative to the polynomial $(B_a)_{a \in A}$.

Proof. Suppose we are given a set S of variables together with $l: B_a \rightarrow S$ and $r: B_c \rightarrow S$ and a map $h: S \rightarrow H((B_a)_{a \in A})$. We calculate as follows.

$$\begin{aligned} s(h \circ l) &= \{h(l(b)) \mid b \in B_a\} \\ &= \{h(x) \mid x \in \text{im}(l)\} \\ &= \{h(x) \mid x \in \text{im}(r)\} \\ &= s(h \circ r) \end{aligned}$$

□

We now fix a set of variables $S := \sum_{a \in A} \mathcal{P}(B_a)$.

Lemma 3.3. *Suppose (X, t) is an algebra for $(B_a)_{a \in A}$ that satisfies all image preserving equations for the set of variables S above. Then there is a unique homomorphism $h : H((B_a)_{a \in A}) \rightarrow X$.*

Proof. In order for h to be a homomorphism, we need it to satisfy the following equation whenever $a \in A$ and $f : B_a \rightarrow H((B_a)_{a \in A})$.

$$h(\{f(b) \mid b \in B_a\}) = t(a, h \circ f)$$

This defines a unique class function by \in -induction as long as the equation above is well defined. That is, given $g : B_c \rightarrow H((B_a)_{a \in A})$ such that $\{f(b) \mid b \in B_a\} = \{g(b) \mid b \in B_c\}$ we need $t(a, h \circ f) = t(c, h \circ g)$.

Note that the lemma is trivial if B_a is inhabited for each a , since then $H((B_a)_{a \in A})$ is empty. Hence we may assume B_a is empty for some a and so X has a canonical element x_0 . Now fix $a, c \in A$ and f and g as above. By induction on rank, we may assume $h(y)$ is already uniquely defined for $y \in \{f(b) \mid b \in B_a\}$. We define a map $k : S \rightarrow X$ as follows.

$$k(a', C) = \begin{cases} h(y) & a' = a, C = f^{-1}(y) \text{ for unique } y \in \{f(b) \mid b \in B_a\} \\ x_0 & \text{otherwise} \end{cases}$$

We next define $l : B_a \rightarrow S$ and $r : B_c \rightarrow S$ as follows.

$$l(b) := (a, \{b' \in B_a \mid f(b') = f(b)\})$$

$$r(b) := (a, \{b' \in B_a \mid f(b') = g(b)\})$$

Finally, it is straightforward to verify that l and r have the same image in S , and that $k \circ l = h \circ f$ and $k \circ r = h \circ g$. Hence it follows from the image preserving equation $t(a, k \circ l) = t(c, k \circ r)$ that $t(a, h \circ f) = t(c, h \circ g)$ as required. \square

Theorem 3.4. *If $H((B_a)_{a \in A})$ is a set, then it is the QW-type for the polynomial $(B_a)_{a \in A}$ with all image preserving equations for S .*

Proof. This follows directly from lemmas 3.2 and 3.3. \square

Theorem 3.5. *If the QW-type for the polynomial $(B_a)_{a \in A}$ with all image preserving equations for S exists, then $H((B_a)_{a \in A})$ is a set.*

Proof. Let (X, t) be the QW-type.

We first define a homomorphism from X to $H((B_a)_{a \in A})$. The essential idea is to use the fact that X is an initial algebra. However, formally speaking we can only apply initiality of X once we know that $H((B_a)_{a \in A})$ is an algebra and in particular a set, which we don't have a priori. Hence we use a trick of approximating $H((B_a)_{a \in A})$ by a sequence of sets.

For each ordinal α we can define an algebra structure on the set $(H((B_a)_{a \in A}) \cap V_\alpha) + 1$ that agrees with the structure s on $H((B_a)_{a \in A})$ when $s(a, f)$ belongs to $H((B_a)_{a \in A}) \cap V_\alpha$ and otherwise sends (a, f) to the 1 component. Note furthermore that this operation satisfies all image preserving equations by lemma 3.2. Hence there is a unique homomorphism $h_\alpha : X \rightarrow (H((B_a)_{a \in A}) \cap V_\alpha) + 1$. Say that $x \in X$ is *defined at α* if $h(x) \in H((B_a)_{a \in A}) \cap V_\alpha$ and undefined otherwise. Note that for $\beta \geq \alpha$ the canonical restriction map $(H((B_a)_{a \in A}) \cap V_\beta) + 1 \rightarrow (H((B_a)_{a \in A}) \cap V_\alpha) + 1$ is a homomorphism. It follows that if x is defined at α , then it is also defined at β and $h_\beta(x) = h_\alpha(x)$. By induction every element x is defined at α for some

ordinal α . Hence this defines a unique homomorphism $X \rightarrow H((B_a)_{a \in A})$. The homomorphism has an inverse by lemma 3.3. Hence $H((B_a)_{a \in A})$ is in bijection with a set, and so a set itself. \square

4. SOME USEFUL PROPOSITIONS

We recall some basic classical set theory that will be useful for the construction of image preserving QW -types. We fill in some of the details, with the remainder left as an exercise for the reader.

Proposition 4.1. *For any ordinals $0 < \alpha < \beta$, there is a canonical surjection $\beta \twoheadrightarrow \alpha$.*

If there is a surjection $X \twoheadrightarrow \beta$ for some set X , then there is also a surjection $X \twoheadrightarrow \alpha$.

Proposition 4.2. *For any well ordered set $(X, <)$ (and in particular for sets of ordinals ordered by \in), there is a unique ordinal β with a unique order isomorphism $(X, <) \cong (\beta, \in)$. We refer to β as the order type of $(X, <)$.*

Proposition 4.3. *For any family of sets $(X_i)_{i \in I}$, there is an ordinal $\aleph((X_i)_{i \in I})$ which is the smallest for which there is no surjection $X_i \twoheadrightarrow \aleph((X_i)_{i \in I})$ for any $i \in I$. It is precisely the set of all ordinals α for which there is a surjection $X_i \twoheadrightarrow \alpha$ for some $i \in I$.*

Proof. Note that whenever $X_i \twoheadrightarrow \alpha$, there is an equivalence relation \sim on X , and a well ordering $<$ on X/\sim such that the order type of $(X/\sim, <)$ is α . However there is clearly a set of such well orders by power set, and so there is a set of all such ordinals α . Since this is a downwards closed set of ordinals, it is an ordinal itself. Since the set cannot contain itself, it is the least ordinal for which there is no surjection from X_i for any i . \square

Proposition 4.4. *If κ is a cardinal number (i.e. an ordinal that is not in bijection with any lower ordinal), then one can define surjections*

- (1) $\kappa \twoheadrightarrow \kappa \times \kappa$
- (2) $\kappa \twoheadrightarrow \kappa^n$ for any $n < \omega$
- (3) $\kappa \twoheadrightarrow \omega \times \kappa$
- (4) $\kappa \twoheadrightarrow \sum_{n < \omega} \kappa^n$

Proof. For 1, see e.g. [Kun09, Theorem I.11.30].

For 3, suppose we are given a bijective pairing function $(-, -): \omega \times \omega \rightarrow \omega$. Any ordinal α can be written uniquely as $\alpha = \lambda + (m, n)$ where λ is a limit ordinal and $m, n \in \omega$. We then decode this as the pair $(m, \lambda + n)$, which clearly gives a bijection.

Deriving the other parts from these two is straightforward. \square

5. THE CONSTRUCTION OF IMAGE PRESERVING QW -TYPES

We now construct image preserving QW -types in **Set**. This is based on Jech's construction of the set of hereditarily countable sets [Jec82].

Definition 5.1. We will define a class function Q from ordinals to sets by recursion on ordinals.

We define $Q(0)$ to be \emptyset and for limit ordinals λ , we define $Q(\lambda)$ to be $\bigcup_{\alpha < \lambda} Q(\alpha)$.

We define $Q(\alpha + 1)$ as follows. Let X be the set of pairs (a, f) where $a \in A$ and $f : B_a \rightarrow Q(\alpha)$. We then take \sim to be the equivalence relation on X generated by identifying $(a_e, t \circ l_e)$ and $(b_e, t \circ r_e)$ whenever $t : V_e \rightarrow Q(\alpha)$ for $e \in E$ and we define $Q(\alpha + 1)$ to be X/\sim .

Remark 5.2. For $\beta \leq \alpha$ we have $Q(\beta) \subseteq Q(\alpha)$, by exploiting the fact that functions are implemented as relations not including explicit reference to their codomain, and noting that for $f : B_a \rightarrow Q(\alpha)$ and $g : B_b \rightarrow Q(\alpha)$ such that $(a, f) \sim (b, g)$, f factors through the inclusion $Q(\beta) \subseteq Q(\alpha)$ if and only if g does.

We now give a series of definitions and lemmas that apply at any stage $\alpha \in \text{On}$.

Definition 5.3. Note that we only identify (a, f) and (b, g) when f and g have the same image in $Q(\alpha)$. Hence we have a well defined image function $\text{im} : Q(\alpha) \rightarrow \mathcal{P}(Q(\alpha))$, such that whenever $x = [(a, f)]$, $\text{im}(x)$ is the image of f in $Q(\alpha)$.

Definition 5.4. Given an element x of $Q(\alpha)$ of the form $[(a, f)]$, we defined the *rank* of x , $\text{rank}(x)$ to be the smallest ordinal β such that f factors through the inclusion $Q(\beta) \subseteq Q(\alpha)$. To check this is a well defined, note that it depends only on the image of f .

Note that $\text{rank}(x) + 1$ is the smallest ordinal β such that $x \in Q(\beta)$.

Definition 5.5. Given a set $X \subseteq Q(\alpha)$, we define the union $\cup X$ by

$$\cup X := \bigcup_{x \in X} \text{im}(x)$$

We define the *transitive closure* of $x \in Q(\alpha)$, $\text{TC}(x)$ by

$$\text{TC}(x) := \bigcup_{1 \leq n < \omega} \cup^n \{x\}$$

Lemma 5.6. For all $x \in Q(\alpha)$, we have the following equation.

$$\text{rank}(x) = \{\text{rank}(y) \mid y \in \text{TC}(x)\}$$

Proof. It is clear that whenever $y \in \text{TC}(x)$ we must have $\text{rank}(y) < \text{rank}(x)$ since this is the case for any $n < \omega$ and any $y \in \cup^n \{x\}$ by induction on n .

It remains to show that for any $\beta < \text{rank}(x)$, we have $\beta = \text{rank}(y)$ for some $y \in \text{TC}(x)$. By the definition of rank , $\text{im}(x)$ cannot be contained in $Q(\beta)$. Hence we must have $x = [(a, f)]$ and $b \in B_a$ such that $fb \notin Q(\beta)$. For this b we have $\beta \leq \text{rank}(fb)$. If $\beta = \text{rank}(fb)$, then $fb \in \cup \{x\} \subseteq \text{TC}(x)$ and so β is as required. Otherwise, $\beta < \text{rank}(fb)$ and so by induction on $\text{rank}(x)$ we may assume $\beta = \text{rank}(y)$ for some $y \in \text{TC}(fb)$. However, $\text{TC}(fb) \subseteq \text{TC}(x)$, so $y \in \text{TC}(x)$ and β is again as required. \square

Definition 5.7. For $x \in Q(\alpha)$, we write $R_n(x)$ for the set $\{\text{rank}(z) \mid z \in \cup^n \{x\}\}$.

Lemma 5.8. For all $x \in Q(\alpha)$,

$$\text{rank}(x) = \bigcup_{1 \leq n < \omega} R_n(x)$$

Proof. By lemma 5.6. \square

Definition 5.9. We define κ to be $\aleph((B_a)_{a \in A})$. We define κ^+ to be the smallest non zero ordinal for which there is no surjection $\kappa \rightarrow \kappa^+$.

Lemma 5.10. *We define for each $x \in Q(\alpha)$ and each $1 \leq n < \omega$, a surjection $F_{x,n}: \kappa^n \rightarrow R_n(x) \cup \{\emptyset\}$.*

Proof. We first consider the case $n = 1$. Suppose that $x = [(a, f)]$. Note that $R_1(x) := \{\text{rank}(fb) \mid b \in B_a\}$ is a set of ordinals, and so it has an order type $\beta \in \text{On}$, and in particular we have a unique order isomorphism with β , say $\theta: \beta \xrightarrow{\cong} R_1(x)$. Furthermore, by definition, there is clearly a surjection from B_a to $R_1(x)$. It follows that $\beta < \kappa$. Hence we can define a canonical surjection $F_{x,1}: \kappa \rightarrow R_1(x) \cup \{\emptyset\}$ as follows.

$$F_{x,1}(\alpha) := \begin{cases} \theta(\alpha) & \alpha < \beta \\ \emptyset & \text{otherwise} \end{cases}$$

Now suppose $n = m + 1$. We fix m ordinals less than κ , say β_1, \dots, β_m and consider the set Y below.

$$Y := \{F_{fb,m}(\beta_1, \dots, \beta_m) \mid b \in B_a\}$$

This is again a set of ordinals with a surjection from B_a for some $a \in A$, and so as before, we have a canonical surjection $G: \kappa \rightarrow Y \cup \{\emptyset\}$. We take $F_{x,n}(\beta_1, \dots, \beta_m, \beta_{m+1})$ to be $G(\beta_{m+1})$. We now simultaneously check that $F_{x,n}$ has the correct codomain and is surjective.

$$\begin{aligned} \text{im}(F_{x,n}) &= \bigcup_{\beta_1, \dots, \beta_m < \kappa} (\{F_{fb,m}(\beta_1, \dots, \beta_m) \mid b \in B_a\} \cup \{\emptyset\}) \\ &= \bigcup_{b \in B_a} \{F_{fb,m}(\beta_1, \dots, \beta_m) \mid \beta_1, \dots, \beta_m < \kappa\} \cup \{\emptyset\} \\ &= \bigcup_{b \in B_a} \{\text{rank}(z) \mid z \in \cup^m \{fb\}\} \cup \{\emptyset\} \\ &= R_n(x) \cup \{\emptyset\} \end{aligned}$$

□

Lemma 5.11. *For any $x \in Q(\alpha)$ we have $\text{rank}(x) < \kappa^+$.*

Proof. First note that this is clear when $\text{rank}(x) = 0$. Hence we may assume for the rest of the proof $\text{rank}(x) > 0$.

By the definition of κ^+ , it suffices to define a surjection $\kappa \rightarrow \text{rank}(x)$. By proposition 4.4 it suffices to define a surjection $\sum_{1 \leq n < \omega} \kappa^n \rightarrow \text{rank}(x)$. However, by lemma 5.8 we can express $\text{rank}(x)$ as $\bigcup_{1 \leq n < \omega} R_n(x)$. Since $\text{rank}(x) > 0$, this is the same as $\bigcup_{1 \leq n < \omega} (R_n(x) \cup \{\emptyset\})$, and so we can just combine the surjections defined in lemma 5.10. □

Theorem 5.12. *All image preserving QW-types exist in Set.*

Proof. We show that $Q(\kappa^+)$ is an initial algebra. We first need to show how to define an algebra structure. Suppose we are given $a \in A$ and a map $f: B_a \rightarrow Q(\kappa^+)$. Then $[(a, f)]$ is an element of $Q(\kappa^+ + 1)$. By lemma 5.11 we have $\text{rank}([(a, f)]) < \kappa^+$, and so f factors through $Q(\beta)$ for some $\beta < \kappa^+$. We can then take $\text{sup}(a, f)$ to be $[(a, f)] \in Q(\beta + 1)$.

We check that this structure respects the equations. Suppose that we are given $g: V_e \rightarrow Q(\kappa^+)$. Note that $g \circ l_e$ and $g \circ r_e$ have the same image, and so $\text{rank}([(a_e, g \circ l_e)]) = \text{rank}([(b_e, g \circ r_e)])$. Hence $[(a_e, g \circ l_e)]$ and $[(b_e, g \circ r_e)]$ must have been first

added at the same stage, $\alpha + 1$. We can now see that they were identified in the definition of $Q(\alpha + 1)$.

Finally, it is clear that for any other algebra structure, we can define a unique homomorphism out of $Q(\kappa^+)$ by recursion on ordinals. \square

6. THE RANKS OF UNORDERED COUNTABLY BRANCHING TREES

It follows from Jech's construction that every hereditarily countable set has rank less than ω_2 . In [Jec82] Jech also showed a converse statement when ω_1 is singular: in this case there is a hereditarily countable set of rank α for every $\alpha < \omega_2$. We will show the analogous result for unordered countably branching trees. We first recall that Jech proved the following lemma as part of the proof of [Jec82, Theorem 2].

Lemma 6.1. *Suppose that ω_1 is singular and let $\alpha < \omega_2$. For each fixed surjection $f : \omega_1 \rightarrow \alpha$ and choice of countable cofinal sequence in ω_1 , we can construct for each limit ordinal $\lambda \leq \alpha$ a choice of countable cofinal subset $Y_\lambda \subseteq \lambda$.*

We can now show the theorem.

Theorem 6.2. *Suppose that ω_1 is singular. Then for every $\alpha < \omega_2$ there is an unordered countably branching tree of rank α .*

Proof. Since $\alpha < \omega_2$ and ω_1 is singular there exists a choice of sets Y_λ as in lemma 6.1 for limit ordinals $\lambda \leq \alpha$.

We construct an unordered countably branching tree t_β for each $\beta \leq \alpha$ by induction. We define $t_0 := \mathbf{leaf}$. For successor ordinals we define $t_{\beta+1} := \mathbf{node}(\lambda n.t_\beta)$. This just leaves the case of t_λ for non zero limit ordinals λ . Since Y_λ is countable there exists a surjection $g : \omega \rightarrow Y_\lambda$. Since Y_λ is cofinal in a limit ordinal it is not finite. Hence there exists a bijection $h : \omega \xrightarrow{\cong} Y_\lambda$, which we can construct from a choice of surjection g as follows:

$$h(0) = g(0)$$

$$h(n+1) = \begin{cases} g(n+1) & g(n+1) \neq h(k) \text{ for all } k \leq n \\ \gamma & \text{otherwise, } \gamma \text{ least s.t. } \gamma \neq h(k) \text{ for all } k \leq n \end{cases}$$

We show how to define t_λ from a choice of bijection h , and then check that it is independent of the particular choice and so well defined.

We define $t_\lambda := \mathbf{node}(\lambda n.t_{h(n)})$. Now suppose that we are given two such bijections $h, h' : \omega \xrightarrow{\cong} Y_\lambda$. Then $\pi := h^{-1} \circ h'$ is a permutation of ω and $h' = h \circ \pi$. Hence $\mathbf{node}(\lambda n.t_{h'(n)})$ and $\mathbf{node}(\lambda n.t_{h(n)})$ are identified by one of the defining equations of unordered countably branching trees, ensuring that t_λ is independent of the choice of h , as required. It is clear that $\text{rank}(t_\beta) = \beta$ for all $\beta \leq \alpha$. \square

Corollary 6.3. *If ω_1 is singular, then the unique homomorphism $h : T \rightarrow T_{\text{Sym}}$ from countably branching trees to unordered countably branching trees is not surjective.*

Proof. Note that h preserves rank. Hence by theorem 6.2 it suffices to show that the rank of every (ordered) countably branching tree is less than ω_1 .

Fix a surjective function $s : \omega \rightarrow \omega + (\omega \times \omega)$. We construct for each ordered tree t a countable enumeration $f(t) : \omega \rightarrow T$ whose image is the union of the set of proper subtrees of t and $\{\mathbf{leaf}\}$. We define $f(\mathbf{leaf})(n) := \mathbf{leaf}$. Now suppose

we are given a tree of the form $t = \text{node}(g)$. By composing with s , it suffices for us to define two functions $f_0 : \omega \rightarrow T$ and $f_1 : \omega \times \omega \rightarrow T$ that jointly enumerate all of the proper subtrees of t (including leaf). We define $f_0(n) := g(n)$ and $f_1(n, m) := f(g(n))(m)$.

By lemma 5.6 the rank of each t is equal to the set of ranks of proper subtrees. However, by the above enumeration this is a countable ordinal and so less than ω_1 , as required. \square

7. CONCLUSION

We have shown that every image preserving QW -type exists in \mathbf{Set} under \mathbf{ZF} without any additional assumptions. Although the proof is somewhat elaborate, we can see from the results of section 6 that some complication is necessary. The naïve construction of defining an equivalence relation on the W -type of the underlying polynomial and quotienting will not work in general. As we saw, both hereditarily countable sets and unordered countably branching trees provide counterexamples when ω_1 is singular.

7.1. Generalisations and limitations of this method. The proof in section 5 made use of the fact that we have a well defined image operation sending each element of the form (a, f) to the image of f . This may not be strictly necessary for the proof, since the key lemma 5.10 only used the sets R_n in definition 5.7 which may be definable in situations without a well defined image operator. So there could be a more general version of the theorem that uses the same key idea. However, it is unclear if there are any interesting examples of QW -type that can be constructed by the same general method, but whose existence isn't already a corollary of theorem 5.12. We therefore leave it as an open problem both to generalise the theorem and to find interesting examples making essential use of the generalisation.

On the other hand there are some examples where the argument here cannot possibly apply. As already discussed in example 2.8, Blass' counterexample cannot be constructed by this method, since it does not provably exist under \mathbf{ZF} . However, there is also a much simpler example of a QW -type that exists, provably in \mathbf{ZF} , but does not seem to be covered by the argument in this paper, namely free monoids. We consider in particular the free monoid on a set with four elements $\{a, b, c, d\}$. As a special case of associativity we have $a(b(cd)) = (ab)(cd)$. However, it is unclear how to define R_1 for this term. Since $\text{rank}(a) = 0$ and $\text{rank}(b(cd)) = 2$, we expect $R_1(a(b(cd))) = \{0, 2\}$. Since $\text{rank}(ab) = \text{rank}(cd) = 1$, we also expect $R_a((ab)(cd)) = \{1, 1\}$. Since R_n play an essential role in lemma 5.10 it seems that it is impossible to construct free monoids by this method, even though they do provably exist in \mathbf{Set} under \mathbf{ZF} .

Another class of examples of QW -types that exist under \mathbf{ZF} , but are not covered directly by the methods of section 5 are W -types with reductions. By their nature they identify two trees of different rank, so we don't expect to have a well defined of rank without the observation in [Swa18] that in the presence of the law of excluded middle they can be viewed as an ordinary W -types of normal forms.

7.2. Related Open Problems. The limitations of the result discussed above naturally lead to the following question.

Open Problem 7.1. *Is there a general construction of QW -types in \mathbf{Set} under \mathbf{ZF} that naturally encompasses all examples known to exist in this setting?*

Our proof of lemma 5.10 makes essential use of the fact that any set of ordinals is order isomorphic to an ordinal, which in turn uses classical logic. This leaves open the case of intuitionistic logic, specifically the following problem.

Open Problem 7.2. *Show all image preserving QW-types exist in Set under the assumptions of **IZF** or find a model of **IZF** where they do not.*

Note that the proof of theorem 6.2 made essential use of the fact that we have equations for all permutations π of ω . It is unclear what happens when we allow some permutations but not all, leading to the following question.

Open Problem 7.3. *We define the QW-type of countably branching weakly unordered trees to have the same underlying polynomial as countably branching trees and equations of the form $\mathbf{node}(\lambda n.v_n) = \mathbf{node}(\lambda n.v_{\pi(n)})$ where π is a finitely supported permutation of ω . Is it provable in **ZF** that the unique homomorphism from countably branching trees to weakly unordered countably branching trees is a surjection?*

REFERENCES

- [AB04] Steven Awodey and Andrej Bauer. Propositions as [types]. *Journal of Logic and Computation*, 14(4):447–471, 2004.
- [ACD⁺18] Thorsten Altenkirch, Paolo Capriotti, Gabe Dijkstra, Nicolai Kraus, and Fredrik Nordvall Forsberg. Quotient inductive-inductive types. In Christel Baier and Ugo Dal Lago, editors, *Foundations of Software Science and Computation Structures*, pages 293–310, Cham, 2018. Springer International Publishing.
- [AK16] Thorsten Altenkirch and Ambrus Kaposi. Type theory in type theory using quotient inductive types. In *Proceedings of the 43rd Annual ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages*, POPL '16, page 18–29, New York, NY, USA, 2016. Association for Computing Machinery.
- [AR01] Peter Aczel and Michael Rathjen. Notes on constructive set theory. Technical Report 40, Institut Mittag-Leffler, 2001.
- [Bla83] Andreas Blass. Words, free algebras, and coequalizers. *Fundamenta Mathematicae*, 117(2):117–160, 1983.
- [CH19] Evan Cavallo and Robert Harper. Higher inductive types in cubical computational type theory. *Proc. ACM Program. Lang.*, 3(POPL), January 2019.
- [CHM18] Thierry Coquand, Simon Huber, and Anders Mörtberg. On higher inductive types in cubical type theory. In *Proceedings of the 33rd Annual ACM/IEEE Symposium on Logic in Computer Science*, LICS '18, pages 255–264, New York, NY, USA, 2018. ACM.
- [FPS20] Marcelo P. Fiore, Andrew M. Pitts, and S. C. Steenkamp. Constructing infinitary quotient-inductive types. In J. Goubault-Larrecq and B. König, editors, *23rd International Conference on Foundations of Software Science and Computation Structures (FoSSaCS 2020)*, volume 12077 of *Lecture Notes in Computer Science*, pages 257–276. Springer, 2020.
- [FPS21] Marcelo P. Fiore, Andrew M. Pitts, and S. C. Steenkamp. Quotients, inductive types, and quotient inductive types. arXiv:2101.02994, January 2021.
- [Git80] M. Gitik. All uncountable cardinals can be singular. *Israel Journal of Mathematics*, 35(1):61–88, Sep 1980.
- [Git85] Moti Gitik. Regular cardinals in models of ZF. *Transactions of the American Mathematical Society*, 290(1):41–68, 1985.
- [Hol14] M. Randall Holmes. On hereditarily small sets in ZF. *Mathematical Logic Quarterly*, 60(3):228–229, 2014.
- [Jec82] Thomas Jech. On hereditarily countable sets. *The Journal of Symbolic Logic*, 47(1):43–47, 1982.
- [Kel80] G.M. Kelly. A unified treatment of transfinite constructions for free algebras, free monoids, colimits, associated sheaves, and so on. *Bulletin of the Australian Mathematical Society*, 22(1):1–83, 1980.

- [Kun09] Kenneth Kunen. *The Foundations of Mathematics*. Studies in Logic: Mathematical Logic and Foundations. College publications, 2009.
- [LS20] Peter LeFanu Lumsdaine and Michael Shulman. Semantics of higher inductive types. *Mathematical Proceedings of the Cambridge Philosophical Society*, 169(1):159–208, 2020.
- [Mai05] Maria Emilia Maietti. Modular correspondence between dependent type theories and categories including pretopoi and topoi. *Mathematical Structures in Computer Science*, 15:1089–1149, 12 2005.
- [MP00] Ieke Moerdijk and Erik Palmgren. Wellfounded trees in categories. *Annals of Pure and Applied Logic*, 104(1):189 – 218, 2000.
- [Soj15] Kristina Sojakova. Higher inductive types as homotopy-initial algebras. In *Proceedings of the 42Nd Annual ACM SIGPLAN-SIGACT Symposium on Principles of Programming Languages*, POPL '15, pages 31–42, New York, NY, USA, 2015. ACM.
- [Swa18] Andrew Wakelin Swan. W-types with reductions and the small object argument. arXiv:1802.07588, February 2018.
- [Uni13] Univalent Foundations Program. *Homotopy Type Theory: Univalent Foundations of Mathematics*. <http://homotopytypetheory.org/book>, Institute for Advanced Study, 2013.
- [vdB12] Benno van den Berg. WISC in indepedent from ZF. Unpublished note available at <https://staff.fnwi.uva.nl/b.vandenberg3/papers/WISC.pdf>, May 2012.