

Lattice-QCD Calculations of TMD Soft Function Through Large-Momentum Effective Theory (Lattice Parton Collaboration (LPC))

Qi-An Zhang,¹ Jun Hua,² Yikai Huo,^{2,3} Xiangdong Ji,^{1,4} Yizhuang Liu,¹ Yu-Sheng Liu,¹ Maximilian Schlemmer,⁵ Andreas Schäfer,⁵ Peng Sun,⁶ Wei Wang,^{2,*} and Yi-Bo Yang^{7,†}

¹*Tsung-Dao Lee Institute, Shanghai Jiao Tong University, Shanghai 200240, China*

²*INPAC, SKLPPC, MOE KLPPC, School of Physics and Astronomy, Shanghai Jiao Tong University, Shanghai 200240, China*

³*Zhiyuan College, Shanghai Jiao Tong University, Shanghai 200240, China*

⁴*Department of Physics, University of Maryland, College Park, MD 20742, USA*

⁵*Institut für Theoretische Physik, Universität Regensburg, D-93040 Regensburg, Germany*

⁶*Nanjing Normal University, Nanjing, Jiangsu, 210023, China*

⁷*CAS Key Laboratory of Theoretical Physics, Institute of Theoretical Physics, Chinese Academy of Sciences, Beijing 100190, China*

(Dated: June 1, 2020)

The transverse-momentum-dependent (TMD) soft function is a key ingredient in QCD factorization of Drell-Yan and other processes with relatively small transverse momentum. We present a lattice QCD study of this function at moderately large rapidity on a 2+1 flavor CLS dynamic ensemble with $a = 0.098$ fm. We extract the rapidity-independent (or intrinsic) part of the soft function through a large-momentum-transfer pseudo-scalar meson form factor and its quasi-TMD wave function using leading-order factorization in large-momentum effective theory. We also investigate the rapidity-dependent part of the soft function—the Collins-Soper evolution kernel—based on the large-momentum evolution of the quasi-TMD wave function.

Introduction. For high-energy processes such as Higgs production at the Large-Hadron Collider, quantum chromodynamics (QCD) factorization and parton distribution functions (PDFs) have been essential for making theoretical predictions [1, 2]. But for processes involving observation of a relatively small transverse momentum, $Q_\perp$ such as in Drell-Yan (DY) production and semi-inclusive deep inelastic scattering, a new non-perturbative quantity called *soft function* is required to capture the physics of non-cancelling soft gluon-radiation at fixed $Q_\perp$ [3–6]. Physically, the soft function in DY is a cross section for a pair of high-energy quark and anti-quark (or gluon) traveling in the opposite light-cone directions to radiate soft gluons of total transverse momentum $Q_\perp$ before they annihilate. Although much progress has been made in calculating the soft function in perturbation theory at $Q_\perp \gg \Lambda_{\text{QCD}}$ [7, 8], it is intrinsically non-perturbative when $Q_\perp$ is $\mathcal{O}(\Lambda_{\text{QCD}})$. Calculating the non-perturbative transverse-momentum-dependent (TMD) soft function from first principles became feasible only recently [9].

The main difficulty in calculating the TMD soft function in lattice QCD is that it involves two light-like Wilson lines along directions $n^\pm = \frac{1}{\sqrt{2}}(1, \vec{0}_\perp, \pm 1)$ in $(t, \perp, z)$ coordinates, making direct simulations in Euclidean space impractical. However, much progress has been made in recent years in calculating physical quantities such as light-cone PDFs using the framework of large-momentum effective theory (LaMET) [10, 11]. The

key observation of LaMET is that the collinear quark and gluon modes, usually represented by light-like field correlators [12–15], can be accessed for large-momentum hadron states. A detailed review of LaMET and its applications to collinear PDFs and other light-cone distributions can be found in Refs.[16, 17]. More recently, some of the present authors have proposed that the TMD soft function can be extracted from a special large-momentum-transfer form factor of either a light meson or a pair of quark-antiquark color sources [9]. Once calculated, the TMD factorization of the Drell-Yan and similar processes can be made with entirely lattice-QCD-computable non-perturbative quantities [18–23].

The TMD soft function is often defined and applied not in momentum space but in transverse coordinate space in terms of the Fourier transformation variable $b_\perp$. In addition, it also depends on the ultraviolet (UV) renormalization scale μ (often defined in dimensional regularization and minimal subtraction or $\overline{\text{MS}}$) and rapidity regulators $Y + Y'$ [9, 12],

$$S(b_\perp, \mu, Y + Y') = e^{(Y+Y')K(b_\perp, \mu)} S_I^{-1}(b_\perp, \mu) \quad (1)$$

where the first factor is related to rapidity evolution [described by the Collin-Soper (CS) kernel K], and the second factor S_I is the intrinsic, rapidity independent, part of the soft contribution. The rapidity-regulator-independent CS-kernel K is found calculable by taking ratio of the quasi-TMDPDF at two different momenta [20–25]. On the other hand, calculating the intrinsic soft function on the lattice has never been attempted before.

In this paper we present the first lattice QCD calculation of the intrinsic soft function S_I with several

* Corresponding author: wei.wang@sjtu.edu.cn

† Corresponding author: ybyang@itp.ac.cn

momenta on a 2+1 flavor CLS ensemble with $a = 0.098$ fm [26], see Table I. In particular we perform simulations of the large-momentum light-meson form factor and quasi-TMD wave functions (TMDWFs), whose ratio gives the intrinsic soft function [9]. The Wilson loop matrix element will be used to remove the linear divergence in the quasi-TMD wave function. The CS kernel, K , can also be calculated from the external momentum dependence of the quasi-TMD wave function [16], and we will calculate it as a by-product. Our result is consistent with that of a quenched lattice study using TMDPDFs [25].




FIG. 1. Illustration of the pseudo-scalar meson form factor F calculated in this work. The initial and final momenta of the pion are large and opposite. The transition “current” is made of two local operators at a fixed spatial separation $b_\perp$. t_{sep} is the time separation between the source and sink of the pion.

Theoretical Framework. The intrinsic soft function (S_I) can be obtained from the QCD factorization of a large-momentum form factor of a non-singlet light pseudo-scalar meson with constituents $\pi = \bar{q}_2 \gamma_5 q_1$, with the transition current made of two quark-bilinears with a fixed transverse separation $\vec{b} = (\vec{n}_\perp b_\perp, 0)$,

$$F(b_\perp, P^z) = \langle \pi(-\vec{P}) | (\bar{q}_1 \Gamma q_1)(\vec{b}) (\bar{q}_2 \Gamma q_2)(0) | \pi(\vec{P}) \rangle_c. \quad (2)$$

Here $q_{1,2}$ are light quark fields of different flavors, and $\vec{P} = (\vec{0}_\perp, P^z)$. The initial and final mesons approach two opposite lightcone directions in the $P^z \rightarrow \infty$ limit. Only the connected diagram is important in the large momentum limit, as illustrated in Fig. 1.

It can be shown that the form factor defined in Eq. (2) is factorizable into the quasi-TMDWF Φ and the intrinsic soft function S_I [9, 16]

$$F(b_\perp, P^z) = S_I(b_\perp) \times \int_0^1 dx dx' H(x, x', P^z) \Phi^\dagger(x', b_\perp, -P^z) \Phi(x, b_\perp, P^z) \quad (3)$$

where H is perturbative hard kernel. The quasi-TMDWF Φ is the Fourier transformation of the coordinate-space

correlation function

$$\begin{aligned} \phi(z, b_\perp, P^z) &= \lim_{\ell \rightarrow \infty} \frac{\phi_\ell(z, b_\perp, P^z, \ell)}{\sqrt{Z_E(2\ell, b_\perp)}}, \\ \phi_\ell(z, b_\perp, P^z, \ell) &= \langle 0 | \bar{q}_1 \left(\frac{z}{2} n^z + \vec{b} \right) \Gamma_\Phi \mathcal{W}(\vec{b}, \ell) q_2 \left(-\frac{z}{2} n^z \right) | \pi(\vec{P}) \rangle. \end{aligned} \quad (4)$$

In the above $\mathcal{W}(\vec{b}, \ell)$ is the spacelike staple-shaped gauge link,

$$\begin{aligned} \mathcal{W}(\vec{b}, \ell) &= \mathcal{P} \exp \left[ig_s \int_{-\ell}^{z/2} ds n^z \cdot A(n^z s + b_\perp) \right] \\ &\times \mathcal{P} \exp \left[ig_s \int_0^{b_\perp} ds n_\perp \cdot A(-\ell n^z + s n_\perp) \right] \\ &\times \mathcal{P} \exp \left[ig_s \int_{-z/2}^{-\ell} ds n^z \cdot A(n^z s) \right], \end{aligned} \quad (5)$$

n^z and $n_\perp$ are the unit vectors in z and transverse directions respectively. $Z_E(2\ell, b_\perp)$ is the vacuum expectation value of a rectangular spacelike Wilson loop with size $2\ell \times b_\perp$ which removes the pinch-pole singularity and Wilson-line self-energy in quasi-TMDWF [9].

Since the UV divergence of the intrinsic soft function is multiplicative [16], the ratio $S_I(b_\perp, 1/a)/S_I(b_{\perp,0}, 1/a)$ calculable on lattice is UV renormalization-scheme independent, where $b_{\perp,0}$ is a reference distance which is taken small enough to be calculated perturbatively. Thus we can obtain the result in the $\overline{\text{MS}}$ scheme through

$$S_{I,\overline{\text{MS}}}(b_\perp, \mu) = \left(\frac{S_I(b_\perp, 1/a)}{S_I(b_{\perp,0}, 1/a)} \right) S_{I,\overline{\text{MS}}}(b_{\perp,0}, \mu) \quad (6)$$

where $S_{I,\overline{\text{MS}}}(b_{\perp,0}, \mu)$ is perturbatively calculable, e.g.,

$$S_{I,\overline{\text{MS}}}(b_\perp, \mu) = 1 - \frac{\alpha_s C_F}{\pi} \ln \frac{\mu^2 b_\perp^2}{4e^{-2\gamma_E}} + \mathcal{O}(\alpha_s). \quad (7)$$

In the present exploratory study, we will consider only the leading order matching in Eq. (3), for which the perturbative kernel is $H(x, x', P^z) = 1/(2N_c) + \mathcal{O}(\alpha_s)$, independent of x and x' . Using $\phi(0, b_\perp, -P^z) = \phi(0, b_\perp, P^z)$ under parity transformation, we obtain

$$S_I(b_\perp) = \frac{2N_c F(b_\perp, P^z)}{|\phi(0, b_\perp, P^z)|^2} + \mathcal{O}(\alpha_s, (1/P^z)^2), \quad (8)$$

where power corrections from finite P^z are ignored. Since P^z is related to the rapidity of the meson, we henceforth replace it by the boost factor $\gamma \equiv E_\pi/m_\pi$. Eq. (6) can be written as

$$S_{I,\overline{\text{MS}}}(b_\perp, \mu) = \frac{F(b_\perp, P^z)}{F(b_{\perp,0}, P^z)} \frac{|\phi(0, b_{\perp,0}, P^z)|^2}{|\phi(0, b_\perp, P^z)|^2} + \mathcal{O}(\alpha_s, \gamma^{-2}). \quad (9)$$

The ratio on the right-hand side of the above expression is independent of the renormalization scale μ since only the leading-order contribution is kept.

On the other hand, the quasi-TMDWF can be used to extract the Collins-Soper kernel K using a method similar to [20]

$$K(b_\perp, \mu) = \frac{1}{\ln(P_1^z/P_2^z)} \ln \left| \frac{C(xP_2^z, \mu) \Phi_{\overline{\text{MS}}}(x, b_\perp, P_1^z, \mu)}{C(xP_1^z, \mu) \Phi_{\overline{\text{MS}}}(x, b_\perp, P_2^z, \mu)} \right| \quad (10)$$

$$= \frac{1}{\ln(P_1^z/P_2^z)} \ln \left| \frac{\int_0^1 dx \Phi(x, b_\perp, P_1^z)}{\int_0^1 dx \Phi(x, b_\perp, P_2^z)} \right| + \mathcal{O}(\alpha_s, \gamma^{-2})$$

$$= \frac{1}{\ln(P_1^z/P_2^z)} \ln \left| \frac{\phi(0, b_\perp, P_1^z)}{\phi(0, b_\perp, P_2^z)} \right| + \mathcal{O}(\alpha_s, \gamma^{-2}). \quad (11)$$

In the second line, again only use the leading order matching kernel $C(xP^z, \mu) = 1 + \mathcal{O}(\alpha_s)$ is used. The renormalization factors for Φ are cancelled. The rapidity-scheme-independent CS kernel K is independent of μ in this approximation because only the leading term has been kept.

While Eqs. (6) and (10) are exact and can be used for precision studied in the future, Eqs. (9) and (11) are the leading-order approximation used in this pioneering work.

TABLE I. Parameters used in the numerical simulation. The first row shows the parameters of the 2+1 flavor clover fermion CLS ensemble (named A654) and the second one shows the number of the A654 configurations and valence pion mass used for this calculation.

β	$L^3 \times T$	a (fm)	c_{sw}	κ_l^{sea}	m_π^{sea} (MeV)
3.34	$24^3 \times 48$	0.098	2.06686	0.13675	333
			N_{cfg}	κ_l^v	m_π^v (MeV)
			868	0.13622	547

Simulation setup. For the present study, we use configurations generated with 2+1 flavor clover fermions and tree-level Symanzik gauge action configuration by the CLS collaboration using periodic boundary conditions [26]. The detailed parameters are listed in Table I. Note that $m_\pi = 547$ MeV instead of 333 MeV is used for valence quarks in order to have a better signal. Physically, the soft function becomes independent of the meson mass for large boost factors γ .

To calculate the form factor in Eq.(2), we generate the wall source propagator,

$$S_w(x, t, t'; \vec{p}) = \sum_{\vec{y}} S(t, \vec{x}; t', \vec{y}) e^{i\vec{p} \cdot (\vec{y} - \vec{x})}, \quad (12)$$

on the Coulomb gauge fixed configurations at $t' = 0$ and t_{sep} for both the initial and final meson states. S is the quark propagator from $(t', \vec{y})$ to $(t, \vec{x})$. Then we can construct the three point function (3pt) corresponding to the

form factor in Eq. (2),

$$C_3(b_\perp, P^z; p^z, t_{\text{sep}}, t) \quad (13)$$

$$= \frac{1}{L^3} \sum_x \text{Tr} \langle S_w^\dagger(\vec{x} + \vec{b}, t, 0; -\vec{p}) \gamma_5 \Gamma S_w(\vec{x} + \vec{b}, t, t_{\text{sep}}; \vec{p}) \times S_w^\dagger(\vec{x}, t, t_{\text{sep}}; -\vec{P} + \vec{p}) \gamma_5 \Gamma S_w(\vec{x}, t, 0; \vec{P} - \vec{p}) \rangle.$$

The quark momentum $\vec{p} = (\vec{0}_\perp, p^z)$, and the relation $\gamma_5 S^\dagger(x, y) \gamma_5 = S(y, x)$ have been applied for the anti-quark propagator. We have tested several choices of Γ , and will use the unity Dirac matrix $\Gamma = I$ as it has the best signal and describes the leading twist light-cone contribution in the large P^z limit. Notice that the $\Gamma = \gamma_4$ case is subleading in large P^z limit, although the excited state contamination might be smaller.

By generating the wall source propagators at all the 48 time slices with quark momentum $p^z = (-2, -1, 0, 1, 2) \times 2\pi/(La)$, we can maximize the statistics of the 3pt function with all the meson momenta P_z from 0 to $8\pi/(La)$ (~ 2.1 GeV) with arbitrary t and t_{sep} . $C_3(b_\perp, P^z, t_{\text{sep}}, t)$ is related to the bare $F(b_\perp, P^z)$ using standard parameterization of 3pt with one excited state,

$$C_3(b_\perp, P^z; p^z, t_{\text{sep}}, t) = \frac{A_w(p_z)^2}{(2E)^2} e^{-E t_{\text{sep}}} [F(b_\perp, P^z) + c_1(e^{-\Delta E t} + e^{-\Delta E(t_{\text{sep}} - t)}) + c_2 e^{-\Delta E t_{\text{sep}}}] \quad (14)$$

A_w is the matrix element of the Coulomb gauge fixed wall (CFW) source pion interpolation field, $E = \sqrt{m_\pi^2 + P^z^2}$ is the pion energy, ΔE is the mass gap between pion and its first excited state, $c_{1,2}$ are parameters for the excited state contamination. Note that the p_z dependence factor A_w^2 will cancel

The same wall source propagators can be used to calculate the two-point function related to the bare quasi-TMDWF,

$$C_2(b_\perp, P^z; p_z, \ell, t) = \frac{1}{L^3 \sqrt{Z_E(2\ell, b_\perp)}} \sum_x \text{Tr} e^{i\vec{P} \cdot \vec{x}}$$

$$\times \langle S_w^\dagger(\vec{x} + \vec{b}, t, 0; -\vec{p}) \mathcal{W}(\vec{b}, \ell) \gamma_5 \Gamma_\Phi S_w(\vec{x}, t, 0; P^z - \vec{p}) \rangle$$

$$= \frac{A_w(p_z) A_p}{2E} e^{-E t} \phi_\ell(0, b_\perp, P^z, \ell) (1 + c_0 e^{-\Delta E t}), \quad (15)$$

where again we parameterize the mixing with one excited state. A_p is the matrix element of the point sink pion interpolation field. It will be removed when we normalize $\phi_\ell(0, b_\perp, P^z, \ell)$ with $\phi_\ell(0, 0, P^z, 0)$. We choose $\Gamma_\Phi = \gamma^t \gamma_5$ to define the wave function amplitude in Eq. (4). Based on the quasi-TMDPDF study in Ref. [25] with the similar staple-shaped gauge link operator, the mixing effect from most of the Dirac matrices are at 1% level except in the cases with certain Dirac matrix which has very large statistical uncertainties. Thus the mixing effect is ignored in this work.

The dispersion relation of the pion state, statistical checks for the measurement histogram, and information on the autocorrection between configurations can be found in the supplemental materials [27].




FIG. 2. Results for the ℓ dependence of the quasi-TMDWF with $z = 0$, and also the square root of the Wilson loop which is used for the subtraction, taking the $\{P^z, b_\perp, t\} = \{6\pi/L, 3a, 6a\}$ case as an example. All the results are normalized with their values at $\ell = 0$.

Numerical Results. We demonstrate the Wilson-line length ℓ dependence of the norm of the quasi-TMDWFs in Fig. 2. As one can see from this figure, with $\{P^z, b_\perp, t\} = \{6\pi/L, 3a, 6a\}$, both the quasi-TMDWF $\phi_\ell(0, b_\perp, P^z, \ell)$ and the square root of the Wilson loop Z_E decay exponentially with the length ℓ , but the subtracted quasi-TMDWF is length independent when $\ell \geq 0.4$ fm. Some other cases with larger P^z , $b_\perp$, and t can be found in the supplemental materials [27]. Based on this observation, we will use $\ell = 7a = 0.686$ fm as asymptotic results for all cases in the following calculation.




FIG. 3. The ratios $C_3(b_\perp, P^z, t_{\text{sep}}, t)/C_2(0, P^z, 0, t_{\text{sep}})$ (data points) which converge to the ground state contribution at $t, t_{\text{sep}} \rightarrow \infty$ (gray band) as function of t_{sep} and t , with $\{P^z, b_\perp\} = \{6\pi/L, 3a\}$. As in this figure, our data in general agree with the predicted fit function (colored bands).

We applied the joint fit of the form factor and quasi-TMDWF with the same P^z and $b_\perp$ with the parameterization in Eqs. (14) and (15), and the ratios $C_3(b_\perp, P^z, t_{\text{sep}}, t)/C_2(0, P^z, 0, t_{\text{sep}})$ with different t_{sep} and t for the $\{P^z, b_\perp\} = \{6\pi/L, 3a\}$ case are shown in Fig. 3, with ground state contribution (gray band) and the fitted results at finite t_2 and t (colored bands). In the

calculation, the excited state contribution is properly described by the fit with $\chi^2/\text{d.o.f.} = 0.6$. The details of the joint fit, and also more fit quality checks are shown in the supplemental materials [27], with similar fitting qualities.




FIG. 4. The intrinsic soft factor as a function of $b_\perp$ with $b_{\perp,0} = a$ as in Eq. (9). With different pion momentum P^z , the results are consistent with each other. The dashed curve shows the result of the 1-loop calculation, see Eq. (7), with the strong coupling $\alpha_s(1/b_\perp)$.

The resulting soft factor as function of $b_\perp$ is plotted in Fig. 4, at $\gamma = 2.17, 3.06$ and 3.98 , which corresponds to $P^z = \{4, 6, 8\}\pi/L = \{1.05, 1.58, 2.11\}$ GeV respectively. As in Fig. 4, the results at different large γ are consistent with each other, demonstrating that the asymptotic limit is stable within errors. We also compare the intrinsic soft function extracted from the lattice to the one-loop result in Eq. (7), with $\alpha_s(\mu = 1/b_\perp)$ evolving from $\alpha_s(\mu = 2 \text{ GeV}) \approx 0.3$. Notice that the $b_\perp$ dependence of the former comes purely from the lattice simulation, while that for the latter is from perturbation theory.

We can see a clear P^z dependence in the quasi-TMDWF $|\phi_\ell(0, b_\perp, P^z, \ell)|$ normalized with $\phi_\ell(0, 0, P^z, 0)$, as in the upper panel of Fig. 5. It is understandable since such a dependence is related to the CS kernel as shown in Eq. (11), up to the possible LaMET matching effects and power corrections of order $1/\gamma^2$. Thus we use Eq. (11) to extract the kernel in the tree level approximation, and compare the result in the lower panel of Fig. 5 with the calculation in Ref. [25] and up to 3-loop perturbative one with $\alpha_s(\mu = 1/b_\perp)$. We estimate the systematic uncertainty by combining in quadrature the statistical errors with contributions from the imaginary part of the quasi-TMDWF, which should be identically zero. For details see the supplemental materials [27], in particular Fig. 11. Our result appears consistent with a quenched calculation based on TMDPDFs from Ref. [25], but the results based on $P_1^z/P_2^z = 3/2$ and $P_1^z/P_2^z = 4/2$ have some differences at certain $b_\perp$ which might come from insufficient statistics at $P^z = 2.11$ GeV, or potential systematic uncertainties.

Summary and Outlook. In this work, we have presented an exploratory lattice calculation of the intrinsic




FIG. 5. Quasi-TMDWF (upper panel) and extracted Collins-Soper kernel (lower panel), as functions of $b_\perp$. The visible P^z dependence of the quasi-TMDWF can be primarily understood by that from the Collins-Soper kernel, as the kernel we obtained with tree level matching is consistent with up to 3-loop perturbative calculations (at small $b_\perp$) with the strong coupling α_s at the scale $1/b_\perp$, and also the non-perturbative result from the pion quasi-TMDPDF. Results from quenched lattice calculations [25] are also shown for comparison.

soft function by simulating the light-meson form factor of four-quark non-local operators and quasi-TMD wave

functions. Our result shows a mild hadron momentum dependence, which allows a future precision study to eliminate the large momentum dependence using perturbative matching [16]. As a reliability check, the agreement between the CS kernel obtained from our quasi-TMDWF result and the previous calculations shows that the systematic uncertainties including the partially quenching effect, the leading perturbative matching and missing power corrections $1/\gamma$ in LaMET expansion might be sub-leading. Still our calculation paves the way towards the first principle predictions of physical cross sections for, e.g., Drell-Yan and Higgs productions at small transverse momentum.

Acknowledgment.— We thank Xu Feng, Jianhui Zhang and Yong Zhao for valuable discussions. We thank the CLS Collaboration for sharing the lattice ensembles used to perform this study. The LQCD calculations were performed using the Chroma software suite [28]. The numerical calculation is supported by Chinese Academy of Science CAS Strategic Priority Research Program of Chinese Academy of Sciences, Grant No. XDC01040100, Center for HPC of Shanghai Jiao Tong University, HPC Cluster of ITP-CAS, and Jiangsu Key Lab for NSLSCS. J. Hua is supported by NSFC under grant No. 11735010 and 11947215. Y.-S. Liu is supported by National Natural Science Foundation of China under grant No.11905126. M. Schlemer and A. Schäfer were supported by the cooperative research center CRC/TRR-55 of DFG. P. Sun is supported by Natural Science Foundation of China under grant No. 11975127 as well as Jiangsu Specially Appointed Professor Program. W. Wang is supported in part by Natural Science Foundation of China under grant No. 11735010, 11911530088, by Natural Science Foundation of Shanghai under grant No. 15DZ2272100. Q.-A. Zhang is supported by the China Postdoctoral Science Foundation and the National Postdoctoral Program for Innovative Talents (Grant No. BX20190207).

-
- [1] R. K. Ellis, W. J. Stirling and B. R. Webber, Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **8**, 1 (1996).
 - [2] H. W. Lin *et al.*, Prog. Part. Nucl. Phys. **100**, 107 (2018) doi:10.1016/j.pnpnp.2018.01.007 [arXiv:1711.07916 [hep-ph]].
 - [3] J. C. Collins and D. E. Soper, Nucl. Phys. B **193**, 381 (1981) Erratum: [Nucl. Phys. B **213**, 545 (1983)]. doi:10.1016/0550-3213(81)90339-4
 - [4] J. C. Collins, D. E. Soper and G. F. Sterman, Nucl. Phys. B **250**, 199 (1985). doi:10.1016/0550-3213(85)90479-1
 - [5] X. d. Ji, J. p. Ma and F. Yuan, Phys. Rev. D **71**, 034005 (2005) doi:10.1103/PhysRevD.71.034005 [hep-ph/0404183].
 - [6] X. d. Ji, J. P. Ma and F. Yuan, Phys. Lett. B **597**, 299 (2004) doi:10.1016/j.physletb.2004.07.026 [hep-ph/0405085].
 - [7] M. G. Echevarria, I. Scimemi and A. Vladimirov, Phys. Rev. D **93**, no. 5, 054004 (2016) doi:10.1103/PhysRevD.93.054004 [arXiv:1511.05590 [hep-ph]].
 - [8] Y. Li and H. X. Zhu, Phys. Rev. Lett. **118**, no. 2, 022004 (2017) doi:10.1103/PhysRevLett.118.022004 [arXiv:1604.01404 [hep-ph]].
 - [9] X. Ji, Y. Liu and Y. S. Liu, arXiv:1910.11415 [hep-ph].
 - [10] X. Ji, Phys. Rev. Lett. **110**, 262002 (2013) doi:10.1103/PhysRevLett.110.262002 [arXiv:1305.1539 [hep-ph]].
 - [11] X. Ji, Sci. China Phys. Mech. Astron. **57**, 1407 (2014) doi:10.1007/s11433-014-5492-3 [arXiv:1404.6680 [hep-ph]].
 - [12] J. Collins, Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **32**, 1 (2011).
 - [13] C. W. Bauer, S. Fleming, D. Pirjol and I. W. Stewart, Phys. Rev. D **63**, 114020 (2001)

- doi:10.1103/PhysRevD.63.114020 [hep-ph/0011336].
- [14] C. W. Bauer and I. W. Stewart, Phys. Lett. B **516**, 134 (2001) doi:10.1016/S0370-2693(01)00902-9 [hep-ph/0107001].
- [15] C. W. Bauer, D. Pirjol and I. W. Stewart, Phys. Rev. D **65**, 054022 (2002) doi:10.1103/PhysRevD.65.054022 [hep-ph/0109045].
- [16] X. Ji, Y. S. Liu, Y. Liu, J. H. Zhang and Y. Zhao, arXiv:2004.03543 [hep-ph].
- [17] K. Cichy and M. Constantinou, Adv. High Energy Phys. **2019**, 3036904 (2019) doi:10.1155/2019/3036904 [arXiv:1811.07248 [hep-lat]].
- [18] X. Ji, P. Sun, X. Xiong and F. Yuan, Phys. Rev. D **91**, 074009 (2015) doi:10.1103/PhysRevD.91.074009 [arXiv:1405.7640 [hep-ph]].
- [19] X. Ji, L. C. Jin, F. Yuan, J. H. Zhang and Y. Zhao, Phys. Rev. D **99**, no. 11, 114006 (2019) doi:10.1103/PhysRevD.99.114006 [arXiv:1801.05930 [hep-ph]].
- [20] M. A. Ebert, I. W. Stewart and Y. Zhao, Phys. Rev. D **99**, no. 3, 034505 (2019) doi:10.1103/PhysRevD.99.034505 [arXiv:1811.00026 [hep-ph]].
- [21] M. A. Ebert, I. W. Stewart and Y. Zhao, JHEP **1909**, 037 (2019) doi:10.1007/JHEP09(2019)037 [arXiv:1901.03685 [hep-ph]].
- [22] X. Ji, Y. Liu and Y. S. Liu, arXiv:1911.03840 [hep-ph].
- [23] A. A. Vladimirov and A. Schfer, Phys. Rev. D **101**, no. 7, 074517 (2020) doi:10.1103/PhysRevD.101.074517 [arXiv:2002.07527 [hep-ph]].
- [24] M. A. Ebert, I. W. Stewart and Y. Zhao, JHEP **2003**, 099 (2020) doi:10.1007/JHEP03(2020)099 [arXiv:1910.08569 [hep-ph]].
- [25] P. Shanahan, M. Wagman and Y. Zhao, arXiv:2003.06063 [hep-lat].
- [26] M. Bruno *et al.*, JHEP **1502**, 043 (2015) doi:10.1007/JHEP02(2015)043 [arXiv:1411.3982 [hep-lat]].
- [27] Supplemental materials.
- [28] R. G. Edwards *et al.* [SciDAC and LHPC and UKQCD Collaborations], Nucl. Phys. Proc. Suppl. **140**, 832 (2005) doi:10.1016/j.nuclphysbps.2004.11.254 [hep-lat/0409003].

SUPPLEMENTAL MATERIALS

A. Simulation checks




FIG. 6. The dispersion relation of the pion state with the pion mass from the 2pt function. The data up to $8\pi/L$ (~ 2 GeV) can be described with the formula $E_\pi = \sqrt{m_\pi^2 + c_1 P^2 + c_2 P^4 a^2}$ with $c_1 = 0.9945(40)$ and $c_2 = -0.0282(27)$. The deviation at $8\pi/L$ from the continuum limit is around 2%.

Fig. 6 shows the dispersion relation with the pion mass we used. The curve shows the fit based on the formula $E_\pi = \sqrt{m_\pi^2 + c_1 P^2 + c_2 P^4 a^2}$, where the last term in the square root parameterizes discretization errors. We used momenta up to $8\pi/L$ (~ 2 GeV). The fit gives results— $c_1 = 0.9945(40)$ and $c_2 = -0.0282(27)$ —that are consistent with the ground state energy calculated from two point function. It indicates only small discretization errors. Thus it is expected that the dispersion relation can recover the standard $E_\pi = \sqrt{m_\pi^2 + P^2}$ in the continuum limit.

Taking the form factor with $P^z = 6\pi/L$, $b_\perp = 3$, $t_2 = 8$ and $t = t_2/2$ as example, Fig. 7 shows the statistical check of the measurements we did. The left panel shows the histogram of 868 (configurations) $\times 48$ (time slides) = 41664 (measurements) with 54 exceptional measurements being dropped in the analysis since they give results deviating from the central ones by much more than 5σ . After we average the measurements over the same configuration, we find that the autocorrelation effect is negligible, since no obvious bin size dependence of the result is observed, as




FIG. 7. Statistical check on the simulation, taking the form factor with $P^z = 6\pi/L$, $b_\perp=3$, $t_2=8$ and $t = t_2/2$ as example. The left panel shows the histogram for all 41610 measurements (868 configurations $\times$ 48 time slides - 54 exceptional measurements), and the right panel shows the bin size dependence after we averaged all the measurements on the same configuration.

shown in the right panel of Fig. 7.




FIG. 8. Figure for the ℓ dependence of $|\phi_\ell(0, b_\perp, P^z, \ell)|$ with $\{P^z, b_\perp, t\} = \{6\pi/L, 3a, 6a\}$ (top left, the case shown in Fig. 2), $\{P^z, b_\perp, t\} = \{8\pi/L, 3a, 6a\}$ (top right), $\{P^z, b_\perp, t\} = \{6\pi/L, 6a, 6a\}$ (bottom left), and $\{P^z, b_\perp, t\} = \{6\pi/L, 3a, 8a\}$ (bottom right).

B. ℓ dependence of TMDWF

In Fig. 8, we give the ℓ dependence of $|\phi_\ell(0, b_\perp, P^z, \ell)|$ for a few more cases, similar to the $\{P^z, b_\perp, t\} = \{6\pi/L, 3a, 6a\}$ case shown in Fig. 2 but with larger P^z , $b_\perp$ and also t .

C. Two-state fit of the form factors

In this work, we use the following joint fit to obtain the norm of the subtracted quasi-TMDWF $|\phi_\ell(0, b_\perp, P^z, \ell)|$ and soft factor $S_I(b_\perp)$ (with $\ell = 7a$),

$$\begin{aligned} \frac{C_3(b_\perp, P^z, t_{\text{sep}}, t)}{C_2(0, P^z, 0, t_{\text{sep}})} &= \frac{|\tilde{\phi}_\ell(0, b_\perp, P^z, \ell)|^2 \tilde{S}_I(b_\perp) + C_1(e^{-\Delta E t} + e^{-\Delta E(t_{\text{sep}} - t)}) + C_2 e^{-\Delta E t_{\text{sep}}}}{1 + C_0 e^{-\Delta E t_{\text{sep}}}}, \\ \frac{C_2(b_\perp, P^z, 0, t)}{C_2(0, P^z, 0, t)} &= \frac{|\tilde{\phi}_\ell(0, b_\perp, P^z, \ell)| e^{\theta(b_\perp, P^z, \ell)} (1 + C_3 e^{-\Delta E t})}{1 + C_0 e^{-\Delta E t}}, \end{aligned} \quad (16)$$

where

$$\tilde{\phi}_\ell(0, b_\perp, P^z, \ell) = \frac{\phi_\ell(0, b_\perp, P^z, \ell)}{\phi_\ell(0, 0, P^z, 0)}, \quad \tilde{S}_I(b_\perp) = \frac{\phi_L(0, 0, P^z, 0) A_w}{2 E A_p} S_I(b_\perp), \quad (17)$$

and $\theta(b_\perp, P^z, \ell)$ is the phase of the quasi-TMDWF. The additional factor in the definition of $\tilde{S}_I$ will be cancelled by $\tilde{S}_I(b_\perp = a)$ in the ratio of Eq. (6).




FIG. 9. The ratios $C_3(b_\perp, P^z, t_{\text{sep}}, t)/C_2(0, P^z, 0, t_{\text{sep}})$ as function of t_{sep} and t , with $\{P^z, b_\perp\} = \{6\pi/L, 1a\}$ (left panel), $\{P^z, b_\perp\} = \{6\pi/L, 3a\}$ (right panel) and $\{P^z, b_\perp\} = \{6\pi/L, 5a\}$ (lower panel).

In Fig. 9, we show the ratios $C_3(b_\perp, P^z, t_{\text{sep}}, t)/C_2(0, P^z, 0, t_{\text{sep}})$ with $P^z = 6\pi/L$, $b_\perp = \{1a, 3a, 5a\}$, compared with the two-state fit predictions (colored bands) and fitted ground state contribution (gray band). All of them show a good agreement between data and fits.

As another check, we also consider the differential summed ratio

$$\begin{aligned} R(b_\perp, P^z, t_{\text{sep}}) &\equiv SR(b_\perp, P^z, t_{\text{sep}}) - SR(b_\perp, P^z, t_{\text{sep}} - 1) = |\tilde{\phi}_\ell(0, b_\perp, P^z, \ell)|^2 \tilde{S}_I(b_\perp) + \mathcal{O}(e^{-\Delta E t_{\text{sep}}}), \\ SR(b_\perp, P^z, t_{\text{sep}}) &\equiv \sum_{0 < t < t_{\text{sep}}} \frac{C_3(b_\perp, P^z, t_{\text{sep}}, t)}{C_2(0, P^z, 0, t_{\text{sep}})}. \end{aligned} \quad (18)$$

As an example, we plot $R(b_\perp, P^z, t_{\text{sep}})$ as function of t_{sep} in Fig. 10 for $\{P^z, b_\perp\} = \{6\pi/L, 3a\}$ and compare it with the standard two-state fit. We can see that the $R(b_\perp, P^z, t_{\text{sep}})$ agree with the ground state contribution from the two state fit at large t_{sep} .




FIG. 10. The ratios $C_3(b_\perp, P^z, t_{\text{sep}}, t)/C_2(0, P^z, 0, t_{\text{sep}})$ as function of t_{sep} and t compared with the differential summed ratio $R(b_\perp, P^z, t_{\text{sep}})$, with $\{P^z, b_\perp\} = \{6\pi/L, 3a\}$.

D. The possible imaginary part in extracting the Collins-Soper kernel




FIG. 11. The real (left panel) and imaginary (right panel) parts of the Collins-Soper kernel when the approximation $C(xP^z, \mu) = 1 + \mathcal{O}(\alpha_s)$ is taken, based on the definition in Eq. (19).

The Collins-Soper kernel with the following definition

$$\begin{aligned}
 K'(b_\perp, \mu) &= \frac{1}{\ln(P_1^z/P_2^z)} \ln \frac{C(xP_2^z, \mu) \Phi_{\overline{\text{MS}}}(x, b_\perp, P_1^z, \mu)}{C(xP_1^z, \mu) \Phi_{\overline{\text{MS}}}(x, b_\perp, P_2^z, \mu)} + \mathcal{O}(\gamma^{-2}) = \frac{1}{\ln(P_1^z/P_2^z)} \ln \frac{\phi(0, b_\perp, P_1^z)}{\phi(0, b_\perp, P_2^z)} + \mathcal{O}(\alpha_s, \gamma^{-2}) \\
 &= \frac{1}{\ln(P_1^z/P_2^z)} \ln \left| \frac{\phi(0, b_\perp, P_1^z)}{\phi(0, b_\perp, P_2^z)} \right| + \frac{i}{\ln(P_1^z/P_2^z)} (\theta(0, b_\perp, P_1^z) - \theta(0, b_\perp, P_2^z)) + \mathcal{O}(\alpha_s, \gamma^{-2})
 \end{aligned} \tag{19}$$

should be real, but the P^z dependence of the phase $\theta(0, b_\perp, P^z) = \tan^{-1} \frac{\phi_{\text{Im}}(0, b_\perp, P^z)}{\phi_{\text{Re}}(0, b_\perp, P^z)}$ can introduce an imaginary part of K' when the approximation $C(xP^z, \mu) = 1 + \mathcal{O}(\alpha_s)$ is employed. Fig. 11 shows the real and imaginary parts as functions of $b_\perp$ with two combinations of P_1/P_2 . The real part $K'_{\text{Re}} = K$ (left panel) corresponds to the definition used in the main text. The non-vanishing imaginary part (right panel) reflects the systematic uncertainty due to imprecise matching. K and $-|K'| = -\sqrt{K^2 + K_{\text{Im}}'^2}$ are still consistent within the statistical uncertainty of K as in Fig. 12.




FIG. 12. The comparison on K and $-|K'| = -\sqrt{K^2 + K_{\text{Im}}'^2}$. They are consistent with each other.

To estimate the inaccurate matching effects, we consider $|K'_{\text{Im}}|$ as a systematic uncertainty and add it with the statistical uncertainty of K in quadrature.