

Remarks on the tangent function from an analytic and a probability point of view.

Zbigniew J. Jurek (University of Wrocław)

February 15, 2021

Abstract. In this note we prove that the function $\tan(1/it)$ is a Voiculescu transform of a free-infinitely divisible distribution, that is, it admits the integral representation associated with Pick functions. Moreover, we found its "counterpart" in the classical infinitely divisible measures expressed by a series of Rademacher variables.

Mathematics Subject Classifications(2020): Primary 60E10, 60E07; Secondary 30D10

Key words and phrases: Pick functions; characteristic functions; infinite divisibility; Laplace transforms; Rademacher variables.

Abbreviated title: *On a tangent function from probability point of view.*

Author's address:

Institute of Mathematics, University of Wrocław, Pl. Grunwaldzki 2/4, 50-384 Wrocław, Poland;

Email: zjjurek@math.uni.wroc.pl ; www.math.uni.wroc.pl/~zjjurek

ORCID Number: 0000-0002-8557-2854

In this note, by quite elementary arguments, we prove that the tangent function $\tan(1/it)$, $t \neq 0$, is a transform of a free-infinitely divisible probability measure. Thus it admits the integral representation (Pick function restricted to the imaginary line); for instance cf. Bondesson (1992), p. 20.

However, if we know that both $1/z$ and $\tan(z)$ are Pick functions, and the property of being Pick function is closed under a superposition, we immediately conclude that $\tan(1/z)$ is a Pick function as well and hence we infer the

representation in question; see F. Gesztesy and E. Tsekanovskii (2000) and references therein. For some explicit examples, see there in Section A, p.121.

Our first aim here is not to use this structural property of Pick functions and the second one is to find for the tangent function its "counterpart part" in the classical infinite divisibility using an isomorphism given in Jurek (2016). More explicitly, we do it via a random integral mapping \mathcal{K} from Jurek (2007) and series representation of a hyperbolic tangent function; cf. Theorem 1.

In Jurek (2020) an analogy between different notions of infinite divisibility was studied and there, in Section 4, are given explicit examples of Pick functions – free-infinite divisible transforms – related to the hyperbolic and Laplace (double exponential) characteristic functions.

1. Notations and terminology.

For a finite Borel measure m on the real line \mathbb{R} and a Borel function g on a positive halfline $[0, \infty)$ we define a *characteristic function* $\phi_m(t)$ and a *Laplace transform* $\mathfrak{L}[g; w]$ as follows

$$\phi_m(t) := \int_{\mathbb{R}} e^{itx} m(dx), \quad t \in \mathbb{R}; \quad \mathfrak{L}[g(x); w] := \int_0^{\infty} g(x) e^{-wx} dx, \quad w > 0. \quad (1)$$

A probability measure $e(m) := e^{-m(\mathbb{R})} \sum_{k=0}^{\infty} \frac{m^{*k}}{k!}$, where $*$ denotes a convolution of measures, is called a *compound Poisson measure*. We will write $\mu = [a, \sigma^2, M]$, if a probability measure μ is *infinitely divisible* (that is: for each $k \geq 2$ there exist measure μ_k such that $\mu_k^{*k} = \mu$) and

$$\phi_{\mu}(t) = \exp\left[ita - \frac{1}{2}\sigma^2 t^2 + \int_{\mathbb{R} \setminus \{0\}} \left(e^{itx} - 1 - \frac{itx}{1+x^2}\right) M(dx)\right], \quad t \in \mathbb{R}; \quad (2)$$

where $a \in \mathbb{R}$ (a shift), σ^2 (a variance of a Gaussian part) and M is a measure satisfying a condition $\int_{\mathbb{R}} \min(1, x^2) M(dx) < \infty$ (so called a *Lévy measure*). The triplet a, σ^2, M is uniquely determined by μ and formula (2) is called *the Lévy-Khintchine representation*. By $(ID, *)$ we denote a convolution semi-group set of all infinitely divisible probability measures. Note that compound Poisson measures $e(m)$ are infinitely divisible and their characteristic functions are equal to $\phi_{e(m)}(t) = \exp \int_{\mathbb{R}} (e^{itx} - 1) m(dx)$.

A stochastic process $Y(t), t \geq 0$, on a probability space (Ω, \mathcal{F}, P) , starting from zero ($Y(0) = 0$ with P.1), with stationary and stochastic independent increments, and continuous in probability is called a *Lévy process*. Hence all increments $Y(t) - Y(s), t > s \geq 0$ have ID distributions and a probability distribution μ_t of random variable $Y(t)$ is equal to μ_1^{*t} . And conversely, each $\mu \in ID$ can be inserted into a Lévy process $Y_{\mu}(t), t \geq 0$, such that probability distribution of $Y_{\mu}(1) = \mu$.

Finally, below we need an random integral mapping

$$(ID, *) \ni \mu \rightarrow \mathcal{K}(\mu) = \mathcal{L}\left(\int_0^\infty s dY_\mu(1 - e^{-s})\right),$$

where $\mathcal{L}(V)$ denotes a probability distribution of a random variable V , that was introduced in Jurek (2007).

Our investigations of the tangent function have the origin in the following series representations:

$$\tanh s = 2s \sum_{k=1}^{\infty} \frac{1}{((2k-1)\pi/2)^2 + s^2} = s \int_{\mathbb{R}} \frac{1}{x^2 + s^2} \left(\sum_{n \in \mathbb{Z}, \text{odd}} \delta_{n\pi/2}(dx) \right) \quad (3)$$

where δ_a is a Dirac measure (point-mass measure concentrated at a); cf. Gradshteyn and Ryzhik (1994), formula **1.421**(2).

2. Results.

Here we prove the following analytic and probabilistic properties of a tangent function $\tan(1/it)$:

Theorem 1. (a) *Let's for $k \in \mathbb{N} := \{1, 2, 3, \dots\}$ define real constants*

$$c_k := ((2k-1)\pi/2)^{-1}, c_{-k} := (2(-k)+1)\pi/2)^{-1}$$

and define discrete measures

$$m(dx) := \sum_{k \in \mathbb{Z}} \frac{x^2}{1+x^2} \delta_{c_k}(dx); \quad M(dx) := \frac{1+x^2}{x^2} m(dx); \quad (4)$$

where δ_c denotes the Dirac measure. Then m is a finite measure with a mass $m(\mathbb{R}) = \tanh(1) \approx 0.76159$, and

$$\int_{\mathbb{R}} \frac{1+itx}{it-x} m(dx) = \tan(1/it) \quad (\equiv -i \tanh(1/t)), \quad t \neq 0; \quad (5)$$

(b) *Since $\int_{(\mathbb{R} \setminus 0)} (x^2 \wedge 1) M(dx) < \infty$ therefore $\mu = [0, 0, M]$ represents a classical infinitely divisible measure with zero shift, zero Gaussian part and Lévy (spectral) measure M . Let ϕ_μ be its characteristic function. Then*

$$it^2 \mathfrak{L}[\log \bar{\phi}_\mu(s); t] = it \log(\phi_{\mathcal{K}(\mu)}(-t^{-1})) = \tan(1/it) \quad (\equiv -i \tanh(1/t)), \quad (6)$$

where $\bar{\phi}_\mu$ is a complex conjugation and $\mathfrak{L}[g; t]$ denotes a Laplace transform of a function g at t .

(c) for a measure m and its characteristic function ϕ_m we have

$$(w^2 - 1)\mathfrak{L}[\phi_m(x) - \tanh(1) \cosh(x); w] = -i \tan\left(\frac{1}{iw}\right), \quad (\equiv -\tanh(1/w)) \quad (7)$$

where \mathfrak{L} stands for a Laplace transform.

(d) A measure $\mu = [0, 0, M] \in (ID, *)$, a "counterpart" of Pick function $\tan(1/it)$ (free-infinite divisible transform), is a probability distribution of a variable X with the following series representation

$$X = \sum_n c_n Y_n < \infty \text{ converges a.s., in distribution and in } L^2, \quad (8)$$

where Y_1, Y_2, \dots are i.i.d. copies of a random sum $Y = \sum_{j=0}^{N_2} r_j$ with N_2, r_1, r_2, \dots independent, where N_2 has Poisson distribution with the parameter 2 and r_j are Rademacher variables with $P(r_j = \pm 1) = 1/2$.

Since a function $\tan(z)$ is analytic one, from (5) via an extension arguments, we have that

Corollary 1. For $z \in \mathbb{C} \setminus \{0\}$

$$\tan(1/z) = \int_{\mathbb{R}} \frac{1 + zx}{z - x} m(dx), \text{ where } m(dx) := \sum_{n \in \mathbb{Z}, \text{ odd}} \frac{x^2}{1 + x^2} \delta_{(n\pi/2)^{-1}}(dx).$$

This representation is also in Gesztesy and Tsekanovskii (2000), Example A.10 and A.11 on p.122 but obtained by different complex analysis arguments. Also note that their examples A.8 and A.9 correspond to stable probability measures with an exponent r . Also $\tan(1/z)$, and much more of free-probability, can be found in Ejsmont and Lehner (2020).

The following lemma gives a probability distribution of the variable Y that appears in the part (d) of Theorem 1 and may be of some independent interest as well.

Lemma 1. Let $N_2, r_1, r_2, \dots, r_k, \dots$ be independent variable where N_2 has a Poisson distribution with a parameter 2 and $r_j = \pm 1$ with probability 1/2 are independent Rademacher variables. Then

$$Y := \sum_{j=1}^{N_2} r_j \stackrel{d}{=} S_+ - S_-, \text{ where } S_{\pm} \text{ are independent Poisson with parameter 1.}$$

Moreover, for $k = 0, 1, 2, \dots$ we have

$$P(Y = k) = P(Y = -k) = e^{-2} \sum_{j=0}^{\infty} \frac{1}{j!} \frac{1}{(k+j)!} \equiv e^{-2} I_k(2); \text{ (a Bessel function).}$$

In particular, $I_0(2) + 2 \sum_{k=1}^{\infty} I_k(2) = e^2$.

(cf. also Remark 2, below).

Remark 1. The above Lemma 1, is a particular case of so called Skellam distributions that are defined as a law of a difference $T_1 - T_2$ of independent Poisson variables with a parameters λ_1, λ_2 , respectively.

3. PROOFS.

Proof of Theorem 1.

Part (a). Noticing that for $k \in N$ we have $c_{-k} = -c_k$ and using (4) we

$$\begin{aligned} \int_{\mathbb{R}} \frac{1+itx}{it-x} m(dx) &= \sum_{k \in Z \setminus \{0\}} \frac{1+itc_k}{it-c_k} \frac{c_k^2}{1+c_k^2} = \sum_{k \in N} \left[\frac{1+itc_k}{it-c_k} + \frac{1+itc_{-k}}{it-c_{-k}} \right] \frac{c_k^2}{1+c_k^2} \\ &= -2it \sum_{k \in N} \frac{1+c_k^2}{t^2+c_k^2} \frac{c_k^2}{1+c_k^2} = -2it \sum_{k \in N} \frac{1}{t^2} \frac{1}{c_k^{-2}+1/t^2} \quad (\text{by(3)}) \\ &= -i2 \frac{1}{t} \sum_{k \in N} \frac{1}{((2k-1)\pi/2)^2 + 1/t^2} = -i \tanh(1/t) = \tan(1/it), \end{aligned}$$

which completes a proof of (4). Furthermore, using the one to the last equality above for $t = 1$, we conclude

$$m(\mathbb{R}) = \sum_{k \in Z} \frac{c_k^2}{1+c_k^2} = 2 \sum_{k \in N} \frac{1}{((2k-1)\pi/2)^2 + 1} = \tanh(1),$$

and this completes a proof of a part (a).

Part (b). In Jurek (2007) a definition of the mapping \mathcal{K} is given by a formula (12) and using Corollary 6 we get first equality in (6). (Also compare Jurek (2016)). Consequently, if

$$(ID, *) \ni \mu \rightarrow \mathcal{K}(\mu) = \mathcal{L}\left(\int_0^\infty s dY_\mu(1 - e^{-s})\right)$$

then its characteristic function is of the form:

$$\log(\phi_{\mathcal{K}(\mu)})(t) = \int_0^\infty \log(\phi_\mu(st)) e^{-s} ds, \quad (\text{by Corollary 4, in Jurek(2007)}).$$

Specifying above for $\mu = [0, 0, M]$, with a symmetric M , we get

$$\begin{aligned}
\log(\phi_{\mathcal{K}([0,0,M])})(t) &= \int_0^\infty \int_{\mathbb{R} \setminus \{0\}} (\cos(tsx) - 1) M(dx) e^{-s} ds \\
&= \int_{\mathbb{R} \setminus \{0\}} \left[\int_0^\infty (\cos(stx) - 1) e^{-s} ds \right] M(dx) = \int_{\mathbb{R} \setminus \{0\}} \left[\frac{1}{t^2 x^2 + 1} - 1 \right] M(dx) \\
&= -t^2 \int_{\mathbb{R} \setminus \{0\}} \frac{x^2}{t^2 x^2 + 1} M(dx) = -t^2 \sum_{k \in Z} \frac{c_k^2}{t^2 c_k^2 + 1} = -t^2 \sum_{k \in Z} \frac{1}{t^2 + c_k^{-2}} = \\
&\quad -t \tanh(t) = it \tan(it).
\end{aligned}$$

Substituting above t by $-1/t$ we get formula (6).

Part (c). Using a measure m , a formula (3), part (a) of Theorem 1 and Laplace transforms of a cosine and hyperbolic cosine functions we get

$$\begin{aligned}
&\mathfrak{L}[\phi_m(x) - \tanh(1) \cosh(x); w] \\
&= \int_0^\infty \left[\sum_{n \in Z, \text{odd}} \cos(wc_n) \frac{c_n^2}{1 + c_n^2} - \cosh(x) \sum_{n \in Z, \text{odd}} \frac{c_n^2}{1 + c_n^2} \right] e^{-wx} dx \\
&= \sum_{n \in Z, \text{odd}} \left[\frac{w}{w^2 + c_n^2} \frac{c_n^2}{1 + c_n^2} - \frac{w}{w^2 - 1} \frac{c_n^2}{c_n^2 + 1} \right] \\
&= \frac{w}{w^2 - 1} \sum_{n \in Z, \text{odd}} \left(\frac{w^2 - 1}{w^2 + c_n^2} - 1 \right) \frac{c_n^2}{1 + c_n^2} = -\frac{1}{w^2 - 1} \frac{1}{w} \sum_{n \in Z, \text{odd}} \frac{c_n^2}{1 + c_n^2 w^{-2}} \\
&= (w^2 - 1)^{-1} (-1) \tanh\left(\frac{1}{w}\right) = (w^2 - 1)^{-1} (-i) \tan\left(\frac{1}{iw}\right),
\end{aligned}$$

which completes a proof of a part (c).

Part(d). First, note that $\cos t$ is a characteristic function of Rademacher variables r_j . Second, variable Y has a compound Poisson distribution with a characteristic function $\exp[2(\cos t - 1)]$. Third, since $\mathbb{E}[Y] = 0$, $Var[Y] = 2$ therefore $\mathbb{E}[X] = 0$ and $Var(X) = \sum_n Var(c_n Y_n) = 2 \sum_n c_n^2 < \infty$. All in all, by Kolmogorov's Three Series Theorem we conclude that series (8) of independent variables converge in all three modes.

Finally, we have

$$\phi_X(t) = \prod_n \exp[2(\cos(c_n t) - 1)] = \exp \int_{\mathbb{R}} (\cos(tx) - 1) M(dx) = \phi_\mu(t),$$

which proves (8) and this completes a proof of Theorem 1.

Proof of Lemma 1.

Let S_+ and S_- denote a number of $(+1)$ and -1 among r_1, r_2, \dots, r_{N_2} , respectively. Then

$$[S_+ = k] \text{ iff } [(N_2 \geq k) \text{ and } (\sum_{j=1}^{N_2} r_j = k(+1) + (N_2 - k)(-1) = 2k - N_2)].$$

Hence using conditioning arguments (S_+ and S_- are N_2 conditionally independent) we get

$$\begin{aligned} P(S_+ = k) &= P((\sum_{j=1}^{N_2} r_j = 2k - N_2) \cap (N_2 \geq k)) = \mathbb{E}[1_{\{N_2 \geq k\}} \binom{N_2}{k} 2^{-N_2}] \\ &= \sum_{j=k}^{\infty} \binom{j}{k} \frac{1}{2^j} e^{-2} \frac{2^j}{j!} = e^{-2} \sum_{j=k}^{\infty} \frac{j!}{k! (j-k)! j!} = \frac{1}{k!} e^{-2} \sum_{n=0}^{\infty} \frac{1}{n!} = \frac{1}{k!} e^{-1}, \end{aligned}$$

and therefore S_+ has a Poisson distribution with a parameter 1.

By a similar argument S_- has also Poisson distribution with a parameter 1 and is independent of S_+ . Finally,

$$\begin{aligned} P(Y = k) &= P(S_+ - S_- = k) = P(S_+ - S_- = -k) \\ &= P((S_+ = k + S_-) \cap (\bigcup_{j=0}^{\infty} (S_- = j))) = \sum_{j=0}^{\infty} P((S_+ = k + j) \cap (S_- = j)) \\ &= \sum_{j=0}^{\infty} P(S_+ = k + j) P(S_- = j) = \sum_{j=0}^{\infty} e^{-1} \frac{1}{(k+j)!} e^{-1} \frac{1}{j!} \equiv e^{-2} I_k(2). \end{aligned}$$

(See Remark 3 below for a definition of a modified Bessel function $I_k(z)$.)

Remark 2. Since $I_k(2)$ are a probability distribution therefore

$$2 \sum_{k=1}^{\infty} I_k(2) + I_0(2) = e^2, \quad \text{where } I_k(2) := \sum_{j=0}^{\infty} \frac{1}{j!} \frac{1}{(k+j)!}, \quad k \geq 0. \quad (9)$$

But here is a straightforward calculation of that fact :

$$\begin{aligned} e^2 &= \sum_{k,l=0}^{\infty} \frac{1}{k!} \frac{1}{l!} = \sum_{k=0}^{\infty} \left(\frac{1}{k!}\right)^2 + 2 \sum_{k=0}^{\infty} \sum_{l=k+1}^{\infty} \frac{1}{k!} \frac{1}{l!} = I_0(2) + 2 \sum_{k=0}^{\infty} \sum_{j=1}^{\infty} \frac{1}{k!} \frac{1}{(k+j)!} \\ &= I_0(2) + 2 \sum_{k=0}^{\infty} \left[\frac{1}{k!} \left(\sum_{j=0}^{\infty} \frac{1}{(k+j)!} - \frac{1}{k!} \right) \right] = I_0(2) + 2 \sum_{j=0}^{\infty} \sum_{k=0}^{\infty} \left(\frac{1}{k!} \frac{1}{(k+j)!} - \left(\frac{1}{k!}\right)^2 \right) \\ &= I_0(2) + 2 \sum_{j=0}^{\infty} (I_j(2) - I_0(2)) = I_0(2) + 2 \sum_{j=1}^{\infty} I_j(2); \end{aligned}$$

which completes a proof of (9).

Remark 3. Recall that

$$I_\nu(z) := \sum_{j=0}^{\infty} \frac{1}{j! \Gamma(\nu + j + 1)} \left(\frac{z}{2}\right)^{\nu+2j}, \quad z \in \mathbb{C},$$

is a *modified Bessel function of the first kind*; cf. Gradshteyn and Ryzhik(1994), formula **8.445**. Moreover, it has the following integral representation

$$I_\nu(z) = \frac{\left(\frac{z}{2}\right)^\nu}{\Gamma(\nu + 1/2)\Gamma(1/2)} \int_{-1}^1 (1 - t^2)^{\nu-1/2} e^{\pm zt} dt; \quad \Re(\nu + 1/2) > 0;$$

Gradshteyn and Ryzhik (1994), formula **8.431(1)**.

Since $\Gamma(1/2) = \sqrt{\pi}$; $\Gamma(k + 1/2) = \sqrt{\pi} 2^{-k} (2k - 1)!!$ then

$$I_k(2) = \frac{2^k}{(2k - 1)!! \pi} \int_{-1}^1 (1 - x^2)^{k-1/2} e^{-2x} dx, \quad \text{for } k \in \mathbb{N},$$

that might be more useful for an explicit calculations.

Acknowledgment. This note was prompted by a very useful discussion with Dr W. Ejsmont (University of Wroclaw) who shared with Author his work in progress and provided a reference to Gesztesy and Tsekanovski (2000).

REFERENCES.

- [1] L. Bondesson (1992), *Generalized gamma convolutions and related classes of distributions and densities*, Springer-Verlag, New York; Lecture Notes in Statistics, vol. 76.
- [2] W. Ejsmont and F. Lehner (2020), The free tangent law, arXiv:2004.02679.
- [3] F. Gesztesy and E. Tsekanovskii (2000), On matrix-valued Herglotz functions, *Math. Nachr.* vol. **218** , pp. 61-138.
- [4] I. S. Gradshteyn and I. M. Ryzhik (1994), *Tables of integrals, series, and products*, Academic Press, San Diego, New York, Boston; 5th Edition.
- [5] Z. J. Jurek (2007), Random integral representations for free-infinitely divisible and tempered stable distributions, *Stat. & Probab. Letters*, **77** no. 4 , pp. 417-425.
Also on: arXiv:1206.3044, [math PR], 14 June 2012.
- [6] Z. J. Jurek (2016), On a method of introducing free-infinitely divisible probability measures, *Demonstratio Mathematica*, **49**,

no 2, pp. 236-251.

Also on: arXiv: 1412.4445, [math PR], 14 December 2014.

[7] Z. J. Jurek (2020), On a relation between classical and free infinitely divisible transforms, *Probab. Math. Stat.*, vol. 40, Fasc. 2, pp. 349-367.

Also on: arXiv:1707.02540 [math PR], 9 July 2017.