# A note on some reduction formulas for the incomplete beta function and the Lerch transcedent

J.L. González-Santander

Received: date / Accepted: date

**Abstract** We derive new reduction formulas for the incomplete beta function and the Lerch transcendent in terms of elementary functions. As an application, we calculate some new integrals. Also, we use these reduction formulas to test the performance of the algorithms devoted to the numerical evaluation of the incomplete beta function.

Keywords Incomplete beta function  $\cdot$  Lerch transcendent  $\cdot$  reduction formulas  $\cdot$  numerical evaluation of special functions

Mathematics Subject Classification (2010) 33B20 · 33B99

### 1 Introduction

The origins of the beta function  $B(\nu, \mu)$  go back to Wallis' attempt of the calculation of  $\pi$  [11]. For this purpose, he evaluated the integral

$$B(\nu,\mu) = \int_0^1 t^{\nu-1} (1-t)^{\mu-1} dt, \qquad (1)$$

where  $\nu$  and  $\mu$  are integers or  $\mu = 1$  and  $\nu$  is rational. Moreover, Wallis suggested that [1, p. 4]

$$\frac{\pi}{4} = \frac{1}{2} \int_0^1 t^{-1/2} \left(1-t\right)^{1/2} dx = \frac{1}{4} \lim_{n \to \infty} \left(\frac{2 \cdot 4 \cdot 6 \cdots 2n}{1 \cdot 3 \cdot 5 \cdots (2n-1)} \frac{1}{\sqrt{n}}\right)^2.$$

This result may have led Euler to consider the integral (1) for  $\nu$  and  $\mu$  not neccessarily integers and its relation to the gamma function. In fact, Euler

J.L. González-Santander

Department of Mathematics. Universidad de Oviedo.

Federico García Lorca 18, 33007 Oviedo, Spain.

Tel.: +34-985-10-3338

E-mail: gonzalezmarjuan @uniovi.es

derived the following relation between the beta and gamma functions [1, Eqn. 1.1.13]:

$$B(\nu,\mu) = \frac{\Gamma(\nu)\Gamma(\mu)}{\Gamma(\nu+\mu)}.$$

A natural generalization of the beta function is the incomplete beta function, defined as [8, Eqn. 8.17.1]

$$B(\nu,\mu,z) = \int_0^z t^{\nu-1} (1-t)^{\mu-1} dt, \qquad 0 \le z \le 1, \ \nu,\mu > 0,$$

where it is straightforward to continue analytically to complex values of  $\nu$ ,  $\mu$ , and z.

Many applications have been developed over time regarding the B  $(\nu, \mu, z)$  function. For instance, in statistics it is used extensively as the probability integral of the beta distribution [5, p. 210-275]. Also, it appears in statistical mechanics for Monte Carlo sampling [6], in the analysis of packings of granular objects [9], and in growth formulas in cosmology [4]. Therefore, in order to evaluate the B  $(\nu, \mu, z)$  function, it is quite interesting to have reduction formulas to simplify its computation, both symbolically and numerically. For instance, when  $\mu = m+1$  is a positive integer (i.e. m = 0, 1, 2, ...), we have the following reduction formula in terms of elementary functions [7, Eqn. 58:4:3]

$$B(\nu, m+1, z) = z^{\nu} \sum_{k=0}^{m} {m \choose k} \frac{(-z)^{k}}{k+\nu}.$$
 (2)

However, when  $\mu = 0$ , the incomplete beta function is given in terms of the Lerch transcendent [7, Eqn. 58:4:4]

$$B(\nu, 0, z) = z^{\nu} \Phi(z, 1, \nu), \quad \nu > 0,$$
(3)

where the Lerch transcendent is defined as [2, Eqn. 1.11(1)]

$$\Phi(z,s,\nu) = \sum_{k=0}^{\infty} \frac{z^k}{(k+\nu)^s}, \quad |z| < 1, \ \nu \neq 0, -1, -2, \dots$$
(4)

It is worth noting that (2) can be proved by induction from (3) and (4), applying the connection formula [7, Eqn. 58:5:3]:

$$B(\nu, \mu, z) = B(\nu + 1, \mu, z) + B(\nu, \mu + 1, z).$$

Nevertheless, reduction formulas for B  $(\nu, 0, z)$  when  $\nu$  is a rational number do not seem to be reported in the most common literature. The aim of this note is just to provide such reduction formulas in terms of elementary functions. As an application, we will calculate some new integrals in terms of elementary functions. Also, we will check that the numerical evaluation of the incomplete beta function is improved with these reduction formulas.

This paper is organized as follows. Section 2 derives reduction formulas for  $B(\nu, 0, z)$ , both for  $\nu$  positive rational, as well as negative rational. Particular

cases for  $\nu$  non-negative integer or  $\nu$  half-integer are also derived. In Section 3, we will apply the reduction formulas derived in Section 2 to calculate some integrals which do not seem to be reported in the most common literature. Further, for particular values of the parameters, the symbolic computation of these integrals is quite accelerated by using the aforementioned reduction formulas. Also, we will use these reduction formulas to numerically test the performance of the algorithm provided in MATHEMATICA to compute the incomplete beta function.

## 2 Main results

First, note that according to (3) and (4),

$$B(\nu, 0, z) = \sum_{k=0}^{\infty} \frac{z^{k+\nu}}{k+\nu},$$
(5)

thus B  $(\nu, 0, z)$  is divergent for non-positive integral values of  $\nu$ . Therefore, we will consider two different cases in this Section:  $\nu \in \mathbb{Q}^+$  and  $\nu \in \mathbb{Q}^- \setminus \{-1, -2, \ldots\}$ .

## 2.1 Case $\nu \in \mathbb{Q}^+$

Consider  $\nu = n + r > 0$  where  $n = \lfloor \nu \rfloor \geq 0$  is the integer part of  $\nu$  and  $0 \leq r \leq 1$ . From (5), we have

$$B(n+r,0,z) = \sum_{k=0}^{\infty} \frac{z^{k+n+r}}{k+n+r} = \sum_{k=n}^{\infty} \frac{z^{k+r}}{k+r}$$
$$= \sum_{k=0}^{\infty} \frac{z^{k+r}}{k+r} - \sum_{k=0}^{n-1} \frac{z^{k+r}}{k+r}.$$
(6)

Set r = 1 in (6) and then apply the Taylor expansion [8, Eqn. 4.6.1]

$$\log(1+z) = -\sum_{k=1}^{\infty} \frac{(-z)^k}{k},$$

to obtain

B 
$$(n+1,0,z) = -\log(1-z) - \sum_{k=1}^{n} \frac{z^k}{k},$$
 (7)  
 $n = 0, 1, 2, \dots$ 

Further, set r = 1/2 in (6) and apply de Taylor expansion [8, Eqn. 4.38.5]

$$\tanh^{-1} z = \sum_{k=0}^{\infty} \frac{z^{2k+1}}{2k+1},\tag{8}$$

to obtain

$$B\left(n+\frac{1}{2},0,z\right) = 2\left(\tanh^{-1}\sqrt{z} - \sum_{k=0}^{n-1}\frac{z^{k+1/2}}{2k+1}\right),$$

$$n = 0, 1, 2, \dots$$
(9)

More generally, set  $r = p/q \in \mathbb{Q}$  in (6) with p < q. Then,

$$B\left(n+\frac{p}{q},0,z\right) = z^{p/q} \sum_{k=0}^{\infty} \frac{z^k}{k+p/q} - \sum_{k=0}^{n-1} \frac{z^{k+p/q}}{k+p/q},$$
(10)

Rewrite the first sum of (10) as a hypergeometric function (see [1, p. 61-62]),

$$\sum_{k=0}^{\infty} \frac{z^k}{k + p/q} = \frac{1}{p/q} \, {}_2F_1\left(\begin{array}{c} 1, p/q \\ 1 + p/q \end{array} \middle| z\right). \tag{11}$$

Apply now the reduction formula [10, Eqn. 7.3.1.131]

$${}_{2}F_{1}\left(\begin{array}{c}1,p/q\\1+p/q\end{array}\middle|z\right)$$

$$=-\frac{p}{q}z^{-p/q}\sum_{k=0}^{q-1}\exp\left(-\frac{2\pi ipk}{q}\right)\log\left(1-z^{1/q}\exp\left(\frac{2\pi ik}{p}\right)\right),\qquad(12)$$

$$p,q=1,2,\ldots;\ p\leq q.$$

Therefore, taking into account (11)-(12), rewrite (10) as the following result.

**Theorem 1** For  $\nu = n + \frac{p}{q} \in \mathbb{Q}^+$ , with  $n = \lfloor \nu \rfloor$  and p < q, the reduction formula

$$B(\nu, 0, z) = z^{\nu} \Phi(z, 1, \nu)$$

$$= -\sum_{k=0}^{q-1} \exp\left(-\frac{2\pi i p k}{q}\right) \log\left(1 - z^{1/q} \exp\left(\frac{2\pi i k}{q}\right)\right) - \sum_{k=0}^{n-1} \frac{z^{k+p/q}}{k+p/q}.$$
(13)

holds true.

Remark 1 Notice that the reduction formula (9) is included in (13), but not (7), which is a singular case.

2.2 Case  $\nu \in \mathbb{Q}^- \backslash \left\{ -1, -2, \ldots \right\}$ 

Consider  $\nu = -n + r < 0$  where  $n = \lfloor |\nu - 1| \rfloor \ge 0$ , and 0 < r < 1. From (5), we have

$$B(-n+r,0,z) = \sum_{k=0}^{\infty} \frac{z^{k-n+r}}{k-n+r} = \sum_{k=-n}^{\infty} \frac{z^{k+r}}{k+r}$$
$$= \sum_{k=0}^{\infty} \frac{z^{k+r}}{k+r} + \sum_{k=1}^{n} \frac{z^{-k+r}}{-k+r}.$$
(14)

Taking r = 1/2 and applying again (8), we have

$$B\left(-n+\frac{1}{2},0,z\right) = 2\left(\tanh^{-1}\sqrt{z} - \sum_{k=1}^{n} \frac{z^{-k+1/2}}{2k-1}\right).$$
 (15)

More generally, take  $r = p/q \in \mathbb{Q}$  with p < q in (14), and apply (11) to obtain the following result.

**Theorem 2** For  $\nu = -n + \frac{p}{q} \in \mathbb{Q}^-$ , with  $n = \lfloor |\nu - 1| \rfloor$  and p < q, the reduction formula

$$B(\nu, 0, z)$$

$$= -\sum_{k=0}^{q-1} \exp\left(-\frac{2\pi i p k}{q}\right) \log\left(1 - z^{1/q} \exp\left(\frac{2\pi i k}{q}\right)\right) + \sum_{k=1}^{n} \frac{z^{p/q-k}}{p/q-k}.$$
(16)

holds true.

Remark 2 Notice that (15) is included in (16) as a particular case. Also, in (16), B  $(\nu, 0, z) \neq z^{\nu} \Phi(z, 1, \nu)$  since (3) does not hold true for  $\nu < 0$ .

## **3** Applications

From the reduction formulas obtained in Section 2, next we calculate some integrals in terms of elementary functions. Also, we will use these reduction formulas as a benchmark for the computation of the incomplete beta function.

#### 3.1 Calculation of integrals

On the one hand, an integral representation of the incomplete beta function is given by [7, Eqn. 58:3:5],

$$B(\nu,\mu,z) = z^{\nu} \int_0^1 t^{\nu-1} (1-zt)^{\mu-1} dt.$$
 (17)

Also, an integral representation of the Lerch transcendent is [2, Eqn. 1.11(3)]

$$\Phi(z, s, \nu) = \frac{1}{\Gamma(s)} \int_0^\infty \frac{t^{s-1} e^{-(\nu-1)t}}{e^t - z} dt, \quad \text{Re } \nu > 0.$$
(18)

Notice that from (3), (17), and (18), we have

$$\int_{0}^{1} \frac{t^{\nu-1}}{1-zt} dt = \int_{0}^{\infty} \frac{e^{-(\nu-1)t}}{e^{t}-z} dt = z^{-\nu} \mathbf{B}(\nu,0,z), \quad (19)$$
  
Re  $\nu > 0.$ 

Therefore, from (13) and (19), we have for  $\nu = n + \frac{p}{q} \in \mathbb{Q}^+$ , with  $n = \lfloor \nu \rfloor$  and p < q,

$$\int_{0}^{1} \frac{t^{\nu-1}}{1-zt} dt = \int_{0}^{\infty} \frac{e^{-(\nu-1)t}}{e^{t}-z} dt$$
(20)

$$= -z^{-n-p/q} \sum_{k=0}^{q-1} \exp\left(-\frac{2\pi i p k}{q}\right) \log\left(1 - z^{1/q} \exp\left(\frac{2\pi i k}{q}\right)\right) - \sum_{k=0}^{n-1} \frac{z^{k-n}}{k+p/q}.$$

On the other hand, in the literature we found [7, Eqn. 58:14:7]

$$\int_0^z \tanh^{2\lambda - 1} t \, dt = \frac{1}{2} \mathbf{B}\left(\lambda, 0, \tanh^2 z\right), \quad \text{Re } \lambda > 0.$$
(21)

Therefore, from (13) and (21), we have for  $\lambda = n + \frac{p}{q} \in \mathbb{Q}^+$ , with  $n = \lfloor \lambda \rfloor$  and p < q,

$$\int_{0}^{z} \tanh^{2\lambda - 1} t \, dt$$

$$= -\frac{1}{2} \sum_{k=0}^{q-1} \exp\left(-\frac{2\pi i p k}{q}\right) \log\left(1 - (\tanh z)^{2/q} \exp\left(\frac{2\pi i k}{q}\right)\right)$$

$$-\frac{1}{2} \sum_{k=0}^{n-1} \frac{(\tanh z)^{2(k+p/q)}}{k+p/q}.$$
(22)

The integral given in (22) generalizes the results found in the literature for  $\lambda = n + 1$  and  $\lambda = n + \frac{1}{2}$  with n = 0, 1, 2, ... [3, Eqns. 2.424.2-3].

It is worth noting that for particular values of  $\lambda$ , the Integrate MATHE-MATICA command is able to compute symbolically the same results as (22), but in a very time-consuming way. For instance, for  $\lambda = \frac{5}{4}$ , we obtain

$$\int_0^z \tanh^{3/2} t \, dt = \tanh^{-1} \left( \sqrt{\tanh z} \right) - 2\sqrt{\tanh z} + \tan^{-1} \left( \sqrt{\tanh z} \right),$$

but the Integrate command takes around 300 times longer than the reduction formula given in (22).



Fig. 1 Evaluation of Im  $(B(\nu, 0, z))$  with MATHEMATICA and (13) with  $\nu = 12.3$ .

#### 3.2 Numerical evaluation

From a numerical point of view, the reduction formulas (13) and (16) are quite useful to plot B ( $\nu$ , 0, z) as a function of  $\nu$  in the real domain. However, for some real values of  $\nu$  and z, we obtain a complex value for B ( $\nu$ , 0, z). In these cases, the imaginary part of B ( $\nu$ , 0, z) is not always easy to compute. Figure 1 shows the plot of Im (B ( $\nu$ , 0, z)) as a function of z for  $\nu = 12.3$ . The reduction formula (13) shows the correct answer, i.e. Im (B ( $\nu$ , 0, z)) =  $-\pi$ , meanwhile the numerical evaluation of Im (B ( $\nu$ , 0, z)) with MATHEMATICA diverges from this result. A similar feature is observed using (16) and a negative value for  $\nu$ . It is worth noting that the equivalent numerical evaluation of Im ( $z^{\nu}\Phi(z, 1, \nu)$ ) with MATHEMATICA yields also  $-\pi$ .

### 4 Conclusions

On the one hand, we have derived in (13) and (16) new expressions for the incomplete beta function B ( $\nu$ , 0, z) and the Lerch transcendent  $\Phi(z, 1, \nu)$  in terms of elementary functions when  $\nu$  is rational and z is complex. Particular formulas for non-negative integers values of  $\nu$  and for half-integer values of  $\nu$  are given in (7) and (9), (15) respectively.

On the other hand, we have calculated the integrals given (20) from the reduction formulas (13) and (16) and the integral representation of the incomplete beta function and the Lerch transcendent. Also, in (22), the integral  $\int_0^z \tanh^{\alpha} t \, dt$  is calculated in terms of elementary functions for  $\alpha \in \mathbb{Q}$  and  $\alpha > -1$ . It is worth noting that (22) accelerates quite significantly the symbolic computation of the latter integral with the aid of computer algebra.

Finally, with the aid of the reduction formulas (13) and (16), we have tested that the numerical algorithm provided by MATHEMATICA sometimes fails to compute the imaginary part of B ( $\nu$ , 0, z). Also, the reduction formulas (13) and (16) are numerically useful to plot B  $(\nu,0,z)$  as a function of  $\nu$  in the real domain.

All the results presented in this paper have been implemented in MATH-EMATICA and can be downloaded from https://bit.ly/2XT7UjK

### Conflict of interest

The authors declare that they have no conflict of interest.

#### References

- 1. Andrews, G.E., Askey, R., Roy, R.: Special functions. Encyclopedia of Mathematics and its Applications, Vol. 71. Cambridge University Press, New York (2004)
- 2. Erdélyi, A.: Higher transcendental functions. McGraw-Hill, New York (1953)
- 3. Gradshteyn, I.S., Ryzhik, I.M.: Table of integrals, series, and products, (7th ed.). Academic press, San Diego (2014)
- 4. Hamilton, A.J.S.: Formulae for growth factors in expanding universes containing matter and a cosmological constant. Mon. Not. R. Astron. Soc. 322(2), 419–425 (2001)
- Johnson, N.L., Kotz, S., Balakrishnan, N.: Continuous Univariate Distributions (2nd ed.), Vol. II. John Wiley & Sons Inc., New York (1995)
- Kofke, D.A.: Comment on "The incomplete beta function law for parallel tempering sampling of classical canonical systems" [J. Chem. Phys. 120, 4119 (2004)]. J. Chem. Phys. 121(2), 1167–1167 (2004)
- 7. Oldham, K. B., Myland, J., Spanier, J.: An atlas of functions: with equator, the atlas function calculator. Springer Science & Business Media, New York (2010)
- Olver, F.W.J., Lozier, D.W., Boisvert, R.F., Clark, C.W., (editors): NIST Handbook of Mathematical Functions. Cambridge University Press, New York (2010)
- Prellberg, T., Owczarek, A.L.: Stacking models of vesicles and compact clusters. J. Stat. Phys. 80, 755–779 (1995)
- Prudnikov, A.P., Brychov, Y.A., Marichev, O.I.: Integrals and Series. Vol. 3. More Special Functions. Gordon and Breach Science Publishers, New York, (1986)
- Wallis, J.: Arithmetica infinitorum. John Wallis, Operum Mathematicorum. Oxford, Leon Lichfield (1656)