

HOW SIMPLIFYING AND FLEXIBLE IS THE SIMPLIFYING ASSUMPTION IN PAIR-COPULA CONSTRUCTIONS - ANALYTIC ANSWERS IN DIMENSION THREE AND A GLIMPSE BEYOND

THOMAS MROZ, SEBASTIAN FUCHS, WOLFGANG TRUTSCHNIG

KEYWORDS

Pair copula, simplifying assumption, conditional distribution, Markov kernel, dependence

ABSTRACT. Motivated by the increasing popularity and the seemingly broad applicability of pair-copula constructions underlined by numerous publications in the last decade, in this contribution we tackle the unavoidable question on how flexible and simplifying the commonly used ‘simplifying assumption’ is from an analytic perspective and provide answers to two related open questions posed by Nagler and Czado in 2016. Aiming at a simplest possible setup for deriving the main results we first focus on the three-dimensional setting. We prove that the family of simplified copulas is flexible in the sense that it is dense in the set of all three-dimensional copulas with respect to the uniform metric d_∞ - considering stronger notions of convergence like the one induced by the metric D_1 , by weak conditional convergence, by total variation, or by Kullback-Leibler divergence, however, the family even turn out to be nowhere dense and hence insufficient for any kind of flexible approximation. Furthermore, returning to d_∞ we show that the partial vine copula is never the optimal simplified copula approximation of a given, non-simplified copula C , and derive examples illustrating that the corresponding approximation error can be strikingly large and extend to more than 28% of the diameter of the metric space. Moreover, the mapping ψ assigning each three-dimensional copula its unique partial vine copula turns out to be discontinuous with respect to d_∞ (but continuous with respect to D_1 and to weak conditional convergence), implying a surprising sensitivity of partial vine copula approximations. The afore-mentioned main results concerning d_∞ are then extended to the general multivariate setting.

1. INTRODUCTION

Pair-copula constructions (most well-known in the context of vine copulas) are a very popular bottom-up approach for constructing high-dimensional copulas out of several bivariate ones; they have a handy graphical representation and can be considered as an ordered sequence of trees. Aiming at a significant reduction of complexity it is usually assumed that the so-called *simplifying assumption*, saying that the copulas of the conditional distribution functions do not depend on the conditioning variables, holds.

Considering the enormous number of scientific contributions working with and applying simplified pair-copulas (see, e.g., [5, 6, 7, 32, 36, 38, 37]) it is quite surprising that, apart from a few critical voices (see, e.g., [2, 8, 15]), no analytic and systematic study on the approximation quality and flexibility of these concepts seems to have been published so far.

After an extensive literature research it seems that the publication coming closest to such a study was written by Spanhel and Kurz [34] who focus mainly on partial vine copulas (special simplified pair-copulas whose conditional distribution functions follow a certain intuitive construction principle) and show that partial vine copulas are optimal w.r.t. Kullback-Leibler divergence if the minimization is performed sequentially, but not

necessarily if the estimation is done jointly. As stated in [34], this “implies that it may not be optimal to specify the true copulas in the first tree” of a simplified pair-copula approximation.

Motivated by the broad applicability of pair-copula constructions, in this contribution we study flexibility and the extent of simplification imposed by the simplifying assumption from an analytic perspective. For the sake of generality of the construction we do not directly assume absolute continuity and work with densities but build the analysis on conditional distributions (Markov kernels). Although most results are established in the three-dimensional setting we also sketch possible extensions to the general multivariate case. We first introduce and discuss the somewhat more general concept of *simplified copulas*, i.e., copulas satisfying the simplifying assumption but do not necessarily follow the hierarchical bottom-up approach. We show that, on the one hand, simplified copulas are very flexible in the sense that they are dense in the family of all three-dimensional copulas with respect to the uniform metric d_∞ - this flexibility, however, gets lost when finer topologies like the one induced by the metric D_1 , by weak conditional convergence, by total variation metric or by the Kullback-Leibler divergence are considered. In fact, we prove that the family of simplified copulas is even *nowhere dense* with respect to either of these four topologies, and, thereby provide an answer to one of the questions posed by Nagler and Czado [27].

Returning to d_∞ we then show that the partial vine copula of a given, non-simplified copula C is *never* the best-possible simplified copula approximation of C (with respect to d_∞). More importantly, the error made by approximation via partial simplified vines may be strikingly large: in the worst case scenario the distance between a three-dimensional copula and its assigned partial vine copula is at least $3/16$ which corresponds to 28.125% of the diameter of the metric space. An analogous result holds in arbitrary dimensions, in this case the worst case distance is at least $1/8$. With these results we answer the question on “how far off can we be by assuming a simplified model?” also posed by Nagler and Czado [27].

Sticking to the analytic perspective we moreover focus on continuity properties of the mapping ψ assigning each three-dimensional copula its unique partial vine copula and show (among other things) that this mapping is *not* continuous with respect to d_∞ . In other words: if $d_\infty(A, B)$ is small then in general we can not infer that $d_\infty(\psi(A), \psi(B))$ is small too. As a direct consequence, although simplified pair-copulas are “highly flexible” ([22]) and partial vine copulas “can yield an approximation that is superior to competing approaches” ([34]), approximations in terms of partial vine copulas can be of very poor quality and lead to wrong conclusions.

The rest of this paper is organized as follows: Section 2 gathers preliminaries and notations that will be used in the sequel. In Section 3 we introduce simplified copulas in dimension $d = 3$, prove that the family of these copulas is dense in the metric space of all copulas with respect to d_∞ (Corollary 3.7) and then show that with respect to either of the afore-mentioned four notions of convergence the family is very small in the sense that it is nowhere dense. In Section 4 we then focus on partial vine copulas and study the afore-mentioned mapping ψ assigning each copula its simplified approximation. We discuss continuity of ψ with respect to different notions of convergence (some lemmata and proofs are moved to the Supplementary to facilitate reading) and provide the afore-mentioned lower bound for the worst-case scenario (Sections 5 and 6). To avoid unnecessary complexity, in the first few sections we proceed as [2, 17, 22, 30, 33] and restrict ourselves to the three-dimensional setting. To underline generality of our findings, however, in Section 7 we extend some of our main results to the general multivariate setting

and discuss the notion of so-called universally simplified copulas. Various examples and graphics illustrate both the obtained results and the ideas underlying the proofs.

2. NOTATION AND PRELIMINARIES

Throughout this paper we will write $\mathbb{I} := [0, 1]$ and let $d \geq 2$ be an integer, which will be kept fixed. Bold symbols will be used to denote vectors, e.g., $\mathbf{x} = (x_1, \dots, x_d) \in \mathbb{R}^d$. The d -dimensional Lebesgue measure will be denoted by λ^d , in case of $d = 1$ we will also simply write λ . We will let \mathcal{C}^d denote the family of all d -dimensional copulas, M will denote the comonotonicity copula, Π the independence copula and, for $d = 2$, W will denote the countermonotonicity copula (we omit the index indicating the dimension since no confusion will arise). For every $C \in \mathcal{C}^d$ the corresponding d -stochastic measure will be denoted by μ_C , i.e. $\mu_C([\mathbf{0}, \mathbf{u}]) = C(\mathbf{u})$ for all $\mathbf{u} \in \mathbb{I}^d$, and \mathcal{P}_C will denote the family of all d -stochastic measures. For more background on copulas and d -stochastic measures we refer to [10, 29]. For every metric space (S, δ) the Borel σ -field on S will be denoted by $\mathcal{B}(S)$.

In what follows Markov kernels will play a prominent role: A *Markov kernel* from \mathbb{R} to $\mathcal{B}(\mathbb{R}^{d-1})$ is a mapping $K : \mathbb{R} \times \mathcal{B}(\mathbb{R}^{d-1}) \rightarrow \mathbb{I}$ such that for every fixed $E \in \mathcal{B}(\mathbb{R}^{d-1})$ the mapping $y \mapsto K(y, E)$ is (Borel-)measurable and for every fixed $y \in \mathbb{R}$ the mapping $E \mapsto K(y, E)$ is a probability measure.

Given a real-valued random variable Y and a real-valued $(d - 1)$ -dimensional random vector \mathbf{X} on a probability space $(\Omega, \mathcal{A}, \mathbb{P})$ we say that a Markov kernel K is a *regular conditional distribution* of \mathbf{X} given Y if

$$K(Y(\omega), E) = \mathbb{E}(\mathbf{1}_E \circ \mathbf{X} | Y)(\omega)$$

holds \mathbb{P} -almost surely for every $E \in \mathcal{B}(\mathbb{R}^{d-1})$. It is well-known that for each random vector (\mathbf{X}, Y) a regular conditional distribution K of \mathbf{X} given Y always exists and is unique for \mathbb{P}^Y -a.e. $y \in \mathbb{R}$. If (\mathbf{X}, Y) has distribution function H (in which case we will also write $(\mathbf{X}, Y) \sim H$ and let μ_H denote the corresponding probability measure on $\mathcal{B}(\mathbb{R}^d)$) we will let K_H denote (a version of) the regular conditional distribution of \mathbf{X} given Y and simply refer to it as *Markov kernel of H* . If $C \in \mathcal{C}^d$ is a copula then we will consider the Markov kernel of C automatically as mapping $K_C : \mathbb{I} \times \mathcal{B}(\mathbb{I}^{d-1}) \rightarrow \mathbb{I}$. Defining the v -section of a set $G \in \mathcal{B}(\mathbb{I}^d)$ as $G_v := \{\mathbf{u} \in \mathbb{R}^{d-1} : (\mathbf{u}, v) \in G\}$ the so-called disintegration theorem yields

$$(2.1) \quad \int_{\mathbb{I}} K_C(v, G_v) \, d\lambda(v) = \mu_C(G)$$

so, in particular, in case of $G = \times_{i=1}^{d-1} G_i$ with $G_i = \mathbb{I}$ for all $i \neq j$ we have

$$\int_{\mathbb{I}} K_C(v, G) \, d\lambda(v) = \lambda(G_j).$$

For more background on conditional expectation and general disintegration we refer to [19, 23].

We call a copula $C \in \mathcal{C}^d$ *completely dependent* (w.r.t. the last coordinate) if there exist λ -preserving transformations $h_1, \dots, h_{d-1} : \mathbb{I} \rightarrow \mathbb{I}$ (i.e., transformations fulfilling $\lambda(h_i^{-1}(F)) = \lambda(F)$ for every $F \in \mathcal{B}(\mathbb{I})$) such that

$$K(y, E) := \mathbf{1}_E(h_1(y), \dots, h_{d-1}(y))$$

is a Markov kernel of C . Since the collection of all completely dependent copulas contains all shuffles of Min, it is dense in $(\mathcal{C}^d, d_\infty)$ (also see [26]). For more properties of complete dependence we refer to [25] as well as to [11] and the references therein.

Markov kernels can be used to define metrics stronger than the standard *uniform metric* d_∞ , defined by

$$d_\infty(C_1, C_2) := \max_{\mathbf{u} \in \mathbb{I}^d} |C_1(\mathbf{u}) - C_2(\mathbf{u})|$$

on \mathcal{C}^d . It is well known that the metric space $(\mathcal{C}^d, d_\infty)$ is compact and that pointwise and uniform convergence of a sequence of copulas $(C_n)_{n \in \mathbb{N}}$ are equivalent (see [10]). Following [11] and defining

$$\begin{aligned} D_1(C_1, C_2) &:= \int_{\mathbb{I}^{d-1}} \int_{\mathbb{I}} |K_{C_1}(v, [\mathbf{0}, \mathbf{u}]) - K_{C_2}(v, [\mathbf{0}, \mathbf{u}])| \, d\lambda(v) d\lambda^{d-1}(\mathbf{u}) \\ D_2(C_1, C_2) &:= \int_{\mathbb{I}^{d-1}} \int_{\mathbb{I}} (K_{C_1}(v, [\mathbf{0}, \mathbf{u}]) - K_{C_2}(v, [\mathbf{0}, \mathbf{u}]))^2 \, d\lambda(v) d\lambda^{d-1}(\mathbf{u}) \\ D_\infty(C_1, C_2) &:= \sup_{\mathbf{u} \in \mathbb{I}^{d-1}} \int_{\mathbb{I}} |K_{C_1}(v, [\mathbf{0}, \mathbf{u}]) - K_{C_2}(v, [\mathbf{0}, \mathbf{u}])| \, d\lambda(v) \end{aligned}$$

it can be shown that D_1, D_2 and D_∞ are metrics generating the same topology on \mathcal{C}^d and that the family of completely dependent copulas is closed with respect to these three metrics. In the sequel we will mainly work with D_1 and refer to [11] for more information on D_2 and D_∞ . The metric space (\mathcal{C}^d, D_1) is complete and separable but not compact.

Viewing copulas in terms of their conditional distributions and considering weak convergence gives rise to what we refer to as weak conditional convergence in the sequel: Consider a sequence of copulas $(C_n)_{n \in \mathbb{N}}$ and a copula C and let $(K_{C_n})_{n \in \mathbb{N}}$ and K_C be (versions of) the corresponding Markov kernels. We will say that $(C_n)_{n \in \mathbb{N}}$ converges *weakly conditional* (w.r.t. the last coordinate) to C if and only if for λ -almost every $v \in \mathbb{I}$ we have that the sequence $(K_{C_n}(v, \cdot))_{n \in \mathbb{N}}$ of probability measures on $\mathcal{B}(\mathbb{I}^{d-1})$ converges weakly to the probability measure $K_C(v, \cdot)$. In the latter case we will write $C_n \xrightarrow{\text{wcc}} C$ (where 'wcc' stands for 'weak conditional convergence').

According to Lemma 5 in [11] weak conditional convergence of $(C_n)_{n \in \mathbb{N}}$ to C implies convergence w.r.t. D_1 but not vice versa (see Example 2.1 below), and convergence w.r.t. D_1 implies convergence in d_∞ but not vice versa.

Example 2.1. For $d \geq 3$, $m \in \mathbb{N}$ and $k \in \{1, \dots, 2^m\}$ define $J_{m,k} := ((k-1)2^{-m}, k2^{-m}]$, set $n = 2^m + k - 2$ and consider the sequence of generalized EFGM copulas $(C_n)_{n \in \mathbb{N}}$ given by

$$C_n(\mathbf{u}, v) := v \prod_{i=1}^{d-1} u_i + f_n(v) \prod_{i=1}^{d-1} u_i(1 - u_i)$$

where $f_{2^m+k-2}(v) := \int_{[0,v]} \mathbf{1}_{J_{m,k}}(t) \, d\lambda(t)$. Then, for every $n \in \mathbb{N}$, the identity

$$K_{C_n}(v, [\mathbf{0}, \mathbf{u}]) = \prod_{i=1}^{d-1} u_i + f'_n(v) \prod_{i=1}^{d-1} u_i(1 - u_i)$$

holds for all $\mathbf{u} \in \mathbb{I}^{d-1}$ and almost all $v \in \mathbb{I}$. Thus, the sequence $(K_{C_n}(v, \cdot))_{n \in \mathbb{N}}$ fails to converge weakly to $K_\Pi(v, \cdot)$ for λ -almost all $v \in \mathbb{I}$, and it follows that $(C_n)_{n \in \mathbb{N}}$ does not

converge weakly conditional to Π . On the other hand, considering

$$\begin{aligned}
& \lim_{m \rightarrow \infty} \sup_{\mathbf{u} \in \mathbb{I}^{d-1}} \int_{\mathbb{I}} |K_{C_{2^m+k-2}}(v, [\mathbf{0}, \mathbf{u}]) - K_{\Pi}(v, [\mathbf{0}, \mathbf{u}])| \, d\lambda(v) \\
&= \lim_{m \rightarrow \infty} \sup_{\mathbf{u} \in \mathbb{I}^{d-1}} \int_{\mathbb{I}} \left| f'_{2^m+k-2}(v) \prod_{i=1}^{d-1} u_i(1-u_i) \right| \, d\lambda(v) \\
&= \lim_{m \rightarrow \infty} \lambda(J_{m,k}) \sup_{\mathbf{u} \in \mathbb{I}^{d-1}} \prod_{i=1}^{d-1} u_i(1-u_i) \\
&= 0
\end{aligned}$$

so $\lim_{n \rightarrow \infty} D_1(C_n, \Pi) = 0$. For a counterexample in the case $d = 2$ we refer to [20].

For any subset $J = \{j_1, \dots, j_{|J|}\} \subseteq \{1, \dots, d\}$ with $2 \leq |J| \leq d$ such that $j_k < j_l$ for all $k, l \in \{1, \dots, |J|\}$ with $k < l$ we let C_J denote the *marginal* copula of C with respect to the coordinates in J . If J only contains two indices i, j then we will sometimes also write C_{ij} instead of $C_{\{i,j\}}$ (no confusion will arise). Weak conditional convergence of a sequence of copulas transfers to marginal copulas:

Theorem 2.2. *Suppose that C, C_1, C_2, \dots are d -dimensional copulas. Then $C_n \xrightarrow{wcc} C$ implies*

$$(C_n)_{J \cup \{d\}} \xrightarrow{wcc} C_{J \cup \{d\}}$$

for every $J \subseteq \{1, \dots, d-1\}$ with $1 \leq |J| \leq d-1$.

Proof. Consider $J \subseteq \{1, \dots, d-1\}$ with $1 \leq |J| \leq d-1$ and w.l.o.g. assume that $J = \{1, \dots, |J|\}$. Disintegration implies that for every copula $C \in \mathcal{C}^d$ there exists some Markov kernel K_C such that C can be expressed as

$$C(\mathbf{u}, v) = \int_{[0, v]} K_C(t, [\mathbf{0}, \mathbf{u}]) \, d\lambda(t)$$

for all $(\mathbf{u}, v) \in \mathbb{I}^{d-1} \times \mathbb{I}$ and some Markov kernel $K_{C_{J \cup \{d\}}}$ such that we have

$$C_{J \cup \{d\}}(\mathbf{s}, v) = \int_{[0, v]} K_{C_{J \cup \{d\}}}(t, [\mathbf{0}, \mathbf{s}]) \, d\lambda(t)$$

for all $(\mathbf{s}, v) \in \mathbb{I}^{|J|} \times \mathbb{I}$. Thus

$$(2.2) \quad K_{C_{J \cup \{d\}}}(t, [\mathbf{0}, \mathbf{s}]) = K_C(t, [\mathbf{0}, \mathbf{s}] \times \mathbb{I}^{d-1-|J|})$$

holds for all $\mathbf{s} \in \mathbb{I}^{|J|}$ and λ -almost all $t \in \mathbb{I}$.

Suppose now that C, C_1, C_2, \dots are as in the theorem. Since projections are continuous, the Continuous Mapping Theorem and the previous identity imply that for λ -almost every $v \in \mathbb{I}$ weak convergence of the sequence $(K_{C_n}(v, \cdot))_{n \in \mathbb{N}}$ to $K_C(v, \cdot)$ implies weak convergence of the sequence $(K_{(C_n)_{J \cup \{d\}}}(v, \cdot))_{n \in \mathbb{N}}$ to $K_{C_{J \cup \{d\}}}(v, \cdot)$, which proves the assertion. \square

We complete this section with two additional notions of convergence considered, e.g., in Spanhel and Kurz [34], the Kullback-Leibler divergence (distance) KL and the total variation metric TV, and describe their relationship with D_1 and d_{∞} . Defining TV on \mathcal{C}^d by

$$TV(C_1, C_2) = \sup_{G \in \mathcal{B}(\mathbb{I}^d)} |\mu_{C_1}(G) - \mu_{C_2}(G)|,$$

convergence with respect to TV implies convergence with respect to D_1 :

Theorem 2.3. *The inequalities*

$$D_1(C_1, C_2) \leq D_\infty(C_1, C_2) \leq 2TV(C_1, C_2)$$

hold for all copulas $C_1, C_2 \in \mathcal{C}^d$. In particular, convergence w.r.t. TV implies convergence w.r.t. D_1 and D_∞ .

Proof. Fix $C_1, C_2 \in \mathcal{C}^d$. For every $\mathbf{u} \in \mathbb{I}^{d-1}$ setting

$$\Lambda_{\mathbf{u}} := \{v \in \mathbb{I} : K_{C_1}(v, [\mathbf{0}, \mathbf{u}]) > K_{C_2}(v, [\mathbf{0}, \mathbf{u}])\} \in \mathcal{B}(\mathbb{I})$$

we get ($\Lambda_{\mathbf{u}}^c := \mathbb{I} \setminus \Lambda_{\mathbf{u}}$)

$$\begin{aligned} 0 &\leq \int_{\mathbb{I}} |K_{C_1}(v, [\mathbf{0}, \mathbf{u}]) - K_{C_2}(v, [\mathbf{0}, \mathbf{u}])| d\lambda(v) \\ &= \int_{\Lambda_{\mathbf{u}}} K_{C_1}(v, [\mathbf{0}, \mathbf{u}]) - K_{C_2}(v, [\mathbf{0}, \mathbf{u}]) d\lambda(v) + \int_{\Lambda_{\mathbf{u}}^c} K_{C_2}(v, [\mathbf{0}, \mathbf{u}]) - K_{C_1}(v, [\mathbf{0}, \mathbf{u}]) d\lambda(v) \\ &= \mu_{C_1}(\Lambda_{\mathbf{u}} \times [\mathbf{0}, \mathbf{u}]) - \mu_{C_2}(\Lambda_{\mathbf{u}} \times [\mathbf{0}, \mathbf{u}]) + \mu_{C_2}(\Lambda_{\mathbf{u}}^c \times [\mathbf{0}, \mathbf{u}]) - \mu_{C_1}(\Lambda_{\mathbf{u}}^c \times [\mathbf{0}, \mathbf{u}]) \\ &= |\mu_{C_1}(\Lambda_{\mathbf{u}} \times [\mathbf{0}, \mathbf{u}]) - \mu_{C_2}(\Lambda_{\mathbf{u}} \times [\mathbf{0}, \mathbf{u}])| + |\mu_{C_1}(\Lambda_{\mathbf{u}}^c \times [\mathbf{0}, \mathbf{u}]) - \mu_{C_2}(\Lambda_{\mathbf{u}}^c \times [\mathbf{0}, \mathbf{u}])| \\ &\leq 2TV(C_1, C_2) \end{aligned}$$

from which the desired inequalities follow immediately. The first inequality has already been proved in [11, Lemma 3]. \square

It is well-known that KL divergence (which is not a metric and only well-defined for absolutely continuous copulas whose density is positive λ^d -almost everywhere) is stronger than TV (see the generalized Pinsker inequality in, e.h., [31]). Altogether we have the following interrelation, where $a \implies b$ indicates the convergence with respect to a implies convergence with respect to b (and the first implication is restricted to those copulas for which KL divergence is well-defined):

$$KL \implies TV \implies D_1 \iff D_\infty \implies d_\infty$$

3. SIMPLIFIED COPULAS

In this section we introduce three-dimensional so-called simplified copulas, i.e., copulas for which the conditional copulas do not depend on the conditioning variable. The enormous importance of this type of copulas is underlined by the fact that every copula can be approximated arbitrarily well with respect to d_∞ by simplified copulas (see Corollary 3.7). On the other hand, we will show that simplified pair-copula constructions may fail to approximate a given dependence structure w.r.t. d_∞ reasonably well (see Example 3.8). Additionally, we will see that the afore-mentioned denseness gets lost entirely when finer topologies or stronger metrics are considered, and prove that for D_1 (Theorem 3.9), for the total variation metric TV (Theorem 3.10), and the Kullback-Leibler (KL) divergence (Theorem 3.11) the family is even nowhere dense.

With very few exceptions, in literature pair-copula constructions are introduced by working with copula densities, i.e., all copulas are assumed to be absolutely continuous. Ensuring that no key idea of the underlying concept is left out and aiming at a setting as general as possible we deviate from this approach and work with Markov kernels instead.

In this and the subsequent three sections all conditioning will be done with respect to the last coordinate, notice that this does not impose any restriction (as can be seen from Theorem 3.10, Theorem 3.11, Remark 5.4 and Section 7).

According to disintegration for every copula $C \in \mathcal{C}^3$ there exists some Markov kernel K_C such that C can be expressed as

$$C(\mathbf{u}, v) = \int_{[0, v]} K_C(t, [\mathbf{0}, \mathbf{u}]) \, d\lambda(t)$$

for all $(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}$. Since K_C is a Markov kernel, for every $\mathbf{u} \in \mathbb{I}^2$ the mapping $t \mapsto K_C(t, [\mathbf{0}, \mathbf{u}])$ is measurable and for almost every $t \in \mathbb{I}$ the mapping $\mathbf{u} \mapsto K_C(t, [\mathbf{0}, \mathbf{u}])$ is a bivariate distribution function with (*conditional*) univariate marginal distribution functions $F_{1|3}(\cdot|t)$ and $F_{2|3}(\cdot|t)$ (conditional on t). Sklar's Theorem implies that for almost every $t \in \mathbb{I}$ there exists some (*conditional*) bivariate copula $C_{12;3}^t$ (conditional on t) satisfying

$$K_C(t, [\mathbf{0}, \mathbf{u}]) = C_{12;3}^t(F_{1|3}(u_1|t), F_{2|3}(u_2|t))$$

for all $\mathbf{u} \in \mathbb{I}^2$ such that the identity

$$(3.1) \quad C(\mathbf{u}, v) = \int_{[0, v]} C_{12;3}^t(F_{1|3}(u_1|t), F_{2|3}(u_2|t)) \, d\lambda(t)$$

holds for all $(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}$.

Remark 3.1.

(1) Since the (*conditional*) univariate marginal distribution functions satisfy $F_{1|3}(1|t) = 1 = F_{2|3}(1|t)$ for every $t \in \mathbb{I}$ the bivariate marginal copulas C_{13} and C_{23} of C satisfy

$$C_{13}(u_1, v) = \int_{[0, v]} C_{12;3}^t(F_{1|3}(u_1|t), F_{2|3}(1|t)) \, d\lambda(t) = \int_{[0, v]} F_{1|3}(u_1|t) \, d\lambda(t)$$

as well as $C_{23}(u_2, v) = \int_{[0, v]} F_{2|3}(u_2|t) \, d\lambda(t)$ for all $(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}$ and their corresponding Markov kernels fulfill

$$(3.2) \quad K_C(t, [0, u_1] \times \mathbb{I}) = K_{C_{13}}(t, [0, u_1]) = F_{1|3}(u_1|t)$$

$$(3.3) \quad K_C(t, \mathbb{I} \times [0, u_2]) = K_{C_{23}}(t, [0, u_2]) = F_{2|3}(u_2|t)$$

for all $\mathbf{u} \in \mathbb{I}^2$ and λ -almost all $t \in \mathbb{I}$ (compare with Equation (2.2)).

- (2) Notice that we choose this different notation for the (*conditional*) univariate distribution functions on purpose since this facilitates comprehending what follows.
- (3) For the copulas corresponding to the conditional bivariate distribution functions $K_C(t, \cdot)$ we write $C_{12;3}^t$ instead of $C_{12|3}^t$ and hence adopt the notation used in the literature (see, e.g., [34]).

The following two observations concerning Equation (3.1) are key:

(O1) the (*conditional*) bivariate copulas $C_{12;3}^t$ may depend on t ;

(O2) since the (*conditional*) univariate marginal distribution functions $F_{1|3}(\cdot|t)$ and $F_{2|3}(\cdot|t)$ may fail to be continuous the (*conditional*) bivariate copulas $C_{12;3}^t$ are not unique in general.

To the best of the authors' knowledge, the second observation has not yet been addressed in the literature which is somehow not surprising considering the fact that pair-copula constructions are usually focused on absolutely continuous copulas.

In the sequel we will study copulas C for which (O1) is not true, i.e., copulas for which the (*conditional*) copulas $C_{12;3}^t$ do not depend on t . We will refer to a copula $C \in \mathcal{C}^3$ as

generalized simplified (with respect to the third coordinate) if there exists some bivariate copula $A \in \mathcal{C}^2$ such that the identity

$$(3.4) \quad C(\mathbf{u}, v) = \int_{[0, v]} A(F_{1|3}(u_1|t), F_{2|3}(u_2|t)) \, d\lambda(t)$$

holds for all $(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}$. In the sequel $\mathcal{C}_{\text{GS}}^3$ will denote the family of all three-dimensional generalized simplified copulas.

The following first results (Theorem 3.2 and Corollary 3.3) imply that the family of generalized simplified copulas is very flexible.

Theorem 3.2. *Every completely dependent three-dimensional copula is generalized simplified.*

Proof. Let $C \in \mathcal{C}^3$ be a completely dependent copula, i.e., assume that there exist λ -preserving functions $h_1, h_2 : \mathbb{I} \rightarrow \mathbb{I}$ such that $K_C(v, E) := 1_E(h_1(v), h_2(v))$ is a Markov kernel of C . Considering

$$K_C(v, [\mathbf{0}, \mathbf{u}]) = 1_{[0, u_1] \times [0, u_2]}(h_1(v), h_2(v)) = 1_{[h_1(v), 1]}(u_1) 1_{[h_2(v), 1]}(u_2)$$

as well as $F_{1|3}(u_1|v) = 1_{[h_1(v), 1]}(u_1)$, $F_{2|3}(u_2|v) = 1_{[h_2(v), 1]}(u_2) \in \{0, 1\}$ it follows that for every copula $A \in \mathcal{C}^2$ the identity

$$C(\mathbf{u}, v) = \int_{[0, v]} A(F_{1|3}(u_1|t), F_{2|3}(u_2|t)) \, d\lambda(t)$$

holds for all $(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}$. This yields $C \in \mathcal{C}_{\text{GS}}^3$. \square

Note that completely dependent copulas are generalized simplified in the broadest sense since Equation (3.4) does not only hold for one or some copulas, it holds for every $A \in \mathcal{C}^2$.

Since the collection of all completely dependent copulas is dense in $(\mathcal{C}^3, d_\infty)$ Theorem 3.2 has the following consequence:

Corollary 3.3. *The collection of all generalized simplified copulas is dense in $(\mathcal{C}^3, d_\infty)$.*

Returning to observation (O2) in what follows we will mainly restrict ourselves to the family of copulas $C \in \mathcal{C}^3$ for which almost all (conditional) univariate marginal distribution functions $F_{1|3}(\cdot|t)$ and $F_{2|3}(\cdot|t)$ are continuous and let \mathcal{C}_c^3 denote the family of all these copulas. According to Sklar's theorem, for every copula $C \in \mathcal{C}_c^3$ the (conditional) bivariate copulas $C_{12;3}^t$ are unique for almost all $t \in \mathbb{I}$. Obviously the family of all absolutely continuous copulas $\mathcal{C}_{\text{ac}}^3$ is a subset of \mathcal{C}_c^3 , so for absolutely continuous copulas the conditional copulas are unique.

We will let $\mathcal{C}_S^3 := \mathcal{C}_{\text{GS}}^3 \cap \mathcal{C}_c^3$ denote the collection of all *simplified* copulas, i.e., the class of all three-dimensional copulas C which are generalized simplified *and* have continuous (conditional) univariate marginal distribution functions $F_{1|3}(\cdot|t)$ and $F_{2|3}(\cdot|t)$. In this case the copula $A \in \mathcal{C}^2$ in Equation 3.4 is unique and equals $C_{12;3}^t$ for almost all $t \in \mathbb{I}$.

Before proceeding we illustrate the above *simplifying assumption* in terms of the (Fréchet) class of all three-dimensional copulas C fulfilling that coordinates 1&3 as well as 2&3 are independent:

Example 3.4. (Class \mathcal{F}_Π^3 of three-dimensional copulas C satisfying $C_{13} = \Pi = C_{23}$)
For $C \in \mathcal{F}_\Pi^3$ we have $F_{1|3}(u_1|t) = u_1$ and $F_{2|3}(u_2|t) = u_2$ for all $\mathbf{u} \in \mathbb{I}^2$ and almost all $t \in \mathbb{I}$ implying $\mathcal{F}_\Pi^3 \subseteq \mathcal{C}_c^3$. If $D \in \mathcal{F}_\Pi^3$ is simplified then there exists some unique bivariate copula

$A \in \mathcal{C}^2$ such that

$$D(\mathbf{u}, v) = \int_{[0,v]} A(F_{1|3}(u_1|t), F_{2|3}(u_2|t)) \, d\lambda(t) = \int_{[0,v]} A(u_1, u_2) \, d\lambda(t) = A(\mathbf{u}) v$$

holds for all $(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}$.

(1) The independence copula $\Pi \in \mathcal{F}_{\Pi}^3$ satisfies

$$\Pi(\mathbf{u}, v) = \int_{[0,v]} \Pi(F_{1|3}(u_1|t), F_{2|3}(u_2|t)) \, d\lambda(t)$$

for all $(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}$. Thus, Π is simplified, obviously $\Pi(\mathbf{u}, v) = \Pi(\mathbf{u}) v$ holds for all $(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}$.

(2) The EFGM copula $C^{\text{EFGM}} \in \mathcal{F}_{\Pi}^3$, given by

$$C^{\text{EFGM}}(\mathbf{u}, v) := \Pi(\mathbf{u}, v) + u_1(1 - u_1) u_2(1 - u_2) v(1 - v)$$

satisfies

$$C^{\text{EFGM}}(\mathbf{u}, v) = \int_{[0,v]} (C^{\text{EFGM}})_{12;3}^t(F_{1|3}(u_1|t), F_{2|3}(u_2|t)) \, d\lambda(t)$$

for all $(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}$, where

$$(C^{\text{EFGM}})_{12;3}^t(\mathbf{u}) = u_1 u_2 + (1 - 2t) u_1(1 - u_1) u_2(1 - u_2)$$

for all $\mathbf{u} \in \mathbb{I}^2$ and almost all $t \in \mathbb{I}$. Thus, C^{EFGM} is non-simplified.

(3) The copula $C^{\text{Cube}} \in \mathcal{F}_{\Pi}^3$ which distributes mass uniformly within the four cubes

$$\begin{array}{cc} (0, \frac{1}{2}) \times (0, \frac{1}{2}) \times (0, \frac{1}{2}) & (\frac{1}{2}, 1) \times (\frac{1}{2}, 1) \times (0, \frac{1}{2}) \\ (0, \frac{1}{2}) \times (\frac{1}{2}, 1) \times (\frac{1}{2}, 1) & (\frac{1}{2}, 1) \times (0, \frac{1}{2}) \times (\frac{1}{2}, 1) \end{array}$$

and has no mass outside these cubes satisfies

$$C^{\text{Cube}}(\mathbf{u}, v) = \int_{[0,v]} (C^{\text{Cube}})_{12;3}^t(F_{1|3}(u_1|t), F_{2|3}(u_2|t)) \, d\lambda(t)$$

for all $(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}$, where $(C^{\text{Cube}})_{12;3}^t = A^1$ for almost all $t \in (0, \frac{1}{2})$ and $(C^{\text{Cube}})_{12;3}^t = A^2$ for almost all $t \in (\frac{1}{2}, 1)$, and the copulas A^1 and A^2 are checkerboard copulas (see [10] for a general definition) whose density is depicted in Figure 1. As a direct consequence C^{Cube} is non-simplified.




FIGURE 1. Mass distribution of the copulas A^1 , A^2 and C^{Cube} from Example 3.4.

In contrast to the afore-mentioned class, some copula families only contain simplified copulas:

Example 3.5. [17, 35]

- (1) All three-dimensional Gaussian and Student t -copulas are simplified.
- (2) The only three-dimensional Archimedean copulas that are simplified are those of Clayton type.

We now focus on empirical copulas, show that they are simplified and then conclude that \mathcal{C}_S^3 is dense in $(\mathcal{C}^3, d_\infty)$ (Corollary 3.7).

Consider a random vector (\mathbf{X}, Y) with continuous univariate marginals and suppose that $(\mathbf{X}_1, Y_1), \dots, (\mathbf{X}_n, Y_n)$ is a sample from (\mathbf{X}, Y) . Since the univariate marginals are continuous w.l.o.g. we can assume that there are no ties. Let \hat{C}_n denote the empirical copula (by which we mean the unique copula determined by trilinear interpolation of the empirical subcopula). Then there exist two permutations σ_1, σ_2 of $\{1, \dots, n\}$ such that the density \hat{c}_n of \hat{C}_n is given by (uniform distribution on n cubes of volume $\frac{1}{n^3}$)

$$\hat{c}_n(u_1, u_2, v) = n^2 \sum_{i=1}^n \mathbb{1}_{I_i^1}(u_1) \mathbb{1}_{I_i^2}(u_2) \mathbb{1}_{V_i}(v)$$

where $I_i^1 = (\frac{\sigma_1(i)-1}{n}, \frac{\sigma_1(i)}{n}]$, $I_i^2 = (\frac{\sigma_2(i)-1}{n}, \frac{\sigma_2(i)}{n}]$ and $V_i = (\frac{i-1}{n}, \frac{i}{n}]$, so the Markov kernel of \hat{C}_n fulfills

$$(3.5) \quad K_{\hat{C}_n}(v, [0, u_1] \times [0, u_2]) = n^2 \sum_{i=1}^n \left(\int_{[0, u_1]} \mathbb{1}_{I_i^1}(t) d\lambda(t) \int_{[0, u_2]} \mathbb{1}_{I_i^2}(s) d\lambda(s) \right) \mathbb{1}_{V_i}(v).$$

Theorem 3.6. *Every three-dimensional empirical copula is simplified.*

Proof. Considering that the (conditional) univariate marginal distribution functions $(\hat{F}_n)_{1|3}(\cdot|v)$, $(\hat{F}_n)_{2|3}(\cdot|v)$ of \hat{C}_n are continuous and given by

$$\begin{aligned} (\hat{F}_n)_{1|3}(u_1|v) &= n \sum_{i=1}^n \left(\int_{[0, u_1]} \mathbb{1}_{I_i^1}(t) d\lambda(t) \right) \mathbb{1}_{V_i}(v) \\ (\hat{F}_n)_{2|3}(u_2|v) &= n \sum_{i=1}^n \left(\int_{[0, u_2]} \mathbb{1}_{I_i^2}(t) d\lambda(t) \right) \mathbb{1}_{V_i}(v) \end{aligned}$$

using Equation (3.5) it follows immediately that $K_{\hat{C}_n}(v, [0, u_1] \times [0, u_2])$ can be expressed as

$$K_{\hat{C}_n}(v, [0, u_1] \times [0, u_2]) = \Pi((\hat{F}_n)_{1|3}(u_1|v), (\hat{F}_n)_{2|3}(u_2|v))$$

from which it follows that \hat{C}_n is simplified. \square

Since the collection of all empirical copulas is dense in $(\mathcal{C}^3, d_\infty)$ (see [9, Proposition 3.2]), Theorem 3.6 has the following consequence (for a stronger and more general result see Corollary 7.2):

Corollary 3.7. *The collection of all simplified copulas is dense in $(\mathcal{C}^3, d_\infty)$.*

Although every copula can be approximated arbitrarily well by simplified copulas a reasonable approximation from the same Fréchet class might not be possible as the following example illustrates:

Example 3.8. (Class \mathcal{F}_Π^3 , cont.)

For the non-simplified copula $C^{\text{Cube}} \in \mathcal{F}_\Pi^3$ introduced in Example 3.4 there exists some $\varepsilon > 0$ such that for every simplified copula $D \in \mathcal{F}_\Pi^3$ we have

$$d_\infty(C^{\text{Cube}}, D) > \varepsilon,$$

which can be shown as follows: Recall that every simplified copula D from this class fulfills $D(\mathbf{u}, v) = A(\mathbf{u})v$ for all $(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}$, where A is some bivariate copula. Furthermore recall that C^{Cube} fulfills

$$C^{\text{Cube}}(\mathbf{u}, v) = \int_{[0, v]} (C^{\text{Cube}})_{12;3}^t(\mathbf{u}) \, d\lambda(t)$$

for all $(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}$, where $(C^{\text{Cube}})_{12;3}^t = A^1$ for almost all $t \in (0, \frac{1}{2})$ and $(C^{\text{Cube}})_{12;3}^t = A^2$ for almost all $t \in (\frac{1}{2}, 1)$, and A^1 and A^2 are bivariate copulas with $A^1 \neq A^2$ (see Example 3.4). Thus,

$$C^{\text{Cube}}(\mathbf{u}, v) = \begin{cases} A^1(\mathbf{u})v & v \in [0, \frac{1}{2}] \\ A^1(\mathbf{u})\frac{1}{2} + A^2(\mathbf{u})(v - \frac{1}{2}) & v \in (\frac{1}{2}, 1] \end{cases}$$

and hence

$$\begin{aligned} & |C^{\text{Cube}}(\mathbf{u}, v) - D(\mathbf{u}, v)| \\ &= \begin{cases} |A^1(\mathbf{u}) - A(\mathbf{u})|v & v \in [0, \frac{1}{2}] \\ |[A^1(\mathbf{u}) - A(\mathbf{u})]\frac{1}{2} + [A^2(\mathbf{u}) - A(\mathbf{u})](v - \frac{1}{2})| & v \in (\frac{1}{2}, 1] \end{cases} \end{aligned}$$

for all $(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}$. If $A = A^1$, then

$$d_\infty(C^{\text{Cube}}, D) \geq |C^{\text{Cube}}(\frac{1}{2}, \frac{3}{4}) - D(\frac{1}{2}, \frac{3}{4})| = \frac{1}{4} |A^2(\frac{1}{2}) - A^1(\frac{1}{2})| = \frac{1}{4} |0 - \frac{1}{2}| = \frac{1}{8}.$$

If $A \neq A^1$ then there exists some $\mathbf{u}^* \in \mathbb{I}^2$ and some $\varepsilon > 0$ with $|A(\mathbf{u}^*) - A^1(\mathbf{u}^*)| > 4\varepsilon$ and hence

$$d_\infty(C^{\text{Cube}}, D) \geq |C^{\text{Cube}}(\mathbf{u}^*, \frac{1}{4}) - D(\mathbf{u}^*, \frac{1}{4})| = \frac{1}{4} |A^1(\mathbf{u}^*) - A(\mathbf{u}^*)| > \varepsilon$$

Thus C^{Cube} can not be approximated arbitrarily well by a simplified copula D from the class $\mathcal{F}_{\mathbb{I}}^3$.

We now focus on the afore-mentioned stronger metrics or finer topologies on \mathcal{C} (or important subclasses). To simplify notation we will write $\mathcal{C}_{\text{ac}, >0}^3$ for the collection of all absolutely continuous copulas with positive density.

Theorem 3.9.

- (1) *The collection of all simplified copulas is nowhere dense in (\mathcal{C}^3, D_1) and $(\mathcal{C}^3, D_\infty)$.*
- (2) *The collection of all simplified copulas is nowhere dense in \mathcal{C}^3 with respect to the topology induced by weak conditional convergence.*
- (3) *The collection of all simplified copulas with positive density is nowhere dense in $(\mathcal{C}_{\text{ac}, >0}^3, D_1)$ and $(\mathcal{C}_{\text{ac}, >0}^3, D_\infty)$.*

Proof. To prove the first assertion assume that the D_1 -closure of the family of all simplified copulas contains an open ball $O_{D_1}(C, r) = \{A \in \mathcal{C}^3 : D_1(A, C) < r\}$ with $C \in \mathcal{C}$ and $r > 0$. Since according to Lemma A.6 non-simplified checkerboard copulas are dense in (\mathcal{C}^3, D_1) we can find a non-simplified checkerboard copula $C^* \in O_{D_1}(C, r)$. Since, by assumption, the D_1 -closure of the family of all simplified copulas contains $O_{D_1}(C, r)$ there exists a sequence $(C_n)_{n \in \mathbb{N}}$ of simplified copulas with $\lim_{n \rightarrow \infty} D_1(C_n, C^*) = 0$, a contradiction to Lemma A.5.

Proceeding analogously yields the second and the third assertion. \square

Theorem 3.9 and Theorem 2.3 imply the following two striking results:

Theorem 3.10.

- (1) *The collection of all simplified copulas is nowhere dense in (\mathcal{C}^3, TV) .*
- (2) *The collection of all simplified copulas with positive density is nowhere dense in $(\mathcal{C}_{ac, >0}^3, TV)$.*

Theorem 3.11. *The collection of all simplified copulas with positive density is nowhere dense in $(\mathcal{C}_{ac, >0}^3, KL)$.*

Theorems 3.9, 3.10 and 3.11 answer the question “How dense does the set of simplified densities lie in the set of all densities?” posed by Nagler and Czado [27] in a complete and definitive manner.

In the same article the authors also pose the question on “how far off can we be by assuming a simplified model?” - one of the main objectives of the subsequent sections is to answer this very question. Notice that, for this purpose, we can restrict ourselves to the metric d_∞ since (according to the afore-mentioned results) simplified copulas are nowhere dense w.r.t. D_1 , D_∞ , TV and KL.

4. SIMPLIFIED PAIR-COPULA CONSTRUCTIONS

Equation (3.1) suggests the construction of a three-dimensional copula in terms of two families of (conditional) univariate marginal distribution functions characterizing the dependence structure between coordinates 1&3 and coordinates 2&3, respectively, and (conditional) bivariate copulas representing the dependence structure between coordinates 1&2 conditional on the third variable. This just-mentioned construction principle is called *vine decomposition* or *pair-copula construction* (see [1, 3]). In case the conditioning variable only enters indirectly through the conditional marginals (as it is the case in Equation (3.4); see, e.g., [18] for an early reference), the pair-copula construction is said to be *simplified* (see [17]).

4.1. Construction principle. Simplified pair-copula constructions are used to approximate the data generating copula (from \mathcal{C}_c^3) by a simplified copula (from \mathcal{C}_S^3) using the following hierarchical bottom-up algorithm based on Equation (3.4):

- (1) Estimation of the (conditional) univariate marginal distribution functions $F_{1|3}(\cdot|t)$ and $F_{2|3}(\cdot|t)$ conditional on t ;
- (2) Estimation of the (conditional) copula A of coordinates 1&2 conditional on variable 3 assuming that the conditioning variable enters only through the arguments of the conditional copula A (simplifying assumption).

The estimation is either done step-by-step or jointly, parametric or non-parametric, for more information we refer to [1, 2, 16, 17, 21, 27, 34] and the references therein. For an additional discussion about estimating conditional copulas satisfying the simplifying assumption (step (2)), we additionally refer to [8, 13, 14, 30].

The 3-dimensional copula resulting from this algorithm is simplified and is said to be a *simplified vine copula* (SVC). Apparently, the above algorithm and thus its output, the SVC, depend on the estimation method used and also on the suitable family of copulas from which the estimators are selected. The above algorithm may certainly provide a reasonable estimator if the (data generating) copula is simplified. The natural question arising at this point, however, is how well an SVC approximates the data generating copula if the latter fails to be simplified. We start with the following example also discussed in [35, Section 5]:

Example 4.1. The EFGM copula $C^{\text{EFGM}} \in \mathcal{F}_{\Pi}^3$ introduced in Example 3.4 is non-simplified. Minimizing the Kullback-Leibler divergence between the conditional copula

and its estimator selected from the family of all bivariate EFGM copulas in step (2) yields the bivariate independence copula as the optimal approximation. The SVC selected by a step-by-step algorithm hence equals the three-dimensional independence copula.

Comparing the data generating copula with its selected SVC yields a d_∞ -distance of $1/64$; this equals 6.25% of the maximal d_∞ -distance of two copulas within the (Fréchet) class of all copulas having pairwise independent marginals (using the results in [28, Section 3.3] it is straightforward to verify that the diameter of this class is $1/4$).

We refer to [2, 17, 35] for more examples and comparisons of the data generating copula with its selected simplified vine copula whereby the quality of the approximations is judged quite differently.

Aiming to obtain more general analytic results concerning the optimality of simplified pair-copula constructions, in what follows we discuss the concept of partial vine copulas.

4.2. Partial vine copulas (PVCs). The basic idea behind a partial vine copula is that the conditional bivariate copulas of the original three-dimensional copula are averaged (see [33, 34]):

Considering that for every $C \in \mathcal{C}_c^3$ the copula $C_{12;3}^t$ is unique for almost every $t \in \mathbb{I}$ it follows that the function $C_p : \mathbb{I}^2 \rightarrow \mathbb{I}$, given by

$$C_p(\mathbf{s}) := \int_{\mathbb{I}} C_{12;3}^t(\mathbf{s}) \, d\lambda(t)$$

is well-defined. In the sequel we will refer to C_p as the *partial copula* of C (also see [4]). Coinciding with the expected conditional copula, the partial copula is often used as an approximation of the conditional copula (see [33, 34] for more information). Given C_p in the above setting the mapping $\psi : \mathcal{C}_c^3 \rightarrow \mathcal{C}_c^3$, given by

$$(\psi(C))(\mathbf{u}, v) := \int_{[0,v]} C_p(F_{1|3}(u_1|t), F_{2|3}(u_2|t)) \, d\lambda(t)$$

is well-defined and assigns to every copula $C \in \mathcal{C}_c^3$ a simplified copula $\psi(C)$. The copula $\psi(C)$ is referred to as the *partial vine copula* of C (with respect to the third coordinate) in the sequel. It is obvious that every partial vine copula is simplified.

The transformation ψ preserves the dependence structure between coordinates 1&3 as well as between coordinates 2&3. The following lemma gathers some additional properties of ψ :

Lemma 4.2. *Suppose that $C \in \mathcal{C}_c^3$. Then the following assertions hold:*

- (1) *The partial vine copula $\psi(C)$ of C satisfies $(\psi(C))_{13} = C_{13}$ as well as $(\psi(C))_{23} = C_{23}$.*
- (2) *If C is simplified then $\psi(C) = C$ holds.*
- (3) *The mapping $\psi : \mathcal{C}_c^3 \rightarrow \mathcal{C}_c^3$ is not injective.*

Proof. Since $F_{1|3}(1|t) = 1 = F_{2|3}(1|t)$ for almost every $t \in \mathbb{I}$ we have

$$\begin{aligned} (\psi(C))_{13}(u_1, v) &= \int_{[0,v]} C_p(F_{1|3}(u_1|t), F_{2|3}(1|t)) \, d\lambda(t) \\ &= \int_{[0,v]} F_{1|3}(u_1|t) \, d\lambda(t) = C_{13}(u_1, v) \end{aligned}$$

for all $(u_1, v) \in \mathbb{I}^2$. The identity $(\psi(C))_{23} = C_{23}$ follows in the same manner. Assertion (2) is trivial and Assertion (3) follows from Example 4.4 below. \square

Example 4.3. (Class \mathcal{F}_{Π}^3 , cont.)

For every $C \in \mathcal{F}_{\Pi}^3$ the identity

$$(\psi(C))(\mathbf{u}, v) = C_{12}(\mathbf{u}) v$$

holds for all $(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}$.

In fact, considering that $F_{1|3}(s_1|t) = s_1$ and $F_{2|3}(s_2|t) = s_2$ hold for all $\mathbf{s} \in \mathbb{I}^2$ and almost all $t \in \mathbb{I}$ we get

$$C_p(\mathbf{s}) = \int_{\mathbb{I}} C_{12;3}^t(F_{1|3}(s_1|t), F_{2|3}(s_2|t)) d\lambda(t) = C(\mathbf{s}, 1) = C_{12}(\mathbf{s})$$

for all $\mathbf{s} \in \mathbb{I}^2$. Having this, the fact that $(\psi(C))(\mathbf{u}, v) = C_p(\mathbf{u}) v = C_{12}(\mathbf{u}) v$ holds for all $(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}$ follows immediately.

As a consequence of Example 4.3, if $C_{13} = \Pi = C_{23}$ and, additionally, $C_{12} = \Pi$, then

$$\psi(C) = \Pi$$

follows although, in general, $C \neq \Pi$. This fact applies in particular to the following copulas:

Example 4.4.

(1) The EFGM copula $C^{\text{EFGM}} \in \mathcal{F}_{\Pi}^3$ introduced in Example 3.4 is non-simplified, satisfies

$$C_{12}^{\text{EFGM}} = C_{13}^{\text{EFGM}} = C_{23}^{\text{EFGM}} = \Pi \quad \text{and} \quad C_p^{\text{EFGM}} = \Pi \quad (\text{also see [33]}),$$

and hence $\psi(C^{\text{EFGM}}) = \Pi \neq C^{\text{EFGM}}$.

(2) The copula $C^{\text{Cube}} \in \mathcal{F}_{\Pi}^3$ introduced in Example 3.4 is non-simplified, satisfies

$$C_{12}^{\text{Cube}} = C_{13}^{\text{Cube}} = C_{23}^{\text{Cube}} = \Pi \quad \text{and} \quad C_p^{\text{Cube}} = \Pi,$$

and hence $\psi(C^{\text{Cube}}) = \Pi \neq C^{\text{Cube}}$.

(3) The copula $C^{\text{RCube}} \in \mathcal{F}_{\Pi}^3$ whose mass is distributed uniformly within the cubes

$$\begin{aligned} & \left(0, \frac{1}{2}\right) \times \left(\frac{1}{2}, 1\right) \times \left(0, \frac{1}{2}\right) && \left(\frac{1}{2}, 1\right) \times \left(0, \frac{1}{2}\right) \times \left(0, \frac{1}{2}\right) \\ & \left(0, \frac{1}{2}\right) \times \left(0, \frac{1}{2}\right) \times \left(\frac{1}{2}, 1\right) && \left(\frac{1}{2}, 1\right) \times \left(\frac{1}{2}, 1\right) \times \left(\frac{1}{2}, 1\right) \end{aligned}$$

and has no mass outside these cubes is non-simplified, satisfies

$$C_{12}^{\text{RCube}} = C_{13}^{\text{RCube}} = C_{23}^{\text{RCube}} = \Pi \quad \text{and} \quad C_p^{\text{RCube}} = \Pi,$$

and hence $\psi(C^{\text{RCube}}) = \Pi \neq C^{\text{RCube}}$.

The copula in (3) is denoted as 'RCube' since it is a reflected version of the copula in (2); both are related to each other via $\mu_{C^{\text{RCube}}} = (\mu_{C^{\text{Cube}}})^T$ where $T : \mathbb{I}^2 \times \mathbb{I} \rightarrow \mathbb{I}^2 \times \mathbb{I}$ is the mapping given by $T(\mathbf{u}, v) := (\mathbf{u}, 1 - v)$ and $(\mu_{C^{\text{Cube}}})^T$ denotes the push-forward of $\mu_{C^{\text{Cube}}}$ via T .

PVCs have been used in [24] to test the simplifying assumption in vine copula models and in [27] to construct a non-parametric estimator for multivariate distributions. In [34] the authors showed that “under regularity conditions, stepwise estimators of pair-copula constructions converge to the PVC irrespective of whether the simplifying assumption holds or not” (see [34, Corollary 6.1]). Nevertheless, this does not need to be true if the estimation is done jointly in a non-simplified setting (see [34, Corollary 6.1]). The authors further proved that “if one sequentially minimizes the Kullback-Leibler divergence related to each tree then the optimal SVC is the PVC” (see [34, Theorem 5.1]). Since, again, this is not necessarily true if the estimation is done jointly in a non-simplified setting (see [34, Theorem 5.2]) the authors conclude that PVCs “may not be the best approximation in the space of SVCs” but are “often the best feasible SVC approximation in practice.”

Motivated by these results in what follows we discuss analytic properties and optimality of simplified pair-copula constructions and focus mainly on partial vine copulas. In Section 5 we calculate the d_∞ -distance between non-simplified copulas and their unique partial vine copulas for different dependence structures, in Section 6 we discuss continuity of ψ with respect to different notions of convergence.

5. OPTIMALITY OF PARTIAL VINE COPULAS

Main objective of this section is to provide an answer to the question “how far off can we be by assuming a simplified model?” posed by Nagler and Czado [27]. We proceed as follows: We first show that partial vine copulas are never the best simplified copula approximation (with respect to d_∞) if the true copula is non-simplified (Theorem 5.1). We then compare non-simplified copulas C with their unique partial vine copulas $\psi(C)$ in different settings and calculate their d_∞ -distance. It turns out that the maximal distance within the family of all copulas with pairwise independent marginals is $1/8$ which corresponds to 50% of the diameter of this class w.r.t. d_∞ . Going even further, we provide an example of a copula $C \in \mathcal{C}^3$ fulfilling $d_\infty(C, \psi(C)) = 3/16$ which, in turn, corresponds to 28.125% of the diameter of $(\mathcal{C}^3, d_\infty)$. In other words, $\psi(C)$ can be far away from C , so working with PVCs must be done with care.

Corollary 3.7 implies that if C does not fulfill the simplifying assumption then the partial vine copula fails to be optimal with respect to d_∞ :

Theorem 5.1. *Suppose that $C \in \mathcal{C}_c^3$ is non-simplified. Then there exists some simplified copula $D \in \mathcal{C}_s^3$ satisfying $d_\infty(C, D) < d_\infty(C, \psi(C))$.*

Proof. Considering $C \in \mathcal{C}_c^3 \setminus \mathcal{C}_s^3$ we have $C \neq \psi(C)$, so setting $0 < d_\infty(C, \psi(C)) =: \varepsilon$ and using Corollary 3.7 yields the desired result. \square

As next step we calculate

$$\sup_{C \in \mathcal{F}_{\Pi}^3} d_\infty(C, \psi(C)),$$

show that the supremum is attained and then characterize all elements in \mathcal{F}_{Π}^3 attaining the maximum. Afterwards we provide a lower bound for

$$\sup_{C \in \mathcal{C}^3} d_\infty(C, \psi(C)).$$

The (dis)continuity results in Section 6 will make it clear why we can not simply use compactness of $(\mathcal{C}^3, d_\infty)$ to conclude that the supremum in the last expression is attained.

5.1. Worst case scenario for the class \mathcal{F}_{Π}^3 . The following theorem holds - notice that the set of maximizers includes the two copulas C^{Cube} and C^{RCube} introduced in Examples 3.4 and 4.4:

Theorem 5.2. *For every copula $C \in \mathcal{F}_{\Pi}^3$ the inequality $d_\infty(C, \psi(C)) \leq \frac{1}{8}$ holds. Moreover, for every $C \in \mathcal{F}_{\Pi}^3$ the following two conditions are equivalent:*

- (a) $d_\infty(C, \psi(C)) = \frac{1}{8}$.
- (b) C satisfies either

$$\begin{aligned} \mu_C \left[\left(0, \frac{1}{2}\right) \times \left(0, \frac{1}{2}\right) \times \left(0, \frac{1}{2}\right) \right] &= \frac{1}{4} = \mu_C \left[\left(0, \frac{1}{2}\right) \times \left(\frac{1}{2}, 1\right) \times \left(\frac{1}{2}, 1\right) \right] \\ \mu_C \left[\left(\frac{1}{2}, 1\right) \times \left(\frac{1}{2}, 1\right) \times \left(0, \frac{1}{2}\right) \right] &= \frac{1}{4} = \mu_C \left[\left(\frac{1}{2}, 1\right) \times \left(0, \frac{1}{2}\right) \times \left(\frac{1}{2}, 1\right) \right] \end{aligned}$$

or

$$\begin{aligned} \mu_C \left[\left(0, \frac{1}{2}\right) \times \left(0, \frac{1}{2}\right) \times \left(\frac{1}{2}, 1\right) \right] &= \frac{1}{4} = \mu_C \left[\left(0, \frac{1}{2}\right) \times \left(\frac{1}{2}, 1\right) \times \left(0, \frac{1}{2}\right) \right] \\ \mu_C \left[\left(\frac{1}{2}, 1\right) \times \left(\frac{1}{2}, 1\right) \times \left(\frac{1}{2}, 1\right) \right] &= \frac{1}{4} = \mu_C \left[\left(\frac{1}{2}, 1\right) \times \left(0, \frac{1}{2}\right) \times \left(0, \frac{1}{2}\right) \right]. \end{aligned}$$

Proof. Consider $C \in \mathcal{F}_{\mathbb{I}}^3$, fix $(\mathbf{u}, v) \in \mathbb{I}^2 \times (0, 1)$ and set

$$k := \frac{1}{v} \int_{[0,v]} C_{12;3}^t(\mathbf{u}) \, d\lambda(t) \quad \text{and} \quad l := \frac{1}{1-v} \int_{[v,1]} C_{12;3}^t(\mathbf{u}) \, d\lambda(t).$$

Then

$$C_{12}(\mathbf{u}) = \int_{\mathbb{I}} C_{12;3}^t(\mathbf{u}) \, d\lambda(t) = kv + l(1-v)$$

and

$$C(\mathbf{u}, v) = \int_{[0,v]} C_{12;3}^t(F_{1|3}(u_1|t), F_{2|3}(u_2|t)) \, d\lambda(t) = \int_{[0,v]} C_{12;3}^t(\mathbf{u}) \, d\lambda(t) = kv$$

Having this and using Example 4.3 yields

$$C(\mathbf{u}, v) - (\psi(C))(\mathbf{u}, v) = kv - C_{12}(\mathbf{u})v = kv - (kv + l(1-v))v = v(1-v)(k-l).$$

Since $W(\mathbf{u}) \leq k \leq M(\mathbf{u})$ as well as $W(\mathbf{u}) \leq l \leq M(\mathbf{u})$ we further have

$$|C(\mathbf{u}, v) - (\psi(C))(\mathbf{u}, v)| = v(1-v)|k-l| \leq v(1-v)d_{\infty}(M, W) \leq \frac{1}{8}$$

Considering $d_{\infty}(C^{\text{Cube}}, \psi(C^{\text{Cube}})) = C^{\text{Cube}}(\frac{1}{2}) - \Pi(\frac{1}{2}) = \frac{1}{4} - \frac{1}{8} = \frac{1}{8}$ we finally obtain

$$\frac{1}{8} = d_{\infty}(C^{\text{Cube}}, \psi(C^{\text{Cube}})) \leq \sup_{C \in \mathcal{F}_{\mathbb{I}}} d_{\infty}(C, \psi(C)) \leq \frac{1}{8}$$

which proves the first assertion.

For proving the stated equivalence we proceed as follows: First suppose that (b) holds. Considering that for $\mathbf{u} = \frac{1}{2}$ and $v = \frac{1}{2}$ we have

$$|k-l| = |2\mu_C[(0, \frac{1}{2}) \times (0, \frac{1}{2}) \times (0, \frac{1}{2})] - 2\mu_C[(0, \frac{1}{2}) \times (0, \frac{1}{2}) \times (\frac{1}{2}, 1)]| = \frac{2}{4}$$

it follows that

$$\frac{1}{8} \geq d_{\infty}(C, \psi(C)) \geq |C(\frac{1}{2}) - (\psi(C))(\frac{1}{2})| = \frac{1}{4}|k-l| = \frac{1}{4} \cdot \frac{2}{4} = \frac{1}{8}$$

so (a) holds and it remains to show that (a) implies (b). First of all notice that

$$\frac{1}{8} = d_{\infty}(C, \psi(C)) = \sup_{(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}} |v(1-v)(k-l)|$$

and that it is straightforward to show that $|k-l|$ is at most $1/2$ and that $1/2$ can only be attained by choosing $u_1 = 1/2 = u_2$ (irrespective of the value of v). In this case either $k = 1/2$ and $l = 0$ or $k = 0$ and $l = 1/2$. Thus,

$$\frac{1}{8} = d_{\infty}(C, \psi(C)) \leq \sup_{v \in \mathbb{I}} |v(1-v)| \cdot \sup_{(\mathbf{u}, v) \in \mathbb{I}^2 \times \mathbb{I}} |k-l| = \frac{1}{2} \sup_{v \in \mathbb{I}} |v(1-v)| = \frac{1}{8}$$

and $v = 1/2$. From the first part of this proof we get

$$\mu_C[(0, \frac{1}{2})^3] = C(\frac{1}{2}, \frac{1}{2}) = k \frac{1}{2} \in \{0, \frac{1}{4}\}$$

as well as

$$\begin{aligned} \mu_C[(0, \frac{1}{2})^3] + \mu_C[(0, \frac{1}{2}) \times (0, \frac{1}{2}) \times (\frac{1}{2}, 1)] &= \mu_C[(0, \frac{1}{2}) \times (0, \frac{1}{2}) \times \mathbb{I}] \\ &= C_{12}(\frac{1}{2}) \\ &= kv + l(1-v) \\ &= \frac{1}{4} \end{aligned}$$

Since for every $C \in \mathcal{F}_{\Pi}^3$ we have

$$\begin{aligned}\mu_C\left[\left(0, \frac{1}{2}\right) \times \mathbb{I} \times \left(0, \frac{1}{2}\right)\right] &= \frac{1}{4} & \mu_C\left[\mathbb{I} \times \left(0, \frac{1}{2}\right) \times \left(0, \frac{1}{2}\right)\right] &= \frac{1}{4} \\ \mu_C\left[\left(0, \frac{1}{2}\right) \times \mathbb{I} \times \left(\frac{1}{2}, 1\right)\right] &= \frac{1}{4} & \mu_C\left[\mathbb{I} \times \left(0, \frac{1}{2}\right) \times \left(\frac{1}{2}, 1\right)\right] &= \frac{1}{4} \\ \mu_C\left[\left(\frac{1}{2}, 1\right) \times \mathbb{I} \times \left(0, \frac{1}{2}\right)\right] &= \frac{1}{4} & \mu_C\left[\mathbb{I} \times \left(\frac{1}{2}, 1\right) \times \left(0, \frac{1}{2}\right)\right] &= \frac{1}{4} \\ \mu_C\left[\left(\frac{1}{2}, 1\right) \times \mathbb{I} \times \left(\frac{1}{2}, 1\right)\right] &= \frac{1}{4} & \mu_C\left[\mathbb{I} \times \left(\frac{1}{2}, 1\right) \times \left(\frac{1}{2}, 1\right)\right] &= \frac{1}{4}\end{aligned}$$

it suffices to distinguish the following two situations:

(i) If $\mu_C\left[\left(0, \frac{1}{2}\right)^3\right] = \frac{1}{4}$ then $\mu_C\left[\left(0, \frac{1}{2}\right) \times \left(0, \frac{1}{2}\right) \times \left(\frac{1}{2}, 1\right)\right] = 0$ and $C_{13} = \Pi = C_{23}$ yields

$$\begin{aligned}\mu_C\left[\left(0, \frac{1}{2}\right) \times \left(\frac{1}{2}, 1\right) \times \left(0, \frac{1}{2}\right)\right] &= 0 & \mu_C\left[\left(0, \frac{1}{2}\right) \times \left(\frac{1}{2}, 1\right) \times \left(\frac{1}{2}, 1\right)\right] &= \frac{1}{4} \\ \mu_C\left[\left(\frac{1}{2}, 1\right) \times \left(0, \frac{1}{2}\right) \times \left(0, \frac{1}{2}\right)\right] &= 0 & \mu_C\left[\left(\frac{1}{2}, 1\right) \times \left(\frac{1}{2}, 1\right) \times \left(0, \frac{1}{2}\right)\right] &= \frac{1}{4} \\ \mu_C\left[\left(\frac{1}{2}, 1\right) \times \left(\frac{1}{2}, 1\right) \times \left(\frac{1}{2}, 1\right)\right] &= 0 & \mu_C\left[\left(\frac{1}{2}, 1\right) \times \left(0, \frac{1}{2}\right) \times \left(\frac{1}{2}, 1\right)\right] &= \frac{1}{4}\end{aligned}$$

(ii) If $\mu_C\left[\left(0, \frac{1}{2}\right)^3\right] = 0$, then $\mu_C\left[\left(0, \frac{1}{2}\right) \times \left(0, \frac{1}{2}\right) \times \left(\frac{1}{2}, 1\right)\right] = \frac{1}{4}$ and $C_{13} = \Pi = C_{23}$ yields

$$\begin{aligned}\mu_C\left[\left(0, \frac{1}{2}\right) \times \left(\frac{1}{2}, 1\right) \times \left(0, \frac{1}{2}\right)\right] &= \frac{1}{4} & \mu_C\left[\left(0, \frac{1}{2}\right) \times \left(\frac{1}{2}, 1\right) \times \left(\frac{1}{2}, 1\right)\right] &= 0 \\ \mu_C\left[\left(\frac{1}{2}, 1\right) \times \left(0, \frac{1}{2}\right) \times \left(0, \frac{1}{2}\right)\right] &= \frac{1}{4} & \mu_C\left[\left(\frac{1}{2}, 1\right) \times \left(\frac{1}{2}, 1\right) \times \left(0, \frac{1}{2}\right)\right] &= 0 \\ \mu_C\left[\left(\frac{1}{2}, 1\right) \times \left(\frac{1}{2}, 1\right) \times \left(\frac{1}{2}, 1\right)\right] &= \frac{1}{4} & \mu_C\left[\left(\frac{1}{2}, 1\right) \times \left(0, \frac{1}{2}\right) \times \left(\frac{1}{2}, 1\right)\right] &= 0,\end{aligned}$$

which completes the proof. \square

Notice that Theorem 5.2 implies the following striking property: The maximal distance of a copula C with pairwise independent marginals and its partial vine copula $\psi(C)$ corresponds to

- 50% of the diameter of the metric space of all copulas with pairwise independent marginals w.r.t. d_∞ ; the diameter of this class equals $1/4$ which can be calculated via [28, Section 3.3].
- 18.75% of the diameter of $(\mathcal{C}^3, d_\infty)$, which is given by $2/3$.

Remark 5.3. An equally striking result can be shown for the metric D_1 : Again working with C^{Cube} it follows that

$$\sup_{C \in \mathcal{F}_{\Pi}^3} D_1(C, \psi(C)) \geq \frac{15}{64} = D_1(C^{\text{Cube}}, \psi(C^{\text{Cube}}))$$

holds. Using the results in [11] we therefore get that the maximal D_1 -distance of a copula $C \in \mathcal{F}_{\Pi}^3$ and its partial vine copula $\psi(C)$ is greater than or equal to 42.1875% of the diameter of the metric space (\mathcal{C}^3, D_1) ; the diameter of this class is at most $5/9$ which can be calculated via [11, Lemma 2].

Remark 5.4. At this point it is worth to mention that C^{Cube} is exchangeable and hence approximating C^{Cube} by $\psi(C^{\text{Cube}})$ leads to equally poor results no matter which coordinate is chosen for the conditioning.

5.2. Worst case scenario for the full class \mathcal{C}_c^3 . We are now going to show that the maximal d_∞ -distance of a copula $C \in \mathcal{C}_c^3$ and its assigned partial vine copula $\psi(C)$ is at least $3/16$ which corresponds to 28.125% of the diameter of the metric space $(\mathcal{C}_c^3, d_\infty)$.

Example 5.5. Consider the intervals $I_i := \left(\frac{i-1}{4}, \frac{i}{4}\right)$ for $i \in \{1, \dots, 4\}$. We use Equation (3.1) in order to construct a three-dimensional non-simplified copula C satisfying that its

conditional copulas $C_{12;3}^t$, $t \in \mathbb{I}$, are identical for all t within each of the four subintervals. To this end, set

$$A^t := \sum_{i=1}^4 D^i \mathbb{1}_{I_i}(t) + \Pi \mathbb{1}_{\{0, \frac{1}{4}, \frac{1}{2}, \frac{3}{4}, 1\}}(t)$$

where the bivariate copulas D^1, \dots, D^4 are the shuffles of W depicted in Figure 2 (for the definition of shuffles we refer to [9, Definition 2.1] and [12, Section 5]).

As next step we construct the (conditional) univariate marginal distribution functions




FIGURE 2. Shuffles D^1, D^2, D^3, D^4 of W as considered in Example 5.5.




FIGURE 3. Densities of the checkerboard copulas B^* (left panel) and B^{**} (right panel).

$F_{1|3}(\cdot|t)$ and $F_{2|3}(\cdot|t)$ (conditional on $t \in \mathbb{I}$) and proceed as follows: Let B^*, B^{**} denote bivariate checkerboard copulas (see [11] for a definition) whose densities $b^*, b^{**} : \mathbb{I}^2 \rightarrow \mathbb{R}$ are given by

$$b^*(u_1, t) := \begin{cases} \frac{1}{2} & (u_1, t) \in (0, \frac{1}{2}) \times I_1 \cup (\frac{1}{2}, 1) \times I_4 \\ 1 & (u_1, t) \in \mathbb{I} \times I_2 \cup \mathbb{I} \times I_3 \\ \frac{3}{2} & (u_1, t) \in (\frac{1}{2}, 1) \times I_1 \cup (0, \frac{1}{2}) \times I_4 \\ 0 & \text{otherwise} \end{cases}$$

and

$$b^{**}(u_2, t) := \begin{cases} \frac{1}{2} & (u_2, t) \in (0, \frac{1}{2}) \times I_2 \cup (\frac{1}{2}, 1) \times I_3 \\ 1 & (u_2, t) \in \mathbb{I} \times I_1 \cup \mathbb{I} \times I_4 \\ \frac{3}{2} & (u_2, t) \in (\frac{1}{2}, 1) \times I_2 \cup (0, \frac{1}{2}) \times I_3 \\ 0 & \text{otherwise} \end{cases},$$

respectively (see Figure 3). Then the Markov kernels of B^* and B^{**} obviously satisfy

$$K_{B^*}(t, [0, 0.5]) = \begin{cases} \frac{1}{4} & t \in I_1 \\ \frac{2}{4} & t \in I_2 \\ \frac{2}{4} & t \in I_3 \\ \frac{3}{4} & t \in I_4 \end{cases} \quad \text{and} \quad K_{B^{**}}(t, [0, 0.5]) = \begin{cases} \frac{2}{4} & t \in I_1 \\ \frac{1}{4} & t \in I_2 \\ \frac{3}{4} & t \in I_3 \\ \frac{2}{4} & t \in I_4. \end{cases}$$

Completing the construction of C we use the copulas A^t , $t \in \mathbb{I}$, as conditional copulas and the Markov kernels K_{B^*} and $K_{B^{**}}$ as (conditional) univariate marginal distribution functions, and set

$$(5.1) \quad C(\mathbf{u}, v) := \int_{[0, v]} A^t \left(K_{B^*}(t, [0, u_1]), K_{B^{**}}(t, [0, u_2]) \right) d\lambda(t).$$

Then $C \in \mathcal{C}_c^3$ is non-simplified, satisfies $C_{12,3}^t = A^t$ for all $t \in I^1 \cup I^2 \cup I^3 \cup I^4$, $C_{13} = B^*$, $C_{23} = B^{**}$, as well as

$$\begin{aligned} C(0.5, 0.5, 1) &= \int_{\mathbb{I}} A^t \left(K_{B^*}(t, [0, 0.5]), K_{B^{**}}(t, [0, 0.5]) \right) d\lambda(t) \\ &= \frac{1}{4} D^1 \left(\frac{1}{4}, \frac{2}{4} \right) + \frac{1}{4} D^2 \left(\frac{2}{4}, \frac{1}{4} \right) + \frac{1}{4} D^3 \left(\frac{2}{4}, \frac{3}{4} \right) + \frac{1}{4} D^4 \left(\frac{3}{4}, \frac{2}{4} \right) \\ &= \frac{1}{4} \left(\frac{1}{4} + \frac{1}{4} + \frac{1}{2} + \frac{1}{2} \right) = \frac{3}{8} \end{aligned}$$

Considering that the partial copula C_p of C is given by $C_p = \frac{1}{4}(D^1 + D^2 + D^3 + D^4)$ the partial vine copula $\psi(C)$ of C satisfies

$$\begin{aligned} (\psi(C))(0.5, 0.5, 1) &= \int_{\mathbb{I}} C_p \left(K_{B^*}(t, [0, 0.5]), K_{B^{**}}(t, [0, 0.5]) \right) d\lambda(t) \\ &= \frac{1}{4} C_p \left(\frac{1}{4}, \frac{2}{4} \right) + \frac{1}{4} C_p \left(\frac{2}{4}, \frac{1}{4} \right) + \frac{1}{4} C_p \left(\frac{2}{4}, \frac{3}{4} \right) + \frac{1}{4} C_p \left(\frac{3}{4}, \frac{2}{4} \right) \\ &= \frac{1}{16} \left(D^1 \left(\frac{1}{4}, \frac{1}{2} \right) + D^2 \left(\frac{1}{4}, \frac{1}{2} \right) + D^3 \left(\frac{1}{4}, \frac{1}{2} \right) + D^4 \left(\frac{1}{4}, \frac{1}{2} \right) \right) \\ &\quad + \frac{1}{16} \left(D^1 \left(\frac{1}{2}, \frac{1}{4} \right) + D^2 \left(\frac{1}{2}, \frac{1}{4} \right) + D^3 \left(\frac{1}{2}, \frac{1}{4} \right) + D^4 \left(\frac{1}{2}, \frac{1}{4} \right) \right) \\ &\quad + \frac{1}{16} \left(D^1 \left(\frac{1}{2}, \frac{3}{4} \right) + D^2 \left(\frac{1}{2}, \frac{3}{4} \right) + D^3 \left(\frac{1}{2}, \frac{3}{4} \right) + D^4 \left(\frac{1}{2}, \frac{3}{4} \right) \right) \\ &\quad + \frac{1}{16} \left(D^1 \left(\frac{3}{4}, \frac{1}{2} \right) + D^2 \left(\frac{3}{4}, \frac{1}{2} \right) + D^3 \left(\frac{3}{4}, \frac{1}{2} \right) + D^4 \left(\frac{3}{4}, \frac{1}{2} \right) \right) \\ &= \frac{1}{16} \left(\frac{1}{4} + \frac{1}{4} + \frac{5}{4} + \frac{5}{4} \right) = \frac{3}{16}, \end{aligned}$$

from which we get $d_\infty(C, \psi(C)) \geq \frac{3}{16}$.

We have therefore proved the following theorem:

Theorem 5.6. *There exists a copula $C \in \mathcal{C}_c^3$ fulfilling $d_\infty(C, \psi(C)) \geq \frac{3}{16}$ and we have*

$$\sup_{C \in \mathcal{C}_c^3} d_\infty(C, \psi(C)) \geq \frac{3}{16}.$$

6. CONTINUITY OF ψ

In this section we discuss continuity properties of the mapping $\psi : \mathcal{C}_c^3 \rightarrow \mathcal{C}_c^3$ assigning every $C \in \mathcal{C}_c^3$ its partial vine copula. Having in mind Lemma 4.2 intuitively one might interpret ψ as projection and therefore think that ψ has to be continuous with respect to d_∞ . It turns out, however, that this interpretation is wrong, we will show that ψ is *not* continuous with respect to d_∞ . Considering stronger topologies than the one induced by d_∞ changes the picture - we will prove that ψ is continuous with respect to weak conditional convergence and with respect to the metric D_1 (under some mild regularity conditions).

6.1. Uniform convergence. The mapping ψ is not continuous with respect to d_∞ - the following result holds:

Theorem 6.1. *Suppose that $C \in \mathcal{C}_c^3$ satisfies $d_\infty(C, \psi(C)) \neq 0$. Then C is a discontinuity point of the the mapping $\psi : \mathcal{C}_c^3 \rightarrow \mathcal{C}_c^3$. In other words: Every non-simplified $C \in \mathcal{C}_c^3$ is a discontinuity point of ψ .*

Proof. Let C be as in the theorem and set $\varepsilon := d_\infty(C, \psi(C)) > 0$. Suppose that $\mathbf{X}_1, \mathbf{X}_2, \dots$ is an i.i.d. sample from $\mathbf{X} \sim C$ and let \widehat{C}_n denote the corresponding empirical copula. With probability one we have that $\mathbf{X}_1, \mathbf{X}_2, \dots$ has no ties and that $(\widehat{C}_n)_{n \in \mathbb{N}}$ converges to C with respect to d_∞ . Considering that empirical copulas are simplified according to Theorem 3.6 and using the triangle inequality it follows immediately that

$$\begin{aligned} \varepsilon &= d_\infty(C, \psi(C)) \\ &\leq d_\infty(C, \psi(\widehat{C}_n)) + d_\infty(\psi(\widehat{C}_n), \psi(C)) \\ &= d_\infty(C, \widehat{C}_n) + d_\infty(\psi(\widehat{C}_n), \psi(C)) \end{aligned}$$

holds for every $n \in \mathbb{N}$. Consequently, since $\lim_{n \rightarrow \infty} d_\infty(\widehat{C}_n, C) = 0$

$$\liminf_{n \rightarrow \infty} d_\infty(\psi(\widehat{C}_n), \psi(C)) \geq \varepsilon$$

follows, implying that ψ is not continuous at C . □

Using convex combinations (of empirical copulas with a non-simplified copula) it is straightforward to verify that the set of all $C \in \mathcal{C}_c^3$ that are non-simplified is dense in $(\mathcal{C}_c^3, d_\infty)$ - Theorem 6.1 therefore has the following corollary:

Corollary 6.2. *The mapping $\psi : \mathcal{C}_c^3 \rightarrow \mathcal{C}_c^3$ is discontinuous on a dense subset of $(\mathcal{C}_c^3, d_\infty)$.*

6.2. Weak conditional convergence. Focusing on weak conditional convergence the mapping ψ behaves more nicely:

Theorem 6.3. *Suppose that C, C_1, C_2, \dots are copulas in \mathcal{C}_c^3 . Then the following assertions hold:*

- (1) $C_n \xrightarrow{wcc} C$ implies $(C_n)_{13} \xrightarrow{wcc} C_{13}$ and $(C_n)_{23} \xrightarrow{wcc} C_{23}$.
- (2) $C_n \xrightarrow{wcc} C$ implies $(C_n)_p \xrightarrow{d_\infty} C_p$.
- (3) $C_n \xrightarrow{wcc} C$ implies $\psi(C_n) \xrightarrow{wcc} \psi(C)$.

Proof. The first assertions follows from Theorem 2.2. To prove the second one we proceed as follows: Since for almost all $v \in \mathbb{I}$ the marginal distribution functions of $K_{C_n}(v, \cdot)$, $n \in \mathbb{N}$, and of $K_C(v, \cdot)$ are continuous, Lemma A.2 implies uniform convergence of the sequence $((C_n)_{12;3}^v)_{n \in \mathbb{N}}$ to $C_{12;3}^v$. For $\mathbf{s} \in \mathbb{I}^2$ we get

$$\begin{aligned} |(C_n)_p(\mathbf{s}) - C_p(\mathbf{s})| &= \left| \int_{\mathbb{I}} (C_n)_{12;3}^t(\mathbf{s}) - C_{12;3}^t(\mathbf{s}) \, d\lambda(t) \right| \\ &\leq \int_{\mathbb{I}} |(C_n)_{12;3}^t(\mathbf{s}) - C_{12;3}^t(\mathbf{s})| \, d\lambda(t) \\ &\leq \int_{\mathbb{I}} d_\infty((C_n)_{12;3}^t, C_{12;3}^t) \, d\lambda(t) \end{aligned}$$

and dominated convergence yields

$$\lim_{n \rightarrow \infty} d_\infty((C_n)_p, C_p) = 0$$

To prove the last assertion notice that for almost all $t \in \mathbb{I}$ we have $(\psi(C))_{12;3}^t = C_p$ as well as $(\psi(C_n))_{12;3}^t = (C_n)_p$ for every $n \in \mathbb{N}$. Hence, using the second assertion it follows that

$$\lim_{n \rightarrow \infty} d_\infty((\psi(C_n))_{12;3}^t, (\psi(C))_{12;3}^t) = \lim_{n \rightarrow \infty} d_\infty((C_n)_p, C_p) = 0$$

holds for almost all $t \in \mathbb{I}$. According to Lemma A.2 it now suffices to show that the marginal distribution functions of the Markov kernels converge weakly, which is, however, an immediate consequence of the fact that $(\psi(C))_{i3} = C_{i3}$ and $(\psi(C_n))_{i3} = (C_n)_{i3}$, $i \in \{1, 2\}$ holds for every $n \in \mathbb{N}$ (see Lemma 4.2). \square

6.3. Convergence with respect to D_1 . We finally discuss D_1 -continuity. Similar to the proof of Theorem 6.3, we first relate D_1 -convergence of copulas to uniform convergence of the corresponding partial copulas. The slightly technical (but straightforward) proof of the following useful lemma is deferred to the appendix:

Lemma 6.4. *Suppose that C, C_1, C_2, \dots are copulas in \mathcal{C}_c^3 . Then the following assertions hold:*

- (1) $C_n \xrightarrow{D_1} C$ implies $(C_n)_{13} \xrightarrow{D_1} C_{13}$ and $(C_n)_{23} \xrightarrow{D_1} C_{23}$.
- (2) $(C_n)_p \xrightarrow{d_\infty} C_p$, $(C_n)_{13} \xrightarrow{D_1} C_{13}$ and $(C_n)_{23} \xrightarrow{D_1} C_{23}$ imply $\psi(C_n) \xrightarrow{D_1} \psi(C)$.

We now show D_1 -continuity of the mapping ψ on the subclass of absolutely continuous copulas satisfying some integrability condition. The following lemma whose proof is deferred to the appendix will be key for proving this result:

Lemma 6.5. *Suppose that C, C_1, C_2, \dots are copulas in \mathcal{C}_c^3 , that C is absolutely continuous and let c_{13}, c_{23} denote the densities of the marginal copulas C_{13}, C_{23} of C . If there exist some constants $p_{13}, p_{23}, p_{123} \in (1, \infty)$ such that*

$$\|c_{13}\|_{p_{13}} < \infty, \quad \|c_{23}\|_{p_{23}} < \infty, \quad \|c_{13} c_{23}\|_{p_{123}} < \infty$$

holds then $C_n \xrightarrow{D_1} C$ implies $(C_n)_p \xrightarrow{d_\infty} C_p$.

Combining the previous two lemmata yields continuity of ψ with respect to D_1 under some mild regularity conditions:

Theorem 6.6. Consider a sequence of copulas $(C_n)_{n \in \mathbb{N}}$ in \mathcal{C}_c^3 and an absolutely continuous copula $C \in \mathcal{C}_c^3$, and let c_{13}, c_{23} denote the densities of the marginal copulas C_{13}, C_{23} of C , respectively. If there exist some constants $p_{13}, p_{23}, p_{123} \in (1, \infty)$ such that

$$\|c_{13}\|_{p_{13}} < \infty, \quad \|c_{23}\|_{p_{23}} < \infty, \quad \|c_{13} c_{23}\|_{p_{123}} < \infty$$

holds then $C_n \xrightarrow{D_1} C$ implies $\psi(C_n) \xrightarrow{D_1} \psi(C)$.

7. RESULTS FOR ARBITRARY DIMENSION

To confirm that the case of dimension three is similar to higher dimension in this section we extend (slightly modified versions of) our main results (Theorem 3.6, Corollary 3.7, Theorem 5.1, Theorem 5.6, Theorem 6.1 and Corollary 6.2) to arbitrary dimensions.

7.1. Simplified copulas. Using disintegration for every copula $C \in \mathcal{C}^d$, every $J \subseteq \{1, \dots, d\}$ with $2 \leq |J| \leq d$ and every $L \subseteq J$ with $1 \leq |L| \leq |J| - 2$, there exists some Markov kernel K_{C_J} such that the lower dimensional marginal copula C_J of C corresponding to the indices of the coordinates of C belonging to J can be expressed as

$$C_J(\mathbf{u}) = \int_{[\mathbf{0}, \mathbf{u}_L]} K_{C_J}(\mathbf{t}, [\mathbf{0}, \mathbf{u}_{J \setminus L}]) \, d\mu_{C_L}(\mathbf{t})$$

for all $\mathbf{u} \in \mathbb{I}^{|J|}$. Thereby $\mathbf{u}_L \in \mathbb{I}^{|L|}$ denotes the vector of coordinates of \mathbf{u} belonging to L , and $\mathbf{u}_{J \setminus L} \in \mathbb{I}^{|J \setminus L|}$ the vector of coordinates of \mathbf{u} belonging to $J \setminus L$. Since K_{C_J} is a Markov kernel, for every $\mathbf{u}_{J \setminus L} \in \mathbb{I}^{|J \setminus L|}$ the mapping $\mathbf{t} \mapsto K_{C_J}(\mathbf{t}, [\mathbf{0}, \mathbf{u}_{J \setminus L}])$ is measurable and, for μ_{C_L} -almost every $\mathbf{t} \in \mathbb{I}^{|L|}$, the mapping $\mathbf{u}_{J \setminus L} \mapsto K_{C_J}(\mathbf{t}, [\mathbf{0}, \mathbf{u}_{J \setminus L}])$ is a multivariate distribution function with (*conditional*) univariate marginal distribution functions $F_{j|L}(\cdot | \mathbf{t})$, $j \in J \setminus L$, (conditional on \mathbf{t}). By Sklar's theorem we get that for almost every $\mathbf{t} \in \mathbb{I}^{|L|}$ there exists some (*conditional*) copula $C_{J \setminus L; L}^{\mathbf{t}}$ (conditional on \mathbf{t}) satisfying

$$K_{C_J}(\mathbf{t}, [\mathbf{0}, \mathbf{u}_{J \setminus L}]) = C_{J \setminus L; L}^{\mathbf{t}}(F_{j_1|L}(u_{j_1} | \mathbf{t}), \dots, F_{j_{|J \setminus L|}|L}(u_{j_{|J \setminus L|}} | \mathbf{t}))$$

for all $\mathbf{u}_{J \setminus L} = (u_{j_1}, \dots, u_{j_{|J \setminus L|}}) \in \mathbb{I}^{|J \setminus L|}$ such that the identity

$$C_J(\mathbf{u}) = \int_{[\mathbf{0}, \mathbf{u}_L]} C_{J \setminus L; L}^{\mathbf{t}}(F_{j_1|L}(u_{j_1} | \mathbf{t}), \dots, F_{j_{|J \setminus L|}|L}(u_{j_{|J \setminus L|}} | \mathbf{t})) \, d\mu_{C_L}(\mathbf{t})$$

holds for all $\mathbf{u} \in \mathbb{I}^{|J|}$.

We will refer to a copula $C \in \mathcal{C}^d$ as *universally simplified* if for every $J \subseteq \{1, \dots, d\}$ with $2 \leq |J| \leq d$ and every $L \subseteq J$ with $1 \leq |L| \leq |J| - 2$ the following properties hold:

(U1) There exists some copula $A \in \mathcal{C}^{|J \setminus L|}$ such that the identity

$$(7.1) \quad C_J(\mathbf{u}) = \int_{[\mathbf{0}, \mathbf{u}_L]} A(F_{j_1|L}(u_{j_1} | \mathbf{t}), \dots, F_{j_{|J \setminus L|}|L}(u_{j_{|J \setminus L|}} | \mathbf{t})) \, d\mu_{C_L}(\mathbf{t})$$

holds for all $\mathbf{u} \in \mathbb{I}^{|J|}$.

(U2) The (*conditional*) univariate marginal distribution functions $F_{j|L}(\cdot | \mathbf{t})$, $j \in J \setminus L$, are continuous for μ_{C_L} -almost all $\mathbf{t} \in \mathbb{I}^{|L|}$.

Notice that every universally simplified three-dimensional copula is simplified in the sense studied in the last sections but not necessarily vice versa. If $C \in \mathcal{C}^d$ is universally simplified then Sklar's theorem implies that the (*conditional*) copulas $C_{J \setminus L; L}^{\mathbf{t}}$ are unique for μ_{C_L} -almost all $\mathbf{t} \in \mathbb{I}^{|L|}$. In what follows we will let \mathcal{C}_c^d denote the family of all d -dimensional copulas having continuous (*conditional*) univariate marginal distribution functions, $\mathcal{C}_{\text{US}}^d$

will denote the family of all d -dimensional universally simplified copulas. Notice that $\Pi \in \mathcal{C}_{\text{US}}^d$ and that the collection of all absolutely continuous copulas is contained in \mathcal{C}_c^d .

As first step we now prove a sharper version of Theorem 3.6 and show that all d -variate empirical copulas (d -linear interpolations) are universally simplified.

Theorem 7.1. *Every d -dimensional empirical copula is universally simplified.*

Proof. Suppose that \mathbf{X} is a d -dimensional random vector with continuous univariate marginals and suppose that $\mathbf{X}_1, \dots, \mathbf{X}_n$ is a sample from \mathbf{X} . W.l.o.g. assume that there are no ties. Letting \hat{C}_n denote the (d -linear interpolation of the) empirical copula there exists unique permutations $\sigma_1, \dots, \sigma_{d-1}$ of $\{1, \dots, n\}$ such that the density \hat{c}_n of \hat{C}_n is given by (uniform distribution on n d -dimensional squares of volume $\frac{1}{n^d}$)

$$\hat{c}_n(\mathbf{u}) = n^{d-1} \sum_{i=1}^n \left(\prod_{j=1}^d \mathbb{1}_{I_i^j}(u_j) \right), \quad \mathbf{u} = (u_1, \dots, u_d) \in \mathbb{I}^d,$$

where $I_i^j = (\frac{\sigma_j(i)-1}{n}, \frac{\sigma_j(i)}{n}]$, $j \in \{1, \dots, d-1\}$, and $I_i^d = (\frac{i-1}{n}, \frac{i}{n}]$ for every $i \in \{1, \dots, n\}$. Since marginals of empirical copulas are empirical copulas too it suffices to prove the result for $J = \{1, \dots, d\}$ and for $1 \leq l \leq d-2$ with $L = \{d-l+1, \dots, d\}$. Considering that the l -dimensional marginal copula $(\hat{C}_n)_L$ of (\hat{C}_n) assigns full mass to the set

$$\bigcup_{i=1}^n \underbrace{\left(\prod_{j=d-l+1}^d I_i^j \right)}_{=: \Omega_i}$$

it is enough to consider $\mathbf{u}_L \in \Omega_{i_0}$ for some $i_0 \in \{1, \dots, n\}$. For such \mathbf{u}_L the Markov kernel (conditioning on the coordinates in L) is given by (straightforward consequence of first considering the conditional density)

$$K_{\hat{C}_n}(\mathbf{u}_L, [0, u_1] \times \dots \times [0, u_{d-l}]) = n^{d-l} \left(\prod_{j=1}^{d-l} \int_{[0, u_j]} \mathbb{1}_{I_{i_0}^j}(x_j) d\lambda(x_j) \right)$$

and the conditional univariate distribution functions $F_{j|L}(u_j|\mathbf{u}_L)$ for every $j \in J \setminus L$ can be expressed as

$$F_{j|L}(u_j|\mathbf{u}_L) = n \int_{[0, u_j]} \mathbb{1}_{I_{i_0}^j}(x_j) d\lambda(x_j).$$

Having this we have shown

$$K_{\hat{C}_n}(\mathbf{u}_L, [0, u_1] \times \dots \times [0, u_{d-l}]) = \Pi_{d-l}(F_{1|L}(u_1|\mathbf{u}_L), \dots, F_{d-l|L}(u_{d-l}|\mathbf{u}_L)),$$

which completes the proof. \square

Since the collection of all empirical copulas is dense in $(\mathcal{C}^d, d_\infty)$ ([9, Proposition 3.2]) Theorem 7.1 has the following immediate consequence:

Corollary 7.2. *The collection of all universally simplified d -dimensional copulas is dense in $(\mathcal{C}^d, d_\infty)$.*

Thus, every copula can be approximated arbitrarily well by universally simplified ones. Given a d -dimensional, non universally simplified copula C , a good uniform approximation by a universally simplified one from the same (Fréchet) class might, however, not exist. The next example illustrates this fact:

Example 7.3. (Family $\mathcal{F}_{\text{Ind}}^d$ of all copulas C satisfying $C(\mathbf{u}, \mathbf{v}) = B(\mathbf{u})\Pi(\mathbf{v})$ for all $(\mathbf{u}, \mathbf{v}) \in \mathbb{I}^3 \times \mathbb{I}^{d-3}$ and some $B \in \mathcal{C}_c^3$ fulfilling $B_{12} = B_{13} = B_{23} = \Pi$.)

First, notice that for every universally simplified copula D in $\mathcal{F}_{\text{Ind}}^d$ there exists some copula $A \in \mathcal{C}^2$ such that, according to Equation (7.1) ($J = \{1, \dots, d\}$ and $L = \{3, \dots, d\}$), the identity

$$\begin{aligned} D(\mathbf{u}, \mathbf{v}) &= \int_{[\mathbf{0}, \mathbf{v}]} A(F_{1|L}(u_1|\mathbf{t}), F_{2|L}(u_2|\mathbf{t})) \, d\mu_{C_L}(\mathbf{t}) \\ &= \int_{[\mathbf{0}, \mathbf{v}]} A(u_1, u_2) \, d\lambda^{d-2}(\mathbf{t}) \\ &= A(\mathbf{u})\Pi(\mathbf{v}) \end{aligned}$$

holds for all $(\mathbf{u}, \mathbf{v}) \in \mathbb{I}^2 \times \mathbb{I}^{d-2}$. Notice that the second equality holds since in case of $\mathbf{X} \sim D$ we have that (X_1, X_3) and (X_4, \dots, X_d) are independent, hence

$$\begin{aligned} \mathbb{P}(X_1 \leq u_1, X_3 \leq u_3, \dots, X_d \leq u_d) &= \mathbb{P}(X_1 \leq u_1, X_3 \leq u_3) \\ &\quad \cdot \mathbb{P}(X_4 \leq u_4, \dots, X_d \leq u_d) \\ &= \mathbb{P}(X_1 \leq u_1) \mathbb{P}(X_3 \leq u_3, \dots, X_d \leq u_d) \end{aligned}$$

for all $\mathbf{u} \in (0, 1)^d$, so X_1 and (X_3, \dots, X_d) are independent and we get $F_{1|L}(u_1|\mathbf{t}) = u_1$ (the same reasoning applies to $F_{2|L}$).

Setting $C(\mathbf{u}, \mathbf{v}) = C^{\text{Cube}}(\mathbf{u})\Pi(\mathbf{v})$ for all $(\mathbf{u}, \mathbf{v}) \in \mathbb{I}^3 \times \mathbb{I}^{d-3}$ and considering Example 3.8 it therefore follows that $d_\infty(C, D) > \varepsilon$ (with ε as in Example 3.8), i.e., it is not possible to approximate C by universally simplified copulas in $\mathcal{F}_{\text{Ind}}^d$ with an error smaller than ε .

7.2. Partial vine copulas (PVC-D). We finally introduce *partial vine copulas* (PVC) belonging to a D -vine structure.

Given $C \in \mathcal{C}_c^d$ the hierarchical construction of a partial vine copula C^{PVC} of D -vine structure may be build as follows:

(S1) the bivariate marginal copulas $(C^{\text{PVC}})_{i,i+1}$ are defined as

$$(C^{\text{PVC}})_{i,i+1} := C_{i,i+1}$$

for all $i \in \{1, \dots, d-1\}$.

(S2) In the second step the bivariate partial copulas $(C_p)_{i,i+2}$, $i \in \{1, \dots, d-2\}$ are defined by

$$(C_p)_{i,i+2}(u_i, u_{i+2}) := \int_{\mathbb{I}} C_{i,i+2;i+1}^{w_{i+1}}(u_i, u_{i+2}) \, d\lambda(w_{i+1})$$

and the three-dimensional marginal copulas $(C^{\text{PVC}})_{i,i+1,i+2}$ with $i \in \{1, \dots, d-2\}$, are constructed via

$$\begin{aligned} &(C^{\text{PVC}})_{i,i+1,i+2}(u_i, u_{i+1}, u_{i+2}) \\ &:= \int_{[0, u_{i+1}]} (C_p)_{i,i+2}((F^{\text{PVC}})_{i|i+1}(u_i|t_{i+1}), (F^{\text{PVC}})_{i+2|i+1}(u_{i+2}|t_{i+1})) \, d\lambda(t_{i+1}) \end{aligned}$$

where the (conditional) univariate distribution functions $(F^{\text{PVC}})_{i|i+1}$ and $(F^{\text{PVC}})_{i+2|i+1}$ correspond to the copulas $(C^{\text{PVC}})_{i,i+1}$ and $(C^{\text{PVC}})_{i+1,i+2}$ from previous steps.

(S3) The bivariate partial copulas $(C_p)_{i,i+3}$ with $i \in \{1, \dots, d-3\}$ are defined by

$$(C_p)_{i,i+3}(u_i, u_{i+3}) := \int_{\mathbb{I}^2} C_{i,i+3;i+1,i+2}^{(w_{i+1}, w_{i+2})}(u_i, u_{i+3}) \, d\mu_{(C^{\text{PVC}})_{i+1,i+2}}(w_{i+1}, w_{i+2})$$

and the 4-dimensional marginal copulas $(C^{\text{PVC}})_{i,i+1,i+2,i+3}$ with $i \in \{1, \dots, d-3\}$ are constructed via

$$\begin{aligned} & (C^{\text{PVC}})_{i,i+1,i+2,i+3}(u_i, u_{i+1}, u_{i+2}, u_{i+3}) \\ & := \int_{[0, u_{i+1}] \times [0, u_{i+2}]} (C_p)_{i,i+3}((F^{\text{PVC}})_{i|i+1,i+2}(u_i | (t_{i+1}, t_{i+2})), (F^{\text{PVC}})_{i+3|i+1,i+2}(u_{i+3} | (t_{i+1}, t_{i+2}))) \\ & \quad d\mu_{(C^{\text{PVC}})_{i+1,i+2}}(t_{i+1}, t_{i+2}), \end{aligned}$$

where the (conditional) univariate distribution functions $(F^{\text{PVC}})_{i|i+1,i+2}$ and $(F^{\text{PVC}})_{i+3|i+1,i+2}$ correspond to the three-dimensional copulas $(C^{\text{PVC}})_{i,i+1,i+2}$ and $(C^{\text{PVC}})_{i+1,i+2,i+3}$ from previous steps.

(S4) The individual steps are continued until one obtains the d -dimensional partial vine copula C^{PVC} of D -vine structure.

The mapping induced by the afore-mentioned procedure will be denoted by $\psi : \mathcal{C}_c^d \rightarrow \mathcal{C}_c^d$, by construction it fulfills

$$\psi(C) = C^{\text{PVC}}.$$

Notice that, by definition, C^{PVC} is simplified with respect to the underlying D -vine structure but may fail to be universally simplified.

Example 7.4. We calculate $\psi(C)$ for the d -dimensional copula $C \in \mathcal{C}_c^d$ given by

$$C(\mathbf{u}, \mathbf{v}) := C^{\text{Cube}}(\mathbf{u}) \Pi(\mathbf{v})$$

for all $(\mathbf{u}, \mathbf{v}) \in \mathbb{I}^3 \times \mathbb{I}^{d-3}$ (see Example 7.3) and show that $\psi(C) = \Pi$ holds.

We start with the following observations:

- C satisfies $C_J = \Pi$ for all $J \subseteq \{1, \dots, d\}$ with $2 \leq |J| \leq d-1$ and $\{1, 2, 3\} \not\subseteq J$.
- C is absolutely continuous.
- the Markov kernel of $C_{\{1, \dots, j\}}$, $j \in \{4, \dots, d\}$, with respect to the coordinates $2, \dots, j-1$ satisfies

$$\begin{aligned} & K_{C_{\{1, \dots, j\}}}((t_2, \dots, t_{j-1}), [0, u_1] \times [0, u_j]) \\ & = \begin{cases} \min(2u_1, 1) u_j & \text{if } (t_2, t_3) \in (0, \frac{1}{2})^2 \cup (\frac{1}{2}, 1)^2 \\ \max(2u_1 - 1, 0) u_j & \text{if } (t_2, t_3) \in (0, \frac{1}{2}) \times (\frac{1}{2}, 1) \cup (\frac{1}{2}, 1) \times (0, \frac{1}{2}) \end{cases} \end{aligned}$$

for almost all $(t_2, \dots, t_{j-1}) \in \mathbb{I}^{j-2}$, hence $C_{1,j;2,\dots,j-1} = \Pi$ and it follows that the partial copula $(C_p)_{1,j;2,\dots,j-1}$ coincides with Π .

We now calculate the partial vine copula $\psi(C)$ step-by-step:

(S1) In the first step we obtain

$$(C^{\text{PVC}})_{i,i+1} = \Pi$$

for all $i \in \{1, \dots, d-1\}$.

(S2) The bivariate partial copulas satisfy

$$(C_p)_{1,3}(u_1, u_3) = \int_{\mathbb{I}} C_{1,3;2}^{w_2}(u_1, u_3) \, d\lambda(w_2) = \frac{1}{2} A^1(u_1, u_3) + \frac{1}{2} A^2(u_1, u_3) = \Pi(u_1, u_3)$$

for all $(u_1, u_3) \in \mathbb{I}^2$ and for every $i \in \{2, \dots, d-2\}$ we get

$$\begin{aligned} (C_p)_{i,i+2}(u_i, u_{i+2}) &= \int_{\mathbb{I}} C_{i,i+2;i+1}^{w_{i+1}}(u_i, u_{i+2}) \, d\lambda(w_{i+1}) = \int_{\mathbb{I}} \Pi(u_i, u_{i+2}) \, d\lambda(w_{i+1}) \\ &= \Pi(u_i, u_{i+2}) \end{aligned}$$

for all $(u_i, u_{i+2}) \in \mathbb{I}^2$. Therefore, the 3-dimensional marginal copulas satisfy

$$\begin{aligned} (C^{\text{PVC}})_{i,i+1,i+2}(u_i, u_{i+1}, u_{i+2}) &= \int_{[0, u_{i+1}]} (C_p)_{i,i+2}((F^{\text{PVC}})_{i|i+1}(u_i|t_{i+1}), (F^{\text{PVC}})_{i+2|i+1}(u_{i+2}|t_{i+1})) \, d\lambda(t_{i+1}) \\ &= \int_{[0, u_{i+1}]} (C_p)_{i,i+2}(u_i, u_{i+2}) \, d\lambda(t_{i+1}) \\ &= \Pi(u_i, u_{i+1}, u_{i+2}) \end{aligned}$$

for all $(u_i, u_{i+1}, u_{i+2}) \in \mathbb{I}^3$ and every $i \in \{1, \dots, d-2\}$.

(S3) In the third step the bivariate partial copulas satisfy

$$\begin{aligned} (C_p)_{i,i+3}(u_i, u_{i+3}) &= \int_{\mathbb{I}^2} C_{i,i+3;i+1,i+2}^{(w_{i+1}, w_{i+2})}(u_i, u_{i+3}) \, d\mu_{(C^{\text{PVC}})_{i+1,i+2}}(w_{i+1}, w_{i+2}) \\ &= \int_{\mathbb{I}^2} \Pi(u_i, u_{i+3}) \, d\lambda^2(w_{i+1}, w_{i+2}) \\ &= \Pi(u_i, u_{i+3}) \end{aligned}$$

for all $(u_i, u_{i+3}) \in \mathbb{I}^2$ and for every $i \in \{1, \dots, d-3\}$. Hence follows that the 4-dimensional marginal copulas satisfy

$$\begin{aligned} (C^{\text{PVC}})_{i,i+1,i+2,i+3}(u_i, u_{i+1}, u_{i+2}, u_{i+3}) &= \int_{[0, u_{i+1}] \times [0, u_{i+2}]} (C_p)_{i,i+3}((F^{\text{PVC}})_{i|i+1,i+2}(u_i|(t_{i+1}, t_{i+2})), (F^{\text{PVC}})_{i+3|i+1,i+2}(u_{i+3}|(t_{i+1}, t_{i+2}))) \\ &\quad d\mu_{(C^{\text{PVC}})_{i+1,i+2}}(t_{i+1}, t_{i+2}) \\ &= \int_{[0, u_{i+1}] \times [0, u_{i+2}]} (C_p)_{i,i+3}(u_i, u_{i+3}) \, d\lambda^2(t_{i+1}, t_{i+2}) \\ &= \Pi(u_i, u_{i+1}, u_{i+2}, u_{i+3}) \end{aligned}$$

for all $(u_1, u_2, u_3, u_4) \in \mathbb{I}^4$.

(S4) Continuing in the same manner we finally arrive at $\psi(C) = C^{\text{PVC}} = \Pi$.

Remark 7.5. Notice that the afore-mentioned construction principle of a partial vine copula of D -vine structure differs from the one introduced in Spanhel and Kurz [34, p.1264]. It is, however, straightforward to verify that constructing the partial vine copula version according to Spanhel and Kurz for the copula C in Example 7.4 yields the same output Π .

7.3. Optimality of partial vine copulas and continuity of δ . Example 7.4 allows to prove the following multivariate version of Theorem 5.6:

Theorem 7.6. *For every $d \geq 3$ there exists a copula $C \in \mathcal{C}_c^d$ fulfilling $d_\infty(C, \psi(C)) \geq \frac{1}{8}$ and we have*

$$\sup_{C \in \mathcal{C}_c^d} d_\infty(C, \psi(C)) \geq \frac{1}{8}.$$

Proof. Again consider the d -dimensional copula $C \in \mathcal{C}_c^d$ studied in Example 7.4. In this case we have $\psi(C) = \Pi$ from which we get

$$\begin{aligned} d_\infty(C, \psi(C)) &\geq C\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \mathbf{1}\right) - \Pi\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}, \mathbf{1}\right) \\ &= C^{\text{Cube}}\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right) - \Pi\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right) = \frac{1}{4} - \frac{1}{8} = \frac{1}{8}, \end{aligned}$$

which implies the stated result. \square

We conclude the paper with the multivariate versions of Theorem 6.1 and Corollary 6.2:

Theorem 7.7. *Suppose that $C \in \mathcal{C}_c^d$ satisfies $d_\infty(C, \psi(C)) \neq 0$. Then C is a discontinuity point of the mapping $\psi : \mathcal{C}_c^d \rightarrow \mathcal{C}_c^d$ assigning every copula its partial D -vine.*

Proof. Proceeding analogous to the proof of Theorem 6.1 and using the fact that empirical copulas are invariant under ψ and converge to the true copula with respect to d_∞ yields the result. \square

Again using convex combinations it is straightforward to verify that the set of all $C \in \mathcal{C}_c^d$ that are not universally simplified is dense in $(\mathcal{C}_c^d, d_\infty)$ - Theorem 7.7 has the following consequence:

Corollary 7.8. *The mapping $\psi : \mathcal{C}_c^d \rightarrow \mathcal{C}_c^d$ is discontinuous on a dense subset of $(\mathcal{C}_c^d, d_\infty)$.*

Remark 7.9. As a consequence of Remark 7.5, all the results presented in Subsection 7.3 (i.e., Theorems 7.6, 7.7 and Corollary 7.8) remain true for the partial vine copula of D -vine structure as discussed in Spanhel and Kurz [34, p.1264]. It is worth mentioning that, although the latter construction principle sequentially minimizes the Kullback-Leibler divergence related to each tree, its outcome can be quite far away from the data generating copula (see Theorem 7.6) which is in line with Theorem 3.10.

APPENDIX A. SUPPLEMENTARY MATERIAL

Lemma A.1.

- (1) *Suppose that F, F_1, F_2, \dots are univariate distribution functions and suppose that F is continuous. Then weak convergence $F_n \rightarrow F$ implies uniform convergence.*
- (2) *Suppose that F, F_1, F_2, \dots are d -dimensional distribution functions ($d \geq 2$) and suppose that F is continuous. Then weak convergence $F_n \rightarrow F$ implies uniform convergence.*

Proof. Since the first statement is well-known and straightforward to verify we focus on the second assertion. Considering that F is continuous the sequence $(F_n)_{n \in \mathbb{N}}$ converges pointwise to F and the same holds true for all univariate marginals. Using Sklar's Theorem, Lipschitz continuity of copulas and statement (1) we get

$$|F_n(\mathbf{x}) - F(\mathbf{x})| \leq \sum_{i=1}^d |(F_n)_i(x_i) - F_i(x_i)|$$

for every $\mathbf{x} \in \mathbb{R}^d$, which completes the proof. \square

Lemma A.2. *Suppose that F, F_1, F_2, \dots are d -dimensional distribution functions with continuous marginals $(F)_i, (F_1)_i, (F_2)_i, \dots$ ($i \in \{1, \dots, d\}$) and copulas C, C_1, C_2, \dots , respectively. Then the following assertions hold:*

- (1) *If $C_n \rightarrow C$ uniformly and $(F_n)_i \rightarrow (F)_i$ weakly then $F_n \rightarrow F$ uniformly.*
- (2) *If $F_n \rightarrow F$ weakly then $C_n \rightarrow C$ uniformly.*

Proof. Since the limits are continuous by assumption, according to Lemma A.1 weak and uniform convergence coincide. We start with proving the first assertion and consider some $\mathbf{x} \in \mathbb{R}^d$. Then Lipschitz continuity of copulas and the triangle inequality yield (we write $\mathbf{F}_n := ((F_n)_1, (F_n)_2, \dots, (F_n)_d)$)

$$\begin{aligned} |F_n(\mathbf{x}) - F(\mathbf{x})| &= |(C_n \circ \mathbf{F}_n)(\mathbf{x}) - (C \circ \mathbf{F})(\mathbf{x})| \\ &\leq |(C_n \circ \mathbf{F}_n)(\mathbf{x}) - (C \circ \mathbf{F}_n)(\mathbf{x})| + |(C \circ \mathbf{F}_n)(\mathbf{x}) - (C \circ \mathbf{F})(\mathbf{x})| \\ &\leq d_\infty(C_n, C) + \sum_{i=1}^d |(F_n)_i(x_i) - (F)_i(x_i)|, \end{aligned}$$

from which the first assertion follows immediately.

To prove the second assertion fix $\mathbf{u} \in \mathbb{I}^d$. Letting $(F_n)_i^{\leftarrow}$ denote the quasi-inverse of $(F_n)_i$ and letting $\mathbf{F}_n^{\leftarrow}$ accordingly denote the vector of quasi-inverses of the univariate marginals yields

$$\begin{aligned} |C_n(\mathbf{u}) - C(\mathbf{u})| &= |(F_n \circ \mathbf{F}_n^{\leftarrow})(\mathbf{u}) - (F \circ \mathbf{F}^{\leftarrow})(\mathbf{u})| \\ &\leq |(F_n \circ \mathbf{F}_n^{\leftarrow})(\mathbf{u}) - (F_n \circ \mathbf{F}^{\leftarrow})(\mathbf{u})| + |(F_n \circ \mathbf{F}^{\leftarrow})(\mathbf{u}) - (F \circ \mathbf{F}^{\leftarrow})(\mathbf{u})| \\ &\leq \sum_{i=1}^d |((F_n)_i \circ (F_n)_i^{\leftarrow})(u_i) - ((F_n)_i \circ (F)_i^{\leftarrow})(u_i)| + |F_n(\mathbf{F}^{\leftarrow}(\mathbf{u})) - F(\mathbf{F}^{\leftarrow}(\mathbf{u}))| \\ &= \sum_{i=1}^d |u_i - ((F_n)_i \circ (F)_i^{\leftarrow})(u_i)| + |F_n(\mathbf{F}^{\leftarrow}(\mathbf{u})) - F(\mathbf{F}^{\leftarrow}(\mathbf{u}))| \\ &= \sum_{i=1}^d |(F)_i((F)_i^{\leftarrow}(u_i)) - (F_n)_i((F)_i^{\leftarrow}(u_i))| + |F_n(\mathbf{F}^{\leftarrow}(\mathbf{u})) - F(\mathbf{F}^{\leftarrow}(\mathbf{u}))| \\ &\leq \sum_{i=1}^d d_\infty((F_n)_i, (F)_i) + \sup_{\mathbf{x} \in \mathbb{R}^d} |F_n(\mathbf{x}) - F(\mathbf{x})|. \end{aligned}$$

This completes the proof. □

Lemma A.3. *Suppose that $C \in \mathcal{C}_{\text{ac}}^3$ is an absolutely continuous copula, and let c_{13}, c_{23} denote the densities of the marginal copulas C_{13}, C_{23} of C , respectively. Then the following inequality holds for every $\tilde{C} \in \mathcal{C}_c^3$:*

$$\begin{aligned} &\int_{\mathbb{I}^2} \int_{\mathbb{I}} |C_{12;3}^t(\mathbf{s}) - \tilde{C}_{12;3}^t(\mathbf{s})| \, d\lambda(t) d\lambda^2(\mathbf{s}) \\ &\leq \int_{\mathbb{I}} \int_{\mathbb{I}^2} |K_C(t, [\mathbf{0}, \mathbf{s}]) - K_{\tilde{C}}(t, [\mathbf{0}, \mathbf{s}])| \left(c_{13}(s_1, t) c_{23}(s_2, t) \right) \, d\lambda^2(\mathbf{s}) d\lambda(t) \\ &\quad + \int_{\mathbb{I}} \int_{\mathbb{I}} |F_{1|3}(s_1|t) - \tilde{F}_{1|3}(s_1|t)| \, c_{13}(s_1, t) \, d\lambda(s_1) d\lambda(t) \end{aligned}$$

$$+ \int_{\mathbb{I}} \int_{\mathbb{I}} |F_{2|3}(s_2|t) - \tilde{F}_{2|3}(s_2|t)| c_{23}(s_2, t) d\lambda(s_2) d\lambda(t)$$

Proof. For $\tilde{C} \in \mathcal{C}_c^3$ and $C \in \mathcal{C}_{ac}^3$ we have

$$\begin{aligned} & \int_{\mathbb{I}^2} \int_{\mathbb{I}} |C_{12;3}^t(\mathbf{s}) - \tilde{C}_{12;3}^t(\mathbf{s})| d\lambda(t) d\lambda^2(\mathbf{s}) \\ &= \int_{\mathbb{I}^2} \int_{\mathbb{I}} \left| K_C(t, [0, F_{1|3}^{\leftarrow}(s_1|t)] \times [0, F_{2|3}^{\leftarrow}(s_2|t)]) \right. \\ & \quad \left. - K_{\tilde{C}}(t, [0, \tilde{F}_{1|3}^{\leftarrow}(s_1|t)] \times [0, \tilde{F}_{2|3}^{\leftarrow}(s_2|t)]) \right| d\lambda(t) d\lambda^2(\mathbf{s}) \\ &\leq \int_{\mathbb{I}^2} \int_{\mathbb{I}} \left| K_C(t, [0, F_{1|3}^{\leftarrow}(s_1|t)] \times [0, F_{2|3}^{\leftarrow}(s_2|t)]) \right. \\ & \quad \left. - K_{\tilde{C}}(t, [0, F_{1|3}^{\leftarrow}(s_1|t)] \times [0, F_{2|3}^{\leftarrow}(s_2|t)]) \right| d\lambda(t) d\lambda^2(\mathbf{s}) \\ & \quad + \int_{\mathbb{I}^2} \int_{\mathbb{I}} \left| K_{\tilde{C}}(t, [0, F_{1|3}^{\leftarrow}(s_1|t)] \times [0, F_{2|3}^{\leftarrow}(s_2|t)]) \right. \\ & \quad \left. - K_{\tilde{C}}(t, [0, \tilde{F}_{1|3}^{\leftarrow}(s_1|t)] \times [0, \tilde{F}_{2|3}^{\leftarrow}(s_2|t)]) \right| d\lambda(t) d\lambda^2(\mathbf{s}) \\ &=: I_1 + I_2. \end{aligned}$$

For every $t \in \mathbb{I}$ define $T^t : \mathbb{I}^2 \rightarrow \mathbb{I}^2$ by

$$T^t(\mathbf{s}) := (F_{1|3}^{\leftarrow}(s_1|t), F_{2|3}^{\leftarrow}(s_2|t)).$$

Then T^t is measurable, obviously satisfies $(T^t)^{-1}(\mathbb{I}^2) = \mathbb{I}^2$, and

$$\begin{aligned} (\lambda^2)^{T^t}([\mathbf{0}, \mathbf{u}]) &= \lambda^2(\{\mathbf{s} \in \mathbb{I}^2 : T^t(\mathbf{s}) \in [\mathbf{0}, \mathbf{u}]\}) \\ &= \lambda^2(\{\mathbf{s} \in \mathbb{I}^2 : s_1 \leq F_{1|3}(u_1|t), s_2 \leq F_{2|3}(u_2|t)\}) \\ &= F_{1|3}(u_1|t) F_{2|3}(u_2|t) \\ &= \int_{[\mathbf{0}, \mathbf{u}]} c_{13}(a_1, t) c_{23}(a_2, t) d\lambda^2(\mathbf{a}) \end{aligned}$$

for every $\mathbf{u} \in \mathbb{I}^2$, implying that $(\lambda^2)^{T^t}$ is absolutely continuous with density $(a_1, a_2) \mapsto c_{13}(a_1, t) c_{23}(a_2, t)$. This yields

$$\begin{aligned} I_1 &= \int_{\mathbb{I}} \int_{\mathbb{I}^2} \left| K_C(t, [0, F_{1|3}^{\leftarrow}(s_1|t)] \times [0, F_{2|3}^{\leftarrow}(s_2|t)]) \right. \\ & \quad \left. - K_{\tilde{C}}(t, [0, F_{1|3}^{\leftarrow}(s_1|t)] \times [0, F_{2|3}^{\leftarrow}(s_2|t)]) \right| d\lambda^2(\mathbf{s}) d\lambda(t) \\ &= \int_{\mathbb{I}} \int_{\mathbb{I}^2} |K_C(t, [\mathbf{0}, \mathbf{s}]) - K_{\tilde{C}}(t, [\mathbf{0}, \mathbf{s}])| d(\lambda^2)^{T^t}(\mathbf{s}) d\lambda(t) \\ &= \int_{\mathbb{I}} \int_{\mathbb{I}^2} |K_C(t, [\mathbf{0}, \mathbf{s}]) - K_{\tilde{C}}(t, [\mathbf{0}, \mathbf{s}])| (c_{13}(s_1, t) c_{23}(s_2, t)) d\lambda^2(\mathbf{s}) d\lambda(t). \end{aligned}$$

Focusing on I_2 , using Sklar's theorem, Lipschitz continuity, and a similar argument as before yields

$$\begin{aligned}
I_2 &= \int_{\mathbb{I}} \int_{\mathbb{I}^2} \left| K_{\tilde{C}}(t, [0, F_{1|3}^{\leftarrow}(s_1|t)] \times [0, F_{2|3}^{\leftarrow}(s_2|t)]) \right. \\
&\quad \left. - K_{\tilde{C}}(t, [0, \tilde{F}_{1|3}^{\leftarrow}(s_1|t)] \times [0, \tilde{F}_{2|3}^{\leftarrow}(s_2|t)]) \right| d\lambda^2(\mathbf{s})d\lambda(t) \\
&\leq \int_{\mathbb{I}} \int_{\mathbb{I}} |\tilde{F}_{1|3}(F_{1|3}^{\leftarrow}(s_1|t)|t) - \tilde{F}_{1|3}(\tilde{F}_{1|3}^{\leftarrow}(s_1|t)|t)| d\lambda(s_1)d\lambda(t) \\
&\quad + \int_{\mathbb{I}} \int_{\mathbb{I}} |\tilde{F}_{2|3}(F_{2|3}^{\leftarrow}(s_2|t)|t) - \tilde{F}_{2|3}(\tilde{F}_{2|3}^{\leftarrow}(s_2|t)|t)| d\lambda(s_2)d\lambda(t) \\
&= \int_{\mathbb{I}} \int_{\mathbb{I}} |\tilde{F}_{1|3}(F_{1|3}^{\leftarrow}(s_1|t)|t) - s_1| d\lambda(s_1)d\lambda(t) \\
&\quad + \int_{\mathbb{I}} \int_{\mathbb{I}} |\tilde{F}_{2|3}(F_{2|3}^{\leftarrow}(s_2|t)|t) - s_2| d\lambda(s_2)d\lambda(t) \\
&= \int_{\mathbb{I}} \int_{\mathbb{I}} |\tilde{F}_{1|3}(F_{1|3}^{\leftarrow}(s_1|t)|t) - F_{1|3}(F_{1|3}^{\leftarrow}(s_1|t)|t)| d\lambda(s_1)d\lambda(t) \\
&\quad + \int_{\mathbb{I}} \int_{\mathbb{I}} |\tilde{F}_{2|3}(F_{2|3}^{\leftarrow}(s_2|t)|t) - F_{2|3}(F_{2|3}^{\leftarrow}(s_2|t)|t)| d\lambda(s_2)d\lambda(t) \\
&= \int_{\mathbb{I}} \int_{\mathbb{I}} |\tilde{F}_{1|3}(s_1|t) - F_{1|3}(s_1|t)| c_{13}(s_1, t) d\lambda(s_1)d\lambda(t) \\
&\quad + \int_{\mathbb{I}} \int_{\mathbb{I}} |\tilde{F}_{2|3}(s_2|t) - F_{2|3}(s_2|t)| c_{23}(s_2, t) d\lambda(s_2)d\lambda(t),
\end{aligned}$$

and the proof is complete. \square

As a direct consequence of Lemma A.3 we obtain the following result:

Lemma A.4. *Suppose that $C \in \mathcal{C}_{\text{ac}}^3$ is an absolutely continuous copula whose density c fulfills $c \leq a \in [1, \infty)$. Then the inequality*

$$J(C, \tilde{C}) := \int_{\mathbb{I}^3} |C_{12;3}^t(\mathbf{s}) - \tilde{C}_{12;3}^t(\mathbf{s})| d\lambda^3(\mathbf{s}, t) \leq a(2 + a)D_{\infty}(C, \tilde{C})$$

holds for every $\tilde{C} \in \mathcal{C}_c^3$.

Proof. Applying Lemma A.3 and Theorem 2.3 yields

$$\begin{aligned}
&\int_{\mathbb{I}^3} |C_{12;3}^t(\mathbf{s}) - \tilde{C}_{12;3}^t(\mathbf{s})| d\lambda^3(\mathbf{s}, t) \\
&\leq \int_{\mathbb{I}} \int_{\mathbb{I}^2} \left| K_C(t, [\mathbf{0}, \mathbf{s}]) - K_{\tilde{C}}(t, [\mathbf{0}, \mathbf{s}]) \right| \left(c_{13}(s_1, t) c_{23}(s_2, t) \right) d\lambda^2(\mathbf{s})d\lambda(t) \\
&\quad + \int_{\mathbb{I}} \int_{\mathbb{I}} |F_{1|3}(s_1|t) - \tilde{F}_{1|3}(s_1|t)| c_{13}(s_1, t) d\lambda(s_1)d\lambda(t)
\end{aligned}$$

$$\begin{aligned}
& + \int_{\mathbb{I}} \int_{\mathbb{I}} |F_{2|3}(s_2|t) - \tilde{F}_{2|3}(s_2|t)| c_{23}(s_2, t) \, d\lambda(s_2) d\lambda(t) \\
& \leq a^2 D_1(C, \tilde{C}) + a D_1(C_{13}, \tilde{C}_{13}) + a D_1(C_{23}, \tilde{C}_{23}) \\
& \leq a^2 D_\infty(C, \tilde{C}) + a D_\infty(C_{13}, \tilde{C}_{13}) + a D_\infty(C_{23}, \tilde{C}_{23}) \\
& \leq a^2 D_\infty(C, \tilde{C}) + 2a D_\infty(C, \tilde{C})
\end{aligned}$$

This proves the assertion. \square

Suppose that $C \in \mathcal{C}^3$ is a checkerboard copula. Then $C \in \mathcal{C}_{ac}^3$. We will say that C has resolution $N \geq 2$ if N is the smallest integer such that (there is a version of) its density c of C is constant on each square of the form $(\frac{i_x-1}{N}, \frac{i_x}{N}) \times (\frac{i_y-1}{N}, \frac{i_y}{N}) \times (\frac{i_z-1}{N}, \frac{i_z}{N})$ with $i_x, i_y, i_z \in \{1, \dots, N\}$. Notice that if C is a checkerboard copula with resolution N then its density c fulfills $c(\mathbf{u}, t) \leq N^2$ for λ^3 -almost all $(\mathbf{u}, t) \in \mathbb{I}^3$. Given a checkerboard copula C with resolution N w.l.o.g. we may assume that the mapping $t \mapsto C_{12;3}^t$ is constant on each interval of the form $[\frac{i-1}{N}, \frac{i}{N})$, $i \in \{1, \dots, N\}$, and define the quantity $\Delta = \Delta(C)$ by

$$\Delta(C) := \max_{i,j \in \{1, \dots, N\}} \int_{\mathbb{I}^2} |C_{12;3}^{t_i}(\mathbf{u}) - C_{12;3}^{t_j}(\mathbf{u})| \, d\lambda^2(\mathbf{u})$$

whereby $t_i = \frac{i-1}{N}$ for every $i \in \{1, \dots, N\}$.

Lemma A.5. *Suppose that $C_1, C_2, C_3, \dots \in \mathcal{C}_S^3$ are simplified copulas and that C is a non-simplified checkerboard copula with resolution $N \geq 2$. Then the quantity $J(C_n, C)$ from Lemma A.4 fulfills*

$$(A.1) \quad J(C_n, C) \geq \Delta(C)/N > 0$$

for every $n \in \mathbb{N}$. As a direct consequence, there is no sequence $(C_n)_{n \in \mathbb{N}}$ in \mathcal{C}_S^3 that converges to C w.r.t. D_∞ (D_1) or weakly conditional.

Proof. Under the assumptions of the lemma we obviously have

$$\begin{aligned}
J(C_n, C) &= \sum_{i=1}^N \frac{1}{N} \int_{\mathbb{I}^2} |C_{12;3}^{t_i}(\mathbf{u}) - (C_n)_{12;3}(\mathbf{u})| \, d\lambda^2(\mathbf{u}) \\
&\geq \frac{1}{N} \max_{i,j \in \{1, \dots, N\}} \int_{\mathbb{I}^2} |C_{12;3}^{t_i}(\mathbf{u}) - C_{12;3}^{t_j}(\mathbf{u})| \, d\lambda^2(\mathbf{u}) \\
&= \frac{\Delta(C)}{N} > 0.
\end{aligned}$$

The second assertion now follows from Lemma A.4 and the fact that $\Delta(C)$ only depends on C and not on n , the assertion concerning weak conditional convergence from the fact that weak conditional convergence implies convergence w.r.t. D_1 . \square

Lemma A.6.

- (1) *The family of all non-simplified checkerboards is dense in $(\mathcal{C}^3, D_\infty)$, in (\mathcal{C}^3, D_1) , and dense in \mathcal{C}^3 endowed with the topology induced by weak conditional convergence.*
- (2) *The family of all non-simplified checkerboards with positive density is dense in the family of all absolutely continuous copulas with positive density w.r.t. D_∞ , w.r.t. D_1 , and w.r.t. the topology induced by weak conditional convergence.*

Proof. To prove the first assertion let $C \in \mathcal{C}^3$ be arbitrary but fixed. Since according to [11] checkerboard copulas are dense in (\mathcal{C}^3, D_1) we can find a sequence $(B_n)_{n \in \mathbb{N}}$ of

checkerboard copulas with $\lim_{n \rightarrow \infty} D_1(B_n, C) = 0$. For every $n \in \mathbb{N}$ let E_n be a non-simplified checkerboard copula with the same resolution and the same $(1, 3)$ - and $(2, 3)$ -marginals as B_n . Setting $C_n := (1 - \frac{1}{n})B_n + \frac{1}{n}E_n$ for every $n \in \mathbb{N}$ yields a sequence $(C_n)_{n \in \mathbb{N}}$ of non-simplified checkerboard copulas. Considering

$$D_1(C_n, C) \leq (1 - \frac{1}{n})D_1(C, B_n) + \frac{1}{n}D_1(C, E_n)$$

it follows that $\lim_{n \rightarrow \infty} D_1(C_n, C) = 0$, which completes the proof of the first assertion concerning D_1 and D_∞ . The assertion concerning weak conditional convergence can be shown analogously: in fact, it is straightforward to extend the bivariate proof in [20, Theorem 3.2] to the three-dimensional setting, hence reusing the convex combination idea and considering $C_n := (1 - \frac{1}{n})B_n + \frac{1}{n}E_n$ yields the desired result.

To prove the second assertion suppose that $C \in \mathcal{C}_{ac}^3$ has positive density. According to the first assertion we can find a sequence $(B_n)_{n \in \mathbb{N}}$ of non-simplified checkerboard copulas with $\lim_{n \rightarrow \infty} D_1(B_n, C) = 0$. Setting $C_n := (1 - \frac{1}{n})B_n + \frac{1}{n}\Pi$ for every $n \in \mathbb{N}$ yields a sequence $(C_n)_{n \in \mathbb{N}}$ of non-simplified checkerboard copulas with positive density. Considering

$$D_1(C_n, C) \leq (1 - \frac{1}{n})D_1(C, B_n) + \frac{1}{n}D_1(C, \Pi)$$

we get $\lim_{n \rightarrow \infty} D_1(C_n, C) = 0$. Since the assertion for weak conditional convergence can be shown analogously, the proof is complete. \square

APPENDIX B. PROOFS

Proof of Lemma 6.4: For every $u \in \mathbb{I}$ we have

$$\begin{aligned} & \int_{\mathbb{I}} |K_{(C_n)_{13}}(v, [0, u]) - K_{C_{13}}(v, [0, u])| \, d\lambda(v) \\ &= \int_{\mathbb{I}} |K_{C_n}(v, [0, u] \times \mathbb{I}) - K_C(v, [0, u] \times \mathbb{I})| \, d\lambda(v) \\ &\leq \sup_{\mathbf{u} \in \mathbb{I}^2} \int_{\mathbb{I}} |K_{C_n}(v, [\mathbf{0}, \mathbf{u}]) - K_C(v, [\mathbf{0}, \mathbf{u}])| \, d\lambda(v) \end{aligned}$$

and hence $D_\infty((C_n)_{13}, C_{13}) \leq D_\infty(C_n, C)$. Since D_1 -convergence is equivalent to D_∞ -convergence (see [11]) this proves (1). We now prove the second assertion. Using Lipschitz continuity of copulas we obtain

$$\begin{aligned} & D_1(\psi(C_n), \psi(C)) \\ &= \int_{\mathbb{I}^2} \int_{\mathbb{I}} |(C_n)_p((F_n)_{1|3}(u_1|v), (F_n)_{2|3}(u_2|v)) - C_p(F_{1|3}(u_1|v), F_{2|3}(u_2|v))| \, d\lambda(v) d\lambda^2(\mathbf{u}) \\ &\leq \int_{\mathbb{I}^2} \int_{\mathbb{I}} |(C_n)_p((F_n)_{1|3}(u_1|v), (F_n)_{2|3}(u_2|v)) - (C_n)_p(F_{1|3}(u_1|v), F_{2|3}(u_2|v))| \, d\lambda(v) d\lambda^2(\mathbf{u}) \\ &\quad + \int_{\mathbb{I}^2} \int_{\mathbb{I}} |(C_n)_p(F_{1|3}(u_1|v), F_{2|3}(u_2|v)) - C_p(F_{1|3}(u_1|v), F_{2|3}(u_2|v))| \, d\lambda(v) d\lambda^2(\mathbf{u}) \\ &\leq \int_{\mathbb{I}^2} \int_{\mathbb{I}} |(F_n)_{1|3}(u_1|v) - F_{1|3}(u_1|v)| + |(F_n)_{2|3}(u_2|v) - F_{2|3}(u_2|v)| \, d\lambda(v) d\lambda^2(\mathbf{u}) \end{aligned}$$

$$\begin{aligned}
& + \int_{\mathbb{I}^2} \int_{\mathbb{I}} |(C_n)_p(F_{1|3}(u_1|v), F_{2|3}(u_2|v)) - C_p(F_{1|3}(u_1|v), F_{2|3}(u_2|v))| \, d\lambda(v) d\lambda^2(\mathbf{u}) \\
& \leq D_1((C_n)_{13}, C_{13}) + D_1((C_n)_{23}, C_{23}) + d_\infty((C_n)_p, C_p)
\end{aligned}$$

from which the assertion follows. \square

Proof of Lemma 6.5: We first have

$$\begin{aligned}
d_\infty((C_n)_p, C_p) &= \sup_{\mathbf{s} \in \mathbb{I}^2} \left| \int_{\mathbb{I}} (C_n)_{12;3}^t(\mathbf{s}) \, d\lambda(t) - \int_{\mathbb{I}} C_{12;3}^t(\mathbf{s}) \, d\lambda(t) \right| \\
&\leq \sup_{\mathbf{s} \in \mathbb{I}^2} \int_{\mathbb{I}} |(C_n)_{12;3}^t(\mathbf{s}) - C_{12;3}^t(\mathbf{s})| \, d\lambda(t) \\
&= D_\infty(B_n, B)
\end{aligned}$$

where the copulas B_n, B are given by

$$B_n(\mathbf{u}, v) := \int_{[0,v]} (C_n)_{12;3}^t(\mathbf{u}) \, d\lambda(t) \quad \text{and} \quad B(\mathbf{u}, v) := \int_{[0,v]} C_{12;3}^t(\mathbf{u}) \, d\lambda(t).$$

Since D_∞ -convergence is equivalent to D_1 -convergence it suffices to prove that $(B_n)_{n \in \mathbb{N}}$ converges to B w.r.t. D_1 , which can be done as follows: Applying Lemma A.3 and Hölder's inequality yields

$$\begin{aligned}
& D_1(B_n, B) \\
&= \int_{\mathbb{I}^2} \int_{\mathbb{I}} |(C_n)_{12;3}^t(\mathbf{s}) - C_{12;3}^t(\mathbf{s})| \, d\lambda(t) d\lambda^2(\mathbf{s}) \\
&\leq \int_{\mathbb{I}} \int_{\mathbb{I}^2} \left| K_C(t, [\mathbf{0}, \mathbf{s}]) - K_{C_n}(t, [\mathbf{0}, \mathbf{s}]) \right| \left(c_{13}(s_1, t) c_{23}(s_2, t) \right) \, d\lambda^2(\mathbf{s}) d\lambda(t) \\
&\quad + \int_{\mathbb{I}} \int_{\mathbb{I}} |F_{1|3}(s_1|t) - (F_n)_{1|3}(s_1|t)| c_{13}(s_1, t) \, d\lambda(s_1) d\lambda(t) \\
&\quad + \int_{\mathbb{I}} \int_{\mathbb{I}} |F_{2|3}(s_2|t) - (F_n)_{2|3}(s_2|t)| c_{23}(s_2, t) \, d\lambda(s_2) d\lambda(t) \\
&\leq \left(\int_{\mathbb{I}^2 \times \mathbb{I}} \left| K_C(t, [\mathbf{0}, \mathbf{s}]) - K_{C_n}(t, [\mathbf{0}, \mathbf{s}]) \right|^p \, d\lambda^3(\mathbf{s}, t) \right)^{\frac{1}{p}} \\
&\quad \cdot \left(\int_{\mathbb{I}^2 \times \mathbb{I}} \left(c_{13}(s_1, t) c_{23}(s_2, t) \right)^{\frac{p}{p-1}} \, d\lambda^3(\mathbf{s}, t) \right)^{\frac{p-1}{p}} \\
&\quad + \left(\int_{\mathbb{I} \times \mathbb{I}} |F_{1|3}(s_1|t) - (F_n)_{1|3}(s_1|t)|^q \, d\lambda^2(s_1, t) \right)^{\frac{1}{q}} \cdot \left(\int_{\mathbb{I} \times \mathbb{I}} |c_{13}(s_1, t)|^{\frac{q}{q-1}} \, d\lambda^2(s_1, t) \right)^{\frac{q-1}{q}}
\end{aligned}$$

$$+ \left(\int_{\mathbb{I} \times \mathbb{I}} |F_{2|3}(s_2|t) - (F_n)_{2|3}(s_2|t)|^r d\lambda^2(s_2, t) \right)^{\frac{1}{r}} \cdot \left(\int_{\mathbb{I} \times \mathbb{I}} |c_{23}(s_2, t)|^{\frac{r}{r-1}} d\lambda^2(s_2, t) \right)^{\frac{r-1}{r}}$$

for all $p, q, r \in (1, \infty)$. The latter expressions are finite by assumption, and the former part is bounded by

$$\begin{aligned} & \left(\int_{\mathbb{I}^2 \times \mathbb{I}} |K_C(t, [\mathbf{0}, \mathbf{s}]) - K_{C_n}(t, [\mathbf{0}, \mathbf{s}])|^p d\lambda^3(\mathbf{s}, t) \right)^{\frac{1}{p}} \\ & \leq \left(\int_{\mathbb{I}^2 \times \mathbb{I}} |K_C(t, [\mathbf{0}, \mathbf{s}]) - K_{C_n}(t, [\mathbf{0}, \mathbf{s}])|^1 d\lambda^3(\mathbf{s}, t) \right)^{\frac{1}{p}} = D_1(C_n, C)^{\frac{1}{p}} \end{aligned}$$

by $D_1((C_n)_{13}, C_{13})^{1/q}$ and $D_1((C_n)_{23}, C_{23})^{1/r}$, respectively. Thus we conclude that $\lim_{n \rightarrow \infty} D_1(B_n, B) = 0$ and hence $\lim_{n \rightarrow \infty} d_\infty((C_n)_p, C_p) = 0$. This proves the assertion. \square

ACKNOWLEDGEMENT

The second and the third author gratefully acknowledge the support of the WISS 2025 project 'IDA-lab Salzburg' (20204-WISS/225/197-2019 and 20102-F1901166-KZP).

REFERENCES

- [1] Aas, K., C. Czado, A. Frigessi, and H. Bakken (2009). Pair-copula constructions of multiple dependence. *Insur. Math. Econ.* 44, 182–198.
- [2] Acar, E., C. Genest, and N. J. (2012). Beyond simplified pair-copula constructions. *J. Multivariate Anal.* 110, 74–90.
- [3] Bedford, T. and R. Cooke (2002). Vines: A new graphical model for dependent random variables. *Ann. Stat.* 30(4), 1031–1068.
- [4] Bergsma, I. (2011). Nonparametric testing of conditional independence by means of the partial copula. Available at <https://arxiv.org/abs/1101.4607v1>.
- [5] Biller, B. and C. G. Corlu (2011). Accounting for parameter uncertainty in large-scale stochastic simulations with correlated inputs. *Oper. Res.* 59, 661–673.
- [6] Chevallier, J., D. Nguyen, and J. C. Reboredo (2019). A conditional dependence approach to co2-energy price relationships. *Energy Economics* 81, 812–821.
- [7] Dalla Valle, L., M. De Giuli, C. Tarantola, and C. Manelli (2016). Default probability estimation via pair copula constructions. *Eur. J. Oper. Res.* 249, 198–311.
- [8] Derumigny, A. and J.-D. Fermanian (2017). About tests of the 'simplifying' assumption for conditional copulas. *Depend. Model.* 5, 154–197.
- [9] Durante, F. and J. Fernández-Sánchez (2010). Multivariate shuffles and approximation of copulas. *Statist. Probab. Lett.* 80, 1827–1834.
- [10] Durante, F. and C. Sempi (2016). *Principles of Copula Theory*. CRC Press, Boca Raton, FL.
- [11] Fernández-Sánchez, J. and W. Trutschnig (2015). Conditioning based metrics on the space of multivariate copulas and their interrelation with uniform and levelwise convergence and iterated function systems. *J. Theoret. Probab.* 28, 1311–1336.
- [12] Fuchs, S., Y. McCord, and K. D. Schmidt (2018). Characterizations of copulas attaining the bounds of multivariate kendall's tau. *J. Optim. Theory Appl.* 178(2), 424–438.

- [13] Gijbels, I., M. Omelka, and N. Veraverbeke (2015a). Estimation of a copula when a covariate affects only marginal distributions. *Scand. J. Stat.* 42, 1109–1126.
- [14] Gijbels, I., M. Omelka, and N. Veraverbeke (2015b). Partial and average copulas and association measures. *Electr. J. Stat.* 9, 2420–2474.
- [15] Gijbels, I., M. Omelka, and N. Veraverbeke (2017). Nonparametric testing for no covariate effects in conditional copulas. *Statistics*. 51, 475–509.
- [16] Hobæk Haff, I. (2013). Parameter estimation for pair-copula constructions. *Bernoulli* 19(2), 462–491.
- [17] Hobæk Haff, I., K. Aas, and A. Frigessi (2010). On the simplified pair-copula construction – simply useful or too simplistic? *J. Multivariate Anal.* 101, 1296–1310.
- [18] Joe, H. (1996). Families of m -variate distributions with given margins and $m(m-1)/2$ bivariate dependence parameters. *Lecture Notes-Monograph Series* 28, 120–141.
- [19] Kallenberg, O. (1997). *Foundations of Modern Probability*. New York: Springer.
- [20] Kasper, T., S. Fuchs, and W. Trutschnig (2020). On weak conditional convergence of bivariate Archimedean and extreme value copulas, and consequences to nonparametric estimation. to appear in *Bernoulli*, Available at <http://www.bernoulli--society.org/index.php/publications/bernoulli--journal/bernoulli--journal--papers>, <https://arxiv.org/abs/2006.07131>.
- [21] Kauermann, G. and C. Schellhase (2014). Flexible pair-copula estimation in d-vines using bivariate penalized splines. *Stat. Comput.* 24, 1081–1100.
- [22] Killiches, M., D. Kraus, and C. Czado (2017). Examination and visualisation of the simplifying assumption for vine copulas in three dimensions. *Aust. N. Z. J. Stat.* 59(1), 95–117.
- [23] Klenke, A. (2007). *Probability Theory - A Comprehensive Course*. Berlin: Springer.
- [24] Kurz, M. and F. Spanhel (2017). Testing the simplifying assumption in high-dimensional vine copulas. Available at <https://arxiv.org/abs/1706.02338>.
- [25] Lancaster, H. O. (1963). Correlation and complete dependence of random variables. *Ann. Math. Statist.* 34(4), 1315–1321.
- [26] Mikusinski, P. and M. Taylor (2010). Some approximations of n -copulas. *Metrika* 72, 385–414.
- [27] Nagler, T. and C. Czado (2016). Evading the curse of dimensionality in nonparametric density estimation with simplified vine copulas. *J. Multivariate Anal.* 151, 69–89.
- [28] Nelsen, R. and M. Úbeda-Flores (2012). How close are pairwise and mutual independence? *Statist. Probab. Lett.* 82, 1823–1828.
- [29] Nelsen, R. B. (2006). *An Introduction to Copulas*. (Second ed.). New York: Springer.
- [30] Portier, F. and J. Segers (2018). On the weak convergence of the empirical conditional copula under a simplifying assumption. *J. Multivariate Anal.* 166, 160–181.
- [31] Reid, M. D. and R. C. Williamson (2009). Generalised Pinsker inequalities. <https://arxiv.org/abs/0906.1244>.
- [32] Rui, Z. (2019). Modelling mortality dependence with regime-switching copulas. *ASTIN Bulletin* 49, 373–407.
- [33] Spanhel, F. and M. Kurz (2016). The partial copula: Properties and associated dependence measures. *Statist. Probab. Lett.* 119, 76–83.
- [34] Spanhel, F. and M. Kurz (2019). Simplified vine copula models: Approximations based on the simplifying assumption. *Electron. J. Stat.* 13, 1254–1291.
- [35] Stöber, J., H. Joe, and C. Czado (2013). Simplified pair copula constructions: Limitations and extensions. *J. Multivariate Anal.* 119, 101–118.

- [36] Xiong, L., K.-X. Yu, and L. Gottschalk (2014). Estimation of the distribution of annual runoff from climatic variables using copulas. *Water Resour. Res.* 50(9), 7134–7152.
- [37] Zhang, Y. and Y. Shao (2020). A numerical strategy to evaluate performance of predictive scores via a copula-based approach. *Stat. Med.* *published online*.
- [38] Zhi, B., X. Wang, and F. Xu (2020). Impawn rate optimisation in inventory financing: A canonical vine copula-based approach. *Int. J. Prod. Econ.* 227, 107659.

February 10, 2021

DEPARTMENT FOR MATHEMATICS, UNIVERSITY OF SALZBURG, HELLBRUNNERSTRASSE 34, A-5020
SALZBURG, AUSTRIA

Email address: thomasmroz@a1.net, sebastian.fuchs@sbg.ac.at, wolfgang.trutschnig@sbg.ac.at