GENERALIZED JACOBSON'S LEMMA IN A BANACH ALGEBRA

HUANYIN CHEN AND MARJAN SHEIBANI

ABSTRACT. Let \mathcal{A} be a Banach algebra, and let $a, b, c \in \mathcal{A}$ satisfying

 $a(ba)^2 = abaca = acaba = (ac)^2a.$

We prove that $1 - ba \in \mathcal{A}^d$ if and only if $1 - ac \in \mathcal{A}^d$. In this case, $(1 - ac)^d = [1 - a(1 - ba)^{\pi}(1 - \alpha(1 + ba))^{-1}bac](1 + ac) + a(1 - ba)^d bac$. This extends the main result on g-Drazin inverse of Corach (Comm. Algebra, **41**(2013), 520–531).

1. INTRODUCTION

Let \mathcal{A} be a complex Banach algebra with identity. An element $a \in \mathcal{A}$ has g-Drazin inverse in case there exists $x \in \mathcal{A}$ such that

$$x = xax, ax = xa, a - a^2x \in \mathcal{A}^{qnil}$$

The preceding x is unique if it exists, we denote it by a^d . Here, $\mathcal{A}^{qnil} = \{a \in \mathcal{A} \mid 1 + ax \in \mathcal{A}^{-1} \text{ whenever } ax = xa\}$, where \mathcal{A}^{-1} stands for the set of all invertible elements of \mathcal{A} .

For any $a, b \in \mathcal{A}$, Jacobson's Lemma for invertibility states that $1-ab \in \mathcal{A}^{-1}$ if and only if $1-ba \in \mathcal{A}^{-1}$ and $(1-ba)^{-1} = 1+b(1-ab)^{-1}a$ (see [8, Lemma 1.4]). Jacobson's Lemma plays an important role in matrix and operator theory. Let $a, b \in \mathcal{A}$. Zhuang et al. proved the Jacobson's Lemma for g-Drazin inverse. That is, it was proved that $1-ab \in \mathcal{A}^d$ if and only if $1-ba \in \mathcal{A}^d$ and

$$(1 - ba)^d = 1 + b(1 - ab)^d a$$

²⁰¹⁰ Mathematics Subject Classification. 15A09, 47A11.

Key words and phrases. Generalized Drazin inverse; Drazin inverse; Group inverse; Generalized Jacobson's Lemma.

(see [14, Theorem 2.3]). Corach et al. generalized Jacobson's Lemma for g-Drain inverse to the case that aba = aca (see [4, Theorem 1]).

The motivation of this paper is to present a new generalized Jacobson's lemma for generalized Drazin inverses. We thereby extend [14, Theorem 2.3] to a wider case. Let $a, b \in \mathcal{A}$. In [2, Theorem 2.2], the authors considered the condition

$$a(ba)^2 = abaca = acaba = (ac)^2 a$$

for Cline's formula, i.e., $ba \in \mathcal{A}^d$ if and only if $ac \in \mathcal{A}^d$ and $(ba)^d = b[(ac)^d]^2 a$. Common local spectral properties for bounded linear operators under the preceding conditions was investigated in [15]. This raises a problem if Jacobson's lemma for generalized Drazin inverse hold under such wider condition. We shall give a confirmative answer to this problem. That is, If $a, b, c, \in \mathcal{A}$ satisfying $a(ba)^2 = abaca = acaba = (ac)^2 a$, we prove that $1 - ba \in \mathcal{A}^d$ if and only if $\beta = 1 - ac \in \mathcal{A}^d$. In this case, $(1-ac)^d = [1-d(-ba)^{\pi}(1-\alpha(1+ba))^{-1}bac](1+ac)+a(1-ba)^d bac$.

The Drazin inverse of $a \in \mathcal{A}$, denoted by a^D , is the unique element a^D satisfying the following three equations

$$a^D = a^D a a^D, a a^D = a^D a, a^k = a^{k+1} a$$

for some $k \in \mathbb{N}$. The small integer k is called the Drazin index of a, and is denoted by i(a). Moreover, we prove the generalized Jacobson's lemma for the Drazin inverse under the preceding condition.

Throughout the paper, all Banach algebra are complex with identity. \mathcal{A}^D and \mathcal{A}^d denote the sets of all Drazin and g-Drazin invertible elements in \mathcal{A} respectively. We use \mathcal{A}^{nil} to denote the set of all nilpotents of Banach algebra \mathcal{A} . \mathbb{C} stands for the field of all complex numbers.

2. GENERALIZED JACOBSON'S LEMMA

We come now to the main result of this paper which will be the tool in our following development.

Lemma 2.1. Let \mathcal{A} be a Banach algebra, let $m \in \mathbb{N}$ and let $a \in \mathcal{A}$. Then a has g-Drazin inverse if and only if there exists $b \in comm(a)$

such that

$$\left[ab - (ab)^2\right]^m = 0, a - a^2b \in \mathcal{A}^{qnil}.$$

In this case,

$$a^{d} = (a + 1 - e)^{-1}e, e = \sum_{i=0}^{m-1} (ab)^{2m-i}(1 - ab)^{i}.$$

Proof. \Longrightarrow Since $a \in \mathcal{A}^d$, there exists $b \in comm(a)$ such that

$$b = bab, a - a^2b \in \mathcal{A}^{qnil}$$

Hence $ab = (ab)^2$. Moreover, we have $e = \sum_{i=0}^{m-1} {\binom{2m}{i}} (ab)^{2m-i} (1-ab)^i = ab$, and then $(a + 1 - e)^{-1}e = (a + 1 - ab)^{-1}ab = (a + 1 - ab)^{-1}(a + 1 - ab)b = b = a^d$, as desired. \Leftarrow Let

$$e = \sum_{i=0}^{m-1} \begin{pmatrix} 2m \\ i \end{pmatrix} (ab)^{2m-i} (1-ab)^i, f = \sum_{i=0}^{m-1} \begin{pmatrix} 2m \\ i \end{pmatrix} (ab)^{2m-i} (1-ab)^i.$$

Since $[ab - (ab)^2]^m = 0$, we have $(ab)^m (1 - ab)^m = 0$, and so $e + f = [ab + (1 - ab)]^{2m} = 1, ef = fe = 0.$

Hence, $e^2 = e^2 + ef = e$. Clearly, a, b and e commute one another. Then we have

$$w := ab - e$$

= $[ab - (ab)^{2}] + (ab)[ab - (ab)^{2}] + \dots + (ab)^{2m-2}[ab - (ab)^{2}]$
- $\sum_{i=1}^{m-1} \sum_{i=0}^{m-1} {2m \choose i} (ab)^{2m-i}(1-ab)^{i}.$

Hence $w^m = 0$. By hypothesis, $a - a^2b = a(1-ab) = a(1-e) - aw \in \mathcal{A}^{qnil}$, and so $a(1-e) \in \mathcal{A}^{qnil}$. Let c = bab. Then

$$ac + 1 - e = 1 + (ab)^{2} - (ab)^{2m} - \sum_{i=1}^{m-1} (ab)^{2m-i} (1 - ab)^{i}$$

$$= 1 + [(ab)^{2} - (ab)^{3}] + [(ab)^{3} - (ab)^{4}]$$

$$+ \dots + [(ab)^{2m-1} - (ab)^{2m}] - \sum_{i=1}^{m-1} (ab)^{2m-i} (1 - ab)^{i}$$

$$\in \mathcal{A}^{-1}.$$

Also we have

$$c(1-e) = bab \left[(1-(ab)^{2m}) - \sum_{i=1}^{m-1} (ab)^{2m-i} (1-ab)^i \in \mathcal{A}^{nil}. \right]$$

We easily check that

$$\begin{array}{rcl} (a+1-e)(c+1-e) &=& ac+a(1-e)+(1-e)c+1-e\\ &=& ac+1-e+a(1-e)+c(1-e)\\ &\in& \mathcal{A}^{-1}. \end{array}$$

Hence $a + (1 - e) \in \mathcal{A}^{-1}$. Therefore $a^d = (a + 1 - e)^{-1}e$, as required.

Theorem 2.2. Let \mathcal{A} be a Banach algebra, and let $a, b, c \in \mathcal{A}$ satisfying

$$a(ba)^2 = abaca = acaba = (ac)^2a.$$

- Then $\alpha = 1 ba \in \mathcal{A}^d$ if and only if $\beta = 1 ac \in \mathcal{A}^d$. In this case, $\beta^d = \left[1 - a\alpha^{\pi}(1 - \alpha(1 + ba))^{-1}bac\right](1 + ac) + a\alpha^d bac.$
- Proof. Let $p = \alpha^{\pi}, x = \alpha^{d}$. Then $1 p\alpha(1 + ba) \in \mathcal{A}^{-1}$. Let $y = \left[1 ap(1 p\alpha(1 + ba))^{-1}bac\right](1 + ac) + axbac$.

Step 1. $[y\beta - (y\beta)^2]^2 = 0$. We see that $y\beta = 1 - (ac)^2 - ap(1 - p\alpha(1 + ba))^{-1}bac[1 - (ac)^2] + axbac(1 - ac).$

We compute that

$$y\beta a = \left[1 - (ac)^2 - ap(1 - p\alpha(1 + ba))^{-1}bac\left[1 - (ac)^2\right] + axbac(1 - ac)\right]a$$

$$= a - (abac - axbac(1 - ac))a - ap(1 - p\alpha(1 + ba))^{-1}[bac - bac(ac)^2]a$$

$$= a - [abac - ax(bac - bacac)]a - ap(1 - p\alpha(1 + ba))^{-1}[ba - bacaba]ca$$

$$= a - [abac - ax(1 - ba)bac]a - ap(1 - p\alpha(1 + ba))^{-1}[1 - (ba)^2]baca$$

$$= a - apbaca - ap(1 - p\alpha(1 + ba))^{-1}p\alpha(1 + ba)baca$$

$$= a - ap(1 - p\alpha(1 + ba))^{-1}[(1 - p\alpha(1 + ba)) + p\alpha(1 + ba)]baca$$

$$= [1 - ap(1 - p\alpha(1 + ba))^{-1}bac]a.$$

Set $z = cac + p(1 - p\alpha(1 + ba))^{-1}bac[1 - (ac)^2] + xbac(1 - ac)$. Then $y\beta = 1 - az$, and so

$$y\beta(1-y\beta) = y\beta az = [1-ap(1-p\alpha(1+ba))^{-1}bac]az = 1-y\beta - ap(1-p\alpha(1+ba))^{-1}bac(1-y\beta).$$

4

Since $acaba = a(caba) = a(caca) = (ac)^2 a = (abac)a = abaca$, we have (baca)(ba) = (ba)(baca), and so $(baca)\alpha = \alpha(baca)$. Hence, (baca)x = x(baca), and then

 $ap(1-p\alpha(1+ba))^{-1}bacaxbac = ap(1-p(1+ba))^{-1}(baca)\alpha xbac = 0.$

Moreover, we have

 $\begin{aligned} &ap(1-p\alpha(1+ba))^{-1}bacy\beta(1-y\beta) \\ &= ap(1-p\alpha(1+ba))^{-1}bac[1-ap(1-p\alpha(1+ba))^{-1}bac(1+ac)]a(1-ca)z \\ &= \left[ap(1-p\alpha(1+ba))^{-1}bac(1+ac) + ap(1-p\alpha(1+ba))^{-2}(baca)bac(1+ac)\right]a(1-ca)z \\ &= \left[ap(1-p\alpha(1+ba))^{-1}(1+ba)bac + ap(1-p\alpha(1+ba))^{-2}(ba)^2(1+ba)bac\right]a(1-ca)z \\ &= \left[ap(1-p\alpha(1+ba))^{-2}\left[p-p\alpha(1+ba) - p(ba)^2\right](1+ba)bac\right]a(1-ca)z \\ &= 0. \end{aligned}$

Hence,

$$y\beta(1-y\beta)^{2} = \begin{bmatrix} 1-y\beta - ap(1-p\alpha(1+ba))^{-1}bac(1-y\beta) \\ 1-y\beta - ap(1-p\alpha(1+ba))^{-1}bac \end{bmatrix} (1-y\beta) \\ = \begin{bmatrix} (1-ap(1-p\alpha(1+ba))^{-1}bac)a - y\beta a \end{bmatrix} z \\ = 0.$$

Therefore $[y\beta - (y\beta)^2]^2 = y^2\beta^2(1 - y\beta)^2 = 0.$ Step 2. $\beta - \beta y\beta \in \mathcal{A}^{qnil}$. Clearly, we have

$$baca(baca)^2 p\alpha (1 - p\alpha (1 + ba))^{-3} = (baca)^3 (1 - p\alpha (1 + ba))^{-3} (\alpha - \alpha^2 \alpha) \in \mathcal{A}^{qnil}$$

By using Cline's formula, we derive

$$(1 - y\beta)a\alpha bacap(1 - p\alpha(1 + ba))^{-2}bac$$

= $ap(1 - p\alpha(1 + ba))^{-1}baca\alpha bacap(1 - p\alpha(1 + ba))^{-2}bac$
= $a(baca)^2p\alpha(1 - p\alpha(1 + ba))^{-3}bac$
 $\in \mathcal{A}^{qnil}.$

By using Cline's formula again, we have

$$\begin{array}{l} \beta(1-y\beta)^{3} \\ = & \beta ap(1-p\alpha(1+ba))^{-1}(baca)p(1-p\alpha(1+bd))^{-1}bac(1-y\beta) \\ = & \beta a(baca)p(1-p\alpha(1+bd))^{-2}bac(1-y\beta) \\ = & (1-ac)a(baca)p(1-p\alpha(1+ba))^{-2}bac(1-y\beta) \\ = & a\alpha bacap(1-p\alpha(1+ba))^{-2}bac(1-y\beta) \\ \in & \mathcal{A}^{qnil}. \end{array}$$

Therefore $\beta(1-y\beta)^3 \in \mathcal{A}^{qnil}$, and so $(\beta-\beta^2 y)^3 \in \mathcal{A}^{qnil}$, This implies that $\beta - \beta^2 y \in \mathcal{A}^{qnil}$, as required.

Step 3. $y \in comm(\beta)$. Set s = ac. Then we have Claim 1. $\beta(axbac) = (axbac)\beta$. We easily check that

(bacsaba)ba = bacacababa = babacacaba == ba(bacsaba),

and so

$$(bacsaba)\alpha = \alpha(bacsaba).$$

Hence (bacsaba)x = x(bacsaba). Obviously,

$$p = (ba)^2 p [1 - p\alpha(1 + ba)]^{-1} = (ba)^4 p [1 - p\alpha(1 + ba)]^{-2}.$$

Then

$$\begin{split} s(apbac) &= sa(ba)^4 p [1 - p\alpha(1 + ba)]^{-2} bac \\ &= s(ac)^2 ababap [1 - p\alpha(1 + ba)]^{-2} bac \\ &= a(bacsaba)bap [1 - p\alpha(1 + ba)]^{-2} bac \\ &= abap [1 - p\alpha(1 + ba)]^{-2} (bacsaba)bac \\ &= abap [1 - p\alpha(1 + ba)]^{-2} bacs(ac)^3 \\ &= abap [1 - p\alpha(1 + ba)]^{-2} (ba)^3 bacs \\ &= a(ba)^4 p [1 - p\alpha(1 + ba)]^{-2} bacs \\ &= (apbac)s. \end{split}$$

Since sabac = (acaba)c = (abaca)c = abacs, we have

$$sa\alpha xbac = sabac - s(apbac) = abacs - (apbac)s = a\alpha xbacs,$$

and so

$$saxbac - sabaxbac = axbacs - abaxbacs.$$

~

On the other hand, we have

$$\begin{split} s(abapbac) &= sa(ba)^5 p[1 - p\alpha(1+ba)]^{-2} bac \\ &= s(ac)^4 abap[1 - p\alpha(1+ba)]^{-2} bac \\ &= ababa(bacsaba)p[1 - p\alpha(1+ba)]^{-2} bac \\ &= ababap[1 - p\alpha(1+ba)]^{-2}(bacsaba)bac \\ &= ababap[1 - p\alpha(1+ba)]^{-2} bacs(ac)^3 \\ &= ababap[1 - p\alpha(1+ba)]^{-2}(ba)^3 bacs \\ &= aba(ba)^4 p[1 - p\alpha(1+ba)]^{-2} bacs \\ &= (abapbac)s. \end{split}$$

Since $sababac = s(ac)^3 = (ac)^3 s = ababacs$, we have $aba\alpha xbacs = saba\alpha xbac.$

6

Then we have

$$ababa\alpha xbacs = ac(aba\alpha xbacs)$$
$$= ac(saba\alpha xbac)$$
$$= sacaba\alpha xbac$$
$$= sababa\alpha xbac,$$

and so

$$aba(1+ba)\alpha xbacs = saba(1+ba)\alpha xbac$$

and then

$$abaxbacs - aba(ba)^2xbacs = sabaxbac - saba(ba)^2xbac.$$

One easily checks that

$$a(ba)^{3}xbacs = ax(ba)^{3}bacs$$

= $axb(ac)^{4}\beta$
= $ax(bacsaba)bac$
= $a(bacsaba)xbac$
= $(ac)^{2}sabaxbac$
= $sa(ba)^{3}xbac$.

This implies that abaxbacs = sabaxbac, and therefore $\beta(axbac) = (axbac)\beta$.

Claim 2. $sap(1 - p\alpha(1 + ba))^{-1}bac(1 + ac) = ap(1 - p\alpha(1 + ba))^{-1}bac(1 + ac)\beta$. Set $t = ap(1 - p\alpha(1 + ba))^{-1}bac(1 + ac)$. Since (bacsaba)ba = ba(bacsaba), we have

$$st = sap(1 - p\alpha(1 + ba))^{-1}bac(1 + ac)$$

= $sa(ba)^4p[1 - p\alpha(1 + ba)]^{-3}bac(1 + ac)$
= $(ac)^3sabap[1 - p\alpha(1 + ba)]^{-3}bac(1 + ac)$
= $a(ba)^4bap[1 - p\alpha(1 + ba)]^{-3}bac(1 + ac)$
= $abap[1 - p\alpha(1 + ba)]^{-3}(ba)^4bac(1 + ac)$
= $abap[1 - p\alpha(1 + ba)]^{-3}b(ac)^5(1 + ac)$

Also we have

$$ts = ap(1 - p\alpha(1 + ba))^{-1}bac(1 + ac)\beta$$

= $ap[1 - p\alpha(1 + ba)]^{-3}(ba)^4bsac(1 + ac)$
= $abap[1 - p\alpha(1 + ba)]^{-3}b(ac)^3sac(1 + ac)$
= $abap[1 - p\alpha(1 + ba)]^{-3}bs(ac)^4(1 + ac)$

Hence, st = ts, and so $\beta t = t\beta$.

Therefore $y \in comm(\beta)$, and then $y = \beta^d$, as required.

 \Leftarrow Since $1 - ac \in \mathbb{R}^d$, it follows by Jacobson's Lemma that $1 - ca \in \mathbb{R}^d$. Applying the preceding discussion, we obtain that $1 - ba \in \mathbb{R}^d$, as desired.

Corollary 2.3. Let R be a Banach algebra, let $\lambda \in \mathbb{C}$, and let $a, b, c \in R$ satisfying

$$a(ba)^2 = abaca = acaba = (ac)^2a.$$

Then $\lambda - ba \in \mathbb{R}^d$ if and only if $\lambda - ac \in \mathbb{R}^d$. In this case,

$$= \begin{cases} (\lambda - ac)^d & \lambda = 0\\ -a((ba)^d)^2c & \lambda = 0\\ \left[1 - a(1 - ba)^{\pi}(\lambda^2 - (\lambda - ba)(\lambda + ba))^{-1}bac\right] & \\ (\lambda + ac) + \frac{1}{\lambda}a(\lambda - ba)^dbac & \lambda \neq 0. \end{cases}$$

Proof. Case 1. $\lambda = 0$. It is obvious by [2, Theorem 2.2].

Case 2. $\lambda \neq 0$. Let $d = \frac{b}{\lambda}$ and $e = \frac{c}{\lambda}$. It is easy to show that $a(da)^2 = adaea = aeada = (ae^2)a$. By virtue of Theorem 2.2, $1 - da \in \mathbb{R}^d$ if and only if $1 - ae \in \mathbb{R}^d$. That is, $\lambda - ba \in \mathbb{R}^d$ if and only if $\lambda - ac \in \mathbb{R}^d$. In this case,

$$\begin{aligned} &(\lambda - ac)^d \\ &= \lambda(1 - ae)^d \\ &= \lambda \left[1 - a(1 - da)^{\pi} (1 - (1 - da)(1 + da))^{-1} dae \right] \\ &(1 + ae) + a(1 - ba)^d dae \\ &= \left[1 - a(1 - ba)^{\pi} (\lambda^2 - (\lambda - ba)(\lambda + ba))^{-1} bac \right] \\ &(\lambda + ac) + \frac{1}{\lambda} a(\lambda - ba)^d bac, \end{aligned}$$

as desired.

Corollary 2.4. Let R be a Banach algebra, let $\lambda \in \mathbb{C}$, and let $a, b, c \in R$ satisfying aba = aca Then $\lambda - ba \in R^d$ if and only if $\lambda - ac \in R^d$. In this case,

$$=\begin{cases} (\lambda - ac)^d & \lambda = 0\\ -a((ba)^d)^2c & \lambda = 0\\ \left[1 - a(1 - ba)^{\pi}(\lambda^2 - (\lambda - ba)(\lambda + ba))^{-1}bac\right] & \\ (\lambda + ac) + \frac{1}{\lambda}a(\lambda - ba)^dbac & \lambda \neq 0. \end{cases}$$

Proof. This is obvious by Corollary 2.3.

3. Drazin inverse

As it is known, $a \in \mathcal{A}^D$ if and only if there exists $x \in R$ such that $x = xax, x \in comm(a), a - a^2x \in \mathcal{A}^{nil}$, and so $a^D = a^d$. For the generalized Jacobson's Lemma for Drazin inverse, we have

Theorem 3.1. Let \mathcal{A} be a Banach algebra, and let $a, b, c \in \mathcal{A}$ satisfying

$$a(ba)^2 = abaca = acaba = (ac)^2a.$$

Then $\alpha = 1 - ba \in \mathcal{A}^D$ if and only if $\beta = 1 - ac \in \mathcal{A}^D$. In this case,

$$= \begin{bmatrix} \beta^D \\ [1 - a\alpha^{\pi}(1 - \alpha(1 + ba))^{-1}bac](1 + ac) + a\alpha^D bac, \\ i(1 - ba) \le i(1 - ac) + 1. \end{bmatrix}$$

Proof. Let $p = \alpha^{\pi}, x = \alpha^{D}$. In view of Theorem 2.1, $\beta \in \mathbb{R}^{d}$ and $\beta^{d} = [1 - an(1 - n\alpha(1 + ba))^{-1}bac](1 + ac) + dxbac.$

We shall prove that
$$\beta^D = \beta^d$$
.

We will suffice to check $\beta - \beta \beta^d \beta \in \mathcal{A}^{nil}$. As in the proof of Theorem 2.2, we have

$$\beta - \beta \beta^d \beta = \beta (1 - \beta^d \beta) = d\alpha bacap (1 - p\alpha (1 + ba))^{-2} bac.$$

In light of [2, Lemma 2.1], we will suffice to prove

$$baca\alpha bacap(1-p\alpha(1+ba))^{-2} \in \mathcal{A}^{nil}$$

Similarly to the discussion in Theorem 2.2, we see that $baca \in comm(\alpha)$, and so $baca, \alpha p$ and $(1 - p\alpha(1 + ba))^{-2}$ commute one another. Set $n = i(\alpha)$. Then

$$\begin{bmatrix} baca\alpha bacdp(1 - p\alpha(1 + ba))^{-2} \end{bmatrix}^n \\ = (baca)^2 (1 - p\alpha(1 + ba))^{-2n} (\alpha - \alpha^2 \alpha^d)^n \\ = 0;$$

hence,

$$(\beta - \beta \beta^d \beta)^{n+1} = a\alpha bacdp(1 - p\alpha(1 + ba))^{-2} [baca\alpha bacap(1 - p\alpha(1 + ba))^{-2}]^n bac = 0.$$

Thus, we have $\beta - \beta \beta^d \beta \in \mathcal{A}^{nil}$. Moreover, we have $i(\beta) \leq i(\alpha) + 1$, as desired. \Box

The group of $a \in \mathcal{A}$ is the unique element $a^{\#} \in \mathcal{A}$ which satisfies $a = aa^{\#}a, a^{\#} = a^{\#}aa^{\#}$. We denote the set of all group invertible elements of \mathcal{A} by $\mathcal{A}^{\#}$. As is well known, $a \in \mathcal{A}^{\#}$ if and only if $a \in \mathcal{A}^{D}$ and i(a) = 1. We are now ready to prove:

Corollary 3.2. Let \mathcal{A} be a Banach algebra, and let $a, b, c \in \mathcal{A}$ satisfying

$$a(ba)^2 = abaca = acaba = (ac)^2a.$$

Then $\alpha = 1 - ba$ has group inverse if and only if $\beta = 1 - ac$ has group inverse. In this case,

$$\beta^{\#} = \left[1 - a\alpha^{\pi}(1 - \alpha(1 + ba))^{-1}bac\right](1 + ac) + a\alpha^{\#}bac.$$

Proof. Since $1-ba \in \mathcal{A}^{\#}$, we have $1-ba \in \mathcal{A}^{D}$. In light of Theorem ???, $1-ac \in \mathcal{A}^{D}$. Let $\alpha = 1-ba$ and $\beta = 1-ac$. Let $p = 1-\alpha\alpha^{D}$. Since $\alpha \in R^{\#}$, we have $\alpha p = \alpha - \alpha^{2}\alpha^{D} = 0$. As in the proof of Theorem 2.2, we have

$$\begin{split} \beta &-\beta\beta^D\beta\\ &= d\alpha bacdp(1-p\alpha(1+ba))^{-2} \big[bacd\alpha bacdp(1-p\alpha(1+ba))^{-2} \big]^n bac\\ &= dbacd\alpha p(1-\alpha(1+ba))^{-2} \big[bacd\alpha bacdp(1-p\alpha(1+ba))^{-2} \big]^n bac\\ &= 0. \end{split}$$

Obviously, $\beta^D \in comm(\beta)$ and $\beta^D = \beta^D \beta \beta^D$. Therefore

$$\beta^{\#} = \beta^{D} \\ = \left[1 - a\alpha^{\pi}(1 - \alpha(1 + ba))^{-1}bac\right](1 + ac) + a(1 - ba)^{\#}bac.$$

This completes the proof.

Example 3.3.

Let

$$A = \begin{pmatrix} 1 & 0 & -1 & 0 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, B = \begin{pmatrix} 1 & 1 & 1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & -1 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ \end{pmatrix},$$
$$C = \begin{pmatrix} 1 & -1 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 1 & 1 & 0 & 1 \\ 0 & 1 & 0 & 0 \end{pmatrix} \in M_4(\mathbb{C}).$$

Then we check that

directly computation, we have

$$(I_4 - AC)^{\#} = \begin{pmatrix} 1 & -2 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$
$$(I_4 - BA)^{\#} = \begin{pmatrix} 2 & 3 & -1 & -3 \\ 0 & 2 & 0 & -1 \\ 1 & 1 & 0 & -1 \\ 0 & 1 & 0 & 0 \end{pmatrix}.$$

References

- C. Benhida and E.H. Zerouali, Local spetral theory of linear operators RS and SR, Integral Equ. Oper. Theory, 54(2006), 1–8.
- [2] H. Chen and M.S. Abdolyousefi, Cline's formula for g-Drazin inverses in a ring, *Filomat*, 33(2019), 2249–2255.
- [3] N. Castro-González; C. Mendes-Araújo and P. Parricio, Generalized inverses of a sum in rings, Bull. Aust. Math. Soc., 82(2010), 156–164.
- [4] G. Corach, Extensions of Jacobson's lemma, Comm. Algebra, 41(2013), 520-531.
- [5] D. Cvetkovic-Ilic and R. Harte, On Jacobson's lemma and Drazin invertibility, Applied Math. Letters, 23(2010), 417–420.

- [6] X. Mary, Weak inverses of products Cline's formula meets Jacobson lemma, J. Algebra Appl., 17(2018), DOI: 10.1142/S021949881850069X.
- [7] V.G. Miller and H. Zguitti, New extensions of Jacobson's lemma and Cline's formula, *Rend. Circ. Mat. Palermo, II. Ser.*, Published online: 09 February 2017, Doi: 10.1007/s12215-017-0298-6.
- [8] D. Mosic, Extensions of Jacobson's lemma for Drazin inverses, Aequat. Math., 91(2017), 419–428.
- [9] K. Yang and X. Fang, Common properties of the operator products in spectral theory, Ann. Funct. Anal., 6(2015), 60–69.
- [10] K. Yang and X. Fang, Common properties of the operator products in local spectral theory, Acta Math. Sin. Engl. Ser., 31(2015), 1715–1724.
- [11] K. Yan; Q. Zeng and Y. Zhu, On Drazin spectral equation for the operator products, *Complex Analysis and Operator Theory*, (2020) 14:12 https://doi.org/10.1007/s11785-019-00979-y.
- [12] K. Yan; Q. Zeng and Y. Zhu, Generalized Jacobson's lemma for Drazin inverses and its applications, *Linear and Multilinear Algebra*, 68(2020), 81–93.
- [13] Q.P. Zeng and H.J. Zhong, New results on common properties of the products AC and BA, J. Math. Anal. Appl., 427(2015), 830–840.
- [14] G. Zhuang; J. Chen and J. Cui, Jacobson's lemma for the generalized Drazin inverse, *Linear Algebra Appl.*, 436(2012), 742–746.
- [15] H. Zguitti, Further common local spectral properties for bounded linear operators, arXiv:1903.00522v1 [math.FA] 1 Mar 2019.

DEPARTMENT OF MATHEMATICS, HANGZHOU NORMAL UNIVERSITY, HANG -ZHOU, CHINA

E-mail address: <huanyinchen@aliyun.com>

WOMEN'S UNIVERSITY OF SEMNAN (FARZANEGAN), SEMNAN, IRAN *E-mail address:* <sheibani@fgusem.ac.ir>