On Frequentist Regret of Linear Thompson Sampling

Nima Hamidi

Department of Statistics, Stanford University, hamidi@stanford.edu

Mohsen Bayati

Graduate School of Business, Stanford University, bayati@stanford.edu

This paper studies the stochastic linear bandit problem, where a decision-maker chooses actions from possibly time-dependent sets of vectors in \mathbb{R}^d and receives noisy rewards. The objective is to minimize regret, the difference between the cumulative expected reward of the decision-maker and that of an oracle with access to the expected reward of each action, over a sequence of T decisions. Linear Thompson Sampling (LinTS) is a popular Bayesian heuristic, supported by theoretical analysis that shows its Bayesian regret is bounded by $\widetilde{\mathcal{O}}(d\sqrt{T})$, matching minimax lower bounds. However, previous studies demonstrate that the frequentist regret bound for LinTS is $\widetilde{\mathcal{O}}(d\sqrt{dT})$, which requires posterior variance inflation and is by a factor of \sqrt{d} worse than the best optimism-based algorithms. We prove that this inflation is fundamental and that the frequentist bound of $\widetilde{\mathcal{O}}(d\sqrt{dT})$ is the best possible, by demonstrating a randomization bias phenomenon in LinTS that can cause linear regret without inflation. We propose a data-driven version of LinTS that adjusts posterior inflation using observed data, which can achieve minimax optimal frequentist regret, under additional conditions. Our analysis provides new insights into LinTS and settles an open problem in the field.

Key words: Linear bandit, Contextual bandit, Thompson sampling, Data-driven exploration

1. Introduction

In recent years, an increasing number of organizations across diverse domains, including but not limited to e-commerce and digital advertising, are embracing the use of online experiments to optimize their decision-making process. However, conducting such experiments involves an opportunity cost, also known as *regret*, caused by exposing some customers to potentially inferior experiences. To reduce this opportunity cost, a growing number of enterprises are turning to multi-armed bandit (MAB) experiments (Scott 2010, 2015, Johari et al. 2017). The MAB approach works by adaptively utilizing the experiment's partially available results and favoring decisions with higher predicted value, or *reward*, thus reducing their regret. The practical motivations for MAB problems, combined with their mathematical richness, have made them the subject of intense study in computer science, economics, operations research, and statistics (Bubeck et al. 2012, Russo et al. 2018, Lattimore and Szepesvari 2019, Slivkins 2019).

This paper aims to answer an open question about one of the key algorithms used in MAB problems, which dates back to Thompson (1933), in a general setting known as the stochastic linear bandit problem with changing action sets. This class includes the standard k-armed bandit problem, as well as the k-armed

1

contextual bandit problem as special cases. In this setting, a decision-maker sequentially selects actions from given action sets and observes the corresponding rewards. The actions, which are vectors in \mathbb{R}^d , can also be thought of as features or context that influence the rewards. The rewards are stochastic and their expectations depend on the actions through a fixed linear function, with an unknown parameter $\Theta^* \in \mathbb{R}^d$. As more decisions are made and their rewards are observed, the reward function can be estimated. The main objective of the decision-maker is to maximize the cumulative expected reward over a sequence of decision epochs. Alternatively, one can measure the expected regret or simply regret, which is the difference between the best achievable cumulative expected reward, obtained by an *oracle* with access to the true expectation of the reward function, and the cumulative expected reward obtained by the decision-maker.

Regret can be measured in either a Bayesian or frequentist fashion. Bayesian regret is used when the unknown parameter Θ^* is random, and the expectations are taken with respect to three sources: (1) the randomness in the reward functions, (2) the unknown parameter Θ^* , and (3) possible randomness introduced by the decision-maker. On the other hand, frequentist regret (also referred to as worst-case regret) is used when the parameter Θ^* is deterministic, and the expectation is only with respect to the (1) and (3).

The main challenge faced by decision-makers is overcoming the *curse of underestimation*, where the reward of the optimal action is underestimated, leading to its permanent discarding. To address this challenge, two approaches have gained considerable attention. The first approach, proposed by Dani et al. (2008), Rusmevichientong and Tsitsiklis (2010) and improved by Abbasi-Yadkori et al. (2011), utilizes *optimism in the face of uncertainty* (based on the Upper Confidence Bound technique due to Lai and Robbins (1985)), and obtains policies with frequentist regret bounds of $\widetilde{\mathcal{O}}(d\sqrt{T})$. As shown by Dani et al. (2008), this approach is minimax optimal up to logarithmic factors. The second approach, introduced by Thompson (1933), arises from a Bayesian heuristic which suggests sampling from the posterior distribution of the reward function, given past observations, and choosing the best action as if this sample were the true reward function. This approach is known as Thompson sampling (TS) or posterior sampling, and although it is Bayesian in nature, it can be applied in the frequentist setting as well. TS is popular in practice due to its simplicity and good empirical performance, as reported by Scott (2010, 2015), Russo et al. (2018).

TS has been extensively studied from a theoretical perspective. For the stochastic linear bandit problem, where the TS heuristic is referred to as LinTS, Russo and Van Roy (2014) established a connection between LinTS and optimistic policies and obtained a Bayesian regret bound of $\widetilde{\mathcal{O}}(d\sqrt{T})$, which is minimax optimal. However, in the frequentist setting, Agrawal and Goyal (2013b) and Abeille et al. (2017) have derived regret bounds of $\widetilde{\mathcal{O}}(d\sqrt{dT})$ for a variant of LinTS (referred to as frequentist LinTS) that samples from a posterior distribution with an *inflated* variance of a factor $\widetilde{\mathcal{O}}(d)$. When d is not a constant, which is often the case in

¹ The notation $\widetilde{\mathcal{O}}(.)$ is defined in Section 2.

modern applications with a large number of customer-specific data that allows *personalizing* the decisions, this bound is far from optimal, being worse by a factor of \sqrt{d} . While it is known in the literature that frequentist LinTS has poor empirical performance due to its conservative over-exploration, it has remained an open question as to whether the inflation is necessary and whether the extra factor for the frequentist regret can be eliminated, see, for instance, (Russo et al. 2018, §8.1.2). Our *main contribution* in this paper is to answer this question negatively. In particular, we construct two families of examples to show that LinTS without inflation suffers from a *randomization bias* phenomenon and can incur linear regret when d grows at least logarithmically in T ($d = \Omega(\log T)$) and at least one of the noise distribution or the prior distribution does not match the one that LinTS assumes.

While the primary focus of this paper is theoretical, the examples we provide offer valuable insights into the inner workings of TS that could prove beneficial for practitioners. In practice, the prior and noise distributions are often unknown or difficult to sample from, necessitating the estimation or approximation of the posterior distribution. However, we demonstrate that even minor discrepancies between the true distributions and their estimates can significantly degrade the performance of TS, a problem that persists even when with noiseless reward function.

This shortcoming of TS can make certain applications vulnerable to adversarial attacks. Specifically, a commonly used assumption in posterior computation, that the set of actions is independent of the true reward function given past observations, can be violated if an adversary with partial knowledge of the true reward function can manipulate the action sets. For instance, consider an established marketplace platform \mathcal{P}_1 with extensive data that is competing with a new platform \mathcal{P}_2 that has limited access to data. \mathcal{P}_2 may use MAB experiments to expedite its learning and decision-making, but \mathcal{P}_1 , with superior knowledge of the true reward function, could participate maliciously in \mathcal{P}_2 's marketplace via intermediary agents and diminish \mathcal{P}_2 's experimental performance. As we will show in Section 3, this malicious participation need not entail active monitoring of \mathcal{P}_2 and can be passively planned in advance, making it a plausible concern.

It is worth highlighting that optimism-based algorithms, such as the OFUL algorithm proposed by Abbasi-Yadkori et al. (2011), do not suffer from the aforementioned randomization bias issue. Building on this insight, we aim to delve into the root cause of the problem with LinTS and present a solution in the form of the TS-AI algorithm. Like LinTS, TS-AI samples from the posterior distribution; however, it adapts the posterior variance based on the observed data, hence the acronym TS-AI, and only inflates it when additional exploration is necessary. This makes TS-AI less prone to over-exploration, while also eliminating the randomization bias that plagues LinTS. We establish that under additional assumptions, TS-AI achieves the frequentist minimax optimal regret. We also present numerical simulations that illustrate the limitations of LinTS, and how TS-AI addresses them. These simulations confirm the advantages of TS-AI and how it retains the benefits of LinTS while overcoming its limitations.

1.1. Other related literature

In the special case of standard MAB problem, TS has been extensively studied, and several research works have established regret bounds for TS that match the minimax lower bounds up to logarithmic factors. Agrawal and Goyal (2012) and Agrawal and Goyal (2013a) provide frequentist regret bounds for TS, while Bubeck and Liu (2013) demonstrate that TS attains optimal Bayesian regret up to constants. Recently, Jin et al. (2020) proposed a modified version of TS that achieves minimax-optimal frequentist regret up to constant terms. In this domain, Phan et al. (2019) and Nie et al. (2018) revealed interesting observations about TS that are related to our work. The former noted that sampling from an approximation of the posterior distribution with constant α -divergence approximation error could result in linear regret. The latter demonstrated that the estimates for the mean rewards have a downward bias when a wide range of bandit algorithms collect the samples. Additionally, the recent work of Bastani et al. (2019) provides a positive result for TS with misspecified prior in a dynamic pricing setting with a large number of parallel bandit problems.

1.2. Organization

We begin by introducing the notations and problem formulation in Section 2. In Section 3, we present two families of examples that demonstrate how LinTS without inflation can incur linear regret. Next, we propose our TS-AI algorithm and provide its theoretical analysis in Sections 4 and 6, and by empirical simulations in Section 5. Finally, we conclude in Section 7, and relegate the proofs to the appendices.

2. Setting and Notation

We begin by introducing the notations that will be used throughout the paper. For any positive integer n, we denote the set $1, 2, \dots, n$ as [n]. Let Σ be a positive semi-definite n by n matrix, and let A be any vector in \mathbb{R}^n . We define the notation $\|A\|_{\Sigma}$ for $\sqrt{A^\top \Sigma A}$. For a matrix \mathbf{M} with singular values $\sigma_1 \geq \dots \geq \sigma_n$, we define its operator norm as $\|\mathbf{M}\|_{\mathrm{op}} := \sigma_1$, and its trace norm (or nuclear norm) as $\|\mathbf{M}\|_* := \sum_{i \in [n]} \sigma_i$. To represent asymptotic upper and lower bounds, we use the standard notations $\mathcal{O}(\cdot)$ and $\Omega(\cdot)$, respectively. When logarithmic terms are suppressed, we use the notations $\widetilde{\mathcal{O}}(\cdot)$ and $\widetilde{\Omega}(\cdot)$, and defer a formal definition to Cormen et al. (2001). Lastly, we denote the cumulative distribution function (CDF) of the standard normal distribution by $\Phi(\cdot)$, and the n-dimensional identity matrix by \mathbb{I}_n .

Let $(\mathcal{A}_t)_{t=1}^T$ be a sequence of T random compact subsets of \mathbb{R}^d where $T \in \mathbb{N}$ is the time horizon. We further assume that $\|A\|_2 \leq \mathbf{a}$ for all $A \in \mathcal{A}_t$ almost surely. A policy π sequentially interacts with the environment in T rounds. At time $t \in [T]$, it receives action set \mathcal{A}_t and chooses an action $\widetilde{A}_t \in \mathcal{A}_t$ and receives a stochastic reward $Y_t = \left\langle \Theta^\star, \widetilde{A}_t \right\rangle + \varepsilon_t$ where ε_t is the reward noise and Θ^\star is an unknown (and potentially random)

vector of parameters. By $A_t^* \in \mathcal{A}_t$ we denote the arm with maximum expected reward. We denote the history of observations up to time t by \mathcal{F}_t . More precisely, we define

$$\mathcal{F}_t := (\mathcal{A}_1, \widetilde{A}_1, Y_1, \cdots, \mathcal{A}_{t-1}, \widetilde{A}_{t-1}, Y_{t-1}, \mathcal{A}_t).$$

In this model, a policy π is formally defined as a (stochastic) function that maps \mathcal{F}_t to an element of \mathcal{A}_t .

We compare policies through their cumulative Bayesian regret defined as

$$\operatorname{Regret}(T,\pi) := \sum_{t=1}^T \mathbb{E} \bigg[\sup_{A \in \mathcal{A}_t} \big\langle \Theta^\star, A \big\rangle - \big\langle \Theta^\star, \widetilde{A}_t \big\rangle \bigg].$$

Recall that the expectation is taken with respect to the entire three sources of randomness in our model, including the prior distribution on Θ^* . The frequentist regret bounds also follow by taking the prior distribution to be the measure that puts all the mass on a single deterministic vector.

3. Bayesian analyses are brittle

In this section, we demonstrate that LinTS may incur linear regret when the assumptions are *slightly* violated. Our analysis reveals that when LinTS employs an inaccurate prior or noise distribution, the Bayesian regret (and frequentist regret) can exhibit a linear growth rate. 2 . To be more specific, we establish that a linear regret can occur, when the dimensionality of the problem satisfies $d = \Omega(\log T)$. We begin by offering an intuitive explanation of these examples in Section 3.1, after which we present the examples in Sections 3.2 and 3.3. The former employs action sets that vary over time, while the latter employs fixed action sets.

3.1. Intuition

Here we construct a vanilla example where an adaptive adversary causes LinTS to fail by adaptively choosing bad action sets. But, this example is chiefly intended to develop the intuition behind our main examples, where *the action sets are selected independently from the history*. Rigorous proofs are provided for the examples in the next subsection.

Next, somewhat counter-intuitively, we first state a positive result, a sufficient condition that leads to a sub-linear regret bound for LinTS. Therefore, a counter-example in which LinTS would fail must violate that sufficient condition which gives us intuition on how to construct counter-examples. We also note that, this result is for the slightly more general version of LinTS where the posterior distribution is inflated by a positive parameter ι (Algorithm 1). We will prove a more general version of this theorem in Section B.

² This does not contradict the minimax optimal bound obtained by Russo and Van Roy (2014). Their analysis assumes that LinTS has access to the true prior and noise distributions, which is a stronger assumption than the one made here.

Algorithm 1 Linear Thompson sampling (LinTS)

Require: Inflation parameter ι .

1: Initialize $\Sigma_1 \leftarrow \lambda \mathbb{I}$ and $\widehat{\Theta}_1 \leftarrow 0$

2: **for** $t = 1, 2, \cdots$ **do**

3: Observe A_t

4: Sample $\widetilde{\Theta}_t \sim \mathcal{N}(\widehat{\Theta}_t, \, \iota^2 \mathbf{\Sigma}_t)$

5: $\widetilde{A}_t \leftarrow \arg\max_{A \in \mathcal{A}_t} \langle A, \widetilde{\Theta}_t \rangle$

6: Observe reward Y_t

7: $\Sigma_{t+1}^{-1} \leftarrow \Sigma_{t}^{-1} + \widetilde{A}_{t} \widetilde{A}_{t}^{\top}$

8: $\widehat{\Theta}_{t+1} \leftarrow \Sigma_{t+1} \left(\Sigma_t^{-1} \widehat{\Theta}_t + \widetilde{A}_t Y_t \right)$

9: end for

Theorem 3.1 If Algorithm 1 satisfies

$$\mathbb{P}\left(\sup_{A \in \mathcal{A}_t} \langle A, \widetilde{\Theta}_t \rangle \ge \sup_{A \in \mathcal{A}_t} \langle A, \Theta^* \rangle \,\middle|\, \Theta^*, \mathcal{F}_t \right) \ge \mathsf{p}\,,\tag{3.1}$$

whenever $\|\widehat{\Theta}_t - \Theta^{\star}\|_{\Sigma_t^{-1}} \leq \rho$, we then have

$$\operatorname{Regret}(T, \pi^{\operatorname{LinTS}}) \le \widetilde{\mathcal{O}}\left(\frac{\rho\iota}{\mathsf{p}}\sqrt{dT}\right).$$
 (3.2)

Recall that the optimal action at time t is denoted by A_t^{\star} . We can now write

$$\sup_{A \in \mathcal{A}_t} \langle A, \widetilde{\Theta}_t \rangle - \sup_{A \in \mathcal{A}_t} \langle A, \Theta^* \rangle \ge \langle A_t^*, \widetilde{\Theta}_t - \Theta^* \rangle.$$

Therefore, a sufficient condition for Eq. (3.1) is that

$$\mathbb{P}\left(\langle A_t^{\star}, \widetilde{\Theta}_t - \Theta^{\star} \rangle \ge 0 \mid \Theta^{\star}, \mathcal{F}_t\right) \ge \mathsf{p}\,,\tag{3.3}$$

whenever $\|\widehat{\Theta}_t - \Theta^{\star}\|_{\Sigma_t^{-1}} \leq \rho$. Next notice that

$$\langle A_t^{\star}, \widetilde{\Theta}_t - \Theta^{\star} \rangle = \langle A_t^{\star}, \widetilde{\Theta}_t - \widehat{\Theta}_t \rangle + \langle A_t^{\star}, \widehat{\Theta}_t - \Theta^{\star} \rangle. \tag{3.4}$$

Looking at the above decomposition, we call $E_t := \Theta^* - \widehat{\Theta}_t$ the *error vector* and $C_t := \widetilde{\Theta}_t - \widehat{\Theta}_t$ the *compensator vector*. The latter name is motivated by the observation that, using Eq. (3.3) and Eq. (3.4), when $\langle A_t^*, C_t \rangle \geq \langle A_t^*, E_t \rangle$ holds then

$$\sup_{A \in \mathcal{A}_t} \langle A, \widetilde{\Theta}_t \rangle \ge \sup_{A \in \mathcal{A}_t} \langle A, \Theta^{\star} \rangle$$

holds as well. Thus, C_t should *compensate* for the underestimation of $\langle A_t^*, \Theta^* \rangle$ caused by E_t . While this inequality is only a *sufficient* condition to obtain Eq. (3.1), we demonstrate how it can be used to deceive LinTS.

An adversary that knows $\widehat{\Theta}_t$ and Θ^* (thereby, E_t) can exploit LinTS by showing an action set of the form $\{0,A\}$ where A satisfies

$$\frac{1}{2}\langle A, E_t \rangle \approx -\langle A, \widehat{\Theta}_t \rangle \approx \langle A, \Theta^{\star} \rangle > 0 = \langle 0, \Theta^{\star} \rangle \,.$$

Therefore, A would be the optimal action (i.e., $A = A_t^*$). In this case, LinTS would choose A if and only if $\langle A, \widetilde{\Theta}_t \rangle > 0$, which would then be approximately equivalent to

$$\langle A, C_t \rangle \ge \frac{1}{2} \langle A, E_t \rangle$$
.

But A can be chosen to be align with $\langle A, E_t \rangle$ which would make it much larger than $\langle A, C_t \rangle$ because C_t is an independent random vector, conditioned on the history. This would allow A to be chosen so that $\langle A, E_t \rangle \approx \|A\|_2 \|E_t\|_2$, whereas $\langle A, C_t \rangle \approx \frac{1}{\sqrt{d}} \|A\|_2 \|C_t\|_2$ with high probability. Therefore, LinTS will select $\widetilde{A}_t = 0$ with a high probability while it is not the optimal arm. Moreover, $\widetilde{A}_t = 0$ reveals no more information about the true parameters. Hence, even if the same action set is shown in the next rounds, LinTS will fail to detect the optimal arm. In the remaining, we will make this intuition more formal.

3.2. Example 1: Noise reduction and changing action sets

In the previous section, we discussed how aligning the arm selection with the error vector can impact the performance. However, when the distributions used to compute the posterior distribution in LinTS do not match the actual distributions, a marginal bias can occur in the error vector E_t . This can have a negative impact on the performance of LinTS when the action set is appropriately chosen. Remarkably, this bias can even occur when the data quality is improved by reducing the noise variance. Importantly, the action sets can be constructed ahead of time without knowledge of the algorithm decisions. In this subsection, we describe our strategy for proving these results. First, we construct small problem instances in which $\widetilde{\Theta}_t$ is marginally biased. We then demonstrate that by combining independent copies of these biased instances, LinTS can incur linear Bayesian regret.

Remark 3.1 We study LinTS as shown in Algorithm 1 with $\iota = 1$ and $\lambda = 1$. This means we study LinTS that does not inflate posterior variance and assumes the noise variance and prior variance are equal to 1. In the example that will be constructed below, the true noise and prior variance will be τ^2 and σ^2 , respectively. Then, the main result of the section will show that when $\tau \neq \sigma$, LinTS achieves linear regret. But note that $\tau \neq \sigma$ means at least one of τ or σ is not equal to 1, which means at least one of noise or prior variance does not match the one that LinTS assumes.

Bias-introducing action sets. In this section, we construct an example in which Θ_t is marginally biased, provided that either the prior distribution or the noise distribution mismatches the one that LinTS uses. Fix $\sigma^2, \tau^2 \geq 0$ and let $\Theta^* \sim \mathcal{N}(0, \sigma^2 \mathbb{I}_2)$ be the vector of unobserved parameters. At time $t \in \{1, 2, 3\}$, we reveal the following action sets to the policy:

$$\mathcal{A}_t := egin{cases} \{e_1\} & ext{if } t = 1, \\ \{e_2\} & ext{if } t = 2, \\ \{e_1, e_2\} & ext{if } t = 3, \end{cases}$$

where e_1 and e_2 are standard basis vectors in \mathbb{R}^2 . For $t \leq 2$, LinTS has only one choice e_t and thus $\widetilde{A}_t = e_t$. Assume that reward $Y_t = \Theta_t^\star + \varepsilon_t$ is revealed to the algorithm where $\varepsilon_t \sim \mathcal{N}(0,\tau^2)$. At time t=3 for the first time, LinTS has two choices. Let i in [2] be such that $\widetilde{A}_3 = e_i$. Then, $Y_3 = \Theta_i^\star + \varepsilon_3$ is provided to the algorithm where $\varepsilon_3 \sim \mathcal{N}(0,\tau^2)$. The following key lemma proves that $\widehat{\Theta}_4$ is marginally biased when $\tau \neq \sigma$.

Lemma 3.1 Let $V = e_1 + e_2$. For any $\sigma, \tau \ge 0$, we have

$$\langle V, \mathbb{E}\left[\widehat{\Theta}_4\right] \rangle = \frac{(\sigma^2 - \tau^2) \beta}{6\sqrt{\sigma^2 + \tau^2 + 2}},$$
 (3.5)

where $\beta := \mathbb{E}[\max\{A, B\}] > 0$ with A and B being two independent standard normal random variables. Furthermore, for a positive constant C, $\widehat{\Theta}_4$ satisfies

$$\mathbb{E}\left[\exp\left(s\langle V,\widehat{\Theta}_4 - \mathbb{E}\left[\widehat{\Theta}_4\right]\rangle\right)\right] \le \exp\left[Cs^2(\sigma + \tau + \sqrt{2})^2\right],\tag{3.6}$$

for all $s \in \mathbb{R}$.

This finding illuminates that a bias emerges in the posterior mean estimate yielded by the LinTS algorithm due to a combination of two factors: distribution mismatch and randomization bias. The former refers to the discrepancy between LinTS's assumption on the true prior variance and the true noise variance, which is shown in Remark 3.1 to be equivalent to $\sigma^2 - \tau^2 \neq 0$. The latter stems from the randomization inherent in LinTS, resulting in the introduction of a positive term β .

We present a proof of Lemma 3.1 in Section A. However, we shall here provide a brief sketch of the proof. Firstly, we demonstrate that $\langle V, \widehat{\Theta}_4 \rangle$ can be expressed as a linear combination of $\langle V, \widehat{\Theta}_3 \rangle$, $\Theta_i^{\star} - \varepsilon_i$, and ε_3 . The first and third terms are unbiased, with a mean of zero, thus our attention is focused on $\mathbb{E}[\Theta_i^{\star} - \varepsilon_i]$. Subsequently, we establish that $\mathbb{E}[\Theta_i^{\star}]$ and $\mathbb{E}[\varepsilon_i]$ are equal to a shared constant, multiplied by $\sigma^2 \mathbb{E}[\widetilde{\Theta}_{3,i}]$ and $\tau^2 \mathbb{E}[\widetilde{\Theta}_{3,i}]$, respectively. Because the expected value of $\widetilde{\Theta}_{3,i}$ is proportional to β , it is now evident to trace back the origin of both $(\sigma^2 - \tau^2)$ and β in Eq. (3.5) to the distribution mismatch and randomization bias.

Stacking biased blocks. By combining independent copies of the above example, we prove that LinTS can choose an incorrect action for at least $\exp(\Omega(d))$ rounds. Let d be a positive integer and define $\Theta^* \sim \mathcal{P}_{\Theta^*} = \mathcal{N}(0, \sigma^2 \mathbb{I}_{2d})$. We will construct a 2d-dimensional linear bandit setting where in the first 3d rounds, the action sets of the type introduced above for each pairs $(\Theta^*_{2i-1}, \Theta^*_{2i})$ for $i \in [d]$ are presented to the algorithm. Namely, define

$$\mathcal{A}_{t} := \begin{cases}
\{e_{t}\} & \text{if } t \leq 2d, \\
\{e_{2(t-2d)-1}, e_{2(t-2d)}\} & \text{if } 2d+1 \leq t \leq 3d, \\
\{0, A\} & \text{otherwise,}
\end{cases}$$
(3.7)

where $A := (\operatorname{sgn}(\tau^2 - \sigma^2)/\sqrt{d}) \cdot \sum_{i=1}^{2d} e_i$. Note that, due to the term $\operatorname{sgn}(\tau^2 - \sigma^2)$, A is in the opposite direction of the marginal bias of $\widehat{\Theta}_{3d+1}$. Therefore, LinTS will be less likely to select A, and this will be an incorrect decision. Formally, the following key lemma, proved in Section A, states that with constant probability, A is the optimal action, while LinTS *perceives* it as suboptimal, with an enormous gap.

Lemma 3.2 For positive constants $p_0 = \frac{1}{2}(1 - \Phi(1))$ and $C_1(\sigma, \tau) := \frac{|\sigma^2 - \tau^2|\beta}{6\sqrt{\sigma^2 + \tau^2 + 2}}$, the following holds

$$\mathbb{P}\Bigg(\big\langle \Theta^{\star}, A \big\rangle \geq \sqrt{2}\sigma \ \ \text{and} \ \ \big\langle \widehat{\Theta}_{3d+1}, A \big\rangle \leq -\frac{C_1(\sigma, \tau)\sqrt{d}}{2} \Bigg) \geq p_0 \,.$$

We denote the event in the above Lemma by \mathcal{B} . Conditional on this event, for all t>3d, the optimal arm is A, and the regret incurred by choosing the action 0 is at least $\sqrt{2}\sigma$. Moreover, let q be the probability of choosing A at t=3d+1. As we will see, when $\tau \neq \sigma$, this probability is exponentially small as a function of d. The probability of selecting A in the next round remains unchanged, whenever A is not chosen. This observation holds up to the first time that A is picked, which can, in turn, take an exponentially long time. By making this argument rigorous, we can state the following proposition which is proved in Section A.

Proposition 3.1 In Example 1, when $\sigma \neq \tau$ and $T \leq \exp(\Omega(d))$, we have $\operatorname{Regret}(T, \pi^{\operatorname{LinTS}}) \geq \Omega(T)$.

An immediately corollary of this result is that when d is comparable to $\log T$, which can naturally occur in practice, LinTS incurs a linear regret.

Corollary 3.1 *In Example 1, when* $d = \Omega(\log T)$ *and* $\sigma \neq \tau$ *, we have*

$$\operatorname{Regret}(T, \pi^{\operatorname{LinTS}}) \ge \Omega(T)$$
.

Drawing upon Remark 3.1, the aforementioned result underscores that a discrepancy between the actual noise or prior distribution and those which LinTS presumes, leads to linear regret. Interestingly, a specific instance of this circumstance arises when $\tau = 0$, thereby highlighting a scenario in which LinTS falters despite being provided with superior-quality data than it assumes.

3.3. Example 2: Mean shift and fixed action sets

In this subsection, we construct an example in which LinTS incurs linear Bayes regret while the action set is fixed over time. Like the previous example, we assume LinTS does not inflate posterior variance and assumes the noise and prior distribution are both standard normal. Let $\mu, \sigma, \tau > 0$ be fixed, and for $d \in \mathbb{N}$, set the prior distribution to be $\mathcal{P}_{\Theta^*} := \mathcal{N}(\mu \mathbf{1}_{3d}, \sigma^2 \mathbb{I}_{3d})$. We now reveal the action set $\mathcal{A}_t := \{0, A', A\}$ to LinTS for all $t \in [T]$ where

$$A' := -\frac{1}{\sqrt{d}} \sum_{i=1}^{d} e_i,$$

$$A := \frac{1}{\sqrt{d}} \sum_{i=d+1}^{3d} e_i - \frac{1}{\sqrt{d}} \sum_{i=1}^{d} e_i.$$
(3.8)

The next proposition, proved in Section A, highlights the key observations about why LinTS fails in this simple setting.

Proposition 3.2 For fixed $\mu, \sigma > 0$, and for sufficiently large d, we have

- 1. $\langle \Theta^*, A' \rangle \leq -\frac{1}{2}\mu\sqrt{d} \leq \frac{1}{2}\mu\sqrt{d} \leq \langle \Theta^*, A \rangle$, with probability at least $\frac{7}{8}$.
- 2. $\widetilde{A}_1 = A'$ with probability $\frac{1}{4}$.
- 3. Conditional on $\widetilde{A}_1 = A'$, $\langle \widehat{\Theta}_2, A \rangle$ and $\langle \widehat{\Theta}_2, A' \rangle$ are less than $-\frac{1}{8}\mu\sqrt{d}$, with probability at least $\frac{15}{16}$.
- 4. Conditional on $\widetilde{A}_1 = A'$, $\widetilde{A}_2 \neq 0$ with probability at most $\exp(-\Omega(d))$.
- 5. For all $T \leq \exp(\Omega(d))$, Regret $(T, \pi^{\text{LinTS}}) \geq \Omega(T\sqrt{d})$.

Remark 3.2 One can slightly modify the proof to obtain a similar result for $\mathcal{P}_{\Theta^*} := \mathcal{N}(0, \sigma^2 \mathbb{I}_{3d} + \rho \mathbf{1}_{3d} \mathbf{1}_{3d}^\top)$. It is easy to see that for any arbitrary constant ρ , the same rate as in Eq. (A.12) is achievable. Also, for $\rho = d^{-\alpha}$ where $\alpha < 1$, one can still get non-trivial results.

An immediate implication of Proposition 3.2 and Remark 3.2 is that if there exists a mismatch between LinTS and the true prior at the mean or variance level in Example 2 with a fixed action set, then LinTS incurs linear regret for an extended exponential period. Similar to Example 1, we obtain the following corollary.

Corollary 3.2 In Example 2, when $d = \Omega(\log T)$ and the true prior \mathcal{P}_{Θ^*} is either $\mathcal{N}(\mu \mathbf{1}_{3d}, \sigma^2 \mathbb{I}_{3d})$ or $\mathcal{P}_{\Theta^*} = \mathcal{N}(0, \sigma^2 \mathbb{I}_{3d} + \rho \mathbf{1}_{3d} \mathbf{1}_{3d}^\top)$, we have

Regret
$$(T, \pi^{\text{LinTS}}) \ge \Omega(T)$$
.

In summary, we have presented a simple setting in which LinTS incurs linear Bayes regret even when the action set is fixed over time.

Remark 3.3 It is worth noting that the design of the action sets in Example 1 (or Example 2) necessitates solely the understanding of the sign of $\tau^2 - \sigma^2$ (or sign of μ). In the context of a competition between two competiting platforms \mathcal{P}_1 and \mathcal{P}_2 from Section 1, \mathcal{P}_1 only needs to be aware of the direction of inconsistency between the true reward distribution and the one which is presumed by \mathcal{P}_2 .

4. Thompson Sampling with Adaptive Inflation

In this section, we present an alternative approach to improve the inflation parameter in LinTS and enhance its performance, subject to additional assumptions. To facilitate a better understanding of our proposed method, we first provide the intuition behind the development of the conditions. These intuitions follow the discussion in Section 3.1, with a focus on exploring the mechanisms that enable LinTS to succeed, rather than those that may cause it to fail. Building on these insights, we introduce Thompson Sampling with Adaptive Inflation (TS-AI), an algorithm that adaptively adjusts its inflation parameter to meet the aforementioned conditions. Finally, we state our informal regret bound for TS-AI and defer its formal proof to Section B.

Note that at any time in LinTS, the posterior mean $\widehat{\Theta}_t$ is the ridge estimator for the parameter Θ^* , given the actions and their observed rewards in prior rounds. Additionally, we define \mathcal{C}_t as the confidence set centered around $\widehat{\Theta}_t$, which is constructed as part of the OFUL algorithm and contains $\widetilde{\Theta}_t$ and Θ^* with high probability. More information on the construction and properties of these confidence sets can be found in (Abbasi-Yadkori et al. 2011).

Assume that $d = \Omega(\log T)$. As in Section 3.1, and for the sake of building intuition, we first restrict our attention to action sets of the form $\{A_t^{\star}, 0\}$ where A_t^{\star} is the optimal arm, i.e., $\langle \Theta^{\star}, A_t^{\star} \rangle > 0$. LinTS chooses A_t^{\star} only if

$$\langle \widetilde{\Theta}_t, A_t^{\star} \rangle > 0.$$
 (4.1)

Compensation inequality. By decomposing the left-hand side of Eq. (4.1) as we did in Eq. (3.4), a sufficient condition for Eq. (4.1) to hold is

$$\langle \widetilde{\Theta}_t - \widehat{\Theta}_t, A_t^* \rangle \ge \langle \Theta^* - \widehat{\Theta}_t, A_t^* \rangle,$$
 (4.2)

which is also equivalent to $\langle C, A_t^{\star} \rangle \geq \langle E, A_t^{\star} \rangle$ with C and E defined in Section 3.1 and illustrated in Figure 1.

OFUL explicitly seeks $\widetilde{\Theta}_t \in \mathcal{C}_t$ that maximizes the left-hand side of Eq. (4.2), and as $\Theta^* \in \mathcal{C}_t$ with high probability, the desired "compensation inequality" holds, and A_t^* is selected with high probability. LinTS, on the other hand, follows a stochastic approach and resorts to a randomly sampled $\widetilde{\Theta}_t$, that with high probability resides in \mathcal{C}_t , to solve Eq. (4.2). Since $\widehat{\Theta}_t$ is the ridge estimator for the collected data thus far, in a fixed




Figure 1 A typical setting for A_t^{\star} , Θ^{\star} , $\widehat{\Theta}_t$, and $\widetilde{\Theta}_t$. The compensation vector $C = \widetilde{\Theta}_t - \widehat{\Theta}_t$ and the error vector $E = \Theta^{\star} - \widehat{\Theta}_t$ are defined in Section 3.1, and O denotes the origin.

design setting (which is not true in our bandit problem with adaptively collected data) the error vector E will be pointing in a random direction. Therefore, provided that A_t^* is independent of E, we have

$$\left|\left\langle E, A_t^{\star} \right\rangle\right| \approx \mathcal{O}\left(\frac{1}{\sqrt{d}} \|E\|_2 \cdot \|A_t^{\star}\|_2\right).$$
 (4.3)

The same expression also holds for $|\langle C, A_t^\star \rangle|$; therefore, the compensation inequality holds with constant probability. To summarize our observation, Eq. (4.3) holds if the error vector E is distributed in a random direction that is independent of the optimal action A_t^\star . The crucial point in the analysis of LinTS in the Bayesian setting is that the error vector is in a random direction whenever LinTS has access to the true prior and noise distribution. In Section 3, nonetheless, we have shown that this condition is violated if LinTS uses an incorrect prior or noise distribution in computing the posterior. Agrawal and Goyal (2013b), Abeille et al. (2017) take a conservative approach and propose to inflate the posterior distribution by a factor of $\widetilde{\mathcal{O}}(\sqrt{d})$ that inflates C by the same factor to ensure $\langle C, A_t^\star \rangle \geq \langle E, A_t^\star \rangle$ holds with constant probability. We present an alternative approach that leverages the randomness of the optimal action to reduce the need for exploration. The following assumption requires the optimal arm (rather than the error vector) to be distributed in a random direction.

Assumption 4.1 Assuming that for any $V \in \mathbb{R}^d$ with $\|V\|_2 = 1$, the inequality

$$\langle A_t^{\star}, V \rangle \le \frac{\nu}{\sqrt{d}} \|A_t^{\star}\|_2 \tag{4.4}$$

holds with a probability of at least $1 - \frac{1}{T^2}$, where ν is a fixed parameter.

Unfortunately, this condition alone does not suffice to reduce the inflation rate of the posterior distribution. To see this, consider a case in which the largest eigenvalue of Σ_t is much larger than the other eigenvalues of

 Σ_t ; thereby, $\|\Sigma_t\|_{\text{op}} \approx \|\Sigma_t\|_*$ and E points to the longest direction of the confidence set. Figure 2 illustrates this situation. In this case, we have

$$\begin{split} \left| \left\langle E, A_t^{\star} \right\rangle \right| &\approx \frac{\left\| E \right\|_2 \cdot \|A_t^{\star}\|_2}{\sqrt{d}} \\ &\approx \frac{\sqrt{d} \left\| \boldsymbol{\Sigma}_t \right\|_{\text{op}} \cdot \|A_t^{\star}\|_2}{\sqrt{d}} \\ &= \sqrt{\left\| \boldsymbol{\Sigma}_t \right\|_{\text{op}} \cdot \left\| A_t^{\star} \right\|_2} \,. \end{split}$$

The second approximation utilizes two observations. First, it exploits the fact that $||E||_2 \approx ||E||_{\Sigma_t^{-1}} \cdot ||E||_{\Sigma_t}$, which is justified by the alignment of E with the longest direction of Σ_t . Second, it relies on the high probability event that E is contained in C_t , implying that $||E||_{\Sigma_t^{-1}}$ is of the order \sqrt{d} .




Figure 2 A thin confidence set.

However, it follows from the definition of LinTS that $\langle \widetilde{\Theta}_t - \widehat{\Theta}_t, A_t^\star \rangle \sim \mathcal{N} \left(0, \iota^2 \| A_t^\star \|_{\Sigma_t}^2 \right)$. Assuming that $\mathbb{E}[A_t^\star A_t^{\star \top}] \approx \mathbb{I}_d$, we realize that $\mathbb{E} \left[\| A_t^\star \|_{\Sigma_t}^2 \right] \approx \| \Sigma_t \|_*$. This suggests $\| A_t^\star \|_{\Sigma_t}$ is proportional to

$$\sqrt{\|\mathbf{\Sigma}_t\|_*/d} \cdot \|A_t^{\star}\|_2$$
.

Now, we can see that Assumption 4.1 is not sufficient for ensuring Eq. (4.2) as we have

$$\left|\left\langle E, A_t^{\star} \right\rangle\right| \approx \sqrt{\|\boldsymbol{\Sigma}_t\|_{\text{op}}} \cdot \|A_t^{\star}\|_2 \gg \sqrt{\frac{\|\boldsymbol{\Sigma}_t\|_*}{d}} \cdot \|A_t^{\star}\|_2 \approx \left|\left\langle C, A_t^{\star} \right\rangle\right|.$$

This observation implies the necessity of the inflation rate of $\widetilde{\mathcal{O}}(\sqrt{d})$ when the eigenvalues of Σ_t differ in magnitude significantly. To make this notion precise, we define the *thinness coefficient* of a positive definite matrix Σ to be

$$\psi(\mathbf{\Sigma}) := \sqrt{\frac{d \cdot \|\mathbf{\Sigma}\|_{\mathrm{op}}}{\|\mathbf{\Sigma}\|_{*}}}.$$

The following assumption requires A_t^{\star} to be distributed in a way that benefits from low thinness.

Assumption 4.2 For $\Psi, \omega > 0$, we have

$$\|A_t^{\star}\|_{\Sigma} \ge \omega \sqrt{\frac{\|\Sigma\|_*}{d}} \cdot \|A_t^{\star}\|_2, \tag{4.5}$$

with probability at least $1 - \frac{1}{T^2}$, for any positive semi-definite matrix Σ with $\psi(\Sigma) \leq \Psi$.

Assumptions 4.1 and 4.2 are sufficient for reducing the inflation parameter, whenever $\psi(\Sigma_t) \leq \Psi$. In the following theorem, we state our regret bound informally. The formal version of this result (Theorem B.1) and its proof can be found in Section B.

Theorem 4.1 (Informal) Assume that $\langle A, \Theta^* \rangle \in [-1, 1]$ for all $A \in \mathcal{A}_t$ almost surely. Then, under Assumptions 4.1 and 4.2, we have

$$\sum_{t=1}^{T} \langle A_t^{\star}, \Theta^{\star} \rangle - \langle \widetilde{A}_t, \Theta^{\star} \rangle \leq \widetilde{\mathcal{O}}(\iota d\sqrt{T}) + 2 \sum_{t=1}^{T} \mathbb{I}(\psi(\Sigma_t) > \Psi),$$

with probability at least $1 - \frac{3}{T}$, provided that the inflation prameter ι of LinTS satisfies $\iota \geq \frac{\nu \Psi}{\omega} \cdot \widetilde{\mathcal{O}}(1)$.

In Section 5, for a concrete example from (Russo and Van Roy 2014), we empirically show that $\psi(\Sigma_t) < \Psi$ holds for a small value of Ψ with high probability in our simulations. However, this condition is *not* a mere property of the environment and depends on the interactions of LinTS with the environment. Nonetheless, notice that $\psi(\Sigma_t)$ is observable, and the policy can intervene if $\psi(\Sigma_t) > \Psi$ for many rounds. This is the main idea behind our TS-AI algorithm presented in Algorithm 2. Note that the parameter ρ_t , formally defined in Eq. (5.1), plays the role of the factor \sqrt{d} inflation as in (Agrawal and Goyal 2013b) and (Abeille et al. 2017). However, in TS-AI, such inflation is only performed when the adaptively calculated thinness parameter $\psi(\Sigma_t)$ is too large. Otherwise, a constant inflation parameter ι is used.

For the same example from (Russo and Van Roy 2014), in Section 6, we will theoretically prove that Assumptions 4.1 and 4.2 hold.

Remark 4.1 (OFUL with smaller confidence intervals) Our proof in Section B reveals that, under Assumptions 4.1 and 4.2, it is possible to improve the performance of the OFUL algorithm by running it with smaller confidence sets that are reduced by a factor of order \sqrt{d} . It is worth noting that while the reduced confidence sets only impact the constants in the regret bound, but they may lead to improve empirical performance of the OFUL algorithm.

Remark 4.2 (Towards relaxing Assumptions 4.1 and 4.2) It is noteworthy that the Assumptions 4.1 and 4.2 primarily serve to facilitate the theoretical analysis. From the proof of Lemma B.1, one requires the inflation parameter ι to satisfy

$$\iota \ge \sup_{V \in \mathbb{S}^{d-1}} \frac{\langle A_t^*, V \rangle \rho \sqrt{\|\Sigma_t\|_{\text{op}}}}{\|A_t^*\|_{\Sigma_t}},\tag{4.6}$$

with a probability of at least $1-T^{-2}$, where \mathbb{S}^{d-1} denotes the d-dimensional unit sphere. Hence, for problems where Assumptions 4.1 and 4.2 may not hold, a data-driven approach to setting the inflation parameter is to follow Eq. (4.6), provided that the structure of the problem allows to bound the supremum on the right-hand side of Eq. (4.6).

Algorithm 2 Thompson Sampling with Adaptive Inflation (TS-AI)

Require: Inflation parameter ι and thinness threshold Ψ .

```
1: Initialize \Sigma_1 \leftarrow \lambda \mathbb{I} and \widehat{\Theta}_1 \leftarrow 0
  2: for t = 1, 2, \cdots do
  3:
              Observe A_t
              if \psi(\Sigma_t) > \Psi then
  4:
                    Sample \widetilde{\Theta}_t \sim \mathcal{N}(\widehat{\Theta}_t, \, \rho_t^2 \, \mathbf{\Sigma}_t) where \rho_t is defined in Eq. (5.1)
  5:
              else
  6:
                   Sample \widetilde{\Theta}_t \sim \mathcal{N}(\widehat{\Theta}_t, \, \iota^2 \mathbf{\Sigma}_t)
  7:
              end if
  8:
             \widetilde{A}_t \leftarrow \arg\max_{A \in A_t} \langle A, \widetilde{\Theta}_t \rangle
              Observe reward Y_t
             \Sigma_{t+1}^{-1} \leftarrow \Sigma_{t}^{-1} + \widetilde{A}_{t} \widetilde{A}_{t}^{\top}
11:
             \widehat{\Theta}_{t+1} \leftarrow \Sigma_{t+1} \left( \Sigma_t^{-1} \widehat{\Theta}_t + \widetilde{A}_t Y_t \right)
12:
13: end for
```

5. Simulations

In this section, we first, in Section 5.1, provide a numerical validation for the examples in Section 3 that demonstrate two scenarios under which LinTS fails to choose the best action for an exponentially long time horizon. Then, in Section 5.2, we compare the performance of our TS-AI with Bayesian and frequentist LinTS in different settings.

5.1. Average failure time of LinTS

We provide two sets of simulations to validate the theoretical predictions of the two examples in Section 3.

Noise reduction example. In this simulation, for each $d \in \{2, 2^2, 2^3, \cdots, 2^{18}\}$, we generate $\Theta^* \sim \mathcal{N}(0, \mathbb{I}_{2d})$ and execute LinTS for 3d rounds using the action sets in Eq. (3.7). The reward for choosing an action $\widetilde{A}_t \in \mathcal{A}_t$ is simply given by $Y_t = \langle \Theta^*, \widetilde{A}_t \rangle$. Therefore, no noise is added to the reward (i.e., $\tau = 0$). After executing LinTS for 3d rounds and obtaining $\widehat{\Theta}_{3d+1}$, we compute the probability that $\langle \widehat{\Theta}_{3d+1}, A \rangle > 0$. Note that we can calculate this probability given that $\widehat{\Theta}_{3d+1}$ is Gaussian and A is a multiple of $\mathbf{1}_{2d}$. Also, recall from Section 3.2 that, the complement of this event is when LinTS incorrectly chooses action 0, and under that scenario, the probability of selecting A in the next round stays the same. Hence, LinTS would be expected to chose 0 for 1/p time periods. Since 0 is suboptimal with probability 1/2, this means LinTS would be expected to choose the suboptimal arm 1/(2p) time periods. We repeat this procedure 100 times to obtain 100 values for p, denoted by $(p_i)_{i=1}^{100}$, and present a boxplot for the values $(1/(2p_i))_{i=1}^{100}$ in Figure 3(a), indicating




Figure 3 Boxplots of $1/p_i$'s in Examples 1 and 2.

the expected number of failures of LinTS versus d. It is evident that, as predicted in Section 3.2, LinTS selects the suboptimal action for at least $\exp(\Omega(d))$ rounds.

Fixed action set example. For given d and μ , we sample $\Theta^* \sim \mathcal{N}(\mu \mathbf{1}_{3d}, \mathbb{I}_{3d})$. Then, we reveal the action set $\mathcal{A}_t = \{0, A, A'\}$ as defined in Eq. (3.8). Then, conditional on $\widetilde{A}_t = A'$, we compute the probability p that the next arm is not 0. Also, recall from Section 3.3 that, under complement of this event, when LinTS incorrectly chooses action 0, this probability does not change in the next round. Hence, LinTS would be expected to fail for 1/p time periods. We repeat this process 100 times to get $(p_i)_{i=1}^{100}$, and as before, we show a boxplot for each value of the varying variable. Figure 3(b) shows the boxplots of $1/p_i$ for $\mu = 0.1$ when d varies between 1 to 2^{18} , and Figure 3(c) illustrates $1/p_i$ for d = 1000 and μ varying between 0 and 1. The results validate the theoretical analysis of Section 3.

5.2. Thinness over time and TS-AI

In this subsection, we present two sets of simulations. Firstly, we investigate the variation of the thinness parameter over time in Section 4 and then compare the performance of TS-AI with both the Bayesian and frequentist versions of LinTS in two different scenarios. The first scenario is a "well-behaved" setting where the Bayesian LinTS does not fail. The second scenario, known as Examples 1 and 2 in Section 3, are the brittle settings where Bayesian LinTS fails.

Scenario I. We consider a setting similar to the simulations section of (Russo and Van Roy 2014). Specifically, for d=50, we generate the parameter vector Θ^* from the standard normal distribution $\mathcal{N}(0,10\mathbb{I}_d)$. At each time step t, we generate k=100 independent vectors from the uniform distribution on the hypercube $[-1/\sqrt{d},1/\sqrt{d}]^d$, which form the action set. We compare the following policies:

- 1. TS-Bayes: Algorithm 1 with no inflation ($\iota = 1$).
- 2. TS-Freq: Algorithm 1 with $\iota = \rho_t$ at time t. This is the version considered by Agrawal and Goyal (2013b) and Abeille et al. (2017).
 - 3. TS-AI: Algorithm 2 with $\iota = 5$ and $\Psi = 2.0$.

Note that for both TS-Freq and TS-AI, we use

$$\rho_t := \sqrt{2\log\left(\frac{\det(\mathbf{\Sigma}_t^{-1})^{\frac{1}{2}}\det(0.1\mathbb{I}_d)^{-\frac{1}{2}}}{0.0001}\right)} + \sqrt{d}.$$
 (5.1)

Each policy chooses \widetilde{A}_t^π for $\pi \in \{\text{TS-Bayes, TS-Freq, TS-AI}\}$, and receives feedback $Y_t^\pi = \left\langle \Theta^\star, \widetilde{A}_t^\pi \right\rangle + \varepsilon_t^\pi$




Figure 4 Scenario I: Thinness and cumulative regret of TS-AI versus Bayesian and Frequentist versions of LinTS in a well-behaved setting.

where ε_t^{π} are i.i.d. standard Gaussian random variables. Next, we compute the thinness parameter for $\Sigma_t^{\pi} = \mathbb{I}/10 + \sum_{j=1}^t \widetilde{A}_j^{\pi} \widetilde{A}_j^{\pi\top}$. We repeat this procedure 20 times. Figure 4(a) displays the thinness of these policies in our experiments. This in particular shows that the thinness stays close to 1 for larger values of t. In other words, the term $\sum_{t=1}^T \mathbb{I}(\psi(\Sigma_t) > \Psi)$ as it appears in Theorem 4.1 (and its formal version, Theorem B.1) is zero for $\Psi < 2$ with high probability. Figure 4(b) shows the cumulative regrets of these policies. Notice that, while TS-AI may inflate the posterior variance by ρ_t , its performance is closer to TS-Bayes than TS-Freq given that the decision to inflate is performed in a more data-driven fashion.

Scenario II. Here, we consider the settings of Examples 1 and 2 from Section 3. For Example 1, we choose d=90 and the only difference is that we select $\sigma^2=1$ while $\tau^2=2$. For Example 2, we choose d=30, there is no variance mismatch ($\sigma^2=\tau^2=2$), but there is mean mismatch (prior mean is 10 while LinTS assumes prior mean is 0). In each case, we repeat the simulation 100 times and show the average regrets, with shaded error bars representing two standard errors, in Figure 5. In both cases, TS-Bayes, as predicted performs poorly but TS-AI nearly ties or outperforms TS-Freq, benefiting from the adaptive inflation.




Figure 5 Scenario II: Thinness and cumulative regret of TS-AI versus Bayesian and Frequentist version of LinTS in the setting of Examples 1 and 2 from Section 3.

6. Justifying Assumptions 4.1 and 4.2 in a Concrete Example

In this section, we prove that parameters ν and ω in Assumptions 4.1 and 4.2 are constants, in the specific example from Russo and Van Roy (2014) that was empirically studied in Section 5.2.

Proofs of Lemmas 6.1 and 6.2 are given in Section C.

Let k be the number of actions at each round. We first start by verifying Assumption 4.2.

Lemma 6.1 Let A be sampled from $\mathrm{Unif}([-1/\sqrt{d},1/\sqrt{d}]^d)$. Then, for any positive definite Σ with $\psi(\Sigma) \leq \Psi$, we have that

$$\mathbb{P}\bigg(\Big|\|A\|_{\mathbf{\Sigma}}^2 - \frac{1}{3d}\|\mathbf{\Sigma}\|_*\Big| \le \frac{1}{6d}\|\mathbf{\Sigma}\|_*\bigg) \le 2\exp\bigg(-\frac{cd}{\Psi^2}\bigg)$$

where c is an absolute constant.

The following corollary is a direct consequence of the above lemma combined with the union bound.

Corollary 6.1 Let A_t^* be the optimal action at time t. Then, for any positive definite Σ with $\psi(\Sigma) \leq \Psi$, we have that

$$\mathbb{P}\left(\left|\|A_t^{\star}\|_{\boldsymbol{\Sigma}}^2 - \frac{1}{3d}\|\boldsymbol{\Sigma}\|_{*}\right| \leq \frac{1}{6d}\|\boldsymbol{\Sigma}\|_{*}\right) \leq 2k \cdot \exp\left(-\frac{cd}{\Psi^2}\right).$$

Specifically, by setting $\Sigma := \mathbb{I}_d$ *, we get that*

$$\mathbb{P}\left(\|A_t^{\star}\|_2^2 \le \frac{1}{6} \quad or \quad \|A_t^{\star}\|_2^2 \ge \frac{1}{2}\right) \le 2k \cdot \exp\left(-cd\right),\tag{6.1}$$

and hence

$$\mathbb{P}\left(\|A_t^{\star}\|_{\Sigma}^2 \ge \frac{1}{3d}\|\Sigma\|_* \cdot \|A_t^{\star}\|_2^2\right) \le 4k \cdot \exp\left(-cd\right),\tag{6.2}$$

Our next lemma asserts that each action satisfies Assumption 4.1 with a constant parameter ν .

Lemma 6.2 Let A be chosen according to $\mathrm{Unif}([-1/\sqrt{d},1/\sqrt{d}]^d)$. Then, for any $V \in \mathbb{R}^d$ and $p \in (0,1)$, we have

$$\mathbb{P}\left(\langle A, V \rangle > \sqrt{\frac{2\log(1/p)}{d}}\right) \le p.$$

Furthermore, using Eq. (6.1), we get that

$$\mathbb{P}\left(\langle A, V \rangle > \sqrt{\frac{12\log(1/p)}{d}} \cdot ||A||_2\right) \le p + 2k \cdot \exp\left(-cd\right).$$

By applying the union bound, we can obtain the following result for the optimal arm. This corollary directly follows from the previous lemma and the application of the union bound.

Corollary 6.2 Let A_t^* be the optimal arm at time t. Then, for any $V \in \mathbb{R}^d$, we have that

$$\mathbb{P}\Bigg(\langle A_t^\star, V \rangle > \sqrt{\frac{12 \log(2kT^2)}{d}} \cdot \|A_t^\star\|_2 \Bigg) \leq \frac{1}{2T^2} + 2k \cdot \exp\left(-cd\right) \,.$$

This means, if d is larger than $\Psi^2 \log(8kT^2)/c$, Assumptions 4.1 and 4.2 are satisfied with

$$\nu := \sqrt{12\log(2kT^2)}$$
 and $\omega := \frac{1}{\sqrt{3}}$.

7. Conclusion and Discussion

This paper focuses on the stochastic linear bandit problem and investigates the Linear Thompson Sampling (LinTS) algorithm. Our goal is to determine if the factor d inflation in the posterior variance of LinTS, necessary to achieve the best-known frequentist regret bound, is essential. By settling an open problem, we show that the factor d inflation is indeed necessary and that the frequentist regret bound of $\widetilde{\mathcal{O}}(d\sqrt{dT})$ is optimal. Additionally, we demonstrate that more data-driven versions of LinTS, which use the observed data to adjust the posterior inflation, can achieve frequentist minimax optimal regret under additional conditions.

While our main results are theoretical, our paper provides insights into the performance of LinTS and identifies potential sources of degradation that may be of interest for practitioners. Our analysis highlights that a even a small mismatch between the prior and true distributions can lead to suboptimal performance, which supports prior literature and emphasizes the importance of careful prior distribution selection. Furthermore, we find that the randomization bias arising from inherent "sampling" nature of the algorithm can be a potential source of degradation in the performance of LinTS. Further understanding the impact of this bias is an intriguing topic for future research, and we hope that our findings will encourage further exploration in this area.

Appendix A: Proofs of Section 3

Prior to commencing the proof, we shall present several fundamental definitions and properties concerning Gaussian and sub-Gaussian random variables. A more thorough discussion on this topic can be found in (Vershynin 2018).

The sub-Gaussian norm of a random variable X by

$$||X||_{\psi_2} = \inf\left\{t > 0: \mathbb{E}\left[e^{\left(\frac{X}{t}\right)^2}\right] \le 2\right\}.$$

For every sub-Gaussian random variable X, there exist positive constants C_{tail} and C_{mgf} , such that

$$\mathbb{P}(X \ge |t|) \le 2 \exp\left(-\frac{C_{\text{tail}} t^2}{\|X\|_{\psi_2}^2}\right). \tag{A.1}$$

$$\mathbb{E}[\exp\left(s(X - \mathbb{E}[X])\right)] \le \exp\left(C_{\text{mgf}} s^2 \|X\|_{\psi_2}^2\right), \quad \text{for all } s \in \mathbb{R}, \tag{A.2}$$

and the Gaussian distribution satisfies

$$\|\mathcal{N}(0,\sigma^2)\|_{\psi_2} \le 2\sigma\,,\tag{A.3}$$

$$\mathbb{P}\left(\mathcal{N}(0,\sigma^2) \ge |t|\right) \le \frac{1}{\sqrt{2\pi}} \frac{\sigma}{|t|} \exp\left(-\frac{t^2}{2\sigma^2}\right). \tag{A.4}$$

We also need the following proposition that is proved in Section C.

Proposition A.1 (Bias decomposition) Let $(X_i)_{i=1}^n$ be a sequence of independent random variables where $X_i \sim \mathcal{N}(0, \sigma_i^2)$. By Y, we denote their sum and let Z be any independent random variable. Then, for any function $g: \mathbb{R} \times \mathbb{R} \to \mathbb{R}$, we have

$$\mathbb{E}[X_i \cdot g(Y, Z)] = \frac{\sigma_i^2}{\sum_{i=1}^n \sigma_i^2} \cdot \mathbb{E}[Y \cdot g(Y, Z)].$$

Proof of Lemma 3.1. Recall that, as stated in Remark 3.1, for notation simplicity, $\lambda = 1$. It follows from the definition of $\widehat{\Theta}_3$ that

$$\widehat{\Theta}_3 = \frac{1}{2} \begin{bmatrix} Y_1 \\ Y_2 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} \Theta_1^{\star} + \varepsilon_1 \\ \Theta_2^{\star} + \varepsilon_2 \end{bmatrix}.$$

Next, at t = 3, the *i*-th entry is updated according to

$$\begin{split} \widehat{\Theta}_{4,i} &= \frac{Y_i + Y_3}{3} \\ &= \frac{2\Theta_i^{\star} + \varepsilon_i + \varepsilon_3}{3} \\ &= \frac{\Theta_i^{\star} + \varepsilon_i}{2} + \frac{\Theta_i^{\star} - \varepsilon_i}{6} + \frac{\varepsilon_3}{3} \\ &= \widehat{\Theta}_{3,i} + \frac{\Theta_i^{\star} - \varepsilon_i}{6} + \frac{\varepsilon_3}{3}. \end{split}$$

Moreover, the other entry remains unchanged. In other words,

$$\widehat{\Theta}_{4,3-i} = \widehat{\Theta}_{3,3-i} .$$

Therefore, setting $V = e_1 + e_2$, we have

$$\begin{split} \left\langle \widehat{\Theta}_{4}, V \right\rangle &= \left\langle \widehat{\Theta}_{4}, V \right\rangle \\ &= \widehat{\Theta}_{4,1} + \widehat{\Theta}_{4,2} \\ &= \widehat{\Theta}_{3,1} + \widehat{\Theta}_{3,2} + \frac{\Theta_{i}^{\star} - \varepsilon_{i}}{6} + \frac{\varepsilon_{3}}{3}. \end{split} \tag{A.5}$$

and in particular

$$\langle \mathbb{E}\left[\widehat{\Theta}_{4}\right], V \rangle = \mathbb{E}\left[\frac{\Theta_{i}^{\star} - \varepsilon_{i}}{6}\right].$$
 (A.6)

We can now compute this expression in terms of the randomization bias coefficient given by

$$\beta := \mathbb{E}[\max\{A,B\}] > 0,$$

where A and B are two independent standard normal random variables. Our main tool in this calculation is the bias decomposition, stated in Proposition A.1. Recall that

$$i = \underset{j \in [1,2]}{\operatorname{arg\,max}} \widetilde{\Theta}_{3,j}.$$

By definition,

$$\widetilde{\Theta}_3 \sim \mathcal{N}\left(0, \left(\frac{\sigma^2 + \tau^2 + 2}{4}\right) \mathbb{I}_2\right).$$

Therefore, we have

$$\mathbb{E}\Big[\widetilde{\Theta}_{3,i}\Big] = \sqrt{\frac{\sigma^2 + \tau^2 + 2}{4}} \cdot \beta.$$

On the other hand, it follows from the symmetry that

$$\begin{split} \mathbb{E} \Big[\widetilde{\Theta}_{3,i} \Big] &= 2 \mathbb{E} \Big[\widetilde{\Theta}_{3,1} \cdot \mathbb{I}(i=1) \Big] \\ &= 2 \mathbb{E} \Big[\widetilde{\Theta}_{3,1} \cdot \mathbb{I}(\widetilde{\Theta}_{3,1} \geq \widetilde{\Theta}_{3,2}) \Big] \,. \end{split} \tag{A.7}$$

Combining Proposition A.1 for the sequence

$$X_1:=\frac{\Theta_1^\star}{2}, \qquad X_2:=\frac{\varepsilon_1}{2}, \qquad \text{and} \qquad X_3:=\widetilde{\Theta}_{3,1}-\widehat{\Theta}_{3,1}\sim \mathcal{N}\bigg(0,\frac{1}{2}\bigg)\,,$$

and $Z = \widetilde{\Theta}_{3,2}$, with Eq. (A.7), we infer that

$$\mathbb{E}\left[\frac{\Theta_{1}^{\star}}{2} \cdot \mathbb{I}(\widetilde{\Theta}_{3,1} \ge \widetilde{\Theta}_{3,2})\right] = \frac{\sigma^{2}}{\sigma^{2} + \tau^{2} + 2} \mathbb{E}\left[\frac{\widetilde{\Theta}_{3,1}}{2} \cdot \mathbb{I}(\widetilde{\Theta}_{3,1} \ge \widetilde{\Theta}_{3,2})\right]$$
$$= \frac{\sigma^{2}}{\sigma^{2} + \tau^{2} + 2} \cdot \frac{\sqrt{\sigma^{2} + \tau^{2} + 2}}{4} \cdot \beta$$
$$= \frac{\sigma^{2}\beta}{4\sqrt{\sigma^{2} + \tau^{2} + 2}}.$$

Consequently, we can write

$$\mathbb{E}[\Theta_i^{\star}] = 2 \,\mathbb{E}\left[\Theta_1^{\star} \cdot \mathbb{I}(\widetilde{\Theta}_{3,1} \ge \widetilde{\Theta}_{3,2})\right]$$

$$= 4 \,\mathbb{E}\left[\frac{\Theta_1^{\star}}{2} \cdot \mathbb{I}(\widetilde{\Theta}_{3,1} \ge \widetilde{\Theta}_{3,2})\right]$$

$$= \frac{\sigma^2 \beta}{\sqrt{\sigma^2 + \tau^2 + 2}}.$$
(A.8)

Similarly, we can conclude that

$$\mathbb{E}[\varepsilon_i] = \frac{\tau^2 \beta}{\sqrt{\sigma^2 + \tau^2 + 2}}.$$
 (A.9)

Combining Eq. (A.6) with Eq. (A.8) and Eq. (A.9), we obtain

$$\langle \mathbb{E}\left[\widehat{\Theta}_4\right], V \rangle = \frac{\left(\sigma^2 - \tau^2\right)\beta}{6\sqrt{\sigma^2 + \tau^2 + 2}}.$$

This equality implies that $\widehat{\Theta}_4$ is directionally biased whenever $\sigma^2 \neq \tau^2$. Finally, Eq. (A.3) and Eq. (A.5) give

$$\left\|\left\langle\widehat{\Theta}_{4},V\right\rangle\right\|_{\psi_{2}}=\left\|\widehat{\Theta}_{3,1}+\widehat{\Theta}_{3,2}+\frac{\Theta_{i}^{\star}-\varepsilon_{i}}{6}+\frac{\varepsilon_{3}}{3}\right\|_{\psi_{2}}$$

$$\leq \|\widehat{\Theta}_{3,1}\|_{\psi_{2}} + \|\widehat{\Theta}_{3,2}\|_{\psi_{2}} + \frac{1}{6} \|\Theta_{i}^{\star}\|_{\psi_{2}} + \frac{1}{6} \|\varepsilon_{i}\|_{\psi_{2}} + \frac{1}{3} \|\varepsilon_{3}\|_{\psi_{2}}$$

$$\leq 2\sqrt{\sigma^{2} + \tau^{2} + 2} + \frac{1}{6} \|\Theta_{i}^{\star}\|_{\psi_{2}} + \frac{1}{6} \|\varepsilon_{i}\|_{\psi_{2}} + \frac{2\tau}{3} .$$

Noting that, and using Eq. (A.3) again,

$$\left\| \Theta_{i}^{\star} \right\|_{\psi_{2}} = \left\| |\Theta_{i}^{\star}| \right\|_{\psi_{2}} \le \left\| |\Theta_{1}^{\star}| + |\Theta_{2}^{\star}| \right\|_{\psi_{2}} \le 2 \left\| |\Theta_{1}^{\star}| \right\|_{\psi_{2}} \le 4\sigma$$

and similarly for ε_i we can show $\|\varepsilon_i\|_{\psi_2} \leq 4\tau$ which means that

$$\|\langle \widehat{\Theta}_4, V \rangle\|_{\psi_2} \le 2\sqrt{\sigma^2 + \tau^2 + 2} + \frac{2}{3}(\sigma + \tau) + \frac{2\tau}{3}$$
$$\le 4(\sigma + \tau + \sqrt{2}).$$

Therefore, we have

$$\|\langle \widehat{\Theta}_4, V \rangle - \mathbb{E}[\langle \widehat{\Theta}_4, V \rangle]\|_{\psi_2} \le \|\langle \widehat{\Theta}_4, V \rangle\|_{\psi_2} + \|\mathbb{E}[\langle \widehat{\Theta}_4, V \rangle]\|_{\psi_2}$$

$$\le 8(\sigma + \tau + \sqrt{2}).$$

This and A.2 imply that the m.g.f. of $\widehat{\Theta}_4 - \mathbb{E}\big[\widehat{\Theta}_4\big]$ satisfies

$$\mathbb{E}\Big[\exp\Big(s\big\langle V, \widehat{\Theta}_4 - \mathbb{E}\big[\widehat{\Theta}_4\big]\big\rangle\Big)\Big] \leq \exp\Big[64C_{\mathrm{mgf}}\,s^2(\sigma + \tau + \sqrt{2})^2\Big]\;,\quad \text{ for all } s \in \mathbb{R}.$$

Proof of Lemma 3.2. Since all the d blocks are decoupled, it follows from Eq. (3.5) that

$$\mathbb{E}\left[\left\langle \widehat{\Theta}_{3d+1}, A \right\rangle\right] = -C_1(\sigma, \tau)\sqrt{d} \tag{A.10}$$

where

$$C_1(\sigma,\tau) = \frac{|\sigma^2 - \tau^2| \cdot \beta}{6\sqrt{\sigma^2 + \tau^2 + 2}}.$$

Assuming $\sigma^2 \neq \tau^2$, we observe that $C_1(\sigma, \tau) > 0$. Moreover, Eq. (3.6) implies that

$$\mathbb{E} \left[\exp \left(s \left(\left\langle \widehat{\Theta}_{3d+1}, A \right\rangle + C_1(\sigma, \tau) \sqrt{d} \right) \right) \right] \leq \exp \left[C s^2 (\sigma + \tau + \sqrt{2})^2 \right], \quad \text{for all } s \in \mathbb{R}$$

which means, for a constant C',

$$\left\|\left\langle \widehat{\Theta}_{3d+1}, A \right\rangle + C_1(\sigma, \tau) \sqrt{d} \right\|_{\psi_2} \le C'(\sigma + \tau + \sqrt{2}).$$

Using this inequality in combination with Eq. (A.1), and Eq. (A.10), we assert the following concentration inequality

$$\mathbb{P}\left(\left\langle \widehat{\Theta}_{3d+1}, A \right\rangle \leq -\frac{C_1(\sigma, \tau)\sqrt{d}}{2}\right) = \mathbb{P}\left(\left\langle \widehat{\Theta}_{3d+1}, A \right\rangle + C_1(\sigma, \tau)\sqrt{d} \leq \frac{C_1(\sigma, \tau)\sqrt{d}}{2}\right)$$

$$\geq 1 - 2 \exp \left[-C_{\text{tail}} \frac{C_1(\sigma, \tau)^2 d}{4C'^2(\sigma + \tau + \sqrt{2})^2} \right] .$$

Next, note that $\langle \Theta^{\star}, A \rangle \sim \mathcal{N}(0, 2\sigma^2),$ and thus, we have

$$\mathbb{P}\Big(\big\langle \Theta^{\star}, A \big\rangle \ge \sqrt{2}\sigma\Big) = 1 - \Phi(1)\,,$$

and for sufficiently large values of d,

$$2\exp\left[-C_{\text{tail}}\frac{C_{1}(\sigma,\tau)^{2}d}{4C'^{2}(\sigma+\tau+\sqrt{2})^{2}}\right] \leq \frac{1}{2}\left(1-\Phi(1)\right),$$

and hence

$$\mathbb{P}\Bigg(\big\langle \Theta^{\star}, A \big\rangle \geq \sqrt{2}\sigma \ \text{ and } \ \big\langle \widehat{\Theta}_{3d+1}, A \big\rangle \leq -\frac{C_1(\sigma, \tau)\sqrt{d}}{2} \Bigg) \geq p_0\,.$$

Proof of Proposition 3.1. First, note that the regret for each block is of order 1 because each of the two actions is equally likely to be selected. Therefore, the regret during the first 3d periods is of order d. This means, unless d = o(T), the regret would already be linear in T. Therefore, in the remaining we assume d = o(T).

For t > 3d, let Z_t be given by

$$Z_t := \begin{cases} 1 & \text{if action } A \text{ is never selected up to time } t, \\ 0 & \text{otherwise.} \end{cases}$$

We now have the following lower bound for the regret of Algorithm 1:

$$\operatorname{Regret}(T, \pi^{\operatorname{LinTS}}, \mathcal{P}_{\Theta^{\star}}) \geq \mathbb{P}(\mathcal{B}) \cdot \mathbb{E}\left[\sqrt{2}\sigma \cdot \sum_{t=3d+1}^{T} Z_{t} \middle| \mathcal{B}\right] \\
\geq \sqrt{2}\sigma p_{0} \cdot \sum_{t=3d+1}^{T} \mathbb{E}[Z_{t} \middle| \mathcal{B}] \\
= \sqrt{2}\sigma p_{0} \cdot \sum_{t=3d+1}^{T} \mathbb{P}(Z_{t} \middle| \mathcal{B}).$$

Define $q := \mathbb{P}(1 - Z_{3d+1} \mid \mathcal{B})$. We get that

$$\mathbb{P}(Z_t \mid \mathcal{B}) = \mathbb{P}(Z_t \mid \mathcal{B}, Z_{t-1}) \cdot \mathbb{P}(Z_{t-1} \mid \mathcal{B})$$
$$= (1 - q) \cdot \mathbb{P}(Z_{t-1} \mid \mathcal{B})$$
$$= (1 - q)^{t-3d}.$$

because $Z_t \mid \mathcal{B}, Z_{t-1}$ has the same distribution as $Z_{3d+1} \mid \mathcal{B}$ since choosing action 0 will not change LinTS's posterior estimate.

Furthermore, it follows from the definition of q and Eq. (A.4) that

$$1 - q \ge \mathbb{P}\left(\left\langle \widetilde{\Theta}_{3d+1} - \widehat{\Theta}_{3d+1}, A \right\rangle \le \frac{C_1(\sigma, \tau)\sqrt{d}}{2}\right)$$
$$\ge \mathbb{P}\left(\mathcal{N}(0, 1) \le \frac{C_1(\sigma, \tau)\sqrt{d}}{2}\right)$$
$$\ge 1 - \exp\left(-C_2(\sigma, \tau)d\right),$$

for a positive constant $C_2(\sigma, \tau)$. By combining the above, we have that

$$\begin{split} \operatorname{Regret}(T, \pi^{\operatorname{LinTS}}, \mathcal{P}_{\Theta^{\star}}) &\geq \sqrt{2}\sigma p_{0} \sum_{t=3d+1}^{T} \left[1 - \exp\left(-C_{2}(\sigma, \tau)d \right) \right]^{t-3d} \\ &= \sqrt{2}\sigma p_{0} \frac{1 - \left[1 - \exp\left(-C_{2}(\sigma, \tau)d \right) \right]^{t-3d}}{\exp\left(-C_{2}(\sigma, \tau)d \right)} \\ &\geq \sqrt{2}\sigma p_{0}(T-3d) \left(1 - \frac{T-3d}{2} \exp\left(-C_{2}(\sigma, \tau)d \right) \right) \,, \end{split}$$

where the last step uses inequality $1-(1-\alpha)^n \geq \alpha[1-(n-1)\alpha/2]$ for any integer n>2 and $\alpha\in(0,1)$. This immediately follows that, whenever $T\leq \exp\Big(C_2(\sigma,\tau)d\Big)$,

Regret
$$(T, \pi^{\text{LinTS}}, \mathcal{P}_{\Theta^*}) \ge \frac{\sqrt{2}\sigma p_0}{2} (T - 3d) \ge \Omega(T)$$
.

In other words, the regret of LinTS grows linearly up to time $\exp\left(C_2(\sigma,\tau)d\right)$.

Proof of Proposition 3.2. Notice that

$$\langle \Theta^{\star}, A' \rangle \sim \mathcal{N} \Big(-\mu \sqrt{d}, \sigma^2 \Big) \quad \text{ and } \quad \langle \Theta^{\star}, A \rangle \sim \mathcal{N} \Big(\mu \sqrt{d}, 3\sigma^2 \Big).$$

Therefore, for a positive constant C, $\langle \Theta^{\star}, A \rangle \geq \frac{1}{2}\mu\sqrt{d}$ and $\langle \Theta^{\star}, A' \rangle \leq -\frac{1}{2}\mu\sqrt{d}$ simultaneously with probability at least $1-2\exp\left(-C\frac{\mu^2d}{\sigma^2}\right)$. This thus implies that A is the optimal arm with high probability. For sufficiently large d, this probability exceeds $\frac{7}{8}$.

On the other hand, at t = 1, LinTS (Algorithm 1) will choose A' with probability $\frac{1}{4}$. This holds true as A' is chosen if and only if

$$\langle \widetilde{\Theta}_1, A' \rangle > 0$$
 and $\langle \widetilde{\Theta}_1, A - A' \rangle < 0$.

The claim follows from the fact these two random variables are two centered and independent normal random variables. In this case, we have

$$\mathbf{\Sigma}_2 = (\mathbb{I}_{3d} + A'A'^{\top})^{-1}$$

$$\begin{split} &= \mathbb{I}_{3d} - \frac{1}{2}A'A'^\top \quad \text{ and } \\ &\widehat{\Theta}_2 = \frac{1}{2}A'Y_1 \\ &= \frac{1}{2}A'(\langle \Theta^\star, A' \rangle + \varepsilon_1) \,. \end{split}$$

Next, we provide an upper bound for the probability that LinTS chooses arm 0 at t = 2. This happens if and only if

$$\left\langle \widetilde{\Theta}_{2},A^{\prime}\right\rangle <0$$
 and $\left\langle \widetilde{\Theta}_{2},A\right\rangle <0.$

Note that

$$\left\langle \widetilde{\Theta}_{2},A^{\prime}\right\rangle \sim\mathcal{N}\left(\frac{Y_{1}}{2},\frac{1}{2}\right)\quad\text{ and }\quad\left\langle \widetilde{\Theta}_{2},A\right\rangle \sim\mathcal{N}\left(\frac{Y_{1}}{2},\frac{5}{2}\right).\tag{A.11}$$

For sufficiently large d, we have

$$\mathbb{P}\bigg(\varepsilon_1 > \frac{1}{4}\mu\sqrt{d}\bigg) \le \frac{1}{16}.$$

From $Y_1 = \langle \Theta^*, A' \rangle + \varepsilon_1$ and the union bound it follows that $\mathbb{P}(\mathcal{B}') \geq 1/16$ where \mathcal{B}' is defined by

$$\mathcal{B}' := \left\{ \langle \Theta^\star, A \rangle \geq \frac{1}{2} \mu \sqrt{d}, \qquad \langle \Theta^\star, A' \rangle \leq -\frac{1}{2} \mu \sqrt{d}, \qquad \widetilde{A}_1 = A', \quad \text{and} \quad Y_1 < -\frac{1}{4} \mu \sqrt{d} \right\} \,.$$

From Eq. (A.11), we can deduce that, if $q:=\mathbb{P}\Big(\widetilde{A}_2\neq 0\ \Big|\ \mathcal{B}'\Big)$,

$$\begin{split} 1 - q &:= \mathbb{P}\Big(\widetilde{A}_2 = 0 \ \Big| \ \mathcal{B}'\Big) \\ &= \mathbb{P}\Big(\big\langle \widetilde{\Theta}_2, A' \big\rangle < 0 \quad \text{and} \quad \big\langle \widetilde{\Theta}_2, A \big\rangle < 0 \ \Big| \ \mathcal{B}'\Big) \\ &\geq 1 - 2 \exp\left(-C\mu^2 d\right) \ , \end{split}$$

for a positive constant d for a positive constant C. Applying the same argument as in proof of Proposition 3.1, we get that

Regret
$$(T, \pi^{\text{LinTS}}, \mathcal{P}_{\Theta^*}) \ge \frac{\mu\sqrt{d}}{32}T\left[1 - \frac{T}{2}\exp(-C_3d)\right]$$
 (A.12)

where C_3 is a constant that only depends on μ , σ , and τ (but not on d). Therefore, for $T \leq \exp(-C_3 d)$, the regret is linear in T.

Appendix B: A Formal Regret Bound for LinTSand Proofs of Section 4

This section is dedicated to our formal analysis of LinTS under our proposed conditions. This also allows us to show that the confidence set in OFUL can also be shrunk significantly under similar conditions. To do so, we adopt the framework in Hamidi and Bayati (2020) to state our results, however we make small changes compared to them. We start by explicitly stating the conditions that we introduced in Section 4. We say that the problem is in a *well-posed* condition at time t if

$$\psi(\mathbf{\Sigma}_{t}) \leq \Psi,$$

$$\|A_{t}^{\star}\|_{\mathbf{\Sigma}_{t}} \geq \omega \sqrt{\frac{\|\mathbf{\Sigma}_{t}\|_{*}}{d}} \cdot \|A_{t}^{\star}\|_{2},$$

$$\left|\left\langle A_{t}^{\star}, \widehat{\Theta}_{t} - \Theta^{\star} \right\rangle\right| \leq \frac{\nu}{\sqrt{d}} \|A_{t}^{\star}\|_{2} \cdot \left\|\widehat{\Theta}_{t} - \Theta^{\star}\right\|_{2},$$
(B.1)

and by \mathbb{W}_t we denote the indicator function of the above event. Now, by a *worth* function we mean a (stochastic) function $\widetilde{\mathsf{M}}_t(\cdot)$ that, given the history \mathcal{F}_t , assigns a real number to each action in the action set \mathcal{A}_t with the additional condition that whenever $\mathbb{W}_t = 1$

$$\mathbb{P}\left(\sup_{A \in \mathcal{A}_t} \frac{|\widetilde{\mathsf{M}}_t(A) - \langle A, \widehat{\Theta}_t \rangle|}{\|A\|_{\Sigma_t}} \le \widetilde{\rho} \,\middle|\, \mathcal{F}_t, \mathcal{A}_t\right) \ge 1 - \frac{1}{T^2} \tag{B.2}$$

for some fixed $\widetilde{\rho} > 1$. We also say that $\widehat{\Theta}_t$ is in a typical condition with respect to Θ^* if

$$\mathbb{P}\left(\sup_{A \in \mathcal{A}_t} \frac{|\langle A, \Theta^{\star} \rangle - \langle A, \widehat{\Theta}_t \rangle|}{\|A\|_{\Sigma_t}} \le \rho^{\star} \,\middle|\, \mathcal{F}_t, \mathcal{A}_t\right) \ge 1 - \frac{1}{T^2} \tag{B.3}$$

for some $\rho^* > 1$ whenever $\mathbb{W}_t = 1$. We also let \mathbb{T}_t^* be the indicator function for this event. Intuitively, ρ^* determines the width of the confidence interval around each $\langle \widehat{\Theta}_t, A \rangle$ so that it contains $\langle \Theta^*, A \rangle$ for all $A \in \mathcal{A}_t$ simultaneously with high probability. Next, we say that the worth function $\widetilde{\mathsf{M}}_t(\cdot)$ is *optimistic* for given realizations \mathcal{F}_t and \mathcal{A}_t if

$$\mathbb{P}\bigg(\sup_{A\in\mathcal{A}_t}\widetilde{\mathsf{M}}_t(A)\geq \sup_{A\in\mathcal{A}_t}\langle A,\Theta^{\star}\rangle \ \bigg|\ \mathcal{F}_t,\mathcal{A}_t\bigg)\geq \mathsf{p}$$

for p > 0 whenever $\mathbb{W}_t = 1$ and $\mathbb{T}_t^{\star} = 1$. This notion of optimism *cannot* hold almost surely as, for instance, $\Theta^{\star} - \widehat{\Theta}_t$ can be arbitrarily large.

Using these notations, we introduce a modified version of Randomized OFUL (ROFUL), introduced in Hamidi and Bayati (2020), that we analyze in this paper. The pseudo-code for this meta algorithm is presented in Algorithm 3. Whenever $\psi(\Sigma_t) > \Psi$, ROFUL is allowed to select any arbitrary action. A natural choice is to choose the action that decreases $\psi(\Sigma_t)$ the most. An alternative is to define $\widetilde{A}_t \leftarrow \arg\max_{A \in \mathcal{A}_t} \widetilde{\mathsf{M}}_t(A)$ as in Line 4. In this case, if one sets $\widetilde{\mathsf{M}}_t(A) = \langle A, \widetilde{\Theta}_t \rangle$ or $\widetilde{\mathsf{M}}_t(A) = \langle A, \widehat{\Theta}_t \rangle + \widetilde{\rho} \|A\|_{\Sigma_t}$, ROFUL becomes LinTS or OFUL respectively. Because, for OFUL, such $\widetilde{\mathsf{M}}_t(A)$ satisfy Eq. (B.2), by definition. For LinTS, noting that $\langle A, \widetilde{\Theta}_t \rangle = \langle A, \widehat{\Theta}_t \rangle + \iota \|A\|_{\Sigma_t} \mathcal{N}(0,1)$, from A.4 follows that Eq. (B.2) holds as long as $\widetilde{\rho}/\iota = \Omega(\sqrt{\log T})$.

Algorithm 3 Randomized OFUL (ROFUL)

Require: Worth functions $(\widetilde{\mathsf{M}}_t)_{t=1}^T$

- 1: **for** $t = 1, 2, \cdots$ **do**
- 2: Observe A_t
- 3: **if** $\psi(\Sigma_t) \leq \Psi$ **then**
- 4: $\widetilde{A}_t \leftarrow \arg \max_{A \in \mathcal{A}_t} \widetilde{\mathsf{M}}_t(A)$
- 5: else
- 6: Choose any $\widetilde{A}_t \in \mathcal{A}_t$
- 7: **end if**
- 8: Observe reward Y_t
- 9: end for

The next two lemmas assert that the optimism holds for LinTS and OFUL. But we first recall the definition of ρ from Abbasi-Yadkori et al. (2011).

$$\rho := 3\sigma \sqrt{d\log(1 + \lambda T \mathbf{a}^2)} + \lambda^{-1/2} \boldsymbol{\theta}, \tag{B.4}$$

where θ is an upper bound for $\|\Theta^*\|_2$ and \mathbf{a} is defined in Section 2. Therefore, $\rho = \widetilde{\mathcal{O}}(\sqrt{d})$. Also, Theorem 1 of Abbasi-Yadkori et al. (2011) gives

$$\mathbb{P}\left(\|\Theta^* - \widehat{\Theta}_t\|_{\Sigma_t^{-1}} \le \rho\right) \ge 1 - \frac{1}{T^3}. \tag{B.5}$$

Let \mathbb{T}_t be the binary indicator for the event in Eq. (B.5). Note that by Cauchy-Schwartz inequality one can easily see that if $\rho^* \geq \rho$, then $\mathbb{T}_t^* \geq \mathbb{T}_t$. In fact, in the rest of this section, one can harmlessly assume $\rho^* = \rho$ and only work with \mathbb{T}_t . We only use separate notation to allow the possibility of $\rho^* < \rho$.

Lemma B.1 (Optimism of LinTS) Set the inflation parameter of LinTS to be

$$\iota := \frac{\nu \Psi}{\omega} \cdot \frac{\rho}{\sqrt{d}}$$

and let $\widetilde{\mathsf{M}}_t(A) := \langle A, \widetilde{\Theta}_t \rangle$. Whenever $\mathbb{W}_t = 1$, $\mathbb{T}_t = 1$, and $\mathbb{T}_t^{\star} = 1$, we have

$$\mathbb{P}\left(\sup_{A \in \mathcal{A}_t} \widetilde{\mathsf{M}}_t(A) \ge \sup_{A \in \mathcal{A}_t} \left\langle A, \Theta^* \right\rangle \,\middle|\, \mathcal{F}_t, \mathcal{A}_t \right) \ge \Phi(-1). \tag{B.6}$$

Lemma B.2 (Optimism of OFUL) Set ι as in Lemma B.1 and $\widetilde{\mathsf{M}}_t(A) := \langle A, \widehat{\Theta}_t \rangle + \iota \|A\|_{\Sigma_t}$. Whenever $\mathbb{W}_t = 1$, $\mathbb{T}_t = 1$, and $\mathbb{T}_t^\star = 1$, we have

$$\mathbb{P}\left(\sup_{A \in \mathcal{A}_t} \widetilde{\mathsf{M}}_t(A) \ge \sup_{A \in \mathcal{A}_t} \left\langle A, \Theta^* \right\rangle \,\middle|\, \mathcal{F}_t, \mathcal{A}_t\right) = 1. \tag{B.7}$$

We establish the proof of Lemma B.1, noting that the proof of Lemma B.2 would be almost identical and marginally simpler.

Proof of Lemma B.1. We have

$$\begin{split} \left\langle \widetilde{\Theta}_{t}, A_{t}^{\star} \right\rangle - \left\langle \Theta^{\star}, A_{t}^{\star} \right\rangle &= \left\langle \widetilde{\Theta}_{t} - \widehat{\Theta}_{t}, A_{t}^{\star} \right\rangle - \left\langle \Theta^{\star} - \widehat{\Theta}_{t}, A_{t}^{\star} \right\rangle \\ &\geq \left\langle \widetilde{\Theta}_{t} - \widehat{\Theta}_{t}, A_{t}^{\star} \right\rangle - \frac{\nu}{\sqrt{d}} \|A_{t}^{\star}\|_{2} \cdot \left\| \widehat{\Theta}_{t} - \Theta^{\star} \right\|_{2} \\ &\geq \left\langle \widetilde{\Theta}_{t} - \widehat{\Theta}_{t}, A_{t}^{\star} \right\rangle - \nu \rho \sqrt{\frac{\|\Sigma_{t}\|_{\text{op}}}{d}} \cdot \|A^{\star}\|_{2}, \end{split}$$

where we used the third equation in Eq. (B.1) first and then Eq. (B.5), combined with $||V||_2 \le ||V||_{\Sigma_{\star}^{-1}} \sqrt{||\Sigma_t||_{\text{op}}}$ for any vector V.

Now, since $\langle \widetilde{\Theta}_t - \widehat{\Theta}_t, A_t^{\star} \rangle \sim \mathcal{N}(0, \iota^2 ||A^{\star}||_{\Sigma_t}^2)$, we have

$$\begin{split} \mathbb{P}\Big(\left\langle \widetilde{\Theta}_{t}, A_{t}^{\star} \right\rangle &\geq \left\langle \Theta^{\star}, A_{t}^{\star} \right\rangle \Big) \geq \mathbb{P}\Bigg(\left\langle \widetilde{\Theta}_{t} - \widehat{\Theta}_{t}, A_{t}^{\star} \right\rangle \geq \nu \rho \sqrt{\frac{\|\boldsymbol{\Sigma}_{t}\|_{\mathrm{op}}}{d}} \cdot \|A^{\star}\|_{2} \Big) \\ &= \Phi\left(-\frac{\nu \rho \sqrt{\frac{\|\boldsymbol{\Sigma}_{t}\|_{\mathrm{op}}}{d}} \cdot \|A^{\star}\|_{2}}{\iota \|A^{\star}\|_{\boldsymbol{\Sigma}_{t}}}\right) \\ &\geq \Phi\left(-\frac{\nu \rho \sqrt{\frac{\|\boldsymbol{\Sigma}_{t}\|_{\mathrm{op}}}{d}} \cdot \|A^{\star}\|_{2}}{\iota \omega \sqrt{\frac{\|\boldsymbol{\Sigma}_{t}\|_{*}}{d}} \cdot \|A^{\star}\|_{2}}\right) \\ &= \Phi\left(-\frac{\nu \rho}{\iota \omega \sqrt{d}} \cdot \sqrt{\frac{d\|\boldsymbol{\Sigma}_{t}\|_{\mathrm{op}}}{\|\boldsymbol{\Sigma}_{t}\|_{*}}}\right) \\ &\geq \Phi(-1) \,, \end{split}$$

where we used the second and third equation in Eq. (B.1). This completes the proof.

We are now ready to state our main result.

Theorem B.1 Let $\widetilde{\mathsf{M}}_t(\cdot)$ be optimistic with parameter p whenever $\mathbb{T}_t = 1$, $\mathbb{T}_t^\star = 1$, and $\mathbb{W}_t = 1$. When Assumptions 4.1 and 4.2 hold, and $\langle A, \Theta^\star \rangle \in [-1,1]$, for all $A \in \mathcal{A}_t$, almost surely. Then, we have that

$$\sum_{t=1}^T \left\langle \Theta^\star, A_t^\star \right\rangle - \left\langle \Theta^\star, \widetilde{A}_t \right\rangle \leq \frac{4(\rho^\star + \widetilde{\rho})\sqrt{T}}{\mathsf{p}} \left(\sqrt{d \log \left(1 + \frac{\lambda T \mathbf{a}^2}{d} \right)} + \sqrt{\log T} \right) + 2 \sum_{t=1}^T \mathbb{I}(\psi(\mathbf{\Sigma}_t) > \Psi) \,,$$

with probability at least $1 - \frac{3}{T}$.

Before describing the proof, we state a direct corollary of Theorem B.1 and Lemma B.1.

Corollary B.1 (LinTS with smaller inflation) Consider LinTS with inflation parameter ι satisfying

$$\iota = \frac{\nu \Psi}{\omega} \cdot \frac{\rho}{\sqrt{d}} \quad and \quad \frac{\widetilde{\rho}}{\iota} = \Omega(\sqrt{\log T}).$$
 (B.8)

Then, when Assumptions 4.1 and 4.2 hold, the regret is at most

$$\frac{4(\rho^* + \widetilde{\rho})\sqrt{T}}{\Phi(-1)} \left(\sqrt{d \log \left(1 + \frac{\lambda T \mathbf{a}^2}{d} \right)} + \sqrt{\log T} \right) + 2 \sum_{t=1}^{T} \mathbb{I}(\psi(\mathbf{\Sigma}_t) > \Psi),$$

with probability at least $1 - \frac{3}{T}$.

The implication of Corollary B.1 is that, under Assumptions 4.1 and 4.2, one can circumvent the need for the \sqrt{d} inflation factor in the posterior of the LinTS algorithm, as the parameter ι grows only logarithmically with respect to T.

By utilizing both Theorem B.1 and Lemma B.2, we can derive a similar corollary for OFUL.

Corollary B.2 (OFUL with smaller confidence intervals) Consider a version of OFUL which is an instance of ROFUL with worth function $\widetilde{\mathsf{M}}_t(A) = \langle A, \widehat{\Theta}_t \rangle + \widetilde{\rho} \|A\|_{\Sigma_t}$ such that $\widetilde{\rho}$ satisfies

$$\widetilde{\rho} = \frac{\nu \Psi}{\omega} \cdot \frac{\rho}{\sqrt{d}} \,.$$

Then, when Assumptions 4.1 and 4.2 hold, the regret is at most

$$4(\rho^{\star} + \widetilde{\rho})\sqrt{T}\left(\sqrt{d\log\left(1 + \frac{\lambda T\mathbf{a}^2}{d}\right)} + \sqrt{\log T}\right) + 2\sum_{t=1}^{T} \mathbb{I}(\psi(\mathbf{\Sigma}_t) > \Psi),$$

with probability at least $1 - \frac{3}{T}$.

Corollary B.2 demonstrates that, under Assumptions 4.1 and 4.2, it is possible to improve the performance of the OFUL algorithm by running it with smaller confidence sets that are reduced by a factor of \sqrt{d} , given that $\tilde{\rho}$ only logarithmically depends on T. It is worth noting that while the reduced confidence sets do not result in better upper bounds due to ρ^* remaining of order $\tilde{\mathcal{O}}(\sqrt{d})$, they may lead to improved empirical performance of the OFUL algorithm.

Proof of Theorem B.1. First, let $(\mathbb{O}_t)_{t\in[T]}$ be the adapted sequence of Bernoulli random variables such that $\mathbb{O}_t = 1$ whenever $\mathbb{T}_t = 1$, $\mathbb{T}_t^* = 1$, $\mathbb{W}_t = 1$, and

$$\sup_{A \in \mathcal{A}_t} \widetilde{\mathsf{M}}_t(A) > \sup_{A \in \mathcal{A}_t} \langle A, \Theta^{\star} \rangle$$

simultaneously. It follows from the definition of optimism that $\mathbb{P}(\mathbb{O}_t = 1 \mid \mathcal{F}_t, \mathcal{A}_t) \geq p$. Let $t \in [T]$ be fixed and assume that $\mathbb{T}_t = 1$, $\mathbb{T}_t^{\star} = 1$, $\mathbb{W}_t = 1$, and $\mathbb{O}_t = 1$. Then, we

$$\begin{split} \left\langle \Theta^{\star}, A_{t}^{\star} \right\rangle - \left\langle \Theta^{\star}, \widetilde{A}_{t} \right\rangle &\leq \widetilde{\mathsf{M}}_{t}(\widetilde{A}_{t}) - \left\langle \Theta^{\star}, \widetilde{A}_{t} \right\rangle \\ &\leq \left(\left\langle \widehat{\Theta}_{t}, \widetilde{A}_{t} \right\rangle + \widetilde{\rho} \|\widetilde{A}_{t}\|_{\Sigma_{t}} \right) - \left(\left\langle \widehat{\Theta}_{t}, \widetilde{A}_{t} \right\rangle - \rho^{\star} \|\widetilde{A}_{t}\|_{\Sigma_{t}} \right) \\ &\leq (\rho^{\star} + \widetilde{\rho}) \|\widetilde{A}_{t}\|_{\Sigma_{t}} \,. \end{split}$$

Furthermore, since $\langle \Theta^{\star}, A_t^{\star} \rangle - \langle \Theta^{\star}, \widetilde{A}_t \rangle \leq 2$ and $\rho^{\star} + \widetilde{\rho} \geq 2$, we have

$$\left\langle \Theta^{\star}, A_{t}^{\star} \right\rangle - \left\langle \Theta^{\star}, \widetilde{A}_{t} \right\rangle \leq 2 \min \left(\frac{\rho^{\star} + \widetilde{\rho}}{2} \| \widetilde{A}_{t} \|_{\Sigma_{t}}, 1 \right) \leq (\rho^{\star} + \widetilde{\rho}) \min \left(\| \widetilde{A}_{t} \|_{\Sigma_{t}}, 1 \right),$$

and,

$$\mathbb{E}[Z_t \mid \mathcal{F}_t, \mathcal{A}_t] \le 0, \tag{B.9}$$

almost surely, where Z_t is defined as

$$Z_t := \mathbb{T}_t \mathbb{T}_t^\star \mathbb{W}_t \left\{ \left\langle \Theta^\star, A_t^\star \right\rangle - \left\langle \Theta^\star, \widetilde{A}_t \right\rangle - \frac{(\rho^\star + \widetilde{\rho}) \mathbb{O}_t}{\mathsf{p}} \cdot \min \left(\|\widetilde{A}_t\|_{\mathbf{\Sigma}_t}, 1 \right) \right\}.$$

It follows from Eq. (B.9) that $\sum_{t=1}^{T} Z_t$ is a super-martingale, and noting that $|Z_t| \leq \frac{2(\rho^* + \tilde{\rho})}{p}$, it follows from Azuma's inequality that

$$\mathbb{P}\left(\sum_{t=1}^{T} Z_{t} \ge \frac{4(\rho^{\star} + \widetilde{\rho})\sqrt{T\log T}}{\mathsf{p}}\right) \le \frac{1}{T}.$$
(B.10)

Next, by applying the Cauchy-Schwartz inequality and Lemma 11 in Abbasi-Yadkori et al. (2011), we deduce that

$$\begin{split} \sum_{t=1}^{T} \left(Z_{t} - \mathbb{T}_{t} \mathbb{T}_{t}^{\star} \mathbb{W}_{t} \left\langle \Theta^{\star}, A_{t}^{\star} - \widetilde{A}_{t} \right\rangle \right) &\geq -\frac{(\rho^{\star} + \widetilde{\rho})}{\mathsf{p}} \sum_{t=1}^{T} \min \left(\|\widetilde{A}_{t}\|_{\Sigma_{t}}, 1 \right) \\ &\geq -\frac{(\rho^{\star} + \widetilde{\rho})}{\mathsf{p}} \sqrt{T \sum_{t=1}^{T} \min \left(\|\widetilde{A}_{t}\|_{\Sigma_{t}}^{2}, 1 \right)} \\ &\geq -\frac{(\rho^{\star} + \widetilde{\rho})}{\mathsf{p}} \sqrt{2T d \log \left(1 + \frac{\lambda T \mathbf{a}^{2}}{d} \right)}. \end{split}$$

By combining the above inequality with Eq. (B.10), we obtain

$$\mathbb{P}\left(\sum_{t=1}^{T} \mathbb{T}_{t} \mathbb{T}_{t}^{\star} \mathbb{W}_{t} \left\langle \Theta^{\star}, A_{t}^{\star} - \widetilde{A}_{t} \right\rangle \geq \frac{4(\rho^{\star} + \widetilde{\rho})\sqrt{T}}{\mathsf{p}} \left(\sqrt{d \log \left(1 + \frac{\lambda T \mathbf{a}^{2}}{d}\right)} + \sqrt{\log T} \right) \right) \leq \frac{1}{T}$$
 (B.11)

We now turn to bounding $(1 - \mathbb{T}_t \mathbb{T}_t^{\star} \mathbb{W}_t) \langle \Theta^{\star}, A_t^{\star} - \widetilde{A}_t \rangle$. Notice that

$$\begin{split} (1 - \mathbb{T}_t \mathbb{T}_t^{\star} \mathbb{W}_t) \left\langle \Theta^{\star}, A_t^{\star} - \widetilde{A}_t \right\rangle &\leq 2 (1 - \mathbb{T}_t \mathbb{T}_t^{\star} \mathbb{W}_t) \\ &= 2 (1 - \mathbb{T}_t \mathbb{T}_t^{\star} \mathbb{W}_t) \mathbb{I}(\psi(\mathbf{\Sigma}_t) \leq \Psi) + 2 (1 - \mathbb{T}_t \mathbb{T}_t^{\star} \mathbb{W}_t) \mathbb{I}(\psi(\mathbf{\Sigma}_t) > \Psi) \\ &\leq 2 (1 - \mathbb{T}_t \mathbb{T}_t^{\star} \mathbb{W}_t) \mathbb{I}(\psi(\mathbf{\Sigma}_t) \leq \Psi) + 2 \mathbb{I}(\psi(\mathbf{\Sigma}_t) > \Psi). \end{split}$$

Also, it follows from Assumptions 4.1 and 4.2 that for any $t \in [T]$

$$\mathbb{P}(\mathbb{T}_t \mathbb{T}_t^{\star} \mathbb{W}_t = 0 \quad \text{and} \quad \psi(\mathbf{\Sigma}_t) \leq \Psi) \leq \frac{2}{T^2} \,,$$

which in combination with the union bound leads to

$$\sum_{t=1}^{T} (1 - \mathbb{T}_t \mathbb{T}_t^{\star} \mathbb{W}_t) \langle \Theta^{\star}, A_t^{\star} - \widetilde{A}_t \rangle \leq 2 \sum_{t=1}^{T} \mathbb{I}(\psi(\Sigma_t) > \Psi)$$

with probability at least $1 - \frac{2}{T}$. Finally, this together with Eq. (B.11) yield

$$\sum_{t=1}^T \left\langle \Theta^\star, A_t^\star - \widetilde{A}_t \right\rangle \leq \frac{4(\rho^\star + \widetilde{\rho})\sqrt{T}}{\mathsf{p}} \left(\sqrt{d \log \left(1 + \frac{\lambda T \mathbf{a}^2}{d} \right)} + \sqrt{\log T} \right) + 2\sum_{t=1}^T \mathbb{I}(\psi(\mathbf{\Sigma}_t) > \Psi) \,,$$

with probability at least $1 - \frac{3}{T}$.

Appendix C: Auxiliary Proofs

Proof of Proposition A.1. It is straightforward to see that $X_i|Y$ follows Gaussian distribution with mean $\frac{\sigma_i^2}{\sum_{i=1}^n \sigma_i^2} \cdot Y$. We thus get

$$\begin{split} \mathbb{E}[X_i \cdot g(Y, Z)] &= \mathbb{E}[\mathbb{E}[X_i \cdot g(Y, Z) \mid Y, Z]] \\ &= \mathbb{E}[\mathbb{E}[X_i \mid Y, Z] \cdot g(Y, Z)] \\ &= \frac{\sigma_i^2}{\sum_{i=1}^n \sigma_i^2} \cdot \mathbb{E}[Y \cdot g(Y, Z)] \,. \end{split}$$

Proof of Lemma 6.1. First of all, it follows from the definition of A that

$$\mathbb{E}\big[\|A\|_{\mathbf{\Sigma}_t}^2\big] = \frac{1}{3d}\operatorname{Tr}(\mathbf{\Sigma}_t) = \frac{1}{3d}\|\mathbf{\Sigma}_t\|_*.$$

It then follows from the Hanson-Wright inequality (e.g., Theorem 6.2.1 in Vershynin (2018)) that

$$\mathbb{P}\left(\left|\|A\|_{\mathbf{\Sigma}}^{2} - \frac{1}{3d}\|\mathbf{\Sigma}\|_{*}\right| \leq \frac{1}{6d}\|\mathbf{\Sigma}\|_{*}\right) \leq \exp\left(-c\min\left\{\frac{\|\mathbf{\Sigma}_{t}\|_{*}^{2}}{\|\mathbf{\Sigma}_{t}\|_{F}^{2}}, \frac{\|\mathbf{\Sigma}_{t}\|_{*}}{\|\mathbf{\Sigma}_{t}\|_{\text{op}}}\right\}\right)$$

for some constant c > 0. Noting that

$$\|\mathbf{\Sigma}_t\|_{\mathrm{op}}\|\mathbf{\Sigma}_t\|_* \geq \|\mathbf{\Sigma}_t\|_{\mathrm{F}}^2,$$

we can simplify the above tail bound to get

$$\mathbb{P}\left(\left|\|A\|_{\boldsymbol{\Sigma}}^2 - \frac{1}{3d}\|\boldsymbol{\Sigma}\|_*\right| \leq \frac{1}{6d}\|\boldsymbol{\Sigma}\|_*\right) \leq \exp\left(-c\frac{\|\boldsymbol{\Sigma}_t\|_*}{\|\boldsymbol{\Sigma}_t\|_{\mathrm{op}}}\right) \leq \exp\left(-\frac{cd}{\Psi^2}\right).$$

Proof of Lemma 6.2. Note that for all s > 0 we have that

$$\mathbb{E}[\exp(s\langle A, V \rangle)] = \prod_{i=1}^{d} \mathbb{E}[\exp(sA_{i}V_{i})]$$

$$\leq \prod_{i=1}^{d} \exp\left(\frac{s^{2}V_{i}^{2}}{2d}\right)$$

$$= \exp\left(\frac{s^{2}\sum_{i=1}^{d}V_{i}^{2}}{2d}\right)$$

$$= \exp\left(\frac{s^{2}}{2d}\right).$$

It thus follows from the Chernoff bound that

$$\mathbb{P}\bigg(\langle A, V \rangle > \sqrt{\frac{2\log(1/p)}{d}}\bigg) \leq p,$$

which is the desired result.

References

Abbasi-Yadkori, Yasin, Dávid Pál, Csaba Szepesvári. 2011. Improved algorithms for linear stochastic bandits. *Advances in Neural Information Processing Systems*. 2312–2320.

Abeille, Marc, Alessandro Lazaric, et al. 2017. Linear thompson sampling revisited. *Electronic Journal of Statistics* **11**(2) 5165–5197.

Agrawal, Shipra, Navin Goyal. 2012. Analysis of thompson sampling for the multi-armed bandit problem. *Conference on learning theory*. 39–1.

Agrawal, Shipra, Navin Goyal. 2013a. Further optimal regret bounds for thompson sampling. Aistats. 99-107.

Agrawal, Shipra, Navin Goyal. 2013b. Thompson sampling for contextual bandits with linear payoffs. *ICML* (3). 127–135.

Bastani, Hamsa, David Simchi-Levi, Ruihao Zhu. 2019. Meta Dynamic Pricing: Transfer Learning Across Experiments arXiv:1902.10918.

Bubeck, Sébastien, Nicolo Cesa-Bianchi, et al. 2012. Regret analysis of stochastic and nonstochastic multi-armed bandit problems. *Foundations and Trends*® *in Machine Learning* **5**(1) 1–122.

Bubeck, Sébastien, Che-Yu Liu. 2013. Prior-free and prior-dependent regret bounds for thompson sampling. *Advances in Neural Information Processing Systems*. 638–646.

Cormen, Thomas H., Charles E. Leiserson, Ronald L. Rivest, Clifford Stein. 2001. *Introduction to Algorithms*. 2nd ed. The MIT Press.

Dani, Varsha, Thomas P. Hayes, Sham M. Kakade. 2008. Stochastic linear optimization under bandit feedback. COLT.

- Hamidi, Nima, Mohsen Bayati. 2020. A general theory of the stochastic linear bandit and its applications. *arXiv* preprint arXiv:2002.05152 URL https://arxiv.org/pdf/2002.05152.pdf.
- Jin, Tianyuan, Pan Xu, Jieming Shi, Xiaokui Xiao, Quanquan Gu. 2020. Mots: Minimax optimal thompson sampling. arXiv preprint arXiv:2003.01803.
- Johari, Ramesh, Pete Koomen, Leonid Pekelis, David Walsh. 2017. Peeking at a/b tests: Why it matters, and what to do about it. Proceedings of the 23rd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining. ACM, New York, NY, USA, 1517–1525. doi:10.1145/3097983.3097992.
- Lai, Tze Leung, Herbert Robbins. 1985. Asymptotically efficient adaptive allocation rules. *Advances in applied mathematics* **6**(1) 4–22.
- Lattimore, Tor, Csaba Szepesvari. 2019. Bandit Algorithms.
- Nie, Xinkun, Xiaoying Tian, Jonathan Taylor, James Zou. 2018. Why adaptively collected data have negative bias and how to correct for it. *International Conference on Artificial Intelligence and Statistics*. 1261–1269.
- Phan, My, Yasin Abbasi Yadkori, Justin Domke. 2019. Thompson sampling and approximate inference. *Advances in Neural Information Processing Systems*. 8804–8813.
- Rusmevichientong, Paat, John N Tsitsiklis. 2010. Linearly parameterized bandits. *Mathematics of Operations Research* **35**(2) 395–411.
- Russo, Daniel, Benjamin Van Roy. 2014. Learning to optimize via posterior sampling. *Mathematics of Operations Research* **39**(4) 1221–1243. doi:10.1287/moor.2014.0650.
- Russo, Daniel J., Benjamin Van Roy, Abbas Kazerouni, Ian Osband, Zheng Wen. 2018. A tutorial on thompson sampling. *Foundations and Trends in Machine Learning* **11**(1) 1–96.
- Scott, Steven L. 2010. A modern bayesian look at the multi-armed bandit. *Applied Stochastic Models in Business and Industry* **26**(6) 639–658.
- Scott, Steven L. 2015. Multi-armed bandit experiments in the online service economy. *Appl. Stoch. Model. Bus. Ind.* **31**(1) 37–45. doi:10.1002/asmb.2104.
- Slivkins, Aleksandrs. 2019. Introduction to multi-armed bandits. *Foundations and Trends*® *in Machine Learning* **12**(1-2) 1–286.
- Thompson, William R. 1933. On the likelihood that one unknown probability exceeds another in view of the evidence of two samples. *Biometrika* **25**(3/4) 285–294.
- Vershynin, Roman. 2018. *High-dimensional probability: An introduction with applications in data science*, vol. 47. Cambridge university press.