A simple direct quantum model which, with no random phase assumptions and with arbitrary initial conditions, evolves to the Boltzman distribution

Michael J. Caola 6 Normanton Rd., Bristol, BS8 2TY, UK caolam@blueyonder.co.uk

November 27, 2024

Abstract

We consider M systems (each an electron in a long square cylinder) uniformly arranged on a ring and with Coulomb interactions. Exact straightforward numerical time-dependent perturbation calculation of a single N-level (≤ 7) system, with no (random) phase assumptions, system show a Boltzman distribution. We exploit the physical ring symmetry and develop several hierarchical physical equation set so of increasing generality and (computation) speed. Given the impressive history of theoretical quantum-mehanical statistical mechanics, our results might seem surprising, but we observe that accurate calculation of correct physical equations should mimic Nature.

1 Introduction

An important part of Statistical Mechanics is to explain how a physical system can tend to an equilibrium state, usually based on its energy structure. Most analysis, both classical and quantum-mechanical, invokes probability assumptions which in the latter are additional to those of the basic non-relativistic Schrodinger wave-function[,,]. We treat the quantum case and in particular wish to understand how a quantum system can (in time) tend to a state

$$\psi = \sum_{n=1}^{N} c_n \psi_n \tag{1}$$

whose (eigen)states ψ_n and (eigen)energies E_n are defined by Hamiltonian H_0 : $H_0\psi_n = E_n\psi_n$, $\omega_n = E_n/\hbar, \Psi_n(\mathbf{r}, \mathbf{t}) = \psi_{\mathbf{n}}(\mathbf{r})\mathbf{e}^{-\mathbf{i}\omega_{\mathbf{n}}\mathbf{t}}$. The expansion coefficient $c_n = c_n(t \to \infty)$ gives the probability w_n that in equilibrium the system is in state ψ_n :

$$w_n = |c_n|^2 = e^{-\beta E_n} / \sum_n e^{-\beta E_n}$$
(2)

This (2) is the Boltzman distribution where $\beta = 1/kT$, T = temperature and k is the Boltzmann constant.

2 Analysis

2.1 Model geometry

Our model is M identical systems uniformly on a ring radius R and normal x. The model is isolated (from the rest of the universe). Each system is an electron in a "matchstick" box which is a long a square $b \ll a$ prism, see fig.1. The physical environment each system m = 1..M is that of its M - 1 neighbours, and inspection/symmetry shows that this is the same for all M systems: every system m = 1..M is in the 'same heat bath'. This 'same heat bath' is essentially a 2D model with finite M, and it would seem that in 1D or 3D only $M = \infty$ is possible.

The energy of an electron in an infinitely deep potential rectangular box, sides (a_x, a_y, a_z) , is

$$E_{n_x,n_y,n_z} = \sqrt{\frac{8}{a_x a_y a_z}} \left(\frac{n_x^2}{a_x^2} + \frac{n_y^2}{a_y^2} + \frac{n_z^2}{a_z^2} \right)$$
(3)

and eignfunctions, $\omega = E/\hbar$,

$$\Psi_{n_x,n_y,n_z}(x,y,z,t) = \psi_{n_x,n_y,n_z}(x,y,z)e^{-i\omega_{n_x,n_y,n_z}t}$$
(4)

$$\psi_{n_x,n_y,n_z}(x,y,z) = \sqrt{\frac{8}{a_x a_y a_z}} \left(\sin \frac{\pi n_x}{a_x} x \sin \frac{\pi n_y}{a_y} y \sin \frac{\pi n_z}{a_z} z \right)$$
(5)

$$\equiv \psi_{n_x}(x)\psi_{n_y}(y)\psi_{n_z}(z) \tag{6}$$

Our matchstick system with $\{a, b \ll a\}$ means that energies $(E_{N,1,1} - E_{1,1,1}) \ll E_{1,2,1}$: the N lowest energy states enjoy a large energy separation from higher ones and it is a well-established quantum common-place that the effects of a perturbation on the system may be accurately calculated using these N states only — we have a N-level system. Thus, with $n_x \equiv n$, we henceforth deal only with states $\psi_{n,1,1}$, n = 1..N

$$\psi_n(x) = \psi_{n,1,1}(x, y, z) = \sqrt{\frac{8}{ab^2}} \left(\sin \frac{\pi n}{a} x \sin \frac{\pi}{b} y \sin \frac{\pi}{b} z \right) \tag{7}$$

The last part of the geometry is, classically stated, the separation $r_{m,m'}$ of electrons at positions x_m in system m from $x_{m'}$ in system m', see Fig.1:

$$r_{m,m'}(x_m, x_{m'}) = \sqrt{(x_m - x_{m'})^2 + 4R^2 \sin^2\left(\frac{\pi(m - m')}{M}\right)}$$
(8)

2.2 Model perturbations

If there are no interactions between systems then system m has wave-function (1)

$$\Psi^m = \sum_{n=1}^N \bar{c}_n^m \Psi_n \tag{9}$$

where probability $\bar{w}_n^m = |\bar{c}_n^m|^2$ is independent of time t, but otherwise arbitrary. In natural fact the electron e_m experiences the Coulomb repulsion V^m of its M-1 neighbours $e_{m'\neq m}$:

$$dV^m = \sum_{m' \neq m}^{N} \frac{e_m de_{m'}}{r_{mm'}} \tag{10}$$

In Eq.(10) we treat $e_m = e$ classically as a point-charge electron, and $e_{m'}$ quantum mechanically: $de_{m'} = e|\psi^{m'}(x_{m'})|^2 dx_{m'}$ is the infinitesimal charge within $dx_{m'}$ of system m', so with $r_{mm'}$ of Eq.(8),

$$V^{m} = \sum_{m' \neq m}^{N} e_{m} \int_{0}^{a} \frac{de_{m'}}{r_{mm'}} = e^{2} \sum_{m' \neq m}^{N} \int_{0}^{a} \frac{|\psi^{m'}(x_{m'})|^{2} dx_{m'}}{\sqrt{(x_{m} - x_{m'})^{2} + 4R^{2} \sin^{2}\left(\frac{\pi(m - m')}{M}\right)}}$$
(11)

With, c.f. Eq.(1,9),

$$\Psi^{m'} = \sum_{n=1}^{N} c_n^{m'} \Psi_n$$
 (12)

we finally have that the time-dependent perturbation on system m = 1..M caused by its M - 1 neighbour systems is

$$V^{m}(x_{m},t) = e^{2} \sum_{n_{1},n_{2}}^{N} \sum_{m' \neq m}^{M} c_{n_{1}}^{m'}(t) c_{n_{2}}^{m'}(t)^{*} \int_{0}^{a} \frac{\Psi_{n_{1}}^{m'}(x_{m'},t)\Psi_{n_{2}}^{m'}(x_{m'},t)^{*} dx_{m'}}{\sqrt{(x_{m}-x_{m'})^{2} + 4R^{2} \sin^{2}\left(\frac{\pi(m-m')}{M}\right)}}$$
(13)

2.3 Perturbation calculations

We summarise nearly 100 year-old time-dependent QM perturbation theory, with general H_0 , ψ_n and V(t). The exact solution to the TDSE $i\hbar\partial\Psi/\partial t = (H_0 + V(t))\Psi$ is $\Psi = \sum_n c_n\Psi_n$, where [L&L]

$$\dot{c}_n = \frac{1}{i\hbar} \sum_{n'} c_{n'} \int \Psi_{n'}^*(\mathbf{r}, t) V(\mathbf{r}, t) \Psi_n(\mathbf{r}, t) d\mathbf{r}.$$
(14)

With correspondence $(H_0, \psi_n, \Psi_n, c_n, V, d\mathbf{r}) \Rightarrow (H_0^m, \psi_n^m, \Psi_{n,1,1}^m, c_n^m, V^m, dxdydz)$, substitution of Eqns(4,7,8) into Eqn(14) and evaluation of the $\int \int ..dydz$ integral give

$$\dot{c}_{n}^{m}(t) = \frac{4e^{2}a^{2}}{i\hbar} \sum_{n',n_{1},n_{2}}^{N} e^{i(\omega_{n}-\omega_{n'}+\omega_{n_{2}}-\omega_{n_{1}})t} c_{n'}^{m}(t) \sum_{m'\neq m}^{M} c_{n_{1}}^{m'}(t) c_{n_{2}}^{m'}(t)^{*} \int_{0}^{a} \int_{0}^{a} \frac{dxdx' \sin(\pi nx/a) \sin(\pi n'x/a) \sin(\pi n_{1}x'/a) \sin(\pi n_{2}x'/a)}{\sqrt{(x-x')^{2}+4R^{2} \sin^{2}\left(\frac{\pi(m-m')}{M}\right)}}$$
(15)

Also, to conventionally manipulate the differential set Eq.(15), we need to replace the integer pair (m, n) by the single integer p: counting in mixed base (M, N) gives p = m + (n-1)M. Eq.(15) then becomes

$$\dot{c}_{p(m,n)} = \frac{4e^2a^2}{i\hbar} \sum_{n',n1,n2}^{N} c_{p(m,n')} \sum_{m'\neq m}^{M} c_{p(m',n_1)} c_{p(m',n_2)}^* e^{i(\omega_n - \omega_{n'} + \omega_{n_2} - \omega_{n_1})t}$$

$$\int_0^a \int_0^a \frac{dxdx' \sin(\pi nx/a) \sin(\pi n'x/a) \sin(\pi n_1 x'/a) \sin(\pi n_2 x'/a)}{\sqrt{(x-x')^2 + 4R^2 \sin^2(\pi (m-m')/M)}}$$

$$p = p(m,n) = m + (n-1)M \qquad m = 1..M \qquad n = 1..N \qquad p = 1..MN$$
(16)

To solve/use Eq.(16) we must give them initial values $\bar{c}_p = c_p(t=0)$, let them solve until time t, and then examine the probabilities $w_p(t) = |c_p(t)|^2$. A typical choice of initial values is $\bar{c}_p = 1/\sqrt{N}$: all initial probabilities are equal, which incidentally corresponds to a temperature $T = \infty$.

Note that Eq.(15) is non-linear, $c_p c_{p'} c_{p''} *$ occurs on the r.h.s. We are still using the traditional linear Schrödinger equation Eq.(14) and the non-linearity arises from our formulation §2.2 of the perturbation V(t).

If we put $d_n^m(t) = c_n^m(t)e^{-i\omega_n t}$, use $\dot{c}_n^m e^{-i\omega_n t} = \dot{d}_n^m + i\omega_n d_n^m$ in (16), and revert $d \to c$, we obtain the 'Interaction Representation' (IR):

$$\dot{c}_{p(m,n)} = -i\omega_n c_{p(m,n)} + \frac{4e^2a^2}{i\hbar} \sum_{n',n1,n2}^N c_{p(m,n')} \sum_{m'\neq m}^M c_{p(m',n_1)} c_{p(m',n_2)}^* \int_0^a \int_0^a \frac{dxdx' \sin(\pi nx/a) \sin(\pi n'x/a) \sin(\pi n_1 x'/a) \sin(\pi n_2 x'/a)}{\sqrt{(x-x')^2 + 4R^2 \sin^2(\pi (m-m')/M)}},$$
(17)

which may be useful in numerical evaluation.

2.4 Symmetry reductions

We shall see that solutions $w_p(t)$ of Eq.(16) are often "steady-state oscillatory". We assume that any physical measurent must last a (small) finite time Δt , so that any measured or inferred physical probability $W_p(t)$ is an an average over many "oscillations",

$$W_p(t) = \frac{1}{\Delta t} \int_t^{t+\Delta t} w_p(t') dt'$$
(18)

One might call (16) coarse-graining, a form of RPA: but this occurs at time t, and there is no RPA from $0 \rightarrow t$ with evolution following (16).

2.4.1 P1: Finite M individual systems, c_n^m

This is Eq.(16) which we recall for convenience:

$$\dot{c}_{p(m,n)} = \frac{4e^2a^2}{i\hbar} \sum_{n',n_1,n_2}^{N} c_{p(m,n')} \sum_{m' \neq m}^{M} c_{p(m',n_1)} c_{p(m',n_2)}^* e^{i(\omega_n - \omega_{n'} + \omega_{n_2} - \omega_{n_1})t} \int_0^a \int_0^a \frac{dxdx' \sin(\pi nx/a) \sin(\pi n'x/a) \sin(\pi n_1 x'/a) \sin(\pi n_2 x'/a)}{\sqrt{(x - x')^2 + 4R^2 \sin^2(\pi (m - m')/M)}}$$
(19)
$$p = p(m,n) = m + (n-1)M \qquad m = 1..M \qquad n = 1..N \qquad p = 1..MN.$$

2.4.2 P2: Finite M similar systems, c_n

We now exploit the physical symmetry of our model to simplify the complex amplitude c_n^m . We start all systems with the same initial probability, $w_n^m(0) = w_n(0)$ and assume/postulate that this holds for all future t, $w_n^m(t) = w_n(t)$. Since the environment of each site m = 1..M is the same and each interacts with the M - 1 others, we can expect the probability to be the same; $w_n^m = w_n^{m'} = w_n$. With $c_n^m \equiv r_n^m e^{i\theta_n^m}$ (pure math polar form, real r and θ , no phys.), this means that $w_n = (r_n^m)^2 \equiv r_n^2$. We postulate that $\theta_n^m = \theta_n(t) + \theta^m(0)$ where θ^m is time-independent, reflecting the static invariance of our ring geometry. We now have $c_n^m(t) = r_n(t)e^{i\theta_n(t)}e^{i\theta^m}$ whose substitution in Eq.(15 or 16) eliminates $e^{i\theta^m}$:

$$c_n(t) = r_n(t)e^{i\theta_n(t)} \tag{20}$$

We can show that in Eq.(16) $\sum_{m'\neq m}^{M} \to \sum_{m'=1}^{M-1}$ (independent of m), we 'normalise' $\int_{0}^{a} \to \int_{0}^{1}$, finally giving

$$\dot{c}_{n} = \frac{4e^{2}}{i\hbar} \sum_{n',n_{1},n_{2}}^{N} c_{n'}c_{n_{1}}c_{n_{2}}^{*}e^{i(\omega_{n}-\omega_{n'}+\omega_{n_{2}}-\omega_{n_{1}})t}$$

$$\int_{0}^{1} \int_{0}^{1} \sum_{m'=1}^{M-1} \frac{dxdx'\sin(\pi nx)\sin(\pi n'x)\sin(\pi n_{1}x')\sin(\pi n_{2}x')}{\sqrt{a^{2}(x-x')^{2}+4R^{2}\sin^{2}(\pi m'/M)}}$$

$$m = 1..M \qquad n = 1..N$$
(21)

2.4.3 P3: Infinite M similar systems, c_n

We are interested in a system in an infinite $M \to \infty$ heat-bath, and the sum

$$\sum_{m'=1}^{M-1} \frac{1}{\sqrt{a^2(x-x')^2 + 4R^2 \sin^2(\pi m'/M)}}$$

in Eq.(20) can be approximated by an integral $\int_1^{\infty} ..dm'$ which Mathematica 7 gives as an Elliptic EK =

$$EllipticK\left(\frac{-4R^2}{a^2(x-x')^2}\right).$$

Eq.(20) then becomes

$$\dot{c}_{n} = \frac{4e^{2}}{i\hbar} \sum_{n',n_{1},n_{2}}^{N} c_{n'} c_{n_{1}} c_{n_{2}}^{*} e^{i(\omega_{n}-\omega_{n'}+\omega_{n_{2}}-\omega_{n_{1}})t}$$
$$\int_{0}^{1} \int_{0}^{1} dx dx' \sin(\pi nx) \sin(\pi n'x) EK\left(\frac{-4R^{2}}{a^{2}(x-x')^{2}}\right) \sin(\pi n_{1}x') \sin(\pi n_{2}x') \qquad (22)$$
$$n = 1..N$$

3 Numerical results

We have implemented the above analysis in Mathematica 12 to give numerical results, particularly the measured probability

$$W_p(t) = \frac{1}{\Delta t} \int_t^{t+\Delta t} w_p(t') dt'.$$
(23)

A typical result for initial $\bar{c}_p = c_p(t=0) = 1/\sqrt{(N)}$, N = 3 and M = 3 of actual (instaneous) $w_p(t)$ and measured (experimental) $W_p(t)$ probabilities is shown in Fig.2. Visually, the intantaneous oscillatory chaos is converted to experimentally observed constant results.

Numerically, with energies $E_n \sim n^2$ the ratio $(E_1 - E_2)/(E_2 - E_3)$ is 0.6 theoretically; physically we use (2) and $Log(w_n)$, see Fig.2, which give (0.78 - 1.10)/(1.10 - 1.59) = 0.65

Similar results are obtained for variations of N, M, a, R, the three cases of §2 and the important initial conditions $c_n^m(t=0)$.

4 Discussion

The preceding may be regarded as a useful direct quantitative pedagogic illustration of the quantum *density matrix* (statistical operator), see Landau & Lifshitz [1].

To broadly summarize: at t = 0 our system is in an arbitrary initial condition, whence it evolves as in (16) (with no random phase assumptions) to a Boltzman distribution.

5 References

[1] Landau L.and Lifschitz E. (1977), Quantum Mechanics 3e, Pergamon Press.

This figure "figg1.jpg" is available in "jpg" format from:

http://arxiv.org/ps/2006.12317v2

This figure "figg2.jpg" is available in "jpg" format from:

http://arxiv.org/ps/2006.12317v2