

# BILINEAR MULTIPLIERS OF SMALL LEBESGUE SPACES

ÖZNUR KULAK AND A.TURAN GÜRKANLI

**ABSTRACT.** Let  $G$  be a locally compact abelian metric group with Haar measure  $\lambda$  and  $\hat{G}$  its dual with Haar measure  $\mu$ , and  $\lambda(G)$  is finite. Assume that  $1 < p_i < \infty$ ,  $p'_i = \frac{p_i}{p_i - 1}$ , ( $i = 1, 2, 3$ ) and  $\theta \geq 0$ . Let  $L^{(p'_i, \theta)}(G)$ , ( $i = 1, 2, 3$ ) be small Lebesgue spaces. A bounded measurable function  $m(\xi, \eta)$  defined on  $\hat{G} \times \hat{G}$  is said to be a bilinear multiplier on  $G$  of type  $[(p'_1; (p'_2; (p'_3)]_\theta$  if the bilinear operator  $B_m$  associated with the symbol  $m$

$$B_m(f, g)(x) = \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) m(s, t) \langle s + t, x \rangle$$

defines a bounded bilinear operator from  $L^{(p'_1, \theta)}(G) \times L^{(p'_2, \theta)}(G)$  into  $L^{(p'_3, \theta)}(G)$ . We denote by  $BM_\theta[(p'_1; (p'_2; (p'_3)]$  the space of all bilinear multipliers of type  $[(p'_1; (p'_2; (p'_3)]_\theta$ . In this paper, we discuss some basic properties of the space  $BM_\theta[(p'_1; (p'_2; (p'_3)]$  and give examples of bilinear multipliers.

## 1. INTRODUCTION

Let  $\Omega$  be locally compact Hausdorff space and let  $(\Omega, B, \mu)$  be finite Borel measure space. The grand Lebesgue space  $L^{p^{\circ}}(\Omega)$  is defined by the norm

$$\|f\|_{p^{\circ}} = \sup_{0 < \varepsilon \leq p-1} \left( \varepsilon \int_{\Omega} |f|^{p-\varepsilon} d\mu \right)^{\frac{1}{p-\varepsilon}}$$

where  $1 < p < \infty$ , [7]. A generalization of the grand Lebesgue spaces are the spaces  $L^{p^{\circ}, \theta}(\Omega)$ ,  $\theta \geq 0$ , defined by the norm

$$\|f\|_{p^{\circ}, \theta, \Omega} = \sup_{0 < \varepsilon \leq p-1} \varepsilon^{\frac{\theta}{p-\varepsilon}} \left( \int_{\Omega} |f|^{p-\varepsilon} d\mu \right)^{\frac{1}{p-\varepsilon}} = \sup_{0 < \varepsilon \leq p-1} \varepsilon^{\frac{\theta}{p-\varepsilon}} \|f\|_{p-\varepsilon};$$

when  $\theta = 0$  the space  $L^{p^{\circ}, 0}(\Omega)$  reduces to Lebesgue space  $L^p(\Omega)$  and when  $\theta = 1$  the space  $L^{p^{\circ}, 1}(\Omega)$  reduces to grand Lebesgue space  $L^{p^{\circ}}(\Omega)$ , (see, [3]). For  $0 < \varepsilon \leq p-1$ ,

$$L^p(\Omega) \subset L^{p^{\circ}, \theta}(\Omega) \subset L^{p-\varepsilon}(\Omega)$$

hold. It is known that the subspace  $C_c^\infty(\Omega)$  is not dense in  $L^{p^{\circ}, \theta}(\Omega)$ , where  $C_c^\infty(\Omega)$  is the space of infinitely differentiable complex valued functions defined on  $\Omega$  with compact support. Its closure of consists of functions  $f \in L^p(\Omega)$  such that

$$\lim_{\varepsilon \rightarrow 0} \varepsilon^{\frac{\theta}{p-\varepsilon}} \|f\|_{p-\varepsilon} = 0, [3].$$

---

2000 *Mathematics Subject Classification.* Primary 46E20; Secondary 42A45.

*Key words and phrases.* Bilinear multipliers, Grand Lebesgue spaces, Small Lebesgue spaces.

For some properties and applications of  $L^p(\Omega)$ , we refer to papers [2], [4], [5] and [6].

Let  $p' = \frac{p}{p-1}$ ,  $1 < p < \infty$ . First consider an auxiliary space namely  $L^{(p',\theta)}(\Omega)$ ,  $\theta \geq 0$ , defined by

$$\|g\|_{(p',\theta)} = \inf_{g=\sum_{k=1}^{\infty} g_k} \left\{ \sum_{k=1}^{\infty} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} \left( \int_{\Omega} |g_k(x)|^{(p-\varepsilon)'} d(x) \right)^{\frac{1}{(p-\varepsilon)'}} \right\}$$

where the functions  $g_k$ ,  $k \in \mathbb{N}$ , being in  $M_0$ , the set of all real valued measurable functions, finite a.e. in  $\Omega$ . After this definition the generalized small Lebesgue spaces have been defined by

$$L^{(p)',\theta}(\Omega) = \left\{ g \in M_0 : \|g\|_{(p)',\theta} < \infty \right\},$$

where

$$\|g\|_{(p)',\theta} = \sup_{\substack{0 \leq \psi \leq |g| \\ \psi \in L^{(p)',\theta}(\Omega)}} \|\psi\|_{(p',\theta)}.$$

For  $\theta = 0$  it is  $\|f\|_{(p',0)} = \|f\|_{(p)',\theta}$ , [2], [6].

Let  $G$  be a locally compact abelian metric group with Haar measure  $\lambda$  and let  $\hat{G}$  be dual group with Haar measure  $\mu$ . The translation and modulation operators are given by

$$T_x f(t) = f(t-x), M_{\xi} f(t) = \langle t, \xi \rangle f(t), t, x \in G, \xi \in \hat{G}.$$

For a function  $f \in L^1(G)$ , the function  $\hat{f}$  defined on  $\hat{G}$  by

$$\hat{f}(\gamma) = \int_G f(x) \langle \gamma, -x \rangle d\lambda(x), \gamma \in \hat{G}$$

is called the Fourier transform of  $f$ , [12].

## 2. MAIN RESULTS

Let  $G$  be a locally compact abelian metric group and  $\hat{G}$  its dual with Haar measures  $\lambda$  and  $\mu$  respectively. Before give the definition of bilinear multiplier on  $G$  of type  $[(p'_1; (p'_2; (p'_3)_{\theta})]$ , we remember that if  $\lambda(G)$  is finite, then  $G$  is compact. Thus the  $\hat{G}$  dual of  $G$  (Pontryagin dual) is a discrete group, and the dual measure on this group is the counting measure. Also since  $G$  is compact abelian metric group is  $\hat{G}$  is countable, [12].

**Definition 1.** Let  $G$  be a locally compact abelian metric group with Haar measure  $\lambda$  and  $\hat{G}$  its dual with Haar measure  $\mu$ , and  $\lambda(G)$  is finite. Assume that  $1 < p_i < \infty$ ,  $p'_i = \frac{p_i}{p_i-1}$ ,  $(i = 1, 2, 3)$  and  $\theta \geq 0$ . We also assume that  $m(s, t)$  is a bounded measurable function on  $\hat{G} \times \hat{G}$ . Consider the bilinear operator  $B_m$  associated with the symbol  $m$

$$B_m(f, g)(x) = \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) m(s, t) \langle s + t, x \rangle,$$

defined for functions  $f, g \in C^\infty(G)$ .  $m$  is said to be a bilinear multiplier on  $G$  of type  $[(p'_1; (p'_2; (p'_3)_\theta)]$ , if there exists  $C > 0$  such that

$$\|B_m(f, g)\|_{(p'_3, \theta)} \leq C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}$$

for all  $f, g \in C^\infty(G)$ . That means  $B_m$  extends to a bounded bilinear operator from  $L^{(p'_1, \theta)}(G) \times L^{(p'_2, \theta)}(G)$  into  $L^{(p'_3, \theta)}(G)$ . We denote by  $BM_\theta[(p'_1; (p'_2; (p'_3))_\theta)]$  the space of all bilinear multipliers of type  $[(p'_1; (p'_2; (p'_3)_\theta)]$  and

$$\|m\|_{[(p'_1; (p'_2; (p'_3)_\theta)]} = \|B_m\|.$$

**Lemma 1.** (Generalized Hölder inequality for generalized small Lebesgue spaces)

Let  $\frac{1}{p'} = \frac{1}{p'_1} + \frac{1}{p'_2}$  and  $r.r' < p' + 1$ . If  $f \in L^{(p'_1, \theta)}(G)$  and  $g \in L^{(p'_2, \theta)}(G)$ , then  $fg \in L^{(r', \theta)}(G)$ . Furthermore

$$\|fg\|_{(r', \theta)} \leq C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}$$

for some  $C > 0$ .

*Proof.* Take any  $f \in C^\infty(G) \subset L^{(p'_1, \theta)}(G)$  and  $g \in C^\infty(G) \subset L^{(p'_2, \theta)}(G)$ . Let  $\frac{1}{p'} = \frac{1}{p'_1} + \frac{1}{p'_2}$  and  $r.r' < p' + 1$ . Since  $1 = \frac{1}{r} + \frac{1}{r'}$ , we have  $r + r' = r.r'$ . Then using the assumption  $r.r' < p' + 1$ , we write  $r + r' < p' + 1$  and so  $r + r' - 1 < p'$ . For a fixed  $0 < \varepsilon \leq r - 1$ , we have  $r' + \varepsilon \leq r' + r - 1 < p'$ . Therefore since  $\mu(G) < \infty$ , we obtain  $L^{p'}(G) \subset L^{r'+\varepsilon}(G)$ . Also we know that  $L^{r'+\varepsilon}(G) \subset L^{(r', \theta)}(G)$ , [2]. Then we have the inclusion  $L^{p'}(G) \subset L^{r'+\varepsilon}(G) \subset L^{(r', \theta)}(G)$ . That means, there exists  $C_1 > 0$  such that

$$(2.1) \quad \|fg\|_{(r', \theta)} \leq C_1 \|fg\|_{p'}.$$

If we apply the Hölder inequality to the right side of (2.1), there exists  $C_2 > 0$  such that

$$(2.2) \quad \|fg\|_{p'} \leq C_2 \|f\|_{p'_1} \|g\|_{p'_2}.$$

On the other hand, since  $L^{(p'_1, \theta)}(G) \subset L^{p'_1}(G)$  and  $L^{(p'_2, \theta)}(G) \subset L^{p'_2}(G)$ , [2], we have

$$(2.3) \quad \|f\|_{p'_1} \leq C_3 \|f\|_{(p'_1, \theta)}$$

and

$$(2.4) \quad \|g\|_{p'_2} \leq C_4 \|g\|_{(p'_2, \theta)}$$

for some  $C_3, C_4 > 0$ . Combining the inequalities (2.1), (2.2), (2.3) and (2.4), we obtain

$$(2.5) \quad \|fg\|_{(r', \theta)} \leq C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}$$

where  $C = C_1 C_2 C_3 C_4$ . Now define the bilinear mapping  $F((f, g)) = fg$ , from  $(C^\infty \times C^\infty)(G)$  to  $L^{(r', \theta)}(G)$ . By (2.5), it is continuous. Since  $(C^\infty \times C^\infty)(G)$  is dense in  $L^{(p'_1, \theta)}(G) \times L^{(p'_2, \theta)}(G)$ , then there exists has a unique continuous bilinear extension of  $F$  denoted  $F^\sim$  from  $L^{(p'_1, \theta)}(G) \times L^{(p'_2, \theta)}(G)$  to  $L^{(r', \theta)}(G)$ . Furthermore, the norm of  $F^\sim$  is equal to the norm of  $F$ . Therefore, for all  $f \in L^{(p'_1, \theta)}(G)$  and  $g \in L^{(p'_2, \theta)}(G)$ , the inequality

$$\|fg\|_{(r', \theta)} = \|F((f, g))\|_{(r', \theta)} \leq C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}$$

is achieved.  $\square$

**Example 1.** Let  $f \in L^{(9,\theta)}(G)$  and  $g \in L^{(10,\theta)}(G)$ . Since  $p'_1 = 9$  and  $p'_2 = 10$ , then  $\frac{1}{p'} = \frac{1}{p'_1} + \frac{1}{p'_2} = \frac{1}{9} + \frac{1}{10}$ , and so  $p' = \frac{90}{19}$ . Also let  $r = 3$  and  $r' = \frac{3}{2}$ . Then  $\frac{1}{r} + \frac{1}{r'} = 1$ . Hence we have  $r.r' = 3.\frac{3}{2} = \frac{9}{2} < \frac{90}{19} + 1 = p' + 1$ . Therefore from the Lemma 1, we obtain that  $fg \in L^{(\frac{3}{2},\theta)}(G)$  and

$$\|fg\|_{(\frac{3}{2},\theta)} \leq C \|f\|_{(9,\theta)} \|g\|_{(10,\theta)}$$

for some  $C > 0$ .

**Theorem 1.** Let  $1 < p_i < \infty$ ,  $p'_i = \frac{p_i}{p_i-1}$ , ( $i = 1, 2, 3$ ) and  $\theta > 0$ . Then  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$  if and only if there exists  $C > 0$  such that

$$\left| \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \hat{h}(s+t) m(s, t) \right| \leq C \|f\|_{(p'_1,\theta)} \|g\|_{(p'_2,\theta)} \|h\|_{(p_3),\theta}$$

for all  $f \in L^{(p'_1,\theta)}(G)$ ,  $g \in L^{(p'_2,\theta)}(G)$  and  $h \in [L^{p_3}]_{(p_3),\theta}$ , where  $[L^{p_3}]_{(p_3),\theta}$  is the closure of  $C^\infty(G)$  in  $L^{p_3),\theta}(G)$ .

*Proof.* Assume that  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ . Let  $f, g \in C^\infty(G) \subset L^1(G)$  and  $h \in C^\infty(G) \cap L^{p_3),\theta}(G)$ . Since  $\mu(G) < \infty$ , we have  $h \in L^1(G)$ . So, we write

$$\begin{aligned} & \left| \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \hat{h}(s+t) m(s, t) \right| \\ &= \left| \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left\{ \int_G h(y) \langle s+t, -y \rangle d\lambda(y) \right\} m(s, t) \right| \\ &= \left| \int_G h(y) B_m(f, g)(-y) d\lambda(y) \right| = \left| \int_G h(y) \tilde{B}_m(f, g)(y) d\lambda(y) \right| \\ (2.6) \quad & \leq \int_G |h(y)| |\tilde{B}_m(f, g)(y)| d\lambda(y) \end{aligned}$$

where  $\lambda$  is Haar measure on  $G$  and  $\tilde{B}_m(f, g)(y) = B_m(f, g)(-y)$ . From the assumption  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ , then  $B_m(f, g) \in L^{(p'_3,\theta)}(G)$ . Since  $G$  is group, we obtain  $\tilde{B}_m(f, g) \in L^{(p'_3,\theta)}(G)$ . By using the Hölder inequality for generalized small Lebesgue spaces and the inequality (2.6), we write

$$\begin{aligned} & \left| \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \hat{h}(s+t) m(s, t) \right| \leq \int_G |h(y)| |\tilde{B}_m(f, g)(y)| d\lambda(y) \\ (2.7) \quad & \leq \|\tilde{B}_m(f, g)\|_{(p'_3,\theta)} \|h\|_{(p_3),\theta} = \|B_m(f, g)\|_{(p'_3,\theta)} \|h\|_{(p_3),\theta} \end{aligned}$$

Also since  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ , there exists  $C > 0$  such that

$$(2.8) \quad \|B_m(f, g)\|_{(p'_3,\theta)} \leq C \|f\|_{(p'_1,\theta)} \|g\|_{(p'_2,\theta)}$$

Combining (2.7) and (2.8), we find

$$\left| \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \hat{h}(s+t) m(s, t) \right| \leq C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \|h\|_{(p_3), \theta}.$$

For the proof of converse, assume that there exists a constant  $C > 0$  such that

$$\left| \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \hat{h}(s+t) m(s, t) \right| \leq C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \|h\|_{(p_3), \theta}$$

for all  $f, g \in C^\infty(G)$  and  $h \in C^\infty(G) \cap L^{p_3), \theta}(G)$ . From the assumption and (2.7), we write

$$(2.9) \quad \left| \int_G h(y) \tilde{B}_m(f, g)(y) d\lambda(y) \right| \leq C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \|h\|_{(p_3), \theta}.$$

Define a function  $\ell$  from  $C^\infty(G) \cap L^{p_3), \theta}(G)$  to  $\mathbb{C}$  such that

$$\ell(h) = \int_G h(y) \tilde{B}_m(f, g)(y) d\lambda(y).$$

It is clear that the function  $\ell$  is linear and bounded by (2.9). Since  $\overline{C^\infty(G) \cap L^{p_3), \theta}(G)} = [L^{p_3}]_{(p_3), \theta}$ . Then  $\ell \in ([L^{p_3}]_{(p_3), \theta})^* = L^{p_3)', \theta}(G) \simeq L^{(p_3'), \theta}(G)$  and by (2.9), we have

$$\begin{aligned} \|B_m(f, g)\|_{(p'_3, \theta)} &= \left\| \tilde{B}_m(f, g) \right\|_{(p'_3, \theta)} = \|\ell\| = \sup_{\|h\|_{(p_3), \theta} \leq 1} \frac{|\ell(h)|}{\|h\|_{(p_3), \theta}} \\ &\leq \sup_{\|h\|_{(p_3), \theta} \leq 1} \frac{C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \|h\|_{p'_3, \omega_3^{-1}}}{\|h\|_{(p_3), \theta}} \leq C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}. \end{aligned}$$

Hence, we obtain  $m \in BM_\theta[(p'_1; (p'_2; (p'_3)]$

□

**Theorem 2.** Let  $m \in BM_\theta[(p'_1; (p'_2; (p'_3)]$ . Then

a)  $T_{(s_0, t_0)} m \in BM_\theta[(p'_1; (p'_2; (p'_3)]$  for each  $(s_0, t_0) \in \hat{G} \times \hat{G}$  and

$$\|T_{(s_0, t_0)} m\|_{[(p'_1; (p'_2; (p'_3)]_\theta} = \|m\|_{[(p'_1; (p'_2; (p'_3)]_\theta}.$$

b)  $M_{t_0}^2 M_{s_0}^1 m \in BM_\theta[(p'_1; (p'_2; (p'_3)]$  for each  $(s_0, t_0) \in G \times G$  and

$$\|M_{t_0}^2 M_{s_0}^1 m\|_{[(p'_1; (p'_2; (p'_3)]_\theta} = \|m\|_{[(p'_1; (p'_2; (p'_3)]_\theta}$$

where  $M_{s_0}^1 m(s, t) = \langle s, s_0 \rangle m(s, t)$  and  $M_{t_0}^2 m(s, t) = \langle t, t_0 \rangle m(s, t)$ .

*Proof.* a) Let  $f, g \in C^\infty(G)$ . Let say  $s - s_0 = u$  and  $t - t_0 = v$ . Then

$$\begin{aligned} B_{T_{(s_0, t_0)} m}(f, g)(x) &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) T_{(s_0, t_0)} m(s, t) \langle s + t, x \rangle \\ &= \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} \hat{f}(u + s_0) \hat{g}(v + t_0) m(u, v) \langle u + s_0 + v + t_0, x \rangle \\ &= \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} \hat{f}(u + s_0) \hat{g}(v + t_0) m(u, v) \langle (s_0 + t_0) + (u + v), x \rangle \end{aligned}$$

$$(2.10) \quad = \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} T_{-s_0} \hat{f}(u) T_{-t_0} \hat{g}(v) m(u, v) \langle s_0 + t_0, x \rangle \langle u + v, x \rangle$$

It is known that

$$\begin{aligned} (M_{-s_0} f) \wedge(u) &= \int_G M_{-s_0} f(y) \langle u, -y \rangle d\lambda(y) \\ &= \int_G \langle -s_0, y \rangle f(y) \langle u, -y \rangle d\lambda(y) = \int_G \langle s_0, -y \rangle f(y) \langle u, -y \rangle d\lambda(y) \\ (2.11) \quad &= \int_G f(y) \langle s_0 + u, -y \rangle d\lambda(y) = \hat{f}(s_0 + u) = T_{-s_0} \hat{f}(u). \end{aligned}$$

Similarly we obtain  $(M_{-t_0} g) \wedge = T_{-t_0} \hat{g}$ . Combining (2.10) and (2.11), we have

$$\begin{aligned} B_{T_{(s_0, t_0)} m}(f, g)(x) &= \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} T_{-s_0} \hat{f}(u) T_{-t_0} \hat{g}(v) m(u, v) \langle s_0 + t_0, x \rangle \langle s + t, x \rangle \\ &= \langle s_0 + t_0, x \rangle \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} (M_{-s_0} f) \wedge(u) (M_{-t_0} g) \wedge(v) m(u, v) \langle s + t, x \rangle \\ (2.12) \quad &= \langle s_0 + t_0, x \rangle B_m(M_{-s_0} f, M_{-t_0} g)(x). \end{aligned}$$

Using the assumption  $m \in BM_\theta[(p'_1; (p'_2; (p'_3)]$  and the equality (2.12) we have

$$\begin{aligned} \|B_{T_{(s_0, t_0)} m}(f, g)\|_{(p'_3, \theta)} &= \|\langle s_0 + t_0, x \rangle B_m(M_{-s_0} f, M_{-t_0} g)\|_{(p'_3, \theta)} \\ &= \|B_m(M_{-s_0} f, M_{-t_0} g)\|_{(p'_3, \theta)} \\ &\leq C \|M_{-s_0} f\|_{(p'_1, \theta)} \|M_{-t_0} g\|_{(p'_2, \theta)} \\ &= C \|\langle -s_0, \cdot \rangle f(\cdot)\|_{(p'_1, \theta)} \|\langle -t_0, \cdot \rangle g(\cdot)\|_{(p'_2, \theta)} \\ &= C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \end{aligned}$$

for some  $C > 0$ . Thus  $T_{(s_0, t_0)} m \in BM_\theta[(p'_1; (p'_2; (p'_3)]$ . Then by definition 1,

$$\|T_{(s_0, t_0)} m\|_{[(p'_1; (p'_2; (p'_3)]_\theta} = \|B_{T_{(s_0, t_0)} m}\|.$$

This implies

$$\begin{aligned} \|T_{(s_0, t_0)} m\|_{[(p'_1; (p'_2; (p'_3)]_\theta} &= \|B_{T_{(s_0, t_0)} m}\| \\ &= \sup \left\{ \frac{\|B_{T_{(s_0, t_0)} m}(f, g)\|_{(p'_3, \theta)}}{\|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}} : \|f\|_{(p'_1, \theta)} \leq 1, \|g\|_{(p'_2, \theta)} \leq 1 \right\} \\ &= \sup \left\{ \frac{\|B_m(M_{-s_0} f, M_{-t_0} g)\|_{(p'_3, \theta)}}{\|M_{-s_0} f\|_{(p'_1, \theta)} \|M_{-t_0} g\|_{(p'_2, \theta)}} : \|M_{-s_0} f\|_{(p'_1, \theta)} \leq 1, \|M_{-t_0} g\|_{(p'_2, \theta)} \leq 1 \right\} \\ &= \|B_m\| = \|m\|_{[(p'_1; (p'_2; (p'_3)]_\theta}. \end{aligned}$$

b) Let  $f, g \in C^\infty(G)$ . We have

$$\begin{aligned}
(2.13) \quad B_{M_{t_0}^2 M_{s_0}^1 m}(f, g)(x) &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) M_{t_0}^2 M_{s_0}^1(m(s, t)) \langle s + t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) M_{t_0}^2(\langle s, s_0 \rangle m(s, t)) \langle s + t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \langle t, t_0 \rangle \langle s, s_0 \rangle m(s, t) \langle s + t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \langle s, s_0 \rangle \hat{f}(s) \langle t, t_0 \rangle \hat{g}(t) m(s, t) \langle s + t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} M_{s_0} \hat{f}(s) M_{t_0} \hat{g}(t) m(s, t) \langle s + t, x \rangle
\end{aligned}$$

If we put  $y + s_0 = u$ , we have

$$\begin{aligned}
(T_{-s_0} f) \hat{\ } (s) &= \int_G T_{-s_0} f(y) \langle s, -y \rangle d\lambda(y) \\
&= \int_G f(y + s_0) \langle s, -y \rangle d\lambda(y) = \int_G f(u) \langle s, -u + s_0 \rangle d\lambda(u) \\
&= \int_G f(u) \langle s, -u \rangle \langle s, s_0 \rangle d\lambda(u) = \langle s, s_0 \rangle \int_G f(u) \langle s, -u \rangle d\lambda(u) \\
&= \langle s, s_0 \rangle \hat{f}(s) = M_{s_0} \hat{f}(s).
\end{aligned}$$

Similarly  $(T_{-t_0} g) \hat{\ } = M_{t_0} \hat{g}$ . Then from (2.13)

$$B_{M_{t_0}^2 M_{s_0}^1 m}(f, g)(x) = \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} (T_{-s_0} f) \hat{\ } (s) (T_{-t_0} g) \hat{\ } (t) m(s, t) \langle s + t, x \rangle = B_m(T_{-s_0} f, T_{-t_0} g)(x).$$

Since  $m \in BM_\theta[(p'_1; (p'_2; (p'_3,$

$$\begin{aligned}
\|B_{M_{t_0}^2 M_{s_0}^1 m}(f, g)\|_{(p'_3, \theta)} &= \|B_m(T_{-s_0} f, T_{-t_0} g)\|_{(p'_3, \theta)} \leq \|B_m\| \|T_{-s_0} f\|_{(p'_1, \theta)} \|T_{-t_0} g\|_{(p'_2, \theta)} \\
&\leq \|B_m\| \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}
\end{aligned}$$

and so  $M_{t_0}^2 M_{s_0}^1 m \in BM_\theta[(p'_1; (p'_2; (p'_3]$ . Finally (2.14), we achieve that

$$\begin{aligned}
\|M_{t_0}^2 M_{s_0}^1 m\|_{[(p'_1; (p'_2; (p'_3)]_\theta} &= \sup \left\{ \frac{\|B_{M_{t_0}^2 M_{s_0}^1 m}(f, g)\|_{(p'_3, \theta)}}{\|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}} : \|f\|_{(p'_1, \theta)} \leq 1, \|g\|_{(p'_2, \theta)} \leq 1 \right\} \\
&= \sup \left\{ \frac{\|B_m(T_{-s_0} f, T_{-t_0} g)\|_{(p'_3, \theta)}}{\|T_{-s_0} f\|_{(p'_1, \theta)} \|T_{-t_0} g\|_{(p'_2, \theta)}} : \|T_{-s_0} f\|_{(p'_1, \theta)} \leq 1, \|T_{-t_0} g\|_{(p'_2, \theta)} \leq 1 \right\} \\
&= \|m\|_{[(p'_1; (p'_2; (p'_3)]_\theta}.
\end{aligned}$$

□

Let  $A$  be an automorphism of  $G$ . The  $\lambda \circ A$  is a nontrivial Haar measure on  $G$ . For any Borel set  $U \subseteq G$ , the modules of  $A$  is defined by  $|A| = \lambda(AU)$  such that  $d\lambda(Ax) = |A| d\lambda(x)$ . Also the adjoint  $A^*$  of  $A$  is an automorphism of  $\hat{G}$ . The adjoint operator  $A^*$  is defined by  $\langle Ax, s \rangle = \langle x, A^*s \rangle$  for  $x \in G$  and  $s \in \hat{G}$ . Furthermore the  $\mu \circ A^*$  is a nontrivial Haar measure on  $\hat{G}$  such that  $d\mu(A^*s) = |A^*| d\mu(s)$ . It is known that  $|A| = |A^*|$ ,  $(A^*)^{-1} = (A^{-1})^*$  and  $|A|^{-1} = |A^{-1}|$ , [1].

**Definition 2.** Let  $A$  be an automorphism of  $G$ . The dilation operator  $D_A^{p'}$  on  $L^{(p',\theta)}(G)$  is defined by

$$D_A^{p'} f(x) = |A|^{\frac{1}{p'}} f(Ax).$$

**Lemma 2.** Let  $A$  be an automorphism of  $G$  and  $f \in L^{(p',\theta)}(G)$ . Then  $D_A^{p'} f \in L^{(p',\theta)}(G)$ . Moreover

$$\left\| D_A^{p'} f \right\|_{(p',\theta)} = |A|^{\frac{1}{p'}} \|f\|_{(p',\theta)} \leq \|f\|_{(p',\theta)}, \text{ if } |A| < 1$$

$$\left\| D_A^{p'} f \right\|_{(p',\theta)} = \|f\|_{(p',\theta)}, \quad \text{if } |A| \geq 1.$$

*Proof.* Let  $A$  be an automorphism of  $G$  and  $f \in L^{(p',\theta)}(G)$ . If we say  $Ax = u$  and use the equality  $|A|^{-1} = |A^{-1}|$ , then

$$\begin{aligned} \left\| D_A^{p'} f \right\|_{(p',\theta)} &= \inf_{\substack{\infty \\ D_A^{p'} f = \sum_{k=1}^{\infty} D_A^{p'} f_k}} \left\{ \sum_{k=1}^{\infty} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} \left( \oint_G \left| D_A^{p'} f_k(x) \right|^{(p-\varepsilon)'} d\lambda(x) \right)^{\frac{1}{(p-\varepsilon)'}} \right\} \\ &= \inf_{\substack{\infty \\ D_A^{p'} f = \sum_{k=1}^{\infty} D_A^{p'} f_k}} \left\{ \sum_{k=1}^{\infty} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} \left( \oint_G \left| |A|^{\frac{1}{p'}} f_k(Ax) \right|^{(p-\varepsilon)'} d\lambda(x) \right)^{\frac{1}{(p-\varepsilon)'}} \right\} \\ &= \inf_{\substack{\infty \\ f = \sum_{k=1}^{\infty} f_k}} \left\{ \sum_{k=1}^{\infty} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} \left( \oint_G \left| |A|^{\frac{1}{p'}} f_k(u) \right|^{(p-\varepsilon)'} d\lambda(A^{-1}u) \right)^{\frac{1}{(p-\varepsilon)'}} \right\} \\ &= \inf_{\substack{\infty \\ f = \sum_{k=1}^{\infty} f_k}} \left\{ \sum_{k=1}^{\infty} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} \left( \oint_G |f_k(u)|^{(p-\varepsilon)'} |A|^{\frac{(p-\varepsilon)'}{p'}} |A|^{-1} d\lambda(u) \right)^{\frac{1}{(p-\varepsilon)'}} \right\} \\ &= \inf_{\substack{\infty \\ f = \sum_{k=1}^{\infty} f_k}} \left\{ \sum_{k=1}^{\infty} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} |A|^{\frac{1}{p'}} |A|^{-\frac{1}{(p-\varepsilon)'}} \left( \oint_G |f_k(u)|^{(p-\varepsilon)'} d\lambda(u) \right)^{\frac{1}{(p-\varepsilon)'}} \right\}. \end{aligned}$$

$$\begin{aligned}
&= \inf_{f=\sum_{k=1}^{\infty} f_k} \left\{ \sum_{k=1}^{\infty} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} |A|^{\frac{1}{p'}} |A|^{\frac{1}{(p-\varepsilon)} - 1} \left( \oint_G |f_k(u)|^{(p-\varepsilon)'} d\lambda(u) \right)^{\frac{1}{(p-\varepsilon)'}} \right\} \\
&= \inf_{f=\sum_{k=1}^{\infty} f_k} \left\{ \sum_{k=1}^{\infty} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} |A|^{1-\frac{1}{p}} |A|^{\frac{1}{(p-\varepsilon)} - 1} \left( \oint_G |f_k(u)|^{(p-\varepsilon)'} d\lambda(u) \right)^{\frac{1}{(p-\varepsilon)'}} \right\} \\
&= \inf_{f=\sum_{k=1}^{\infty} f_k} \left\{ \sum_{k=1}^{\infty} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} |A|^{\frac{1}{(p-\varepsilon)} - \frac{1}{p}} \left( \oint_G |f_k(u)|^{(p-\varepsilon)'} d\lambda(u) \right)^{\frac{1}{(p-\varepsilon)'}} \right\} \\
&\quad (2.15) \\
&= \inf_{f=\sum_{k=1}^{\infty} f_k} \left\{ \sum_{k=1}^{\infty} \inf_{0 < \varepsilon < p-1} |A|^{\frac{1}{(p-\varepsilon)} - \frac{1}{p}} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} \left( \oint_G |f_k(u)|^{(p-\varepsilon)'} d\lambda(u) \right)^{\frac{1}{(p-\varepsilon)'}} \right\}.
\end{aligned}$$

Assume that  $|A| < 1$  and  $0 < \varepsilon < p-1$ , since  $\inf_{0 < \varepsilon < p-1} |A|^{\frac{1}{(p-\varepsilon)} - \frac{1}{p}} = |A|^{1-\frac{1}{p}} = |A|^{\frac{1}{p'}}$ . Then by the last inequality and (2.15), we have

$$\begin{aligned}
\|D_A^{p'} f\|_{(p',\theta)} &= \inf_{f=\sum_{k=1}^{\infty} f_k} \left\{ \sum_{k=1}^{\infty} \inf_{0 < \varepsilon < p-1} |A|^{\frac{1}{(p-\varepsilon)} - \frac{1}{p}} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} \left( \oint_G |f_k(u)|^{(p-\varepsilon)'} d\lambda(u) \right)^{\frac{1}{(p-\varepsilon)'}} \right\} \\
&= \inf_{f=\sum_{k=1}^{\infty} f_k} \left\{ \sum_{k=1}^{\infty} |A|^{\frac{1}{p'}} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} \left( \oint_G |f_k(u)|^{(p-\varepsilon)'} d\lambda(u) \right)^{\frac{1}{(p-\varepsilon)'}} \right\} \\
&= |A|^{\frac{1}{p'}} \inf_{f=\sum_{k=1}^{\infty} f_k} \left\{ \sum_{k=1}^{\infty} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} \left( \oint_G |f_k(u)|^{(p-\varepsilon)'} d\lambda(u) \right)^{\frac{1}{(p-\varepsilon)'}} \right\} = |A|^{\frac{1}{p'}} \|f\|_{(p',\theta)}.
\end{aligned}$$

Thus  $D_A^{p'} f \in L^{(p',\theta)}(G)$ .

Let  $|A| \geq 1$ , and let  $0 < \varepsilon < p-1$ . Since  $\inf_{0 < \varepsilon < p-1} |A|^{\frac{1}{(p-\varepsilon)} - \frac{1}{p}} = 1$ , by (2.15), we have

$$\|D_A^{p'} f\|_{(p',\theta)} = \inf_{f=\sum_{k=1}^{\infty} f_k} \left\{ \sum_{k=1}^{\infty} \inf_{0 < \varepsilon < p-1} |A|^{\frac{1}{(p-\varepsilon)} - \frac{1}{p}} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} \left( \oint_G |f_k(u)|^{(p-\varepsilon)'} d\lambda(u) \right)^{\frac{1}{(p-\varepsilon)'}} \right\}$$

$$= \inf_{f=\sum_{k=1}^{\infty} f_k} \left\{ \sum_{k=1}^{\infty} \inf_{0 < \varepsilon < p-1} \varepsilon^{-\frac{\theta}{p-\varepsilon}} \left( \int_G |f_k(u)|^{(p-\varepsilon)'} d\lambda(u) \right)^{\frac{1}{(p-\varepsilon)'}} \right\} = \|f\|_{(p',\theta)}.$$

Thus  $D_A^{p'} f \in L^{(p',\theta)}(G)$ .  $\square$

**Theorem 3.** Let  $A$  be an automorphism of  $G$  and  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ . If  $\frac{1}{q} = \frac{1}{p'_1} + \frac{1}{p'_2} - \frac{1}{p'_3}$ , then  $\tilde{D}_{A^*}^q m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ , where  $\tilde{D}_{A^*}^q m(s, t) = |A^*|^{\frac{1}{q}} m(A^*s, A^*t)$ . Furthermore

$$\|\tilde{D}_{A^*}^q m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \leq \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]}.$$

*Proof.* Take any  $f \in L^{(p'_1,\theta)}(G)$  and  $g \in L^{(p'_2,\theta)}(G)$ . We know by Lemma 2 that  $D_A^{p_1} f \in L^{(p'_1,\theta)}(G)$  and  $D_A^{p_2} g \in L^{(p'_2,\theta)}(G)$ . If we put  $A^*s = u$  and  $A^*t = v$ , then  $d\mu(u) = |A^*| d\mu(s)$  and  $d\mu(v) = |A^*| d\mu(t)$ . From the assumption  $\frac{1}{q} = \frac{1}{p'_1} + \frac{1}{p'_2} - \frac{1}{p'_3}$ , we have

$$\begin{aligned}
B_{\tilde{D}_{A^*}^q m}(f, g)(x) &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \tilde{D}_{A^*}^q m(s, t) \langle s+t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) |A^*|^{\frac{1}{q}} m(A^*s, A^*t) \langle s+t, x \rangle \\
&= |A^*|^{-1} \sum_{u \in \hat{G}} |A^*|^{-1} \sum_{v \in \hat{G}} \hat{f}(A^{*-1}u) \hat{g}(A^{*-1}v) |A^*|^{\frac{1}{q}} m(u, v) \langle A^{*-1}u + A^{*-1}v, x \rangle \\
&= |A^*|^{-2} \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} \hat{f}(A^{*-1}u) \hat{g}(A^{*-1}v) |A^*|^{\frac{1}{q}} m(u, v) \langle (A^{-1})^*(u+v), x \rangle \\
&= |A^*|^{-2} \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} \hat{f}(A^{*-1}u) \hat{g}(A^{*-1}v) |A^*|^{\frac{1}{q}} m(u, v) \langle u+v, A^{-1}x \rangle \\
&= |A^*|^{-2} \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} \hat{f}(A^{*-1}u) \hat{g}(A^{*-1}v) |A^*|^{\frac{1}{p'_1} + \frac{1}{p'_2} - \frac{1}{p'_3}} m(u, v) \langle u+v, A^{-1}x \rangle \\
&= |A^*|^{-2} \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} \hat{f}(A^{*-1}u) \hat{g}(A^{*-1}v) |A^*|^{1 - \frac{1}{p'_1} + 1 - \frac{1}{p'_2} - \frac{1}{p'_3}} m(u, v) \langle u+v, A^{-1}x \rangle \\
&= |A^*|^{-\frac{1}{p'_3}} |A^*|^{-2} |A^*|^2 \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} |A^*|^{-\frac{1}{p'_1}} \hat{f}(A^{*-1}u) |A^*|^{-\frac{1}{p'_2}} \hat{g}(A^{*-1}v) m(u, v) \langle u+v, A^{-1}x \rangle \\
(2.16) \quad &= |A^*|^{-\frac{1}{p'_3}} \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} D_{A^{*-1}}^{p'_1} \hat{f}(u) D_{A^{*-1}}^{p'_2} \hat{g}(v) m(u, v) \langle u+v, A^{-1}x \rangle.
\end{aligned}$$

On the other hand, if we say that  $Ay = s$ , then we have

$$\begin{aligned}
(D_A^{p'_1} f)^\wedge(u) &= \int_G D_A^{p'_1} f(y) \langle u, -y \rangle d\lambda(y) \\
&= \int_G |A|^{\frac{1}{p'_1}} \hat{f}(Ay) \langle u, -y \rangle d\lambda(y) = \int_G |A|^{\frac{1}{p'_1}} f(s) \langle u, -A^{-1}s \rangle d\lambda(A^{-1}s)
\end{aligned}$$

$$\begin{aligned}
&= \int_G |A|^{\frac{1}{p'_1}} f(s) \langle u, -A^{-1}s \rangle |A^{-1}| d\lambda(y) = \int_G |A|^{\frac{1}{p'_1}-1} f(s) \langle u, -A^{-1}s \rangle d\lambda(s) \\
&= \int_G |A|^{-\frac{1}{p_1}} f(s) \langle (A^{-1})^* u, -s \rangle d\lambda(y) = |A^*|^{-\frac{1}{p_1}} \int_G f(s) \langle A^{*-1}u, -s \rangle d\lambda(s) \\
&\quad = |A^*|^{-\frac{1}{p_1}} \hat{f}(A^{*-1}u) = D_{A^{*-1}}^{p_1} \hat{f}(u).
\end{aligned}$$

Similarly we achieve  $(D_A^{p'_2} g)^\wedge = D_{A^{*-1}}^{p_2} \hat{g}$ . Then from (2.16), we obtain

$$\begin{aligned}
B_{\tilde{D}_{A^*}^q m}(f, g)(x) &= |A^*|^{-\frac{1}{p'_3}} \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} D_{A^{*-1}}^{p_1} \hat{f}(u) D_{A^{*-1}}^{p_2} \hat{g}(u) m(u, v) \langle u + v, A^{-1}x \rangle \\
&= |A^*|^{-\frac{1}{p'_3}} \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} (D_A^{p'_1} f)^\wedge(u) (D_A^{p'_2} g)^\wedge(v) m(u, v) \langle u + v, A^{-1}x \rangle \\
&= |A|^{-\frac{1}{p'_3}} B_m(D_A^{p'_1} f, D_A^{p'_2} g)(A^{-1}x) = D_{A^{-1}}^{p'_3} B_m(D_A^{p'_1} f, D_A^{p'_2} g)(x).
\end{aligned}$$

Since  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ , from Lemma 2, we have

$$\begin{aligned}
\|B_{\tilde{D}_{A^*}^q m}(f, g)\|_{(p'_3, \theta)} &= \|D_{A^{-1}}^{p'_3} B_m(D_A^{p'_1} f, D_A^{p'_2} g)\|_{(p'_3, \theta)} \leq \|B_m(D_A^{p'_1} f, D_A^{p'_2} g)\|_{(p'_3, \theta)} \\
&\leq \|B_m\| \|D_A^{p'_1} f\|_{(p'_1, \theta)} \|D_A^{p'_2} g\|_{(p'_2, \theta)} \\
&\leq \|B_m\| \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \\
(2.17) \quad &= \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}.
\end{aligned}$$

Thus we obtain  $\tilde{D}_{A^*}^q m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ . Also by (2.17),

$$\|\tilde{D}_{A^*}^q m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \leq \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]}.$$

□

**Theorem 4.** Let  $A$  be an automorphism of  $G$  and  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$  such that  $m(A^*s, A^*t) = m(s, t)$ , where  $\frac{1}{q} = \frac{1}{p'_1} + \frac{1}{p'_2} - \frac{1}{p'_3}$ . Then

$$\frac{1}{p'_1} + \frac{1}{p'_2} = \frac{1}{p'_3}.$$

*Proof.* Assume that  $f \in L^{(p'_1, \theta)}(G)$  and  $g \in L^{(p'_2, \theta)}(G)$ . Since  $\frac{1}{q} = \frac{1}{p'_1} + \frac{1}{p'_2} - \frac{1}{p'_3}$ , then by Theorem 3

$$(2.18) \quad B_{\tilde{D}_{A^*}^q m}(f, g)(x) = D_{A^{-1}}^{p'_3} B_m(D_A^{p'_1} f, D_A^{p'_2} g)(x), \quad x \in G.$$

On the other hand, we write

$$\begin{aligned}
&D_{A^{-1}}^{p'_3} B_m(D_A^{p'_1} f, D_A^{p'_2} g)(x) = \\
&= |A|^{-\frac{1}{p'_3}} \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} (D_A^{p'_1} f)^\wedge(u) (D_A^{p'_2} g)^\wedge(v) m(u, v) \langle u + v, A^{-1}x \rangle \\
&= |A|^{-\frac{1}{p'_3}} \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} D_{A^{*-1}}^{p_1} \hat{f}(u) D_{A^{*-1}}^{p_2} \hat{g}(v) m(u, v) \langle u + v, A^{-1}x \rangle
\end{aligned}$$

$$= |A|^{-\frac{1}{p_3}} \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} |A^*|^{-\frac{1}{p_1}} \hat{f}(A^{*-1}u) |A^*|^{-\frac{1}{p_2}} \hat{g}(A^{*-1}v) m(u, v) \langle A^{*-1}(u+v), x \rangle.$$

We make the substitution  $A^{*-1}u = s$ ,  $A^{*-1}v = t$ . Using  $\mu(A^*\hat{G}) = |A^*|\mu(\hat{G})$ ,  $m(A^*s, A^*t) = m(s, t)$ ,  $|A| = |A^*|$  and  $(A^*)^{-1} = (A^{-1})^*$ , we have

$$\begin{aligned} & D_{A^{-1}}^{p'_3} B_m \left( D_A^{p'_1} f, D_A^{p'_2} g \right) (x) = \\ &= |A|^{-\frac{1}{p_3} - \frac{1}{p_1} - \frac{1}{p_2}} |A^*| \sum_{s \in \hat{G}} |A^*| \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) m(A^*s, A^*t) \langle s+t, x \rangle \\ &= |A|^{-\frac{1}{p_3} - \frac{1}{p_1} + 1 - \frac{1}{p_2} + 1} \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) m(s, t) \langle s+t, x \rangle \\ &\quad (2.19) \quad = |A|^{\frac{1}{p'_1} + \frac{1}{p'_2} - \frac{1}{p'_3}} B_m(f, g)(x). \end{aligned}$$

Hence by (2.18) and (2.19), we have

$$(2.20) \quad B_m(f, g)(x) = |A|^{-\left(\frac{1}{p'_1} + \frac{1}{p'_2} - \frac{1}{p'_3}\right)} B_{\tilde{D}_{A^*}^q m}(f, g)(x).$$

Since  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ , by Theorem 3, we have  $\tilde{D}_{A^*}^q m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$  and

$$\left\| \tilde{D}_{A^*}^q m \right\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \leq \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]}.$$

Then, by (2.20) and Theorem 3,

$$\begin{aligned} & \|B_m(f, g)\|_{(p'_3, \theta)} = |A|^{-\left(\frac{1}{p'_1} + \frac{1}{p'_2} - \frac{1}{p'_3}\right)} \left\| B_{\tilde{D}_{A^*}^q m}(f, g) \right\|_{(p'_3, \theta)} \\ & \leq |A|^{-\left(\frac{1}{p'_1} + \frac{1}{p'_2} - \frac{1}{p'_3}\right)} \left\| \tilde{D}_{A^*}^q m \right\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \\ & \leq |A|^{-\left(\frac{1}{p'_1} + \frac{1}{p'_2} - \frac{1}{p'_3}\right)} \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \\ & = |A|^{-\left(\frac{1}{p'_1} + \frac{1}{p'_2} - \frac{1}{p'_3}\right)} \|B_m\| \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}. \end{aligned}$$

Since this inequality holds for any  $|A|$ , one needs  $\frac{1}{p'_1} + \frac{1}{p'_2} = \frac{1}{p'_3}$ .  $\square$

**Theorem 5.** Let  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ .

a) If  $\Phi \in \ell^1(\hat{G} \times \hat{G})$ , then  $\Phi * m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$  and

$$\|\Phi * m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \leq \|\Phi\|_{\ell^1} \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]}.$$

b) If  $\Phi \in L^1(G \times G)$ , then  $\Phi^\wedge m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$  and

$$\|\Phi^\wedge m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \leq \|\Phi\|_1 \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]}.$$

c) Let  $\frac{1}{p'} = \frac{1}{p'_1} + \frac{1}{p'_2}$ ,  $p_3 \cdot p'_3 < p' + 1$  and  $m(s, t) = a$ . Then  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ .

*Proof.* **a)** Take any  $f, g \in C^\infty(G)$ . Then

$$\begin{aligned}
B_{\Phi*m}(f, g)(x) &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) (\Phi * m)(s, t) \langle s + t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left( \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} m(s - u, t - v) \Phi(u, v) \right) \langle s + t, x \rangle \\
&= \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} \left( \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) m(s - u, t - v) \langle s + t, x \rangle \right) \Phi(u, v) \\
(2.21) \quad &= \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} B_{T_{(u,v)}m}(f, g)(x) \Phi(u, v).
\end{aligned}$$

Since  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ , by Theorem 2 we have  $T_{(u,v)}m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ . Using the equality (2.21), we write

$$\begin{aligned}
\|B_{\Phi*m}(f, g)\|_{(p'_3, \theta)} &\leq \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} \|B_{T_{(u,v)}m}(f, g)(x) \Phi(u, v)\|_{(p'_3, \theta)} \\
&\leq \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} |\Phi(u, v)| \|T_{(u,v)}m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \\
(2.22) \quad &= \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \|\Phi\|_{\ell^1} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} < \infty.
\end{aligned}$$

Hence  $\Phi * m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ , and by (2.22)

$$\|\Phi * m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \leq \|\Phi\|_{\ell^1} \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]}.$$

**b)** Let  $\Phi \in L^1(G \times G)$ . Take any  $f, g \in C^\infty(G)$ . Then we have

$$\begin{aligned}
B_{\hat{\Phi}m}(f, g)(x) &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \Phi^m(s, t) \langle s + t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left( \int_G \int_G \Phi(u, v) \langle s, -u \rangle \langle t, -v \rangle d\lambda(u) d\lambda(v) \right) m(s, t) \langle s + t, x \rangle \\
&= \int_G \int_G \Phi(u, v) \left( \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \langle s, -u \rangle \langle t, -v \rangle m(s, t) \langle s + t, x \rangle \right) d\lambda(u) d\lambda(v) \\
&= \int_G \int_G \Phi(u, v) \left( \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) M_{-v}^2 M_{-u}^1 m(s, t) \langle s + t, x \rangle \right) d\lambda(u) d\lambda(v) \\
(2.23) \quad &= \int_G \int_G \Phi(u, v) B_{M_{-v}^2 M_{-u}^1}(f, g)(x) d\lambda(u) d\lambda(v).
\end{aligned}$$

Since  $m \in BM_\theta [(p'_1; (p'_2; (p'_3))]$ , by Theorem 2 we have  $M_{-v}^2 M_{-u}^1 m \in BM_\theta [(p'_1; (p'_2; (p'_3))]$ , where  $M_{-v}^2$  and  $M_{-u}^1$  are modulation operators. Then by the equality (2.23), we obtain

$$\begin{aligned}
& \|B_{\Phi^\wedge m}(f, g)\|_{(p'_3, \theta)} \leq \int_G \int_G \left\| \Phi(u, v) B_{M_{-v}^2 M_{-u}^1}(f, g) \right\|_{(p'_3, \theta)} d\lambda(u) d\lambda(v) \\
& \leq \int_G \int_G |\Phi(u, v)| \|M_{-v}^2 M_{-u}^1 m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} d\lambda(u) d\lambda(v) \\
& = \int_G \int_G |\Phi(u, v)| \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} d\lambda(u) d\lambda(v) \\
(2.24) \quad & = \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \|\Phi\|_1 \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}.
\end{aligned}$$

Thus  $\Phi^\wedge m \in BM_\theta [(p'_1; (p'_2; (p'_3))]$ . By (2.24), we achieve that

$$\|\Phi^\wedge m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \leq \|\Phi\|_1 \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]}.$$

c) Take any  $f \in L^{(p'_1, \theta)}(G)$  and  $g \in L^{(p'_2, \theta)}(G)$ . Then by Lemma 1, we have

$$\begin{aligned}
& \|B_m(f, g)\|_{(p'_3, \theta)} = \left\| \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) m(s, t) \langle s + t, x \rangle \right\|_{(p'_3, \theta)} \\
& = \left\| a \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \langle s + t, x \rangle \right\|_{(p'_3, \theta)} = |a| \left\| \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \langle s, x \rangle \langle t, x \rangle \right\|_{(p'_3, \theta)} \\
(2.25) \quad & = |a| \left\| \left( \sum_{s \in \hat{G}} \hat{f}(s) \langle s, x \rangle \right) \left( \sum_{t \in \hat{G}} \hat{g}(t) \langle t, x \rangle \right) \right\|_{(p'_3, \theta)} = |a| \|fg\|_{(p'_3, \theta)}.
\end{aligned}$$

Then by Lemma 1, and (2.25) we obtain

$$\|B_m(f, g)\|_{(p'_3, \theta)} \leq C |a| \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}.$$

Thus  $m \in BM_\theta [(p'_1; (p'_2; (p'_3))]$ . □

**Corollary 1.** Let  $\frac{1}{p'} = \frac{1}{p'_1} + \frac{1}{p'_2}$ ,  $p_3 \cdot p'_3 < p' + 1$ . If  $\Phi \in L^1(G \times G)$ , then  $\Phi^\wedge \in BM_\theta [(p'_1; (p'_2; (p'_3))]$ .

*Proof.* Let  $\frac{1}{p'} = \frac{1}{p'_1} + \frac{1}{p'_2}$ ,  $p_3 \cdot p'_3 < p' + 1$ . If we take  $m(s, t) = 1$  in Theorem 5 (c), we have  $m \in BM_\theta [(p'_1; (p'_2; (p'_3))]$ . Since  $\Phi \in L^1(G \times G)$ , by Theorem 5 (b), we obtain  $\Phi^\wedge = \Phi^\wedge m \in BM_\theta [(p'_1; (p'_2; (p'_3))]$ . □

**Proposition 1.** let  $A$  be an automorphism of  $G$  and let  $m \in BM_\theta [(p'_1; (p'_2; (p'_3))]$ . If  $\Psi \in \ell^1(\hat{G}, |A^*|^{-\frac{1}{q}} d\mu)$  such that  $\frac{1}{q} = \frac{1}{p'_1} + \frac{1}{p'_2} - \frac{1}{p'_3}$ , then

$$m_\Psi(s, t) = \sum_{u \in A^* \hat{G}} m(A^* s, A^* t) \Psi(u) \in BM_\theta [(p'_1; (p'_2; (p'_3))].$$

Moreover,

$$\|m_\Psi\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \leq \|\Psi\|_{\ell^1(\hat{G}, |A^*|^{-\frac{1}{q}} d\mu)} \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]}.$$

*Proof.* Let  $f, g \in C^\infty(G)$ . Then

$$\begin{aligned}
B_{m_\Psi}(f, g)(x) &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) m_\Psi(s, t) \langle s + t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left\{ \sum_{u \in A^* \hat{G}} m(A^* s, A^* t) \Psi(u) \right\} \langle s + t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left\{ \sum_{u \in A^* \hat{G}} |A^*|^{-\frac{1}{q}} |A^*|^{\frac{1}{q}} m(A^* s, A^* t) \Psi(u) \right\} \langle s + t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left\{ \sum_{u \in A^* \hat{G}} |A^*|^{-\frac{1}{q}} \tilde{D}_{A^*}^q m(s, t) \Psi(u) \right\} \langle s + t, x \rangle \\
&= \sum_{u \in A^* \hat{G}} \left( \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \tilde{D}_{A^*}^q m(s, t) \langle s + t, x \rangle \right) |A^*|^{-\frac{1}{q}} \Psi(u) \\
&= \sum_{u \in A^* \hat{G}} B_{\tilde{D}_{A^*}^q m}(f, g) \Psi(u) |A^*|^{-\frac{1}{q}} \Psi(u).
\end{aligned}$$

Since  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ ,  $\tilde{D}_{A^*}^q m \in BM(v_{s_1}, v_{s_2}, v_{s_3})$  by Theorem 3, thus we observe that

$$\begin{aligned}
\|B_{m_\Psi}(f, g)\|_{(p'_3, \theta)} &\leq \sum_{u \in A^* \hat{G}} \|B_{\tilde{D}_{A^*}^q m}(f, g)\|_{(p'_3, \theta)} |A^*|^{-\frac{1}{q}} \Psi(u) \\
&\leq \sum_{u \in A^* \hat{G}} \|B_{\tilde{D}_{A^*}^q m}\| \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} |A^*|^{-\frac{1}{q}} \Psi(u) \\
&= \sum_{u \in A^* \hat{G}} \|\tilde{D}_{A^*}^q m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} |A^*|^{-\frac{1}{q}} \Psi(u) \\
&\leq \sum_{u \in A^* \hat{G}} \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} |A^*|^{-\frac{1}{q}} \Psi(u) \\
&= \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \sum_{u \in A^* \hat{G}} |A^*|^{-\frac{1}{q}} \Psi(u) \\
&= \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \sum_{u \in \hat{G}} |A^*|^{-\frac{1}{q}} \Psi(u) \\
&= \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \|\Psi\|_{\ell^1(\hat{G}, |A^*|^{-\frac{1}{q}} d\mu)}.
\end{aligned}$$

Hence,  $m_\Psi \in BM_\theta[(p'_1; (p'_2; (p'_3))]$  and

$$\|m_\Psi\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \leq \|\Psi\|_{\ell^1(\hat{G}, |A^*|^{-\frac{1}{q}} d\mu)} \|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]}.$$

□

**Proposition 2.** Let  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ . If  $U_1, U_2$  are bounded measurable sets in  $\hat{G}$ , then

$$h(s, t) = \frac{1}{\mu(U_1 \times U_2)} \sum_{u \in U_1} \sum_{v \in U_2} m(s + u, t + v) \in BM_\theta[(p'_1; (p'_2; (p'_3))].$$

*Proof.* Let  $f, g \in C^\infty(G)$ . We observe that

$$\begin{aligned} B_h(f, g)(x) &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) h(s, t) \langle s + t, x \rangle = \\ &= \frac{1}{\mu(U_1 \times U_2)} \sum_{u \in U_1} \sum_{v \in U_2} \left\{ \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) m(s + u, t + v) \langle s + t, x \rangle \right\} \\ &= \frac{1}{\mu(U_1 \times U_2)} \sum_{u \in U_1} \sum_{v \in U_2} B_{T_{(-u, -v)m}}(f, g)(x). \end{aligned}$$

By Theorem 2, we have

$$\begin{aligned} \|B_h(f, g)\|_{(p'_3, \theta)} &\leq \frac{1}{\mu(U_1 \times U_2)} \sum_{u \in U_1} \sum_{v \in U_2} \|B_{T_{(-u, -v)m}}(f, g)\|_{(p'_3, \theta)} \\ &\leq \frac{1}{\mu(U_1 \times U_2)} \sum_{u \in U_1} \sum_{v \in U_2} \|T_{(-u, -v)m}\|_{[(p'_1; (p'_2; (p'_3))]} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \\ &= \frac{1}{\mu(U_1 \times U_2)} \|m\|_{[(p'_1; (p'_2; (p'_3))]_0} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \\ &= \|m\|_{[(p'_1; (p'_2; (p'_3))]_\theta} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}. \end{aligned}$$

Finally, we achieve  $h \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ .  $\square$

**Proposition 3.** Let  $\frac{1}{p'} = \frac{1}{p'_1} + \frac{1}{p'_2}$ ,  $p_3, p'_3 < p' + 1$  and let  $A, B$  be automorphisms of  $G$ . If  $\lambda \in M(G)$  and  $m(s, t) = \hat{\lambda}(A^*s + B^*t)$ , then  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ . Moreover,

$$\|m\|_{[(p'_1; (p'_2; (p'_3))]_\theta} \leq C \|\lambda\|$$

for some  $C > 0$ .

*Proof.* Let  $f, g \in C^\infty(G)$ . Then

$$\begin{aligned} B_m(f, g)(x) &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) m(s, t) \langle s + t, x \rangle \\ &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left\{ \int_G \langle A^*s + B^*t, -y \rangle d\lambda(y) \right\} \langle s + t, x \rangle \\ &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left\{ \int_G \langle A^*s, -y \rangle \langle B^*t, -y \rangle d\lambda(y) \right\} \langle s, x \rangle \langle t, x \rangle \\ &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left\{ \int_G \langle s, A(-y) \rangle \langle t, A(-y) \rangle d\lambda(y) \right\} \langle s, x \rangle \langle t, x \rangle \\ &= \int_G \left\{ \sum_{s \in \hat{G}} \hat{f}(s) \langle s, A(-y) \rangle \langle s, x \rangle \right\} \left\{ \sum_{t \in \hat{G}} \hat{g}(t) \langle t, A(-y) \rangle \langle t, x \rangle \right\} d\lambda(y) \\ &= \int_G \left\{ \sum_{s \in \hat{G}} \hat{f}(s) \langle s, A(-y) + x \rangle \right\} \left\{ \sum_{t \in \hat{G}} \hat{g}(t) \langle t, A(-y) + x \rangle \right\} d\lambda(y) \end{aligned}$$

$$\begin{aligned}
&= \int_G f(x + A(-y)) g(x + A(-y)) d\lambda(y) \\
(2.26) \quad &= \int_G f(x - Ay) g(x - Ay) d\lambda(y).
\end{aligned}$$

By (2.26) and Lemma 1, we have

$$\begin{aligned}
\|B_m(f, g)\|_{(p'_3, \theta)} &\leq \int_G \|f(x - Ay) g(x - Ay)\|_{(p'_3, \theta)} d|\lambda|(y) \\
&\leq \int_G C \|f(\cdot - Ay)\|_{(p'_1, \theta)} \|g(\cdot - Ay)\|_{(p'_2, \theta)} d|\lambda|(y) \\
&= C \int_G \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} d|\lambda|(y) \\
(2.27) \quad &= C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \int_G d|\lambda|(y) = C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \|\lambda\|.
\end{aligned}$$

Since  $\lambda \in M(G)$ , then by (2.27)  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ . Thus we have

$$\|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} = \sup \left\{ \frac{\|B_m(f, g)\|_{(p'_3, \theta)}}{\|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}} : \|f\|_{(p'_1, \theta)} \leq 1, \|g\|_{(p'_2, \theta)} \leq 1 \right\} \leq C \|\lambda\|.$$

□

It is known that the unit operator  $I$  is an automorphism of  $G$ . It is easy to see the conjugate  $I^*$  of  $I$  is an unit operator from  $\hat{G}$  into itself. It is continuous, one-to-one and onto. Thus,  $I^*$  becomes an automorphism of  $\hat{G}$ . Similarly one can easily show that  $-I$  and its conjugate  $-I^*$  are authomorphisms of  $G$  and  $\hat{G}$  respectively. Since  $\hat{\lambda}(A^*s + B^*t) = \hat{\lambda}(s \mp t)$ , in Proposition 3, one can get  $m(s, t) = \hat{\lambda}(s \mp t)$ . As an application of this result we can give the following example.

**Example 2.** If  $\lambda \in M(G)$  and  $m(s, t) = \hat{\lambda}(s \mp t)$ , then  $\hat{\lambda} \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3))]$  and

$$\|\hat{\lambda}\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \leq C \|\lambda\|_1, \quad C > 0$$

for  $\frac{1}{p'_1} + \frac{1}{p'_2} = \frac{1}{p'}$  and  $p'_3 p_3 < p' + 1$ .

**Theorem 6.** Let  $\frac{1}{p'} = \frac{1}{p'_1} + \frac{1}{p'_2}$ ,  $p_3 \cdot p'_3 < p' + 1$ . If  $m(s, t) = \hat{\Psi}_1(s) \hat{\Phi}(s, t) \hat{\Psi}_2(t)$  such that  $\Phi \in L^1(G \times G)$  and  $\Psi_1, \Psi_2 \in L^1(G)$ , then  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ .

*Proof.* Let  $f, g, h \in C^\infty(G)$ . For the proof of this Theorem we will use Theorem 1. Then by Fubini Theorem

$$\left| \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \hat{h}(s + t) m(s, t) \right| =$$

$$\begin{aligned}
&= \left| \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left\{ \int_G h(y) \langle s+t, -y \rangle d\lambda(y) \right\} m(s, t) \right| \\
&= \left| \int_G h(y) \left\{ \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \hat{\Psi}_1(s) \hat{\Phi}(s, t) \hat{\Psi}_2(t) \langle s+t, -y \rangle \right\} d\lambda(y) \right| \\
&= \left| \int_G h(y) \left\{ \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} (f * \Psi_1)^\wedge(s) (g * \Psi_2)^\wedge(t) \hat{\Phi}(s, t) \langle s+t, -y \rangle \right\} d\lambda(y) \right| \\
(2.28) \quad &\leq \int_G |h(y) B_{\hat{\Phi}}(f * \Psi_1, g * \Psi_2)(-y)| d\lambda(y).
\end{aligned}$$

On the other hand  $L^{(p'_1, \theta)}(G)$  and  $L^{(p'_2, \theta)}(G)$  are Banach convolution module over  $L^1(G)$ , [6]. So we have  $f * \Psi_1 \in L^{(p'_1, \theta)}(G)$  and  $g * \Psi_2 \in L^{(p'_2, \theta)}(G)$ . Also by Corollary 1,  $\hat{\Phi} \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ . Therefore we achieve  $B_{\hat{\Phi}}(f * \Psi_1, g * \Psi_2) \in L^{(p'_3, \theta)}(G)$ . By Theorem 2.5 in [2] and the inequality (2.28), we obtain that

$$\begin{aligned}
&\left| \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \hat{h}(s+t) m(s, t) \right| \leq \|h\|_{p_3, \theta} \|B_{\hat{\Phi}}(f * \Psi_1, g * \Psi_2)\|_{(p'_3, \theta)} \\
&\leq \|h\|_{p_3, \theta} \|B_{\hat{\Phi}}\| \|f * \Psi_1\|_{(p'_1, \theta)} \|g * \Psi_2\|_{(p'_2, \theta)} \\
&\leq \|h\|_{p_3, \theta} \|B_{\hat{\Phi}}\| \|\Psi_1\|_1 \|f\|_{(p'_1, \theta)} \|\Psi_2\|_1 \|g\|_{(p'_2, \theta)}.
\end{aligned}$$

Let  $C = \|B_{\hat{\Phi}}\| \|\Psi_1\|_1 \|\Psi_2\|_1$ . Then

$$\left| \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \hat{h}(s+t) m(s, t) \right| \leq C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \|h\|_{p_3, \theta}.$$

Hence by Theorem 1, we achieve  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ .  $\square$

**Example 3.** If  $K \in L^1(G)$  then  $m(s, t) = \hat{K}(s-t)$  defines a bilinear multiplier in  $BM_\theta[(p'_1; (p'_2; (p'_3))]$  and

$$\|m\|_{[(p'_1; (p'_2; (p'_3)))]_\theta} \leq C \|K\|_1, \quad C > 0$$

for  $\frac{1}{p'} = \frac{1}{p'_1} + \frac{1}{p'_2}$ ,  $p_3 \cdot p'_3 < p' + 1$ . Indeed for  $f, g \in C^\infty(G) \subset L^1(G)$ , since

$$f(x-y) = \sum_{s \in \hat{G}} \hat{f}(s) \langle s, x-y \rangle$$

and

$$g(x+y) = \sum_{t \in \hat{G}} \hat{g}(t) \langle t, x+y \rangle,$$

one has

$$B_m(f, g)(x) = \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) m(s, t) \langle s+t, x \rangle$$

$$\begin{aligned}
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left( \int_G K(y) \langle s-t, -y \rangle d\lambda(y) \right) \langle s+t, x \rangle \\
&= \int_G \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) K(y) \langle s-t, -y \rangle \langle s+t, x \rangle d\lambda(y) \\
&= \int_G \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) K(y) \langle s, -y \rangle \langle -t, -y \rangle \langle s, x \rangle \langle t, x \rangle d\lambda(y) \\
&= \int_G \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) K(y) \langle s, -y \rangle \langle t, y \rangle \langle s, x \rangle \langle t, x \rangle d\lambda(y) \\
&= \int_G \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) K(y) \langle s, x-y \rangle \langle t, x+y \rangle d\lambda(y) \\
&= \int_G \left( \sum_{s \in \hat{G}} \hat{f}(s) \langle s, x-y \rangle \right) \left( \sum_{t \in \hat{G}} \hat{g}(t) \langle t, x+y \rangle \right) K(y) d\lambda(y) \\
(2.29) \quad &= \int_G f(x-y) g(x+y) K(y) d\lambda(y).
\end{aligned}$$

Then from (2.29) and by Lemma 1,

$$\begin{aligned}
&\|B_m(f, g)\|_{(p'_3, \theta)} \leq \int_G \|f(x-y) g(x+y)\|_{(p'_3, \theta)} |K(y)| d\lambda(y) \\
&\leq C \int_G \|f(x-y)\|_{(p'_1, \theta)} \|g(x+y)\|_{(p'_2, \theta)} |K(y)| d\lambda(y),
\end{aligned}$$

for some  $C > 0$ . Since  $\|T_{-x}f(-y)\|_{(p'_1, \theta)} = \|f(y)\|_{(p'_1, \theta)}$  and  $\|T_{-x}g(y)\|_{(p'_2, \theta)} = \|g(y)\|_{(p'_2, \theta)}$  by Theorem 2, then

$$\begin{aligned}
&\|B_m(f, g)\|_{(p'_3, \theta)} \leq C \int_G \|T_{-x}f(-y)\|_{(p'_1, \theta)} \|T_{-x}g(y)\|_{(p'_2, \theta)} |K(y)| d\lambda(y) \\
(2.30) \quad &= C \int_G \|\tilde{f}\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} |K(y)| d\lambda(y) = C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \|K\|_1 \\
&= C_1 \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)},
\end{aligned}$$

where  $C_1 = C \|K\|_1$  and  $\tilde{f}(y) = f(-y)$ . Thus  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$ . Finally by using (2.30), we obtain

$$\|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} = \sup \left\{ \frac{\|B_m(f, g)\|_{(p'_3, \theta)}}{\|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}} : \|f\|_{(p'_1, \theta)} \leq 1, \|g\|_{(p'_2, \theta)} \leq 1 \right\} \leq C \|K\|_1.$$

**Definition 3.** Let  $1 < p_i < \infty$ ,  $p'_i = \frac{p_i}{p_i - 1}$ , ( $i = 1, 2, 3$ ) and  $\theta > 0$ . We denote by  $\tilde{M}_\theta[(p'_1; (p'_2; (p'_3))]$  the space of measurable functions  $M : \hat{G} \rightarrow \mathbb{C}$  such that  $m(s, t) = M(s - t) \in BM_\theta[(p'_1; (p'_2; (p'_3))]$  that is to say

$$B_M(f, g)(x) = \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) M(s - t) \langle s + t, x \rangle$$

extends to bounded bilinear map from  $L^{(p'_1, \theta)}(G) \times L^{(p'_2, \theta)}(G)$  to  $L^{(p'_3, \theta)}(G)$ . We denote  $\|M\|_{[(p'_1; (p'_2; (p'_3))_\theta]} = \|B_M\|$ .

**Example 4.** Let  $G$  be a locally compact abelian metric group and let  $0_G$  be the unit of  $G$ . Take the bilinear Hardy-Littlewood maximal function on  $G$ ;

$$M(f, g)(x) = \sup_{r > 0} \frac{1}{\lambda(B(0_G, r))} \int_{B(0_G, r)} |f(x - y) g(x + y)| d\lambda(y)$$

for all  $f, g \in L^1_{loc}(G)$ , where  $B(0_G, r)$  is open ball in  $G$ . The Hardy-Littlewood maximal function is bounded from  $L^{(p'_1, \theta)}(G) \times L^{(p'_2, \theta)}(G)$  to  $L^{(p'_3, \theta)}(G)$  whenever  $\frac{1}{p'} = \frac{1}{p'_1} + \frac{1}{p'_2}$  and  $p_3 \cdot p'_3 < p' + 1$ .

Take the function

$$M(y) = \frac{1}{\lambda(B(0_G, r))} \chi_{B(0_G, r)}(y).$$

Since  $M \in L^1(G)$ , by Proposition 4,  $M$  defines a bilinear multiplier in  $\tilde{M}_\theta[(p'_1; (p'_2; (p'_3))]$  and

$$(2.31) \quad \|B_M(f, g)\|_{(p'_3, \theta)} \leq C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \|K\|_1 = C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}$$

where

$$\begin{aligned} B_M(f, g)(x) &= \int_G f(x - y) g(x + y) M(y) d\lambda(y) \\ &= \frac{1}{\lambda(B(0_G, r))} \int_{B(0_G, r)} f(x - y) g(x + y) d\lambda(y) \end{aligned}$$

for all  $r > 0$ . By (2.31), we write

$$(2.32) \quad M(f, g)(x) = \sup_{r > 0} \frac{1}{\lambda(B(0_G, r))} \int_{B(0_G, r)} |f(x - y) g(x + y)| d\lambda(y) = \sup_{r > 0} B_M(|f|, |g|).$$

So by (2.31) we have

$$(2.33) \quad \|B_M(|f|, |g|)\|_{(p'_3, \theta)} \leq C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}$$

for all  $r > 0$ . Then by (2.32) and (2.33), we obtain

$$\|M(f, g)\|_{(p'_3, \theta)} \leq C \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}.$$

Therefore  $M(f, g)$  is bounded from  $L^{(p'_1, \theta)}(G) \times L^{(p'_2, \theta)}(G)$  to  $L^{(p'_3, \theta)}(G)$ .

**Proposition 4.** Let  $M \in \ell^1(\hat{G})$ . Then for all  $f \in L^{(p'_1, \theta)}(G)$  and  $g \in L^{(p'_2, \theta)}(G)$

$$B_M(f, g)(x) = \int_G f(x - y) g(x + y) M^\vee(y) dy.$$

*Proof.* Let  $f, g \in C^\infty(G)$ . Then

$$\begin{aligned}
B_M(f, g)(x) &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) M(s-t) \langle s+t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left( \int_G M^\vee(y) \langle s-t, -y \rangle d\lambda(y) \right) \langle s+t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left( \int_G M^\vee(y) \langle s-t, -y \rangle \langle s+t, x \rangle d\lambda(y) \right) \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left( \int_G \check{M}(y) \langle s, -y \rangle \langle t, y \rangle \langle s, x \rangle \langle t, x \rangle d\lambda(y) \right) \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left( \int_G M^\vee(y) \langle s, x-y \rangle \langle t, x+y \rangle d\lambda(y) \right) \\
&= \int_G \check{M}(y) \left( \sum_{s \in \hat{G}} \hat{f}(s) \langle s, x-y \rangle \right) \left( \sum_{t \in \hat{G}} \hat{g}(t) \langle t, x+y \rangle \right) d\lambda(y) \\
&= \int_G f(x-y) g(x+y) M^\vee(y) d\lambda(y).
\end{aligned}$$

□

**Proposition 5.** Let  $K \in \ell^1(\hat{G})$ . Then the following equalities are satisfied;

- a)  $B_{T_{y_1+y_2}K}(f, g) = M_{y_1-y_2}B_K(M_{-y_1}f, M_{y_2}g)$ ,  $y_1, y_2 \in \hat{G}$ ,
- b)  $B_{M_y K}(f, g) = B_K(T_{-y}f, T_y g)$ ,  $y \in \hat{G}$ .

*Proof.* a) Let  $f, g \in C^\infty(G)$  and let  $y_1, y_2 \in \hat{G}$ . If we make the substitutions  $s - y_1 = u$  and  $t + y_2 = v$ , then we have

$$\begin{aligned}
B_{T_{y_1+y_2}K}(f, g)(x) &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) T_{y_1+y_2}K(s-t) \langle s+t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) K(s-t - y_1 - y_2) \langle s+t, x \rangle \\
&= \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} \hat{f}(u+y_1) \hat{g}(v-y_2) K(u-v) \langle u+v+y_1-y_2, x \rangle \\
&= \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} T_{-y_1} \hat{f}(u) T_{y_2} \hat{g}(v) K(u-v) \langle u+v, x \rangle \langle y_1-y_2, x \rangle \\
&= \langle y_1 - y_2, x \rangle \sum_{u \in \hat{G}} \sum_{v \in \hat{G}} (M_{-y_1}f)^\wedge(u) (M_{y_2}g)^\wedge(v) K(u-v) \langle u+v, x \rangle \\
&= \langle y_1 - y_2, x \rangle B_K(M_{-y_1}f, M_{y_2}g)(x) = M_{y_1-y_2}B_K(M_{-y_1}f, M_{y_2}g)(x)
\end{aligned}$$

- b) Let  $f, g \in C^\infty(G)$  and let  $y \in \hat{G}$ . Then

$$B_{M_y K}(f, g)(x) = \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) M_y K(s-t) \langle s+t, x \rangle$$

$$\begin{aligned}
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \langle s - t, y \rangle K(s - t) \langle s + t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \langle s, y \rangle \langle -t, y \rangle K(s - t) \langle s + t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \langle s, y \rangle \hat{f}(s) \langle t, -y \rangle \hat{g}(t) K(s - t) \langle s + t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} M_y \hat{f}(s) M_{-y} \hat{g}(t) K(s - t) \langle s + t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} (T_{-y} f)^{\wedge}(s) (T_y g)^{\wedge}(t) K(s - t) \langle s + t, x \rangle = B_K(T_{-y} f, T_y g)(x).
\end{aligned}$$

□

**Theorem 7.** Let  $K \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3))]$ .

a) If  $\Phi \in \ell^1(\hat{G})$ , then  $\Phi * K \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3))]$  and

$$\|\Phi * K\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \leq \|\Phi\|_{\ell^1} \|K\|_{[(p'_1; (p'_2; (p'_3))_\theta]}.$$

b) If  $\Phi \in L^1(G)$ , then  $\hat{\Phi} K \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3))]$  and

$$\|\hat{\Phi} K\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \leq \|\Phi\|_1 \|K\|_{[(p'_1; (p'_2; (p'_3))_\theta]}.$$

*Proof.* a) Take any  $f, g \in C^\infty(G)$  and use Proposition 5

$$\begin{aligned}
B_{\Phi * K}(f, g)(x) &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) (\Phi * K)(s - t) \langle s + t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left( \sum_{u \in \hat{G}} K(s - t - u) \Phi(u) \right) \langle s + t, x \rangle \\
&= \sum_{u \in \hat{G}} \left( \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) K(s - t - u) \langle s + t, x \rangle \right) \Phi(u) \\
&= \sum_{u \in \hat{G}} \left( \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) T_u K(s - t) \langle s + t, x \rangle \right) \Phi(u) \\
&= \sum_{u \in \hat{G}} B_{T_u K}(f, g)(x) \Phi(u) = \sum_{u \in \hat{G}} B_{T_{u+0_{\hat{G}}} K}(f, g)(x) \Phi(u)
\end{aligned}$$

By Proposition 5 and the last equality

$$\begin{aligned}
&B_{\Phi * K}(f, g)(x) = \sum_{u \in \hat{G}} B_{T_{u+0_{\hat{G}}} K}(f, g)(x) \Phi(u) \\
&\stackrel{(2.34)}{=} \sum_{u \in \hat{G}} M_{u-0_{\hat{G}}} B_K(M_{-u} f, M_{0_{\hat{G}}} g)(x) \Phi(u) = \sum_{u \in \hat{G}} M_{u-0_{\hat{G}}} B_K(M_{-u} f, M_{0_{\hat{G}}} g)(x) \Phi(u).
\end{aligned}$$

Since  $K \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3))]$ , by (2.34), we have

$$\|B_{\Phi * K}(f, g)\|_{(p'_3, \theta)} \leq \sum_{u \in \hat{G}} \|M_{u-0_{\hat{G}}} B_K(M_{-u} f, M_{0_{\hat{G}}} g)(x) \Phi(u)\|_{(p'_3, \theta)}$$

$$\begin{aligned}
&= \sum_{u \in \hat{G}} \|B_K(M_{-u}f, M_{0,\hat{G}}g)(x)\Phi(u)\|_{(p'_3,\theta)} \\
&\leq \sum_{u \in \hat{G}} |\Phi(u)| \|K\|_{[(p'_1;(p'_2;(p'_3)]_\theta} \|M_{-u}f\|_{(p'_1,\theta)} \|g\|_{(p'_2,\theta)} \\
(2.35) \quad &= \|M\|_{[(p'_1;(p'_2;(p'_3)]_\theta} \|\Phi\|_{\ell^1} \|f\|_{(p'_1,\theta)} \|g\|_{(p'_2,\theta)} < \infty.
\end{aligned}$$

Hence  $\Phi * K \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3]]$ . Finally by (2.35), we obtain

$$\|\Phi * K\|_{[(p'_1;(p'_2;(p'_3)]_\theta} \leq \|\Phi\|_{\ell^1} \|K\|_{[(p'_1;(p'_2;(p'_3)]_\theta}.$$

b) Let  $f, g \in C^\infty(G)$ . Then by Proposition 5

$$\begin{aligned}
B_{\hat{\Phi}K}(f, g)(x) &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \hat{\Phi}K(s-t) \langle s+t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \left( \int_G \Phi(u) \langle s-t, -u \rangle d\lambda(u) \right) K(s-t) \langle s+t, x \rangle \\
&= \int_G \Phi(u) \left( \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) \langle s-t, -u \rangle K(s-t) \langle s+t, x \rangle \right) d\lambda(u) \\
&= \int_G \Phi(u) \left( \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) M_{-u}K(s-t) \langle s+t, x \rangle \right) d\lambda(u) \\
(2.36) \quad &= \int_G \Phi(u) B_{M_{-u}K}(f, g)(x) d\lambda(u) = \int_G \Phi(u) B_K(T_u f, T_{-u}g)(x) d\lambda(u).
\end{aligned}$$

Since  $K \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3]]$ , by (2.36), we obtain

$$\begin{aligned}
\|B_{\hat{\Phi}K}(f, g)\|_{(p'_3,\theta)} &\leq \int_G \|\Phi(u) B_K(T_u f, T_{-u}g)\|_{(p'_3,\theta)} d\lambda(u) \\
&\leq \int_G |\Phi(u)| \|K\|_{[(p'_1;(p'_2;(p'_3)]_\theta} \|T_u f\|_{(p'_1,\theta)} \|T_{-u}g\|_{(p'_2,\theta)} d\lambda(u) \\
&= \int_G |\Phi(u)| \|K\|_{[(p'_1;(p'_2;(p'_3)]_\theta} \|f\|_{(p'_1,\theta)} \|g\|_{(p'_2,\theta)} d\lambda(u) \\
(2.37) \quad &= \|K\|_{[(p'_1;(p'_2;(p'_3)]_\theta} \|\Phi\|_1 \|f\|_{(p'_1,\theta)} \|g\|_{(p'_2,\theta)} < \infty.
\end{aligned}$$

Finally  $\hat{\Phi}K \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3]]$  and by (2.37)

$$\|\hat{\Phi}K\|_{[(p'_1;(p'_2;(p'_3)]_\theta} \leq \|\Phi\|_1 \|K\|_{[(p'_1;(p'_2;(p'_3)]_\theta}.$$

□

**Proposition 6.** Let  $\Phi \in L^1(G)$  and  $M \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3]]$ . Then  $m(s, t) = M(s-t) \hat{\Phi}(s+t) \in BM_\theta[(p'_1; (p'_2; (p'_3]]$  and

$$\|m\|_{[(p'_1;(p'_2;(p'_3)]_\theta} \leq \|\Phi\|_1 \|M\|_{[(p'_1;(p'_2;(p'_3)]_\theta}.$$

*Proof.* Let  $f, g \in C^\infty(G)$ , Then for all  $x \in G$ , we have

$$\begin{aligned}
B_m(f, g)(x) &= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) M(s-t) \hat{\Phi}(s+t) \langle s+t, x \rangle \\
&= \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) M(s-t) \left( \int_G \Phi(u) \langle s+t, -u \rangle d\lambda(u) \right) \langle s+t, x \rangle \\
&= \int_G \Phi(u) \left( \sum_{s \in \hat{G}} \sum_{t \in \hat{G}} \hat{f}(s) \hat{g}(t) M(s-t) \langle s+t, x-u \rangle \right) d\lambda(u) \\
(2.38) \quad &= \int_G \Phi(u) B_M(f, g)(x-u) d\lambda(u) = \Phi * B_M(f, g)(x).
\end{aligned}$$

On the other hand Since  $L^{(p'_3, \theta)}(G)$  is a Banach convolution module over  $L^1(G)$ , [6] and  $M \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3))]$ , by (2.38) we have

$$\begin{aligned}
\|B_m(f, g)\|_{(p'_3, \theta)} &= \|\Phi * B_M(f, g)\|_{(p'_3, \theta)} \leq \|B_M(f, g)\|_{(p'_3, \theta)} \|\Phi\|_1 \\
(2.39) \quad &\leq \|\Phi\|_1 \|M\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} < \infty.
\end{aligned}$$

Hence  $m \in BM_\theta[(p'_1; (p'_2; (p'_3))]$  and by (2.39)

$$\|m\|_{[(p'_1; (p'_2; (p'_3))_\theta]} \leq \|\Phi\|_1 \|M\|_{[(p'_1; (p'_2; (p'_3))_\theta]}.$$

□

**Proposition 7.** Let  $K \in \ell^1(\hat{G})$  be non-zero function and let  $K \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3))]$ . If  $A$  is an automorphism of  $G$  and if  $\frac{1}{q} = \frac{1}{p'_1} + \frac{1}{p'_2} - \frac{1}{p'_3}$ , then there exists  $C > 0$  such that

$$\left| \int_G K^\vee(u) d\lambda(u) \right| \leq C |A|^{\frac{1}{p'_3}} \|K\|_{[(p'_1; (p'_2; (p'_3))_\theta]} , \quad |A| < 1$$

and

$$\left| \int_G K^\vee(u) d\lambda(u) \right| \leq C |A|^{-\frac{1}{q}} \|K\|_{[(p'_1; (p'_2; (p'_3))_\theta]} , \quad |A| \geq 1.$$

*Proof.* Let  $A$  be any automorphism of  $G$ . Define a function  $f : G \rightarrow \mathbb{C}$  by  $f(x) = \langle \gamma, Ax \rangle$  for fixed  $\gamma \in \hat{G}$ . By Proposition 4, we write

$$\begin{aligned}
B_K(f, f)(x) &= \int_G f(x-y) f(x+y) K^\vee(y) dy \\
&= \int_G \langle \gamma, A(x-y) \rangle \langle \gamma, A(x+y) \rangle K^\vee(y) dy \\
&= \int_G \langle \gamma, Ax \rangle \langle \gamma, -Ay \rangle \langle \gamma, Ax \rangle \langle \gamma, Ay \rangle K^\vee(y) dy
\end{aligned}$$

$$(2.40) \quad = \int_G \langle \gamma, Ax \rangle \langle \gamma, Ax \rangle K^\vee(y) dy = \langle \gamma, Ax \rangle \langle \gamma, Ax \rangle \int_G K^\vee(y) dy.$$

Using (2.40) and making the substitution  $Ax = u$ , we have

$$\begin{aligned} \|B_K(f, f)\|_{p'_3} &= \left( \int_G |B_K(f, f)(x)|^{p'_3} d\lambda(x) \right)^{\frac{1}{p'_3}} \\ &= \left( \int_G \left| \langle \gamma, Ax \rangle \langle \gamma, Ax \rangle \int_G K^\vee(y) dy \right|^{p'_3} d\lambda(x) \right)^{\frac{1}{p'_3}} \\ &= \left| \int_G K^\vee(y) dy \right| \left( \int_G |\langle \gamma, Ax \rangle \langle \gamma, Ax \rangle|^{p'_3} d\lambda(x) \right)^{\frac{1}{p'_3}} \\ &= \left| \int_G K^\vee(y) dy \right| \left( \int_G |\langle \gamma, u \rangle \langle \gamma, u \rangle|^{p'_3} |A|^{-1} d\lambda(u) \right)^{\frac{1}{p'_3}} \\ (2.41) \quad &= |A|^{-\frac{1}{p'_3}} \left| \int_G K^\vee(y) dy \right| \left( \int_G d\lambda(u) \right)^{\frac{1}{p'_3}} = |A|^{-\frac{1}{p'_3}} \lambda(G)^{\frac{1}{p'_3}} \left| \int_G K^\vee(y) dy \right|. \end{aligned}$$

On the other hand we can write

$$f(x) = \langle \gamma, Ax \rangle = |A|^{-\frac{1}{p'_1}} |A|^{\frac{1}{p'_1}} \langle \gamma, Ax \rangle = |A|^{-\frac{1}{p'_1}} D_A^{p'_1} \gamma(x).$$

Let  $|A| \geq 1$ . By Lemma 2, we obtain

$$\begin{aligned} \|f\|_{(p'_1, \theta)} &= \left\| |A|^{-\frac{1}{p'_1}} D_A^{p'_1} \gamma \right\|_{(p'_1, \theta)} = |A|^{-\frac{1}{p'_1}} \left\| D_A^{p'_1} \gamma \right\|_{(p'_1, \theta)} \\ &= |A|^{-\frac{1}{p'_1}} \|\gamma\|_{(p'_1, \theta)}. \end{aligned}$$

Since  $L^{p'_1+\varepsilon}(G) \subset L^{(p'_1, \theta)}(G)$ , [2], there exists  $C_1 > 0$  such that  $\|\gamma\|_{(p'_1, \theta)} \leq C_1 \|\gamma\|_{p'_1+\varepsilon}$ . Then

$$\begin{aligned} \|f\|_{(p'_1, \theta)} &= |A|^{-\frac{1}{p'_1}} \|\gamma\|_{(p'_1, \theta)} \leq C_1 |A|^{-\frac{1}{p'_1}} \|\gamma\|_{p'_1+\varepsilon}. \\ &= C_1 |A|^{-\frac{1}{p'_1}} \left( \int_G |\langle \gamma, x \rangle|^{p'_1+\varepsilon} d\lambda(x) \right)^{\frac{1}{p'_1+\varepsilon}} \\ (2.42) \quad &= C_1 |A|^{-\frac{1}{p'_1}} \left( \int_G d\lambda(x) \right)^{\frac{1}{p'_1+\varepsilon}} = C_1 |A|^{-\frac{1}{p'_1}} \lambda(G)^{\frac{1}{p'_1+\varepsilon}} < \infty. \end{aligned}$$

Similarly there exists  $C_2 > 0$  such that

$$(2.43) \quad \|f\|_{(p'_2, \theta)} \leq C_2 |A|^{-\frac{1}{p'_2}} \lambda(G)^{\frac{1}{p'_2+\varepsilon}} < \infty.$$

Using the assumption  $K \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3)]$  and the inequalities (2.42) and (2.43), we obtain

$$(2.44) \quad \begin{aligned} \|B_K(f, f)\|_{(p'_3, \theta)} &\leq \|K\|_{[(p'_1; (p'_2; (p'_3)]_\theta} \|f\|_{(p'_1, \theta)} \|f\|_{(p'_2, \theta)} \\ &\leq C_1 C_2 |A|^{-\frac{1}{p'_1} - \frac{1}{p'_2}} \lambda(G)^{\frac{1}{p'_1 + \varepsilon} + \frac{1}{p'_2 + \varepsilon}} \|K\|_{[(p'_1; (p'_2; (p'_3)]_\theta}. \end{aligned}$$

Since  $L^{(p'_3, \theta)}(G) \subset L^{p'_3}(G)$ , there exists  $C_3 > 0$  such that

$$(2.45) \quad \|B_K(f, f)\|_{p'_3} \leq C_3 \|B_K(f, f)\|_{(p'_3, \theta)}.$$

Then by (2.44) and (2.45), we have

$$(2.46) \quad \|B_K(f, f)\|_{p'_3} \leq C_1 C_2 C_3 |A|^{-\frac{1}{p'_1} - \frac{1}{p'_2}} \lambda(G)^{\frac{1}{p'_1 + \varepsilon} + \frac{1}{p'_2 + \varepsilon}} \|K\|_{[(p'_1; (p'_2; (p'_3)]_\theta}.$$

Also by (2.41) and (2.46), we achieve

$$\begin{aligned} \lambda(G)^{\frac{1}{p'_3}} |A|^{-\frac{1}{p'_3}} \left| \int_G K^\vee(y) dy \right| &= \|B_K(f, f)\|_{p'_3} \leq \\ &\leq C_1 C_2 C_3 |A|^{-\frac{1}{p'_1} - \frac{1}{p'_2}} \lambda(G)^{\frac{1}{p'_1 + \varepsilon} + \frac{1}{p'_2 + \varepsilon}} \|K\|_{[(p'_1; (p'_2; (p'_3)]_\theta}. \end{aligned}$$

This implies

$$\left| \int_G K^\vee(y) dy \right| \leq C |A|^{\frac{1}{p'_3} - \frac{1}{p'_1} - \frac{1}{p'_2}} \|K\|_{[(p'_1; (p'_2; (p'_3)]_\theta} = C |A|^{-\frac{1}{q}} \|K\|_{[(p'_1; (p'_2; (p'_3)]_\theta},$$

where  $C = C_1 C_2 C_3 \lambda(G)^{\frac{1}{p'_1 + \varepsilon} + \frac{1}{p'_2 + \varepsilon} - \frac{1}{p'_3}}$ .

Now let  $|A| < 1$ . By Lemma 2, we have

$$\begin{aligned} \|f\|_{(p'_1, \theta)} &= \left\| |A|^{-\frac{1}{p'_1}} D_A^{p'_1} \gamma \right\|_{(p'_1, \theta)} = |A|^{-\frac{1}{p'_1}} \left\| D_A^{p'_1} \gamma \right\|_{(p'_1, \theta)} \\ &= |A|^{-\frac{1}{p'_1}} |A|^{\frac{1}{p'_1}} \|\gamma\|_{(p'_1, \theta)} = \|\gamma\|_{(p'_1, \theta)}. \end{aligned}$$

Then Since  $L^{p'_1 + \varepsilon}(G) \subset L^{(p'_1, \theta)}(G)$ , we achieve

$$\|f\|_{(p'_1, \theta)} = \|\gamma\|_{(p'_1, \theta)} \leq C_1 \|\gamma\|_{p'_1 + \varepsilon}.$$

$$(2.47) \quad = C_1 \lambda(G)^{\frac{1}{p'_1 + \varepsilon}} < \infty,$$

for some  $C_1 > 0$ . Similarly we have

$$(2.48) \quad \|f\|_{(p'_2, \theta)} \leq C_2 \lambda(G)^{\frac{1}{p'_2 + \varepsilon}} < \infty,$$

for some  $C_2 > 0$ . Again using the assumption  $K \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3)]$  and the inequalities (2.47) and (2.48), we obtain

$$(2.49) \quad \begin{aligned} \|B_K(f, f)\|_{(p'_3, \theta)} &\leq \|K\|_{[(p'_1; (p'_2; (p'_3)]_\theta} \|f\|_{(p'_1, \theta)} \|f\|_{(p'_2, \theta)} \\ &\leq C_1 C_2 \lambda(G)^{\frac{1}{p'_1 + \varepsilon} + \frac{1}{p'_2 + \varepsilon}} \|K\|_{[(p'_1; (p'_2; (p'_3)]_\theta}}. \end{aligned}$$

Thus by (2.45) and (2.49), we have

$$(2.50) \quad \|B_K(f, f)\|_{p'_3} \leq C_1 C_2 C_3 \lambda(G)^{\frac{1}{p'_1 + \varepsilon} + \frac{1}{p'_2 + \varepsilon}} \|K\|_{[(p'_1; (p'_2; (p'_3)]_\theta}}.$$

Using (2.41) and (2.50), we achieve

$$\begin{aligned} \lambda(G)^{\frac{1}{p'_3}} |A|^{-\frac{1}{p'_3}} \left| \int_G K^\vee(y) dy \right| &= \|B_K(f, f)\|_{p'_3} \leq \\ &\leq C_1 C_2 C_3 \lambda(G)^{\frac{1}{p'_1+\varepsilon} + \frac{1}{p'_2+\varepsilon}} \|K\|_{[(p'_1; (p'_2; (p'_3)_\theta)]}. \end{aligned}$$

Then

$$\left| \int_G K^\vee(y) dy \right| \leq C_1 C_2 C_3 |A|^{\frac{1}{p'_3}} \lambda(G)^{\frac{1}{p'_1+\varepsilon} + \frac{1}{p'_2+\varepsilon} - \frac{1}{p'_3}} \|K\|_{[(p'_1; (p'_2; (p'_3)_\theta)]} = C |A|^{\frac{1}{p'_3}} \|K\|_{[(p'_1; (p'_2; (p'_3)_\theta)]},$$

where  $C = C_1 C_2 C_3 \lambda(G)^{\frac{1}{p'_1+\varepsilon} + \frac{1}{p'_2+\varepsilon} - \frac{1}{p'_3}}$ .  $\square$

**Proposition 8.** Let  $K \in \ell^1(\hat{G})$  be non-zero function and let  $K \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3))]$ . If  $A$  is an automorphism of  $\hat{G}$  satisfying  $|A| > 1$ , then

$$\frac{1}{p'_3} \geq \frac{1}{p'_1} + \frac{1}{p'_2}.$$

*Proof.* Assume that  $\frac{1}{p'_3} < \frac{1}{p'_1} + \frac{1}{p'_2}$ . By Proposition 5,

$$\begin{aligned} \|B_{T_{-u}K}(f, g)\|_{(p'_3, \theta)} &= \|B_{T_{-u+0}\hat{G}K}(f, g)\|_{(p'_3, \theta)} \\ &= \|M_{-u-0}\hat{G}B_K(M_u f, M_{0\hat{G}}g)\|_{(p'_3, \theta)} = \|B_K(M_u f, g)\|_{(p'_3, \theta)} \\ &\leq \|K\|_{[(p'_1; (p'_2; (p'_3)_\theta)]} \|M_u f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)} \\ &= \|K\|_{[(p'_1; (p'_2; (p'_3)_\theta)]} \|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}, \end{aligned}$$

and so  $T_{-u}K \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3))]$  for all  $u \in \hat{G}$ . On the other hand

$$\begin{aligned} \|T_{-u}K\|_{[(p'_1; (p'_2; (p'_3)_\theta)]} &= \|B_{T_{-u}K}\| \\ &= \sup \left\{ \frac{\|B_{T_{-u}K}(f, g)\|_{(p'_3, \theta)}}{\|f\|_{(p'_1, \theta)} \|g\|_{(p'_2, \theta)}} : \|f\|_{(p'_1, \theta)} \leq 1, \|g\|_{(p'_2, \theta)} \leq 1 \right\} \\ &= \sup \left\{ \frac{\|M_{-u}B_K(M_u f, M_{0\hat{G}}g)\|_{(p'_3, \theta)}}{\|M_u f\|_{(p'_1, \theta)} \|M_{0\hat{G}}g\|_{(p'_2, \theta)}} : \|M_u f\|_{(p'_1, \theta)} \leq 1, \|M_{0\hat{G}}g\|_{(p'_2, \theta)} \leq 1 \right\} \\ &= \|B_K\| = \|K\|_{[(p'_1; (p'_2; (p'_3)_\theta)]}. \end{aligned}$$

Thus, by Proposition 7, there exists  $C > 0$  such that

$$\begin{aligned} \left| \int_G (T_{-u}K)^\vee(y) d\lambda(y) \right| &\leq C |A|^{-\frac{1}{q}} \|T_{-u}K\|_{[(p'_1; (p'_2; (p'_3)_\theta)]} \\ (2.51) \quad &= C |A|^{-\frac{1}{q}} \|K\|_{[(p'_1; (p'_2; (p'_3)_\theta)]}. \end{aligned}$$

Since  $T_{-u}K \in \ell^1(\hat{G})$ , from (2.51) we write

$$\begin{aligned} |K(u)| &= |T_{-u}K(0_{\hat{G}})| = \left| ((T_{-u}K)^\vee)^\wedge(0_{\hat{G}}) \right| = \left| \int_G (T_{-u}K)^\vee(y) \langle 0_{\hat{G}}, -y \rangle d\lambda(y) \right| \\ (2.52) \quad &= \left| \int_G (T_{-u}K)^\vee(y) d\lambda(y) \right| \leq C |A|^{-\frac{1}{q}} \|K\|_{[(p'_1; (p'_2; (p'_3)]_\theta} \end{aligned}$$

for all  $u \in \hat{G}$ . Since  $\frac{1}{p'_3} < \frac{1}{p'_1} + \frac{1}{p'_2}$ , then  $-\frac{1}{q} < 0$ . Thus the right side of (2.52) approaches to zero for  $|A| \rightarrow \infty$ . This implies  $K = 0$ . But this is a contradiction with the assumption  $K \neq 0$ . Then the assumption  $\frac{1}{p'_3} < \frac{1}{p'_1} + \frac{1}{p'_2}$  is not true. Therefore we conclude  $\frac{1}{p'_3} \geq \frac{1}{p'_1} + \frac{1}{p'_2}$ .  $\square$

**Corollary 2.** Let  $\frac{1}{p'_3} < \frac{1}{p'_1} + \frac{1}{p'_2}$ . If  $A$  is an automorphism of  $G$  satisfying  $|A| > 1$ . Then  $\tilde{M}_\theta[(p'_1; (p'_2; (p'_3)] = \{0\}$ .

*Proof.* Take any  $K \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3]$ . Let  $\Psi \in L^1(G)$  such that  $0 \neq \Psi^\wedge \in \ell^1(\hat{G})$ . By Theorem 7, we have  $\Psi^\wedge K \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3]$ . On the other hand, since  $K \in \tilde{M}_\theta[(p'_1; (p'_2; (p'_3]$ ,  $K$  is bounded function. Then we have  $\Psi^\wedge K \in \ell^1(\hat{G})$ . Since  $\Psi^\wedge K \in \ell^1(\hat{G}) \cap \tilde{M}_\theta[(p'_1; (p'_2; (p'_3]$  and  $\frac{1}{p'_3} < \frac{1}{p'_1} + \frac{1}{p'_2}$ , by Proposition 8, we have  $\Psi^\wedge K = 0$ . Also since  $\Psi^\wedge$  is non-zero function, we obtain  $K = \{0\}$ . This completes the proof.  $\square$

## REFERENCES

- [1] Benedetto John J., Benedetto Robert L. A wavelet theory for local fields and related groups, The Journal of Geometric Analysis, 14, 423–456, 2004.
- [2] Fiorenza A. Duality and reflexivity in grand Lebesgue spaces, Collect. Math. 2000; 51 (2) : 131 – 148.
- [3] Greco L, Iwaniec T, Sbordone C. Inverting the p-harmonic operator, Manuscripta Math.1997; 92 : 259 – 272.
- [4] Gürkanlı AT, Kulak Ö, Sandıkçı A. The spaces of bilinear multipliers of weighted Lorentz type modulation spaces, Georgian Math. J. 2016; 23(3), 351–362.
- [5] Gürkanlı AT, Inclusion and the approximate identities of the generalized grand Lebesgue spaces. Turk J Math. 2018; 42 : 3195 – 3203.
- [6] Gürkanlı AT, Multipliers of Grand and Small Lebesgue Spaces, arXiv:1903.06743v1, 2019
- [7] Iwaniec T, Sbordone C. On the integrability of the Jacobian under minimal hypotheses, Arc. Rational Mech. Anal. 1992; 119 : 129 – 143.
- [8] Kulak Ö, Gürkanlı AT. Bilinear multipliers of weighted Lebesgue spaces and variable exponent Lebesgue spaces, J. Inequal. Appl. 2013; 259.
- [9] Kulak Ö, Gürkanlı AT. Bilinear multipliers of weighted Wiener amalgam spaces and variable exponent Wiener amalgam spaces, J. Inequal. Appl. 2014; 476.
- [10] Y. Katznelson, An Introduction to Harmonic Analysis, John Wiley and Sons, Inc., New York, 1968.
- [11] L. Nachbin, The Haar integral. Robert E. Krieger Publishing Company, Huntington, New York 1976.
- [12] W. Rudin, Fourier Analysis on Groups, Interscience, New York, 1999.
- [13] W. Zelazko, On the algebras  $L^p$  of locally compact groups, Colloquium Mathematicum, Vol. VIII, Fasc. 1, (1961), 115–120.

AMASYA UNIVERSITY, FACULTY OF ARTS AND SCIENCES, DEPARTMENT OF MATHEMATICS, AMASYA,  
TURKEY

*E-mail address:* oznur.kulak@amasya.edu.tr

ISTANBUL AREL UNIVERSITY, FACULTY OF SCIENCE AND LETTERS, DEPARTMENT OF MATHE-  
MATICS AND COMPUTER SCIENCES, İSTANBUL, TURKEY

*Current address:*  $\ddot{\text{y}}$

*E-mail address:* turangurkanli@arel.edu.tr