A note on the phase transition for independent alignment percolation

Marcelo Hilário¹ and Daniel Ungaretti²

^{1,2}Universidade Federal de Minas Gerais

July 2, 2020

Abstract

We study the independent alignment percolation model on \mathbb{Z}^d introduced by Beaton, Grimmett and Holmes [arXiv:1908.07203]. It is a model for random intersecting line segments defined as follows. First the sites of \mathbb{Z}^d are independently declared occupied with probability p and vacant otherwise. Conditional on the configuration of occupied vertices, consider the set of all line segments that are parallel to the coordinate axis, whose extremes are occupied vertices and that do not traverse any other occupied vertex. Declare independently the segments on this set open with probability λ and closed otherwise. All the edges that lie on open segments are also declared open giving rise to a bond percolation model in \mathbb{Z}^d . We show that for any $d \geq 2$ and $p \in (0, 1]$ the critical value for λ satisfies $\lambda_c(p) < 1$ completing the proof that the phase transition is non-trivial over the whole interval (0, 1]. We also show that the critical curve $p \mapsto \lambda_c(p)$ is continuous at p = 1, answering a question posed by the authors in [arXiv:1908.07203].

1 Definition of the model

Alignment percolation has been recently introduced by Beaton, Grimmett and Holmes [2] as a model of random line segments in the hypercubic lattice $\mathbb{L}^d = (\mathbb{Z}^d, \mathbb{E}^d)$ with $d \ge 2$. There, the authors define two versions of the model, referred to as the 'one-choice model' and the 'independent model' as we revisit below.

Fix $d \geq 2$ and let $\Omega := \{0,1\}^{\mathbb{Z}^d}$ be the set of configurations ω on \mathbb{Z}^d . Given $\omega = \{\omega(v); v \in \mathbb{Z}^d\} \in \Omega$ a site $v \in \mathbb{Z}^d$ is said occupied when $\omega(v) = 1$ and vacant when $\omega(v) = 0$. For any parameter $p \in (0,1]$ the independent Bernoulli site percolation model is the measure \mathbb{P}_p^s on Ω under which $\{\omega(v); v \in \mathbb{Z}^d\}$ is a family of independent Bernoulli random variables with mean p. In other words, any site $v \in \mathbb{Z}^d$ independently is occupied or vacant with probability p and 1 - p, respectively.

Denote $\eta(\omega) = \{v \in \mathbb{Z}^d; \ \omega(v) = 1\}$. For a fixed configuration ω , we say that a pair of sites $v_1, v_2 \in \mathbb{Z}^d$ is *feasible* when v_1 and v_2 differ only at a single coordinate and both v_1 and v_2 belong to η but no other site in the line segment v_1v_2 that connects v_1 and v_2 belongs to η . We denote the set of feasible pairs by $F(\eta)$ and define a random graph $G = G(\omega)$ whose vertex set is the set of occupied vertices in ω and whose edges are the feasible pairs; that is, $G(\omega) = (\eta, F(\eta))$.

In [2] two bond percolation models on G are defined by specifying measures on $\{0,1\}^{F(\eta)}$:

One-choice model. For every site $v \in \eta$ choose a feasible pair $f_v = vu \in F(\eta)$ uniformly among all 2*d* available possibilities. Declare $e = uv \in F(\eta)$ open if either $e = f_u$ or $e = f_v$. Otherwise, declare it closed.

Independent model. Fix $\lambda \in [0, 1]$ and declare each edge $e \in F(\eta)$ to be open independently with probability λ .

Both versions described above can be regarded as dependent bond percolation models on $(\mathbb{Z}^d, \mathbb{E}^d)$ by declaring an edge $e \in \mathbb{E}^d$ open if and only if the unique edge of $F(\eta)$ that contains e is open. This gives rise to a random element σ in $\Sigma := \{0, 1\}^{\mathbb{E}^d}$ which in turn induces a random subgraph of \mathbb{Z}^d . We are mainly interested in studying connectivity properties of this random graph.

Here we focus in the independent version of the model. In this context, let P_{λ}^{ω} denote the measure on Σ induced by independent Bernoulli percolation on $\{0,1\}^{F(\eta)}$ with parameter λ . The distribution of the pair $\xi := (\omega, \sigma)$ is the probability measure $\mathbb{P}_{p,\lambda,d}$ on $\Xi := \Omega \times \Sigma$ satisfying

$$\mathbb{P}_{p,\lambda,d}(A \times B) := \int_A P_\lambda^\omega(B) \, \mathrm{d}\mathbb{P}_p^{\mathrm{s}}(\omega) \quad A \subset \Omega, \, B \subset \Sigma \text{ measurable},$$

where Ω and Σ are endowed with their respective cylinder σ -fields. We will often drop the dependency in d from the notation and write simply $\mathbb{P}_{p,\lambda}$. In what follows we may abuse notation and write simply $\xi(v)$ and $\xi(e)$ in order to refer to the state of a site v and an edge e, that is, $\omega(v)$ and $\sigma(e)$, respectively.

The set of *open edges* of \mathbb{L}^d is denoted by $\mathcal{O} = \{e \in \mathbb{E}^d; \xi(e) = 1\}$ and the set of *closed edges* by $\mathcal{C} = \mathbb{E}^d \setminus \mathcal{O}$. A site $v \in \mathbb{Z}^d$ percolates if it belongs to an infinite connected component of \mathcal{O} . We say that \mathcal{O} percolates if there is a site v that percolates.

A standard coupling shows that the model is monotone in λ , see Section 2. Thus, it is reasonable to expect that the model exhibits a non-trivial phase transition in parameter λ . Define

$$\theta(p,\lambda,d) := \mathbb{P}_{p,\lambda,d}(o \text{ percolates}) \text{ and } \lambda_c(p,d) := \sup\{\lambda \ge 0; \ \theta(p,\lambda,d) = 0\},$$

where o denotes the origin in \mathbb{Z}^d . We denote by $p_c^{\text{site}}(d)$ and $p_c^{\text{bond}}(d)$ the critical points for site and bond independent percolation on \mathbb{L}^d . It follows from [2, Theorem 2.4] that $\lambda_c(p,d) \geq \frac{p}{2d-1} > 0$ and also that there is a universal $p^* > 0$ such that

$$\lambda_c(p,d) < 1$$
 if $p < p^*$ or $p > p_c^{\text{site}}(d)$.

Since $p_c^{\text{site}}(d) \sim \frac{1}{2d}$ and is strictly decreasing on d we know there is some $d_0(p^*)$ such that for $d \geq d_0(p^*)$ we have $\lambda_c(p,d) \in (0,1)$. However, since we do not have a good control on p^* it may be the case that $p^* < p_c^{\text{site}}(d)$. Therefore, their result does not rule out that $\lambda_c(p,d) = 1$ on $[p^*, p_c^{\text{site}}(d)]$. In Theorem 1 we bridge this gap providing non-trivial upper bounds on $\lambda_c(p,d)$ for every $p \in (0,1]$. For $d \geq 3$ it is not hard to find an upper bound which holds uniformly over all the interval (0,1] by a simple comparison to Bernoulli percolation on the hexagonal lattice. For the case d = 2, that seems to be more delicate, we employ renormalization methods in order to obtain the result.

Theorem 1. Let $p_c^{\bigcirc} = 1 - 2\sin(\pi/18)$ be the critical point for independent bond percolation on the hexagonal lattice¹. For any $p \in (0, 1]$ and $d \ge 3$ we have that

$$\lambda_c(p,d) \le p_c^{\bigcirc}.\tag{1}$$

Moreover, for every $p \in (0,1]$ there is $\lambda_0(p) \in (0,1)$ such that $\lambda_c(p,2) \leq \lambda_0$.

In [2] the authors conjecture that the function $p \mapsto \lambda_c(p, d)$ is non-increasing and continuous, as suggested by numerical simulation. As already mentioned there, one difficulty in obtaining results in this direction stems from the fact that it seems hard to couple the model with different values of p and still keep control of how the connectivity changes, see Section 2. Currently, there is not much progress in proving strict monotonicity and continuity of $\lambda_c(\cdot, d)$, however, our renormalization arguments can be used to prove continuity as p approaches 1.

¹For the hexagonal lattice the critical point is known exactly, see [5, Chapter 3]



Figure 1: For $d \ge 3$, Theorem 1 complements Figure 2.1 of [2] with an explicit upper bound; the critical curve is inside the highlighted region. For d = 2 we do not have explicit bounds but we ensure the critical curve is non-trivial in the interval $[p^*, p_c^{\text{site}}(2)]$.

Theorem 2. Fix $d \ge 2$. For every $\lambda < p_c^{\text{bond}}(d)$ there exists $p_0(\lambda, d) \in (0, 1)$ such that

$$\mathbb{P}_{p,\lambda,d}(\mathcal{O} \ percolates) = 0 \quad for \ every \ p > p_0.$$
⁽²⁾

In particular, the function $p \mapsto \lambda_c(p, d)$ is continuous at p = 1.

Theorem 2 answers affirmatively a question posed in [2] (see the paragraph preceding Remark 2.6 therein).

Let us say a word about the proofs of Theorems 1 and 2. As mentioned above, the proof of Theorem 1 for $d \geq 3$ is obtained by embedding the hexagonal lattice in \mathbb{Z}^d in an appropriate way. This comparison does not work for d = 2 and to prove that $\lambda_c(\cdot, 2) < 1$ we develop a multiscale renormalization.

Renormalization is a classical technique that has been successfully applied to a vast range of areas in mathematics and physics. Instead of providing a complete account on the applicability of the method, we would just like to mention that it has been successfully used to study models in which correlations decay suitably with distance. To mention a few examples, it has been employed to show the existence of phase transition on dependent percolation e.g. random interlacements [9] and Poisson cylinder model [10], to study the long-range behavior of random walks on random environments [3] and on detection problems [1].

The structure of the argument, as we use here, can be summarized in the following steps.

S1. (Recurrence inequality) Define a sequence of scales L_k increasing sufficiently fast and consider a family of events $\{A_k(x); k \in \mathbb{N}, x \in \mathbb{Z}^d\}$ where each $A_k(x)$ is measurable with respect to the restriction of the process to a finite region having diameter of order L_k around x. Moreover the family is required to be *cascading*, meaning that the occurrence of $A_{k+1}(x)$ implies the occurrence of events $A_k(x')$ and $A_k(x'')$ for points x' and x'' well-separated in scale k but close enough in scale k + 1. Defining $q_{L_k}(p, \lambda, d) := \mathbb{P}_{p,\lambda,d}(A_k(x))$ which does not depend on x (due to translation invariance) one can deduce that

$$q_{L_{k+1}}(p,\lambda,d) \le g\left(\frac{L_{k+1}}{L_k}\right) \cdot \left[q_{L_k}(p,\lambda,d)^2 + \operatorname{error}_k(p,\lambda,d)\right],\tag{3}$$

where g is some positive real function and the error term is small due to the decay of correlations and the choice of L_k . In summary, inequality (3) relates the probabilities of the events A_k at two successive scales by using the structure of cascading events and decay of correlations.

S2. (Inductive step) Assuming that $q_{L_{k_0}}(p, \lambda, d) \leq f(L_{k_0})$ for some choice of $k_0(p, \lambda, d)$ and positive real function f(x) tending to zero as $x \to \infty$, we show inductively that

$$q_{L_k}(p,\lambda,d) \le f(L_k) \quad \text{for every } k \ge k_0.$$
 (4)

S3. (Trigger) Find a choice of p, λ, d, k_0 so that $q_{L_{k_0}}(p, \lambda, d) \leq f(L_{k_0})$. Once we fix such parameters, we can trigger the induction in (4) and conclude that $q_{L_k}(p, \lambda, d)$ indeed decays at least as fast as $f(L_k)$ as we increase k.

The proofs of Theorem 1 for d = 2 and of Theorem 2 follow essentially the structure above. The sequence of events $A_k(x)$ are either the existence of an open path or a dual path crossing a box of size L_k around x. Apart from the difference in the definition of the events $A_k(x)$, the trigger step S3 is also performed differently. For Theorem 1 we fix $p \in (0, 1)$ and take λ sufficiently close to 1, whereas for Theorem 2 we fix $\lambda < p_c^{\text{bond}}(d)$ and take p sufficiently close to 1. Our application is reasonably simple and showcases the power of multiscale renormalization. This technique can be used in similar percolation problems as long as it presents a good decay of correlations which entails a small error term in (3), see Lemma 1.

Throughout the text, we will use c or C to indicate positive constants whose values may change from line to line. Numbered constants like c_0, c_1, \ldots or k_0, k_1, \ldots will have their values fixed at the first time they appear and will remain fixed. We may indicate the dependence of constants on other parameters, for instance, $c_i(d)$ is a constant whose value depend on d but not on any other parameter of the model.

Let us conclude this section by summarizing the structure of the paper. On Section 2 we discuss some properties of the model: monotonicity, uniqueness of the infinite cluster, the lattice condition, and decay of correlations. On Section 3 we give the multiscale framework that will be used for proving Theorems 1 and 2, and then complete their proofs on Sections 3.1 and 3.2, respectively.

2 Properties of the independent model

In this section we collect some useful properties fulfilled by independent alignment. Some of them were were already mentioned in [2].

Let us first describe some monotonicity properties. Since we are concerned with percolation of the subgraph induced by the set of open edges, we will compare two configurations $\xi, \xi' \in \Xi$ using the natural partial order in Σ , that is,

$$\xi \preccurlyeq \xi'$$
 if and only if $\xi(e) \le \xi'(e), \ \forall e \in \mathbb{E}^d$.

Recall that $\mathbb{P}_{p,\lambda}$ is the underlying probability distribution of the model and denote by $\xi_{p,\lambda}$ a random element distributed as $\mathbb{P}_{p,\lambda}$. We may possibly omit one of the parameters in $\xi_{p,\lambda}$ when there is no risk of confusion.

For a fixed $p \in (0, 1]$ the $\xi_{p,\lambda}$ can be constructed in a monotone way in λ (with respect to this partial order) using the standard coupling. For a fixed $\lambda \in [0, 1]$ the picture is more complex. This was already mentioned on [2] but we elaborate on this discussion here.

Proposition 1. Fix $\lambda \in (0, 1]$ and denote by ξ_p a random element of Ξ with distribution given by $\mathbb{P}_{p,\lambda}$. For any $0 < p_1 < p_2 \leq 1$ we have that almost surely neither $\xi_{p_1} \preccurlyeq \xi_{p_2}$ nor $\xi_{p_2} \preccurlyeq \xi_{p_1}$.

Proof. Recall that an event $A \subset \Sigma$ is said *increasing* if $x \in A$ and $y \succeq x$ implies $y \in A$. By Strassen's Theorem [8, Theorem 2.4], there exists a coupling of (ξ_{p_1}, ξ_{p_2}) such that $\xi_{p_1} \succeq \xi_{p_2}$ if and only if we have $\mathbb{P}_{p_1}(A) \ge \mathbb{P}_{p_2}(A)$ for any increasing event A.

The inequality $\mathbb{P}_{p_1}(A) \geq \mathbb{P}_{p_2}(A)$ actually holds for some increasing events. For example, if for any finite $E \subset \mathbb{E}^d$ we denote by $\xi(E)$ the restriction of ξ to the edges of E and write $\xi(E) \equiv 1$ if $\xi(e) = 1$ for every edge in E then we claim that

$$\mathbb{P}_{p_1,\lambda}(\xi(E) \equiv 1) \ge \mathbb{P}_{p_2,\lambda}(\xi(E) \equiv 1)$$
 for every finite $E \subset \mathbb{E}^d$.

In order to prove this, we use a standard coupling for the underlying site percolation configuration. Let $\{U_v; v \in \mathbb{Z}^d\}$ be a family of i.i.d. random variables with the uniform distribution on [0, 1] and for $p \in (0, 1]$ define $\eta_p \in \Omega$ by $\eta_p(v) := \mathbb{1}\{U_v \leq p\}$ for every $v \in \mathbb{Z}^d$. On Ω we have a partial ordering defined by $\eta \preccurlyeq \eta'$ if and only if $\eta(v) \le \eta'(v)$ for every $v \in \mathbb{Z}^d$. Notice that the coupling above makes $\eta_{p_1} \preccurlyeq \eta_{p_2}$. Denote by $t(\eta_p, E)$ the number of feasible pairs of η_p that contain at least one edge that belongs to E. We have

$$\mathbb{P}_{p,\lambda}(\xi(E) \equiv 1) = \mathbb{E}\left[\mathbb{P}_{p,\lambda}(\xi(E) \equiv 1 \mid \eta_p)\right] = \mathbb{E}\left[\lambda^{t(\eta_p, E)}\right].$$

Adding new sites to a configuration can only increase $t(\cdot, E)$. Thus, we have

$$\mathbb{P}_{p_1,\lambda}(\xi(E) \equiv 1) = \mathbb{E}\big[\lambda^{t(\eta_{p_1},E)}\big] \ge \mathbb{E}\big[\lambda^{t(\eta_{p_2},E)}\big] = \mathbb{P}_{p_2,\lambda}(\xi(E) \equiv 1).$$

On the other hand, the inequality may fail for some increasing events A. In fact, as already noted in [2] the increasing event

 $A = \{o \text{ has an open incident edge}\}$

has probability $p[1 - (1 - \lambda)^{2d}] + (1 - p)[1 - (1 - \lambda)^d]$ which is increasing in p.

Conjecture 2.5 of [2] states that the function $\lambda_c(\cdot, d)$ should be continuous and strictly increasing. Proposition 1 does not invalidate this conjecture but shows that to compare the model for different values of p may not be a simple task.

Another known property of the model is that whenever \mathcal{O} percolates, the infinite cluster is unique $\mathbb{P}_{p,\lambda,d}$ -a.s. The proof of this result follows from the classical argument of Burton and Keane [4], and is part of [2, Theorem 2.4]. This property may be useful, for instance to prove continuity of the critical curve but we will not use it in this paper.

We now discuss other properties that were not presented in [2]. Denote by μ the marginal of $\mathbb{P}_{p,\lambda}$ on Σ . Moreover, for $x, y \in \Sigma$ define $x \lor y, x \land y \in \Sigma$ as

$$(x \lor y)(e) := \max\{x(e), y(e)\}$$
 and $(x \land y)(e) := \min\{x(e), y(e)\}$ for every $e \in \mathbb{E}^d$.

It is worth noticing that μ fails to satisfy the so-called *lattice condition*

$$\mu(x \vee y)\mu(x \wedge y) \ge \mu(x)\mu(y) \quad \text{for every } x, y \in \Sigma.$$
(5)

Indeed, consider $E = \{\pm e_1, \pm e_2\} \subset \mathbb{E}^d$ and the configurations on $\{0, 1\}^E$ given by

$$x(e) = \mathbb{1}\{\pm e_1\}(e)$$
 and $y(e) = \mathbb{1}\{e_1, e_2\}(e)$.

It is straightforward to check that

$$\begin{aligned} \mu(x \lor y)\mu(x \land y) &- \mu(x)\mu(y) \\ &= \left[p\lambda^3(1-\lambda)\right] \cdot \left[p\lambda(1-\lambda)^3\right] - \left[p\lambda^2(1-\lambda)^2 + (1-p)\lambda(1-\lambda)\right] \cdot \left[p\lambda^2(1-\lambda)^2\right] < 0. \end{aligned}$$

Proving (5) is the usual strategy to show that the measure μ satisfies the FKG inequality, also known as positive association. However, the FKG inequality is not equivalent to the lattice condition; for background on positive association and its relation to the lattice condition we refer to [6, Chapter 2]. This raises the question whether the model satisfies the FKG inequality.

Although independent alignment percolation model does not exhibit great monotonicity properties, it does present fast correlation decay with distance. This fact will allow us to implement the multiscale renormalization approach outlined in Section 1.

For $x \in \mathbb{Z}^d$ and L > 0 let us define

$$B(x,L) := \left\{ z \in \mathbb{Z}^d; \ \|z - x\|_{\infty} \le L \right\} \quad \text{and} \quad \partial B(x,L) := \left\{ z \in \mathbb{Z}^d; \ \|z - x\|_{\infty} = \lfloor L \rfloor \right\}.$$

For $B \subset \mathbb{Z}^d$, we say that an event A is supported on edges of B if A belongs to the smallest σ -algebra that makes $\{\xi(e); e = \{u, v\} \in \mathbb{E}^d, u, v \in B\}$ measurable.

Lemma 1 (Decay of correlations). Let L > 0 and $p \in (0, 1)$. For $x_1, x_2 \in \mathbb{Z}^d$ with $D = ||x_1 - x_2||_{\infty} > 2L$, let A_i , i = 1, 2, be events supported on edges of $B_i := B(x_i, L)$, respectively. Then,

$$\left|\operatorname{Cov}_{p,\lambda}(A_1, A_2)\right| \le 4 \cdot (2L+1)^{d-1} \cdot e^{-\alpha(p) \cdot (D-2L)}$$
 (6)

with $\alpha(p) := -\log(1-p) > 0.$

Proof. The assumption on D implies $dist(B_1, B_2) = D - 2\lfloor L \rfloor \ge 1$ and, in particular, $B_1 \cap B_2 = \emptyset$. Choose the smallest $1 \le t \le d$ such that $D = |x_{1,t} - x_{2,t}|$ and notice that we can separate B_1 from B_2 by a hyperplane orthogonal to e_t the *t*-th vector of the canonical basis of \mathbb{R}^d . Denote by $\pi_t(B)$ the projection of a set $B \subset \mathbb{R}^d$ into the subspace e_t^{\perp} (perpendicular to e_t) and define

$$\Pi_t = \pi_t(B_1) \cap \pi_t(B_2).$$

If we have $\Pi_t = \emptyset$ then A_1 and A_2 are independent and their covariance is zero. Otherwise, for every $z \in \Pi_t$ define $I_z = [a_z, b_z] \cap \mathbb{Z}^d$ as the unique line segment supported on the line $\pi_t^{-1}(z)$ with $a_z \in B_1, b_z \in B_2$ and such that $(I_z \setminus \{a_z, b_z\}) \cap (B_1 \cup B_2) = \emptyset$. Define the event

$$C = \{ \exists z \in \Pi_t \text{ such that } \omega(v) = 0, \, \forall v \in I_z \}$$

$$\tag{7}$$

and notice that on C^c the state of edges on B_1 and B_2 are independent. Indeed, for $\omega \in C^c$ notice that if $F_i(\omega)$ denotes the feasible edges intersecting B_i and $f_i = \{u_i, v_i\} \in F_i(\omega)$ then $|f_1 \cap f_2| \leq 1$. Thus, omitting the dependency on p and λ we have that

$$\mathbb{P}(A_1 \cap A_2 \cap C^c) = \mathbb{P}(A_1 \cap A_2 \mid C^c)\mathbb{P}(C^c) = \mathbb{P}(A_1 \mid C^c)\mathbb{P}(A_2 \mid C^c)\mathbb{P}(C^c).$$

Using twice that for any event E one has $|\mathbb{P}(E) - \mathbb{P}(E \cap C^c)| \leq \mathbb{P}(C)$, we can bound

$$\begin{aligned} \left| \mathbb{P}(A_1)\mathbb{P}(A_2) - \mathbb{P}(A_1 \cap C^c)\mathbb{P}(A_2 \cap C^c) \right| \\ &= \left| \mathbb{P}(A_1)[\mathbb{P}(A_2) - \mathbb{P}(A_2 \cap C^c)] + \mathbb{P}(A_2 \cap C^c)[\mathbb{P}(A_1) - \mathbb{P}(A_1 \cap C^c)] \right| \\ &\leq (1 + \mathbb{P}(C^c)) \cdot \mathbb{P}(C), \end{aligned}$$

which leads to the estimate

$$\begin{aligned} \left| \operatorname{Cov}(A_1, A_2) \right| &= \left| \mathbb{P}(A_1 \cap A_2) - \mathbb{P}(A_1) \mathbb{P}(A_2) \right| \\ &\leq \mathbb{P}(C) + (1 + \mathbb{P}(C^c)) \mathbb{P}(C) + \left| \mathbb{P}(A_1 \cap A_2 \cap C^c) - \mathbb{P}(A_1 \cap C^c) \mathbb{P}(A_2 \cap C^c) \right| \\ &= (2 + \mathbb{P}(C^c)) \mathbb{P}(C) + \left| \mathbb{P}(A_1 \mid C^c) \mathbb{P}(A_2 \mid C^c) \cdot \left[\mathbb{P}(C^c) - \mathbb{P}(C^c)^2 \right] \right| \\ &\leq 2\mathbb{P}(C) + \mathbb{P}(C^c) \mathbb{P}(C) + \mathbb{P}(C^c) - \mathbb{P}(C^c)^2 \\ &= 4\mathbb{P}(C) - 2\mathbb{P}(C)^2. \end{aligned}$$

Finally, we use the union bound and the fact that $|\Pi_t| \leq (2L+1)^{d-1}$ to estimate

$$\mathbb{P}(C) \le (2L+1)^{d-1} \cdot (1-p)^{D-2L}.$$

Remark 1. Lemma 1 is stated for $\mathbb{P}_{p,\lambda}$. However, it can be adapted for the 'one-choice model' and more generally for any measures P^{ω} on the feasible pairs $F(\eta)$ satisfying that random variables $\{\xi(f_i); i \in I\}$ are mutually independent for any endpoint-disjoint family $\{f_i; i \in I\} \subset F(\eta)$. To see why this is true, just consider instead of the event C appearing in (7) the event

$$\hat{C} = \{ \exists f_1 \in F_1(\omega), f_2 \in F_2(\omega) \text{ such that } |f_1 \cap f_2| \ge 1 \}.$$

To prevent \tilde{C} from occurring we just have to ensure each line I_z with $z \in \pi_t(B_1) \cup \pi_t(B_2) =: \Pi_t$ contains 2 sites in $\eta(\omega)$ lying in the region between the two boxes. The decay of correlations will follow once one notes that $|\Pi_t| \leq 2(2L+1)^{d-1}$ and changing $\alpha(p)$ accordingly. Moreover, we notice that the bound on (6) is independent of λ .

3 Multiscale renormalization

In this section we build a multiscale renormalization scheme that allow us to prove Theorem 1 in the case d = 2 and Theorem 2. A key step will be the use of the correlation decay in Lemma 1.

We start by defining the sequence of scales L_0, L_1, \ldots along which we analise the model. Let

$$L_0 := 10^4$$
 and $L_{k+1} = L_k^{3/2}$, (8)

that is, $L_k = L_0^{(3/2)^k}$ is a sequence growing super-exponentially fast. This same sequence is used in the proofs of both Theorem 1 in the case d = 2 and Theorem 2. The family of cascading events $\{A_k(x)\}_{x \in \mathbb{Z}^d, k \in \mathbb{Z}_+}$ needs to be defined properly for each proof, but $A_k(x)$ will always be supported in the annular region $B(x, 10L_k) \setminus B(x, L_k)$. However once these events are defined, Steps **S1** and **S2** in Section 1 can be carried on exactly the same way in both proofs.

For each k, let $\mathcal{L}_k^1 \subset \partial B(o, L_{k+1})$ and $\mathcal{L}_k^2 \subset \partial B(o, 5L_{k+1})$ be minimal collections of points such that

$$\partial B(o, L_{k+1}) \subset \bigcup_{x \in \mathcal{L}_n^1} B(x, L_k) \text{ and } \partial B(o, 5L_{k+1}) \subset \bigcup_{x \in \mathcal{L}_n^2} B(x, L_k).$$

The following lemma gives bounds on the size of \mathcal{L}_k^i .

Lemma 2. There are constants c = c(d) and C = C(d) such that for i = 1, 2 we have

$$c \cdot \left(\frac{L_{k+1}}{L_k}\right)^{d-1} \le |\mathcal{L}_k^i| \le C \cdot \left(\frac{L_{k+1}}{L_k}\right)^{d-1}.$$
(9)

Proof. We prove the result for \mathcal{L}_k^1 and the same reasoning applies for \mathcal{L}_k^2 . Notice that any $B(x, L_k)$ with $x \in \partial B(o, L_{k+1})$ satisfies

$$\lfloor L_k \rfloor^{d-1} \le |B(x, L_k) \cap \partial B(o, L_{k+1})| \le d \cdot (2\lfloor L_k \rfloor + 1)^{d-1}.$$
(10)

Since $\{B(x, L_k); x \in \mathcal{L}_k^1\}$ covers $\partial B(o, L_{k+1})$, the upper bound on (10) implies that

$$2d \cdot (2\lfloor L_{k+1} \rfloor + 1)^{d-1} = |\partial B(o, L_{k+1})| \le |\mathcal{L}_k^1| \cdot d \cdot (2\lfloor L_k \rfloor + 1)^{d-1}$$

and thus the lower bound on (9) follows. For the upper bound consider $\mathcal{N}_0 = \emptyset$ and, inductively, while $\{B(x, \frac{1}{2}L_k); x \in \mathcal{N}_j\}$ is a disjoint collection and there is some

$$y \in \partial B(o, L_{k+1}) \setminus \bigcup_{x \in \mathcal{N}_j} B(x, \frac{1}{2}L_k),$$

define $\mathcal{N}_{j+1} := \mathcal{N}_j \cup \{y\}$. This process ends in finitely many steps, producing a set \mathcal{N} satisfying

$$2d \cdot (2\lfloor L_{k+1} \rfloor + 1)^{d-1} = |\partial B(o, L_{k+1})| \ge |\mathcal{N}| \cdot \lfloor \frac{1}{2} L_k \rfloor^{d-1}$$

using (10). Finally, just notice that we can take $\mathcal{L}_k^1 \subset \mathcal{N}$, since for any $y \in \partial B(o, L_{k+1})$ we have

$$\operatorname{dist}(y, B(x, \frac{1}{2}L_k)) < \frac{1}{2}L_k \quad \text{for some } x \in \mathcal{N} \implies ||x - y||_{\infty} \le L_k.$$

We require two properties for our events $A_k(x)$.

P1. $\mathbb{P}_{p,\lambda,d}(A_k(x))$ does not dependent upon the choice of x which allows us to define

$$q_k(p,\lambda,d) := \mathbb{P}_{p,\lambda,d}(A_k(o)).$$

P2. The occurrence of event $A_{k+1}(o)$ implies that there are $x_i \in \mathcal{L}_k^i$ such that $A_k(x_i)$ also occur, for i = 1, 2. Heuristically, we say that events $A_k(x)$ are cascading since if an event occurs on scale k + 1 it must have occurred on two well-separated regions for the previous scale.

Properties **P1** and **P2** imply the validity of Steps **S1** and **S2**.

Lemma 3 (Recurrence Inequality). Let $\{A_k(x); x \in \mathbb{Z}^d, k \in \mathbb{N}\}\$ be a collection of events on Ξ satisfying Properties P1 and P2. Then,

$$q_{k+1}(p,\lambda,d) \le c_0(d) \cdot L_k^{d-1} q_k(p,\lambda,d)^2 + c_1(d) \cdot L_k^{2d-2} e^{-3\alpha(p) \cdot L_k^{3/2}},\tag{11}$$

where c_0 and c_1 are positive constants depending only on d.

Proof. The boxes $B(x_i, 10L_k)$ given by property **P2** are well-separated since

$$\|x_1 - x_2\|_{\infty} \ge \lfloor 5L_{k+1} \rfloor - \lfloor L_{k+1} \rfloor \ge 4\lfloor L_{k+1} \rfloor \quad \text{for every } x_i \in \mathcal{L}_k^i.$$

Using the fact that $A_k(x)$ is supported on edges inside $B(x, 10L_k)$, we have by Lemma 1 that there exists a constant $c_2 = c_2(d) > 0$ such that

$$|\operatorname{Cov}_{p,\lambda}(A_k(x_1), A_k(x_2))| \le c_2 \cdot L_k^{d-1} \cdot e^{-\alpha(p)(4\lfloor L_{k+1} \rfloor - 20L_k)} \le c_2 \cdot L_k^{d-1} \cdot e^{-3\alpha(p)L_{k+1}}$$

for every k. Lemma 2 implies that $|\mathcal{L}_k^i| \leq c_3 \cdot \frac{L_{k+1}}{L_k} = c_3 \cdot L_k^{(d-1)/2}$ for some constant $c_3(d) > 0$. Thus, we can write

$$q_{k+1}(p,\lambda,d) \leq \sum_{x_1 \in \mathcal{L}_k^1, x_2 \in \mathcal{L}_k^2} \mathbb{P}_{p,\lambda,d}(A_k(x_1) \cap A_k(x_2))$$

$$\leq |\mathcal{L}_k^1| \cdot |\mathcal{L}_k^2| \cdot \left[q_k(p,\lambda,d)^2 + c_2(d) \cdot L_k^{d-1} e^{-3\alpha(p)L_k^{3/2}}\right]$$

$$\leq c_3(d)^2 \cdot L_k^{d-1} q_k(p,\lambda,d)^2 + c_3(d)^2 \cdot c_2(d) \cdot L_k^{2d-2} e^{-3\alpha(p) \cdot L_k^{3/2}}.$$
 (12)

The proof is finished by defining $c_0 = c_3^2$ and $c_1 = c_3^2 \cdot c_2$.

Lemma 4 (Inductive step). Let $\{A_k(x); x \in \mathbb{Z}^d, k \in \mathbb{N}\}\$ be a collection of events on Ξ satisfying Properties P1 and P2. There is $k_0 = k_0(p, d)$ such that if for some $k_1 \ge k_0$ one has

$$q_{k_1}(p,\lambda,d) \le L_{k_1}^{-2d} \tag{13}$$

then

$$q_k(p,\lambda,d) \le L_k^{-2d} \quad holds \text{ for every } k \ge k_1.$$
(14)

Proof. Choose $k_0(d, p)$ as the smallest integer k such that

$$c_0(d)L_k^{-1} \le \frac{1}{2}$$
 and $c_1(d) \cdot L_k^{5d-2}e^{-3\alpha(p)\cdot L_k^{3/2}} \le \frac{1}{2}$

where $c_0(d), c_1(d)$ are given by Lemma 3. This is always possible since both left-hand sides tend to zero when $k \to \infty$. Suppose there is some $k(p, \lambda, d) \ge k_0$ satisfying the inequality in (14). Then, we can write

$$\begin{aligned} \frac{q_{k+1}(p,\lambda,d)}{L_{k+1}^{-2d}} &\leq c_0(d) \cdot L_k^{d-1+3d} q_k(p,\lambda,d)^2 + c_1(d) \cdot L_k^{2d-2+3d} e^{-3\alpha(p) \cdot L_k^{3/2}} \\ &\leq c_0(d) \cdot L_k^{4d-1} L_k^{-4d} + c_1(d) \cdot L_k^{5d-2} e^{-3\alpha(p) \cdot L_k^{3/2}} \\ &\leq 1, \end{aligned}$$

by our choice of k_0 . This means that the inequality in (14) carries on to the next scale k + 1. The result follows by induction.

3.1 Phase transition for alignment percolation

We first consider the case $d \ge 3$, in which no renormalization is needed. Indeed, a construction that appeared in [7] fits very nicely to this case.

Proof of Theorem 1, case $d \ge 3$. We show that it is possible embed the hexagonal lattice in \mathbb{Z}^d in such a way that the state of their edges are independent. Notice that it suffices to prove the claim on the case d = 3, since for larger d one can run the same argument on the subset $\{x \in \mathbb{Z}^d; x_i = 0, \forall 4 \le i \le d\}$.

Thus, let us consider d=3 and for a fixed $k\in\mathbb{Z}$ we write

 $V_1 := \{(x, y, z) \in \mathbb{Z}^3; x + y + z = k\}$ and $V_2 := \{(x, y, z) \in \mathbb{Z}^3; x + y + z = k + 1\}.$



Figure 2: Copy of the hexagonal lattice with the property that every edge is independent. In the picture above, we used k = 6 to obtain $V_1 = \{\bullet\}$ and $V_2 = \{\blacksquare\}$. Solid lines represent edges of the subgraph and dashed lines are present to help visualizing the lattice as a subset of \mathbb{R}^3 .

The desired subgraph of \mathbb{L}^3 is obtained by considering the vertex set $V_1 \cup V_2$ and the edges uv with $u \in V_1$, $v \in V_2$ and $||u - v||_{\infty} = 1$. As shown in Figure 2, this defines a subgraph of \mathbb{Z}^3 that is isomorphic to the hexagonal lattice. Under the measure P_{λ}^{ω} , the state of each edge is independent since a line parallel to one of the canonical directions intersects the planes V_1 and V_2 in precisely one point each. Therefore, the alignment percolation restricted to such a graph is just independent Bernoulli percolation whose critical point equals $p^{\bigcirc} = 1 - 2\sin(\pi/8)$.

We now focus on the case d = 2. Consider the dual graph $(\mathbb{L}^2)^*$ of \mathbb{L}^2 , defined as the graph with vertex set $\mathbb{Z}^2 + (1/2, 1/2)$ and edge set $(\mathbb{E}^2)^*$ connecting sites at Euclidean distance 1. Therefore, $(\mathbb{L}^2)^*$ and \mathbb{L}^2 are isomorphic. Moreover, to each edge e^* in $(\mathbb{E}^2)^*$ it corresponds a unique edge e in \mathbb{E}^2 , so that e and e^* intersect at right angle. Given any configuration $\xi \in \Xi$, we denote \mathcal{O}^* (resp. \mathcal{C}^*) the set of dual edges whose corresponding primal edge is open (resp. closed). Notice however that, unlike when we consider independent bond percolation, for alignment percolation the distributions of \mathcal{O}^* and \mathcal{O} are not the same. Notice also that any event can be defined in terms of the statuses of either the primal or the dual edges.

Define for $x \in \mathbb{Z}^2$ events

 $A_k(x) := \{ \text{there is an open circuit on } B(x, 10L_k) \setminus B(x, L_k) \text{ surrounding } B(x, L_k) \}^c,$

which can be seen as the event on which there exists a dual path of C^* edges from the inside of $B(x, L_k)$ to the outside of $B(x, 10L_k)$. The key fact is that the events $A_k(x)$ satisfy Properties **P1** and **P2**, allowing us to use Lemmas 3 and 4.

Lemma 5 (Trigger for Theorem 1). For any $p \in (0,1)$ there are $k_0(p) > 0$ and $\lambda_0(p) \in (0,1)$ such that for every $\lambda \ge \lambda_0$ and $k \ge k_0$

$$q_k(p,\lambda) \leq L_k^{-4}.$$

Proof. Let $k_0(p)$ be given as in Lemma 4. Now, let us check that we can take λ sufficiently close to 1 in order to ensure that $q_{k_0}(p,\lambda) \leq L_{k_0}^{-4}$. Let $N(k_0)$ be the total number of edges from \mathbb{E}^d with some extremity in $B(o, 10L_{k_0})$ and define $\lambda_0 = (1 - L_{k_0}^{-4})^{N(k_0)^{-1}}$. Notice that we can write

$$q_{k_0}(p,\lambda) = \mathbb{P}_{p,\lambda}(A_{k_0}(o)) \le 1 - \mathbb{E}_{p,\lambda}\left[\mathbb{P}_{p,\lambda}(\text{all edges inside } B(o,10L_{k_0}) \text{ are open } \mid \eta)\right].$$

Since k_0 is fixed, the number of edges of $F(\eta)$ that have some extremity inside $B(o, 10L_{k_0})$ is bounded from above by $N(k_0)$. Thus, we have for every $\lambda \ge \lambda_0$,

$$q_{k_0}(p,\lambda) \le 1 - \lambda^{N(k_0)} \le 1 - \lambda_0^{N(k_0)} = L_{k_0}^{-4}$$

The result follows from Lemma 4.

Proof of Theorem 1, case d = 2. We now show that for any $\lambda \geq \lambda_0(p)$ we have that \mathcal{O} percolates $\mathbb{P}_{p,\lambda}$ -a.s. Since we are working on the plane, we have that if \mathcal{O} does not percolate there must be a sequence γ_n of disjoint circuits in $(\mathbb{L}^2)^*$ that go around the origin with $\gamma_n \subset \mathcal{C}^*$ and

$$\operatorname{dist}\left(o, \gamma_n \cap (\mathbb{R}^+ \times \{0\})\right) \to \infty.$$
(15)

For each $k \geq 0$ consider the points $\{x_{k,i}\} \subset \mathbb{R}^+ \times \{0\}$ defined by

$$x_{k,i} := (\lceil 10L_k \rceil + (i-1)2\lfloor L_k \rfloor, 0) \quad \text{for } 1 \le i \le s_k,$$

where s_k is determined below. Notice that

$$B(x_{k,i}, 10L_k) \subset \mathbb{R}^+ \times \mathbb{R}$$
 and boxes $B(x_{k,i}, L_k)$ and $B(x_{k,i+1}, L_k)$ are adjacent.



Figure 3: Multiscale renormalization for Theorem 1, case d = 2. If \mathcal{O} does not percolate, circuits $\gamma_n \subset \mathcal{C}^*$ must intersect boxes $B(x_{k,i}, L_k)$ with large k, as exemplified by the highlighted box.

Choosing s_k as the minimum index i such that $x_{k,i} \ge x_{k+1,1}$ ensures that the family of boxes $\{B(x_{k,i}, L_k); k \ge 0, 1 \le i \le s_k\}$ covers the half-line $(\lfloor 10L_0 \rfloor, \infty) \times \{0\}$. Moreover, the sequence s_k satisfies

$$s_k \le 5 \cdot \frac{L_{k+1}}{L_k} = 5L_k^{1/2}.$$

Choose some ordering \tilde{A}_l for the collection of events $\{A_k(x_{k,i}); k \ge 0, 1 \le i \le s_k\}$. The existence of the sequence of circuits $\gamma_n \subset \mathcal{C}^*$ satisfying (15) implies

$$\mathbb{P}_{p,\lambda}(\{\mathcal{O} \text{ percolates}\}^c) \leq \mathbb{P}_{p,\lambda}(\tilde{A}_l, \text{ i.o.}).$$

However, we have by Lemma 5 that

$$\sum_{l\geq 1} \mathbb{P}_{p,\lambda}(\tilde{A}_l) = \sum_{k\geq 0} \sum_{i=1}^{s_k} q_k(p,\lambda) \le \sum_{k\geq 0} 5 \cdot L_k^{-7/2} < \infty$$

and Borel-Cantelli lemma implies that $\mathbb{P}_{p,\lambda}(\tilde{A}_l, \text{ i.o.}) = 0$. We conclude that for $\lambda \geq \lambda_0(p)$ the set \mathcal{O} percolates $\mathbb{P}_{p,\lambda}$ -a.s.

3.2 Continuity at p = 1

We now employ a similar renormalization argument to prove Theorem 2. The sequence of scales we choose is still (8), but now we consider events

$$A_k(x) = \{ B(x, L_k) \text{ is connected to } \partial B(x, 10L_k) \text{ by edges in } \mathcal{O} \}$$
$$=: \{ B(x, L_k) \leftrightarrow \partial B(x, 10L_k) \}.$$

Once again, it is straightforward to check that events $A_k(x)$ satisfy properties **P1** and **P2**.

Proof of Theorem 2. Fix $\lambda < p_c^{\text{bond}}(d)$. We first need to complete the trigger for events $A_k(x)$. We can assume without loss of generality that $p \in [1/2, 1]$ and notice that we can adapt the argument in Lemma 4 to get k_0 depending only on d, since the error term in (11) is decreasing in p as can be shown by using the fact that $\alpha(p) := -\log(1-p) > 0$, as given by Lemma 1. Denoting by

V = V(d,k) the set of vertices of $B(o, 10L_k)$ and by \mathbb{P}_{λ}^{b} the independent bond percolation measure on \mathbb{L}^{d} , we can write

$$q_k(p,\lambda,d) \leq \mathbb{P}_{p,\lambda}(\xi(V) \neq 1) + \mathbb{P}_{p,\lambda}(\xi(V) \equiv 1, B(o, L_k) \leftrightarrow \partial B(o, 10L_k))$$
$$= (1 - p^{|V|}) + p^{|V|} \cdot \mathbb{P}_{\lambda}^{\mathrm{b}}(B(o, L_k) \leftrightarrow \partial B(o, 10L_k)).$$

Since $\lambda < p_c^{\text{bond}}(d)$ we have exponential decay for the radius of an open cluster, cf. [5, Theorem 5.4]. Thus, there is a positive constant $\psi(\lambda)$ such that

$$\mathbb{P}^{\mathrm{b}}_{\lambda}\big(B(o, L_k) \leftrightarrow \partial B(o, 10L_k)\big) \leq |\partial B(o, L_k)| \cdot \mathbb{P}^{\mathrm{b}}_{\lambda}\big(0 \leftrightarrow \partial B(o, 9L_k)\big) \leq c(d)L_k^{d-1} \cdot \exp[-\psi(\lambda)L_k].$$

Let us now pick $\tilde{k}_0(\lambda, d) > 0$ such that

$$c(d)L_k^{d-1} \cdot \exp[-\psi(\lambda)L_k] \le \frac{1}{2}L_k^{-2d} \quad \text{for } k \ge \tilde{k}_0(\lambda, d).$$

Taking $k_1(\lambda, d) = \max\{\tilde{k}_0(\lambda, d), k_0(d)\}$, we have

$$q_{k_1}(p,\lambda,d) \le \left(1 - p^{|V(k_1)|}\right) + \frac{1}{2}L_{k_1}^{-2d}.$$
(16)

Since λ and d are fixed we can pick $p_0 = p_0(\lambda, d)$ such that $1 - p^{|V(k_1)|} \leq \frac{1}{2}L_{k_1}^{-2d}$, for every $p \geq p_0(\lambda, d)$. Plugging into (16) we get

$$q_{k_1}(p,\lambda,d) \le L_{k_1}^{-2d} \quad \text{for } p \ge p_0(\lambda,d), \tag{17}$$

concluding the trigger step. This implies that there is no percolation for these values of p, λ and d. In fact, from (17) and Lemma 4 we get we get $q_k(p, \lambda, d) \leq L_k^{-2d}$ whenever $k \geq k_1$ and since $\theta(p, \lambda, d) \leq \lim_{k \to \infty} q_k(p, \lambda, d) = 0$ we have

$$\theta(p,\lambda,d) = 0 \quad \text{for } p \ge p_0,$$
(18)

that implies (2). In other words, this means that for any $\lambda < p_c^{\text{bond}}(d)$ the critical curve restricted to the interval $[p_0(\lambda, d), 1]$ must be above the horizontal segment of height λ .

On the other hand, a consequence of [2, Theorem 2.4 (*iii*)] is that for $\lambda > p_c^{\text{bond}}(d)$ the critical curve must be below λ on some interval $[\tilde{p}_0(\lambda, d), 1]$, since the curve giving the upper bound near p = 1 is continuous, recall Figure 1. The continuity of $\lambda_c(\cdot, d)$ at p = 1 follows.

References

- R. Baldasso, A. Teixeira (2018) How can a clairvoyant particle escape the exclusion process?, Annales de l'Institut Henri Poincaré, Probabilités et Statistiques.
- [2] N. Beaton, G. Grimmett, M. Holmes (2019) Alignment percolation, preprint arXiv:1908.07203.
- [3] O. Blondel, M.R. Hilário, A. Teixeira (2018) Random walks on dynamical random environment with non-uniform mixing, arXiv:1805.09750. To appear in The Annals of Probability.
- [4] R.M. Burton, M. Keane (1989) Density and uniqueness in percolation, Comm. Math. Phys. 121, no. 3, 501–505.
- [5] G. Grimmett (1999) *Percolation*, Die Grundlehren der mathematischen Wissenschaften in Einzeldarstellungen, Springer.
- [6] G. Grimmett (2006) *The Random-Cluster Model*, Grundlehren der mathematischen Wissenschaften, Springer Berlin Heidelberg.

- [7] M.R. Hilário, V. Sidoravicius (2019) Bernoulli line percolation, Stochastic Processes and their Applications, 129 (12), pp. 5037-5072.
- [8] T. Liggett (2012) Interacting particle systems, Springer Science & Business Media
- [9] A.-S. Sznitman (2010) Vacant set of random interlacements and percolation, Annals of Mathematics, volume 171, Issue 3, pp. 2039–2087.
- [10] J. Tykesson, D. Windisch (2012) Percolation in the vacant set of Poisson cylinders, Probability theory and related fields, 154(1-2), pp. 165–191.