

Totally umbilic surfaces in hyperbolic 3-manifolds of finite volume

Colin Adams

William H. Meeks III

Alvaro K. Ramos.*

July 10, 2020

Abstract

We construct for every connected surface S of finite negative Euler characteristic and every $H \in [0, 1)$, a hyperbolic 3-manifold $N(S, H)$ of finite volume and a proper, two-sided, totally umbilic embedding $f: S \rightarrow N(S, H)$ with mean curvature H . Conversely, we prove that a complete, totally umbilic surface with mean curvature $H \in [0, 1)$ embedded in a hyperbolic 3-manifold of finite volume must be proper and have finite negative Euler characteristic.

Mathematics Subject Classification: Primary 53A10, Secondary 49Q05, 53C42.

Key words and phrases: Calabi-Yau problem, hyperbolic 3-manifolds of finite volume, totally umbilic surfaces.

1 Introduction.

In this manuscript we develop the theory of totally umbilic surfaces in hyperbolic 3-manifolds of finite volume; all spaces considered here are assumed to be complete and connected and all surfaces in them will be assumed to be complete, connected and embedded, unless otherwise stated.

Theorem 1.1. *Let Σ be a totally umbilic surface with mean curvature $H_\Sigma \geq 0$ in a hyperbolic 3-manifold N of finite volume. Then:*

1. Σ is proper in N .
2. Σ has positive Euler characteristic if and only if Σ is a geodesic sphere. In particular, Σ is not diffeomorphic to a plane or a projective plane.
3. Σ has zero Euler characteristic if and only if N is non-compact and $H_\Sigma = 1$. In this case, Σ is a flat torus or a flat Klein bottle that is contained in some cusp end of N .
4. Σ has negative Euler characteristic if and only if it has finite negative Euler characteristic if and only if $H_\Sigma \in [0, 1)$. Furthermore:

*The second and third authors were partially supported by CNPq - Brazil, grant no. 400966/2014-0.

- (a) Σ has finite area $A(\Sigma) = \frac{2\pi}{H_\Sigma^2 - 1} \chi(\Sigma)$, where $\chi(\Sigma)$ is the Euler characteristic of Σ .
- (b) If $H_\Sigma > 0$, then, for every $H \in (0, H_\Sigma)$, there is a totally umbilic surface with mean curvature H in the ambient isotopy class of Σ .
- (c) If $H_\Sigma > 0$, then there is a totally geodesic surface Σ_0 in the isotopy class of Σ ; also, Σ is diffeomorphic to Σ_0 if Σ_0 is two-sided and Σ is diffeomorphic to the two-sided cover of Σ_0 if Σ_0 is one-sided.

The next theorem characterizes the admissible topological types of totally umbilic surfaces in hyperbolic 3-manifolds of finite volume with mean curvature in $[0, 1)$. It is a direct consequence of Theorem 1.1 and Theorem 5.1.

Theorem 1.2 (Admissible Topology Theorem). *A surface S appears topologically as a totally umbilic surface with mean curvature $H \in [0, 1)$ in some hyperbolic 3-manifold of finite volume if and only if S has finite negative Euler characteristic.*

Our construction of a hyperbolic 3-manifold of finite volume with a given admissible, two-sided, totally geodesic surface depends on the Switch Move Theorem [2, Theorem 4.1] and the Switch Move Gluing Operation [2, Theorem 5.1] from our previous study of modifications of hyperbolic 3-manifolds that are link complements; see Theorem 3.1 for the statement of the Switch Move Theorem. More specifically, for each admissible surface S , we apply these theorems to construct a finite volume hyperbolic 3-manifold $N(S)$ with an order-2 isometry whose fixed point set is two-sided and contains a component Σ diffeomorphic to S ; see Theorem 4.1 for additional topological properties satisfied by $N(S)$ and Σ . In Section 5, we apply geometric arguments to prove that for any $T > 0$, there is a finite cover $N_T(S)$ of $N(S)$ together with a lift $\tilde{\Sigma}$ of Σ , so that one of the two T -parallel surfaces to $\tilde{\Sigma}$ in $N_T(S)$ is a properly embedded, totally umbilic surface diffeomorphic to S with mean curvature $\tanh(T)$; crucial in these arguments is the property that the fundamental group of a hyperbolic 3-manifold of finite volume is LERF (see Definition 5.3).

2 The proof of Theorem 1.1.

In this section, we explain why a totally umbilic surface in a hyperbolic 3-manifold of finite volume must be proper and then show how Theorem 1.1 follows from this properness property. In order to carry out these proofs, we need the following definition, which will also be used in Section 5 to construct admissible totally umbilic surfaces that are parallel to totally geodesic ones.

Definition 2.1. Let N be a Riemannian 3-manifold and $f: S \rightarrow N$ be a two-sided embedding with image Σ and unitary normal vector field η . We define, for $t > 0$, the t -parallel surface to Σ as the image Σ_t of the immersion

$$f_t: S \rightarrow N, \quad x \mapsto \exp(t\eta(f(x))).$$

Thus,

$$\Sigma_t = \{\exp(t\eta(p)) \mid p \in \Sigma\}.$$

Proof of Theorem 1.1. We first prove item 1 of the theorem. Since it is well-known that totally umbilic surfaces with mean curvature $H \geq 1$ in a hyperbolic 3-manifold of finite volume N are either flat tori or Klein bottles of constant mean curvature 1 in cusp ends of N or geodesic spheres, item 1 holds for $H \geq 1$.

Next, we consider the totally geodesic case. For a given manifold M , let $\widehat{T}(M)$ denote the bundle of unoriented tangent two-planes of M . Then, the next result follows from the work of Shah [10] (also see Ratner [9] and Payne [8]). The phrase “immersed surface $f(S)$ ” in a 3-manifold N is used to indicate that the image surface $f(S)$ of an immersion $f: S \rightarrow N$ may have points of self-intersection.

Theorem 2.2. *Let $f: S \rightarrow N$ be a complete, totally geodesic immersion of a surface S to a hyperbolic 3-manifold N of finite volume. Then, either $f(S)$ is a properly immersed surface of finite area or $f_*(\widehat{T}(S))$ is dense in $\widehat{T}(N)$.*

In fact, Theorem 2.2 can be seen to hold by the following discussion. Let $f: S \rightarrow N$ be as stated and assume that S is endowed with a hyperbolic metric. Then, both S and N are examples of locally symmetric spaces of rank one. By the last statement of [8, Theorem 1.1], the closure of the image $f(S)$ is a totally geodesic submanifold of N , which in the context of this theorem means $f(S)$ is either proper or it is dense in N . If $f(S)$ is not proper, then, by [10, Theorem D], $f_*(\widehat{T}(S))$ is dense in $\widehat{T}(N)$ and the theorem holds.

Let Σ be a totally geodesic surface in a hyperbolic 3-manifold N of finite volume. Observe that the closure of Σ in N is a minimal lamination of N of class C^α , for all $\alpha \in (0, 1)$. Therefore, $\widehat{T}(\Sigma)$ is not dense in $\widehat{T}(N)$ and so Theorem 2.2 implies that Σ must have finite area and be proper in N .

To finish the proof of item 1, let Σ be a totally umbilic surface in N with mean curvature $H \in (0, 1)$. Assume Σ is oriented with respect to a unit normal field η pointing towards its mean convex side. Let $T = \tanh^{-1}(H) > 0$ and consider the T -parallel immersion $f_T: \Sigma \rightarrow N$ with immersed image surface Σ_T . Then, Σ_T is a complete totally geodesic immersed surface.

We claim that Σ_T is proper. Otherwise, Σ_T has infinite area and Theorem 2.2 implies that $\widehat{T}(\Sigma_T)$ is dense in $\widehat{T}(N)$. Since Σ_T has bounded norm of its second fundamental form, Σ_T intersects itself transversely in a dense set of points in N . Let $\Pi: \mathbb{H}^3 \rightarrow N$ denote the universal Riemannian covering map and let $\widetilde{\Sigma}_T^1, \widetilde{\Sigma}_T^2 \subset \mathbb{H}^3$ be two components of $\Pi^{-1}(\Sigma_T)$ corresponding to two lifts of Σ_T that intersect transversely in \mathbb{H}^3 . Let C_1, C_2 be the respective boundary circles of $\widetilde{\Sigma}_T^1, \widetilde{\Sigma}_T^2$, intersecting transversely in the boundary sphere at infinity of \mathbb{H}^3 . Consider two respective lifts $\widetilde{\Sigma}_1, \widetilde{\Sigma}_2$ of Σ in \mathbb{H}^3 with the same circles C_1, C_2 at infinity. Then, $\widetilde{\Sigma}_1$ intersects $\widetilde{\Sigma}_2$ transversely along a proper arc, which implies that Σ is not embedded, a contradiction. It also follows from this argument that Σ_T must be a proper totally geodesic (embedded) surface in N .

To show that Σ is proper, there are two cases to consider. First, assume that Σ_T is two-sided and oriented with respect to the unitary normal field $\widehat{\eta}$ corresponding to the opposite orientation from the induced immersion f_T . Then, as Σ is connected, it is the image of the T -parallel immersion $\widehat{f}_T: \Sigma_T \rightarrow N$, which must be a proper map, since the inclusion map of Σ_T in N is proper and the distance between any two corresponding points $x \in \Sigma_T, \widehat{f}_T(x) \in \Sigma$ is bounded by T . On the other hand, if Σ_T is one-sided, we may pass to the (proper) two-sheeted, two-sided cover of Σ_T

and repeat the same argument, finishing the proof of item 1 of Theorem 1.1.

As already observed, a surface Σ appears as a totally umbilic surface with mean curvature $H_\Sigma \geq 1$ in a hyperbolic manifold N of finite volume if and only if Σ is a geodesic sphere with $H_\Sigma > 1$ or it is a flat torus or a flat Klein bottle in a cusp end of N when $H_\Sigma = 1$. On the other hand, if Σ is a totally umbilic surface with $H_\Sigma \in [0, 1)$, which must be proper by item 1, then Corollary 4.7 of [7] implies that the Euler characteristic of Σ is negative, completing the proof of items 2 and 3.

Note that the main statement of item 4 follows from the above discussion and the fact that any properly immersed, infinite topology surface with constant mean curvature $H \in [0, 1)$ in a hyperbolic 3-manifold of finite volume has unbounded norm of its second fundamental form; see item 4 of [7, Theorem 1.3] for this unboundedness property.

The other statements of item 4 will be explained next. Suppose that Σ is a totally umbilic surface in a hyperbolic 3-manifold N of finite volume, with finite negative Euler characteristic and $H_\Sigma \in [0, 1)$. Then, item 4a follows immediately from [7, Corollary 4.7].

Next, we prove items 4b and 4c by showing that for any $t \in (0, T)$, the t -parallel immersion $f_t: \Sigma \rightarrow N$ is injective, where $T = \tanh^{-1}(H_\Sigma)$. Recall, from the proof of item 1, that the image surface $\Sigma_T = f_T(\Sigma)$ is a totally geodesic (embedded) surface in N . Let $\tilde{\Sigma}_T \subset \mathbb{H}^3$ be a component of $\Pi^{-1}(\Sigma_T)$ and let $\tilde{\Sigma}$ be a component of $\Pi^{-1}(\Sigma)$ with the same boundary circle at infinity as $\tilde{\Sigma}_T$. Let $W \subset \mathbb{H}^3$ be the closed region with boundary $\tilde{\Sigma} \cup \tilde{\Sigma}_T$. Since both Σ and Σ_T are embedded, if σ is a covering transformation of Π , then either σ maps W to itself, in which case σ leaves invariant each surface in W parallel to $\tilde{\Sigma}$, or $\sigma(W) \cap W = \tilde{\Sigma}_T$, in which case σ is a glide reflection along $\tilde{\Sigma}_T$ or a loxodromic transformation of \mathbb{H}^3 with respect to a geodesic γ in $\tilde{\Sigma}_T$ that has order-two rotational part about γ , or $\sigma(W) \cap W = \emptyset$. It follows that f_t is injective for all $t \in (0, T)$ and f_T is injective if and only if Σ_T is two-sided. In the case where Σ_T is one-sided, then the induced immersion $f_T: \Sigma \rightarrow \Sigma_T$ is a double covering of Σ_T , and items 4b and 4c follow. \square

3 Results on hyperbolic link complements.

In this section, we present some results concerning the hyperbolicity of link complements in 3-manifolds that will be used to construct the totally geodesic examples described in Theorem 4.1.

Theorem 3.1 (Switch Move Theorem [2, Theorem 4.1]). *Let L be a link in a 3-manifold M such that $M \setminus L$ admits a complete hyperbolic metric of finite volume. Let $\alpha \subset M$ be a compact arc which intersects L transversely in its two distinct endpoints, and such that $\text{int}(\alpha)$ is a properly embedded geodesic of $M \setminus L$. Let \mathcal{B} be a closed ball in M containing α in its interior and such that $\mathcal{B} \cap L$ is composed of two arcs in L , as in Figure 1(a). Let L' be the resulting link in M obtained by replacing $L \cap \mathcal{B}$ by the components as appearing in Figure 1 (b). Then $M \setminus L'$ admits a complete hyperbolic metric of finite volume.*

A consequence of the Switch Move Theorem is the following Untwisted Chain Theorem.




Figure 1: The Switch Move replaces the arcs g and g' by the tangle $\gamma_1 \cup \gamma_2 \cup C$.




Figure 2: For any positive integer k , replacing the trivial component C in (a) with the untwisted chain with $2k + 1$ components as in (b) preserves hyperbolicity of the complement.




Figure 3: The Untwisted Chain Theorem with $k = 1$ can be repeated in the highlighted subball to obtain any odd number of components.

Corollary 3.2 (Untwisted Chain Theorem). *Let L be a link in a 3-manifold M such that the link complement $M \setminus L$ admits a complete hyperbolic metric of finite volume. Suppose that there is a sphere S in M bounding a ball \mathcal{B} that intersects L as in Figure 2 (a). For any positive integer k , let L' be the resulting link in M obtained by replacing $L \cap \mathcal{B}$ by the untwisted chain with $2k + 1$ components as in Figure 2 (b). Then $M \setminus L'$ admits a complete hyperbolic metric of finite volume.*

Proof. We first prove Corollary 3.2 when $k = 1$. Let L , \mathcal{B} and M be as stated and let $D \subset \mathcal{B}$ be a two-punctured disk in $\mathcal{B} \setminus L$ with $\partial D = C$, where C is the circle component of $\mathcal{B} \cap L$. Since D is an incompressible three-punctured sphere in the hyperbolic 3-manifold $M \setminus L$, then [1, Theorem 3.1] gives that, up to isotopy, D is totally geodesic. Hence, there exists a compact arc $\alpha \subset \bar{D}$ in the closure of D , such that its interior $\tilde{\alpha}$ is a proper geodesic in the hyperbolic metric of $M \setminus L$. Moreover, α is transverse to L with the endpoints of α contained on C and α separates the punctures of D . After applying the Switch Move Theorem in a neighborhood of α , we obtain a link L' as in Figure 3, such that $M \setminus L'$ admits a complete, hyperbolic metric of finite volume. The general case follows by induction on k , as indicated in Figure 3. \square

Theorem 3.3 (Theorem 1.1 of [3]). *Let S be a closed (possibly non-orientable) surface with $\chi(S) \leq 0$. Then, there exists a link L in $S \times (0, 1)$ such that:*

- a. *If $\chi(S) < 0$, $(S \times [0, 1]) \setminus L$ admits a complete hyperbolic metric of finite volume with totally geodesic boundary.*
- b. *If $\chi(S) = 0$, $(S \times (0, 1)) \setminus L$ admits a complete hyperbolic metric of finite volume.*

Remark 3.4. In fact, the link L given by Theorem 3.3 can be any *fully alternating link*, see [3]. A fully alternating link L in $S \times (0, 1)$ is a link that admits a projection to S that is *alternating* in the

sense that the link can be oriented so that two consecutive crossings have distinct over/under signs and is *full* in the sense that every component of the complement of the projected image of L on S is a disk.

We also note that Theorem 3.3 is proved in [6] in the case where S is orientable, without obtaining a totally geodesic boundary when $\chi(S) < 0$. The proof in [6] could be extended to show these additional facts and it uses different techniques than those applied in [3].

4 Construction of hyperbolic 3-manifolds with totally geodesic surfaces.

Theorem 4.1. *Let S be a surface with finite negative Euler characteristic. There exists a finite volume hyperbolic 3-manifold N and a proper, two-sided embedding $f: S \rightarrow N$ with totally geodesic image Σ . Moreover:*

1. *If S is closed (resp. orientable), N is closed (resp. orientable).*
2. *If e_1 and e_2 are distinct ends of Σ , then N contains disjoint cusp ends C_1, C_2 such that, for $i = 1, 2$, $\Sigma \cap C_i$ is an annular representative of e_i .*
3. *Σ is a two-sided component of the fixed point set of an order-two isometry of N .*

Proof. Let S be as stated. To prove Theorem 4.1, we construct a complete hyperbolic 3-manifold of finite volume N together with an order-two diffeomorphism φ that has a two-sided fixed point set containing a component Σ diffeomorphic to S . By the Mostow rigidity theorem, after changing coordinates by a diffeomorphism isotopic to the identity, we may assume that φ is an isometry of N , from where it follows that Σ is totally geodesic and item 3 holds.

The proof breaks up into cases which are treated separately.

Case 4.2. *S is an n -punctured sphere with $n \geq 3$.*

Proof. For $n \geq 3$, let $L_n \subset \mathbb{R}^3$ be the $2n$ -component untwisted chain link; the L_4 version appears in Figure 4. Then, by adding to \mathbb{R}^3 the point at infinity and considering $L_n \subset \mathbb{S}^3$, $\mathbb{S}^3 \setminus L_n$ has an explicit hyperbolic metric of finite volume, as described in Example 6.8.7 of [11].

Thinking of the compactified xy -plane as a sphere $\mathcal{S} \subset \mathbb{S}^3$, we can view every other component of L_n as being contained in \mathcal{S} with the remaining components perpendicular to \mathcal{S} and symmetric with respect to reflection R through \mathcal{S} . Then, the restriction of R to the link complement $\mathbb{S}^3 \setminus L_n$ is an order-two isometry of the hyperbolic metric described above, with two-sided fixed point set $\mathcal{S} \setminus L_n$. This fixed point set contains an n -punctured sphere S_n , where the n punctures come from the n components of L_n in \mathcal{S} . It follows directly that S_n is totally geodesic and satisfies the statements of the theorem. \square

Case 4.3. *S is an n -punctured projective plane with $n \geq 2$.*




Figure 4: The untwisted chain link L_4 .

Proof. Fix $n \geq 2$ and let $k = n - 1$. Let L_{2k} be the $4k$ -component untwisted chain link in \mathbb{S}^3 . As in Case 4.2, \mathcal{S} denotes the compactification of the xy -plane and we assume that every other component of L_{2k} is contained in \mathcal{S} . Furthermore, we assume that the $2k$ components of L_{2k} that are not contained in \mathcal{S} are perpendicular to \mathcal{S} and lie on the unit sphere \mathbb{S}^2 centered at the origin of \mathbb{R}^3 and that L_{2k} is invariant under the inversion $\phi(\mathbf{x}) = -\mathbf{x}$ through the origin and the reflection Φ through \mathbb{S}^2 .

Let $Z \subset \mathbb{S}^3$ denote the z -axis with the point at infinity. Define $J_k = L_{2k} \cup Z$ and let $M = \mathbb{S}^3 \setminus J_k$. Then, M is a $2k$ -cover of $\mathbb{S}^3 \setminus B_3$, where B_3 is the Borromean ring with three components. Since $\mathbb{S}^3 \setminus B_3$ admits a complete hyperbolic metric of finite volume (see [11, Section 3.4]), there exists a complete hyperbolic metric g on M . Moreover, ϕ and Φ restrict to isometries of (M, g) .

Note that $\mathbb{S}^2 \setminus J_k$ contains a connected component \tilde{S} which is a $(2k + 2)$ -punctured sphere, where $2k$ punctures come from L_{2k} and the other two from $\mathbb{S}^2 \cap Z$. By construction, $\Phi|_{\tilde{S}} = \text{Id}_{\tilde{S}}$, and so \tilde{S} is totally geodesic.

Let $N = M/\phi$. Since ϕ is a fixed-point free, orientation reversing order-two isometry of (M, g) , N is a non-orientable hyperbolic manifold that is double covered by M . Since $\phi|_{\mathbb{S}^2}$ is the antipodal map and $\phi(\tilde{S}) = \tilde{S}$, the surface $S = \tilde{S}/\phi$ is a $(k + 1)$ -punctured projective plane in N . Also, since Φ and ϕ commute with each other, the map Φ descends to N as an order-two isometry of N which contains S in its fixed point set. Since $k + 1 = n$ and S satisfies the properties stated by Theorem 4.1, this proves Case 4.3. \square

Case 4.4. S is closed.

Proof. For the following construction, see Figure 5 (a). Consider \mathbb{S}^1 to be the unit circle in the yz -plane and let $P = S \times \mathbb{S}^1$. Let $\mathbb{S}_+^1 = \mathbb{S}^1 \cap \{z \geq 0\}$, $\mathbb{S}_-^1 = \mathbb{S}^1 \cap \{z \leq 0\}$, $M_1 = S \times \mathbb{S}_+^1$ and $M_2 = S \times \mathbb{S}_-^1$. Then, M_1, M_2 are subsets of P glued along their boundary surfaces $S_1 = S \times \{(-1, 0)\}$ and $S_2 = S \times \{(1, 0)\}$. Let $R: P \rightarrow P$ be the reflective symmetry interchanging M_1 with M_2 ; the fixed point set of R is $S_1 \cup S_2$.

By Theorem 3.3, there exists a link L in $\text{int}(M_1)$ such that $M_1 \setminus L$ admits a finite volume hyperbolic metric with totally geodesic boundary $S_1 \cup S_2$. Let $L' = R(L) \subset \text{int}(M_2)$ and $\Gamma =$




Figure 5: (a) is the manifold $P = S \times \mathbb{S}^1$ with the link L and its reflection $R(L)$. (b) is the manifold $P = S' \times \mathbb{S}^1$ with the link L_1 that creates a puncture to S_1 .

$L \cup L'$. Then, $P \setminus \Gamma$ admits a complete hyperbolic metric g for which R is an isometry and the surfaces S_1, S_2 are totally geodesic surfaces forming the fixed point set of R .

Note that $P \setminus \Gamma$ is orientable if and only if S is orientable. Moreover, after performing an appropriate Dehn filling in the ends of $P \setminus \Gamma$ in a symmetric manner with respect to R , we obtain a closed hyperbolic 3-manifold N , where S_1 and S_2 are each as stated in Theorem 4.1. \square

Case 4.5. S is S' punctured one time, where S' is a closed surface with $\chi(S') < 0$.

Proof. The starting point to this case is the closed manifold $P = S' \times \mathbb{S}^1$. As in Case 4.4, there exists a link $\Gamma = L \cup R(L)$, where L is a link in the interior of $M_1 = S' \times \mathbb{S}_+^1$, such that $P \setminus \Gamma$ is hyperbolic, the reflection R restricts to an isometry and the fixed point set of R consists of the two totally geodesic surfaces $S_1 = S' \times \{(-1, 0)\}$ and $S_2 = S' \times \{(1, 0)\}$.

In the hyperbolic metric of $M_1 \setminus L$, let α_1 be a minimizing geodesic ray from S_1 to L . Then, α_1 is proper, perpendicular to S_1 and α_1 does not intersect S_2 . Let $\alpha = \alpha_1 \cup R(\alpha_1)$. Then, α is a complete geodesic of $P \setminus \Gamma$ and the closure of α in P admits a neighborhood $\mathcal{B} \subset P$ with the following properties: $R(\mathcal{B}) = \mathcal{B}$, $\mathcal{B} \cap S_1 = E$ is a disk, $\mathcal{B} \cap S_2 = \emptyset$ and \mathcal{B} intersects Γ in two arcs $g \subset L$ and $g' = R(g) \subset R(L)$. Then, \mathcal{B} satisfies the hypothesis of Theorem 3.1. Then, we can replace the arcs $g \cup g'$ in $\mathcal{B} \cap \Gamma$ by a tangle $\gamma_1 \cup \gamma_2 \cup C$ as in Figure 1 (b) to form a new link $L_1 \subset P$, where we may choose C as a circle in $\text{int}(E)$ that bounds a disk in E , punctured once by α (see Figure 5 (b)). Then, Theorem 3.1 gives that $P \setminus L_1$ admits a complete, hyperbolic metric of finite volume, and since we may choose the arcs γ_1, γ_2 to be invariant under R , it follows that R restricts to an isometry φ . Note that the fixed point set of φ contains three connected components, one being S_2 and the other two being the connected components of $S_1 \setminus (C \cup \gamma_1 \cup \gamma_2)$, one a

three-punctured sphere and the other diffeomorphic to S and satisfying the conclusions required by Theorem 4.1. \square

Case 4.6. S is a torus or a Klein bottle punctured once.

Proof. Let T be either a torus or a Klein bottle and let P_1 be the product manifold $T \times [0, 1]$. Let Γ_1 be a link in $\text{int}(P_1)$ such that $\text{int}(P_1 \setminus \Gamma_1)$ is hyperbolic, as given by item b of Theorem 3.3. Let $P = T \times [-1, 1]$ and $R: P \rightarrow P$ be the reflection $R(x, t) = (x, -t)$; take P_2 as the reflected image of P_1 in P , with respective hyperbolic link $\Gamma_2 = R(\Gamma_1) \subset P_2$. Also, let α_1 be a complete geodesic in the hyperbolic metric of $\text{int}(P_1 \setminus \Gamma_1)$ with one endpoint in $T \times \{0\}$ and another endpoint in a component L_1 of Γ_1 and let $\alpha_2 = R(\alpha_1)$.

Let $\alpha \subset P$ be the concatenation of α_1 and α_2^{-1} . Then, α is an arc with one endpoint in L_1 and another endpoint in $L_2 = R(L_1)$. Let \mathcal{B} be a regular neighborhood of α in P , invariant under R and that intersects $\Gamma = \Gamma_1 \cup \Gamma_2$ in two arcs $g \subset L_1$ and $g' = R(g) \subset L_2$ and intersects T in a disk Δ . Let Γ' be the link in $\text{int}(P)$ obtained from Γ by replacing $g \cup g'$ in \mathcal{B} by the tangle $\gamma_1 \cup \gamma_2 \cup C$ as in Figure 1 (b), where $C \subset \Delta$. Then, Theorem 5.1 in [2] implies that the interior of the link complement $P \setminus \Gamma'$ admits a complete hyperbolic metric of finite volume. Furthermore, we may choose $\gamma_1 \cup \gamma_2 \cup C$ so that it is invariant under R . Hence, R restricts to an isometry φ of the hyperbolic metric of $P \setminus \Gamma'$. This proves Case 4.6, since the fixed point set of φ contains a three-punctured sphere D bounded by C in T and a component $T \setminus \overline{D}$ diffeomorphic to S that satisfies the properties required by Theorem 4.1. \square

The remaining cases to treat in Theorem 4.1 are those where S is diffeomorphic to S' punctured $n \geq 2$ times, where S' is a closed surface with $\chi(S') \leq 0$. If $\chi(S') < 0$, let $P = S' \times \mathbb{S}^1$ with respective hyperbolic link L_1 as given by Case 4.5. Let $\mathcal{B} \subset P$ be the ball where the Switch Move Theorem was applied. Then, in $P \setminus L_1$, \mathcal{B} satisfies the hypothesis of the Untwisted Chain Theorem, Corollary 3.2. Then, we can replace the circle component C of $\mathcal{B} \cap L_1$ by an untwisted chain with $2n - 1$ components as in Figure 2 (b), where every other component is in the interior of $D = \mathcal{B} \cap S_1$ and the remaining components lie in \mathcal{B} and are symmetric with respect to R to create a hyperbolic link $L_n \subset P$. Again, the reflection R restricts to an isometry φ of the hyperbolic metric of $P \setminus L_n$, and the fixed point set of φ contains S_2 , $n - 1$ three-punctured spheres and a surface diffeomorphic to S , which finishes the proof of Theorem 4.1 when $\chi(S') < 0$. The proof for the case when S' is a torus or a Klein bottle is analogous. \square

5 The Proof of Theorem 1.2.

To finish the proof of Theorem 1.2, it suffices to prove Theorem 5.1 below.

Theorem 5.1. *Let $H \in [0, 1)$ and suppose that S is a connected surface of finite topology and negative Euler characteristic. Then, there exists a hyperbolic 3-manifold N of finite volume and a proper, two-sided embedding $f: S \rightarrow N$, with image Σ having mean curvature H and satisfying:*

1. Σ is totally umbilic.

2. Σ does not separate N .
3. If S is closed (resp. orientable), then N can be chosen to be closed (resp. orientable).
4. Each end e of Σ admits an annular end representative E which is embedded in a cusp end \mathcal{C}_e of N . In addition, if e and f are two distinct ends of Σ , the respective cusp ends $\mathcal{C}_e, \mathcal{C}_f$ are distinct.

To prove Theorem 5.1, we make use of the totally geodesic examples provided by Theorem 4.1 to construct hyperbolic manifolds of finite volume with two-sided, totally umbilic surfaces which are properly embedded with any admissible finite topology and mean curvature $H \in (0, 1)$. In order to define these associated totally umbilic examples, we recall, from Definition 2.1, that if N is a 3-manifold and $f: S \rightarrow N$ is a two-sided embedding with image Σ and unitary normal vector field η , then the t -parallel surface to Σ is the image Σ_t of the immersion $f_t: S \rightarrow N$ defined by $f_t(x) = \exp(t\eta(f(x)))$. Note that the t -parallel surface Σ_t to a totally geodesic surface Σ in a hyperbolic 3-manifold N is totally umbilic and has mean curvature $\tanh(t) \in (0, 1)$. For convenience, we will often assume that the domain of the immersion f_t is Σ instead of the abstract surface S . Note that the ambient distance of two respective points $x \in \Sigma$ and $f_t(x) \in \Sigma_t$ is uniformly bounded by t .

Lemma 5.2. *Let $f: S \rightarrow N$ satisfy the properties given by Theorem 4.1, and assume that N has $m \geq 0$ ends. Then, for any $T > 0$:*

1. *There exists a pairwise disjoint collection of cusp end representatives of the ends of N , $\{\mathcal{C}_1(T), \dots, \mathcal{C}_m(T)\}$, such that for any $t \in (0, T]$, f_t is injective on $f^{-1}(\cup_{i=1}^m \mathcal{C}_i(T))$.*
2. *For $t \in (0, T]$ sufficiently small, the immersions $f_t: S \rightarrow N$ are injective.*
3. *If $\Lambda = \{t > 0 \mid f_t \text{ is not injective}\} \neq \emptyset$, then $t_0 = \inf \Lambda > 0$ and there exists a closed geodesic in N , meeting Σ orthogonally with length $4t_0$.*

Proof. First, we notice that if S is compact, the proofs of items 1 and 2 are immediate.

Assume now that S is noncompact and let e_1, \dots, e_n be the ends of $\Sigma = f(S)$. An immediate consequence of item 2 of Theorem 4.1 is that there exist pairwise disjoint annular end representatives $E_1, \dots, E_n \subset \Sigma$ and a collection $\{\mathcal{C}_1, \dots, \mathcal{C}_m\}$ of pairwise disjoint cusp ends of N such that, after possibly passing to subends, it holds, for $1 \leq i \leq n$, that $\Sigma \cap \mathcal{C}_i = E_i$ with $\partial \mathcal{C}_i$ intersecting Σ orthogonally and, for $n+1 \leq j \leq m$, $\mathcal{C}_j \cap \Sigma = \emptyset$.

For a given $i \in \{1, \dots, n\}$, consider the cusp end \mathcal{C}_i . There exists a family of compact surfaces $\{\mathcal{T}_i(s)\}_{s \geq 0}$ (if \mathcal{C}_i is orientable, each $\mathcal{T}_i(s)$ is a torus, otherwise each $\mathcal{T}_i(s)$ is a Klein bottle) arising from the descent of parallel horospheres of \mathbb{H}^3 via the universal covering projection, parameterized by $s = \text{dist}(\mathcal{T}_i(s), \mathcal{T}_i(0) = \partial \mathcal{C}_i)$ such that $\cup_{s \geq 0} \mathcal{T}_i(s) = \mathcal{C}_i$. For any $\lambda > 0$, let

$$\mathcal{C}_i(\lambda) = \cup_{s \geq \lambda} \mathcal{T}_i(s), \quad N(\lambda) = N \setminus (\cup_{i=1}^n \mathcal{C}_i(\lambda)), \quad (1)$$

and

$$E_i(\lambda) = E_i \cap \mathcal{C}_i(\lambda), \quad \Sigma(\lambda) = \Sigma \cap N(\lambda). \quad (2)$$

The assumption that $\Sigma \cap \mathcal{C}_i = E_i$, for $1 \leq i \leq n$, implies that $E_i(\lambda) = \Sigma \cap \mathcal{C}_i(\lambda)$. In particular, for every $\lambda > 0$, we can express Σ as a disjoint union $\Sigma = (\cup_{i=1}^n E_i(\lambda)) \cup \Sigma(\lambda)$.

Fix $T > 0$, $t \in (0, T]$ and $i \in \{1, 2, \dots, n\}$. Using the universal covering of \mathcal{C}_i by a horoball of \mathbb{H}^3 , it is easy to see that, for any $s > T$, E_i is orthogonal to $\mathcal{T}_i(s)$ and $E_i \cap \mathcal{T}_i(s)$ is a geodesic of $\mathcal{T}_i(s)$, which is injectively mapped by f_t to a geodesic of $\mathcal{T}_i(h_t(s))$, where $h_t(s) \in (s - t, s)$ and the function h_t is increasing; hence $f_t|_{E_i(T)}$ is injective. In order to finish the proof of item 1, just note that $f_t(E_i(T)) \subset \mathcal{C}_i$ and, if $i \neq j$, $\mathcal{C}_i \cap \mathcal{C}_j = \emptyset$.

Next, we prove the second statement of the lemma. Since $\overline{\Sigma(3T)}$ is compact, then there exists some $\varepsilon(T) \in (0, T)$ such that $f_t|_{\Sigma(3T)}$ is injective for all $t \in (0, \varepsilon(T))$. We claim that f_t is injective for all $t \in (0, \varepsilon(T))$.

Assume that $f_t(x) = f_t(y) = p$, for some $x, y \in \Sigma$ and $t \in (0, \varepsilon(T))$. If $x \neq y$, since $f_t|_{\Sigma(3T)}$ is injective and $\{E_1(3T), E_2(3T), \dots, E_n(3T), \Sigma(3T)\}$ is a partition of Σ , there exists some $i \in \{1, 2, \dots, n\}$ such that either $x \in E_i(3T)$ or $y \in E_i(3T)$. In particular, $p \in \mathcal{C}_i(2T)$. Without loss of generality, assume that $x \in E_i(3T)$. Item 1 and the fact that $t \in (0, T)$ gives that $f_t|_{E_i(T)}$ is injective. Moreover, $E_i(3T) \subset E_i(T)$, and therefore $y \notin E_i(T)$. There are two possibilities: either $y \in E_j(T)$ for $j \neq i$ or $y \in \Sigma(T)$. If $y \in E_j(T)$, then $p \in \mathcal{C}_j$, which is impossible since $\mathcal{C}_j \cap \mathcal{C}_i = \emptyset$. On the other hand, if $y \in \Sigma(T)$, then $p \in N(2T)$. However, $N(2T) \cap \mathcal{C}_i(2T) = \emptyset$, and this contradiction proves item 2.

We next prove item 3. First, item 2 gives that $t_0 > 0$, and our next argument shows that f_{t_0} is not injective.

Choose $\lambda > t_0$. By the definition of t_0 , there exist a sequence $\{t_k\}_{k \in \mathbb{N}} \subset [t_0, \lambda)$, $t_k \rightarrow t_0$ and points $x_k, y_k \in \Sigma$, $x_k \neq y_k$, such that $f_{t_k}(x_k) = f_{t_k}(y_k) = p_k$. From the definition of f_{t_k} and from the assumption that $t_k < \lambda$ for all $k \in \mathbb{N}$, the triangle inequality implies that $d_N(x_k, y_k) < 2\lambda$, for all $k \in \mathbb{N}$.

By item 1, it follows that $f_t|_{E_i(\lambda)}$ is injective for all $i \in \{1, \dots, n\}$ and all $t \in (0, \lambda)$. Therefore, $x_k, y_k \in \Sigma(3\lambda)$ for all $k \in \mathbb{N}$, since $x_k \in E_i(3\lambda) \subset E_i(\lambda)$ gives that $y_k \in E_i(\lambda)$ and vice versa. The compactness of $\overline{\Sigma(3\lambda)}$ implies that, up to subsequences, there are $x, y \in \overline{\Sigma(3\lambda)}$ such that $x_k \rightarrow x$ and $y_k \rightarrow y$. By the continuity of $(t, z) \mapsto f_t(z)$, it follows that $f_{t_0}(x) = f_{t_0}(y) = p$. Moreover, $\{f_t|_{\overline{\Sigma(3\lambda)}}\}_{t \in [0, \lambda]}$ is a smooth, compact family of immersions of the compact surface $\overline{\Sigma(3\lambda)}$ into N . Hence, there exists an $\varepsilon > 0$ such that for any $t \in [0, \lambda]$ and any $z \in \overline{\Sigma(3\lambda)}$, $f_t|_{B_{\Sigma(z, \varepsilon)}}$ is injective, which implies that $d_{\Sigma}(x_k, y_k) \geq \varepsilon$, thus $x \neq y$. This proves that f_{t_0} is not injective.

Let $U \ni x$ and $V \ni y$ be two disjoint open disks of Σ such that the restrictions $f_{t_0}|_U$ and $f_{t_0}|_V$ are injective. It follows that $f_{t_0}(U)$ and $f_{t_0}(V)$ are embedded disks intersecting at some point $p = f_{t_0}(x) = f_{t_0}(y)$. Note that the fact that f_t is injective for all $t \in (0, t_0)$ gives that the intersection of $f_t(U)$ and $f_t(V)$ is tangential at p . Consider the two oriented geodesic rays $\gamma_x = \{f_t(x)\}_{t \in [0, t_0]}$ and $\gamma_y = \{f_t(y)\}_{t \in [0, t_0]}$. Then the concatenation γ of γ_x with γ_y^{-1} is a smooth geodesic arc in N , with length $2t_0$ and meeting Σ orthogonally at the points x and y . Finally, if $\varphi: N \rightarrow N$ is the order-two isometry containing Σ in its fixed point set, then $\tilde{\gamma} = \gamma \cup \varphi(\gamma)$ is a closed geodesic with length $4t_0$ and meeting Σ orthogonally, proving item 3 of Lemma 5.2. \square

Lemma 5.2 gives properly embedded, totally umbilic surfaces for small values of H . In order to finish the proof of Theorem 5.1, we apply a technical result stating that the fundamental groups

of hyperbolic 3-manifolds of finite volume satisfy the following definition.

Definition 5.3 (Locally Extendable Residually Finite group). *A group G is called LERF if for every finitely generated subgroup K of G and any $g \notin K$, there exists a representation $\sigma: G \rightarrow F$ from G to a finite group F such that $\sigma(g) \notin \sigma(K)$.*

The above definition can be extended as follows. *A group G is LERF if and only if for every finitely generated subgroup K of G and any finite subset $\mathcal{F} = \{g_1, \dots, g_n\} \subset G$ such that $\mathcal{F} \cap K = \emptyset$, there exists a representation $\sigma: G \rightarrow F$ from G to a finite group F such that $\sigma(\mathcal{F}) \cap \sigma(K) = \emptyset$. Indeed, if G is LERF and \mathcal{F} and K are as above, for each $i \in \{1, \dots, n\}$, there exists a representation $\sigma_i: G \rightarrow F_i$ from G to a finite group F_i such that $\sigma_i(g_i) \notin \sigma_i(K)$. Let $\sigma = \sigma_1 \times \dots \times \sigma_n: G \rightarrow F = F_1 \times \dots \times F_n$, then σ is a representation as claimed. This equivalent definition will be used in the proof of Theorem 5.1.*

By a series of recent works (see [4] for a complete list of appropriate references) on group theoretical properties of fundamental groups of hyperbolic 3-manifolds, one has the following result.

Theorem 5.4. *If N is a hyperbolic 3-manifold of finite volume, then $\pi_1(N)$ is LERF.*

The case where N is orientable is treated in [4, Corollary 4.2.3]. However, from the discussion in the book [4], it was not clear to us if Theorem 5.4 applies to the non-orientable case. For this reason, we next explain how this property follows from the orientable case. Recall, from [4], that a group G is LERF if and only if any finitely generated subgroup K of G is closed in the profinite topology of G . Let N be a non-orientable hyperbolic 3-manifold of finite volume and let \tilde{N} be its oriented 2-sheeted cover. Then $\pi_1(\tilde{N})$ is LERF. Since $\pi_1(\tilde{N})$ can be viewed as a finitely generated, index-2 subgroup of $\pi_1(N)$, $\pi_1(\tilde{N})$ is closed in the profinite topology of $\pi_1(N)$. Let $K \subset \pi_1(N)$ be a finitely generated subgroup. There are two cases to consider: either $K \subset \pi_1(\tilde{N})$, in which case K is closed in $\pi_1(N)$ since $\pi_1(\tilde{N})$ is LERF and closed in $\pi_1(N)$, or $K \not\subset \pi_1(\tilde{N})$, and there exists some $a \in K$, $a \notin \pi_1(\tilde{N})$; it then follows that $K = \left(K \cap \pi_1(\tilde{N})\right) \cup a \left(K \cap \pi_1(\tilde{N})\right)$ is the union of two closed sets, thus K is closed in $\pi_1(N)$. Therefore, $\pi_1(N)$ is LERF.

With the above discussion in mind, we now continue with the proof of Theorem 5.1. Fix a surface S of finite topology, with negative Euler characteristic and $T > 0$. Our goal (precisely stated in Lemma 5.5 below) is to produce a hyperbolic 3-manifold N_T , together with a two-sided, proper embedding $\tilde{f}: S \rightarrow N_T$ with totally geodesic image $\tilde{\Sigma}$ such that, for each $t \in (0, T]$, the related parallel immersion $\tilde{f}_t: S \rightarrow N_T$ is injective. Since the image surface $\tilde{\Sigma}_t = \tilde{f}_t(S)$ is totally umbilic with mean curvature $H = \tanh(t)$, $\lim_{T \rightarrow \infty} \tanh(T) = 1$ and T is arbitrary, Lemma 5.5 proves Theorem 5.1.

By Theorem 4.1, there exists a hyperbolic 3-manifold of finite volume N , together with a two-sided proper embedding $f: S \rightarrow N$, with totally geodesic image Σ and an order-two isometry $\varphi: N \rightarrow N$ such that $\varphi|_{\Sigma} = \text{Id}_{\Sigma}$. Using this particular example, we prove next lemma.

Lemma 5.5. *For each $T > 0$, there exists a finite Riemannian covering space $\Pi_T: N_T \rightarrow N$ satisfying:*

1. The embedding $f: S \rightarrow N$ lifts to an embedding $\tilde{f}: S \rightarrow N_T$.
2. The order-two isometry $\varphi: N \rightarrow N$ lifts to an order-two isometry $\tilde{\varphi}: N_T \rightarrow N_T$ such that $\tilde{\varphi}|_{\tilde{\Sigma}} = \text{Id}_{\tilde{\Sigma}}$, where $\tilde{\Sigma} = f(S)$.
3. The t -parallel surfaces to $\tilde{\Sigma}$ are embedded for all $t \in (0, T]$.

Proof. Fix $T > 0$. Since N is a hyperbolic 3-manifold of finite volume, there are only a finite number of smooth closed geodesics in N with length less than $L = 5T$. In particular, the (possibly empty) collection \mathcal{G}_L of prime, smooth closed geodesics in N that are orthogonal to $\Sigma \subset N$ at some point and with length less than L is finite.

Note that if \mathcal{G}_L were empty, then Lemma 5.5 follows directly from item 3 of Lemma 5.2, by letting $N_T = N$, $\Pi_T = \text{Id}$, $f = f$ and $\tilde{\varphi} = \varphi$. Thus, we next assume that

$$\mathcal{G}_L = \{\gamma_1, \gamma_2, \dots, \gamma_{n(L)}\} \neq \emptyset.$$

For each $j \in \{1, \dots, n(L)\}$, let $p_j \in \Sigma \cap \gamma_j$ be a point where γ_j meets Σ orthogonally. The geodesic $\gamma_j \in \mathcal{G}_L$ is invariant under the isometry φ ; furthermore, $\varphi|_{\gamma_j}$ reverses the orientation of γ_j and fixes the point p_j .

Let $i: \Sigma \rightarrow N$ be the inclusion map. Choosing p_1 as a base point, we let $i_*: \pi_1(\Sigma, p_1) \rightarrow \pi_1(N, p_1)$ be the induced homomorphism on fundamental groups. We let

$$K_1 = i_*(\pi_1(\Sigma, p_1)) \subset \pi_1(N, p_1)$$

be the image of the finitely generated group $\pi_1(\Sigma, p_1)$. Since $p_1 \in \gamma_1$, γ_1 can be considered to represent a nontrivial element $[\gamma_1] \in \pi_1(N, p_1)$. Moreover, since Σ is totally geodesic in a hyperbolic 3-manifold, i_* is injective and, for any positive integer l , $[\gamma_1]^l \notin K_1$.

Fix a positive integer k sufficiently large so that

$$k \cdot \text{Length}(\gamma_1) \geq L \tag{3}$$

and let $\mathcal{F} = \{[\gamma_1], [\gamma_1]^2, \dots, [\gamma_1]^k\}$. Then, $\mathcal{F} \cap K_1 = \emptyset$ and, since $\pi_1(N, p_1)$ is LERF, there exists a representation $\sigma: \pi_1(N, p_1) \rightarrow F_1$, from $\pi_1(N, p_1)$ to a finite group F_1 , such that $\sigma(\mathcal{F}) \cap \sigma(K_1) = \emptyset$. Let $\widehat{K}_1, \widetilde{K}_1$ be the subgroups of $\pi_1(N, p_1)$ defined by

$$\widehat{K}_1 = \sigma^{-1}(\sigma(K_1)), \quad \widetilde{K}_1 = \widehat{K}_1 \cap \varphi_*(\widehat{K}_1).$$

Note that $K_1 \subset \widetilde{K}_1$, since $K_1 \subset \widehat{K}_1$ and φ_* fixes all elements of K_1 . Also, for $l \in \{1, 2, \dots, k\}$, $[\gamma_1]^l \notin \widehat{K}_1$, hence $[\gamma_1]^l \notin \widetilde{K}_1$.

Next we show that \widetilde{K}_1 has finite index in $\pi_1(N, p_1)$. Since $\widehat{K}_1 \supset \ker(\sigma)$ and F_1 is finite, the index of \widehat{K}_1 in $\pi_1(N, p_1)$ is finite. Moreover, $\varphi_*: \pi_1(N, p_1) \rightarrow \pi_1(N, p_1)$ is a group isomorphism; hence the index of $\varphi_*(\widehat{K}_1)$ in $\pi_1(N, p_1)$ is also finite. Then, since the intersection of two subgroups of finite index also has finite index, the claim follows.

Let $\Pi_1: (N_1, q_1) \rightarrow (N, p_1)$ be the Riemannian covering space of (N, p_1) with image subgroup $(\Pi_1)_*(\pi_1(N_1, q_1)) = \widetilde{K}_1$. Note that Π_1 is a finite covering, since the index of \widetilde{K}_1 in $\pi_1(N, p_1)$

is finite; in particular, N_1 is a hyperbolic manifold of finite volume and, if N is closed (resp. orientable), N_1 is also closed (resp. orientable).

Since $\varphi_*(\tilde{K}_1) = \tilde{K}_1$, then, by the lifting criterion, the maps $i: \Sigma \rightarrow N$, $\varphi: N \rightarrow N$ have respective lifts

$$i_1: (\Sigma, p_1) \rightarrow (N_1, q_1), \quad \varphi_1: (N_1, q_1) \rightarrow (N_1, q_1).$$

Let Σ_1 denote the embedded, totally geodesic image surface of the injective immersion i_1 , and note that Σ_1 is two-sided and contained in the fixed point set of the order-two isometry φ_1 .

Consider the (possibly empty) collection $\mathcal{G}_L^1 = \{\gamma_1^1, \gamma_2^1, \dots, \gamma_{n_1(L)}^1\}$ of smooth, prime closed geodesics in N_1 that have length less than L and are orthogonal to Σ_1 at some point. Then, Π_1 induces an injective map from \mathcal{G}_L^1 to \mathcal{G}_L , hence $n_1(L) \leq n(L)$. We next prove that this construction yields $n_1(L) < n(L)$.

Claim 5.6. *The image set of geodesics $\Pi_1(\mathcal{G}_L^1) = \{\Pi_1(\gamma_j^1) \mid j = 1, \dots, n_1(L)\}$ forms a subset of $\{\gamma_1, \gamma_2, \dots, \gamma_{n(L)}\}$, which does not include γ_1 . In particular, $n_1(L) < n(L)$.*

Proof. Arguing by contradiction, suppose, after possibly reordering, that $\gamma_1 = \Pi_1(\gamma_1^1)$ as a set. Then, γ_1^1 is the lift of a certain smallest power J of γ_1 , which implies that the length of γ_1^1 is equal to $J \cdot \text{Length}(\gamma_1)$. However, for $l \in \{1, \dots, k\}$, $[\gamma_1]^l \notin \tilde{K}_1$; hence, none of the powers of γ_1 less than or equal to k lift. Then $J > k$, and (3) implies that the length of γ_1^1 is larger than L , which is a contradiction. \square

By induction, Claim 5.6 allows us to produce a finite Riemannian cover $\Pi_T: N_T \rightarrow N$ such that:

1. The embedding $f: S \rightarrow N$ lifts to an embedding $\tilde{f}: S \rightarrow N_T$, with totally geodesic image surface $\tilde{\Sigma}$.
2. There are no smooth closed geodesics in N_T with length less than L and intersecting $\tilde{\Sigma}$ orthogonally at some point.
3. $\varphi: N \rightarrow N$ lifts to an order-two isometry $\tilde{\varphi}: N_T \rightarrow N_T$ and $\tilde{\varphi}|_{\tilde{\Sigma}} = \text{Id}_{\tilde{\Sigma}}$.

Note that \tilde{f} and $\tilde{\varphi}$ satisfy the hypothesis of Lemma 5.2. In particular, since $\tilde{\Sigma}$ was constructed in such a way that there are no smooth closed geodesics in N_T with length less than $L = 5T$ intersecting $\tilde{\Sigma}$ orthogonally, item 3 of Lemma 5.2 implies that the t -parallel surfaces to $\tilde{\Sigma}$ are embedded for all $t \in (0, T]$, which completes the proof of Lemma 5.5. \square

As already explained, Theorem 5.1 now follows from Theorem 4.1 and Lemma 5.5, since in the above construction the t -parallel surface $\tilde{\Sigma}_t = \tilde{f}_t(S)$ is a properly embedded surface in N_T satisfying the conditions 1, 2, 3 and 4 of Theorem 5.1 for $H = \tanh(t) \in (0, \tanh(T)]$.

References

- [1] C. Adams. Thrice-punctured spheres in hyperbolic 3-manifolds. *Trans. Amer. Math. Soc.*, 287:645–656, 1985.
- [2] C. Adams, W. H. Meeks III, and A. Ramos. Modifications preserving hyperbolicity of link complements. preprint at <http://arxiv.org/pdf/2007.01739>.
- [3] C. Adams, C. Albers Riera, B. Haddock, Z. Li, D. Nishida, and B. Reinoso. Hyperbolicity of links in thickened surfaces. Preprint at <http://arxiv.org/pdf/1802.05770>.
- [4] M. Aschenbrenner, S. Friedl, and H. Wilton. *3-manifold Groups*. EMS series of lectures in mathematics. European Mathematical Society, 2015. ISBN 9783037191545.
- [5] P. Collin, L. Hauswirth, and H. Rosenberg. Minimal surfaces in finite volume hyperbolic 3-manifolds and in $M \times S$, M a finite area hyperbolic surface. *Amer. J. Math.*, 140(4):1075–1112, 2018. DOI 10.1353/ajm.2018.0024.
- [6] J. Howie and J. Purcell. Geometry of alternating links on surfaces. *To appear in Trans. of the A. Math. Soc.* preprint at <https://arxiv.org/pdf/1712.01373.pdf>.
- [7] W. H. Meeks III and A. K. Ramos. Properly immersed surfaces in hyperbolic 3-manifolds. *J. Differential Geom.*, 112(2):233–261, 2019. DOI 10.4310/jdg/1559786424 <https://doi.org/10.4310/jdg/1559786424>, MR3960267.
- [8] T. Payne. Closures of totally geodesic immersions into locally symmetric spaces of noncompact type. *Proc. Amer. Math. Soc.*, 127(3):829–833, 1999. MR1468202, Zbl 0936.53034, DOI 10.1090/S0002-9939-99-04552-9.
- [9] M. Ratner. Raghunathan’s topological conjecture and distributions of unipotent flows. *Duke Math. J.*, 63(1):235–280, 1991. MR1106945, Zbl 0733.22007, DOI 10.1215/S0012-7094-91-06311-8.
- [10] N. Shah. Closures of totally geodesic immersions in manifolds of constant negative curvature. In *Group theory from a geometrical viewpoint (Trieste, 1990)*, pages 718–732. World Sci. Publ., River Edge, NJ, 1991. MR1170382, Zbl 0846.53041.
- [11] W. P. Thurston. *The geometry and topology of three manifolds*. Princeton Lecture Notes.

Colin Adams at cadams@williams.edu

Department of Mathematics and Statistics, Williams College, Williamstown, MA 01267

William H. Meeks, III at profmeeks@gmail.com

Mathematics Department, University of Massachusetts, Amherst, MA 01003

Álvaro K. Ramos at alvaro.ramos@ufrgs.br

Departamento de Matemática Pura e Aplicada, Universidade Federal do Rio Grande do Sul, Brazil