HYPERBOLIC SELF AVOIDING WALK

ITAI BENJAMINI AND CHRISTOFOROS PANAGIOTIS

ABSTRACT. Expected ballisticity of a continuous self avoiding walk on hyperbolic spaces \mathbb{H}^d is established.

1. Introduction

Consider continuous n steps random walk on \mathbb{H}^d , where the next step is chosen uniformly and independently on the unit sphere around the current location. Condition this uniform product measure on sequences in which the distance between any pair of vertices is bigger than c, for some fixed 0 < c < 1.

We will write \mathbb{P}_n for this measure, which will be called the SAW measure, and \mathbb{E}_n for the expectation with respect to this measure. We denote by x_0, x_1, \ldots, x_n the vertices of the walk in order of appearance and d the distance function.

Theorem 1.1. There exists a constant C = C(d, c) > 0 such that $\mathbb{E}_n d(x_0, x_n) > Cn$ for every $n \geq 1$.

For background on hyperbolic geometry see e.g. [3]. For background on self avoiding walks see [1]. The papers [5, 6, 7, 8] contains results of the speed of self avoiding walks.

2. Proof

Proof. Let us first recall that there is a constant δ fixed throughout such that for every geodesic triangle in \mathbb{H}^d with sides α, β, γ we have that α lies in the δ neighbourhood of $\beta \cup \gamma$.

We will start by proving the following geometric lemma.

Date: 25.3.2020.

Lemma 2.1. There is a constant $c_1 > 0$ such that for every finite set A of vertices in \mathbb{H}^d with the property that the distance between any two of its points is at least c, at least $c_1|A|$ vertices of A are at distance at most (1-c)/2 from the boundary of the convex hull of A.

Proof. Notice that the convex hull K of A coincides with the convex hull of $A \cap \partial K$. According to [2], there is a constant t = t(d) such that the volume of K is at most $t|A \cap \partial K|$. For every x in $B := \{x \in A \mid d(x, \partial K) > (1-c)/2\}$, the ball of radius (1-c)/2 around x is contained in K, hence the volume of K is greater than t'|B| for a certain constant t'>0. The assertion follows now easily.

For a set of vertices A, we will write $\mathcal{H}(A)$ for the convex hull of A. It follows from Lemma (2.1) that there is a constant R > 0 such that for every SAW x of length n, the number of indices i with $d(x_i, \partial \mathcal{H}(x)), d(x_j, \partial \mathcal{H}(x)) \leq (1-c)/2$ for some $i+1 \leq j \leq i+R$ is of order n. As there are finitely many choices for j-i, the number of indices i with $d(x_i, \partial \mathcal{H}(x)), d(x_j, \partial \mathcal{H}(x)) \leq (1-c)/2$ is of order n, where $1 \leq r = r(A) \leq R$ is the number that appears most often. In fact, we can assume that r is as large as we want. In particular, we can always choose $r > 3\delta + 1 - c$. The reason for making this choice will become clear later.

For any number $\mathcal{C} > 0$, we define $\mathcal{A}_i(\mathcal{C})$ to be the event that x_i has distance at most \mathcal{C} from the geodesic between x_0 and x_n . We also define \mathcal{B}_i to be the event $d(x_i, \partial \mathcal{H}(x)), d(x_j, \partial \mathcal{H}(x)) \leq (1-c)/2$. Our aim is to utilize the above observation in order construct SAWs that satisfy $\mathcal{A}_i(\mathcal{C})$.

Lemma 2.2. There are universal constants 0 < C < 1, C > 0 such that $\mathbb{P}_n(\mathcal{A}_i(C)) \geq C\mathbb{P}_n(\mathcal{B}_i)$ for every large enough m and every $i \geq 1$.

Proof. Consider a SAW $x \in \mathcal{B}_i$. Our aim is to define a family of walks x' for which the position of x'_0, \ldots, x'_i and the relative position between the points x'_{i+r}, \ldots, x'_n is fixed and coincides with that for x, and furthermore, the event $\mathcal{A}_i(\mathcal{C})$ occurs.

To this end, let z_i , z_{i+r} be the points in the boundary of $\mathcal{H}(x_0, \ldots, x_i)$, $\mathcal{H}(x_{i+r}, \ldots, x_n)$ closest to x_i , x_{i+r} , respectively, and write d for the distance between z_i and z_{i+r} . Consider two separating hyperplanes H_1 and H_2 that pass through z_i and z_{i+r} , respectively. In what follows, the position of

 z_{i+r} and x_{i+r}, \ldots, x_n will change but for convenience we will keep the same notation throughout the proof.

Notice that we can rotate $\mathcal{H}(x_{i+r},\ldots,x_n)$ around z_{i+r} and then around z_i so that H_2 lies in the complement of the open ball of radius d around z_i and has distance d from H_1 . Now we can apply another isometry to H_2 , if necessary, so that H_1 and H_2 are at distance

$$r - d(x_i, z_i) - d(x_{i+r}, z_{i+r}) \ge r + c - 1 > 3\delta$$

apart and the distance is attained by the pair z_i , z_{i+r} . Adding the geodesic walk $y_i = x_i, \ldots, y_{i+r} = x_{i+r}$ of length r that connects x_i to x_{i+r} results in a SAW of length n because any $y_{i+1}, \ldots, y_{i+r-1}$ is at distance at least 1 - (1-c)/2 > c from H_1 , H_2 . We can now construct a family of walks with the desired properties by choosing the jth step, $j = i+1, \ldots, i+r$ suitably, while keeping the distance between H_1 and H_2 close to $r - d(x_i, z_i) - d(x_{i+r})$ and larger than 3δ . In fact, these steps can be chosen from some domes D_j , $j = i, \ldots, i+r$ of positive area that differ only by a rotation.

It follows from the next lemma that all these SAWs satisfy $\mathcal{A}_i(\mathcal{C})$. Since the domes D_j have positive measure, the probability of $\mathcal{A}_i(\mathcal{C})$ conditioned on x_0, \ldots, x_i and the relative position between the points x_{i+r}, \ldots, x_n is bounded from below by a uniform positive constant, whenever $x \in \mathcal{B}_i$. Taking expectation we obtain the desired assertion.

In the following lemma, we will write \overline{xy} for the geodesic between points x and y in \mathbb{H}^d and $d(A, B) := \inf_{x \in A, y \in B} d(x, y) \ge \delta$ for the distance between subsets of \mathbb{H}^d .

Lemma 2.3. Consider two bi-infinite geodesics A, B in \mathbb{H}^d with $d(A, B) > 3\delta$. Assume that $x_0 \in A$ and $y_0 \in B$ realize the distance d(A, B), then the following holds: for any $x \in A$, $y \in B$ we have $d(\overline{xy}, \overline{x_0y_0}) < 2\delta$.

Proof. Let $x \in A, y \in B$ and write γ, γ' for the geodesics starting at x_0 and ending at x, y, respectively, parametrized by arc-length.

Consider an R > 0 such that $d(\gamma(R), \gamma'(R)) = 2\delta$. We claim that $d(\gamma'(R), \overline{xy}) \leq \delta$. Indeed, suppose to the contrary that $d(\gamma'(R), \overline{xy}) > \delta$. Since $d(\gamma'(R), \gamma \cup \overline{xy}) < \delta$, there exists some $t \in \gamma$ such that $d(\gamma'(R), t) < \delta$.

Hence $R - \delta < d(x_0, t) < R + \delta$, which implies that $d(\gamma(R), t) < \delta$. Thus $d(\gamma(R), \gamma'(R)) < 2\delta$ and this contradiction proves the claim.

Now for the triangle with vertices x_0, y_0, y we have that $d(\gamma'(R), \overline{x_0y_0} \cup \overline{y_0y}) < \delta$. But $d(\gamma'(R), \overline{y_0y}) > \delta$ which shows that $d(\gamma'(R), \overline{x_0y_0}) < \delta$ and completes the proof.

Using Lemma (2.2) and the fact that on any SAW of length n, the number of indices i such that \mathcal{B}_i occurs is of order n, we obtain that in a SAW of length n, the expected number of indices i, for which x_i is within distance \mathcal{C} to the geodesic between x_0 and x_n is of order n. Since the open balls of radius c/2 around the vertices of the SAW are disjoint, we get that the \mathcal{C} -neighbourhood of the geodesic has expected area of order n. But the area of the \mathcal{C} -neighbourhood of a geodesic of length k is of order k for any $k \geq 1$. This completes the proof of the theorem.

3. Open problems

In this section we will state some open problems. Given some $\varepsilon = \varepsilon(n) > 0$, we consider continuous n steps SAWs on \mathbb{H}^d , where now the steps are chosen uniformly from the sphere of radius ε and we condition on sequences in which the distance between any pair of vertices is bigger than $c\varepsilon$, for some fixed 0 < c < 1. We are interested in the behaviour of these SAWs as ε tends to 0. We will refer to the n step model as the (n, ε) -SAW. Let us write $\mathbb{E}_{n,\varepsilon}$ for the expectation with respect to this measure.

Conjecture 3.1. $\lim_{n\to\infty} \mathbb{E}_{n,1/n}(d(x_0,x_n))=0$

Conjecture 3.2. There exists a constant $1 > \beta = \beta(d, c) > 0$ such that

$$\lim_{n\to\infty} \mathbb{E}_{n,n^{-\beta}}(d(x_0,x_n))$$

exists and is not zero (possibly up to logarithmic factors).

We expect that β is increasing in c because when the self-avoidance restrictions are stronger, the walk tends to move further away from the origin.

Let us mention the observation motivating these conjectures. Given $\varepsilon > 0$, let $\frac{1}{\varepsilon}\mathbb{H}^d$ denote \mathbb{H}^d with the metric scaled by a. It is clear that the (n,ε) -SAW on \mathbb{H}^d coincides with the (n,1)-SAW on $\frac{1}{\varepsilon}\mathbb{H}^d$. Moreover, $d(x_0,x_n)=$

 $\varepsilon d_{\varepsilon}(x_0, x_n)$, where d_{ε} is the distance function on $\frac{1}{\varepsilon}\mathbb{H}^d$. As ε tends to 0, the curvature of $\frac{1}{\varepsilon}\mathbb{H}^d$ tends to 0 as well and $\frac{1}{\varepsilon}\mathbb{H}^d$ looks more and more like \mathbb{R}^d . This indicates that for certain values of n and ε , the (n, ε) -SAW on \mathbb{H}^d behaves like (n, 1)-SAW on \mathbb{R}^d . In fact, we can make a more precise prediction for the values of n and ε for which this holds. On balls of size smaller than $\frac{1}{\varepsilon}$, $\frac{1}{\varepsilon}\mathbb{H}^d$ looks similar to \mathbb{R}^d , while on balls of size larger than $\frac{1}{\varepsilon}\mathbb{H}^d$, the two spaces look different. If the (n, 1)-SAW on \mathbb{R}^d is typically at distance n^{β} , then this indicates that the (n, 1)-SAW on $n^{-\beta}\mathbb{H}^d$ behaves like the (n, 1)-SAW on \mathbb{R}^d . This reasoning leads naturally to the above questions.

As c tends to 0, we expect that β tends to 1/2 because in the limit we obtain a random walk. Also, for $d \geq 5$, we expect that β is always equal to 1/2, as in this case, the scaling limit of SAW on \mathbb{Z}^d is the Brownian motion [4].

References

- R. Bauerschmidt, H. Duminil-Copin, J. Goodman, and G. Slade, Lectures on self-avoiding walks. (English summary) Probability and statistical physics in two and more dimensions, 395-467, Clay Math. Proc., 15, Amer. Math. Soc., Providence, RI, 2012.
- [2] I. Benjamini and R. Eldan, Convex hulls in the hyperbolic space. Geom. Dedicata 160 (2012) 365-371.
- [3] D. Burago, Y. Burago and S. Ivanov, A course in metric geometry. Graduate Studies in Mathematics, 33. American Mathematical Society, Providence, RI, 2001. xiv+415 pp.
- [4] T. Hara and G. Slade, Self-avoiding walk in five or more dimensions I. The critical behaviour. Comm. Math. Phys. 147 (1992) 101-136
- [5] H. Duminil-Copin and A. Hammond, Self-avoiding walk is sub-ballistic. Comm. Math. Phys. 324 (2013) 401-423.
- [6] Z. Li, Positive speed self-avoiding walks on graphs with more than one end. arXiv:1612.02464
- [7] N. Madras and C. Wu, Self-avoiding walks on hyperbolic graphs. Combin. Probab. Comput. 14 (2005) 523-548.
- [8] C. Panagiotis Self-avoiding walks and polygons on hyperbolic graphs. arXiv:1908.00127