

Efficient emulators for scattering using eigenvector continuation

R. J. Furnstahl,^{1,*} A. Garcia,^{1,†} P. J. Millican,^{1,‡} and Xilin Zhang^{1,§}

¹*Department of Physics, The Ohio State University, Columbus, OH 43210, USA*

(Dated: October 21, 2021)

Eigenvector continuation (EC) has been shown to accurately and efficiently reproduce ground states for targeted sets of Hamiltonian parameters. It uses as variational basis vectors the corresponding ground-state eigensolutions from selected other sets of parameters. Here we extend the EC approach to scattering using the Kohn variational principle. We first test it using a model for S-wave nucleon-nucleon scattering and then demonstrate that it also works to give accurate predictions for non-local potentials, charged-particle scattering, complex optical potentials, and higher partial waves. These proofs-of-principle validate EC as an effective emulator for applying Bayesian inference to parameter estimation constrained by scattering observables.

OVERVIEW

Bayesian inference is increasingly favored for uncertainty quantification in nuclear physics calculations (e.g., see [1–5]), but the computational requirements can be substantial. In particular, Bayesian parameter estimation generally requires Monte Carlo sampling of the parameter space, with many evaluations of the likelihood with different parameters. Each evaluation may be sufficiently expensive that a full parameter estimation is infeasible. Eigenvector continuation (EC) [6, 7] has already shown that it can be used as an efficient and accurate emulator [8] to ameliorate this problem. In applying an emulator, one trains computer models of the relevant calculations using a representative set of parameters and then samples for other parameters from the model instead of full calculations. Efficient and effective EC emulators for nuclear bound-state properties and transitions have been demonstrated for many-body calculations using chiral effective field theory (χ EFT) Hamiltonians [8, 9].

We would also like to have fast EC emulators for scattering, e.g., for treating reactions and for few-body scattering used to constrain χ EFT low-energy constants [10]. The variational method for ground-state energies is well known from elementary quantum mechanics. In addition, there are variational formulations of scattering, such as those by Schwinger and Kohn (see Refs. [11–14] and references therein). The conventional applications in scattering are for two-body scattering, usually in partial waves, but the literature contains adaptations to three-body scattering, including nucleon-deuteron scattering [15], a process of particular interest for χ EFT [10]. Here we merge EC and the Kohn variational principle and explore how well it works using a series of model calculations, starting with two-body scattering in partial waves.

FORMALISM

Consider a Hamiltonian $\hat{H}(\boldsymbol{\theta}) = \hat{T} + \hat{V}(\boldsymbol{\theta})$ with adjustable parameters $\boldsymbol{\theta}$. For example, the vector $\boldsymbol{\theta}$ could

be the depth of a simple square well or the full set of low-energy constants for an effective field theory. EC is a variational method that employs a non-orthogonal basis composed of eigenvectors from different parameter sets $\{\boldsymbol{\theta}_i\}$ of the Hamiltonian. For calculating the ground state of $\hat{H}(\boldsymbol{\theta})$, the trial wave function is

$$|\psi_{\text{trial}}\rangle = \sum_{i=1}^{N_b} c_i |\psi_{\text{gs}}(\boldsymbol{\theta}_i)\rangle, \quad (1)$$

where $|\psi_{\text{gs}}(\boldsymbol{\theta}_i)\rangle$ is the ground-state eigenvector of $\hat{H}(\boldsymbol{\theta}_i)$. (The dependence of $|\psi_{\text{trial}}\rangle$ on $\boldsymbol{\theta}$ is suppressed for notational convenience.) The N_b $\boldsymbol{\theta}_i$ s are chosen either systematically or randomly to span a particular range of values, see below. The effectiveness of the EC basis can be understood by an analytic continuation analysis [6, 16].

The variational principle for the ground-state energy states that the expectation value of $\hat{H}(\boldsymbol{\theta})$ in the trial state, subject to the condition that $|\psi_{\text{trial}}\rangle$ is normalized, is stationary:

$$\delta[\langle\psi_{\text{trial}}|\hat{H}(\boldsymbol{\theta})|\psi_{\text{trial}}\rangle - \lambda(\langle\psi_{\text{trial}}|\psi_{\text{trial}}\rangle - 1)] = 0. \quad (2)$$

The stationary solution given Eq. (1) is a generalized eigenvalue problem yielding Lagrange multiplier λ_{min} , which is an upper bound to E_{gs} , and the $\{c_i\}$ provide an approximation to $|\psi_{\text{gs}}(\boldsymbol{\theta})\rangle$ [6–9].

For the extension of EC to scattering we use the Kohn variational principle (KVP) [13, 17]. There are many variational methods for scattering, but the KVP is particularly straightforward to adapt to EC in a form similar to (2). Let us start with the goal of finding the phase shift $\delta_\ell(E)$ at energy E for nonrelativistic two-body scattering in an uncoupled partial-wave channel with angular momentum ℓ and short-range forces only. In coordinate space, $\hat{T} \rightarrow -\nabla^2/2\mu$ with $\hbar = 1$ and reduced mass μ , and we allow $\hat{V}(\boldsymbol{\theta})$ to be local or nonlocal.

We take the trial wave function for the extended EC to be

$$|\psi_{\text{trial}}\rangle = \sum_{i=1}^{N_b} c_i |\psi_E(\boldsymbol{\theta}_i)\rangle, \quad (3)$$

where $|\psi_E(\boldsymbol{\theta}_i)\rangle$ is the partial-wave solution for the Schrödinger equation with Hamiltonian $\hat{H}(\boldsymbol{\theta}_i)$ at energy $E > 0$, normalized such that for every i ,

$$u_{\ell,E}^{(i)}(r) \xrightarrow{r \rightarrow \infty} \frac{1}{p} \sin\left(pr - \ell \frac{\pi}{2}\right) + \frac{\mathcal{K}_\ell^{(i)}(E)}{p} \cos\left(pr - \ell \frac{\pi}{2}\right). \quad (4)$$

Here $p = \sqrt{2\mu E}$, the scattering wave function is decomposed as

$$\langle \mathbf{r} | \psi_E(\boldsymbol{\theta}_i) \rangle = \frac{u_{\ell,E}^{(i)}(r)}{r} Y_{\ell m}(\Omega_r), \quad (5)$$

and $\mathcal{K}_\ell^{(i)}(E) = \tan \delta_\ell^{(i)}(E)$ is the partial-wave K matrix element [13] for $\hat{H}(\boldsymbol{\theta}_i)$ at energy E .

The KVP asserts that [13]

$$\beta[|\psi_{\text{trial}}\rangle] = \tau_{\text{trial}} - 2\mu \langle \psi_{\text{trial}} | \hat{H}(\boldsymbol{\theta}) - E | \psi_{\text{trial}} \rangle, \quad (6)$$

subject to the radial part of $\langle \mathbf{r} | \psi_{\text{trial}} \rangle$ being normalized as in (4) with $\mathcal{K}_\ell^{(i)}(E)/p \rightarrow \tau_{\text{trial}}$, will be a stationary approximation to $[\mathcal{K}_\ell(E)]_{\text{exact}}$ (i.e., it is accurate to second order in the difference of the exact and trial wave functions although not an upper bound in general). The normalization condition for $|\psi_{\text{trial}}\rangle$ is fulfilled if $\sum_{i=1}^{N_b} c_i = 1$, which can be imposed with a Lagrange multiplier λ . Substituting (3) into (6) with this constraint term and requiring the derivatives with respect to c_i and λ to be zero yields a simple matrix inversion problem with solution

$$c_i = \sum_{j=1}^{N_b} (\Delta \tilde{U})_{ij}^{-1} \frac{1}{p} (\mathcal{K}_\ell^{(j)}(E) - \lambda), \quad (7)$$

$$\lambda = \frac{\sum_{i,j=1}^{N_b} (\Delta \tilde{U})_{ij}^{-1} (\mathcal{K}_\ell^{(j)}(E) - 1)}{\sum_{i,j=1}^{N_b} (\Delta \tilde{U})_{ij}^{-1}}, \quad (8)$$

where

$$\Delta \tilde{U}_{ij} \equiv 2\mu \langle \psi_E(\boldsymbol{\theta}_i) | 2\hat{V}(\boldsymbol{\theta}) - \hat{V}(\boldsymbol{\theta}_i) - \hat{V}(\boldsymbol{\theta}_j) | \psi_E(\boldsymbol{\theta}_j) \rangle. \quad (9)$$

In obtaining (9) we have used that $(\hat{H}(\boldsymbol{\theta}_i) - E) |\psi_E(\boldsymbol{\theta}_i)\rangle = 0$ for every i . Finally, the stationary approximation to the exact partial-wave K matrix is

$$[\mathcal{K}_\ell(E)]_{\text{exact}} \approx \sum_{i=1}^{N_b} c_i \mathcal{K}_\ell^{(i)}(E) - \frac{p}{2} \sum_{i,j=1}^{N_b} c_i \Delta \tilde{U}_{ij} c_j. \quad (10)$$

Thus the approximation is given by a weighted average of the K matrices from the basis Hamiltonians with a correction term.

Note that the validity of the KVP relies only on the cancellation of $\delta\tau_{\text{trial}}$ with surface terms arising from the variation of $\langle \psi_{\text{trial}} | \hat{H}(\boldsymbol{\theta}) - E | \psi_{\text{trial}} \rangle$, which is satisfied by Coulomb, non-local, and complex potentials, as well as for coupled channels. (When the Coulomb potential is present, the asymptotic behavior of the scattering wave function is different from Eq. (4). For complex potentials, the $\langle \psi_E(\boldsymbol{\theta}_i) |$ factors in Eqs. (6) and (9) need to be applied with time reversal [18]. See the discussion in the Supplementary Material (SM).) It is worth pointing out that any long-range potential in $\hat{H}(\boldsymbol{\theta})$ independent of $\boldsymbol{\theta}$, such as Coulomb, will cancel from $\Delta \tilde{U}_{ij}$ in Eq. (9) and one needs only to evaluate the matrix element within the range of the remaining potentials, which simplifies calculations. Also note that Eq. (9) can be evaluated in momentum space or any other convenient basis. More details on the derivation of Eqs. (7)–(10) are given in the SM.

The matrix $\Delta \tilde{U}$ to be inverted may be expected to be increasingly ill-conditioned as the basis size N_b increases. Even for conventional applications of the KVP, there will be ill-conditioning issues for certain values of E , giving rise to so-called “Kohn anomalies” [19]. The often-recommended remedy is to use a complex formulation (involving the S matrix rather than the K matrix), which mostly avoids the problem [20–22]. Here we also have ill-conditioning, but at all E for sufficiently large N_b . We find, however, that a simple regularization of the smallest singular values of $\Delta \tilde{U}$ is sufficient to ameliorate the ill-conditioning [23, 24]. This can be done by adding a small value to the diagonal of $\Delta \tilde{U}$ (called a nugget in this context, but cf. Tikhonov regularization [23, 24]) or by using the pseudo-inverse in (8). Because we can accurately calculate test results, we can verify the efficacy of the regularization. In the following calculations, the nugget is chosen to be between 10^{-10} and 10^{-8} to optimize—by hand—those EC estimations with an ill-conditioning problem.

EC FOR A MODEL OF NN SCATTERING

We use the so-called “Minnesota potential” [25], which was developed to reproduce 1S_0 and 3S_1 nucleon-nucleon (NN) scattering phase shifts with a simple functional form, as a test example to explore the application of EC for scattering. The potential is a sum of local Gaussian terms, without Coulomb interaction or coupled channels. Each S-wave channel has a repulsive short-range term and an attractive term with longer range:

$$V_{1S_0}(r) \equiv V_{0R} e^{-\kappa_R r^2} + V_{0s} e^{-\kappa_s r^2}, \quad (11)$$

$$V_{3S_1}(r) \equiv V_{0R} e^{-\kappa_R r^2} + V_{0t} e^{-\kappa_t r^2} \quad (12)$$




FIG. 1. (a) Scattering wave functions for the Minnesota 1S_0 potential in (11) with $E = 50$ MeV. The dot-dashed curves are for four choices of $\theta_i = \{V_{0R}, V_{0s}\}$ that comprise the EC trial basis, the dashed curve is for the exact values from Ref. [25], and the solid curve is the EC prediction. The curves have a common crossing point at the value of r where the second term in (4) is zero. (b) Scattering phase shifts for the same parameter sets and the EC prediction.




FIG. 2. Sampled points in the parameter space for the Minnesota potential in the 1S_0 channel. The best parameter set from Ref. [25] is a star, four values for the EC trial basis are circles, and the crosses are test point, which are either interpolations (blue) or extrapolations (red).

The best values from Ref. [25] are 200., -91.85, and -178 MeV for V_{0R} , V_{0s} , and V_{0t} , and 1.487, 0.465, and 0.639 fm^{-2} for κ_R , κ_s and κ_t .

To illustrate how EC works, values are chosen for $\theta_i = \{V_{0R}, V_{0s}\}$ for $i = 1$ to 4, to form a trial basis for EC calculations in the 1S_0 channel. These points in the parameter space are (0., -291.85), (100., 8.15), (300., -191.85), and (300., 8.15). Figure 1(a) shows the scattering wave functions at $E = 50$ MeV from the four basis potentials (blue dot-dashed lines), the exact wave function corresponding to the best value parameters at the same energy (red dashed), and the wave function from EC based on the four-potential basis (black solid line). It is evident that with four basis elements the EC wave function agrees very well with the exact wave function. Figure 1(b) shows the corresponding phase shifts.

Again, the exact result is very well reproduced by the EC prediction, even though the wave functions and phase shifts of individual basis elements are significantly different.

Next we make a more global study with the same potential. For each channel, we vary the two potential strengths by ± 100 MeV about the best values, and scan the 2-dimensional parameter space by comparing the EC -phase shift with the exact phase shift. The values for the $\theta_i = \{V_{0R}, V_{0s}\}$ parameters are randomly drawn using Latin-hypercube sampling [26], as used in EC bound-state studies [8, 9]. A range of basis sizes N_b have been explored. For those parameter values, we compute scattering phase shifts and wave functions by directly solving the Schrödinger equation using an R -matrix package [27], which serves as input for the subsequent EC calculations using Eqs. (7)–(9). To explore the predictive power of the EC, we randomly sampled 200 points from the two-dimensional space, and for each made EC predictions as well as direct calculations using the R -matrix package, whose phase-shift calculation, as we checked, has precision—i.e., relative error—better than 10^{-8} with the order of 10^2 mesh points used therein. Comparing these results indicates the accuracy of the EC emulator.

An example of the parameter sets for this comparison protocol is shown in Fig. 2, where the sampled points in the V_{0R} - V_{0s} parameter space (for the 1S_0 channel) are shown. The trial basis points ($N_b = 4$) are blue circles, the tested sample points are blue crosses if within the convex hull of the basis points (for these the EC calculations are considered to be interpolations) and otherwise are red crosses (these EC calculations are extrapolations), and finally the best-value point is a red star.

In Fig. 3, the mean values of the relative error (in absolute value) of the EC calculations for the interpolated sample points (left panel) and the extrapolated points




FIG. 3. Relative errors between EC predictions with $\theta_i = \{V_{0R}, V_{0s}\}$ and direct calculations of $p/K_\ell(E) = p \cot \delta(E)$ for the Minnesota potential in the 1S_0 channel. This is the mean of the errors for (a) interpolated and (b) extrapolated parameter sets as shown in Fig. 2. The size of the nugget used for inverting $\Delta\tilde{U}$ is 10^{-10} here.




FIG. 4. Same as Fig. 3 but for a four-dimensional parameter space with $\theta_i = \{V_{0R}, \kappa_R, V_{0s}, \kappa_s\}$. The nugget is set to 10^{-9} here.

(right panel) are plotted against the scattering energy E (in the center of mass frame) for three calculations using $N_b = 4, 6$, and 8 basis elements. (The errors are in the value of $p/K_\ell(E) = p \cot \delta(E)$. Since the relative error is tiny here, other functions of δ have almost the same relative errors.) With a basis size of 4 , the EC calculation can reproduce the phase shift to better than 0.1 percent at almost all energies. The accuracy improves to be better than 10^{-4} , and for most energies it reaches 10^{-6} , with $N_b = 6$. For $N_b = 8$, the $\Delta\tilde{U}$ matrix becomes ill-conditioned, but after regularizing the small singular values by adding a nugget (10^{-10}) to the diagonal of this matrix when computing the matrix inversion in (7) and (8), the accuracy of these calculations is comparable to the $N_b = 6$ case. We also computed the standard deviations of the absolute value of the relative errors, and found them to be similar in size to the mean values. It is interesting to note in this case that EC works equally well for interpolated and extrapolated points.

To explore a higher-dimensional parameter set, we vary both the potential strength (± 100 MeV about the best values) and the two Gaussian widths κ_R and κ_s within a $\pm 50\%$ range about their best values. So now $\theta_i = \{V_{0R}, \kappa_R, V_{0s}, \kappa_s\}$.¹ For this demonstration, we uniformly sample 1000 test points within the four-dimensional parameter space. Figure 4 shows the parallel error information to Fig. 3, with a nugget of 10^{-9} size. For the interpolated points, the accuracy improves from 10^{-3} – 10^{-4} to 10^{-6} or better as N_b increases from 6 to 10 . For $N_b = 14$, the ill-conditioning issues require the use of a nugget but its accuracy is similar to $N_b = 10$. The results for the extrapolated parameter sets are worse

¹ Note that because the κ parameters do not appear linearly in the Hamiltonian, one can no longer make a single set of matrix elements calculations for all of the test parameter sets. In other contexts this might be a relevant computational disadvantage.

than the interpolated results for $N_b = 6$, but become as accurate with $N_b = 10$ and larger. Again, the standard deviations of the relative errors are comparable to their mean values. The parallel results for the 3S_1 channel are similar for a large enough trial basis (see the SM).

OTHER EXAMPLES: p - α AND α -Pb




FIG. 5. The relative errors for $\tan \delta(E)$ in the p - α $P_{3/2}$ channel in the two-dimensional space $\theta_i = \{V_{p\alpha,1}^{(0)}, \beta_1\}$. The nugget is set to 10^{-8} here.

To explore the effectiveness of EC for non-local potentials, the inclusion of a Coulomb potential, and for higher-partial waves, we use proton- α scattering in the $S_{1/2}$ and $P_{3/2}$ channels as examples, with the non-local potential [28]:

$$V_\ell(r', r) = V_{p\alpha, \ell}^{(0)} r'^\ell r^\ell e^{-\beta_\ell (r' + r)}. \quad (13)$$

The best values are: $V_{p\alpha,0}^{(0)} = -168.28$ MeV, $\beta_0 = 0.8$ fm $^{-1}$, $V_{p\alpha,1}^{(0)} = -291.26$ MeV, and $\beta_1 = 1.25$ fm $^{-1}$. The Coulomb potential takes the point-charge form. For each of the two channels, we vary both the potential strengths $V_{p\alpha, \ell}^{(0)}$ around its best values ± 100 MeV and the width parameters β_ℓ around its best values $\pm 50\%$. As a representative case, the relative errors for interpolated points in the $P_{3/2}$ channel are plotted in Fig. 5 for several basis sizes (additional plots for the $S_{1/2}$ channel are given in the SM). The nugget for both channels is set to be 10^{-8} . The performance of EC is again excellent except at some isolated energies, and these exceptions are not at the same energies for different basis sizes.

The Kohn variational approach also applies to complex potentials, which are extensively used in optical potentials for nuclear scattering and reactions. To test the EC for these applications, we use a Wood-Saxon optical potential constructed for describing α - ^{208}Pb low-energy

scattering [29]:

$$V(r) = V_0 f(r, R_R, a_R) + iW_0 f(r, R_I, a_I), \quad (14)$$

with $f(r, R, a) \equiv (1 + \exp(r - R)/a)^{-1}$. We take $V_0 = -100$ MeV, $W_0 = -10$ MeV, $R_R = R_I = 8.36$ fm, and $a_R = a_I = 0.58$ fm as the best value [27]. The Coulomb potential is simplified as for point charges [27].




FIG. 6. The relative errors for $\tan \delta(E)$ in the α - ^{208}Pb $\ell = 20$ channel in the two-dimensional space $\theta_i = \{V_0, W_0\}$. Note that δ is complex. The nugget is set to 10^{-10} here.

We vary the V_0 and W_0 parameters in different partial waves (so $\theta_i = \{V_0, W_0\}$). In Fig. 6, the size of relative errors for the $\ell = 20$ channel is shown as a representative example. The results for $\ell = 0$ are shown in the SM. (Note that the scattering phase shift is complex here. The vertical axis of the plot is for the modulus of the relative error.) The lower end of the energy range is chosen such that the Sommerfeld parameter η is less about 10, because the numerical calculation of Coulomb functions in the R -matrix package becomes unreliable for larger values [27, 30]. The upper end of the energy range is chosen to match Ref. [29]. The nugget used in $\Delta \tilde{U}$'s inversion is set to 10^{-10} in both $L = 20$ and $L = 0$ calculations. With 10 basis elements, the relative accuracy for interpolated points is no worse than 10^{-4} , while increasing N_b further improves it to 10^{-5} or better. Again, the errors for interpolated and extrapolated points are similar, and the standard deviations are similar in size to the mean values.

Based on these results for p - α and α -Pb scatterings, we expect EC could play an important role in fitting potential parameters for nuclear scattering and coupled-channel reactions.

SUMMARY AND OUTLOOK

We have extended the eigenvector continuation method to scattering using the Kohn variational principle. The EC enables accurate calculations of observables

for any parameter set θ given calculations of scattering wave functions and K -matrix elements from a limited number N_b of parameter sets θ_i . Unlike the bound-state application of EC, for scattering the KVP does not give an upper bound to observables but is only guaranteed to give stationary results. Nevertheless, for good trial functions the KVP has been demonstrated in the literature to give accurate results for a wide range of applications [14]. An EC basis provides a very effective trial function and its application to the KVP is simple, involving only the inversion of the matrix defined in Eq. (9). Issues of ill-conditioning with increasing basis size are successfully treated with simple regularizations.

Here we have provided representative results from a wide range of tests of the EC for scattering using model problems. These include multi-dimensional parameter sets, both local and non-local potentials, charged-particle scattering, and complex optical potentials. In all cases shown here and in all our other tests to date, the EC is found to be effective with moderate basis sizes both for interpolated and extrapolated parameter sets. We are working to formulate a robust uncertainty quantification and to develop a procedure for determining the optimal regularization parameter for ill-conditioning, which has thus far been fixed empirically.

The success of the EC enables the development of efficient emulators for scattering. In subsequent work we will demonstrate the application to coupled channels in both coordinate-space and momentum-space (which is a straightforward generalization of the presentation here) and set up the application to Nd scattering [15, 17]. It would be also interesting to apply our method to fit an NN potential to the NN energy spectra from Lattice QCD calculations, since the eigenenergies and phase shifts are directly connected.

This work was supported in part by the National Science Foundation under Grant Nos. PHY-1614460 and PHY-1913069 and by the NUCLEI SciDAC Collaboration under Department of Energy MSU subcontract RC107839-OSU

* furnstahl.1@osu.edu

† garcia.823@osu.edu

‡ millican.7@osu.edu

§ zhang.10038@osu.edu

- [1] R. J. Furnstahl, D. R. Phillips, and S. Wesolowski, *J. Phys. G* **42**, 034028 (2015), [arXiv:1407.0657](#).
- [2] X. Zhang, K. M. Nolle, and D. R. Phillips, *Phys. Lett. B* **751**, 535 (2015), [arXiv:1507.07239](#).
- [3] S. Wesolowski, R. J. Furnstahl, J. A. Melendez, and D. R. Phillips, *J. Phys. G* **46**, 045102 (2019), [arXiv:1808.08211](#).
- [4] M. Catacora-Rios, G. King, A. Lovell, and F. Nunes, *Phys. Rev. C* **100**, 064615 (2019).

- [5] L. Neufcourt, Y. Cao, S. Giuliani, W. Nazarewicz, E. Olsen, and O. B. Tarasov, *Phys. Rev. C* **101**, 014319 (2020), [arXiv:1910.12624 \[nucl-th\]](#).
- [6] D. Frame, R. He, I. Ipsen, D. Lee, D. Lee, and E. Rrapaj, *Phys. Rev. Lett.* **121**, 032501 (2018), [arXiv:1711.07090](#).
- [7] D. K. Frame, *Ab Initio Simulations of Light Nuclear Systems Using Eigenvector Continuation and Auxiliary Field Monte Carlo*, Ph.D. thesis (2019), [arXiv:1905.02782](#).
- [8] S. König, A. Ekström, K. Hebeler, D. Lee, and A. Schwenk, (2019), [arXiv:1909.08446](#).
- [9] A. Ekström and G. Hagen, *Phys. Rev. Lett.* **123**, 252501 (2019), [arXiv:1910.02922 \[nucl-th\]](#).
- [10] H. Witała, J. Golak, R. Skibiński, K. Topolnicki, E. Epelbaum, K. Hebeler, H. Kamada, H. Krebs, U. Meißner, and A. Nogga, *Few Body Syst.* **60**, 19 (2019).
- [11] M. L. Goldberger and K. M. Watson, *Collision Theory* (Wiley, New York, 1964).
- [12] R. G. Newton, *Scattering theory of waves and particles* (Dover, 2002).
- [13] J. R. Taylor, *Scattering Theory: The Quantum Theory of Nonrelativistic Collisions* (Dover, 2006).
- [14] R. Nesbet, *Variational methods in electron-atom scattering theory*, Physics of atoms and molecules (Plenum Press, 1980).
- [15] A. Kievsky, *Nucl. Phys.* **A624**, 125 (1997), [arXiv:nucl-th/9706061](#).
- [16] A. Sarkar and D. Lee, (2020), [arXiv:2004.07651 \[nucl-th\]](#).
- [17] W. Kohn, *Phys. Rev.* **74**, 1763 (1948).
- [18] M. Kamimura, *Progress of Theoretical Physics Supplement* **62**, 236 (1977).
- [19] C. Schwartz, *Phys. Rev.* **124**, 1468 (1961).
- [20] J. Z. H. Zhang, S. Chu, and W. H. Miller, *The Journal of Chemical Physics* **88**, 6233 (1988).
- [21] R. R. Lucchese, *Phys. Rev. A* **40**, 6879 (1989).
- [22] S. K. Adhikari, *Journal of Computational Physics* **103**, 415 (1992).
- [23] A. Neumaier, *SIAM Review* **40**, 636 (1998).
- [24] H. Engl, M. Hanke, and A. Neubauer, *Regularization of Inverse Problems*, Mathematics and Its Applications (Springer Netherlands, 1996).
- [25] D. Thompson, M. Lemere, and Y. Tang, *Nuclear Physics A* **286**, 53 (1977).
- [26] B. Tang, *Journal of the American Statistical Association* **88**, 1392 (1993).
- [27] P. Descouvemont, *Comput. Phys. Commun.* **200**, 199 (2016), [arXiv:1510.03540 \[nucl-th\]](#).
- [28] S. Ali, A. Ahmad, and N. Ferdous, *Rev. Mod. Phys.* **57**, 923 (1985).
- [29] G. Goldring, M. Samuel, B. Watson, M. Bertin, and S. Tabor, *Physics Letters B* **32**, 465 (1970).
- [30] A. Barnett, *Computer Physics Communications* **27**, 147 (1982).
- [31] R. Barrett, B. Robson, and W. Tobocman, *Rev. Mod. Phys.* **55**, 155 (1983), [Erratum: *Rev. Mod. Phys.* **56**, 567–567 (1984)].

Supplementary Material for Eigenvector continuation for scattering

R. J. Furnstahl,¹ A. Garcia,¹ P. J. Millican,¹ Xilin Zhang,¹

¹*Department of Physics, The Ohio State University, Columbus, Ohio 43210, USA*

KOHN VARIATIONAL METHOD

To have a self-contained presentation, we briefly review the KVP [13, 17] for single-channel scattering. The generalization to coupled-channel processes is straightforward [17, 18]. For a specified partial wave ℓ with local potential $V(r)$, the KVP stationary functional (6) with the trial radial wave function u_t (we suppress the dependence on ℓ and energy E) reduces to [13]

$$\beta[u_t] = \tau_{\text{trial}} - \int_0^\infty dr u_t(r) D u_t(r). \quad (\text{S1})$$

Here, p is the asymptotic momentum ($p = \sqrt{2\mu E}$) and

$$D \equiv -\frac{d^2}{dr^2} + \frac{\ell(\ell+1)}{r^2} + U(r) - p^2, \quad (\text{S2})$$

with $U(r) \equiv 2\mu V(r)$. The value of τ_{trial} is extracted from the asymptotic behavior of the trial wave function,

$$u_t(r) \xrightarrow{r \rightarrow \infty} \frac{1}{p} \sin(pr - \frac{1}{2}\ell\pi) + \tau_{\text{trial}} \cos(pr - \frac{1}{2}\ell\pi), \quad (\text{S3})$$

where the sine term sets the normalization. This normalization convention goes hand-in-hand with the particular form of the $\beta[u_t]$ functional, which needs to be modified for other normalizations.

For the exact radial function u_{exact} , we have

$$D u_{\text{exact}}(r) = 0, \quad (\text{S4})$$

and therefore from (S3) and (4) with $\tau_{\text{trial}} \rightarrow \tau_{\text{exact}}$,

$$\beta[u_{\text{exact}}] = \frac{1}{p} [\mathcal{K}_\ell(E)]_{\text{exact}} = \frac{1}{p} [\tan \delta_\ell(E)]_{\text{exact}}. \quad (\text{S5})$$

To see that β is a stationary functional, we write

$$u_t(r) = u_{\text{exact}}(r) + \delta u(r) \quad (\text{S6})$$

and substitute into $\delta\beta = \beta[u_t] - \beta[u_{\text{exact}}]$. Using (S4) and keeping only to first order in δu , we obtain

$$\delta\beta = \delta\tau - \int_0^\infty dr u_{\text{exact}}(r) D \delta u(r) + \mathcal{O}(\delta u^2). \quad (\text{S7})$$

We want to act D to the left to take advantage of (S4), which requires partially integrating the derivatives in D . This yields only surface terms, to which we can use the asymptotic forms:

$$\delta\beta = \delta\tau + (u_{\text{exact}} \delta u' - u'_{\text{exact}} \delta u) \Big|_0^\infty + \mathcal{O}(\delta u^2). \quad (\text{S8})$$

The lower limit does not contribute because

$$u_{\text{exact}}(0) = u_t(0) = \delta u(0) = 0, \quad (\text{S9})$$

while the upper limit, after using Eq. (S3) for u_{exact} and u_t , yields $-\delta\tau[\sin^2(pr - \frac{1}{2}\ell\pi) + \cos^2(pr - \frac{1}{2}\ell\pi)] = -\delta\tau$ (with two other terms canceling), so in the end

$$\delta\beta = 0 + \mathcal{O}(\delta u^2). \quad (\text{S10})$$

Thus, the functional $\beta[u_t]$ approximates τ_{exact} at its stationary point up to $\mathcal{O}(\delta u^2)$.

If the long-range Coulomb potential is present, the asymptotic behavior of the radial basis functions and of u_t differs by the argument in the sine and cosine functions, namely,

$$\sin(pr - \frac{1}{2}\ell\pi) \rightarrow \sin(pr - \frac{1}{2}\ell\pi - \eta \ln 2pr + \sigma_\ell), \quad (\text{S11})$$

and similarly for the cos function. Here η is the so-called Sommerfeld parameter and σ_ℓ the pure Coulomb phase shift [13]. Then the phase shift δ_ℓ would be the shorter-range interaction-induced phase shift, i.e., the total phase shift with σ_ℓ subtracted. (Note that the total phase shift here is measured with respect to the incoming and outgoing spherical waves in the form of $\exp[\pm i(pr - \eta \ln 2pr)]$.)

For a general non-local potential, the above derivation still holds, except that the integral involving $V(r)$ in Eq. (S1) needs to be changed to a double integration, i.e., $\int dr dr' u_t(r) V(r, r') u_t(r')$. It should be emphasized that the potential in the function may be complex in general, but in this case no complex conjugation is to be applied to u_t in the integral [18]. The applicability of KVP for complex potentials is important for this approach to be used in optical potential model fitting.

The generalization to coupled channels can be found e.g., in Ref. [18, 31]. The central step involves finding that the variation of the integral in the definition of the β functional comes from the end points/surface terms of the integral (using Green's theorem), which exactly cancel the variation of the first term in Eq. (S1), which now turns into the K-matrix \mathcal{K} [31].

ADAPTING EC TO SCATTERING

To adapt EC to the scattering problem, we do not solve an energy eigenvalue problem but instead we find solutions at a specified scattering energy $E = p^2/2\mu$. We still have a set of N_b Hamiltonians specified by $\{\theta_i\}$ but now for each one we find the scattering solution for energy E and the corresponding phaseshift. This will be

our basis. This identifies the $\tau_i = \mathcal{K}_\ell^{(i)}/p$ value in Eq. (4) for each basis wave function. A trial wave function as in (3), which we write as

$$u_t(r) = \sum_i c_i u_i(r; E), \quad (\text{S12})$$

with each u_i normalized according to (4), will have the asymptotic form

$$u_t(r) \xrightarrow{r \rightarrow \infty} \left(\sum_{i=1}^N c_i \right) \frac{1}{p} \sin(pr - \frac{1}{2}\ell\pi) + \left(\sum_{i=1}^N c_i \tau_i \right) \cos(pr - \frac{1}{2}\ell\pi), \quad (\text{S13})$$

because the u_i all have the same p . Matching, we find the constraint:

$$\sum_{i=1}^N c_i = 1. \quad (\text{S14})$$

So we substitute (S12) into (S1) and require β to be stationary with respect to the c_i s, subject to the constraint (S14), which we incorporate using a Lagrange multiplier λ . The functional is

$$\begin{aligned} \beta[u_t] &= \sum_j c_j \tau_j(E) - \sum_{jk} c_j c_k \int_0^\infty dr u_j(r; E) \left[-\frac{d^2}{dr^2} + \frac{l(l+1)}{r^2} + 2\mu V(r; \boldsymbol{\theta}) - p^2 \right] u_k(r; E) \\ &= \sum_j c_j \tau_j - \sum_{jk} c_j c_k \int_0^\infty dr u_j(r; E) (2\mu) [V(r; \boldsymbol{\theta}) - V_k(r)] u_k(r; E). \end{aligned} \quad (\text{S15})$$

To get the second line we have added and subtracted $V_k(r)$ inside the integral and used that each of the $u_j(r)$ wave functions are eigenstates with their corresponding V_j but all the same E :

We define

$$\Delta U_{jk} \equiv \int_0^\infty dr u_j(r; E) (2\mu) [V(r; \boldsymbol{\theta}) - V_k(r)] u_k(r; E). \quad (\text{S16})$$

Note that this is not a symmetric matrix. Then the functional to make stationary is:

$$\tau_{\text{exact}} \approx \beta[u_t] = \sum_j c_j \tau_j - \sum_{jk} c_j \Delta U_{jk} c_k. \quad (\text{S17})$$

We want to be this stationary under the constraint that the sum of the c_j coefficients is one. So

$$\frac{\partial}{\partial c_i} \left[\sum_j c_j \tau_j - \sum_{jk} c_j c_k \Delta U_{jk} - \lambda \left(\sum_j c_j - 1 \right) \right] = 0, \quad (\text{S18})$$

$$\frac{\partial}{\partial \lambda} \left[\sum_j c_j \tau_j - \sum_{jk} c_j c_k \Delta U_{jk} - \lambda \left(\sum_j c_j - 1 \right) \right] = 0. \quad (\text{S19})$$

The second equation just gives us back the constraint. The first line, for each i , gives

$$\tau_i - c_k \Delta U_{ik} - c_j \Delta U_{ji} - \lambda = 0 \quad (\text{S20})$$

or

$$\sum_j (\Delta U^\top + \Delta U)_{ij} c_j \equiv \sum_j \Delta \tilde{U}_{ij} c_j \quad (\text{S21})$$

$$= \tau_i - \lambda, \quad (\text{S22})$$

which defines $\Delta \tilde{U}_{ij}$ as in (9). Finally,

$$c_j = \sum_i (\Delta \tilde{U})_{ji}^{-1} (\tau_i - \lambda), \quad (\text{S23})$$

in agreement with (7).

We can identify λ by summing this last equation over j and setting it equal to 1, then solving for λ :

$$\lambda = \frac{\sum_{ij} (\Delta \tilde{U})_{ji}^{-1} \tau_i - 1}{\sum_{ij} (\Delta \tilde{U})_{ji}^{-1}}, \quad (\text{S24})$$

in agreement with (8). To get our estimate of $\tau_{\text{exact}} = [\mathcal{K}_\ell(E)]_{\text{exact}}/p$ as in (10), we substitute from (S23) and (S24) into (S17).

ADDITIONAL RESULTS

Here, we provide additional results in the partial wave channels not already presented. The setups for these calculations are the same as for their counterparts in the main text. Figures S1 and S2 provide results for NN scattering in the 3S_1 channel with the Minnesota potential by

varying $\theta_i = \{V_{0R}, V_{0s}\}$ and $\theta_i = \{V_{0R}, V_{0s}, \kappa_R, \kappa_R\}$, respectively. Figure S3 shows the results for p - α scattering in the S-wave channel, in parallel to the P-wave results in the main text. Figure S4 plots results for S-wave α - ^{208}Pb scattering. In these plots, both mean values and

the standard deviations (std) of the relative errors (absolute values) are plotted against E . As mentioned in the main text and shown in these plots, the std values are similar to the mean values in general.




FIG. S1. Relative errors between EC predictions and direct calculations of $p/K_e(E) = p \cot \theta(E)$ for the Minnesota potential in the 3S_1 channel. To regularize the $\Delta\tilde{U}$ matrix for its inversion, the nugget is set to 10^{-9} here. The top plots show the means of errors for sampled parameter sets in Fig. 2 while the bottom plots show the standard deviations of these errors.




FIG. S2. Same as Fig. S1 but for the 4-dimensional parameter space $\theta_i = \{V_{0R}, \kappa_R, V_{0s}, \kappa_s\}$. The nugget for $\Delta\tilde{U}$'s inversion is set to 10^{-8} .




FIG. S3. The relative errors for $\tan \delta(E)$ in the p - α $S_{1/2}$ channel in the two-dimensional space $\theta_i = \{V_{p\alpha,0}^{(0)}, \beta_0\}$. The nugget is set to 10^{-8} here.




FIG. S4. The relative errors for $\tan \delta(E)$ in the α - ^{208}Pb $\ell = 0$ channel in the two-dimensional space $\theta_i = \{V_0, W_0\}$. Note that δ is complex. The nugget is set to 10^{-10} here.