

An identity for the sum of inverses of odd divisors of n in terms of the number of representations of n as a sum of r squares

Sumit Kumar Jha

Abstract. Let

$$\sum_{\substack{d|n \\ d \equiv 1(2)}} \frac{1}{d}$$

denote the sum of inverses of odd divisors of a positive integer n , and let $c_r(n)$ be the number of representations of n as a sum of r squares where representations with different orders and different signs are counted as distinct. The aim is of this note is to prove the following interesting combinatorial identity:

$$\sum_{\substack{d|n \\ d \equiv 1(2)}} \frac{1}{d} = \frac{1}{2} \sum_{r=1}^n \frac{(-1)^{n+r}}{r} \binom{n}{r} c_r(n).$$

Mathematics Subject Classification (2010). 11A05, 11P99.

Keywords. Sum of divisors; Sum of squares; Partial Bell polynomials; Faà di Bruno's formula; Jacobi's theta function.

1. Main result

In the following, let

$$\sum_{\substack{d|n \\ d \equiv 1(2)}} \frac{1}{d}$$

denote the sum of inverses of odd divisors of a positive integer n .

Definition 1.1. [2, formula 7.324] Let $\theta(q)$ be the following infinite product

$$\theta(q) := \prod_{j=1}^{\infty} \frac{1-q^j}{1+q^j} = \sum_{n=-\infty}^{\infty} (-1)^n q^{n^2}$$

where $|q| < 1$.

Definition 1.2. For any positive integer r define $c_r(n)$ by

$$\theta(q)^r = \sum_{n=0}^{\infty} c_r(n) (-1)^n q^n$$

where $c_r(n)$ be the number of representations of n as a sum of r squares where representations with different orders and different signs are counted as distinct.

Our aim is to derive the following identity.

Theorem 1. For all positive integers n we have

$$\sum_{\substack{d|n \\ d \equiv 1(2)}} \frac{1}{d} = \frac{1}{2} \sum_{r=1}^n \frac{(-1)^{n+r}}{r} \binom{n}{r} c_r(n). \quad (1)$$

We require following two lemmas for our proof.

Lemma 1. For all positive integers n we have

$$2 \sum_{\substack{d|n \\ d \equiv 1(2)}} \frac{1}{d} = \frac{1}{n!} \sum_{k=1}^n (-1)^k (k-1)! B_{n,k} \left(\theta'(0), \theta''(0), \dots, \theta^{(n-k+1)}(0) \right) \quad (2)$$

where $B_{n,k} \equiv B_{n,k}(x_1, x_2, \dots, x_{n-k+1})$ are the partial Bell polynomials defined by [1, p. 134]

$$B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = \sum_{\substack{1 \leq i \leq n, \ell_i \in \mathbb{N} \\ \sum_{i=1}^n i \ell_i = n \\ \sum_{i=1}^n \ell_i = k}} \frac{n!}{\prod_{i=1}^{n-k+1} \ell_i!} \prod_{i=1}^{n-k+1} \left(\frac{x_i}{i!} \right)^{\ell_i}.$$

Proof. It is easy to see that

$$\begin{aligned} \log(\theta(q)) &= \sum_{j=1}^{\infty} \log(1 - q^j) - \sum_{j=1}^{\infty} \log(1 + q^j) \\ &= - \sum_{j=1}^{\infty} \sum_{l=1}^{\infty} \frac{q^{lj}}{l} + \sum_{j=1}^{\infty} \sum_{l'=1}^{\infty} \frac{q^{l'j} (-1)^{l'}}{l'} \\ &= - \sum_{n=1}^{\infty} q^n \left(\sum_{d|n} \frac{1 - (-1)^d}{d} \right). \end{aligned}$$

Let $f(q) = \log q$. Using Faà di Bruno's formula [1, p. 137] we have

$$\frac{d^n}{dq^n} f(\theta(q)) = \sum_{k=1}^n f^{(k)}(\theta(q)) \cdot B_{n,k} \left(\theta'(q), \theta''(q), \dots, \theta^{(n-k+1)}(q) \right). \quad (3)$$

Since $f^{(k)}(q) = \frac{(-1)^{k-1} (k-1)!}{q^k}$ and $\theta(0) = 1$, letting $q \rightarrow 0$ in the above equation gives us Equation (2). \square

Lemma 2. We have, for positive integers n, k ,

$$B_{n,k} \left(\theta'(0), \theta''(0), \dots, \theta^{(n-k+1)}(0) \right) = (-1)^n \frac{n!}{k!} \sum_{r=1}^k (-1)^{k-r} \binom{k}{r} c_r(n) \quad (4)$$

Proof. We start with the generating function for the partial Bell polynomials [1, Equation (3a') on p. 133], and proceed as follows

$$\begin{aligned} \sum_{n=k}^{\infty} B_{n,k} \left(\theta'(0), \theta''(0), \dots, \theta^{(n-k+1)}(0) \right) \frac{q^n}{n!} &= \frac{1}{k!} \left(\sum_{j=1}^{\infty} \theta^{(j)}(0) \frac{q^j}{j!} \right)^k \\ &= \frac{1}{k!} (\theta(q) - 1)^k \\ &= \frac{1}{k!} \sum_{r=0}^k (-1)^{k-r} \binom{k}{r} \theta(q)^r \\ &= \frac{1}{k!} \sum_{r=0}^k (-1)^{k-r} \binom{k}{r} \sum_{n=0}^{\infty} (-1)^n c_r(n) q^n \end{aligned}$$

to conclude Equation (4). \square

Proof of Theorem 1. We combine Equation (2) and Equation (4) to obtain

$$\begin{aligned} 2 \sum_{\substack{d|n \\ d \equiv 1(2)}} \frac{1}{d} &= (-1)^n \sum_{k=1}^n \frac{1}{k} \sum_{r=1}^k (-1)^r \binom{k}{r} c_r(n) \\ &= (-1)^n \sum_{r=1}^n (-1)^r c_r(n) \sum_{k=r}^n \frac{1}{k} \binom{k}{r} \\ &= (-1)^n \sum_{r=1}^n \frac{(-1)^r}{r} \binom{n}{r} c_r(n), \end{aligned}$$

where we used the identity $\sum_{k=r}^n \frac{1}{k} \binom{k}{r} = \frac{1}{r} \binom{n}{r}$ which can be proved using the Pascal's formula

$$\binom{k}{r-1} = \binom{k+1}{r} - \binom{k}{r}.$$

This concludes proof of our main result Equation (1). \square

References

- [1] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, D. Reidel Publishing Co., Dordrecht, 1974.
- [2] N. J. Fine, *Basic Hypergeometric Series and Applications*, American Mathematical Soc., 1988.

Sumit Kumar Jha
International Institute of Information Technology
Hyderabad-500 032, India
e-mail: kumarjha.sumit@research.iiit.ac.in