## SHARP WEYL LAWS WITH SINGULAR POTENTIALS

#### RUPERT L. FRANK AND JULIEN SABIN

ABSTRACT. We consider the Laplace—Beltrami operator on a three-dimensional Riemannian manifold perturbed by a potential from the Kato class and study whether various forms of Weyl's law remain valid under this perturbation. We show that a pointwise Weyl law holds, modified by an additional term, for any Kato class potential with the standard sharp remainder term. The additional term is always of lower order than the leading term, but it may or may not be of lower order than the sharp remainder term. In particular, we provide examples of singular potentials for which this additional term violates the sharp pointwise Weyl law of the standard Laplace-Beltrami operator. For the proof we extend the method of Avakumović to the case of Schrödinger operators with singular potentials.

#### 1. Introduction and main results

In this paper we investigate the question to what extent the Weyl law in its various forms for the Laplace–Beltrami operator on a compact manifold remains valid when a singular potential is added.

To be more precise, let (M, g) be a three-dimensional compact Riemannian manifold without boundary. We comment on the three-dimensionality assumption in Remark 4.5 below. Our results are valid under not very restrictive regularity assumptions on M, namely  $C^4$ smoothness, as discussed in Remark 4.6, but they are also new in the  $C^{\infty}$  setting.

We denote by  $-\Delta_g$  the (nonnegative) Laplace–Beltrami operator on (M, g). Its spectrum is discrete and we denote by  $\mathbb{1}(-\Delta_g \leq t)(x, y)$  the integral kernel of its spectral projection  $\mathbb{1}(-\Delta_g \leq t)$  corresponding to eigenvalues  $\leq t$ . Moreover,

$$N(t, -\Delta_g) = \operatorname{Tr} \mathbb{1}(-\Delta_g \leqslant t) = \int_M \mathbb{1}(-\Delta_g \leqslant t)(x, x) \, dv_g(x)$$

denotes the number of its eigenvalues  $\leq t$ , counting multiplicities.

We are interested in four different forms of Weyl's law concerning the limit  $t \to \infty$ , namely

- integrated Weyl law:  $N(t, -\Delta_g) = \frac{t^{3/2}}{6\pi^2} \operatorname{Vol}_g(M) + o(t^{3/2})$
- pointwise Weyl law:  $\mathbb{1}(-\Delta_g \leqslant t)(x,x) = \frac{t^{3/2}}{6\pi^2} + o(t^{3/2})$  uniformly in  $x \in M$
- sharp integrated Weyl law:  $N(t, -\Delta_g) = \frac{t^{3/2}}{6\pi^2} \operatorname{Vol}_g(M) + \mathcal{O}(t)$
- sharp pointwise Weyl law:  $\mathbb{1}(-\Delta_g \leqslant t)(x,x) = \frac{t^{3/2}}{6\pi^2} + \mathcal{O}(t)$  uniformly in  $x \in M$ .

Date: July 8, 2020.

<sup>© 2020</sup> by the authors. This paper may be reproduced, in its entirety, for non-commercial purposes.

Clearly, the pointwise versions imply the integrated versions and the sharp versions imply the regular versions. The integrated Weyl law was originally proved by Weyl [61] in Euclidean space and its pointwise version in Euclidean space is due to Carleman [11, 12]. The extension of the pointwise Weyl law to manifolds is due to Minakshisundaram and Pleijel [44]. The sharp pointwise Weyl law for closed manifolds is due to Avakumović [4], Levitan [40], and Hörmander [29]. The adjective 'sharp' refers to the fact that there are manifolds, for instance the round sphere or, more generally, Zoll manifolds, for which the remainder  $\mathcal{O}(t)$  cannot be improved. On the other hand, there is a large literature on improved Weyl laws with o(t) remainders for certain manifolds, both in integrated [19, 37, 28, 60, 26, 32, 5] and in pointwise [49, 57] form. These improved Weyl laws will not play a major role in this paper and we only comment on them in Remark 6.6 below.

Let  $V: M \to \mathbb{R}$  belong to the Kato class, that is, it is measurable and

$$\lim_{\varepsilon \to 0} \sup_{x \in M} \int_{d_g(x,y) < \varepsilon} \frac{|V(y)|}{d_g(x,y)} \, dv_g(y) = 0.$$

Under this assumption  $-\Delta_g + V$  can be defined via a quadratic form as a selfadjoint, bounded below operator with discrete spectrum. When V is smooth, it is well-known [29] that all four versions of Weyl's law remain valid when  $-\Delta_g$  is replaced by  $-\Delta_g + V$ . Our goal in this paper is to investigate to what extent this is true for general Kato class V. We will find both preservation and violation, depending on the specific form of Weyl's law. Let us summarize our main results.

**Theorem 1.1** (Pointwise Weyl law for Kato class potentials). Let  $V: M \to \mathbb{R}$  be in the Kato class. Then, as  $t \to +\infty$  and uniformly in  $x \in M$ 

$$\mathbb{1}(-\Delta_g + V \leqslant t)(x, x) = \frac{t^{3/2}}{6\pi^2} + o(t^{3/2}).$$

On the other hand, we show that there is no  $\varepsilon > 0$  such that the term  $o(t^{3/2})$  in Theorem 1.1 can be replaced by  $\mathcal{O}(t^{(3-\varepsilon)/2})$  for all Kato class V and, in particular, we show that the sharp form of the pointwise Weyl law is *not* valid for all Kato class potentials. In fact, in Proposition 6.4 we will show that the sharp pointwise Weyl law fails for the potential

$$V(x) = \gamma \frac{\chi(d_g(x, x_0))}{d_g(x, x_0)^{2-\eta}}$$
(1.1)

with  $\eta \in (0,1)$ ,  $x_0 \in M$ ,  $\gamma \in \mathbb{R} \setminus \{0\}$  and  $\chi$  a cut-off function which is  $\equiv 1$  near zero. More precisely, we will see that there is an explicit singular term in  $\mathbb{1}(-\Delta_g + V \leq t)(x,x)$  that lives at scale  $t^{-1/2}$  around  $x_0$  and has size  $t^{(3-\eta)/2}$ . We refer to this proposition and Lemma 6.5 for a detailed analysis of this example.

Next, we show that the sharp pointwise Weyl law remains valid for V satisfying a mild additional regularity condition.

**Theorem 1.2** (Sharp pointwise Weyl law). Assume that  $V: M \to \mathbb{R}$  satisfies for some  $\varepsilon' > 0$ ,

$$\sup_{x \in M} \int_{d_g(y,x) < \varepsilon'} \frac{|V(y)|}{d_g(y,x)^2} \, dv_g(y) < \infty. \tag{1.2}$$

Then, as  $t \to +\infty$  and uniformly in  $x \in M$ ,

$$\mathbb{1}(-\Delta_g + V \leqslant t)(x, x) = \frac{t^{3/2}}{6\pi^2} + \mathcal{O}(t).$$

Note that the condition in this theorem is, in particular, satisfied if  $V \in L^q(M)$  for some q > 3. For comparison, for any q < 3 there is an  $\eta \in (0,1)$  such that (1.1) belongs to  $L^q$ , so q = 3 is the threshold for the validity of the sharp pointwise Weyl law on the  $L^q$  scale.

Finally, we discuss the integrated forms of Weyl's law. Clearly, Theorem 1.1 yields this in the form of an  $o(t^{3/2})$  remainder. Remarkably, however, despite the possible failure of the sharp pointwise Weyl law, the integrated form remains valid in the strong form for arbitrary Kato class potentials. This was recently shown by Huang and Sogge [30]. Here we give an independent proof of their result and extend it to potentials V that are sums of Kato class and  $L^{3/2}(M)$  functions. Note that the Kato class and  $L^{3/2}(M)$  share the same critical scaling behavior and that neither one is contained in the other one.

**Theorem 1.3** (Sharp integrated Weyl law). Assume that  $V: M \to \mathbb{R}$  is the sum of a function in the Kato class and a function in  $L^{3/2}(M)$ . Then, as  $t \to +\infty$ ,

$$N(t, -\Delta_g + V) = \frac{t^{3/2}}{6\pi^2} + \mathcal{O}(t).$$
 (1.3)

This concludes the summary of our main results. Let us now comment on the motivation for these results and on their method of proofs.

Potentials with singularities appear naturally in physics, most notably the Coulomb case  $|x|^{-1}$  in three dimensions. Spectral asymptotics in this case, both in integrated [31, 52, 53, 58] and in pointwise form [20, 21], have been extensively studied. As we will explain below, however, our asymptotic regime is different and we find phenomena of another nature.

Our motivation to consider Kato class potentials is two-fold. First, it is natural to investigate the validity of Weyl's law under rather minimal assumptions on the potential V for which the Schrödinger operator  $-\Delta_g + V$  can be defined as selfadjoint, bounded below operator. Both membership to the Kato class and to  $L^{3/2}(M)$  are almost optimal conditions in this respect. For instance, singularities of the type  $|x|^{-\beta}$  for  $\beta < 2$  are allowed for both classes. Note that the exponent  $\beta = 2$  is critical with respect to the scaling of the Laplacian and if, in this case, the coefficient of the singularity is too large negative, then the corresponding quadratic form is not bounded from below. We mention in passing (see Remark A.1) that neither the Kato class is contained in  $L^{3/2}$ , nor the other way around.

Second, the property that makes the Kato class more natural for our analysis than  $L^{3/2}$  is its important property that all eigenfunctions of  $-\Delta_g + V$  with Kato class V are bounded. In fact, this class is almost sharp with respect to this property, in the sense that the fact that  $e^{-t(-\Delta_g+V)}$  maps  $L^2$  to  $L^{\infty}$  essentially implies that V belongs to the Kato class [1]. In contrast, eigenfunctions for potentials in  $L^{3/2}$  need not be bounded. Since boundedness of eigenfunctions is an obvious requirement for a pointwise Weyl law to hold, the Kato class appears naturally in this context.

When comparing our results with other results about Weyl's law with singular potentials, it is important to distinguish between the 'large eigenvalue regime'  $-\Delta_q + V \leq t$ ,  $t \to +\infty$ ,

which we consider here, and the 'semiclassical regime'  $-h^2\Delta_g + V \leq 0$ ,  $h \to 0$ . The large eigenvalue regime corresponds to a semiclassical regime  $-h^2\Delta_g - 1 + h^2V \leq 0$ ,  $h = t^{-1/2}$ , with a small parameter in front of the nonconstant part of the potential. Effects of singularities have been studied in details in the semiclassical regime but, as far as we know, their study in the large eigenvalue regime has only begun very recently in [8, 30].

The analysis in the semiclassical regime is, in part, motivated by that of the Schrödinger operator  $-\Delta - Z^{4/3}\varphi(Z^{1/3}x)$  in  $L^2(\mathbb{R}^3)$  with a potential satisfying  $\varphi(y) \sim |y|^{-1}$  as  $y \to 0$ , which appears in atomic physics in connection with Thomas–Fermi theory. Via rescaling we see that it corresponds to the semiclassical regime with  $h = Z^{-1/3}$ . It is well known that the singularity of  $\varphi$  has an effect, called the Scott correction, on the semiclassical expansion of the Riesz means of order one (meaning that instead of counting eigenvalues, we sum the negative eigenvalues). On the other hand, this effect does not appear when looking at the number of eigenvalues less than a fixed energy level E < 0 [27]. This phenomenon was thoroughly studied in [56] where it was shown, in particular, that singularities  $|x|^{-\beta}$  with  $\beta < 2$  do not affect the local eigenvalue counting function to order  $\mathcal{O}(h^{-d+1})$ . We emphasize that the results [27, 56] involve (at least locally) integrated Weyl laws and that it is not clear to us whether these methods can make assertions about pointwise Weyl laws. Moreover, the specific powerlike form of the singularity seems to be important, whereas Kato class potentials can have much more complicated singularities. Violations of the main term in the Weyl law in the semi-classical regime have also been studied, see for instance [38, 7].

Let us return to the large eigenvalue regime studied in this paper. One of our key findings, which is behind Theorems 1.1, 1.2 and 1.3, is that there is an additional term in the pointwise Weyl law. Theorem 1.1 says that this term is  $o(t^{3/2})$  for all Kato class potentials, Theorem 1.2 says that it is  $\mathcal{O}(t)$  under the additional assumption (1.2) and Theorem 1.3 says that it is  $\mathcal{O}(t)$  when integrated over M for all Kato class potentials. On the other hand, for the potential (1.1) we show that this term contributes a constant times  $t^{(3-\eta)/2}$  at the singularity.

We have not seen this additional term discussed in the literature before. This term is neither of semiclassical origin nor of spectral origin, in contrast to the term giving rise to the Scott correction [31, 52, 53], for instance.

Besides drawing attention to this additional term in the pointwise Weyl law, a goal of this paper is to advertise the method that we use to prove such results. It is the method of Avakumović [4], which was developed in the 1950's but seems to have been forgotten in the last decades. As we will demonstrate, this method, which was originally developed for the case V=0, is a powerful tool (at least in three dimensions) to get very precise results on Weyl laws and is well suited to extensions to singular potentials V. For comparison, it is not clear to us how to adapt the approaches of the related works [56, 30] to obtain pointwise Weyl laws. In particular, we hope that our result can be used as a benchmark for the proof of pointwise Weyl laws with singular potentials by other methods, in the sense that such other methods must reveal the additional terms violating the standard Weyl law that we uncover in case of a strong enough singularity.

In the remaining part of this introduction we now discuss Avakumović's method and our new additional ingredients.

Avakumović's method relies on two ingredients, namely Tauberian theorems and parametrix estimates. The Tauberian theorems used by Avakumović concern a higher order Stieltjes transform instead of the Fourier transform, as in the work of Levitan [40] and then in many works following [29]. Consequently, one will seek a parametrix for the resolvent (or its powers) for spectral parameters far away from the spectrum and not for the wave propagator.

In connection with the parametrix estimates, two of Avakumović's fundamental insights are the following. First, an exponentially small remainder in these estimates is acceptable in the Tauberian theorems. This observation immediately leads to the remainder bound  $\mathcal{O}(t^{(N-1)/2}\operatorname{dist}(x,\partial\Omega)^{-1})$  in the case of an open set  $\Omega\subset\mathbb{R}^N$  with Dirichlet boundary conditions; see [3]. The second key insight is that some error terms which are not exponentially small are themselves (higher order) Stieltjes transforms and therefore acceptable in a sufficiently precise version of the Tauberian theorems.

Our main task is to prove corresponding parametrix estimates in the case where V is added. In fact, we will use the same parametrix for the Green's function as Avakumović does, so the task will be to bound the effect of V on the difference between the true Green's function and the parametrix. Several terms in this difference can be estimated by modifying Avakumović's arguments. In fact, in the case of bounded potentials V, this has already been done by Bojanić in [10]. When V is unbounded, however, there are other terms which are not as small as the previous ones and which need to be treated separately. These are the terms that lead to the 'additional term' in the pointwise Weyl law, mentioned above, and that may lead to a violation of the sharp pointwise Weyl law.

This analysis eventually leads to the decomposition

$$(-\Delta_g + V + \lambda)^{-2}(x, x) = \frac{1}{8\pi\sqrt{\lambda}} - \int_0^\infty \frac{t^{3/2} r_0^V(t, x) + R^V(t, x)}{(t + \lambda)^3} dt + \text{exponentially small as } \lambda \to +\infty.$$
 (1.4)

The term  $R^V(t,x)$  here is of size  $\mathcal{O}(t)$  and can be dealt with as in Avakumović's paper. The important new contribution is the term  $t^{3/2}r_0^V(t,x)$ , which is defined through an explicit, absolutely convergent power series and encompasses the singular contributions coming from V. This is the term that ultimately give rise to violations of the sharp pointwise Weyl law. A careful analysis of the structural properties of this term eventually leads to the various versions of Weyl's law in Theorems 1.1, 1.2 and 1.3.

One can get from the decomposition (1.4) to a corresponding decomposition of the spectral density by a Tauberian theorem for a higher order Stieltjes transform. Note that

$$(-\Delta_g + V + \lambda)^{-2}(x, x) = 2 \int_0^\infty \frac{\mathbb{1}(-\Delta_g + V \leqslant t)(x, x)}{(t + \lambda)^3} dt, \qquad \frac{1}{8\pi\sqrt{\lambda}} = 2 \int_0^\infty \frac{t^{3/2}/(6\pi^2)}{(t + \lambda)^3} dt.$$

The Tauberian theorem gives

$$\mathbb{1}(-\Delta_g + V \leqslant t)(x, x) = \frac{t^{3/2}}{6\pi^2} - \frac{1}{2}r_0^V(t, x)t^{3/2} + \mathcal{O}(t). \tag{1.5}$$

Due to the presence of the term  $t^{3/2}r_0^V(t,x)$ , we need a more precise Tauberian theorem than Avakumović. Fortunately for us, however, such a theorem can be established using techniques that Avakumović introduced in a different context in [2], relying on the Tauberian theorem of Ingham [33] and Karamata [34]. Since we have not been able to find the relevant Tauberian theorem in the literature, we have decided to add a complete proof in Sections 2 and 3.

The precise statement of the decomposition in (1.4) appears in Proposition 5.1 and its proof takes up all of Section 5. We think it is worthwhile to precede this proof by our interpretation of Avakumović's result for V=0 in Section 4. One difference is that we find it more convenient to work with  $(-\Delta_g + \lambda)^{-2}(x,x)$  than with Avakumović's

$$\lim_{y \to x} \left( (-\Delta + \lambda)^{-1}(x, y) - \lim_{\lambda' \to 0} \left( (-\Delta + \lambda')^{-1}(x, y) - \frac{1}{\lambda' \operatorname{Vol}_g M} \right) \right).$$

The derivation of Theorems 1.1, 1.2 and 1.3 from decomposition (1.5), as well as the counterexamples to the strong pointwise Weyl law appear in Section 6.

In a final Section 7 we sketch an application of these ideas to the case of subsets of Euclidean space and in an appendix we recall relevant results about the Kato class.

**Acknowledgements:** Partial support through US National Science Foundation grant DMS-1363432 (R.L.F.) and ANR DYRAQ ANR-17-CE40-001 (J. S.) are acknowledged.

### 2. A Tauberian theorem for the Laplace transform

In this and in the next section we give a proof of the Tauberian theorem that is required for the proof of our main result. The proof of this theorem, which concerns the Stieltjes transform, relies on another Tauberian theorem, namely one for the Laplace transform. The latter, which is due to Ingham [33, Thm. III] and Karamata [35, Satz II] (see also [34]) is the topic of this section. For our application we need a nonasymptotic version of this theorem.

We denote by  $BV_{loc}[0,\infty)$  the set of functions on  $[0,\infty)$  whose total variation on any compact subinterval of  $[0,\infty)$  is finite. Let  $A \in BV_{loc}[0,\infty)$  be such that  $A(u) = \mathcal{O}(e^{\alpha u})$  as  $u \to +\infty$  for all  $\alpha > 0$ . Under these assumptions, the Laplace transform

$$f(s) := \int_0^\infty e^{-us} dA(u) \tag{2.1}$$

is well-defined and analytic on Re s>0. Here, dA denotes the Lebesgue–Stieljes measure on  $[0,\infty)$  associated to A in such a way that  $dA([0,t))=A(t_-)-A(0)$ . This is equivalent to the definition of the Lebesgue–Stieljes measure [13, Chap. 4] when A is extended by A(0) on  $(-\infty,0)$ . We will often assume that A(0)=0 and we emphasize that this does not imply an assumption on  $A(0_+)$ .

The crucial assumption on f is that  $\frac{f(s)-a}{s}$  has a continuous extension to s=it for  $t \in [-T,T]$  for some  $a \in \mathbb{C}$  (which of course implies that a=f(0)), and that A has some generalized monotonicity property.

The following theorem is the main result of this section.

**Theorem 2.1.** Let  $A \in BV_{loc}[0,\infty)$  with A(0) = 0 and  $A(u) = \mathcal{O}(e^{\alpha u})$  as  $u \to +\infty$  for all  $\alpha > 0$ . Assume that there are  $c, \delta > 0$  such that for all  $u + \delta \geqslant v \geqslant u \geqslant 0$  we have

 $A(v)-A(u)\geqslant -c$  and assume that there are  $a\in\mathbb{C}$  and T>0 such that  $\varphi(s):=\frac{f(s)-a}{s}$ , with f from (2.1), can be continuously extended to s=it for  $t\in[-T,T]$ . Then, for all  $\omega>0$  and with an absolute constant C,

$$|A(\omega)| \leqslant C \left( |a| + \int_{-T}^{T} |\varphi(it)| dt + \left( 1 + \frac{1}{\delta T} \right) c \right).$$

As we already mentioned, this theorem is due to Ingham [33, Thm. III] and Karamata [35, Satz II] (see also [34]). A slightly weaker form that, however, is *not* sufficient for our purposes, appears in [59, Sec. 22.37]. In these works the result is only stated in an asymptotic form, whereas we need a nonasymptotic form. Such a version is probably well-known to experts and it is stated in [25, Hilfss. 6] without proof and with reference to the unpublished work [45], to which we do not have access. We also note that in [4, Sec. 1.6] Avakumović sketches how this result under the additional assumption  $A(\omega) = \mathcal{O}(\sqrt{\omega})$  can be deduced from known results. While the required nonasymptotic forms of these known results are probably also known to experts, we have not been able to find an appropriate reference. In view of this situation we believe it is useful to include a self-contained and complete proof of Theorem 2.1.

Remark 2.2. Avakumović [4] uses the following consequence of Theorem 2.1: Let  $A \in BV_{loc}[0,\infty)$  with A(0)=0 and  $A(u)=\mathcal{O}(e^{\alpha u})$  for all  $\alpha>0$ . Assume that there is a c>0 such that for all  $0 \le u \le v \le u+1$  we have  $A(v)-A(u) \ge -c$  and assume that

$$s \mapsto \int_0^\infty e^{-su} A(u) \, du =: \varphi(s)$$

can be extended analytically to  $|s| < \varepsilon$  for some  $\varepsilon > 0$ . Then for all  $\omega > 0$ ,

$$|A(\omega)| \leqslant C \left( \int_{-\varepsilon/2}^{\varepsilon/2} |\varphi(it)| dt + \left(1 + \frac{2}{\varepsilon}\right) c \right)$$

where C is an absolute constant. Indeed, since  $\varphi(s) = f_A(s)/s$ , where  $f_A$  is the Laplace transform of A, this follows from Corollary 2.1 with a = 0,  $T = \varepsilon/2$  and  $\delta = 1$ .

The proof of Theorem 2.1 uses the Jackson kernel, whose properties are summarized in the following lemma.

**Lemma 2.3.** For any  $\eta \in [-1,1]$ , define

$$\ell(\eta) := \begin{cases} -\frac{4}{3}(1 - 2|\eta|)^3 + \frac{8}{3}(1 - |\eta|)^3 & \text{if } |\eta| \le 1/2, \\ \frac{8}{3}(1 - |\eta|)^3 & \text{if } 1/2 \le |\eta| \le 1. \end{cases}$$

Then for all  $y \in \mathbb{R}$ ,

$$K(y) := \int_{-1}^{1} \ell(\eta) e^{-i\eta y} d\eta = \left(\frac{\sin(y/4)}{y/4}\right)^{4}$$

and, in particular, K > 0 almost everywhere and K, |y|K,  $y^2K \in L^1(\mathbb{R})$ .

Proof of Theorem 2.1. We follow closely the proof of [35, Satz II]. First, we extend A by zero on  $(-\infty, 0)$ . The property  $A(v) - A(u) \ge -c$  then holds for all  $u \le v \le u + \delta$ .

Step 1. As a preliminary step we note that by iterating the assumption on A, we obtain for all  $u \leq v$ 

$$A(v) - A(u) \geqslant -c\left(1 + \frac{v - u}{\delta}\right). \tag{2.2}$$

Step 2. We now show that

$$\left| \int_0^\infty TK(T(v-u))A(u) \, du \right| \leqslant \frac{8\pi}{3}|a| + \frac{4}{3} \int_{-T}^T |\varphi(it)| \, dt =: \Theta. \tag{2.3}$$

For all  $s \in \mathbb{C}$  with  $\operatorname{Re} s > 0$ , we have by integration by parts

$$\frac{f(s)}{s} = \int_0^\infty e^{-su} A(u) \, du$$

and therefore

$$\varphi(s) = \frac{f(s) - a}{s} = \int_0^\infty (A(u) - a)e^{-su} du.$$

Now let  $v \in \mathbb{R}$ ,  $\sigma > 0$  and  $t \in [-T, T]$ . We multiply the previous formula at  $s = \sigma + it$  by  $\ell(t/T)e^{ivt}$  and integrate over  $t \in [-T, T]$ . By Fubini, it follows that

$$\int_{-T}^{T} \varphi(\sigma + it)\ell(t/T)e^{ivt} dt = \int_{0}^{\infty} TK(T(v - u)) (A(u) - a) e^{-\sigma u} du.$$

Since  $\varphi(s)$  is continuous up to Re s=0, we can take the limit  $\sigma \to 0$  of the left side. On the right side, we write

$$A(u) - a = \left(A(u) + c\left(1 + \frac{u}{\delta}\right)\right) - \left(a + c\left(1 + \frac{u}{\delta}\right)\right).$$

For the second parenthesis on the right side we can apply dominated convergence, using the fact that  $(1+|y|)K(y) \in L^1$ . For the first parenthesis we apply monotone convergence, using (2.2), A(0) = 0 and  $K \ge 0$ . In this way, we obtain in the limit  $\sigma \to 0$ 

$$\int_{-T}^{T} \varphi(it)\ell(t/T)e^{ivt} dt = \int_{0}^{\infty} TK(T(v-u)) (A(u)-a) du.$$

Using

$$\int_0^\infty TK(T(v-u))\,du = \int_{-\infty}^{Tv} K(y)\,dy \leqslant \int_{\mathbb{R}} K(y)\,dy = \frac{8\pi}{3}\,,$$

and  $0 \le \ell(\eta) \le 4/3$ , we deduce (2.3).

Step 3. Let us introduce

$$B(u) := -\inf \{ A(u') : 0 \le u' \le u \}.$$

Since A(0) = 0 we have  $B \ge 0$  and, clearly,

$$A(u) \geqslant -B(u)$$
 for all  $u \geqslant 0$ . (2.4)

For later purposes, we note that the assumption on A implies

$$B(v) - B(u) \leqslant c$$
 for all  $0 \leqslant u \leqslant v \leqslant u + \delta$ ,

and therefore, by iteration,

$$B(v) - B(u) \leqslant c \left(1 + \frac{v - u}{\delta}\right)$$
 for all  $0 \leqslant u \leqslant v$ . (2.5)

Step 4. We now prove an upper bound on A in terms of B, namely,

$$A(v) \leqslant B(v) + c\left(1 + \frac{M_1}{\delta T}\right) + \frac{3}{4\pi}\Theta \quad \text{for all } v \geqslant 0,$$
 (2.6)

with  $M_1 = \int_0^\infty y K(y) \, dy / \int_0^\infty K(y) \, dy$ . Indeed, by (2.3) and (2.2), we have for any  $v \geqslant 0$ ,

$$\begin{split} \Theta \geqslant & \int_0^v TK(T(v-u))A(u)\,du + \int_v^\infty TK(T(v-u))A(u)\,du \\ \geqslant & -B(v)\int_0^v TK(T(v-u))\,du + A(v)\int_v^\infty TK(T(v-u))\,du \\ & -c\int_v^\infty TK(T(v-u))\left(1 + \frac{u-v}{\delta}\right)du \\ \geqslant & -\int_0^\infty K(y)\,dy\left(B(v) - A(v) - c\left(1 + \frac{M_1}{\delta T}\right)\right), \end{split}$$

which is the same as (2.6).

Step 5. Next, we prove that B is bounded, namely,

$$B(v) \leqslant M\left(\Theta + c\left(1 + \frac{1}{\delta T}\right)\right)$$
 (2.7)

for an absolute constant M. Indeed, by (2.3), (2.2), (2.6) and (2.5) we have for any  $v \ge 0$ ,

$$\begin{split} -\Theta &\leqslant \int_{0}^{v} TK(T(v-T^{-1}-u))A(u)\,du + \int_{v}^{\infty} TK(T(v-T^{-1}-u))A(u)\,du \\ &\leqslant A(v)\int_{-\infty}^{v} TK(T(v-T^{-1}-u))\,du + c\int_{-\infty}^{v} TK(T(v-T^{-1}-u))\left(1+\frac{v-u}{\delta}\right)du \\ &+ \int_{v}^{\infty} TK(T(v-T^{-1}-u))B(u)\,du \\ &+ \left(c\left(1+\frac{M_{1}}{\delta T}\right)+\pi^{-1}\Theta\right)\int_{v}^{\infty} TK(T(v-T^{-1}-u))\,du \\ &\leqslant A(v)\int_{-\infty}^{v} TK(T(v-T^{-1}-u))\,du + c\int_{-\infty}^{v} TK(T(v-T^{-1}-u))\left(1+\frac{v-u}{\delta}\right)du \\ &+ B(v)\int_{v}^{\infty} TK(T(v-T^{-1}-u))\,du + c\int_{v}^{\infty} TK(T(v-T^{-1}-u))\left(1+\frac{u-v}{\delta}\right)du \\ &+ \left(c\left(1+\frac{M_{1}}{\delta T}\right)+\pi^{-1}\Theta\right)\int_{v}^{\infty} TK(T(v-T^{-1}-u))\,du \,. \end{split}$$

This implies, with some absolute constant  $M_2$ ,

$$A(v) \int_{-\infty}^{v} TK(T(v - T^{-1} - u)) du$$

$$\geqslant -B(v) \int_{v}^{\infty} TK(T(v - T^{-1} - u)) du - M_2 \left(\Theta + c\left(1 + \frac{1}{\delta T}\right)\right),$$

which is the same as

$$A(v) \int_{-1}^{\infty} K(y) dy \geqslant -B(v) \int_{-\infty}^{-1} K(y) dy - M_2 \left(\Theta + c \left(1 + \frac{1}{\delta T}\right)\right).$$

Since B is nondecreasing, this implies

$$-B(v)\int_{-1}^{\infty} K(y) dy \geqslant -B(v)\int_{-\infty}^{-1} K(y) dy - M_2\left(\Theta + c\left(1 + \frac{1}{\delta T}\right)\right),$$

which is the same as (2.7) with  $M = M_2 / \int_{-1}^{1} K(y) dy$ .

Step 6. The bound in the theorem now follows from (2.3), (2.6) and (2.7).

#### 3. A Tauberian theorem for the Stieltjes transform

In this section we will deduce a Tauberian theorem for the Stieltjes transform from the one for the Laplace transform proved in the previous section. The Stieltjes transform or, more precisely, a higher order version of it arises naturally in our context through powers of the resolvent of elliptic operators.

**Theorem 3.1.** Let  $\delta, t_0 > 0$ . Let  $B_0, B_1, B_2 : [0, \infty) \to \mathbb{R}$  be such that  $B_0$  is bounded,  $B_1$  is nondecreasing and right-continuous, and assume that

$$C_{B_2} := \sup_{t \geqslant 0} \frac{|B_2(t)|}{t_0 + t}, \quad -C_{B_0} := \inf_{0 \leqslant u \leqslant v \leqslant u + \delta} \min\{u(B_0(v^2) - B_0(u^2)), 0\}$$

are finite. Assume also that  $u \mapsto u^3 B_0(u^2)$ ,  $u \mapsto B_2(u^2)$  are right-continuous and belong to  $BV_{loc}[0,\infty)$ . Assume that there are  $C_0 > 0$ ,  $\Lambda > 0$  and  $\varepsilon_0 \in (0,\eta)$  such that for all  $\lambda \geqslant \Lambda$ ,

$$\left| \int_0^\infty \frac{B_0(t)t^{3/2} + B_1(t) + B_2(t)}{(t+\lambda)^3} dt \right| \leqslant C_0 \frac{e^{-\varepsilon_0\sqrt{\lambda}}}{\lambda}.$$

Then  $B_0(t) t^{3/2} + B_1(t) + B_2(t) = \mathcal{O}(t)$  as  $t \to +\infty$ . More precisely, one has for all  $t \ge 0$ ,

$$|B_0(t) t^{3/2} + B_1(t) + B_2(t)| \le \mathcal{B} \varepsilon_0^{-1} (t + \varepsilon_0^{-2})$$

where, for some universal constant C,

$$\mathcal{B} \leqslant C\varepsilon_0^3 \left( 1 + \varepsilon_0 \sqrt{\Lambda} e^{\varepsilon_0 \sqrt{\Lambda}/2} \right) \left( \|B_0\|_{L^{\infty}} \Lambda^{3/2} + |B_1(0)| + C_{B_2} t_0 (1 + \delta^{-1} \varepsilon_0^{-1}) + C_{B_2} \Lambda \right)$$

$$+ C\varepsilon_0^3 \Lambda \left( \varepsilon_0 \sqrt{\Lambda} e^{\varepsilon_0 \sqrt{\Lambda}/2} + \varepsilon_0^{-2} \Lambda^{-1} \right) C_0$$

$$+ C \left( \varepsilon_0 + \delta^{-1} \right) \left( C_{B_0} + \|B_0\|_{L^{\infty}} \left( \delta + \varepsilon_0^{-1} \right) + C_{B_2} \left( 1 + \ln(1 + \delta \varepsilon_0) \right) \right).$$

This theorem generalizes [59, Thm. 22.38], whose proof seems to be influenced by Avakumović's work [2]. In fact, [59, Thm. 22.38] corresponds to the special case where  $B_0$  is a negative constant, where  $B_2 \equiv 0$  and where A vanishes near the origin. For our application it will be crucial that  $B_0$  is not required to be constant.

We will typically apply this theorem in the situation where  $\Lambda = C_1/\varepsilon_0^2$ ,  $\delta = \varepsilon_0^{-1}$  and  $t_0 = 0$ . In this case the bound on  $\mathcal{B}$  simplifies to

$$\mathcal{B} \leqslant C\left(\left(1 + C_1^2 e^{\sqrt{C_1}/2}\right) \left(\|B_0\|_{L^{\infty}} + \varepsilon_0^3 |B_1(0)| + \varepsilon_0 C_{B_2} + \varepsilon_0 C_0\right) + \varepsilon_0 C_{B_0}\right). \tag{3.1}$$

*Proof.* To enhance the readability of the proof, we present here only the proof of the qualitative assertion  $\mathcal{O}(t)$  in the theorem, without keeping track of the precise dependence of the constants. The details of these more precise bounds are deferred to Appendix B. We abbreviate

$$A(t) := B_0(t) t^{3/2} + B_1(t) + B_2(t)$$
.

Step 1. We show that, for all t > 0.

$$|A(t)| \le C(1+t^{3/2}),$$
 (3.2)

see (B.1) for a quantitative version. The bounds on  $B_0$  and  $B_2$  imply that

$$\int_0^\infty \frac{B_0(t)t^{3/2}}{(t+\lambda)^3} dt \leqslant C\lambda^{-1/2}, \ \int_0^\infty \frac{B_2(t)}{(t+\lambda)^3} dt \leqslant C(\lambda^{-2} + \lambda^{-1}),$$

so, in view of the exponential decay of the Stieltjes transform of A, we deduce that

$$\int_0^\infty \frac{B_1(t) - B_1(0)}{(t+\lambda)^3} dt \leqslant C(\lambda^{-1/2} + \lambda^{-2}).$$

By the monotonicity of  $B_1$ , we deduce that for all  $\lambda > 0$ ,

$$(B_1(\lambda) - B_1(0))/(8\lambda^2) \leqslant \int_{\lambda}^{\infty} \frac{B_1(t) - B_1(0)}{(t+\lambda)^3} dt \leqslant C(\lambda^{-1/2} + \lambda^{-2}),$$

which gives  $|B_1(t)| \leq C(1+t^{3/2})$  and therefore also (3.2).

Step 2. We claim that

$$s \mapsto \int_0^\infty e^{-su} A(u^2) du$$
 is analytic in  $\{\operatorname{Re} s > 0\} \cup \{|s| < \varepsilon_0\}$ . (3.3)

This qualitative assertion is supplemented in Appendix B by the quantitative bounds (B.5) for  $s = \sigma > 0$  and (B.6) for  $|s| \leq \varepsilon_0/2$ .

In order to prove (3.3) we use the representation formula (recall the Yukawa potential)

$$e^{-s\sqrt{t}} = \frac{1}{\pi} \int_0^\infty \frac{\sin(s\sqrt{\lambda})}{\lambda + t} d\lambda.$$

This implies that for all s, t > 0,

$$\frac{e^{-s\sqrt{t}}}{\sqrt{t}} = \int_0^\infty \frac{\lambda^{3/2} \kappa(s\sqrt{\lambda})}{(t+\lambda)^3} d\lambda \tag{3.4}$$

with

$$\kappa(z) := \frac{8 \sin z - z \cos z}{\pi z^3} \quad \text{for all } z \in \mathbb{C}.$$
 (3.5)

Using (3.4) we can write

$$\int_0^\infty e^{-su} A(u^2) \, du = \frac{1}{2} \int_0^\infty \frac{e^{-s\sqrt{t}}}{\sqrt{t}} A(t) \, dt = \frac{1}{2} \int_0^\infty \lambda^{3/2} \, \kappa(s\sqrt{\lambda}) \int_0^\infty \frac{A(t)}{(t+\lambda)^3} \, dt \, d\lambda \, .$$

The fact that  $\kappa$  is entire and satisfies the bound

$$|\kappa(z)| \leqslant Ce^{|z|} \quad \text{for all } z \in \mathbb{C} \,,$$
 (3.6)

together with the assumed exponential decay for the Stieltjes transform implies (3.3).

Step 3. In order to apply Theorem 2.1, we would like to deal with functions vanishing near the origin and we define, with a parameter  $u_0 > 0$  to be specified later,

$$g_0(u) := \begin{cases} A((u - u_0)^2) - A(0) & \text{if } u \geqslant u_0, \\ 0 & \text{if } u < u_0. \end{cases}$$

We claim that

$$s \mapsto \int_0^\infty e^{-su} dg_0(u)$$
 is analytic in  $\{\operatorname{Re} s > 0\} \cup \{|s| < \varepsilon_0\}$  (3.7)

and

$$\left| \int_0^\infty e^{-tu} dg_0(u) \right| \leqslant C e^{-tu_0} (1 + t^{-3}) \tag{3.8}$$

for all t > 0. These qualitative assertions are supplemented by the quantitative bounds (B.7) for  $s = \sigma > 0$  and (B.8) for  $|s| \leq \varepsilon_0/2$ .

To prove (3.7) and (3.8) we define  $g(u) := A(u^2) - A(0)$ . Then, using g(0) = 0, we see that

$$\int_0^\infty e^{-su} dg(u) = s \int_0^\infty e^{-su} g(u) \, du = s \int_0^\infty e^{-su} A(u^2) \, du - A(0) \, .$$

According to (3.3) this is analytic in  $\{\text{Re } s > 0\} \cup \{|s| < \varepsilon_0\}$ . Moreover, from (3.2) we obtain the rough bound, for all t > 0,

$$\left| \int_0^\infty e^{-tu} dg(u) \right| \leqslant C(1 + t^{-3})$$

In terms of the function  $g_0$ , we have

$$\int_0^\infty e^{-su} dg_0(u) = \int_{u_0}^\infty e^{-su} dg(u - u_0) = \int_0^\infty e^{-s(u + u_0)} dg(u) = e^{-su_0} \int_0^\infty e^{-su} dg(u).$$

Therefore the analyticity properties and the bound of the corresponding integral with g imply (3.7) and (3.8).

Step 4. We apply Theorem 2.1 to the function

$$h(u) := \int_{[0,u]} \frac{dg_0(w)}{w^2}.$$

Note that, since  $g_0$  vanishes for  $u \leq u_0$ , the function h is well-defined, belongs to  $h \in BV_{loc}[0,\infty)$  and satisfies h(0) = 0. Moreover, its Laplace transform

$$f(s) := \int_0^\infty e^{-su} dh(u)$$

is well-defined for Re s > 0 by (3.8) and satisfies

$$f(s) = \int_0^\infty \frac{e^{-su}}{u^2} dg_0(u) = \int_s^\infty \int_{s'}^\infty \int_0^\infty e^{-s''u} dg_0(u) ds'' ds'.$$
 (3.9)

Therefore, in view of (3.7), f is analytic in  $\{\text{Re } s > 0\} \cup \{|s| < \varepsilon_0\}$ . This qualitative assertion is supplemented with the quantitative bounds (B.9) on f(0) and (B.10) on (f(s) - f(0))/s for  $|s| \leq \varepsilon_0/2$ .

Step 5. We claim that the function g satisfies the almost monotonicity property  $h(v) - h(u) \ge -c$  for  $u + \delta \ge v \ge u \ge 0$ . The proof is not complicated, but somewhat lengthy and we defer it to Appendix B; see (B.12).

Step 6. According to Steps 4 and 5, the function h satisfies all the assumptions of Theorem 2.1 and we deduce that  $h(u) = \mathcal{O}(1)$  as  $u \to +\infty$ . Note now that since  $g_0$  is right-continuous,

$$\int_0^x uh(u) \, du = \int_{[0,x]} \frac{dg_0(v)}{v^2} \int_v^x u \, du = \frac{1}{2}x^2 h(x) - \frac{1}{2}g_0(x).$$

Hence,  $g_0(x) = x^2 h(x) - 2 \int_0^x u h(u) du = \mathcal{O}(x^2)$  as  $x \to +\infty$ , meaning that  $A(u) = \mathcal{O}(u)$  as  $u \to +\infty$ . A quantitative version of this bound, valid for all u > 0 is proved in (B.14). This completes the proof of Theorem 3.1.

### 4. The method of Avakumović

We now apply Theorem 3.1 to prove the sharp Weyl laws in 3D, inspired by the method of Avakumović. Let (M, g) a three-dimensional smooth compact Riemannian manifold without boundary, and let  $-\Delta_g$  the (non-negative) Laplace-Beltrami operator on (M, g). For  $\lambda > 0$  and  $x, y \in M$  with  $x \neq y$ , denote the Green's function by

$$G_{\lambda}(x,y) = (-\Delta_q + \lambda)^{-1}(x,y).$$

The key result to obtain a sharp Weyl law is the following Avakumović-type result:

**Proposition 4.1.** There are constants  $C_1, C_2, C_3 > 0$ ,  $\varepsilon_0 > 0$ , and a function  $R : [0, \infty) \times M \to \mathbb{R}$  such that for all  $x \in M$  and for all  $\lambda \geq C_1/\varepsilon_0^2$ ,

$$\left| \partial_{\lambda} G_{\lambda}(x,x) + \frac{1}{8\pi\sqrt{\lambda}} - \int_{0}^{\infty} \frac{R(t,x)}{(t+\lambda)^{3}} dt \right| \leqslant C_{2} \frac{e^{-\varepsilon_{0}\sqrt{\lambda}/4}}{\lambda}$$

and for all  $x \in M$ ,  $R(\cdot, x)$  is  $C^1$  on  $[0, \infty)$  and for all  $t \ge 0$ ,

$$|R(t,x)| \leqslant C_3 t.$$

More precisely, the constants  $C_1, C_2, C_3$  and  $\varepsilon_0$  can be chosen as

$$C_1 = C \mathcal{M}_{\rho/2} \,, \qquad C_2 = \frac{C \,\mathcal{M}_{\rho/2}^{3/2} \,\mathcal{M}_{\rho/2}'}{\rho} \,, \qquad C_3 = \frac{C \,\mathcal{M}_{\rho/2}^{1/2} \,\mathcal{M}_{\rho/2}'}{\rho} \,, \qquad \varepsilon_0 = \frac{\rho}{C \,\mathcal{M}_{\rho/2}^{1/2}} \,,$$

where C is an absolute constant,  $\rho$  is the injectivity radius of M and  $\mathcal{M}_{\rho/2}$  and  $\mathcal{M}'_{\rho/2}$  are explicit constants, defined below, which depend in a scale invariant way on M.

Corollary 4.2. We have, as  $\lambda \to \infty$ , uniformly in  $x \in M$ ,

$$\mathbb{1}(-\Delta_g \leqslant \lambda)(x,x) = \frac{\lambda^{3/2}}{6\pi^2} + \mathcal{O}(\lambda).$$

More precisely, we have for all  $x \in M$  and all  $t \ge 0$ ,

$$\left| \mathbb{1}(-\Delta_g \leqslant \lambda)(x, x) - \frac{\lambda^{3/2}}{6\pi^2} \right| \leqslant C_0 \rho^{-1} \left( t + \rho^{-2} \right),$$

where  $C_0$  depends only on upper bounds on  $\mathcal{M}_{\rho/2}$ ,  $\mathcal{M}'_{\rho/2}$  and  $\rho^3/\operatorname{Vol}_g M$ .

Proof of Corollary 4.2 assuming Proposition 4.1. Define for all  $t \ge 0$ 

$$A(t,x) := \mathbb{1}(-\Delta_g \leqslant t)(x,x) - t^{3/2}/(6\pi^2) + R(t,x)/2.$$

Let  $\lambda_n$  denote the eigenvalues of  $-\Delta_g$  in nondecreasing order and repeated according to multiplicities and  $\varphi_n$  the corresponding normalized eigenfunctions. Then

$$\mathbb{1}(-\Delta_g \leqslant t)(x,x) = \sum_{\lambda_n \leqslant t} |\varphi_n(x)|^2$$

and thus

$$\int_0^\infty \frac{\mathbb{1}(-\Delta_g \leqslant t)(x,x)}{(\lambda+t)^3} dt = \frac{1}{2} \sum_{n>0} \frac{|\varphi_n(x)|^2}{(\lambda_n+\lambda)^2} = -\frac{1}{2} \partial_\lambda G_\lambda(x,x).$$

Next, we have

$$\int_0^\infty \frac{t^{3/2}}{(\lambda + t)^3} dt = \frac{3\pi}{8\sqrt{\lambda}},$$

so that, by Proposition 4.1,

$$\left| \int_0^\infty \frac{A(t,x)}{(t+\lambda)^3} dt \right| = \left| -\frac{1}{2} \partial_\lambda G_\lambda(x,x) - \frac{1}{16\pi\sqrt{\lambda}} + \frac{1}{2} \int_0^\infty \frac{R(t,x)}{(t+\lambda)^3} dt \right| \leqslant C_2 \frac{e^{-\varepsilon_0\sqrt{\lambda}/4}}{\lambda}$$

for all  $\lambda \geq C_1/\varepsilon_0^2$ . The result then follows from Theorem 3.1 in the simplified form (3.1) with  $B_0 \equiv -(6\pi)^{-1}$ ,  $B_1(t) = \mathbb{1}(-\Delta_g \leq t)(x,x)$ ,  $B_2(t) = R(t,x)/2$ . Note, in particular, that the almost monotonicity assumption on  $B_0$  in that theorem is satisfied with  $C_{B_0} = 0$  and any  $\delta \geq 0$ . In order to obtain the claimed bound we apply Theorem 3.1 with the choice  $\delta = \varepsilon_0^{-1}$ . (The parameters  $\varepsilon_0$  in Theorem 3.1 and in Proposition 4.1 have essentially the same meaning, up to a multiplicative constant.) We also use the fact that  $B_1(0) = 1/\operatorname{Vol}_g M$ .  $\square$ 

4.1. **Proof of Proposition 4.1.** We first recall the construction of a local parametrix for  $G_{\lambda}$ . Let us denote by  $\rho > 0$  the injectivity radius of M. Then, for any  $y \in M$ , the exponential  $\exp_y$  at y is well-defined on the set  $\{v \in T_yM, |v| < \rho\}$ , and maps it to the set  $\{x \in M, d_g(x,y) < \rho\}$ . As explained in Avakumović [4], in Minakshisundaram–Pleijel [44], or in [6, Sec. III.E.3], for any  $(x,y) \in M \times M$  with  $d_g(x,y) < \rho$ , the function

$$\theta(x,y) := |\det T_{\exp_y^{-1}(x)} \exp_y|,$$

where T is the tangent map, is such that  $\theta(y,y) = 1$  and such that the Riemannian volume in normal coordinates around y is  $\theta(\exp_y(v), y) dv$ , where dv is the Lebesgue measure on  $T_yM$ . The function  $U_0 := \theta^{-1/2}$  is then smooth on  $V_\rho := \{(x,y) \in M \times M, d_g(x,y) < \rho\}$ .

In order to quantify the dependence of our error estimates on the geometry of the manifold, we introduce for  $\varepsilon \in (0, \rho]$ 

$$\mathcal{M}_{\varepsilon} := \sup_{(x,y) \in V_{\varepsilon}} \frac{U_0(x,y) + \varepsilon^2 |(-\Delta_g)_x U_0(x,y)|}{\min\{U_0(x,y)^2, U_0(y,x)^2\}} \quad \text{and} \quad \mathcal{M}'_{\varepsilon} := \sup_{(x,y) \in V_{\varepsilon}} U_0(x,y) \,.$$

Let  $\chi \in C_0^{\infty}(\mathbb{R})$  with  $\chi \equiv 1$  on [-1/2, 1/2] and  $\chi \equiv 0$  outside of (-1, 1). We consider  $\chi$  as fixed and consider  $L^{\infty}$  bounds on  $\chi$ ,  $\chi'$  and  $\chi''$  as absolute constants. For  $\varepsilon \in (0, \rho]$  and  $x \neq y$ , we define

$$T_{\lambda,\varepsilon}(x,y) := \frac{e^{-\sqrt{\lambda}d_g(x,y)}}{4\pi d_g(x,y)} U_0(x,y) \chi(\varepsilon^{-1}d_g(x,y))$$

and note that

$$||T_{\lambda,\varepsilon}||_{L_x^{\infty}L_y^2} \leqslant \frac{C}{\lambda^{1/4}}.$$
(4.1)

Next, let

$$\gamma_{\lambda,\varepsilon}(x,y) := T_{\lambda,\varepsilon}(x,y) - G_{\lambda}(x,y)$$

and compute using the expression of the Laplace-Beltrami operator in normal coordinates [6, Sec. II.G.V]

$$\begin{split} R_{\lambda,\varepsilon}(x,y) &:= (-\Delta_g + \lambda)_x \gamma_{\lambda,\varepsilon}(x,y) \\ &= \frac{e^{-\sqrt{\lambda} d_g(x,y)}}{4\pi d_g(x,y)} \chi(\varepsilon^{-1} d_g(x,y)) (-\Delta_g)_x U_0(x,y) \\ &- \frac{e^{-\sqrt{\lambda} d_g(x,y)}}{4\pi d_g(x,y)} U_0(x,y) \left(\varepsilon^{-2} \chi''(\varepsilon^{-1} d_g(x,y)) - 2\varepsilon^{-1} \sqrt{\lambda} \chi'(\varepsilon^{-1} d_g(x,y))\right). \end{split}$$

This implies

$$|R_{\lambda,\varepsilon}(x,y)| \leqslant C\mathcal{M}_{\varepsilon} \frac{e^{-\frac{\sqrt{\lambda}}{2}d_g(x,y)}}{\varepsilon^2 d_g(x,y)} \min\{U_0(x,y)^2, U_0(y,x)^2\} \mathbb{1}(d_g(x,y) < \varepsilon)$$

and, consequently,

$$||R_{\lambda,\varepsilon}||_{L_x^{\infty}L_y^1} + ||R_{\lambda,\varepsilon}||_{L_y^{\infty}L_x^1} \leqslant \frac{C\mathcal{M}_{\varepsilon}}{\varepsilon^2\lambda} \quad \text{and} \qquad ||R_{\lambda,\varepsilon}||_{L_x^{\infty}L_y^2} + ||R_{\lambda,\varepsilon}||_{L_y^{\infty}L_x^2} \leqslant \frac{C\mathcal{M}_{\varepsilon}\mathcal{M}_{\varepsilon}'}{\varepsilon^2\lambda^{1/4}}. \quad (4.2)$$

Applying  $(-\Delta_g + \lambda)^{-1}$  to the equation defining  $R_{\lambda,\varepsilon}$ , we obtain the integral equation

$$\gamma_{\lambda,\varepsilon}(x,y) = \int_M T_{\lambda,\varepsilon}(x,z) R_{\lambda,\varepsilon}(z,y) \, dv_g(z) - \int_M \gamma_{\lambda,\varepsilon}(x,z) R_{\lambda,\varepsilon}(z,y) \, dv_g(z).$$

Formally, this equation can be solved by iteration and one obtains the series representation

$$\gamma_{\lambda,\varepsilon}(x,y) = \sum_{n\geqslant 1} (-1)^{n+1} \gamma_{\lambda,\varepsilon}^{(n)}(x,y)$$

with

$$\gamma_{\lambda,\varepsilon}^{(n)}(x,y) := \int_M dv_g(z_1) \cdots \int_M dv_g(z_n) T_{\lambda,\varepsilon}(x,z_1) R_{\lambda,\varepsilon}(z_1,z_2) \cdots R_{\lambda,\varepsilon}(z_n,y).$$

We now prove bounds on  $\gamma_{\lambda,\varepsilon}^{(n)}$  which show, in particular, that the above series converges provided  $\lambda \varepsilon^2$  is large enough. By the Cauchy-Schwarz inequality with respect to the  $z_1$ -integration we obtain

$$|\gamma_{\lambda,\varepsilon}^{(n)}(x,y)| \leqslant ||T_{\lambda,\varepsilon}||_{L_x^{\infty}L_y^2} ||R_{\lambda,\varepsilon}||_{L_y^{\infty}L_x^2} ||R_{\lambda,\varepsilon}||_{L_x^{\infty}L_x^1}^{n-1}. \tag{4.3}$$

Inserting the bounds from (4.1) and (4.2) we find that for all  $n \ge 1$  and all  $(x, y) \in M \times M$ ,

$$|\gamma_{\lambda,\varepsilon}^{(n)}(x,y)| \leqslant \frac{C\mathcal{M}_{\varepsilon}\mathcal{M}_{\varepsilon}'}{\varepsilon^2\lambda^{1/2}} \left(\frac{A\mathcal{M}_{\varepsilon}}{\varepsilon^2\lambda}\right)^{n-1}.$$

This implies that the Neumann series defining  $\gamma_{\lambda,\varepsilon}$  converges for  $\lambda \varepsilon^2 > A\mathcal{M}_{\varepsilon}$  and for, say,  $\lambda \varepsilon^2 \geqslant 2A\mathcal{M}_{\varepsilon}$  we have

$$|\gamma_{\lambda,\varepsilon}(x,y)| \leqslant \frac{C\mathcal{M}_{\varepsilon}\mathcal{M}'_{\varepsilon}}{\varepsilon^2\lambda^{1/2}}.$$

This implies that  $\gamma_{\lambda,\varepsilon}$ , as an operator, maps  $L^1$  to  $L^\infty$ , and hence also  $L^2$  to  $L^2$ . To show that we indeed have  $G_{\lambda} = T_{\lambda,\varepsilon} - \gamma_{\lambda,\varepsilon}$  (where now  $\gamma_{\lambda,\varepsilon}$  is defined as the sum of the above convergent series), we notice that by construction we have  $(-\Delta_g + \lambda)_x (T_{\lambda,\varepsilon} - \gamma_{\lambda,\varepsilon})(x,y) = \delta_y$ , or, as operators,  $(-\Delta_g + \lambda)(T_{\lambda,\varepsilon} - \gamma_{\lambda,\varepsilon})f = f$  for all  $f \in C_0^\infty(M)$ . Since there is a unique bounded operator on  $L^2$  that satisfies this relation, we indeed have the identity  $G_{\lambda} = T_{\lambda,\varepsilon} - \gamma_{\lambda,\varepsilon}$ .

In order to prove Proposition 4.1 we need to take the structure of  $\gamma_{\lambda,\varepsilon}^{(n)}$  more precisely into account and we split

$$\gamma_{\lambda,\varepsilon}^{(n)}(x,y) = a_{\lambda,\varepsilon}^{(n)}(x,y) + b_{\lambda,\varepsilon}^{(n)}(x,y)$$

with

$$a_{\lambda,\varepsilon}^{(n)}(x,y) := \int_{d_g(z_n,y)<\varepsilon/2} dv_g(z_n) \int_{d_g(z_{n-1},z_n)<\varepsilon/2} dv_g(z_{n-1}) \cdots \int_{d_g(z_1,z_2)<\varepsilon/2} dv_g(z_1) \times T_{\lambda,\varepsilon}(x,z_1) R_{\lambda,\varepsilon}(z_1,z_2) \cdots R_{\lambda,\varepsilon}(z_n,y).$$

**Lemma 4.3.** There are A, C > 0 and for every  $n \ge 1$  and every  $\varepsilon \in (0, \rho)$  there is a function  $r_{\varepsilon}^{(n)} : [0, \infty) \times M \to \mathbb{R}$  such that for all  $x \in M$  and all  $\lambda > 0$ 

$$\partial_{\lambda} a_{\lambda,\varepsilon}^{(n)}(x,x) = -\int_{0}^{\infty} \frac{r_{\varepsilon}^{(n)}(t,x)}{(t+\lambda)^{3}} dt$$

and for all  $x \in M$ , all  $n \ge 1$  and all  $\varepsilon \in (0, \rho)$ ,  $r_{\varepsilon}^{(n)}(\cdot, x)$  is  $C^1$  on  $[0, \infty)$  and for all t > 0

$$|r_{\varepsilon}^{(n)}(t,x)| \leqslant C\mathcal{M}_{\rho/2}\mathcal{M}'_{\rho/2}\frac{\varepsilon t}{\rho^2} \left(\frac{A\mathcal{M}_{\rho/2}\varepsilon^2}{\rho^2}\right)^{n-1}.$$

*Proof.* Due to the localization of the integrals, we have

$$a_{\lambda,\varepsilon}^{(n)}(x,y) := \int_{d_g(z_n,y)<\varepsilon/2} dv_g(z_n) \int_{d_g(z_{n-1},z_n)<\varepsilon/2} dv_g(z_{n-1}) \cdots \int_{d_g(z_1,z_2)<\varepsilon/2} dv_g(z_1) \times \chi_{\lambda,\varepsilon}^{(n)}(x,z_1,\ldots,z_n,y)$$

with

$$\chi_{\lambda,\varepsilon}^{(n)}(x,z_1,\ldots,z_n,y) = \frac{e^{-\sqrt{\lambda}(d_g(x,z_1)+\cdots+d_g(z_n,y))}}{(4\pi)^{n+1}} \times \frac{U_0(x,z_1)\chi(\varepsilon^{-1}d_g(x,z_1))\Delta U_0(z_1,z_2)\cdots\Delta U_0(z_n,y)}{d_g(x,z_1)\cdots d_g(z_n,y)}.$$

We now use the representation formula (3.4) with  $\kappa$  defined by (3.5) to write

$$\partial_{\lambda} \chi_{\lambda,\varepsilon}^{(n)}(x,z_{1},\ldots,z_{n},x) = -\frac{1}{2(4\pi)^{n+1}} \int_{0}^{\infty} dt \, \frac{t^{3/2}}{(t+\lambda)^{3}} \, \kappa(\sqrt{t} \, (d_{g}(x,z_{1}) + \cdots + d_{g}(z_{n},x))) \times \\ \times \, (d_{g}(x,z_{1}) + \cdots + d_{g}(z_{n},x)) \, \frac{U_{0}(x,z_{1})\chi(\varepsilon^{-1}d_{g}(x,z_{1}))\Delta U_{0}(z_{1},z_{2}) \cdots \Delta U_{0}(z_{n},x)}{d_{g}(x,z_{1}) \cdots d_{g}(z_{n},x)}.$$

This yields the formula in the lemma with

$$r_{\varepsilon}^{(n)}(t,x) := \frac{t^{3/2}}{2(4\pi)^{n+1}} \int_{d_g(z_n,x)<\varepsilon/2} dv_g(z_n) \int_{d_g(z_{n-1},z_n)<\varepsilon/2} dv_g(z_{n-1}) \cdots \int_{d_g(z_1,z_2)<\varepsilon/2} dv_g(z_1) \times (d_g(x,z_1) + \cdots + d_g(z_n,x)) \kappa(\sqrt{t} (d_g(x,z_1) + \cdots + d_g(z_n,x))) \times \frac{U_0(x,z_1)\chi(\varepsilon^{-1}d_g(x,z_1))\Delta U_0(z_1,z_2)\cdots\Delta U_0(z_n,x)}{d_g(x,z_1)\cdots d_g(z_n,x)}.$$

In order to prove the bound in the lemma, we use the bound

$$|\kappa(r)| \leqslant Cr^{-1}$$
 for all  $r > 0$ , (4.4)

and obtain

$$|r_{\varepsilon}^{(n)}(t,x)| \leqslant \frac{Ct}{2(4\pi)^{n+1}} \int_{d_{g}(z_{n},x)<\varepsilon/2} dv_{g}(z_{n}) \int_{d_{g}(z_{n-1},z_{n})<\varepsilon/2} dv_{g}(z_{n-1}) \cdots \int_{d_{g}(z_{1},z_{2})<\varepsilon/2} dv_{g}(z_{1})$$

$$\times \frac{|U_{0}(x,z_{1})\chi(\varepsilon^{-1}d_{g}(x,z_{1}))\Delta U_{0}(z_{1},z_{2})\cdots\Delta U_{0}(z_{n},x)|}{d_{g}(x,z_{1})\cdots d_{g}(z_{n},x)}$$

$$\leqslant \frac{Ct}{2(4\pi)^{n+1}} \mathcal{M}_{\rho/2}^{n}(\rho/2)^{-2n} \int dv_{g}(z_{n}) \int dv_{g}(z_{n-1})\cdots \int dv_{g}(z_{1})$$

$$\times \frac{\mathbb{1}(d_{g}(x,z_{1})<\varepsilon)U_{0}(x,z_{1})}{d_{g}(x,z_{1})} \frac{\mathbb{1}(d_{g}(z_{1},z_{2})<\varepsilon/2)U_{0}(z_{2},z_{1})^{2}}{d_{g}(z_{1},z_{2})} \times$$

$$\times \cdots \times \frac{\mathbb{1}(d_{g}(z_{n},x)<\varepsilon/2)U_{0}(x,z_{n})^{2}}{d_{g}(z_{n},x)}.$$

We bound the right side similarly as in (4.3) using

$$\left\| \frac{\mathbb{1}(d_g(x,y) < \varepsilon/2)U_0(y,x)^2}{d_g(x,y)} \right\|_{L_y^\infty L_x^1} \leqslant C\varepsilon^2, \quad \left\| \frac{\mathbb{1}(d_g(x,y) < \varepsilon/2)U_0(y,x)}{d_g(x,y)} \right\|_{L_y^\infty L_x^2} \leqslant C\sqrt{\varepsilon}.$$

We obtain

$$|r_{\varepsilon}^{(n)}(t,x)| \leqslant C\mathcal{M}_{\rho/2}\mathcal{M}'_{\rho/2}\frac{\varepsilon t}{\rho^2} \left(\frac{A\mathcal{M}_{\rho/2}\varepsilon^2}{\rho^2}\right)^{n-1},$$

as claimed. Using that  $\kappa'$  is bounded, one can similarly show that  $r_{\varepsilon}^{(n)}(\cdot, x)$  is  $C^1$  on  $[0, \infty)$  with a bound  $|\partial_t r_{\varepsilon}^{(n)}(t, x)| \leq C_{\rho, \varepsilon} \sqrt{t} (1 + \sqrt{t}) n^2 (A_{\rho} \varepsilon^2)^{n-1}$ .

**Lemma 4.4.** There exists A > 0 and C > 0 such that for all  $x \in M$ , all  $\lambda > 0$ , all  $n \ge 1$ , and all  $\varepsilon \in (0, \rho)$  we have

$$|\partial_{\lambda} b_{\lambda,\varepsilon}^{(n)}(x,x)| \leqslant C \mathcal{M}_{\varepsilon}' \, \mathcal{M}_{\varepsilon} \, \frac{e^{-\varepsilon\sqrt{\lambda}/4}}{\varepsilon\lambda} \left(\frac{A \mathcal{M}_{\varepsilon}}{\varepsilon^{2}\lambda}\right)^{n-1}.$$

*Proof.* We first bound the integrand of  $b_{\lambda,\varepsilon}^{(n)}$  by absolute values and then bound

$$1 - \mathbb{1}(d_g(z_n, x) < \varepsilon/2, \ d_g(z_{n-1}, z_n) < \varepsilon/2, \dots, d_g(z_1, z_2) < \varepsilon/2) \leqslant \sum_{k=1}^n \mathbb{1}(d_g(z_k, z_{k+1}) \geqslant \varepsilon/2),$$

where we put  $z_{n+1} = x$  on the right side. In this way we can bound  $|\partial_{\lambda} b_{\lambda,\varepsilon}^{(n)}(x,x)|$  by a sum of n(n+1) integrals. The factor of n comes from the constraint  $\mathbb{1}(d_g(z_k,z_{k+1}) \geq \varepsilon/2)$ ,  $k=1,\ldots,n$ , and the factor n+1 comes from the product rule when differentiating with respect to  $\lambda$  the product of n+1 functions.

The factors without  $\lambda$  derivatives can be bounded as before by

$$|T_{\lambda,\varepsilon}(x,y)| \leqslant C \frac{e^{-\sqrt{\lambda}d_g(x,y)}}{d_g(x,y)} U_0(x,y) \mathbb{1}(d_g(x,y) < \varepsilon),$$
  

$$|R_{\lambda,\varepsilon}(x,y)| \leqslant C \mathcal{M}_{\varepsilon} \frac{e^{-\frac{1}{2}\sqrt{\lambda}d_g(x,y)}}{\varepsilon^2 d_g(x,y)} \min\{U_0(x,y)^2, U_0(y,x)^2\} \mathbb{1}(d_g(x,y) < \varepsilon).$$

Using the explicit expression for  $T_{\lambda,\varepsilon}$  and  $R_{\lambda,\varepsilon}$  one easily finds that

$$|\partial_{\lambda} T_{\lambda,\varepsilon}(x,y)| \leqslant C \frac{e^{-\sqrt{\lambda} d_g(x,y)}}{\sqrt{\lambda}} U_0(x,y) \mathbb{1}(d_g(x,y) < \varepsilon),$$
  

$$|\partial_{\lambda} R_{\lambda,\varepsilon}(x,y)| \leqslant C \mathcal{M}_{\varepsilon} \frac{e^{-\sqrt{\lambda} d_g(x,y)}}{\varepsilon^2 \sqrt{\lambda}} \min\{U_0(x,y)^2, U_0(y,x)^2\} \mathbb{1}(d_g(x,y) < \varepsilon).$$

We see that the effect of the derivative  $\partial_{\lambda}$  is to multiple the bounds by a factor of  $d_g/\sqrt{\lambda}$  and, due to the localization of the integrals, this can be bounded by  $\varepsilon/\sqrt{\lambda}$ . Moreover, we see that the difference between the bounds involving  $R_{\lambda,\varepsilon}$  and  $T_{\lambda,\varepsilon}$  (and their derivatives) is a factor  $\varepsilon^{-2}$  in the former. This shows that, again with the convention  $z_{n+1} = x$ ,

$$\begin{split} |\partial_{\lambda}b_{\lambda,\varepsilon}^{(n)}(x,x)| &\leqslant C(n+1)\mathcal{M}_{\varepsilon}^{n} \frac{1}{\varepsilon^{2n}} \frac{\varepsilon}{\sqrt{\lambda}} \sum_{k=1}^{n} \int dv_{g}(z_{n}) \cdots \int dv_{g}(z_{1}) \mathbb{1}(d_{g}(z_{k},z_{k+1}) \geqslant \varepsilon/2) \\ &\times \frac{e^{-\frac{1}{2}\sqrt{\lambda}d_{g}(x,z_{1})}}{d_{g}(x,z_{1})} U_{0}(x,z_{1}) \mathbb{1}(d_{g}(x,z_{1}) < \varepsilon) \\ &\times \frac{e^{-\frac{1}{2}\sqrt{\lambda}d_{g}(z_{1},z_{2})}}{d_{g}(z_{1},z_{2})} \min\{U_{0}(z_{1},z_{2})^{2}, U_{0}(z_{2},z_{1})^{2}\} \mathbb{1}(d_{g}(z_{1},z_{2}) < \varepsilon) \cdots \\ &\times \frac{e^{-\frac{1}{2}\sqrt{\lambda}d_{g}(z_{n},z_{n+1})}}{d_{g}(z_{n},z_{n+1})} \min\{U_{0}(z_{n},z_{n+1})^{2}, U_{0}(z_{n+1},z_{n})^{2}\} \mathbb{1}(d_{g}(z_{n},z_{n+1}) < \varepsilon). \end{split}$$

For all  $1 \leq k \leq n$  we bound

$$\left| \int dv_g(z_1) \cdots \int dv_g(z_n) K_0(x, z_1) K_1(z_1, z_2) \cdots K_n(z_n, y) \right|$$

$$\leq \|K_0\|_{L_x^{\infty} L_y^2} \left( \prod_{j=1}^{k-1} \|K_j\|_{L_x^{\infty} L_y^1}^{1/2} \|K_j\|_{L_y^{\infty} L_x^1}^{1/2} \right) \|K_k\|_{L_y^{\infty} L_x^2} \left( \prod_{\ell=k+1}^{n} \|K_{\ell}\|_{L_y^{\infty} L_x^1} \right),$$

which is proved by factorizing the kernel as

$$\left(K_0 K_1^{1/2} \cdots K_{k-1}^{1/2} K_{k+1}^{1/2} \cdots K_n^{1/2}\right) \times \left(K_1^{1/2} \cdots K_{k-1}^{1/2} K_k K_{k+1}^{1/2} \cdots K_n^{1/2}\right)$$

(suppressing the arguments for simplicity) and applying the Cauchy–Schwarz inequality. This bound allows us to always put the factor involving the pair  $(x, z_1)$  (which only has a single  $U_0$ ) and the factor involving  $(z_k, z_{k+1})$  (which has the additional cut-off away from the diagonal) into  $L^2$ . Using (4.1), (4.2) together with

$$\left\| \frac{e^{-\frac{1}{2}\sqrt{\lambda}d_g(x,y)}}{d_g(x,y)} \min\{U_0(x,y)^2, U_0(y,x)\}^2 \mathbb{1}(\varepsilon/2 \leqslant d_g(x,y) < \varepsilon) \right\|_{L_y^{\infty}L_x^2} \leqslant \frac{C\mathcal{M}'_{\varepsilon}}{\lambda^{1/4}} e^{-\frac{1}{4}\varepsilon\sqrt{\lambda}},$$

yields the bound

$$|\partial_{\lambda}b_{\lambda,\varepsilon}^{(n)}(x,x)| \leqslant Cn(n+1)\mathcal{M}_{\varepsilon}'\mathcal{M}_{\varepsilon}\frac{e^{-\varepsilon\sqrt{\lambda}/4}}{\varepsilon\lambda}\left(\frac{A\mathcal{M}_{\varepsilon}}{\varepsilon^{2}\lambda}\right)^{n-1}.$$

The term n(n+1) here can be dropped by increasing the absolute constant A. This completes the proof of the lemma.

Proof of Proposition 4.1. With  $r_{\varepsilon}^{(n)}$  from Lemma 4.3 let  $R_{\varepsilon}(t,x) = \sum_{n\geqslant 1} (-1)^{n+1} r_{\varepsilon}^{(n)}(t,x)$ . According to that lemma, if we choose  $\varepsilon = \varepsilon_0 = \rho/\sqrt{2A\mathcal{M}_{\rho/2}}$ , then the series defining  $R_{\varepsilon}(t,x)$  converges and for all  $x\in M$ ,  $R_{\varepsilon}(\cdot,x)$  is  $C^1$  on  $[0,\infty)$  and for all t>0,

$$|R_{\varepsilon}(t,x)| \leqslant C \,\mathcal{M}_{\rho/2}^{1/2} \,\mathcal{M}_{\rho/2}' \,\frac{t}{\rho}.$$

We may assume that the absolute constant in the definition of  $\varepsilon$  satisfies  $A \ge 2$ . This implies that  $\varepsilon_0 \le \rho/2$ , since  $\mathcal{M}_{\varepsilon} \ge 1$ , which in turn is a consequence of  $U_0(x,x) = 1$ .

This bound on  $R_{\varepsilon}$  and the bound from Lemma 4.4 imply that  $\gamma_{\lambda,\varepsilon}$  can be differentiated with respect to  $\lambda$ , that this derivative is given by differentiating its series expansion termwise

and that for all  $\lambda \geqslant 2A'\mathcal{M}_{\rho/2}/\varepsilon^2$ 

$$\left| \partial_{\lambda} \gamma_{\lambda,\varepsilon}(x,x) + \int_{0}^{\infty} \frac{R_{\varepsilon}(t,x)}{(t+\lambda)^{3}} dt \right| = \left| \sum_{n\geqslant 1} (-1)^{n+1} \left( \gamma_{\lambda,\varepsilon}^{(n)}(x,x) - a_{\lambda,\varepsilon}^{(n)}(x,x) \right) \right|$$

$$= \left| \sum_{n\geqslant 1} (-1)^{n+1} b_{\lambda,\varepsilon}^{(n)}(x,x) \right|$$

$$\leqslant C \mathcal{M}_{\varepsilon} \, \mathcal{M}'_{\varepsilon} \, \frac{e^{-\varepsilon\sqrt{\lambda}/4}}{\varepsilon\lambda}$$

$$\leqslant C \mathcal{M}_{\rho/2}^{3/2} \, \mathcal{M}'_{\rho/2} \frac{e^{-\varepsilon\sqrt{\lambda}/4}}{\rho\lambda},$$

where in the last line we used the fact that  $\mathcal{M}_{\varepsilon}$  and  $\mathcal{M}'_{\varepsilon}$  are nondecreasing in  $\varepsilon$  and the choice of  $\varepsilon$ . On the other hand, by definition of  $\gamma_{\lambda,\varepsilon}$  and a computation of  $\partial_{\lambda}T_{\lambda,\varepsilon}(x,x)$  we have

$$\partial_{\lambda}G_{\lambda}(x,x) = \partial_{\lambda}T_{\lambda,\varepsilon}(x,x) - \partial_{\lambda}\gamma_{\lambda,\varepsilon}(x,x) = -\frac{1}{8\pi\sqrt{\lambda}} - \partial_{\lambda}\gamma_{\lambda,\varepsilon}(x,x).$$

Combining this identity with the bound on  $\partial_{\lambda}\gamma_{\lambda,\varepsilon}$ , we obtain the bound in the proposition.

Remark 4.5. The above proof is specific to three space dimensions because of the particular form of the resolvent kernel of the Laplacian in three dimensions,

$$(-\Delta_{\mathbb{R}^3} + \lambda)^{-1}(x,y) = \frac{e^{-\sqrt{\lambda}|x-y|}}{4\pi|x-y|}.$$

In particular, we use in a crucial way that  $e^{\sqrt{\lambda}|x-y|}(-\Delta_{\mathbb{R}^3}+\lambda)^{-1}(x,y)$  is independent of  $\lambda$ , so that the terms  $a^{(n)}$  defined above can be written as exact Stieltjes transforms. This fact is not true anymore in other dimensions: for instance, in dimension five one has

$$(-\Delta_{\mathbb{R}^5} + \lambda)^{-1}(x, y) = \frac{e^{-\sqrt{\lambda}|x-y|}}{8\pi^2|x-y|^3} \left(1 + \sqrt{\lambda}|x-y|\right)$$

and it is not clear how to deal with the term  $1 + \sqrt{\lambda}|x - y|$  on the right. In the footnote on p. 328 of [4], Avakumović mentions that in higher dimensions, one should use the heat kernel instead of Green's functions, but without any further details. We were not able to reconstruct his argument in higher dimensions.

Remark 4.6 (Regularity needed on g). The previous arguments shows that we need uniform bounds on  $\Delta U_0$ , so, in particular, the assumption  $U_0 \in C^2$  suffices. By the definition of  $U_0$ , this means that the exponential map is  $C^3$ , and since the exponential map may be seen as a time-one flow of an ODE whose vector field depends on the Christoffel symbols (that involve one derivative of g), we see that overall this argument requires  $g \in C^4$ . It is interesting to compare this regularity with the one found in [17, 55] for the (weaker)  $L^{\infty}$  bounds for eigenfunctions or quasimodes, where they only need essentially  $C^2$  regularity (which is the minimal regularity needed for geodesics to exist). The additional two derivatives needed here can be explained by the choice of the parametrix. For  $L^{\infty}$  bounds of eigenfunctions,

one may replace the function  $U_0$  by 1 in the parametrix  $T_{\lambda,\varepsilon}$  above, which gives bounds of the same order for  $R_{\lambda,\varepsilon}$  (and hence no information on  $\Delta U_0$  is indeed). We, however, need more than mere bounds for  $R_{\lambda,\varepsilon}$ , we need that  $e^{\sqrt{\lambda}d_g(x,y)}R_{\lambda,\varepsilon}(x,y)$  is independent of  $\lambda$  for  $d_g(x,y) < \varepsilon/2$ , which motivates the introduction of  $U_0$  in the parametrix.

# 5. Adding singular potentials

The previous construction is quite robust and allows the inclusion of singular potentials. Indeed, let  $V: M \to \mathbb{R}$  belonging to the *Kato class* defined as the set of all measurable V such that

$$\lim_{r \to 0_+} \|V\|_{\mathcal{K}(r)} = 0,$$

where

$$||V||_{\mathcal{K}(r)} = \sup_{x \in M} \int_{d_g(x,y) < r} \frac{|V(y)|}{d_g(x,y)} dv_g(y).$$

Then  $-\Delta_g + V$  defines a bounded below quadratic form, as we recall below in Appendix A. Define for  $\lambda > -\inf \operatorname{spec}(-\Delta_g + V)$  and  $x, y \in M$ ,

$$G_{\lambda}^{V}(x,y) := (-\Delta_{q} + V + \lambda)^{-1}(x,y).$$

**Proposition 5.1.** There are constants  $C_1, \ldots, C_6 \geqslant 1$  and for every Kato class  $V: M \to \mathbb{R}$  there is a constant  $\varepsilon_0 > 0$  such that for all  $\varepsilon \in (0, \varepsilon_0]$  there are functions  $R_{\varepsilon}^V, r_{0,\varepsilon}^V: (0, \infty) \times M \to \mathbb{R}$  with the following properties. For all  $x \in M$ , all  $\lambda \geqslant C_1/\varepsilon^2$ , and all  $\varepsilon \in (0, \varepsilon_0]$ 

$$\left| \partial_{\lambda} G_{\lambda}^{V}(x,x) + \frac{1}{8\pi\sqrt{\lambda}} - \int_{0}^{\infty} \frac{t^{3/2} r_{0,\varepsilon}^{V}(t,x) + R_{\varepsilon}^{V}(t,x)}{(t+\lambda)^{3}} dt \right| \leqslant C_{2} \frac{e^{-\varepsilon\sqrt{\lambda}/4}}{\varepsilon\lambda},$$

and for all  $x \in M$ , all  $\varepsilon \in (0, \varepsilon_0]$ , all t > 0,

$$\left| R_{\varepsilon}^{V}(t,x) \right| \leqslant C_{3} \rho^{-2} \varepsilon t, \qquad \left| r_{0,\varepsilon}^{V}(t,x) \right| \leqslant C_{4} \|V\|_{\mathcal{K}(\varepsilon)}.$$

Moreover, for all  $x \in M$ , all  $0 < t \le t'$ , and all  $\varepsilon \in (0, \varepsilon_0]$ ,

$$\left| r_{0,\varepsilon}^{V}(t,x) - r_{0,\varepsilon}^{V}(t',x) \right| \leqslant C_5 \|V\|_{\mathcal{K}(\varepsilon)} \frac{\sqrt{t'} - \sqrt{t}}{\sqrt{t}}. \tag{5.1}$$

$$\left| t^{-3/2} R_{\varepsilon}^{V}(t,x) - (t')^{-3/2} R_{\varepsilon}^{V}(t',x) \right| \leqslant C_5 \frac{\varepsilon^2}{\rho^2} \frac{\sqrt{t'} - \sqrt{t}}{\sqrt{t}}. \tag{5.2}$$

The function  $r_{0,\varepsilon}^V$  is explicitly given by (5.11) and (5.8). The constant  $\varepsilon_0$  satisfies

$$C_6 \|V\|_{\mathcal{K}(\varepsilon_0)} \leqslant 1$$
,  $C_6 \varepsilon_0 \leqslant \rho$ ,

and the constants  $C_1, \ldots, C_6$  depend only on  $\mathcal{M}_{\rho/2}$  and  $\mathcal{M}'_{\rho/2}$ .

Before proving this proposition let us derive the following important consequence which is the technical main result of this paper.

**Corollary 5.2.** Let  $V: M \to \mathbb{R}$  be in the Kato class and let  $\varepsilon_0 > 0$  be as in Proposition 5.1. Then, for all  $\varepsilon \in (0, \varepsilon_0]$  we have, as  $\lambda \to \infty$ , uniformly in  $x \in M$ ,

$$\mathbb{1}(-\Delta_g + V \leqslant t)(x, x) = \frac{t^{3/2}}{6\pi^2} - \frac{1}{2} r_{0,\varepsilon}^V(t, x) t^{3/2} + \mathcal{O}^{(\varepsilon)}(t), \quad t \to +\infty$$
 (5.3)

where the  $\mathcal{O}^{(\varepsilon)}$  is uniform in  $x \in M$  (but depends on  $\varepsilon$ ). More precisely, we have for all  $x \in M$  and all  $t \ge 0$ ,

$$\left| \mathbb{1}(-\Delta_g + V \leqslant t)(x, x) - \frac{t^{3/2}}{6\pi^2} + \frac{1}{2} r_{0,\varepsilon}^V(t, x) t^{3/2} \right| \leqslant C_0 \varepsilon^{-1} (t + \varepsilon^{-2}),$$

where  $C_0$  depends only on upper bounds on  $\mathcal{M}_{\rho/2}$  and  $\mathcal{M}'_{\rho/2}$ .

Proof of Corollary 5.2 assuming Proposition 5.1. Define for all  $t \ge 0$ 

$$A(t,x) := \mathbb{1}(-\Delta_q + V \leq t)(x,x) - t^{3/2}/(6\pi^2) + r_{0,\varepsilon}^V(t,x) t^{3/2}/2 + R_{\varepsilon}^V(t,x)/2$$
.

As in the proof of Corollary 4.2, by the spectral theorem for the operator  $-\Delta_g + V$  we find

$$\int_0^\infty \frac{\mathbb{1}(-\Delta_g + V \leqslant t)(x, x)}{(\lambda + t)^3} dt = -\frac{1}{2} \,\partial_\lambda G_\lambda^V(x, x) \,.$$

Continuing to argue as in that proof, we have by Proposition 5.1,

$$\left| \int_0^\infty \frac{A(t,x)}{(t+\lambda)^3} dt \right| = \left| -\frac{1}{2} \partial_\lambda G_\lambda(x,x) - \frac{1}{16\pi\sqrt{\lambda}} + \frac{1}{2} \int_0^\infty \frac{t^{3/2} r_{0,\varepsilon}^V(t,x) + R_\varepsilon^V(t,x)}{(t+\lambda)^3} dt \right|$$

$$\leqslant C_2 \frac{e^{-\varepsilon\sqrt{\lambda}/4}}{\varepsilon\lambda}$$

for all  $\lambda \geq C_1/\varepsilon^2$ . The result then follows from Theorem 3.1 in the simplified form (3.1) with  $B_0 \equiv -(6\pi)^{-1} + r_{0,\varepsilon}^V(t,x)/2$ ,  $B_1(t) = \mathbb{1}(-\Delta_g + V \leq t)(x,x)$  and  $B_2(t) = R_{\varepsilon}^V(t,x)/2$ . Note that the assumptions on  $B_0$  and  $B_2$  required for Theorem 3.1 are satisfied by the bounds in Proposition 5.1. For instance, the bounds (5.1) and (5.2) imply that  $u \mapsto u^3 r_{0,\varepsilon}^V(u^2,x)$  and  $u \mapsto R_{\varepsilon}^V(u^2,x)$  are locally Lipschitz on  $[0,\infty)$ , and hence continuous and locally of bounded variation on  $[0,\infty)$ . Moreover, inequality (5.1) implies that the almost monotonicity assumption on  $B_0$  in Theorem 3.1 is satisfied for any  $\delta \geq 0$  with  $C_{B_0} \leq C_5 ||V||_{\mathcal{K}(\varepsilon)} \delta$ . We apply Theorem 3.1 for any  $\varepsilon \in (0,\varepsilon_0]$  with the choice  $\delta = \varepsilon^{-1}$ . In view of the bounds in Proposition 5.1 this gives

$$\left| \mathbb{1}(-\Delta_g + V \leqslant t)(x, x) - \frac{t^{3/2}}{6\pi^2} + \frac{1}{2} r_{0,\varepsilon}^V(t, x) t^{3/2} \right| \leqslant C_0 \left( 1 + \varepsilon^3 B_1(0) \right) \varepsilon^{-1} \left( t + \varepsilon^{-2} \right),$$

where  $C_0$  depends only on upper bounds on  $\mathcal{M}_{\rho/2}$  and  $\mathcal{M}'_{\rho/2}$ . To complete the proof of the corollary, it suffices to show that

$$B_1(0) = \mathbb{1}(-\Delta + V \le 0)(x, x) \le C\varepsilon_0^{-3}$$
. (5.4)

Since each negative eigenvalue  $E_n$  of  $-\Delta_g + V$  satisfies  $-\lambda_0 < E_n \le 0$  with  $\lambda_0 := 2C_1/\varepsilon_0^2$  (indeed, since the resolvent was shown to exist for all  $\lambda \ge C_1/\varepsilon_0^2$ , this means in particular that the spectrum of  $-\Delta_g + V$  is above  $-C_1/\varepsilon_0^2$ ), we have

$$\lambda_0^{-2} \mathbb{1}(-\Delta + V \leq 0)(x, x) \leq (-\Delta_g + V + \lambda_0)^{-2}(x, x) = -\partial_\lambda G_{\lambda_0}^V(x, x)$$
.

On the other hand, Proposition 5.1 implies that

$$-\partial_{\lambda} G_{\lambda_0}^{V}(x,x) \leqslant \frac{1}{8\pi\sqrt{\lambda_0}} + \int_0^{\infty} \frac{t^{3/2} C_4 \|V\|_{\mathcal{K}(\varepsilon)} + C_3 \rho^{-2} \varepsilon_0 t}{(t+\lambda_0)^3} dt + C_2 \frac{e^{-\varepsilon_0\sqrt{\lambda_0}/4}}{\varepsilon_0\lambda_0} \leqslant C\lambda_0^{-1/2}$$

with a constant C depending only on upper bounds on  $\mathcal{M}_{\rho/2}$  and  $\mathcal{M}'_{\rho/2}$ . Multiplying by  $\lambda_0^2$  and recalling the definition of  $\lambda_0$ , we obtain (5.4). This completes the proof of the corollary.

5.1. **Proof of Proposition 5.1.** We will see that the free parametrix  $T_{\lambda,\varepsilon}$  is still a good parametrix for  $G_{\lambda}^{V}$ . With  $T_{\lambda,\varepsilon}$  and  $R_{\lambda,\varepsilon}$  defined in the previous section, depending on a parameter  $\varepsilon \in (0, \rho]$  to be determined, we set

$$\gamma_{\lambda,\varepsilon}^V := T_{\lambda,\varepsilon} - G_{\lambda}^V$$

and

$$R_{\lambda,\varepsilon}^V := (-\Delta_q + V + \lambda)\gamma_{\lambda,\varepsilon}^V = R_{\lambda,\varepsilon} + VT_{\lambda,\varepsilon}$$
.

We thus have the integral representation

$$\gamma_{\lambda,\varepsilon}^{V}(x,y) = \int_{M} T_{\lambda,\varepsilon}(x,z) R_{\lambda,\varepsilon}^{V}(z,y) \, dv_{g}(z) - \int_{M} \gamma_{\lambda,\varepsilon}^{V}(x,z) R_{\lambda,\varepsilon}^{V}(z,y) \, dv_{g}(z)$$

and, at least formally, the series representation

$$\gamma_{\lambda,\varepsilon}^{V}(x,y) = \sum_{n\geqslant 1} (-1)^{n+1} \gamma_{\lambda,\varepsilon}^{(n,V)}(x,y)$$

with

$$\gamma_{\lambda,\varepsilon}^{(n,V)}(x,y) := \int_M dv_g(z_1) \cdots \int_M dv_g(z_n) T_{\lambda,\varepsilon}(x,z_1) R_{\lambda,\varepsilon}^V(z_1,z_2) \cdots R_{\lambda,\varepsilon}^V(z_n,y) .$$

When trying to prove pointwise bounds on  $\gamma_{\lambda,\varepsilon}^{(n,V)}$  by the same method as in the previous section, one runs into the problem that, while there are sufficiently good bounds on  $\|VT_{\lambda,\varepsilon}\|_{L_y^\infty L_x^1}$ , see (5.6) below, the norm  $\|VT_{\lambda,\varepsilon}\|_{L_y^\infty L_x^2}$  might be infinite for Kato class potentials. One can, however, obtain integrated bounds since  $T_{\lambda,\varepsilon} \in L_y^\infty L_x^2$ , showing that the series defining  $\gamma_{\lambda,\varepsilon}^V$  converges in  $L_y^\infty L_x^2$  meaning that  $\gamma_{\lambda,\varepsilon}^V$  maps  $L^2$  to  $L^\infty$  and hence  $L^2$  to  $L^2$ . Again by uniqueness of the resolvent, this shows the identity  $G_\lambda^V = T_{\lambda,\varepsilon} - \gamma_{\lambda,\varepsilon}^V$ , where  $\gamma_{\lambda,\varepsilon}^V$  is defined as the sum of the above convergent series.

The way around the impasse of obtaining a pointwise convergent series is to extract the most singular term from  $\gamma_{\lambda,\varepsilon}^{(n,V)}$ . Note that, since  $R_{\lambda,\varepsilon}^V$  is a sum of two terms, namely,  $R_{\lambda,\varepsilon}$  and  $VT_{\lambda,\varepsilon}$ , the quantity  $\gamma_{\lambda,\varepsilon}^{(n,V)}$  can be written as a sum of  $2^n$  terms. The most singular term is the one where all factors are  $VT_{\lambda,\varepsilon}$ , that is,

$$\sigma_{\lambda,\varepsilon}^{(n,V)}(x,y) := \int_M dv_g(z_1) \cdots \int_M dv_g(z_n) T_{\lambda,\varepsilon}(x,z_1) V(z_1) T_{\lambda,\varepsilon}(z_1,z_2) \cdots V(z_n) T_{\lambda,\varepsilon}(z_n,y) .$$

Before discussing this singular term, let us derive bounds on the difference  $\gamma_{\lambda,\varepsilon}^{(n,V)} - \sigma_{\lambda,\varepsilon}^{(n,V)}$  that show, in particular, that the series  $\sum_{n\geqslant 1} (-1)^{n+1} \left(\gamma_{\lambda,\varepsilon}^{(n,V)} - \sigma_{\lambda,\varepsilon}^{(n,V)}\right)$  converges. We write

symbolically (identifying the kernels with operators)

$$\gamma_{\lambda,\varepsilon}^{(n,V)} - \sigma_{\lambda,\varepsilon}^{(n,V)} = \sum_{k=1}^{n} T_{\lambda,\varepsilon} (VT_{\lambda,\varepsilon})^{k-1} R_{\lambda,\varepsilon} (R_{\lambda,\varepsilon}^{V})^{n-k} = \sum_{k=1}^{n} (T_{\lambda,\varepsilon}V)^{k-1} T_{\lambda,\varepsilon} R_{\lambda,\varepsilon} (R_{\lambda,\varepsilon}^{V})^{n-k}.$$

Moreover, by using the Cauchy–Schwarz inequality in the integral connecting  $T_{\lambda,\varepsilon}$  and  $R_{\lambda,\varepsilon}$  we find that

$$\left\| (T_{\lambda,\varepsilon}V)^{k-1} T_{\lambda,\varepsilon} R_{\lambda,\varepsilon} (R_{\lambda,\varepsilon}^V)^{n-k} \right\|_{L_x^{\infty} L_y^{\infty}} \leq \|T_{\lambda,\varepsilon}V\|_{L_x^{\infty} L_y^{1}}^{k-1} \|T_{\lambda,\varepsilon}\|_{L_x^{\infty} L_y^{2}} \|R_{\lambda,\varepsilon}\|_{L_y^{\infty} L_x^{2}}^{V} \|R_{\lambda,\varepsilon}^V\|_{L_x^{\infty} L_x^{1}}^{n-k}.$$
(5.5)

Combining this inequality with the bounds (4.2), (4.1) and the obvious bounds

$$||VT_{\lambda,\varepsilon}||_{L_{y}^{\infty}L_{x}^{1}} \leqslant C\mathcal{M}_{\varepsilon}'||V||_{\mathcal{K}(\varepsilon)}$$
 and  $||T_{\lambda,\varepsilon}V||_{L_{x}^{\infty}L_{y}^{1}} \leqslant C\mathcal{M}_{\varepsilon}'||V||_{\mathcal{K}(\varepsilon)},$  (5.6)

we obtain

$$\left| \gamma_{\lambda,\varepsilon}^{(n,V)}(x,y) - \sigma_{\lambda,\varepsilon}^{(n,V)}(x,y) \right| \leqslant C \frac{\mathcal{M}_{\varepsilon}\mathcal{M}_{\varepsilon}'}{\varepsilon^{2}\lambda^{1/2}} \left( A \left( \mathcal{M}_{\varepsilon}' \|V\|_{\mathcal{K}(\varepsilon)} + \frac{\mathcal{M}_{\varepsilon}}{\varepsilon^{2}\lambda} \right) \right)^{n-1}.$$

(Similarly as in the previous section, a factor of n can be dropped by enlarging the absolute constant A.) This is the desired bound on  $\gamma_{\lambda,\varepsilon}^{(n,V)}(x,y) - \sigma_{\lambda,\varepsilon}^{(n,V)}(x,y)$ .

To proceed, we decompose

$$\gamma_{\lambda,\varepsilon}^{(n,V)}(x,y) - \sigma_{\lambda,\varepsilon}^{(n,V)}(x,y) = a_{\lambda,\varepsilon}^{(n,V)}(x,y) + b_{\lambda,\varepsilon}^{(n,V)}(x,y), \qquad (5.7)$$

where  $a_{\lambda,\varepsilon}^{(n,V)}$  and  $b_{\lambda,\varepsilon}^{(n,V)}$  are defined in the same way as in the previous sections, namely, in  $a_{\lambda,\varepsilon}^{(n,V)}$  all n integrations with respect to the variables  $z_k$ ,  $k=1,\ldots,n$ , are restricted to  $d_g(z_k,z_{k+1})<\varepsilon/2$ , with  $z_{n+1}=y$ . The following two lemmas are the analogues of Lemmas 4.3 and 4.4.

**Lemma 5.3.** There are A, C > 0 and for every  $n \ge 1$  and  $\varepsilon \in (0, \rho]$  there is a function  $r_{\varepsilon}^{(n,V)}: (0,\infty) \times M \to \mathbb{R}$  such that for all  $x \in M$ 

$$\partial_{\lambda} a_{\lambda,\varepsilon}^{(n,V)}(x,x) = -\int_{0}^{\infty} \frac{r_{\varepsilon}^{(n,V)}(t,x)}{(t+\lambda)^{3}} dt$$

and for all  $x \in M$ , all t > 0, all  $n \ge 1$  and all  $\varepsilon \in (0, \rho]$ 

$$|r_{\varepsilon}^{(n,V)}(t,x)| \leqslant C \mathcal{M}_{\rho/2} \mathcal{M}_{\varepsilon}' \frac{\varepsilon t}{\rho^2} \left( A \left( \mathcal{M}_{\varepsilon}' ||V||_{\mathcal{K}(\varepsilon)} + \frac{\mathcal{M}_{\rho/2} \varepsilon^2}{\rho^2} \right) \right)^{n-1}.$$

*Proof.* The proof is similar to that of Lemma 4.3 and we only sketch the main steps. Decomposing the integrand similarly as in the proof of the bound on  $\gamma_{\lambda,\varepsilon}^{(n,V)} - \sigma_{\lambda,\varepsilon}^{(n,V)}$  and using the same representation formula as in the proof of Lemma 4.3 we obtain the claimed formula

with

$$r_{\varepsilon}^{(n,V)}(t,x) := \frac{t^{3/2}}{2(4\pi)^{n+1}} \sum_{k=1}^{n} \int_{d_{g}(z_{n},x)<\varepsilon/2} dv_{g}(z_{n}) \int_{d_{g}(z_{n-1},z_{n})<\varepsilon/2} dv_{g}(z_{n-1}) \cdots \int_{d_{g}(z_{1},z_{2})<\varepsilon/2} dv_{g}(z_{1})$$

$$\times (d_{g}(x,z_{1}) + \cdots + d_{g}(z_{n},x)) \kappa(\sqrt{t} (d_{g}(x,z_{1}) + \cdots + d_{g}(z_{n},x)))$$

$$\times \frac{U_{0}(x,z_{1})\chi(\varepsilon^{-1}d_{g}(x,z_{1}))}{d_{g}(x,z_{1})} \left( \prod_{\ell=1}^{k-1} V(z_{\ell}) \frac{U_{0}(z_{\ell},z_{\ell+1})}{d_{g}(z_{\ell},z_{\ell+1})} \right) \frac{\Delta U_{0}(z_{k},z_{k+1})}{d_{g}(z_{k},z_{k+1})}$$

$$\times \left( \prod_{m=k+1}^{n} \frac{\Delta U_{0}(z_{m},z_{m+1}) + V(z_{m})U_{0}(z_{m},z_{m+1})}{d_{g}(z_{m},z_{m+1})} \right).$$

Here we use the convention  $z_{n+1} = x$ . Using (4.4) we obtain

$$\left| r_{\varepsilon}^{(n,V)}(t,x) \right| \leqslant \frac{Ct}{2(4\pi)^{n+1}} \sum_{k=1}^{n} \int_{d_{g}(z_{n},x) < \varepsilon/2} dv_{g}(z_{n}) \int_{d_{g}(z_{n-1},z_{n}) < \varepsilon/2} dv_{g}(z_{n-1}) \cdots \int_{d_{g}(z_{1},z_{2}) < \varepsilon/2} dv_{g}(z_{1}) \right| \\
\times \frac{U_{0}(x,z_{1}) \mathbb{1}(d_{g}(x,z_{1}) < \varepsilon)}{d_{g}(x,z_{1})} \left( \prod_{\ell=1}^{k-1} |V(z_{\ell})| \frac{U_{0}(z_{\ell},z_{\ell+1})}{d_{g}(z_{\ell},z_{\ell+1})} \right) \frac{|\Delta U_{0}(z_{k},z_{k+1})|}{d_{g}(z_{k},z_{k+1})} \\
\times \left( \prod_{m=k+1}^{n} \frac{|\Delta U_{0}(z_{m},z_{m+1})| + |V(z_{m})|U_{0}(z_{m},z_{m+1})}{d_{g}(z_{m},z_{m+1})} \right).$$

Using the bound (5.5) we can estimate the k-th summand on the right side by

$$\begin{split} & \left\| \frac{U_0(x,y)\mathbb{1}(d_g(x,y) < \varepsilon)}{d_g(x,y)} V(y) \right\|_{L_x^{\infty} L_y^1}^{k-1} \\ & \times \left\| \frac{U_0(x,y)\mathbb{1}(d_g(x,y) < \varepsilon/2)}{d_g(x,y)} \right\|_{L_x^{\infty} L_y^2} \frac{\mathcal{M}_{\rho/2}}{(\rho/2)^2} \left\| \frac{U_0(y,x)^2 \mathbb{1}(d_g(x,y) < \varepsilon/2)}{d_g(x,y)} \right\|_{L_y^{\infty} L_x^2} \\ & \times \left( \frac{\mathcal{M}_{\rho/2}}{(\rho/2)^2} \left\| \frac{U_0(y,x)^2 \mathbb{1}(d_g(x,y) < \varepsilon/2)}{d_g(x,y)} \right\|_{L_y^{\infty} L_x^1} + \left\| V(x) \frac{U_0(x,y) \mathbb{1}(d_g(x,y) < \varepsilon/2)}{d_g(x,y)} \right\|_{L_y^{\infty} L_x^1} \right)^{n-k}. \end{split}$$

Using similar bounds as before one sees that this is bounded by a constant times

$$(A\mathcal{M}'_{\varepsilon}||V||_{\mathcal{K}(\varepsilon)})^{k-1} \sqrt{\varepsilon} \frac{\mathcal{M}_{\rho/2}\mathcal{M}'_{\varepsilon} \sqrt{\varepsilon}}{\rho^{2}} \left( A \left( \frac{\mathcal{M}_{\rho/2} \varepsilon^{2}}{\rho^{2}} + \mathcal{M}'_{\varepsilon}||V||_{\mathcal{K}(\varepsilon)} \right) \right)^{n-k}.$$

This leads to the claimed bound.

**Lemma 5.4.** There are A, C > 0 such that for all  $x \in M$ , all  $\lambda > 0$ , all  $n \ge 1$ , and all  $\varepsilon \in (0, \rho]$  we have

$$\left| \partial_{\lambda} b_{\lambda,\varepsilon}^{(n,V)}(x,x) \right| \leqslant C \mathcal{M}_{\varepsilon}' \mathcal{M}_{\varepsilon} \frac{e^{-\varepsilon\sqrt{\lambda}/4}}{\varepsilon\lambda} \left( A \left( \mathcal{M}_{\varepsilon}' \|V\|_{\mathcal{K}(\varepsilon)} + \frac{\mathcal{M}_{\varepsilon}}{\varepsilon^{2}\lambda} \right) \right)^{n-1}.$$

*Proof.* We recall that the integrand of  $\gamma_{\lambda,\varepsilon}^{(n,V)} - \sigma_{\lambda,\varepsilon}^{(n,V)}$  can be written symbolically as a sum of terms of the form  $(T_{\lambda,\varepsilon}V)^{k-1}T_{\lambda,\varepsilon}R_{\lambda,\varepsilon}(R_{\lambda,\varepsilon}^V)^{n-k}$  for some  $1 \leqslant k \leqslant n$ , which will be fixed from now on. When taking the  $\lambda$  derivative, by the product rule it can fall on any one of the

n+1 terms. Next, as in the proof of Lemma 4.4, we first take absolute values of the integral kernels and then multiply them by  $\mathbb{1}(d_g(z_\ell, z_{\ell+1}) \geqslant \varepsilon/2)$  for some  $1 \leqslant \ell \leqslant n$ , which will also be fixed from now on. Thus,  $\partial_{\lambda} b_{\lambda,\varepsilon}^{(n,V)}(x,x)$  can be bounded by a sum of  $n^2(n+1)$  terms.

Each of the terms  $T_{\lambda,\varepsilon}$ ,  $R_{\lambda,\varepsilon}$ ,  $\partial_{\lambda}R_{\lambda,\varepsilon}$  and  $\partial_{\lambda}T_{\lambda,\varepsilon}$  can be bounded as in the proof of Lemma 4.4 and we see again that the effect of the  $\lambda$  derivative can be bounded by multiplying the bound without this derivative by a factor of  $\varepsilon/\sqrt{\lambda}$ . Our task is thus to bound

$$\frac{\varepsilon}{\sqrt{\lambda}} \int_{d_{g}(z_{n},x)<\varepsilon} dv_{g}(z_{n}) \int_{d_{g}(z_{n-1},z_{n})<\varepsilon} dv_{g}(z_{n-1}) \cdots \int_{d_{g}(z_{1},z_{2})<\varepsilon} dv_{g}(z_{1}) \, \mathbb{1}(d_{g}(z_{\ell},z_{\ell+1}) \geqslant \varepsilon/2) 
\times \, \mathbb{1}(d_{g}(x,z_{1})<\varepsilon) \left( \prod_{m=0}^{k-2} \frac{e^{-\sqrt{\lambda}d_{g}(z_{m},z_{m+1})}}{d_{g}(z_{m},z_{m+1})} U_{0}(z_{m},z_{m+1}) |V(z_{m+1})| \right) 
\times \frac{e^{-\sqrt{\lambda}d_{g}(z_{k-1},z_{k})}}{d_{g}(z_{k-1},z_{k})} U_{0}(z_{k-1},z_{k}) \frac{\mathcal{M}_{\varepsilon}}{\varepsilon^{2}} \frac{e^{-\frac{1}{2}\sqrt{\lambda}d_{g}(z_{k},z_{k+1})}}{d_{g}(z_{k},z_{k+1})} U_{0}(z_{k+1},z_{k})^{2} 
\times \prod_{m'=k+1}^{n} \left( \frac{\mathcal{M}_{\varepsilon}}{\varepsilon^{2}} \frac{e^{-\frac{1}{2}\sqrt{\lambda}d_{g}(z_{m'},z_{m'+1})}}{d_{g}(z_{m'},z_{m'+1})} U_{0}(z_{m'+1},z_{m'})^{2} + |V(z_{m'})| \frac{e^{-\sqrt{\lambda}d_{g}(z_{m'},z_{m'+1})}}{d_{g}(z_{m'},z_{m'+1})} U_{0}(z_{m'},z_{m'+1}) \right),$$

with the convention  $z_0 = x = z_{n+1}$ . As in (5.5) we apply the Cauchy–Schwarz inequality in the  $z_k$  variable and obtain

$$\frac{\varepsilon}{\sqrt{\lambda}} \mathcal{F} \left\| \frac{e^{-\sqrt{\lambda}d_g(x,y)}}{d_g(x,y)} U_0(x,y) \mathbb{1}(d_g(x,y) < \varepsilon) V(y) \right\|_{L_x^{\infty} L_y^1}^{k-1}$$

$$\times \left\| \frac{e^{-\sqrt{\lambda}d_g(x,y)}}{d_g(x,y)} U_0(x,y) \mathbb{1}(d_g(x,y) < \varepsilon) \right\|_{L_x^{\infty} L_y^2} \frac{\mathcal{M}_{\varepsilon}}{\varepsilon^2} \left\| \frac{e^{-\frac{1}{2}\sqrt{\lambda}d_g(x,y)}}{d_g(x,y)} U_0(y,x)^2 \mathbb{1}(d_g(x,y) < \varepsilon) \right\|_{L_y^{\infty} L_x^2}$$

$$\times \left( \left\| \frac{\mathbb{1}(d_g(x,y) < \varepsilon)}{d_g(x,y)} \left( \frac{\mathcal{M}_{\varepsilon}}{\varepsilon^2} e^{-\frac{1}{2}\sqrt{\lambda}d_g(x,y)} U_0(y,x)^2 + |V(x)| U_0(x,y) e^{-\sqrt{\lambda}d_g(x,y)} \right) \right\|_{L_y^{\infty} L_x^1} \right)^{n-k}$$

with

$$\mathcal{F} = \begin{cases} \left\| \frac{e^{-\sqrt{\lambda}d_g(x,y)}}{d_g(x,y)} U_0(x,y) \mathbb{1}(\varepsilon/2 \leqslant d_g(x,y) < \varepsilon) V(y) \right\|_{L_x^\infty L_y^1} / \left\| \frac{e^{-\sqrt{\lambda}d_g(x,y)}}{d_g(x,y)} U_0(x,y) \mathbb{1}(d_g(x,y) < \varepsilon) V(y) \right\|_{L_x^\infty L_y^1} \\ \text{if } \ell \leqslant k-2 \,, \\ \left\| \frac{e^{-\sqrt{\lambda}d_g(x,y)}}{d_g(x,y)} U_0(x,y) \mathbb{1}(\varepsilon/2 \leqslant d_g(x,y) < \varepsilon) \right\|_{L_x^\infty L_y^2} / \left\| \frac{e^{-\sqrt{\lambda}d_g(x,y)}}{d_g(x,y)} U_0(x,y) \mathbb{1}(d_g(x,y) < \varepsilon) \right\|_{L_x^\infty L_y^2} \\ \text{if } \ell = k-1 \,, \\ \left\| \frac{e^{-\frac{1}{2}\sqrt{\lambda}d_g(x,y)}}{d_g(x,y)} U_0(y,x)^2 \mathbb{1}(\varepsilon/2 \leqslant d_g(x,y) < \varepsilon) \right\|_{L_y^\infty L_x^2} / \left\| \frac{e^{-\frac{1}{2}\sqrt{\lambda}d_g(x,y)}}{d_g(x,y)} U_0(y,x)^2 \mathbb{1}(d_g(x,y) < \varepsilon) \right\|_{L_y^\infty L_x^2} \\ \text{if } \ell = k \,, \\ \left\| \frac{\mathbb{1}(\varepsilon/2 \leqslant d_g(x,y) \leqslant \varepsilon)}{d_g(x,y)} \left( \frac{\mathcal{M}_x}{\varepsilon^2} e^{-\frac{1}{2}\sqrt{\lambda}d_g(x,y)} U_0(y,x)^2 + |V(x)| U_0(x,y) e^{-\sqrt{\lambda}d_g(x,y)} \right) \right\|_{L_y^\infty L_x^1} \\ / \left\| \frac{\mathbb{1}(d_g(x,y) \leqslant \varepsilon)}{d_g(x,y)} \left( \frac{\mathcal{M}_\varepsilon}{\varepsilon^2} e^{-\frac{1}{2}\sqrt{\lambda}d_g(x,y)} U_0(y,x)^2 + |V(x)| U_0(x,y) e^{-\sqrt{\lambda}d_g(x,y)} \right) \right\|_{L_y^\infty L_x^1} \\ \text{if } \ell \geqslant k+1 \,. \end{cases}$$

Using similar bounds as before we obtain the upper bound

$$C\frac{\varepsilon}{\sqrt{\lambda}} \mathcal{F}' \left( A \mathcal{M}'_{\varepsilon} \|V\|_{\mathcal{K}(\varepsilon)} \right)^{k-1} \frac{1}{\lambda^{1/4}} \frac{\mathcal{M}_{\varepsilon} \mathcal{M}'_{\varepsilon}}{\varepsilon^{2} \lambda^{1/4}} \left( A \left( \mathcal{M}'_{\varepsilon} \|V\|_{\mathcal{K}(\varepsilon)} + \frac{\mathcal{M}_{\varepsilon}}{\varepsilon^{2} \lambda} \right) \right)^{n-k}$$

with

ith 
$$\mathcal{F}' = \begin{cases} \left\| \frac{e^{-\sqrt{\lambda}d_g(x,y)}}{d_g(x,y)} U_0(x,y) \mathbb{1}(\varepsilon/2 \leqslant d_g(x,y) < \varepsilon) V(y) \right\|_{L_x^{\infty} L_y^1} / (\mathcal{M}'_{\varepsilon} ||V||_{\mathcal{K}(\varepsilon_{\lambda})}) \\ \text{if } \ell \leqslant k-2 \,, \\ \left\| \frac{e^{-\sqrt{\lambda}d_g(x,y)}}{d_g(x,y)} U_0(x,y) \mathbb{1}(\varepsilon/2 \leqslant d_g(x,y) < \varepsilon) \right\|_{L_x^{\infty} L_y^2} / \lambda^{-1/4} \\ \text{if } \ell = k-1 \,, \\ \left\| \frac{e^{-\frac{1}{2}\sqrt{\lambda}d_g(x,y)}}{d_g(x,y)} U_0(y,x)^2 \mathbb{1}(\varepsilon/2 \leqslant d_g(x,y) < \varepsilon) \right\|_{L_y^{\infty} L_x^2} / (\mathcal{M}'_{\varepsilon} \lambda^{-1/4}) \\ \text{if } \ell = k \,, \\ \left\| \frac{1(\varepsilon/2 \leqslant d_g(x,y) < \varepsilon)}{d_g(x,y)} \left( \frac{\mathcal{M}_{\varepsilon}}{\varepsilon^2} e^{-\frac{1}{2}\sqrt{\lambda}d_g(x,y)} U_0(y,x)^2 + |V(x)| U_0(x,y) e^{-\sqrt{\lambda}d_g(x,y)} \right) \right\|_{L_y^{\infty} L_x^1} / (\mathcal{M}'_{\varepsilon} ||V||_{\mathcal{K}(\varepsilon_{\lambda})} + \frac{\mathcal{M}_{\varepsilon}}{\varepsilon^2 \lambda}) \\ \text{if } \ell \geqslant k+1 \,. \end{cases}$$

Since  $\mathcal{F}' \leqslant Ce^{-\varepsilon\sqrt{\lambda}/4}$ , we obtain the claimed bound.

This concludes our discussion of the terms  $a_{\lambda,\varepsilon}^{(n,V)}(x,x)$  and  $b_{\lambda,\varepsilon}^{(n,V)}(x,x)$  in the decomposition (5.7). We now focus on the term  $\sigma_{\lambda,\varepsilon}^{(n,V)}(x,x)$ . Using the representation formula (3.4) with the function  $\kappa$  from (3.5) we can write

$$\partial_{\lambda} \sigma_{\lambda,\varepsilon}^{(n,V)}(x,x) = -\int_{0}^{\infty} \frac{t^{3/2} r_{0,\varepsilon}^{(n,V)}(t,x)}{(t+\lambda)^{3}} dt$$

with

$$r_{0,\varepsilon}^{(n,V)}(t,x) := \frac{1}{2(4\pi)^{n+1}} \int dv_g(z_n) \int dv_g(z_{n-1}) \cdots \int dv_g(z_1) \times \kappa(\sqrt{t}(d_g(x,z_1) + \cdots + d_g(z_n,x))) \ (d_g(x,z_1) + \cdots + d_g(z_n,x)) \times \frac{U_0(x,z_1)\chi(\varepsilon^{-1}d_g(x,z_1))}{d_g(x,z_1)} V(z_1) \frac{U_0(z_1,z_2)\chi(\varepsilon^{-1}d_g(z_1,z_2))}{d_g(z_1,z_2)} \cdots V(z_n) \frac{U_0(z_n,x)\chi(\varepsilon^{-1}d_g(z_n,x))}{d_g(z_n,x)} .$$
(5.8)

To bound  $r_{0,\varepsilon}^{(n,V)}$ , we use the Rodnianski–Schlag bound (A.1). A small variation of this argument, combined with the bound

$$|\kappa(r)| \leqslant C \quad \text{for all } r > 0,$$
 (5.9)

gives

$$|r_{0,\varepsilon}^{(n,V)}(t,x)| \leqslant C\mathcal{M}'_{\varepsilon} \left(A\mathcal{M}'_{\varepsilon} ||V||_{\mathcal{K}(\varepsilon)}\right)^{n}.$$
 (5.10)

Note that this shows that  $t^{3/2}r_{0,\varepsilon}^{(n,V)}$  is  $\mathcal{O}(t^{3/2})$ , while the other term  $r_{\varepsilon}^{(n,V)}$  was shown before to be  $\mathcal{O}(t)$ . What makes the term  $r_{0,\varepsilon}^{(n,V)}$  well-behaved is the following bound on its variation.

**Lemma 5.5.** There are constant C and A such that for all  $0 < t \le t'$ , all  $x \in M$ , all  $n \ge 1$ , and all  $\varepsilon \in (0, \rho]$ 

$$\left| r_{0,\varepsilon}^{(n,V)}(t,x) - r_{0,\varepsilon}^{(n,V)}(t',x) \right| \leqslant C \frac{\sqrt{t'} - \sqrt{t}}{\sqrt{t}} \mathcal{M}'_{\varepsilon} \left( A \mathcal{M}'_{\varepsilon} \|V\|_{\mathcal{K}(\varepsilon)} \right)^{n}$$

and

$$\begin{split} \left| t^{-3/2} r_{\varepsilon}^{(n,V)}(t,x) - (t')^{-3/2} r_{\varepsilon}^{(n,V)}(t',x) \right| \\ \leqslant C \frac{\sqrt{t'} - \sqrt{t}}{\sqrt{t}} \, \mathcal{M}_{\rho/2} \, \mathcal{M}_{\varepsilon}' \, \frac{\varepsilon^2}{\rho^2} \left( A \left( \mathcal{M}_{\varepsilon}' \|V\|_{\mathcal{K}(\varepsilon)} + \frac{\mathcal{M}_{\rho/2} \, \varepsilon^2}{\rho^2} \right) \right)^{n-1}. \end{split}$$

The proof of this lemma relies on the following property of the function  $\kappa$  from (3.5).

**Lemma 5.6.** There is a C > 0 such that for all  $v \ge u > 0$  and all  $r \ge 0$  we have

$$|\kappa(rv) - \kappa(ru)| \leqslant \frac{C(v-u)}{u}.$$

*Proof of Lemma 5.6.* By the mean value theorem, we have

$$\kappa(rv) - \kappa(ru) = \kappa'(\xi)r(v-u)$$

for some  $ru \leqslant \xi \leqslant rv$ . Using  $|\kappa'(t)| \leqslant C \min\{1, t^{-2}\}$  for all  $t \geqslant 0$ , we obtain

$$|\kappa'(\xi)| \leqslant C \min\{1, (ru)^{-2}\}.$$

Since  $\min\{1, (ru)^{-2}\} \leqslant (ru)^{-1}$ , we obtain the inequality in the lemma.

Proof of Lemma 5.5. We bound

$$\left| r_{0,\varepsilon}^{(n,V)}(t,x) - r_{0,\varepsilon}^{(n,V)}(t',x) \right| \leqslant \frac{1}{2(4\pi)^{n+1}} \int dv_g(z_n) \int dv_g(z_{n-1}) \cdots \int dv_g(z_1) \times \\
\times \left| \kappa(\sqrt{t}(d_g(x,z_1) + \cdots + d_g(z_n,x))) - \kappa(\sqrt{t'}(d_g(x,z_1) + \cdots + d_g(z_n,x))) \right| \times \\
\times \left( d_g(x,z_1) + \cdots + d_g(z_n,x) \right) \times \\
\times \frac{U_0(x,z_1)\chi(\varepsilon^{-1}d_g(x,z_1))}{d_g(x,z_1)} V(z_1) \frac{U_0(z_1,z_2)\chi(\varepsilon^{-1}d_g(z_1,z_2))}{d_g(z_1,z_2)} \cdots V(z_n) \frac{U_0(z_n,x)\chi(\varepsilon^{-1}d_g(z_n,x))}{d_g(z_n,x)} .$$

By Lemma 5.6 we have

$$\left|\kappa(\sqrt{t}(d_g(x,z_1)+\cdots+d_g(z_n,x)))-\kappa(\sqrt{t'}(d_g(x,z_1)+\cdots+d_g(z_n,x)))\right|\leqslant C\frac{\sqrt{t'}-\sqrt{t}}{\sqrt{t}}.$$

The first bound in the lemma now follows from the Rodnianski–Schlag bound (A.1). The second bound in the lemma is simpler to prove. We begin similarly as before, using Lemma 5.6. This leads to a similar expression as the one we estimated in the proof of Lemma 5.3, but with an additional factor  $t^{-1} \frac{\sqrt{t'} - \sqrt{t}}{\sqrt{t}} \left( d_g(x, z_1) + \cdots + d_g(z_n, x) \right)$  under the integral. Because of the restriction in the domain of integration, this factor is bounded by  $t^{-1} \frac{\sqrt{t'} - \sqrt{t}}{\sqrt{t}} (1 + n/2)\varepsilon$ , we arrive at the second bound in the lemma.

Finally, we are ready to give the

Proof of Proposition 5.1. We denote by A the maximum of the constants with this name appearing in Lemma 5.3, Lemma 5.4 and Lemma 5.5. By increasing A even further, we may assume that  $A \geqslant 1$ . We set  $\varepsilon_1 := \rho/\sqrt{4A\mathcal{M}_{\rho/2}}$  and let  $\varepsilon_2 > 0$  be maximal with the property that  $\|V\|_{\mathcal{K}(\varepsilon_2)} \leqslant 1/(4A\mathcal{M}'_{\rho/2})$ . In the following we assume that  $0 < \varepsilon \leqslant \varepsilon_0 := \min\{\varepsilon_1, \varepsilon_2\}$ . We note that  $\varepsilon \leqslant \varepsilon_1 \leqslant \rho/2$  and therefore  $\mathcal{M}'_{\varepsilon} \leqslant \mathcal{M}'_{\rho/2}$  and  $\mathcal{M}_{\varepsilon} \leqslant \mathcal{M}_{\rho/2}$ . Then, by Lemma 5.3, the series

$$R_\varepsilon^V(t,x) := \sum_{n\geqslant 1} (-1)^{n+1} r_\varepsilon^{(n,V)}(t,x)$$

converges and satisfies

$$|R_{\varepsilon}^{V}(t,x)| \leqslant C\mathcal{M}_{\rho/2}\mathcal{M}'_{\rho/2}\frac{\varepsilon t}{\rho^{2}}$$

Moreover, also the series

$$r_{0,\varepsilon}^{V}(t,x) := \sum_{n\geqslant 1} (-1)^{n+1} r_{0,\varepsilon}^{(n,V)}(t,x)$$
(5.11)

converges and satisfies, by (5.10),

$$\left|r_{0,\varepsilon}^{V}(t,x)\right| \leqslant C \left(\mathcal{M}_{\rho/2}'\right)^{2} \|V\|_{\mathcal{K}(\varepsilon)}$$

as well as, by Lemma 5.5, if  $0 < t \le t'$ ,

$$\left| r_{0,\varepsilon}^V(t,x) - r_{0,\varepsilon}^V(t',x) \right| \leqslant C \frac{\sqrt{t'} - \sqrt{t}}{\sqrt{t}} \left( \mathcal{M}'_{\rho/2} \right)^2 \|V\|_{\mathcal{K}(\varepsilon)}$$

and

$$\left| t^{-3/2} R_{\varepsilon}^{V}(t,x) - (t')^{-3/2} R_{\varepsilon}^{V}(t',x) \right| \leqslant C \frac{\sqrt{t'} - \sqrt{t}}{\sqrt{t}} \mathcal{M}_{\rho/2} \mathcal{M}_{\varepsilon}' \frac{\varepsilon^{2}}{\rho^{2}}.$$

Provided that  $\lambda \geqslant 4A\mathcal{M}_{\rho/2}/\varepsilon^2$ , we obtain from Lemma 5.4

$$\left| \partial_{\lambda} \gamma_{\lambda,\varepsilon}^{V}(t,x) + \int_{0}^{\infty} \frac{R_{\varepsilon}^{V}(t,x) + t^{3/2} r_{0,\varepsilon}^{V}(t,x)}{(t+\lambda)^{3}} dt \right|$$

$$= \left| \sum_{n\geqslant 1} (-1)^{n+1} \left( \partial_{\lambda} \gamma_{\lambda,\varepsilon}^{(n,V)}(x,x) - \partial_{\lambda} \sigma_{\lambda,\varepsilon}^{(n,V)}(x,x) - \partial_{\lambda} a_{\lambda,\varepsilon}^{(n,V)}(x,x) \right) \right|$$

$$= \left| \sum_{n\geqslant 1} (-1)^{n+1} \partial_{\lambda} b_{\lambda,\varepsilon}^{(n,V)}(x,x) \right|$$

$$\leqslant C \mathcal{M}_{\varepsilon}' \mathcal{M}_{\varepsilon} \frac{e^{-\varepsilon\sqrt{\lambda}/4}}{\varepsilon\lambda} .$$

Combining this bound with the identity

$$\partial_{\lambda} G_{\lambda}^{V}(x,x) = \partial_{\lambda} T_{\lambda,\varepsilon}^{V}(x,x) - \partial_{\lambda} \gamma_{\lambda,\varepsilon}^{V}(x,x) = -\frac{1}{8\pi\sqrt{\lambda}} - \partial_{\lambda} \gamma_{\lambda,\varepsilon}^{V}(x,x) ,$$

we obtain the bound in the proposition.

## 6. Consequences and discussion of Corollary 5.2

In this section we show that the main results stated in the introduction are consequences of Corollary 5.2.

6.1. **Pointwise Weyl laws.** We begin with the proof of the pointwise Weyl law for Kato class potentials.

*Proof of Theorem 1.1.* By Corollary 5.2 and Proposition 5.1 we have for all  $\varepsilon \in (0, \varepsilon_0)$ 

$$\limsup_{t \to \infty} t^{-3/2} \sup_{x \in M} \left| \mathbb{1}(-\Delta_g + V \leqslant t)(x, x) - \frac{t^{3/2}}{6\pi^2} \right| \leqslant \frac{1}{2} \limsup_{t \to \infty} \sup_{x \in M} \left| r_{0, \varepsilon}^V(t, x) \right| \leqslant C \|V\|_{\mathcal{K}(\varepsilon)},$$

with C independent of  $\varepsilon$ . By definition of the Kato class, this implies the theorem by letting  $\varepsilon \to 0$ .

We next state and prove a pointwise spectral cluster bound, valid for Kato class potentials. This bound appears in [8] under the additional assumption that  $V \in L^{3/2}(M)$ .

**Theorem 6.1** (Spectral cluster bounds). Let  $V: M \to \mathbb{R}$  be in the Kato class. Then uniformly in  $x \in M$  and t, T > 0,

$$\left| \mathbb{1} \left( t < -\Delta_g + V \leqslant (\sqrt{t} + \sqrt{T})^2 \right) (x, x) \right| \leqslant C \left( t T^{1/2} + t + T^{3/2} + 1 \right).$$

*Proof.* According to Corollary 5.2, we have, uniformly in  $x \in M$  and for a fixed  $\varepsilon \in (0, \varepsilon_0)$ 

$$\mathbb{1}\left(t < -\Delta_g + V \leqslant (\sqrt{t} + \sqrt{T})^2\right)(x, x) 
= \mathbb{1}\left(-\Delta_g + V \leqslant (\sqrt{t} + \sqrt{T})^2\right)(x, x) - \mathbb{1}\left(-\Delta_g + V \leqslant t\right)(x, x) 
= \frac{(\sqrt{t} + \sqrt{T})^3 - t^{3/2}}{6\pi^2} + \frac{1}{2}\left(r_{0,\varepsilon}^V((\sqrt{t} + \sqrt{T})^2, x)(\sqrt{t} + \sqrt{T})^3 - r_{0,\varepsilon}^V(t, x)t^{3/2}\right) + \mathcal{O}(t + T) 
= \frac{1}{2}r_{0,\varepsilon}^V((\sqrt{t} + \sqrt{T})^2, x)\left((\sqrt{t} + \sqrt{T})^3 - t^{3/2}\right) + \frac{1}{2}\left(r_{0,\varepsilon}^V((\sqrt{t} + \sqrt{T})^2, x) - r_{0,\varepsilon}^V(t, x)\right)t^{3/2} 
+ \mathcal{O}(t\sqrt{T} + T^{3/2} + t + T).$$

According to Proposition 5.1, we have, uniformly in  $x \in M$ ,

$$r_{0,\varepsilon}^{V}((\sqrt{t}+\sqrt{T})^{2},x)\left((\sqrt{t}+\sqrt{T})^{3}-t^{3/2}\right)=\mathcal{O}(t\sqrt{T}+T^{3/2})$$

and

$$r_{0,\varepsilon}^{V}((\sqrt{t}+\sqrt{T})^{2},x)-r_{0,\varepsilon}^{V}(t,x)=\mathcal{O}(t^{-1/2}T^{1/2})$$

Inserting this into the above bound, we obtain the assertion of the theorem.

Next, we prove the sharp pointwise version of Weyl's law under an additional regularity assumption.

Proof of Theorem 1.2. According to Corollary 5.2 it suffices to prove that

$$\sup_{x \in M} \left| r_{0,\varepsilon}^V(t,x) \right| \leqslant C t^{-1/2} \,.$$

We shall show that this holds for a certain choice of the parameter  $\varepsilon$  in the definition of  $r_{0,\varepsilon}^V$ .

Applying the bound (4.4) in the definition (5.8) we obtain

$$\left| r_{0,\varepsilon}^{(n,V)}(t,x) \right| \leqslant \frac{C t^{-1/2}}{2(4\pi)^{n+1}} \int dv_g(z_n) \cdots \int dv_g(z_1) \\
\times \frac{U_0(x,z_1)|\chi(\varepsilon^{-1}d_g(x,z_1))|}{d_g(x,z_1)} |V(z_1)| \frac{U_0(z_1,z_2)|\chi(\varepsilon^{-1}d_g(z_1,z_2))|}{d_g(z_1,z_2)} \cdots |V(z_n)| \frac{U_0(z_n,x)|\chi(\varepsilon^{-1}d_g(z_n,x))|}{d_g(z_n,x)} \\
\leqslant C t^{-1/2} A^n (\mathcal{M}'_{\varepsilon})^{n+1} \int dv_g(z_n) \int dv_g(z_{n-1}) \cdots \int dv_g(z_1) \\
\times \frac{\mathbb{1}(d_g(x,z_1) < \varepsilon)}{d_g(x,z_1)} |V(z_1)| \frac{\mathbb{1}(d_g(z_1,z_2) < \varepsilon)}{d_g(z_1,z_2)} \cdots |V(z_n)| \frac{\mathbb{1}(d_g(z_n,x) < \varepsilon)}{d_g(z_n,x)}.$$

We write  $z_0 = z_{n+1} = x$  and

$$\prod_{k=0}^{n} \frac{1}{d_g(z_k, z_{k+1})} = \prod_{\ell=0}^{n} \frac{d_g(z_\ell, z_{\ell+1})^{1/n}}{\prod_{k=0}^{n} d_g(z_k, z_{k+1})^{1/n}},$$

so by Hölder's inequality, the above n-fold integral is bounded by

$$\begin{split} & \prod_{\ell=0}^{n} \left( \int dv_{g}(z_{n}) \cdots \int dv_{g}(z_{1}) \frac{d_{g}(z_{\ell}, z_{\ell+1})^{(n+1)/n}}{\prod_{k=0}^{n} d_{g}(z_{k}, z_{k+1})^{(n+1)/n}} \prod_{m=0}^{n} \mathbb{1}(d_{g}(z_{m}, z_{m+1}) < \varepsilon) |V(z_{1})| \cdots |V(z_{n})| \right)^{1/(n+1)} \\ & \leqslant \left( \sup_{x \in M} \int dv_{g}(y) \frac{\mathbb{1}(d_{g}(x, y) < \varepsilon)}{d_{g}(x, y)^{(n+1)/n}} |V(y)| \right)^{n} \\ & \leqslant \left( \sup_{x \in M} \int dv_{g}(y) \frac{\mathbb{1}(d_{g}(x, y) < \varepsilon)}{d_{g}(x, y)^{2}} |V(y)| \right) \left( \sup_{x \in M} \int dv_{g}(y) \frac{\mathbb{1}(d_{g}(x, y) < \varepsilon)}{d_{g}(x, y)} |V(y)| \right)^{n-1}. \end{split}$$

Thus, we have shown that

$$\left| r_{0,\varepsilon}^{(n,V)}(t,x) \right| \leqslant C t^{-1/2} (\mathcal{M}_{\varepsilon}')^2 \left( \sup_{x \in M} \int dv_g(y) \frac{\mathbb{1}(d_g(x,y) < \varepsilon)}{d_g(x,y)^2} |V(y)| \right) \left( A \mathcal{M}_{\varepsilon}' \|V\|_{\mathcal{K}(\varepsilon)} \right)^{n-1}.$$

We now choose  $\varepsilon \leqslant \varepsilon'$  to have the supremum on the right finite and  $\varepsilon > 0$  so small that  $||V||_{\mathcal{K}(\varepsilon)} \leqslant 1/(2A\mathcal{M}'_{\varepsilon})$  to have exponential decay in n. By (5.11) we obtain, after summation over n,

$$\left| r_{0,\varepsilon}^{V}(t,x) \right| \leqslant C t^{-1/2} (\mathcal{M}_{\varepsilon}')^{2} \left( \sup_{x \in M} \int dv_{g}(y) \frac{\mathbb{1}(d_{g}(x,y) < \varepsilon)}{d_{g}(x,y)^{2}} |V(y)| \right).$$

This proves the theorem.

By Hölder's inequality the assumption in Theorem 1.2 is satisfied if  $V \in L^q$  for some q > 3. Thus, we obtain

Corollary 6.2. Assume that  $V \in L^q(M)$  for some q > 3. Then, uniformly in  $x \in M$ ,

$$\mathbb{1}(-\Delta_g + V \leqslant t)(x, x) = \frac{t^{3/2}}{6\pi^2} + \mathcal{O}(t)$$
.

6.2. Violation of the sharp pointwise Weyl law. While Corollary 6.2 shows that under the assumption  $V \in L^q$  for some q > 3 the pointwise Weyl law holds with the same remainder bound  $\mathcal{O}(t)$  as in the case without V, we now show that there are potentials  $V \in L^q(M)$  for some  $3/2 < q \le 3$  for which this remainder estimate is violated. These examples also show that for Kato class potentials the remainder  $o(t^{3/2})$  in Theorem 1.1 cannot be improved to  $\mathcal{O}(t^{3/2-\varepsilon})$  for some  $\varepsilon > 0$  valid for all Kato class V.

Note that if  $V \in L^q$  for some q > 3/2, then V is in the Kato class, so Corollary 5.2 is applicable. Therefore our strategy to proving a violation of the  $\mathcal{O}(t)$  bound will be to show that the bound  $r_{0,\varepsilon}^V(t,x) = \mathcal{O}(t^{-1/2})$  can be violated. Recall that  $r_{0,\varepsilon}^V$  is defined in (5.11) as an infinite sum of terms  $r_{0,\varepsilon}^{(n,V)}$ . As a first step towards showing the violation of an  $\mathcal{O}(t^{-1/2})$  bound on  $r_{0,\varepsilon}^V$ , we show that such a violation can come from at most a finite number of terms  $r_{0,\varepsilon}^{(n,V)}$ .

**Lemma 6.3.** Let  $V \in L^q(M)$  for some q > 3/2 and set

$$N_q := 1 + \left\lfloor \frac{q}{2q - 3} \right\rfloor .$$

Then, uniformly in  $x \in M$ ,

$$\sum_{n\geqslant N_a} |r_0^{(n,V)}(t,x)| = \mathcal{O}(t^{-1/2}),$$

provided  $\varepsilon$  in the definition of  $r_{0,\varepsilon}^{(n,V)}$  is chosen small enough.

As a consequence, there is only a finite number of additional terms in the corrected Weyl law (5.3). This number of terms grows as the singularity of the potential gets worse. Note that if q > 3, then  $N_q = 1$  and the statement of the lemma is a consequence of the proof of Theorem 1.2.

*Proof.* During the proof of Theorem 1.2 we have shown the bound

$$\left| r_{0,\varepsilon}^{(n,V)}(t,x) \right| \leqslant C t^{-1/2} \mathcal{M}_{\varepsilon}' \left( A \mathcal{M}_{\varepsilon}' \sup_{x \in M} \int dv_g(y) \frac{\mathbb{1}(d_g(x,y) < \varepsilon)}{d_g(x,y)^{(n+1)/n}} |V(y)| \right)^n.$$

We bound

$$\int dv_g(y) \frac{\mathbb{1}(d_g(x,y) < \varepsilon)}{d_g(x,y)^{(n+1)/n}} |V(y)| \leq ||V||_{L^q} \left( \int dv_g(y) \frac{\mathbb{1}(d_g(x,y) < \varepsilon)}{d_g(x,y)^{(n+1)q'/n}} \right)^{1/q'}.$$

Under the assumption (n+1)q'/n < 3, which is the same as  $n \ge N_q$ , we have

$$\int dv_g(y) \frac{\mathbb{1}(d_g(x,y) < \varepsilon)}{d_g(x,y)^{(n+1)q'/n}} \leq \sup_{\substack{d_g(y,x) < \varepsilon \\ \\ \leq C' \varepsilon^{3-(n+1)q'/n}}} U_0(x,y)^{-2} \int_{|z| < \varepsilon} \frac{dz}{|z|^{(n+1)q'/n}} \sup_{\substack{d_g(y,x) < \varepsilon \\ \\ d_g(y,x) < \varepsilon}} U_0(x,y)^{-2}.$$

Note that the constant C' here can be chosen independently of n. Thus,

$$\left| r_{0,\varepsilon}^{(n,V)}(t,x) \right| \leqslant C t^{-1/2} \varepsilon^{-1} \mathcal{M}_{\varepsilon}' \left( A \mathcal{M}_{\varepsilon}' \|V\|_{L^{q}} (C')^{1/q'} \varepsilon^{3/q'-1} \sup_{d_{g}(y,x) < \varepsilon} U_{0}(x,y)^{-2/q'} \right)^{n}.$$

Note that 3/q' - 1 = 2 - 3/q > 0. Therefore, by choosing  $\varepsilon$  small enough, the bound is exponentially decaying in n and we obtain the assertion in the lemma.

The following example shows that the additional terms  $r_{0,\varepsilon}^{(n,V)}$  can cause a violation of the sharp pointwise Weyl law.

**Proposition 6.4.** Let  $\eta \in (0,1)$ ,  $x_0 \in M$ ,  $\gamma \in \mathbb{R}$  and let  $V: M \to \mathbb{R}$  defined by

$$V(x) = \gamma \frac{\chi(d_g(x, x_0))}{d_g(x, x_0)^{2-\eta}}, \quad x \in M.$$

Then

$$\mathbb{1}(-\Delta_g + V \leqslant t)(x,x) = \frac{t^{3/2}}{6\pi^2} - \frac{1}{2} \sum_{1 \leqslant n < 1/\eta} (-1)^{n+1} r_{0,\varepsilon}^{(n,V)}(t,x) t^{3/2} + \mathcal{O}(t).$$

Moreover, for all  $n < 2/\eta$  we have

$$\sup_{x \in M} |r_{0,\varepsilon}^{(n,V)}(t,x)| \leqslant Ct^{-n\eta/2}. \tag{6.1}$$

and, in geodesic normal coordinates around  $x_0$ , for all  $y \in T_{x_0}M \sim \mathbb{R}^3$ ,

$$\lim_{t \to +\infty} t^{n\eta/2} r_{0,\varepsilon}^{(n,V)}(t,y/\sqrt{t}) = \frac{\gamma^n}{2(4\pi)^{n+1}} \int_{\mathbb{R}^3} dz_1 \cdots \int_{\mathbb{R}^3} dz_n \times \frac{\kappa(|z_1| + |z_1 - z_2| + \cdots + |z_{n-1} - z_n| + |z_n|) (|z_1| + |z_1 - z_2| + \cdots + |z_{n-1} - z_n| + |z_n|)}{|z_1||z_1 - z_2| \cdots |z_{n-1} - z_n||z_n||z_1 + y|^{2-\eta} \cdots |z_n + y|^{2-\eta}},$$
(6.2)

where the right side is finite.

After the proof of this proposition we will discuss the singular term for n=1 in more detail and show, in particular, that this term is not identically zero for  $\gamma \neq 0$ . This shows that the sharp pointwise Weyl law can be violated for a Kato class potential. We also note that as the potential becomes more singular (that is, as  $\eta$  decreases to zero), the number of 'additional' terms becomes arbitrarily large.

*Proof.* Since the potential V belongs to  $L^q$  for all  $q < 3/(2 - \eta)$ , we may apply Lemma 6.3 and obtain

$$\sup_{x \in M} \sum_{n \geqslant 1 + |1/\eta|} |r_0^{(n,V)}(t,x)| = \mathcal{O}(t^{-1/2}).$$

Here we used the fact that  $\frac{q}{2q-3} \searrow \frac{1}{\eta}$  as  $q \nearrow 3/(2-\eta)$ , and therefore  $N_q = 1 + \left\lfloor \frac{q}{2q-3} \right\rfloor = 1 + \lfloor 1/\eta \rfloor$  for all q below but sufficiently close to  $3/(2-\eta)$ .

This, together with Corollary 5.2 implies that

$$\mathbb{1}(-\Delta_g + V \leqslant t)(x,x) = \frac{t^{3/2}}{6\pi^2} - \frac{1}{2} \sum_{1 \leqslant n \leqslant \lfloor 1/n \rfloor} (-1)^{n+1} r_{0,\varepsilon}^{(n,V)}(t,x) t^{3/2} + \mathcal{O}(t).$$

This proves the first claim in the proposition, except in the case where  $1/\eta$  is an integer and  $n = 1/\eta$ . The fact that this term can also be included in the remainder term follows from (6.1) (noting that  $1/\eta < 2/\eta$ ), which will be proved below.

From now on, we concentrate on proving (6.1) and (6.2). It is easy to see that the limit is independent of  $\varepsilon$ . Therefore we may assume in the following that  $\varepsilon$  is so small (depending on n) and t is so large (depending on y) that all points in the support of the integrand of the n-fold integral defining  $r_{0,\varepsilon}^{(n,V)}$  are contained in  $B_{\rho/2}(x_0)$ . Consequently we can compute the integral defining  $r_0^{(n,V)}(t,y)$  in geodesic normal coordinates around  $x_0$ . We identify  $x_0$  with the origin,  $T_{x_0}M$  with  $\mathbb{R}^3$  and  $B_{\rho/2}(x_0)$  with  $\mathcal{U}$ . We recall that in these coordinates,  $dv_g(z) = \theta(z,0) dz$ . We will also identify  $\exp_{x_0}(y)$  with y and y with the Euclidean metric to lighten the notation. In particular, we have  $d_g(x_0,y) = |y|$  for all y. We also use the following representation of the Riemannian distance:

$$d_g(z, z') = \sqrt{\langle z - z', G(z, z')(z - z') \rangle} =: |z - z'|_{z, z'}, \quad z, z' \in \mathcal{U},$$

for some smooth family (G(z,z')) of positive definite symmetric matrices with

$$G(z, z) = (D_z \exp_{x_0})^T g(\exp_{x_0}(z)) D_z \exp_{x_0},$$

see for instance [18, Eq. (4.3)]. In particular, the map  $u \mapsto |u|_{z,z'}$  is homogeneous of degree 1. Changing variables  $z_j = (y + \zeta_j)/\sqrt{t}$  in the integral defining  $r^{(n,V)}(t,y)$ , we find (with the convention  $\zeta_0 = 0 = \zeta_{n+1}$ )

$$\begin{split} r_{0,\varepsilon}^{(n,V)}(t,y) &= \frac{t^{-n\eta/2} \, \gamma^n}{2(4\pi)^{n+1}} \int_{\mathbb{R}^3} d\zeta_n \int_{\mathbb{R}^3} d\zeta_{n-1} \cdots \int_{\mathbb{R}^3} d\zeta_1 \\ &\left( \prod_{m=1}^n \frac{\theta(0,(y+\zeta_m)/\sqrt{t}) \, \chi(|\zeta_{m-1}-\zeta_m|/(\varepsilon\sqrt{t}))}{\theta((y+\zeta_{m-1})/\sqrt{t},(y+\zeta_m)/\sqrt{t})^{1/2}} \right) \frac{\chi(|\zeta_n|_{(y+\zeta_n)/\sqrt{t},y/\sqrt{t}}/(\varepsilon\sqrt{t}))}{\theta((y+\zeta_n)/\sqrt{t},y/\sqrt{t})^{1/2}} \\ &\times \kappa \left( \sum_{m=0}^n |\zeta_m-\zeta_{m+1}|_{(y+z_m)/\sqrt{t},(y+\zeta_{m+1})/\sqrt{t}} \right) \frac{\sum_{m=0}^n |\zeta_m-\zeta_{m+1}|_{(y+z_m)/\sqrt{t},(y+\zeta_{m+1})/\sqrt{t}}}{\prod_{m=0}^n |\zeta_m-\zeta_{m+1}|_{(y+z_m)/\sqrt{t},(y+\zeta_{m+1})/\sqrt{t}}} \\ &\times \prod_{m=1}^n \frac{\chi(|y+\zeta_m|/\sqrt{t})}{|y+\zeta_m|^{2-\eta}} \, . \end{split}$$

We now let  $t \to \infty$  and use dominated convergence. Note that the integrand in the above formula tends pointwise to the integrand in (6.2) since for all  $y, \zeta, \zeta', \zeta'' \in \mathbb{R}^3$ ,

$$|\zeta|_{(y+\zeta')/\sqrt{t},(y+\zeta'')/\sqrt{t}} \to |\zeta|_{0,0} = |\zeta|$$
 as  $t \to +\infty$ ,

as well as  $\theta(0,0) = 1 = \chi(0)$ .

To get the integrable majorant, one can use

$$(1/C)|z-z'| \leqslant d_g(z,z') \leqslant C|z-z'|$$
 for all  $z,z' \in \mathcal{U}$ ,

as well as  $1/C \le \theta \le C$  and  $\chi \le C$ . The required bound then follows immediately from the bound  $|\kappa(r)| \le C(1+|r|)^{-2}$  and the fact that, if  $n < 2/\eta$ ,

$$\int_{\mathbb{R}^3} \cdots \int_{\mathbb{R}^3} \frac{\sum_{m=0}^n |z_m - z_{m+1}| dz_1 \cdots dz_n}{(1 + \sum_{m=0}^n |z_m - z_{m+1}|)^2 \left( \prod_{m=0}^n |z_m - z_{m+1}| \right) |z_1 + y|^{2-\eta} \cdots |z_n + y|^{2-\eta}} < +\infty,$$

where again  $z_0 = 0 = z_{n+1}$ . This fact may be proved inductively, using the estimates

$$\begin{split} \sup_{z''} \int_{\mathbb{R}^3} \frac{dz}{(1+a+|z|+|z-z'|)^2|z||z+z''|^{2-\eta}} &\leqslant \frac{C}{(1+a+|z'|)^{2-\eta}} \,, \\ \sup_{z''} \int_{\mathbb{R}^3} \frac{dz}{(1+a+|z|+|z-z'|)^\beta |z-z'||z+z''|^{2-\eta}} &\leqslant \frac{C}{(1+a+|z'|)^{\beta-\eta}} \,, \end{split}$$

which hold for all  $z' \in \mathbb{R}^3$ , all a > 0, and all  $\beta > \eta$  for some C > 0 independent of z', a. The claimed dependence of the right sides on a follows simply by scaling.

This completes the proof of (6.2). The same argument yields the uniform bound (6.1).  $\square$ 

In the remainder of this subsection, we discuss the limiting profile that arises in the case n = 1 in (6.2). We set

$$\Xi_{\eta}(y) := \frac{1}{16 \,\pi^2} \int_{\mathbb{R}^3} dz \, \frac{\kappa(2|z|)}{|z| \, |z+y|^{2-\eta}} \,,$$

so that (6.2) reads  $\lim_{t\to+\infty} t^{\eta/2} r_{0,\varepsilon}^{(1,V)}(t,y/\sqrt{t}) = \gamma \Xi_{\eta}(y)$  and, by Corollary 5.2,

$$\mathbb{1}(-\Delta_g + V \leqslant t)(y/\sqrt{t}, y/\sqrt{t}) = \frac{t^{3/2}}{6\pi^2} - \frac{\gamma t^{(3-\eta)/2}}{2} \Xi_{\eta}(y) + o(t^{(3-\eta)/2}).$$

Note that due to the oscillation of the function  $\kappa$  it is not a priori clear whether  $\Xi_{\eta}$  is not identically zero. However, physical intuition suggests that places where the potential V assumes large positive values repel 'particles' and conversely, places where V is large negative attract 'particles'. Thus, we expect the particle density at the singularity,  $\mathbb{1}(-\Delta_g + V \leq t)(x_0, x_0)$ , to be decreasing in  $\gamma$  and therefore  $\Xi_{\eta}(0) > 0$ . The following lemma confirms this and also provides the asymptotic behavior of  $\Xi_{\eta}$  at infinity.

**Lemma 6.5.** Let  $\eta \in (0,1)$ . Then

$$\Xi_{\eta}(0) = \frac{1}{\pi^2} \frac{2^{1-\eta}}{3-\eta} \frac{\cos(\pi\eta/2)}{1-\eta} \Gamma(\eta) > 0$$

and

$$\lim_{|y|\to\infty}|y|^{2-\eta}\,\Xi_\eta(y)=\frac{1}{2\pi^2}\,.$$

Numerical integration indicates that  $\Xi_{\eta}$  is positive and decreasing, but not convex.

*Proof.* We have

$$16\pi^{2}\Xi_{\eta}(0) = 2^{-\eta}4\pi \int_{0}^{\infty} dr \frac{\kappa(r)}{r^{1-\eta}} = 2^{5-\eta} \int_{0}^{\infty} dr \frac{\sin r - r \cos r}{r^{4-\eta}} = \frac{2^{5-\eta}}{3-\eta} \int_{0}^{\infty} dr \frac{\sin r}{r^{2-\eta}}.$$

Here, when integrating by parts we used the fact that  $\lim_{r\to 0} r^{-3+\eta} (\sin r - r \cos r) = 0$ . We cannot integrate by parts again if we want to have the integral absolutely convergent at infinity. We write

$$\int_0^\infty dr \, \frac{\sin r}{r^{2-\eta}} = \operatorname{Im} \, \int_0^\infty dr \, \frac{e^{ir} - 1}{r^{2-\eta}} \, .$$

The term 1 was subtracted to make the integral converge at the origin. Thus, we are integrating the analytic function  $(e^{iz}-1)/e^{(2-\eta)\log z}$  along the positive real axis. Since the

integrand is  $O(|z|^{-2+\eta})$  as  $|z| \to \infty$  on the upper half plane and  $O(|z|^{-1+\eta})$  as  $|z| \to 0$ , we can move the contour to the positive imaginary axis and obtain

$$\int_0^\infty dr \, \frac{e^{ir} - 1}{r^{2-\eta}} = i \int_0^\infty dt \, \frac{e^{-t} - 1}{t^{2-\eta} e^{(1-\eta/2)i\pi}} = i e^{-(1-\eta/2)i\pi} \Gamma(-1+\eta) \,.$$

In the last formula we have used the standard method of analytic continuation of the gamma function. Using

$$ie^{-(1-\eta/2)i\pi}\Gamma(-1+\eta) = \frac{ie^{i\pi\eta/2}}{1-\eta}\Gamma(\eta)$$
,

we obtain the claimed formula for  $\Xi_{\eta}(0)$ .

In order to compute the large y asymptotics we set s := |y| and compute

$$16 \pi^{2} \Xi_{\eta}(y) = 2^{-\eta} 2\pi \int_{0}^{\infty} dr \, r \, \kappa(r) \int_{-1}^{1} \frac{dt}{(r^{2} - 4rst + 4s^{2})^{1-\eta/2}}$$

$$= \frac{2^{-\eta} \pi}{\eta} \int_{0}^{\infty} dr \, \frac{\kappa(r)}{s} \left( (r + 2s)^{\eta} - |r - 2s|^{\eta} \right)$$

$$= \frac{2^{3-\eta}}{\eta} s^{-1+\eta} \int_{0}^{\infty} d\rho \, \sin(s\rho) \, F(\rho) \,,$$

where

$$F(\rho) := \rho \int_{\rho}^{\infty} \frac{d\rho'}{(\rho')^3} \left( (\rho' + 2)^{\eta} - |\rho' - 2|^{\eta} \right).$$

Here we integrated by parts once. We shall show that

$$\int_0^\infty d\rho \, \sin(s\rho) \, F(\rho) = \eta \, 2^{\eta} \, s^{-1} + o(s^{-1}) \,,$$

which implies the asymptotics in the lemma. Integrating by parts once and using the integrability of F' on  $[1, +\infty)$  and the Riemann–Lebesgue lemma, we obtain

$$\int_{1}^{\infty} d\rho \sin(s\rho) F(\rho) = \frac{\cos s}{s} F(1) + o(s^{-1}).$$

This term is cancelled to leading order by

$$\int_0^1 d\rho \, \sin(s\rho) \, \rho F(1) = -\frac{\cos s}{s} \, F(1) + \frac{\sin s}{s^2} \, F(1) \, .$$

This leaves us with

$$\int_0^1 d\rho \sin(s\rho) \left( F(\rho) - \rho F(1) \right) = \int_0^1 d\rho \sin(s\rho) \rho \int_\rho^1 \frac{d\rho'}{(\rho')^3} \left( (\rho' + 2)^\eta - |\rho' - 2|^\eta \right)$$
$$= \eta 2^\eta \int_0^1 d\rho \sin(s\rho) \rho \int_\rho^1 \frac{d\rho'}{(\rho')^2} \left( 1 + (\rho')^2 g(\rho') \right),$$

where

$$g(\rho') := \eta^{-1} \, 2^{-\eta} (\rho')^{-3} \left( (\rho' + 2)^{\eta} - |\rho' - 2|^{\eta} - \eta \, 2^{\eta} \, \rho' \right).$$

Using the fact that  $g \in C^1[0,1]$  one finds after two integrations by part that

$$\int_0^1 d\rho \sin(s\rho) \, \rho \int_\rho^1 d\rho' \, g(\rho') = \mathcal{O}(s^{-2}) \,.$$

Finally, by an explicit computation,

$$\int_0^1 d\rho \sin(s\rho) \, \rho \int_\rho^1 \frac{d\rho'}{(\rho')^2} = \frac{1}{s} + \mathcal{O}(s^{-2}) \, .$$

Collecting all these bounds, we obtain the claimed asymptotics of  $\int d\rho \sin(s\rho) F(\rho)$ .

6.3. Integrated Weyl laws. Next, we show that, while the term  $r_{0,\varepsilon}^V$  can lead to a violation of the pointwise sharp Weyl law, the integrated version of the sharp Weyl law remains valid for Kato class potentials. In fact, it even holds for potentials which are sums of Kato class and  $L^{3/2}(M)$  functions.

Proof of Theorem 1.3. We begin with the case where V belongs to the Kato class (which was treated by a different method in [30]). Clearly, we have

$$N(t, -\Delta_g + V) = \int_M \mathbb{1}(-\Delta_g + V \leqslant t)(x, x) dx.$$

In view of the uniform pointwise asymptotics in Corollary 5.2, it suffices to show that

$$\int_{M} r_{0,\varepsilon}^{V}(t,x) dx = \mathcal{O}(t^{-1/2}) \quad \text{as } t \to +\infty.$$

The parameter  $\varepsilon$  which enters the definition of  $r_{0,\varepsilon}^V$  is fixed throughout this proof. From (4.4) we deduce that for all  $a, b, x \in M$  and all  $\alpha \ge 0$ ,

$$\int_{\substack{d_g(x,a)<\varepsilon\\d_g(x,b)<\varepsilon}} \kappa(\sqrt{t}(d_g(x,a)+\alpha+d_g(b,x))) \frac{d_g(x,a)+\alpha+d_g(b,x)}{d_g(a,x)d_g(x,b)} dv_g(x)$$

$$\leqslant \frac{C}{\sqrt{t}} \int_{\substack{d_g(x,a)<\varepsilon\\d_g(x,b)<\varepsilon}} \frac{dx}{d_g(a,x)d_g(x,b)} \leqslant \frac{C'\varepsilon}{\sqrt{t}}.$$

We do not track the dependence of the constants in this proof on the geometry of M. Inserting this bound into the definition of (5.11) we easily obtain that for all  $n \ge 1$ 

$$\int_{M} \left| r_0^{(n,V)}(t,x) \right| dx \leqslant C'' \varepsilon (A \|V\|_{\mathcal{K}(\varepsilon)})^{n-1} \|V\|_{L^1} t^{-1/2}.$$

Decreasing  $\varepsilon$  if necessary we can assume that  $A||V||_{\mathcal{K}(\varepsilon)} \leq 1/2$  and then we obtain the claimed bound by summing over n.

Let us now consider the general case where  $V = V_1 + V_2$  with  $V_1$  in the Kato class and  $V_2 \in L^{3/2}(M)$ . We make use of the fact that for any self-adjoint lower A and B one has  $N(0, A + B) \leq N(0, A) + N(0, B)$ . This implies that for any  $\delta \in (0, 1]$ ,

$$N(0, -(1+\delta)\Delta_g + V_1 - t) - N(0, -\delta\Delta_g - V_2) \leqslant N(0, -\Delta_g + V - t)$$
  
 
$$\leqslant N(0, -(1-\delta)\Delta_g + V_1 - t) + N(0, -\delta\Delta_g + V_2).$$

By the first part of the proof, we have for any  $\delta \in (0, 1/2]$ ,

$$N(0, -(1 \pm \delta)\Delta_g + V_1 - t) = N((1 \pm \delta)^{-1}t, -\Delta_g + (1 \pm \delta)^{-1}V_1) = \frac{t^{3/2}}{6\pi^2} + \mathcal{O}(\delta t^{3/2} + t).$$

(Here we use the fact that the bounds in Corollary 5.2 and in the first part of the proof hold for  $(1 \pm \delta)^{-1}V$  instead of V with constants independent of  $\delta \in (0, 1/2]$ .)

In order to deal with the contribution from  $V_2$ , we shall use the CLR inequality

$$N(0, -\Delta_g + W) \leqslant C \int_M (W(x) - E_*)_-^{3/2} dx$$
(6.3)

with constants C and  $E_*$  depending only on M. This bound is essentially known and we give some references after the proof. In our case it implies that

$$N(0, -\delta \Delta_g \pm V_2) = N(0, -\Delta_g \pm \delta^{-1}V_2) \leqslant C \int_M (\delta^{-1}V_2(x) - E_*)_-^{3/2} dx = \mathcal{O}(\delta^{-3/2}).$$

Choosing  $\delta = t^{-3/5}$  we get  $\delta t^{3/2} = \delta^{-3/2} = t^{9/10} = \mathcal{O}(t)$ . Combining the previous bounds we obtain the assertion of the theorem.

Since we have not been able to find the CLR inequality (6.3) stated explicitly in the literature, let us briefly comment on its proof. Inequalities of this type go back to Cwikel [16], Lieb [42, 43] and Rozenblum [47, 48] in the Euclidean case. Further proofs in that case were given in [22, 41, 14, 23]. The proofs in [47, 41] extend to the case of a manifold. For instance, [24], which is based on [41] (see also [39, 9]), implies that the Sobolev inequality

$$\int_{M} (|\nabla_{g} u|^{2} + E_{*}|u|^{2}) dv_{g}(x) \geqslant S \left( \int_{M} |u|^{6} dv_{g}(x) \right)^{1/3},$$

which is valid with some positive constants S and  $E_*$  depending on M, implies (6.3) with  $C \leq \sqrt{e}S^{-3/2}$ . (Conversely, it is elementary to see that (6.3) implies the Sobolev inequality with constant S satisfying  $C \geq S^{-3/2}$ .) Alternatively, one can deduce (6.3) from [23, Thm. 3.2 and Lem. 3.4]. This needs as an ingredient the bound  $\mathbb{1}(-\Delta_g \leq E)(x,x) \leq C'(E+E_*)^{3/2}$  for all  $E \geq 0$  and  $x \in M$ , which itself follows from a rough version of the Weyl law for  $-\Delta_g$ .

Remark 6.6. The argument in the previous proof based on (6.3) shows that if  $V \in L^{3/2}(M)$  and  $N(t, -\Delta_g) = t^{3/2}/(6\pi^2) + o(t)$ , then also  $N(t, -\Delta_g + V) = t^{3/2}/(6\pi^2) + o(t)$ . More precisely, if  $|N(t, -\Delta_g) - t^{3/2}/(6\pi^2)| \le r(t)$  for a nondecreasing function r, then the previous proof shows that, if  $V \in L^{3/2}(M)$ , then

$$|N(t, -\Delta_q + V) - t^{3/2}/(6\pi^2)| \le r(2t) + Ct^{9/10}$$
.

Since the CLR inequality is also valid in any dimension  $N \ge 3$ , the previous argument extends to this case with the remainder  $\mathcal{O}(t^{9/10})$  replaced by  $\mathcal{O}(t^{N^2/(2(N+2))})$ . For instance, when M has nonpositive sectional curvatures, then this bound and Bérard's theorem yield

$$N(t, -\Delta_g + V) = \frac{|\mathbb{S}^{N-1}|}{N(2\pi)^N} t^{N/2} + \mathcal{O}(t^{(N-1)/2}/\ln t)$$

for  $V \in L^{N/2}(M)$ . The same result, but for Kato class V, was shown in [30, Thm. 1.3]. On the other hand, our argument is not strong enough to reproduce the  $L^{3/2}$  analogue of the  $\mathcal{O}(t^{3/4})$  bound for  $M = (\mathbb{R}/\mathbb{Z})^3$  and Kato class V, which is shown in [30, Thm. 1.4].

We expect that  $N(t, -\Delta_g) = t^{3/2}/(6\pi^2) + o(t)$  implies  $N(t, -\Delta_g + V) = t^{3/2}/(6\pi^2) + o(t)$  for all Kato class V, but we do not know how to prove it.

#### 7. Euclidean domains

In this section we will show that the previous construction extends easily to domains in  $\mathbb{R}^3$ . Let  $\Omega \subsetneq \mathbb{R}^3$  be an open set. We emphasize that no regularity assumption on  $\partial\Omega$  and no finite measure assumption are imposed, except in Corollary 7.7. Let us denote by  $-\Delta_{\Omega}^D$  the Dirichlet Laplacian on  $\Omega$ . Let V a Kato class potential on  $\Omega$ . The starting point is a parametrix for the Green's function

$$G^V_{\lambda}(x,y) := (-\Delta^D_{\Omega} + V + \lambda)^{-1}(x,y), \ x \neq y \in \Omega.$$

which will be the Green's function for same operator on  $\mathbb{R}^N$ , thus getting rid of the boundary:

$$T_{\lambda}^{V}(x,y) := (-\Delta_{\mathbb{R}^3} + V \mathbb{1}_{\Omega} + \lambda)^{-1}(x,y), \ x \neq y \in \mathbb{R}^N.$$

We begin with an estimate on this Green's function.

**Lemma 7.1.** There exist C > 0 and  $\Lambda > 0$  such that for all  $x \neq y \in \mathbb{R}^N$  and for all  $\lambda \geqslant \Lambda$ , we have

$$|T_{\lambda}^{V}(x,y)| \leqslant C \frac{e^{-\frac{\sqrt{\lambda}}{2}|x-y|}}{|x-y|}.$$

*Proof.* Using the resolvent formula, we expand  $T_{\lambda}^{V} = T_{\lambda}^{0} + \sum_{n \geq 1} T_{\lambda}^{(n,V)}$ , where

$$T_{\lambda}^{(n,V)}(x,y) = \frac{1}{(4\pi)^{n+1}} \int_{\Omega} dz_1 \cdots \int_{\Omega} dz_n \frac{e^{-\sqrt{\lambda}(|x-z_1|+|z_1-z_2|+\cdots+|z_n-y|)}}{|x-z_1||z_1-z_2|\cdots|z_n-y|} V(z_1) \cdots V(z_n).$$

By the triangle inequality, we have for all  $z_1, \ldots, z_n$ ,

$$e^{-\frac{\sqrt{\lambda}}{2}(|x-z_1|+|z_1-z_2|+\cdots+|z_n-y|)} \le e^{-\frac{\sqrt{\lambda}}{2}|x-y|}$$

and

$$\frac{|x-y|}{|x-z_1||z_1-z_2|\cdots|z_n-y|} \leqslant \frac{|x-z_1|+|z_1-z_2|+\cdots+|z_n-y|}{|x-z_1||z_1-z_2|\cdots|z_n-y|} 
= \frac{1}{|z_1-z_2|\cdots|z_n-y|} + \frac{1}{|x-z_1||z_2-z_3|+\cdots+|z_n-y|} 
+ \cdots + \frac{1}{|x-z_1||z_1-z_2|\cdots|z_{n-1}-z_n|}.$$

As a consequence, we deduce that

$$\sup_{x \neq y} |x - y| e^{\frac{\sqrt{\lambda}}{2}|x - y|} |T_{\lambda}^{(n,V)}(x,y)| \leq \frac{n+1}{(4\pi)^{n+1}} \left( \sup_{x \in \mathbb{R}^N} \int_{\Omega} \frac{e^{-\frac{\sqrt{\lambda}}{2}|x - z|}}{|x - z|} |V(z)| \, dz \right)^n$$

$$\leq C \left( (n+1) (A||V||_{\mathcal{K}(8/\sqrt{\lambda})})^n,$$

where in the last line we used Lemma A.2.

Remark 7.2. By the maximum principle, we know that  $0 \le G_{\lambda}^0 \le T_{\lambda}^0$  for all  $\lambda \ge 0$  (pointwise). Hence, the same proof using the resolvent identity for  $G_{\lambda}^V$  shows that the same bound

$$|G_{\lambda}^{V}(x,y)| \leqslant C \frac{e^{-\frac{\sqrt{\lambda}}{2}|x-y|}}{|x-y|} \tag{7.1}$$

holds for  $\lambda$  large enough.

We can now estimate the error between  $G_{\lambda}^{V}$  and  $T_{\lambda}^{V}$ .

**Lemma 7.3.** There are C>0 and  $\Lambda>0$  such that for all  $x\neq y\in\Omega$  and  $\lambda\geqslant\Lambda$  we have

$$\left|G_{\lambda}^{V}(x,y)-T_{\lambda}^{V}(x,y)\right|\leqslant C\frac{e^{-\frac{\sqrt{\lambda}}{2}d(x,\partial\Omega)}}{d(x,\partial\Omega)}.$$

*Proof.* For all  $x \in \Omega$ , the function  $\gamma_{\lambda}^{V} := G_{\lambda}^{V} - T_{\lambda}^{V}$  satisfies the equation

$$\begin{cases} (-\Delta + V + \lambda)_y \gamma_\lambda^V(x, \cdot) = 0 & \text{in } \Omega, \\ \gamma_\lambda^V(x, \cdot) = -T_\lambda^V(x, \cdot) & \text{on } \partial \Omega. \end{cases}$$

Let us show that there are C>0 and  $\Lambda>0$  such that for all  $x\in\Omega$  and  $\lambda\geqslant\Lambda$  we have

$$\|\gamma_{\lambda}^{V}(x,\cdot)\|_{L^{\infty}(\Omega)} \leq C \|\gamma_{\lambda}^{V}(x,\cdot)\|_{L^{\infty}(\partial\Omega)}$$
.

Since  $\|\gamma_{\lambda}^{V}(x,\cdot)\|_{L^{\infty}(\partial\Omega)} = \|T_{\lambda}^{V}(x,\cdot)\|_{L^{\infty}(\partial\Omega)}$ , this together with Lemma 7.1 gives the result. To prove the desired bound, let us introduce the function  $\widetilde{\gamma} := -G_{\lambda}^{0}(V\gamma_{\lambda}^{V}(x,\cdot))$ , so that

$$\begin{cases} (-\Delta + \lambda)\widetilde{\gamma} = -V\gamma_{\lambda}^{V}(x, \cdot) & \text{in } \Omega, \\ \widetilde{\gamma} = 0 & \text{on } \partial\Omega. \end{cases}$$

Then, the function  $\gamma_{\lambda}^{V}(x,\cdot) - \widetilde{\gamma}$  satisfies the equation

$$\begin{cases} (-\Delta + \lambda)(\gamma_{\lambda}^{V}(x,\cdot) - \widetilde{\gamma}) = 0 & \text{in } \Omega, \\ \gamma_{\lambda}^{V}(x,\cdot) - \widetilde{\gamma} = \gamma_{\lambda}^{V}(x,\cdot) & \text{on } \partial \Omega. \end{cases}$$

By the maximum principle, this implies that

$$\|\gamma_{\lambda}^{V}(x,\cdot) - \widetilde{\gamma}\|_{L^{\infty}(\Omega)} \leq \|\gamma_{\lambda}^{V}(x,\cdot)\|_{L^{\infty}(\partial\Omega)}$$

and hence

$$\|\gamma_{\lambda}^{V}(x,\cdot)\|_{L^{\infty}(\Omega)} \leqslant \|\gamma_{\lambda}^{V}(x,\cdot)\|_{L^{\infty}(\partial\Omega)} + \|\widetilde{\gamma}\|_{L^{\infty}(\Omega)}.$$

On the other hand, we have

$$\|\widetilde{\gamma}\|_{L^{\infty}(\Omega)} = \|G_{\lambda}^{0}V\gamma_{\lambda}^{V}(x,\cdot)\|_{L^{\infty}(\Omega)} \leqslant \|G_{\lambda}^{0}V\|_{L^{\infty}\to L^{\infty}} \|\gamma_{\lambda}^{V}(x,\cdot)\|_{L^{\infty}(\Omega)}.$$

Since, again by the maximum principle,  $0 \leqslant G_{\lambda}^{0} \leqslant T_{\lambda}^{0}$  pointwise, we deduce using Lemma A.2 that

$$||G_{\lambda}^{0}V||_{L^{\infty}\to L^{\infty}} \leqslant ||T_{\lambda}^{0}V||_{L^{\infty}\to L^{\infty}} \leqslant C||V||_{\mathcal{K}(2/\sqrt{\lambda})}.$$

For  $\lambda > 0$  large enough so that  $C\|V\|_{\mathcal{K}(2/\sqrt{\lambda})} < 1$ , we deduce that

$$\|\gamma_{\lambda}^{V}(x,\cdot)\|_{L^{\infty}(\Omega)} \leqslant (1 - C\|V\|_{\mathcal{K}(2/\sqrt{\lambda})})^{-1} \|\gamma_{\lambda}^{V}(x,\cdot)\|_{L^{\infty}(\partial\Omega)},$$

which is the desired result.

**Corollary 7.4.** There are C > 0 and  $\Lambda > 0$  such that for all  $x \in \Omega$  and all  $\lambda \geqslant \Lambda$  we have

$$\left|\partial_{\lambda}G_{\lambda}^{V}(x,x)-\partial_{\lambda}T_{\lambda}^{V}(x,x)\right|\leqslant C\frac{e^{-\frac{\sqrt{\lambda}}{2}d(x,\partial\Omega)}}{\lambda d(x,\partial\Omega)}.$$

*Proof.* We first remark that  $\partial_{\lambda}G_{\lambda}^{V}=-(G_{\lambda}^{V})^{2}$ , where the product on the right side is in the sense of operators. Thus,

$$\partial_{\lambda}G_{\lambda}(x,x) = -\int_{\Omega} |G_{\lambda}^{V}(x,y)|^{2} dy.$$

This and the corresponding formula for  $T_{\lambda}^{V}$  gives

$$\left|\partial_{\lambda}G_{\lambda}^{V}(x,x) - \partial_{\lambda}T_{\lambda}^{V}(x,x)\right| \leqslant \int_{\Omega} |G_{\lambda}^{V}(x,y) - T_{\lambda}^{V}(x,y)|(|G_{\lambda}^{V}(x,y)| + |T_{\lambda}^{V}(x,y)|) dy.$$

Using Lemma 7.1, (7.1) and Lemma 7.3, we deduce that

$$\begin{aligned} \left| \partial_{\lambda} G_{\lambda}^{V}(x,x) - \partial_{\lambda} T_{\lambda}^{V}(x,x) \right| &\leqslant C \frac{e^{-\frac{\sqrt{\lambda}}{2}d(x,\partial\Omega)}}{d(x,\partial\Omega)} \int_{\Omega} \frac{e^{-\frac{\sqrt{\lambda}}{2}|x-y|}}{|x-y|} \, dy \\ &\leqslant C \frac{e^{-\frac{\sqrt{\lambda}}{2}d(x,\partial\Omega)}}{\lambda d(x,\partial\Omega)} \,, \end{aligned}$$

which is the desired result.

It will also be useful to have the following uniform bound.

**Lemma 7.5.** There are C > 0 and  $\Lambda > 0$  such that for all  $x \in \Omega$  and all  $\lambda \geqslant \Lambda$  we have

$$|\partial_{\lambda}G_{\lambda}^{V}(x,x)| \leqslant \frac{C}{\sqrt{\lambda}}.$$

*Proof.* We have seen that

$$\sup_{x \in \Omega} |\partial_{\lambda} G_{\lambda}^{V}(x,x)| = \sup_{x \in \Omega} \int_{\Omega} |G_{\lambda}^{V}(x,y)|^{2} dy = \|G_{\lambda}^{V}\|_{L^{2} \to L^{\infty}}^{2}.$$

Now using the resolvent identity as operators

$$G_{\lambda}^{V} = G_{\lambda}^{0} - G_{\lambda}^{0} V G_{\lambda}^{V}$$

we deduce that

$$||G_{\lambda}^{V}||_{L^{2}\to L^{\infty}} \leq ||G_{\lambda}^{0}||_{L^{2}\to L^{\infty}} + ||G_{\lambda}^{0}V||_{L^{\infty}\to L^{\infty}} ||G_{\lambda}^{V}||_{L^{2}\to L^{\infty}}.$$

As in the proof of Lemma 7.3 we have  $||G_{\lambda}^{0}V||_{L^{\infty}\to L^{\infty}} \leqslant C||V||_{\mathcal{K}(2/\sqrt{\lambda})} < 1$  for all large  $\lambda$ , so

$$||G_{\lambda}^{V}||_{L^{2}\to L^{\infty}} \leq (1-C||V||_{\mathcal{K}(2/\sqrt{\lambda})})^{-1}||G_{\lambda}^{0}||_{L^{2}\to L^{\infty}}.$$

Using again the pointwise inequality  $0 \leqslant G_{\lambda}^{0} \leqslant T_{\lambda}^{0}$ , we deduce the desired bound.

One can then use the same strategy as in the previous section to decompose  $\partial_{\lambda}T_{\lambda}^{V}(x,x)$ , and we obtain

**Proposition 7.6.** Let  $V: \Omega \to \mathbb{R}$  in the Kato class. Then there is an  $\varepsilon_0 > 0$  such that for all  $\varepsilon \in (0, \varepsilon_0]$  one has

$$\mathbb{1}(-\Delta_D^{\Omega} + V \leqslant t)(x, x) = \frac{t^{3/2}}{6\pi^2} - \frac{1}{2}r_{0, \varepsilon}^V(t, x) t^{3/2} + \mathcal{O}^{(\varepsilon)}\left(\frac{t}{d(x, \partial\Omega)}\right), \quad t \to +\infty$$

where the  $\mathcal{O}^{(\varepsilon)}$  is uniform in  $x \in \Omega$  (but depends on  $\varepsilon$ ) and where

$$r_{0,\varepsilon}^{V}(t,x) = \sum_{n\geq 1} (-1)^n r_{0,\varepsilon}^{(n,V)}(t,x)$$

with  $r_{0,\varepsilon}^{(n,V)}(t,x)$  defined as in the previous section. More precisely, one has for all  $x \in \Omega$ ,  $t \ge 0$  and  $\varepsilon \in (0,\varepsilon_0]$ ,

$$\left| \mathbb{1}(-\Delta_D^{\Omega} + V \leqslant t)(x, x) - \frac{t^{3/2}}{6\pi^2} + \frac{1}{2}r_{0, \varepsilon}^V(t, x) t^{3/2} \right| \leqslant C_0 \alpha^{-1} (t + \alpha^{-2}),$$

where  $\alpha = \min\{d(x, \partial\Omega), \varepsilon\}$  and where  $C_0$  is an absolute constant.

Proof. We only sketch the main steps in the proof. We apply again Theorem 3.1 in the simplified from (3.1) with  $B_0(t) = -(6\pi^2)^{-1} + \frac{1}{2}r_{0,\varepsilon}^V(t,x)$  and  $B_1(t) = \mathbb{1}(-\Delta_D^\Omega + V \leqslant t)(x,x)$ . Note that no term  $C_{B_2}$  arises in this situation because the remainder  $R_{\lambda,\varepsilon}$  in the parametrix for V=0 vanishes in a ball of radius  $\varepsilon/2$ . Thus, we have  $C_{B_2}=0$  and  $t_0=0$ . The number  $\varepsilon_0$  in the theorem is chosen so that  $\|V\|_{\mathcal{K}(\varepsilon_0)}$  is smaller than an absolute constant, so that the series defining  $r_{0,\varepsilon}^V$  converges and so that Corollary 7.4 is applicable for all  $\lambda \geqslant C/\varepsilon_0^{-2}$ . This, together with the Rodnianski–Schlag bound (A.1), shows that  $\|B_0\|_{L^\infty}$  is bounded by an absolute constant and that  $B_0$  satisfies the almost monotonicity condition for any  $\delta \geqslant 0$  with constant  $C_{B_0}$  bounded by an absolute constant times  $\delta$ . Next, we have  $0 \leqslant B_1(0) \leqslant C\varepsilon_0^{-3}$  with an absolute constant C. This follows as in the proof of Corollary 5.2 from Lemma 7.5. Finally, the decomposition of  $\partial_\lambda T_\lambda^V(x,x)$  leads to an error  $(\varepsilon\lambda)^{-1}e^{-\varepsilon\sqrt{\lambda}/4}$  for  $\lambda \geqslant C\varepsilon^{-2}$  for any  $\varepsilon \in (0,\varepsilon_0]$ . On the other hand, Corollary 7.4 leads to an error  $(d(x,\partial\Omega)\lambda)^{-1}e^{-d(x,\partial\Omega)\sqrt{\lambda}/2}$  for  $\lambda \geqslant C\varepsilon_0^{-2}$ . Together, this gives an error  $(\alpha\lambda)^{-1}e^{-\alpha\sqrt{\lambda}/4}$  for  $\lambda \geqslant C\varepsilon^{-2}$  for any  $\varepsilon \in (0,\varepsilon_0]$ . Thus, we apply Theorem 3.1 with  $\varepsilon_0 = \alpha/2$ ,  $\Lambda = C\alpha^{-2}$  and  $C_0 = C\alpha^{-1}$ . With these choices, the corollary is a consequence of (3.1).

**Corollary 7.7.** Assume that  $\Omega$  has finite measure and satisfies  $|\{x \in \Omega : d(x, \partial\Omega) \leq \delta\}| \leq C\delta$  for all sufficiently small  $\delta > 0$ . Let  $V : \Omega \to \mathbb{R}$  be in the Kato class. Then, one has

$$N(t, -\Delta_{\Omega}^{D} + V) = \frac{t^{3/2}}{6\pi^{2}} |\Omega| + \mathcal{O}(t \log t), \quad t \to +\infty.$$

For V=0 this corollary is due to Courant [15]. Still for V=0, the error estimate  $\mathcal{O}(t \log t)$  was improved by Seeley [51] to the sharp order  $\mathcal{O}(t)$ . We have not tried to adapt Seeley's proof to the case  $V \neq 0$ .

*Proof.* We write  $N(t, -\Delta_{\Omega}^D + V) = \int_{\Omega} \mathbbm{1}(-\Delta_{\Omega}^D + V \leqslant t)(x, x) \, dx$  and split the integral into the region  $\{x \in \Omega: \ d(x, \partial\Omega) > 1/\sqrt{t}\}$  and its complement.

In the set  $\{x \in \Omega : d(x, \partial\Omega) > 1/\sqrt{t}\}$  we use the bound of Proposition 7.6 together with

$$\int_{\Omega} \frac{\mathbb{1}(d(x,\partial\Omega) > 1/\sqrt{t})}{d(x,\partial\Omega)} dx \leqslant C \log t, \tag{7.2}$$

and obtain

$$\begin{split} \int_{\Omega} \mathbb{1}(d(x,\partial\Omega) > 1/\sqrt{t}) \mathbb{1}(-\Delta_{\Omega}^{D} + V \leqslant t)(x,x) \, dx \\ &= \frac{t^{3/2}}{6\pi^{2}} |\{x \in \Omega, \ d(x,\partial\Omega) > 1/\sqrt{t})\}| + \mathcal{O}(t \log t) \\ &= \frac{t^{3/2}}{6\pi^{2}} (|\Omega| - \mathcal{O}(t^{-1/2})) + \mathcal{O}(t \log t) \\ &= \frac{t^{3/2}}{6\pi^{2}} |\Omega| + \mathcal{O}(t \log t). \end{split}$$

To prove (7.2) under the assumptions of the corollary, we write, using the coarea formula and integration by parts,

$$\int_{\Omega} \frac{\mathbb{1}(d(x,\partial\Omega) > 1/\sqrt{t})}{d(x,\partial\Omega)} dx$$

$$= \int_{1/\sqrt{t}}^{\infty} \mathcal{H}^{2}(\{x \in \Omega : d(x,\partial\Omega) = s\}) \frac{ds}{s}$$

$$= \int_{1/\sqrt{t}}^{\infty} |\{x \in \Omega : d(x,\partial\Omega) \leqslant s\}| \frac{ds}{s^{2}} - |\{x \in \Omega : d(x,\partial\Omega) \leqslant 1/\sqrt{t}\}| \sqrt{t}$$

Since  $|\{x \in \Omega : d(x, \partial\Omega) \leq s\}|$  is  $\leq Cs$  for all sufficiently small s and  $\leq |\Omega|$  for all s, we obtain (7.2).

In the set  $\{x \in \Omega : d(x, \partial\Omega) \leq 1/\sqrt{t}\}$  we use the uniform bound of Lemma 7.5, which implies that for all  $\lambda$  large enough one has

$$\frac{1}{(t+\lambda)^2}\mathbb{1}(-\Delta_{\Omega}^D + V \leqslant t)(x,x) \leqslant -\partial_{\lambda}G_{\lambda}^V(x,x) \leqslant \frac{C}{\sqrt{\lambda}}.$$

Taking  $\lambda = t$  gives for all t large enough,

$$\mathbb{1}(-\Delta_{\Omega}^D + V \leqslant t)(x, x) \leqslant Ct^{3/2}.$$

and hence

$$\int_{\Omega} \mathbb{1}(d(x,\partial\Omega) \leqslant 1/\sqrt{t}) \mathbb{1}(-\Delta_{\Omega}^{D} + V \leqslant t)(x,x) \, dx \leqslant C \frac{t^{3/2}}{\sqrt{t}} = Ct \, .$$

Combining the bounds in the two different sets we obtain the claimed asymptotics.  $\Box$ 

### APPENDIX A. SCHRÖDINGER OPERATORS WITH KATO-CLASS POTENTIALS

In this appendix we work in general dimensions  $N \ge 1$ , since this presents no extra effort and since the results might be useful elsewhere. The Kato class was explicitly introduced by Kato [36] in the context of selfadjointness of Schrödinger operators (see also [50]), and developed by Aizenman and Simon [1] as a natural class of potentials to study Schrödinger semigroups. For a comprehensive review, we refer to [54].

## A.1. Kato class potentials. Let

$$w_N(r) = \begin{cases} r^{-(N-2)} & \text{if } N \geqslant 3, \\ \log(1/r) & \text{if } N = 2, \\ 1 & \text{if } N = 1, \end{cases}$$

and set, for any measurable function  $V: \mathbb{R}^N \to \mathbb{R}$ ,

$$||V||_{\mathcal{K}(r)} := \sup_{x \in \mathbb{R}^N} \int_{|y-x| < r} |V(y)| w_N(|x-y|) \, dy.$$

We say that  $V: \mathbb{R}^N \to \mathbb{R}$  belongs to the Kato class if it is measurable and

$$\lim_{r\to 0} \|V\|_{\mathcal{K}(r)} = 0.$$

Remark A.1. When  $N \geqslant 2$ , any potential  $V \in L^p(\mathbb{R}^N) + L^\infty(\mathbb{R}^N)$  for p > N/2 belongs to the Kato class by the Hölder inequality (one can even replace  $L^p(\mathbb{R}^N)$  by  $L^p_{\text{unif}}(\mathbb{R}^N)$ ). However, the Kato class and  $L^{N/2}$  are distinct, and neither one is included in the other. For instance,  $x \mapsto \mathbb{1}(|x| \leqslant 1)/(|x|^2 \log |x|) \in L^{N/2}$  but does not belong to the Kato class; and  $x \mapsto \mathbb{1}(|x| \leqslant 2)(|x|-1)^{-\alpha}$  for  $2/N < \alpha < 1$  belongs to the Kato class but not to  $L^{N/2}$ .

For  $\lambda > 0$  let us introduce the Green's function of the Laplacian,

$$G_{\lambda}(x-y) := (-\Delta + \lambda)^{-1}(x,y) = \frac{1}{(2\pi)^N} \int_{\mathbb{R}^N} \frac{e^{i\xi \cdot (x-y)}}{|\xi|^2 + \lambda} d\xi, \quad x, y \in \mathbb{R}^N.$$

**Lemma A.2.** Let V a Kato class potential. Then, for all  $\lambda \ge 1$  we have

$$||V * G_{\lambda}||_{L^{\infty}} \leqslant C||V||_{\mathcal{K}(2/\sqrt{\lambda})}.$$

The proof is based on the following bounds on the Green's function  $G_{\lambda}$ .

**Lemma A.3.** There is a C > 0 such that for any  $\lambda > 0$  and  $x \neq y \in \mathbb{R}^N$ , the function  $G_{\lambda}$  satisfies: if  $|x - y| \leq 1/\sqrt{\lambda}$ ,

$$|G_{\lambda}(x-y)| \leqslant \begin{cases} \frac{C}{|x-y|^{N-2}} & \text{if } N \geqslant 3, \\ C\log\frac{1}{\sqrt{\lambda}|x-y|} & \text{if } N=2, \\ \frac{C}{\sqrt{\lambda}} & \text{if } N=1, \end{cases}$$

and if  $|x - y| \ge 1/\sqrt{\lambda}$ ,

$$|G_{\lambda}(x-y)| \leqslant C\lambda^{\frac{N-2}{2}} \frac{e^{-\sqrt{\lambda}|x-y|}}{(\sqrt{\lambda}|x-y|)^{\frac{N-1}{2}}} = C\frac{e^{-\sqrt{\lambda}|x-y|}}{|x-y|^{N-2}} (\sqrt{\lambda}|x-y|)^{\frac{N-3}{2}}.$$

This lemma is well-known and there are several ways to prove it. One is by writing  $(|\xi|^2 + \lambda)^{-1}$  as a superposition of Gaussians  $e^{-t|\xi|^2}$  and using the Fourier transform of the Gaussian together with some straightforward estimates. Another way is to recognize  $G_{\lambda}(x-y)$  as an inverse power of |x-y| times a Bessel function of the third kind and using standard estimates for these.

Proof of Lemma A.2. Let  $x \in \mathbb{R}^N$ . Let us split

$$\int_{\mathbb{R}^N} |V(y)| G_{\lambda}(x-y) \, dy = \int_{|x-y| \leqslant 1/\sqrt{\lambda}} |V(y)| G_{\lambda}(x-y) \, dy$$

$$+ \sum_{k \geqslant 1} \int_{k/\sqrt{\lambda} < |x-y| \leqslant (k+1)/\sqrt{\lambda}} |V(y)| G_{\lambda}(x-y) \, dy.$$

Using the first bounds of Lemma A.3, we deduce that

$$\int_{|x-y| \leqslant 1/\sqrt{\lambda}} |V(y)| G_{\lambda}(x-y) \, dy \leqslant ||V||_{\mathcal{K}(1/\sqrt{\lambda})} \sup_{|z| \leqslant 1/\sqrt{\lambda}} \frac{G_{\lambda}(z)}{w_N(|z|)},$$

and since for all  $|z| \leq 1/\sqrt{\lambda}$  we have

$$\frac{G_{\lambda}(z)}{w_{N}(|z|)} \leqslant \begin{cases} C & \text{if } N \geqslant 3, \\ C\frac{\log\frac{1}{\sqrt{\lambda}|z|}}{\log\frac{1}{|z|}} = C(1 + \frac{\log\frac{1}{\sqrt{\lambda}}}{\log\frac{1}{|z|}}) \leqslant 2C & \text{if } N = 2, \\ \frac{C}{\sqrt{\lambda}} \leqslant C & \text{if } N = 1, \end{cases}$$

we deduce that

$$\int_{|x-y| \leqslant 1/\sqrt{\lambda}} |V(y)| G_{\lambda}(x-y) \, dy \leqslant C \|V\|_{\mathcal{K}(1/\sqrt{\lambda})}.$$

For any  $k \ge 1$ , split the annulus

$$A = \{ z \in \mathbb{R}^N : k/\sqrt{\lambda} < |z| \leqslant (k+1)/\sqrt{\lambda} \} = A \cap \bigcup_{j=1}^{M(k,\lambda)} B(z_j, 2/\sqrt{\lambda})$$

in  $M(k,\lambda) \leqslant Ck^{N-1}$  balls of radius  $2/\sqrt{\lambda}$ . Thus, we have

$$\int_{k/\sqrt{\lambda} < |x-y| \leqslant (k+1)/\sqrt{\lambda}} |V(y)| G_{\lambda}(x-y) \, dy \leqslant \sum_{j=1}^{M(k,\lambda)} \int_{\substack{|x-y-z_j| \leqslant 2/\sqrt{\lambda} \\ x-y \in A}} |V(y)| G_{\lambda}(x-y) \, dy$$

$$\leqslant ||V||_{\mathcal{K}(2/\sqrt{\lambda})} \sum_{j=1}^{M(k,\lambda)} \sup_{\substack{|z-z_j| \leqslant 2/\sqrt{\lambda} \\ x \in A}} \frac{G_{\lambda}(z)}{w_N(|z-z_j|)}.$$

Since  $r \mapsto 1/w_N(r)$  is an increasing function, for all  $|z-z_j| \leqslant 2/\sqrt{\lambda}$  we have

$$\frac{1}{w_N(z-z_j)} \leqslant \frac{1}{w_N(2/\sqrt{\lambda})},$$

while, since  $G_{\lambda}$  is radially decreasing,

$$\sum_{j=1}^{M(k,\lambda)} \sup_{\substack{|z-z_j| \leqslant 2/\sqrt{\lambda} \\ z \in A}} \frac{G_{\lambda}(z)}{w_N(|z-z_j|)} \leqslant CM(k,\lambda) \frac{G_{\lambda}(k/\sqrt{\lambda})}{w_N(2/\sqrt{\lambda})}$$

$$\leqslant Ck^{N-1} \frac{\lambda^{(N-2)/2}}{w_N(2/\sqrt{\lambda})} \frac{e^{-k}}{k^{N-2}} k^{(N-3)/2}$$

$$\leqslant Ck^{(N-1)/2} e^{-k}.$$

Summing over k, this leads to the result.

As a consequence of Lemma A.3 we now prove the following quadratic form estimate, following [54, Rk. (2), p. 459].

**Lemma A.4.** Let V in the Kato class. Then, for all  $u \in C_0^{\infty}(\mathbb{R}^N)$  and all  $\lambda \geqslant 1$  we have

$$\int_{\mathbb{R}^N} V|u|^2 dx \leqslant C||V||_{\mathcal{K}(2/\sqrt{\lambda})} \left( \int_{\mathbb{R}^N} |\nabla u|^2 dx + \lambda \int_{\mathbb{R}^N} |u|^2 dx \right).$$

Note that, since the Kato class is included in  $L^1_{loc}(\mathbb{R}^N)$ , the left-side of the previous inequality is well-defined. Moreover, the inequality shows that the quadratic form  $u\mapsto \langle u, Vu\rangle$  extends continuously from  $C_0^\infty(\mathbb{R}^N)$  to  $H^1(\mathbb{R}^N)$ , and that it is relatively bounded with respect to the quadratic form of  $-\Delta$  with relative bound zero (by taking  $\lambda$  large enough). By the KLMN theorem, the quadratic form  $u\mapsto \int (|\nabla u|^2+V|u|^2)\,dx$  defined on  $H^1(\mathbb{R}^N)$  is thus associated to a unique bounded below self-adjoint operator with domain included in  $H^1(\mathbb{R}^N)$ . This is the definition of  $-\Delta + V$  for V in the Kato class.

*Proof.* We prove the estimate for bounded, compactly supported V. Then, replacing V by  $V_n = \mathbb{1}(|V| \leq n)\mathbb{1}(|x| \leq n)V$  and taking the limit  $n \to \infty$  by monotone convergence on the left side and using  $||V_n||_{\mathcal{K}(r)} \leq ||V||_{\mathcal{K}(r)}$  on the right side leads to the result. Now let v a Schwartz function. Define for all  $z \in \mathbb{C}$ ,

$$v_z = |v|^z \frac{v}{|v|}$$

and consider the function

$$\varphi: s \in S \mapsto \left\langle |V|^s v_{2s}, (-\Delta + \lambda)^{-1} |V|^{1-s} v_{2(1-s)} \right\rangle$$

defined on the strip  $S = \{z \in \mathbb{C} : 0 \leq \operatorname{Re} z \leq 1\}$ . Due to the bound for all  $s \in S$  and for all  $x \in \mathbb{R}^N$ ,

$$||V(x)|^s v_{2s}(x)| \leq \mathbb{1}_{\sup V} (1 + ||V||_{L^{\infty}}) (1 + ||v||_{L^{\infty}}^2),$$

we deduce that  $||V|^s v_{2s}||_{L^2} \leq C$ , and since  $||(-\Delta + \lambda)^{-1}||_{L^2 \to L^2} \leq 1$ , we deduce that  $\varphi$  is well-defined, analytic on the interior of S, continuous and bounded on S. Now for s = it we have, by Lemma A.2

$$|\varphi(it)| \leq ||V|^{it} v_{2it}||_{L^{\infty}} ||(-\Delta + \lambda)^{-1}|V|^{1-it}||_{L^{\infty} \to L^{\infty}} ||v_{2(1-it)}||_{L^{1}}$$

$$\leq ||G_{\lambda} * |V|||_{L^{\infty}} ||v||_{L^{2}}^{2}$$

$$\leq C||V||_{\mathcal{K}(2/\sqrt{\lambda})} ||v||_{L^{2}}^{2}.$$

For s = 1 + it we have, again by Lemma A.2

$$\begin{aligned} |\varphi(1+it)| &\leqslant \|v_{2(1+it)}\|_{L^{1}} \|(-\Delta+\lambda)^{-1}|V|^{1+it}\|_{L^{\infty}\to L^{\infty}} \||V|^{-it}v_{-2it}\|_{L^{\infty}} \\ &\leqslant \|G_{\lambda}*|V|\|_{L^{\infty}} \|v\|_{L^{2}}^{2} \\ &\leqslant C\|V\|_{\mathcal{K}(2/\sqrt{\lambda})} \|v\|_{L^{2}}^{2}. \end{aligned}$$

Using Hadamard's three line lemma, we deduce that

$$\|(-\Delta + \lambda)^{-1/2}|V|^{1/2}v\|_{L^2}^2 = |\varphi(1/2)| \leqslant C\|V\|_{\mathcal{K}(2/\sqrt{\lambda})}\|v\|_{L^2}^2.$$

As a consequence, for any Schwartz function w,

$$\langle |V|^{1/2}(-\Delta+\lambda)^{-1/2}w,v\rangle \leqslant C\|V\|_{\mathcal{K}(2/\sqrt{\lambda})}^{1/2}\|v\|_{L^2}\|w\|_{L^2}.$$

Taking the supremum over all v shows that

$$||V|^{1/2}(-\Delta+\lambda)^{-1/2}w||_{L^2} \leqslant C||V||_{\mathcal{K}(2/\sqrt{\lambda})}^{1/2}||w||_{L^2}.$$

Choosing  $w = (-\Delta + \lambda)^{1/2}u$ , we obtain the result.

Finally, we recall the useful bound of Rodnianski and Schlag [46, Lemma 2.5] which is proved for  $M = \mathbb{R}^3$  and  $\varepsilon = +\infty$  but carries easily to the more general following situation:

**Lemma A.5.** Let M a three-dimensional Riemannian manifold and V a Kato class potential on M. Then, for all  $n \ge 1$  and all  $\varepsilon > 0$  we have

$$\sup_{x_0, x_{n+1} \in M \times M} \int_{M_{\varepsilon}(x_0, x_{n+1})} \frac{\prod_{j=1}^n |V(x_j)|}{\prod_{j=0}^n d_g(x_j, x_{j+1})} \sum_{\ell=0}^n d_g(x_\ell, x_{\ell+1}) dv_g(x_1) \cdots dv_g(x_n)$$

$$\leqslant (n+1) \|V\|_{\mathcal{K}(\varepsilon)}^n, \quad (A.1)$$

where  $M_{\varepsilon}(x_0, x_{n+1}) := \{(x_1, \dots, x_n) \in M^n, \forall j = 0, \dots, n, d_g(x_j, x_{j+1}) < \varepsilon \}$  and

$$||V||_{\mathcal{K}(\varepsilon)} := \sup_{x \in M} \int_{d_g(x,y) < \varepsilon} \frac{|V(y)|}{d_g(x,y)} \, dv_g(y).$$

A.2. The case of bounded domains or compact manifolds. If we replace  $\mathbb{R}^N$  by an open set  $\Omega \subset \mathbb{R}^N$  or a compact Riemannian manifold M, one can still define the Kato class by replacing  $\sup_{x \in \mathbb{R}^N}$  by  $\sup_{x \in \Omega}$  or  $\sup_{x \in M}$ , and in the case of a manifold by additionally replacing each |x-y| by the Riemannian distance  $d_g(x,y)$ . In both cases, the same strategy as above shows that V is infinitesimally form bounded with respect to  $-\Delta_{\Omega}^D$  or  $-\Delta_g$ , using in both cases the fact that the Green's functions  $(-\Delta_{\Omega}^D + \lambda)^{-1}(x,y)$  or  $(-\Delta_g + \lambda)^{-1}(x,y)$  can be controlled by the one on  $\mathbb{R}^N$ . This last point can be proved by the maximum principle in the case of a Euclidean domain, or using the estimates of Section 4.1 on a Riemannian manifold. Indeed, it can be shown that the function  $\gamma_{\lambda,\varepsilon}$  satisfies

$$|\gamma_{\lambda,\varepsilon}(x,y)| \leqslant Ce^{-\sqrt{\lambda}d_g(x,y)/4}$$

for  $\varepsilon^2 \lambda \geqslant A'$ , defining

$$B = \sup_{x,y \in M} e^{\sqrt{\lambda} d_g(x,y)/4} |\gamma_{\lambda,\varepsilon}(x,y)|$$

and estimating B using the integral equation satisfied by  $\gamma_{\lambda,\varepsilon}$ . The smallness comes from the estimate

$$\left| \int_{M} \gamma_{\lambda,\varepsilon}(x,z) R_{\lambda,\varepsilon}(z,y) \, dv_g(z) \right| \leqslant C B e^{-\sqrt{\lambda} d_g(x,y)/4} \int_{M} \frac{e^{-\sqrt{\lambda} d_g(z,y)/4}}{\varepsilon^2 d_g(z,y)} \, dv_g(z) \leqslant C e^{-\sqrt{\lambda} d_g(x,y)/4} \frac{B}{\varepsilon^2 \lambda}.$$

The infinitesimal form boundedness also proves that  $-\Delta + V$  either on an open set of finite measure in  $\mathbb{R}^N$  or on a compact manifold has discrete spectrum, since for  $\lambda > 0$  large enough we have

$$-\Delta + V + \lambda + 1 = \varepsilon(-\Delta) + V + \lambda + (1 - \varepsilon)(-\Delta) + 1 \geqslant (1 - \varepsilon)(-\Delta + 1)$$

in the sense of quadratic forms, hence the operator

$$(-\Delta + 1)^{1/2}(-\Delta + V + \lambda + 1)^{-1/2}$$

is bounded. Since the operator  $(-\Delta+1)^{-1/2}$  is compact, this shows that  $(-\Delta+V+\lambda+1)^{-1/2}$  is compact and hence  $-\Delta+V$  has discrete spectrum (and is bounded below).

# Appendix B. Details in the proof of Theorem 3.1

In this appendix we keep track of the various constants in the proof of Theorem 3.1. It will be convenient to abbreviate

$$A_0 := ||B_0||_{L^{\infty}} \Lambda^{3/2} + |B_1(0)| + C_{B_2}(t_0 + \Lambda) + C_0 \Lambda.$$

Step 1. We show that A satisfies the following pointwise bound for all  $t \ge 0$ ,

$$|A(t)| \le C \left( \|B_0\|_{L^{\infty}} t^{3/2} + (C_{B_2} + C_0)t + A_0 \right). \tag{B.1}$$

The starting point of the proof is the bound

$$|A(t)| \le ||B_0||_{L^{\infty}} t^{3/2} + \max\{|B_1(0)|, |B_1(t)|\} + C_{B_2}(t_0 + t)$$
(B.2)

for all  $t \ge 0$ , which follows from the fact that  $B_1$  is nondecreasing. The claimed inequality (B.1) therefore follows if we can show that for all  $t \ge 0$ ,

$$|B_1(t)| \le C \left( ||B_0||_{L^{\infty}} t^{3/2} + (C_{B_2} + C_0)t + A_0 \right). \tag{B.3}$$

Since  $B_1$  is nondecreasing, we have for all  $\lambda > 0$ 

$$\int_0^\infty \frac{B_1(t) - B_1(0)}{(t+\lambda)^3} dt \geqslant \int_\lambda^\infty \frac{B_1(t) - B_1(0)}{(t+\lambda)^3} dt \geqslant \frac{B_1(\lambda) - B_1(0)}{8\lambda^2}.$$
 (B.4)

On the other hand, using

$$\int_0^\infty \frac{|B_0(t)|t^{3/2}}{(t+\lambda)^3} dt \leqslant C \|B_0\|_{L^\infty} \lambda^{-1/2} \quad \text{and} \quad \int_0^\infty \frac{|B_2(t)|}{(t+\lambda)^3} dt \leqslant C C_{B_2}(t_0 \lambda^{-2} + \lambda^{-1}),$$

we obtain because of the assumption on A that for all  $\lambda \geqslant \Lambda$ ,

$$\int_0^\infty \frac{B_1(t) - B_1(0)}{(t+\lambda)^3} dt = \int_0^\infty \frac{A(t) - B_0(t)t^{3/2} - B_2(t) - B_1(0)}{(t+\lambda)^3} dt$$

$$\leq C \left( |B_1(0)|\lambda^{-2} + ||B_0||_{L^\infty} \lambda^{-1/2} + C_{B_2}(t_0\lambda^{-2} + \lambda^{-1}) + C_0 \frac{e^{-\varepsilon_0\sqrt{\lambda}}}{\lambda} \right).$$

Combining this with (B.4) we obtain for all  $\lambda \geqslant \Lambda$ ,

$$B_1(\lambda) - B_1(0) \leqslant C \left( |B_1(0)| + ||B_0||_{L^{\infty}} \lambda^{3/2} + C_{B_2}(t_0 + \lambda) + C_0 \lambda e^{-\varepsilon_0 \sqrt{\lambda}} \right).$$

This, together with the bound

$$|B_1(\lambda)| \le (B_1(\lambda) - B_1(0)) + |B_1(0)| \le (B_1(\max\{\lambda, \Lambda\}) - B_1(0)) + |B_1(0)|,$$

yields (B.3). This completes the proof of (B.1).

Step 2. Next, we derive bounds on  $\int_0^\infty e^{-su} A(u^2) du$  both for  $s = \sigma > 0$  and for  $|s| \le \varepsilon_0/2$ . First, using (B.1) we obtain for all  $\sigma > 0$ 

$$\left| \int_0^\infty e^{-\sigma u} A(u^2) \, du \right| \leqslant C \left( \|B_0\|_{L^\infty} \sigma^{-4} + (C_{B_2} + C_0) \sigma^{-3} + A_0 \sigma^{-1} \right). \tag{B.5}$$

In order to derive a bound for  $|s| \leq \varepsilon_0/2$ , we first note that because of (B.1) we have, for all  $\lambda > 0$ ,

$$\int_0^\infty \frac{|A(t)|}{(t+\lambda)^3} dt \leqslant C \left( \|B_0\|_{L^\infty} \lambda^{-1/2} + (C_{B_2} + C_0) \lambda^{-1} + A_0 \lambda^{-2} \right).$$

Combining this estimate with (3.6) we deduce that for all  $|s| \leq \varepsilon_0/2$ ,

$$\int_0^{\Lambda} \lambda^{3/2} |\kappa(s\sqrt{\lambda})| \int_0^{\infty} \frac{|A(t)|}{(t+\lambda)^3} dt d\lambda \leqslant C\sqrt{\Lambda} e^{\varepsilon_0\sqrt{\Lambda}/2} A_0.$$

On the other hand, using the assumption on A and again (3.6) we deduce that for all  $|s| \leq \varepsilon_0/2$ ,

$$\int_{\Lambda}^{\infty} \lambda^{3/2} |\kappa(s\sqrt{\lambda})| \left| \int_{0}^{\infty} \frac{A(t)}{(t+\lambda)^{3}} dt \right| d\lambda \leqslant CC_{0} \int_{\Lambda}^{\infty} \lambda^{1/2} e^{-\varepsilon_{0}\sqrt{\lambda}} e^{|s|\sqrt{\lambda}} d\lambda \leqslant CC_{0} \varepsilon_{0}^{-3}.$$

To summarize, we have for all  $|s| \leq \varepsilon_0/2$ ,

$$\left| \int_0^\infty e^{-su} A(u^2) \, du \right| \leqslant C \left( \sqrt{\Lambda} e^{\varepsilon_0 \sqrt{\Lambda}/2} A_0 + C_0 \, \varepsilon_0^{-3} \right). \tag{B.6}$$

Step 3. Next, we derive bounds on  $\int_0^\infty e^{-su} dg_0(u)$  both for  $s = \sigma > 0$  and for  $|s| \leq \varepsilon_0/2$ . We first consider the expression with g instead of  $g_0$ . Using  $|A(0)| \leq |B_1(0)| + C_{B_2}t_0$ , we obtain immediately from (B.5) and (B.6) that for all  $\sigma > 0$ 

$$\left| \int_0^\infty e^{-\sigma u} dg(u) \right| \le C \left( \|B_0\|_{L^\infty} \sigma^{-3} + (C_{B_2} + C_0) \sigma^{-2} + A_0 \right)$$

and for all  $|s| \leq \varepsilon_0/2$ 

$$\left| \int_0^\infty e^{-su} dg(u) \right| \leqslant C \left( \varepsilon_0 \sqrt{\Lambda} e^{\varepsilon_0 \sqrt{\Lambda}/2} A_0 + C_0 \varepsilon_0^{-2} + |B_1(0)| + C_{B_2} t_0 \right).$$

This implies that for all  $\sigma > 0$ 

$$\left| \int_0^\infty e^{-\sigma u} dg_0(u) \right| \leqslant C e^{-\sigma u_0} \left( \|B_0\|_{L^\infty} \sigma^{-3} + (C_{B_2} + C_0) \sigma^{-2} + A_0 \right)$$
 (B.7)

and for all  $|s| \leq \varepsilon_0/2$ 

$$\left| \int_0^\infty e^{-su} dg_0(u) \right| \leqslant C e^{\varepsilon_0 u_0/2} \left( \varepsilon_0 \sqrt{\Lambda} e^{\varepsilon_0 \sqrt{\Lambda}/2} A_0 + C_0 \varepsilon_0^{-2} + |B_1(0)| + C_{B_2} t_0 \right). \tag{B.8}$$

Step 4. Now we prove bounds on f(0) and f'(s) for  $|s| \leq \varepsilon_0/2$ . According to (3.9), f is the second antiderivative of  $s \mapsto \int_0^\infty e^{-su} dg_0(u)$ , normalized to vanish at infinity. To estimate the antiderivative for  $|s| \leq \varepsilon_0/2$ , we choose a path of integration which goes from s to  $\varepsilon_0/2$  and from  $\varepsilon_0/2$  to  $+\infty$  in straight lines. To estimate it for  $\sigma > 0$ , we simply integrate from

 $\sigma$  to  $+\infty$  in a straight line. Using the bounds from the previous step we find easily that for all  $\sigma > 0$ ,

$$|f'(\sigma)| \le Ce^{-\sigma u_0} \Big( ||B_0||_{L^{\infty}} \sigma^{-2} + (C_{B_2} + C_0)\sigma^{-1} + u_0^{-1} A_0 \Big)$$

and for all  $|s| \leq \varepsilon_0/2$ ,

$$|f'(s)| \leq C\varepsilon_0 e^{\varepsilon_0 u_0/2} \left( \varepsilon_0 \sqrt{\Lambda} e^{\varepsilon_0 \sqrt{\Lambda}/2} A_0 + C_0 \varepsilon_0^{-2} + |B_1(0)| + C_{B_2} t_0 \right) + C e^{-\varepsilon_0 u_0/2} \left( ||B_0||_{L^{\infty}} \varepsilon_0^{-2} + (C_{B_2} + C_0) \varepsilon_0^{-1} + u_0^{-1} A_0 \right).$$

Integrating once more, we obtain from these two bounds that

$$|f(0)| \leq C\varepsilon_0^2 e^{\varepsilon_0 u_0/2} \left( \varepsilon_0 \sqrt{\Lambda} e^{\varepsilon_0 \sqrt{\Lambda}/2} A_0 + C_0 \varepsilon_0^{-2} + |B_1(0)| + C_{B_2} t_0 \right)$$

$$+ C e^{-\varepsilon_0 u_0/2} \left( ||B_0||_{L^{\infty}} \varepsilon_0^{-1} + (C_{B_2} + C_0) \left( 1 + \ln_{+\frac{1}{\varepsilon_0 u_0}} \right) + (1 + u_0 \varepsilon_0) u_0^{-2} A_0 \right).$$
 (B.9)

This is the desired bound on a = f(0). Moreover, if we define  $\varphi(s) = (f(s) - a)/s$ , then for all  $|s| \leq \varepsilon_0/2$ ,

$$|\varphi(s)| \leqslant \sup_{|s'| \leqslant \varepsilon_0/2} |f'(s')| \leqslant C\varepsilon_0 e^{\varepsilon_0 u_0/2} \left( \varepsilon_0 \sqrt{\Lambda} e^{\varepsilon_0 \sqrt{\Lambda}/2} A_0 + C_0 \varepsilon_0^{-2} + |B_1(0)| + C_{B_2} t_0 \right) + C e^{-\varepsilon_0 u_0/2} \left( ||B_0||_{L^{\infty}} \varepsilon_0^{-2} + (C_{B_2} + C_0) \varepsilon_0^{-1} + u_0^{-1} A_0 \right).$$
(B.10)

Thus, setting  $T = \varepsilon_0/2$ , we obtain

$$|a| + \int_{-T}^{T} |\varphi(it)| dt \leqslant C\varepsilon_0^2 e^{\varepsilon_0 u_0/2} \left( \varepsilon_0 \sqrt{\Lambda} e^{\varepsilon_0 \sqrt{\Lambda}/2} A_0 + C_0 \varepsilon_0^{-2} + |B_1(0)| + C_{B_2} t_0 \right)$$

$$+ Ce^{-\varepsilon_0 u_0/2} \left( ||B_0||_{L^{\infty}} \varepsilon_0^{-1} + (C_{B_2} + C_0) \left( 1 + \ln_{+} \frac{1}{\varepsilon_0 u_0} \right) + (1 + \varepsilon_0 u_0) u_0^{-2} A_0 \right).$$
 (B.11)

Step 5. We now show that the bound  $h(v) - h(u) \ge -c$  holds for all  $0 \le u \le v \le u + \delta$  with

$$c \leq C \left( C_{B_0} + ||B_0||_{L^{\infty}} (\delta + u_0) + C_{B_2} \left( 1 + \ln(1 + \delta u_0^{-1}) + t_0 u_0^{-2} \right) \right).$$
 (B.12)

To prove this, we decompose

$$g_0(u) = \widetilde{B_0}(u)u^3 + \widetilde{B_1}(u) + \widetilde{B_2}(u)$$

where

$$\widetilde{B_0}(u) = \begin{cases} B_0((u - u_0)^2) & \text{if } u \geqslant u_0, \\ B_0(0) & \text{if } u < u_0, \end{cases}, \quad \widetilde{B_1}(u) = \begin{cases} B_1((u - u_0)^2) - B_1(0) & \text{if } u \geqslant u_0, \\ 0 & \text{if } u < u_0, \end{cases}$$

$$\widetilde{B_2}(u) = \begin{cases} ((u - u_0)^3 - u^3)B_0((u - u_0)^2) + B_2((u - u_0)^2) - B_2(0) & \text{if } u \geqslant u_0, \\ -u^3B_0(0) & \text{if } u < u_0, \end{cases}$$

We have, for all  $0 \le u \le v$ ,

$$h(v) - h(u) = \int_{(u,v]} \frac{dg_0(w)}{w^2} = \int_{(u,v]} \frac{d(\widetilde{B_0}(w)w^3)}{w^2} + \int_{(u,v]} \frac{d\widetilde{B_1}(w)}{w^2} + \int_{(u,v]} \frac{d\widetilde{B_2}(w)}{w^2}$$
(B.13)

and we discuss separately the three terms on the right side. The function  $\widetilde{B_1}$  is non-decreasing since  $B_1$  is non-decreasing and therefore the second term is nonnegative.

For the first term in (B.13) we have if  $u \leq v \leq u + \delta$ , using the right-continuity of  $w \mapsto w^3 \widetilde{B_0}(w^2)$ ,

$$\int_{(u,v]} \frac{d(\widetilde{B_0}(w)w^3)}{w^2} = \widetilde{B_0}(v)v - \widetilde{B_0}(u)u + 2\int_u^v \widetilde{B_0}(w) dw$$

$$= \widetilde{B_0}(v)(v-u) + (\widetilde{B_0}(v) - \widetilde{B_0}(u))u + 2\int_u^v \widetilde{B_0}(w) dw$$

$$\geqslant -3\|B_0\|_{L^{\infty}}\delta + (\widetilde{B_0}(v) - \widetilde{B_0}(u))u.$$

If  $u \leq u_0$ , we simply bound

$$(\widetilde{B_0}(v) - \widetilde{B_0}(u))u \geqslant -2\|B_0\|_{L^{\infty}}u_0$$
.

If  $u > u_0$  we use the assumption on  $B_0$  to deduce that

$$\widetilde{B_0}(v) - \widetilde{B_0}(u) \geqslant -C_{B_0}/(u - u_0),$$

and hence

$$(\widetilde{B}_0(v) - \widetilde{B}_0(u))u = (\widetilde{B}_0(v) - \widetilde{B}_0(u))(u - u_0) + (\widetilde{B}_0(v) - \widetilde{B}_0(u))u_0$$
  
 $\geqslant -C_{B_0} - 2||B_0||_{L^{\infty}}u_0.$ 

To summarize, for all  $0 \le u \le v \le u + \delta$ ,

$$\int_{(u,v]} \frac{d(\widetilde{B_0}(w)w^3)}{w^2} \geqslant -(C_{B_0} + ||B_0||_{L^{\infty}}(3\delta + 2u_0)).$$

For the third term in (B.13) we first note that if  $u < u_0$ , then

$$\int_{(u,\min\{v,u_0\}]} \frac{dB_2(w)}{w^2} = -3B_0(0) \left(\min\{v,u_0\} - u\right).$$

If  $v \leq u + \delta$ , the right side is  $\geq -3\|B_0\|_{L^{\infty}}\delta$ . Thus, in the following we will bound the integral under the assumption that  $u \geq u_0$ . Since  $\widetilde{B_2}$  is right-continuous, we write

$$\int_{(u,v]} \frac{d\widetilde{B_2}(w)}{w^2} = \frac{\widetilde{B_2}(v)}{v^2} - \frac{\widetilde{B_2}(u)}{u^2} + 2 \int_u^v \frac{\widetilde{B_2}(w)}{w^3} dw.$$

We use  $|u^3 - (u - u_0)^3| \le Cu_0(u_0^2 + u^2) \le Cu_0u^2$  to get

$$|\widetilde{B_2}(u)| \le C \left( u_0 u^2 ||B_0||_{L^{\infty}} + C_{B_2} (t_0 + (u - u_0)^2) \right).$$

Therefore, since  $u_0 \leq u$ ,

$$\frac{|\widetilde{B_2}(u)|}{u^2} \leqslant C\left(u_0 \|B_0\|_{L^{\infty}} + \frac{C_{B_2}t_0}{u^2} + \frac{C_{B_2}(u - u_0)^2}{u^2}\right) \leqslant C\left(u_0 \|B_0\|_{L^{\infty}} + \frac{C_{B_2}t_0}{u_0^2} + C_{B_2}\right)$$

and, using

$$\int_{u}^{u+\delta} \frac{(w-u_0)^2}{w^3} dw \leqslant C \left(1 + \ln\left(1 + \frac{\delta}{u_0}\right)\right),\,$$

also

$$\int_{u}^{v} \frac{|\widetilde{B}_{2}(w)|}{w^{3}} dw \leqslant C \left( u_{0} \|B_{0}\|_{L^{\infty}} \ln \frac{v}{u} + C_{B_{2}} t_{0} \left( \frac{1}{u^{2}} - \frac{1}{v^{2}} \right) + C_{B_{2}} \left( 1 + \ln \left( 1 + \frac{\delta}{u_{0}} \right) \right) \right)$$

$$\leqslant C \left( u_{0} \|B_{0}\|_{L^{\infty}} \ln \left( 1 + \delta u_{0}^{-1} \right) + C_{B_{2}} \left( 1 + \ln \left( 1 + \delta u_{0}^{-1} \right) + t_{0} u_{0}^{-2} \right) \right).$$

To summarize, for all  $0 \leqslant u \leqslant v \leqslant u + \delta$ ,

$$\int_{(u,v]} \frac{d\widetilde{B_2}(w)}{w^2} \geqslant -C\left(\|B_0\|_{L^{\infty}} (u_0 + \delta) + C_{B_2} \left(1 + \ln(1 + \delta u_0^{-1}) + t_0 u_0^{-2}\right)\right).$$

Combining our bounds on the three terms in (B.13) gives (B.12).

Step 6. Applying Theorem 2.1 to h with  $T = \varepsilon_0/2$  and using the bounds (B.11) and (B.12), we find that for all  $u \ge 0$  we have

$$|h(u)| \leq C\varepsilon_0^2 e^{\varepsilon_0 u_0/2} \left( \varepsilon_0 \sqrt{\Lambda} e^{\varepsilon_0 \sqrt{\Lambda}/2} A_0 + C_0 \varepsilon_0^{-2} + |B_1(0)| + C_{B_2} t_0 \right)$$

$$+ C e^{-\varepsilon_0 u_0/2} \left( ||B_0||_{L^{\infty}} \varepsilon_0^{-1} + (C_{B_2} + C_0) \left( 1 + \ln_+ \frac{1}{\varepsilon_0 u_0} \right) + (1 + \varepsilon_0 u_0) u_0^{-2} A_0 \right)$$

$$+ C \left( 1 + \delta^{-1} \varepsilon_0^{-1} \right) \left( C_{B_0} + ||B_0||_{L^{\infty}} (\delta + u_0) + C_{B_2} \left( 1 + \ln(1 + \delta u_0^{-1}) + t_0 u_0^{-2} \right) \right)$$

$$=: \Theta.$$

Thus, if  $x \geqslant u_0$ ,

$$|A((x-u_0)^2) - A(0)| = |g_0(x)| = \left| x^2 h(x) - 2 \int_0^x u h(u) \, du \right| \le 2\Theta x^2$$

and thus, if  $u \ge 0$ ,

$$|A(u)| \le |A(0)| + 2\Theta(u_0 + \sqrt{t})^2 \le |B_1(0)| + C_{B_2}t_0 + 4\Theta(u_0^2 + t)$$
.

Note that with the choice  $u_0 = \varepsilon_0^{-1}$  we get

$$\Theta \leqslant C\varepsilon_0^2 \left( \left( 1 + \varepsilon_0 \sqrt{\Lambda} e^{\varepsilon_0 \sqrt{\Lambda}/2} \right) A_0 + C_0 \varepsilon_0^{-2} \right) 
+ C \left( 1 + \delta^{-1} \varepsilon_0^{-1} \right) \left( C_{B_0} + \|B_0\|_{L^{\infty}} \left( \delta + \varepsilon_0^{-1} \right) + C_{B_2} \left( 1 + \ln(1 + \delta \varepsilon_0) + t_0 \varepsilon_0^2 \right) \right)$$

and therefore, finally,

$$|A(u)| \leq C\varepsilon_0^2 \left( \left( 1 + \varepsilon_0 \sqrt{\Lambda} e^{\varepsilon_0 \sqrt{\Lambda}/2} \right) A_0 + C_0 \varepsilon_0^{-2} \right) \left( t + \varepsilon_0^{-2} \right) + C \left( 1 + \delta^{-1} \varepsilon_0^{-1} \right) \left( C_{B_0} + ||B_0||_{L^{\infty}} \left( \delta + \varepsilon_0^{-1} \right) + C_{B_2} \left( 1 + \ln(1 + \delta \varepsilon_0) + t_0 \varepsilon_0^2 \right) \right) \left( t + \varepsilon_0^{-2} \right).$$
(B.14)

This proves the bound claimed in Theorem 3.1.

### References

- [1] M. AIZENMAN AND B. SIMON, Brownian motion and Harnack inequality for Schrödinger operators, Comm. Pure Appl. Math., 35 (1982), pp. 209–273.
- [2] V. G. AVAKUMOVIĆ, Bemerkung über einen Satz des Herrn T. Carleman, Math. Z., 53 (1950), pp. 53–58.
- [3] ——, Über die Eigenfunktionen der Schwingungsgleichung, Acad. Serbe Sci. Publ. Inst. Math., 4 (1952), pp. 95–96.

- [4] ——, Über die Eigenfunktionen auf geschlossenen Riemannschen Mannigfaltigkeiten, Math. Z., 65 (1956), pp. 327–344.
- [5] P. H. BÉRARD, On the wave equation on a compact Riemannian manifold without conjugate points, Math. Z., 155 (1977), pp. 249–276.
- [6] M. BERGER, P. GAUDUCHON, AND E. MAZET, Le spectre d'une variété riemannienne, Lecture Notes in Mathematics, Vol. 194, Springer-Verlag, Berlin-New York, 1971.
- [7] M. S. BIRMAN AND A. LAPTEV, The negative discrete spectrum of a two-dimensional Schrödinger operator, Comm. Pure Appl. Math., 49 (1996), pp. 967–997.
- [8] M. D. Blair, Y. Sire, and C. D. Sogge, Quasimode, eigenfunction and spectral projection bounds for Schrödinger operators on manifolds with critically singular potentials, The Journal of Geometric Analysis, (2019), pp. 1–38.
- [9] P. Blanchard, J. Stubbe, and J. Rezende, New estimates on the number of bound states of Schrödinger operators, Lett. Math. Phys., 14 (1987), pp. 215–225.
- [10] R. Bojanić, Propriétés asymptotiques des solutions des équations différentielles lineáries, Srpska Akad. Nauka Zb. Rad. Mat. Inst., 35(3) (1953), pp. 213–254.
- [11] T. Carleman, Propriétés asymptotiques des fonctions fondamentales des membranes vibrantes, Comptes Rendus des Mathématiciens Scandinaves á Stockholm, (1934), pp. 14–18.
- [12] T. Carleman, Uber die asymptotische verteilung der eigenwerte partielle differentialgleichungen, Berichten der mathematisch-physisch Klasse der Sächsischen Akad. der Wissenschaften zu Leipzig, LXXXVIII Band, Sitsung, 15 (1936).
- [13] M. CARTER AND B. VAN BRUNT, *The Lebesgue-Stieltjes integral*, Undergraduate Texts in Mathematics, Springer-Verlag, New York, 2000. A practical introduction.
- [14] J. G. CONLON, A new proof of the Cwikel-Lieb-Rosenbljum bound, Rocky Mountain J. Math., 15 (1985), pp. 117–122.
- [15] R. COURANT, Uber die Eigenwerte bei den Differentialgleichungen der mathematischen Physik, Math. Z., 7 (1920), pp. 1–57.
- [16] M. CWIKEL, Weak type estimates for singular values and the number of bound states of Schrödinger operators, Ann. of Math. (2), 106 (1977), pp. 93–100.
- [17] H. Donnelly, Bounds for eigenfunctions of the Laplacian on compact Riemannian manifolds, J. Funct. Anal., 187 (2001), pp. 247–261.
- [18] D. Dos Santos Ferreira, C. E. Kenig, and M. Salo, On L<sup>p</sup> resolvent estimates for Laplace-Beltrami operators on compact manifolds, Forum Math., 26 (2014), pp. 815–849.
- [19] J. J. Duistermaat and V. W. Guillemin, The spectrum of positive elliptic operators and periodic bicharacteristics, Invent. Math., 29 (1975), pp. 39–79.
- [20] C. Fefferman and L. Seco, *The density in a one-dimensional potential*, Adv. Math., 107 (1994), pp. 187–364.
- [21] ——, The density in a three-dimensional radial potential, Adv. Math., 111 (1995), pp. 88–161.
- [22] C. L. FEFFERMAN, The uncertainty principle, Bull. Amer. Math. Soc. (N.S.), 9 (1983), pp. 129–206.
- [23] R. L. Frank, Cwikel's theorem and the CLR inequality, J. Spectr. Theory, 4 (2014), pp. 1–21.
- [24] R. L. Frank, E. H. Lieb, and R. Seiringer, Equivalence of Sobolev inequalities and Lieb-Thirring inequalities, in XVIth International Congress on Mathematical Physics, World Sci. Publ., Hackensack, NJ, 2010, pp. 523–535.
- [25] W. Gromes, Über das asymptotische Verhalten der Spektralfunktion elliptischer Systeme, Math. Z., 118 (1970), pp. 254–270.
- [26] D. R. Heath-Brown, *Lattice points in the sphere*, in Number theory in progress, Vol. 2 (Zakopane-Kościelisko, 1997), de Gruyter, Berlin, 1999, pp. 883–892.
- [27] B. Helffer, A. Knauf, H. Siedentop, and R. Weikard, On the absence of a first order correction for the number of bound states of a Schrödinger operator with Coulomb singularity, Comm. Partial Differential Equations, 17 (1992), pp. 615–639.

- [28] E. HLAWKA, Über Integrale auf konvexen Körpern. I, Monatsh. Math., 54 (1950), pp. 1–36.
- [29] L. HÖRMANDER, The spectral function of an elliptic operator, Acta Math., 121 (1968), pp. 193–218.
- [30] X. Huang and C. D. Sogge, Weyl formulae for Schrödinger operators with critically singular potentials, arXiv preprint arXiv:2005.10323, (2020).
- [31] W. Hughes, Thermodynamics for Coulomb systems: a problem at vanishing particle densities, J. Statist. Phys., 41 (1985), pp. 975–1013.
- [32] M. N. Huxley, Exponential sums and lattice points. III, Proc. London Math. Soc. (3), 87 (2003), pp. 591–609.
- [33] A. E. Ingham, On Wiener's Method in Tauberian Theorems, Proc. London Math. Soc. (2), 38 (1935), pp. 458–480.
- [34] J. KARAMATA, Weiterführung der N. Wienerschen Methode, Math. Z., 38 (1934), pp. 701–708.
- [35] J. KARAMATA, Ueber einen satz von H. Heilbronn und E. Landau, Publ. Inst. Math. Beograd, 5 (1936), pp. 28–38.
- [36] T. Kato, Schrödinger operators with singular potentials, Israel J. Math., 13 (1972), pp. 135–148 (1973).
- [37] E. Landau, Vorlesungen über zahlentheorie, 1927.
- [38] A. Laptev, Asymptotics of the negative discrete spectrum of a class of Schrödinger operators with large coupling constant, Proc. Amer. Math. Soc., 119 (1993), pp. 481–488.
- [39] D. LEVIN AND M. SOLOMYAK, The Rozenblum-Lieb-Cwikel inequality for Markov generators, J. Anal. Math., 71 (1997), pp. 173–193.
- [40] B. M. LEVITAN, On the asymptotic behavior of the spectral function of a self-adjoint differential equation of the second order, Izvestiya Akad. Nauk SSSR. Ser. Mat., 16 (1952), pp. 325–352.
- [41] P. Li and S. T. Yau, On the Schrödinger equation and the eigenvalue problem, Comm. Math. Phys., 88 (1983), pp. 309–318.
- [42] E. Lieb, Bounds on the eigenvalues of the Laplace and Schroedinger operators, Bull. Amer. Math. Soc., 82 (1976), pp. 751–753.
- [43] E. H. LIEB, The number of bound states of one-body Schroedinger operators and the Weyl problem, in Geometry of the Laplace operator (Proc. Sympos. Pure Math., Univ. Hawaii, Honolulu, Hawaii, 1979), Proc. Sympos. Pure Math., XXXVI, Amer. Math. Soc., Providence, R.I., 1980, pp. 241–252.
- [44] S. MINAKSHISUNDARAM AND A. . PLEIJEL, Some properties of the eigenfunctions of the Laplace-operator on Riemannian manifolds, Canad. J. Math., 1 (1949), pp. 242–256.
- [45] D. Pauly, Über die bilinearentwicklung der greenschen funktionen, (1965). Staatsexamensarbeit, Aachen.
- [46] I. Rodnianski and W. Schlag, Time decay for solutions of Schrödinger equations with rough and time-dependent potentials, Invent. Math., 155 (2004), pp. 451–513.
- [47] G. V. ROZENBLJUM, Distribution of the discrete spectrum of singular differential operators, Dokl. Akad. Nauk SSSR, 202 (1972), pp. 1012–1015.
- [48] ——, Distribution of the discrete spectrum of singular differential operators, Izv. Vysš. Učebn. Zaved. Matematika, (1976), pp. 75–86.
- [49] Y. G. Safarov, Asymptotics of a spectral function of a positive elliptic operator without a nontrapping condition, Funktsional. Anal. i Prilozhen., 22 (1988), pp. 53–65, 96.
- [50] M. SCHECHTER, Spectra of partial differential operators, North-Holland Publishing Co., Amsterdam-London; American Elsevier Publishing Co., Inc., New York, 1971. North-Holland Series in Applied Mathematics and Mechanics, Vol. 14.
- [51] R. SEELEY, A sharp asymptotic remainder estimate for the eigenvalues of the Laplacian in a domain of  $\mathbb{R}^3$ , Adv. in Math., 29 (1978), pp. 244–269.
- [52] H. SIEDENTOP AND R. WEIKARD, On the leading energy correction for the statistical model of the atom: interacting case, Comm. Math. Phys., 112 (1987), pp. 471–490.
- [53] ——, On the leading correction of the Thomas-Fermi model: lower bound, Invent. Math., 97 (1989), pp. 159–193. With an appendix by A. M. Klaus Müller.

- [54] B. Simon, Schrödinger semigroups, Bull. Amer. Math. Soc. (N.S.), 7 (1982), pp. 447–526.
- [55] H. F. SMITH, Spectral cluster estimates for C<sup>1,1</sup> metrics, Amer. J. Math., 128 (2006), pp. 1069–1103.
- [56] A. V. Sobolev, Discrete spectrum asymptotics for the Schrödinger operator with a singular potential and a magnetic field, Rev. Math. Phys., 8 (1996), pp. 861–903.
- [57] C. D. SOGGE AND S. ZELDITCH, Riemannian manifolds with maximal eigenfunction growth, Duke Math. J., 114 (2002), pp. 387–437.
- [58] J. P. Solovej and W. L. Spitzer, A New Coherent States Approach to Semiclassics Which Gives Scott's Correction, Commun. Math. Phys., 241 (2003), pp. 383–420.
- [59] E. C. Titchmarsh, Eigenfunction expansions associated with second-order differential equations. Vol. 2, Oxford, at the Clarendon Press, 1958.
- [60] A. Walfisz, Über Gitterpunkte in vierdimensionalen Ellipsoiden, Math. Z., 72 (1959/60), pp. 259–278.
- [61] H. Weyl, Über die asymptotische verteilung der eigenwerte, Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse, 1911 (1911), pp. 110–117.
- (R. L. Frank) Mathematisches Institut, Ludwig-Maximilians-Universität München, Theresienstr. 39, 80333 München, Germany, and Mathematics 253-37, Caltech, Pasadena, CA 91125, USA

E-mail address: r.frank@lmu.de, rlfrank@caltech.edu

(J. Sabin) Laboratoire de Mathématiques d'Orsay, Univ. Paris-Sud, CNRS, Université Paris-Saclay, 91405 Orsay, France

E-mail address: julien.sabin@universite-paris-saclay.fr