Kolmogorov widths of the intersection of two finite-dimensional balls

A.A. Vasil'eva

In this paper we study the problem on estimating the Kolmogorov widths of the set $B_{p_0}^m \cap \nu B_{p_1}^m$ in l_q^m .

First we recall some definitions.

Let X be a normed space, and let $M \subset X$, $n \in \mathbb{Z}_+$. The Kolmogorov *n*-width of the set M in the space X is defined as

$$d_n(M, X) = \inf_{L \in \mathcal{L}_n(X)} \sup_{x \in M} \inf_{y \in L} ||x - y||;$$

here $\mathcal{L}_n(X)$ is the family of subspaces in X of dimension at most n.

Let $1 \leq p < \infty$. We denote by l_p^m the linear space \mathbb{R}^m with the norm

$$||(x_1, \ldots, x_m)||_{l_p^m} = \left(\sum_{j=1}^m |x_j|^p\right)^{1/p};$$

by B_p^m we denote the unit ball in l_p^m . For $p = \infty$ the norm is defined as

 $||(x_1, \ldots, x_m)||_{l_{\infty}^m} = \max\{|x_1|, \ldots, |x_m|\}.$

The problem on estimating the widths $d_n(B_p^m, l_q^m)$ was studied by A. Pietsch, M.I. Stesin, B.S. Kashin, E.D. Gluskin, A.Yu. Garnaev (see, e.g., [1–7]). Here we use and formulate these results for some particular cases. First we introduce notation for order equalities and inequalities. Let X, Y be sets, $f_1, f_2 : X \times Y \to \mathbb{R}_+$. We write $f_1(x, y) \underset{y}{\leq} f_2(x, y)$ (or $f_2(x, y) \underset{y}{\geq} f_1(x, y)$) if for each $y \in Y$ there is c(y) > 0 such that $f_1(x, y) \leqslant c(y)f_2(x, y)$ for all $x \in X$; $f_1(x, y) \underset{y}{\asymp} f_2(x, y)$ if $f_1(x, y) \underset{y}{\lesssim} f_2(x, y)$ and $f_2(x, y) \underset{y}{\leq} f_1(x, y)$.

Theorem A. [6] Let $1 \leq p \leq q < \infty$, $0 \leq n \leq m/2$.

1. Let
$$1 \leq q \leq 2$$
. Then $d_n(B_p^m, l_q^m) \underset{p,q}{\asymp} 1$.
2. Let $2 < q < \infty$, $\lambda_{pq} = \min\left\{1, \frac{1/p - 1/q}{1/2 - 1/q}\right\}$. Then
 $d_n(B_p^m, l_q^m) \underset{p,q}{\asymp} \min\{1, n^{-1/2}m^{1/q}\}^{\lambda_{pq}}$

Theorem B. [3,4]. Let $1 \leq q \leq p \leq \infty$, $0 \leq n \leq m$. Then

$$d_n(B_p^m, l_q^m) = (m-n)^{1/q-1/p}$$

The problem on estimating the Kolmogorov widths of intersections of finitedimensional balls naturally appears when estimating the Kolmogorov widths of intersections of function classes (see, e.g., [8–10]). Notice that in [8] the intersection of the arbitrary family of balls was considered, but the order estimate for the widths was obtained only for m = 2n. In [10] the problem on estimating the Kolmogorov widths of some function classes in a weighted L_q -space was studied; these classes were defined by constraints on the weighted L_{p_1} -norm of the highest-order derivatives and on the weighted L_{p_0} -norm of the functions. The lower estimates for the widths were reduced to estimating the widths of B^m_{∞} and B^m_1 in l^m_q ; the upper estimates were reduced to estimating the widths of $B^m_{p_0}$, $B^m_{p_1}$, B^m_2 and B^m_q (here the particular case of Theorem 2 from [8] was applied). The problem on estimating the widths of $B^m_{p_0} \cap \nu B^m_{p_1}$ was not studied in [10].

Let $1 \leq p_1 < p_0 \leq \infty$. First we notice that $B_{p_0}^m \cap \nu B_{p_1}^m = \nu B_{p_1}^m$ for $\nu \leq 1$, and $B_{p_0}^m \cap \nu B_{p_1}^m = B_{p_0}^m$ for $\nu \geq m^{1/p_1 - 1/p_0}$. Hence, it suffices to consider the case $1 < \nu < m^{1/p_1 - 1/p_0}$. Further, $B_{p_1}^m \subset B_{p_0}^m \cap \nu B_{p_1}^m \subset B_{p_0}^m$ for $\nu \geq 1$. Therefore, Theorem A implies that for $n \leq m/2$

$$d_n(B_{p_0}^m \cap \nu B_{p_1}^m, l_q^m) \underset{p,q}{\asymp} 1 \quad \text{if } 1 \leqslant p_1 \leqslant p_0 \leqslant q \leqslant 2,$$
$$d_n(B_{p_0}^m \cap \nu B_{p_1}^m, l_q^m) \underset{p,q}{\asymp} \min\{1, n^{-1/2} m^{1/q}\} \quad \text{if } 1 \leqslant p_1 \leqslant p_0 \leqslant 2 < q < \infty.$$

Now, we estimate the Kolmogorov widths for other cases. Notice that it suffices to consider $\nu = k^{1/p_1-1/p_0}$, $k = 1, \ldots, m$.

Theorem 1. Let $p_0 > p_1$, $m \in \mathbb{N}$, $n \in \mathbb{Z}_+$, $n \leqslant m/2$, $1 \leqslant k \leqslant m$, $\nu = k^{1/p_1 - 1/p_0}$.

1. Let $1 \leq p_1 < 2 < p_0 < q < \infty$. Then

$$d_{n}(B_{p_{0}}^{m} \cap \nu B_{p_{1}}^{m}, l_{q}^{m}) \underset{p_{0}, p_{1}, q}{\asymp} \begin{cases} 1, & \text{for } n \leqslant m^{2/q}, \\ (n^{-1/2}m^{1/q})^{\frac{1/p_{0}-1/q}{1/2-1/q}}, & \text{for } m^{\frac{2}{q}} \leqslant n \leqslant k^{1-\frac{2}{q}}m^{\frac{2}{q}}, \\ k^{\frac{1}{2}-\frac{1}{p_{0}}}n^{-\frac{1}{2}}m^{\frac{1}{q}}, & \text{for } k^{1-\frac{2}{q}}m^{\frac{2}{q}} \leqslant n \leqslant m/2. \end{cases}$$

2. Let $2 \leq p_1 < p_0 < q < \infty$. Then

$$d_{n}(B_{p_{0}}^{m}\cap\nu B_{p_{1}}^{m}, l_{q}^{m}) \underset{p_{0},p_{1},q}{\asymp} \begin{cases} 1, & for \quad n \leqslant m^{2/q}, \\ (n^{-1/2}m^{1/q})^{\frac{1/p_{0}-1/q}{1/2-1/q}}, & for \quad m^{\frac{2}{q}} \leqslant n \leqslant k^{1-\frac{2}{q}}m^{\frac{2}{q}}, \\ k^{\frac{1}{p_{1}}-\frac{1}{p_{0}}} \left(n^{-\frac{1}{2}}m^{\frac{1}{q}}\right)^{\frac{1/p_{1}-1/q}{1/2-1/q}}, & for \quad k^{1-\frac{2}{q}}m^{\frac{2}{q}} \leqslant n \leqslant m/2 \end{cases}$$

3. Let $2 \leq p_1 \leq q \leq p_0$. Then

$$d_n(B_{p_0}^m \cap \nu B_{p_1}^m, l_q^m) \underset{p_0, p_1, q}{\asymp} \begin{cases} k^{\frac{1}{q} - \frac{1}{p_0}}, & for \quad n \leqslant k^{1 - \frac{2}{q}} m^{\frac{2}{q}}, \\ k^{\frac{1}{p_1} - \frac{1}{p_0}} \left(n^{-\frac{1}{2}} m^{\frac{1}{q}} \right)^{\frac{1/p_1 - 1/q}{1/2 - 1/q}}, & for \quad k^{1 - \frac{2}{q}} m^{\frac{2}{q}} \leqslant n \leqslant m/2 \end{cases}$$

4. Let $1 \leq p_1 < 2 < q \leq p_0$. Then

$$d_n(B_{p_0}^m \cap \nu B_{p_1}^m, l_q^m) \underset{p_0, p_{1,q}}{\asymp} \begin{cases} k^{\frac{1}{q} - \frac{1}{p_0}}, & \text{for} \quad n \leqslant k^{1 - \frac{2}{q}} m^{\frac{2}{q}}, \\ k^{\frac{1}{2} - \frac{1}{p_0}} n^{-\frac{1}{2}} m^{\frac{1}{q}}, & \text{for} \quad k^{1 - \frac{2}{q}} m^{\frac{2}{q}} \leqslant n \leqslant m/2 \end{cases}$$

5. Let $q \leq 2$, $1 \leq p_1 < q < p_0$. Then

$$d_n(B_{p_0}^m \cap \nu B_{p_1}^m, l_q^m) \underset{p_0, p_1, q}{\asymp} k^{\frac{1}{q} - \frac{1}{p_0}}.$$

6. Let $1 \leq q \leq p_1 < p_0 \leq \infty$. Then

$$d_n(B_{p_0}^m \cap \nu B_{p_1}^m, l_q^m) \underset{p_0, p_1, q}{\asymp} k^{1/p_1 - 1/p_0} m^{1/q - 1/p_1}$$

Given $k \in \{1, \ldots, m\}$, we denote $\hat{x}_j = 1$ for $1 \leq j \leq k$, $\hat{x}_j = 0$ for $k+1 \leq j \leq m$,

$$V_k = \operatorname{conv} \{ (\varepsilon_1 \hat{x}_{\sigma(1)}, \ldots, \varepsilon_m \hat{x}_{\sigma(m)}) : \varepsilon_j = \pm 1, \ 1 \leq j \leq m, \ \sigma \in S_m \};$$

here S_m is the permutation group on *m* elements. Notice that $V_1 = B_1^m$, $V_m = B_{\infty}^m$. We need the following E.D. Gluskin's results.

Theorem C. [5] There is a non-increasing function $a(\cdot) : [2, \infty) \to (0, 1]$ with the following property: if $q \ge 2$, $1 \le k \le m$, $n \le a(q)m^{\frac{2}{q}}k^{1-\frac{2}{q}}$, then

$$d_n(V_k, l_q^m) \underset{q}{\gtrsim} k^{1/q}.$$
 (1)

Also notice that arguing as in [5] for q = 2 we obtain the inequality

$$d_n(V_k, l_q^m) \ge \min_{x \in \mathbb{R}} \left(k - 2\sqrt{\frac{kn}{m}} x + x^2 \right)^{1/2} = k^{1/2} \left(1 - \frac{n}{m} \right)^{1/2}.$$
 (2)

Theorem D. [11] Let $1 < q \leq 2$, $n \leq m/2$. Then $d_n(V_k, l_q^m) \gtrsim k^{1/q}$.

Proof of Theorem 1. For $p_1 \leq q \leq p_0$ we define $\lambda \in (0, 1)$ by the equation $\frac{1}{q} = \frac{1-\lambda}{p_1} + \frac{\lambda}{p_0}$. Then

$$B_{p_0}^m \cap \nu B_{p_1}^m \subset \nu^{1-\lambda} B_q^m = k^{\frac{1}{q} - \frac{1}{p_0}} B_q^m.$$
(3)

It follows from Hölder's inequality; it is also the particular case of Theorem 2 from [8].

If $p_1 \leq 2 \leq p_0$, we define $\tilde{\lambda} \in (0, 1)$ by the equation $\frac{1}{2} = \frac{1-\tilde{\lambda}}{p_1} + \frac{\tilde{\lambda}}{p_0}$; we similarly obtain

$$B_{p_0}^m \cap \nu B_{p_1}^m \subset \nu^{1-\tilde{\lambda}} B_2^m = k^{\frac{1}{2} - \frac{1}{p_0}} B_2^m.$$
(4)

Let us prove the upper estimates for the Kolmogorov widths. In cases 1 and 2 for $n \leq k^{1-\frac{2}{q}}m^{\frac{2}{q}}$ we use the inclusion $B_{p_0}^m \cap \nu B_{p_1}^m \subset B_{p_0}^m$. In cases 3 and 4 for $n \leq k^{1-\frac{2}{q}}m^{\frac{2}{q}}$ we apply (3); the same inclusion we use in case 5 for $n \leq m/2$.

In cases 1 and 4 for $n \ge k^{1-\frac{2}{q}}m^{\frac{2}{q}}$ we apply (4). In cases 2 and 3 for $n \ge k^{1-\frac{2}{q}}m^{\frac{2}{q}}$ we use the inclusion $B_{p_0}^m \cap \nu B_{p_1}^m \subset \nu B_{p_1}^m$, as well as in case 6 for $n \leq m/2$. It remains to apply Theorems A and B.

Now we obtain the lower estimate. Notice that

$$k^{-\frac{1}{p_0}}V_k \subset B^m_{p_0} \cap \nu B^m_{p_1},\tag{5}$$

$$k^{1/p_1 - 1/p_0} m^{-1/p_1} B^m_{\infty} \subset B^m_{p_0} \cap \nu B^m_{p_1}.$$
 (6)

First we consider q > 2. Let $a(\cdot) \in (0, 1]$ be as in Theorem C. If $a(q)k^{1-\frac{2}{q}}m^{\frac{2}{q}} \leq n \leq a(q)m$, there is a number $\tilde{q} \in [2, q]$ such that

$$n = a(q)k^{1-\frac{2}{\tilde{q}}}m^{\frac{2}{\tilde{q}}} \leqslant a(\tilde{q})k^{1-\frac{2}{\tilde{q}}}m^{\frac{2}{\tilde{q}}}.$$

Then

$$d_n(B_{p_0}^m \cap \nu B_{p_1}^m, l_q^m) \stackrel{(5)}{\geqslant} k^{-\frac{1}{p_0}} d_n(V_k, l_q^m) \stackrel{(1)}{\underset{q}{\gtrsim}} k^{\frac{1}{q} - \frac{1}{p_0}} \cdot m^{\frac{1}{q} - \frac{1}{\tilde{q}}} \underset{q}{\asymp} k^{\frac{1}{2} - \frac{1}{p_0}} n^{-\frac{1}{2}} m^{\frac{1}{q}}.$$

If $a(q)m \leq n \leq m/2$, we have

$$d_n(B_{p_0}^m \cap \nu B_{p_1}^m, l_q^m) \stackrel{(5)}{\geqslant} k^{-\frac{1}{p_0}} d_n(V_k, l_q^m) \geqslant k^{-\frac{1}{p_0}} m^{\frac{1}{q} - \frac{1}{2}} d_n(V_k, l_2^m) \stackrel{(2)}{\underset{p_0, p_1, q}{\gtrsim}} k^{\frac{1}{2} - \frac{1}{p_0}} m^{\frac{1}{q} - \frac{1}{2}}.$$

Hence, we obtain the lower estimate for $a(q)k^{1-\frac{2}{q}}m^{\frac{2}{q}} \leqslant n \leqslant m/2$ in cases 1 and 4. Now we set $\tilde{k} = \left[(a(q)^{1/2}n^{-1/2}m^{1/q})^{\frac{1}{1/q-1/2}} \right]$. If $a(q)k^{1-\frac{2}{q}}m^{\frac{2}{q}} \leqslant n \leqslant a(q)m$,

we have $k \leq \tilde{k} \leq m$; this yields the inequality $\nu \tilde{k}^{1/p_0-1/p_1} \leq 1$ and the inclusion $\nu \tilde{k}^{-1/p_1} V_{\tilde{k}} \subset B_{p_0}^m \cap \nu B_{p_1}^m$. Since $n \leq a(q) \tilde{k}^{1-\frac{2}{q}} m^{\frac{2}{q}}$,

$$d_n(B_{p_0}^m \cap \nu B_{p_1}^m, l_q^m) \ge d_n(\nu \tilde{k}^{-1/p_1} V_{\tilde{k}}, l_q^m) \overset{(1)}{\underset{q}{\gtrsim}} \nu \tilde{k}^{1/q-1/p_1} \underset{p_0, p_1, q}{\asymp} k^{\frac{1}{p_1} - \frac{1}{p_0}} (n^{-1/2} m^{1/q})^{\frac{1/p_1 - 1/q}{1/2 - 1/q}}.$$

If $a(q)m \leq n \leq m/2$, we use (6) and get

$$d_n(B_{p_0}^m \cap \nu B_{p_1}^m, \, l_q^m) \ge k^{1/p_1 - 1/p_0} m^{-1/p_1} d_n(B_{\infty}^m, \, l_q^m) \gtrsim_{p_0, p_1, q} k^{1/p_1 - 1/p_0} m^{1/q - 1/p_1}$$

by Theorem B. Hence, we obtain the lower estimate for $a(q)k^{1-\frac{2}{q}}m^{\frac{2}{q}} \leq n \leq m/2$ in cases 2 and 3.

Now we consider the case $n \leq a(q)k^{1-\frac{2}{q}}m^{\frac{2}{q}}$. Let $\tilde{k} = \left[(a(q)^{1/2}n^{-1/2}m^{1/q})^{\frac{1}{1/q-1/2}} \right]$. Then $\tilde{k} \leq k$; i.e., $\tilde{k}^{1/p_1-1/p_0} \leq \nu$. This yields the inclusion $\tilde{k}^{-1/p_0}V_{\tilde{k}} \subset B_{p_0}^m \cap \nu B_{p_1}^m$. In addition, $n \leq a(q)\tilde{k}^{1-\frac{2}{q}}m^{\frac{2}{q}}$. Therefore,

$$d_n(B_{p_0}^m \cap \nu B_{p_1}^m, l_q^m) \ge d_n(\tilde{k}^{-1/p_0}V_{\tilde{k}}, l_q^m) \underset{q}{\overset{(1)}{\underset{p_0}{\underset{p_0}{\atop{p_0}{\atop{p_1}}}}} \tilde{k}^{1/q-1/p_0} \underset{p_0,p_1,q}{\asymp} \min(1, n^{-1/2}m^{1/q})^{\frac{1/p_0-1/q}{1/2-1/q}}.$$

Hence, we obtain the lower estimate in cases 1, 2.

Further,

$$d_n(B_{p_0}^m \cap \nu B_{p_1}^m, \, l_q^m) \stackrel{(5)}{\geqslant} d_n(k^{-1/p_0}V_k, \, l_q^m) \stackrel{(1)}{\underset{q}{\gtrsim}} k^{1/q-1/p_0}$$

This yields the lower estimate in cases 3, 4.

For $q \leq 2$, $p_0 > q > p_1 \ge 1$ the inequality $d_n(B_{p_0}^m \cap \nu B_{p_1}^m, l_q^m) \gtrsim k^{1/q-1/p_0}$ follows

from (5) and Theorem D.

For $1 \leq q \leq p_1 < p_0 \leq \infty$ we apply the inclusion (6) together with Theorem В.

REFERENCES

- [1] B.S. Kashin, "The diameters of octahedra", Usp. Mat. Nauk 30:4 (1975), 251-252 (in Russian).
- [2] B.S. Kashin, "Diameters of some finite-dimensional sets and classes of smooth functions", Math. USSR-Izv., 11:2 (1977), 317–333.
- [3] A. Pietsch, "s-numbers of operators in Banach space", Studia Math., 51 (1974), 201 - 223.
- [4] M.I. Stesin, "Aleksandrov diameters of finite-dimensional sets and of classes of smooth functions", Dokl. Akad. Nauk SSSR, 220:6 (1975), 1278–1281 [Soviet Math. Dokl.].
- [5] E.D. Gluskin, "On some finite-dimensional problems of the theory of diameters", Vestn. Leningr. Univ., 13:3 (1981), 5–10 (in Russian).
- [6] E.D. Gluskin, "Norms of random matrices and diameters of finite-dimensional sets", Math. USSR-Sb., 48:1 (1984), 173–182.
- [7] A.Yu. Garnaev and E.D. Gluskin, "On widths of the Euclidean ball", Dokl. Akad. Nauk SSSR 277:5 (1984), 1048–1052 [Sov. Math. Dokl. 30 (1984), 200– 204|.
- [8] E.M. Galeev, "The Kolmogorov diameter of the intersection of classes of periodic functions and of finite-dimensional sets", Math. Notes, 29:5 (1981), 382-388.

- [9] E.M. Galeev, "Widths of functional classes and finite-dimensional sets", *Vladikavkaz. Mat. Zh.*, **13**:2 (2011), 3–14.
- [10] A.A. Vasil'eva, "Kolmogorov widths of weighted Sobolev classes on a multidimensional domain with conditions on the derivatives of order r and zero", arXiv:2004.06013v2.
- [11] E. D. Gluskin, "Intersections of a cube with an octahedron are poorly approximated by subspaces of small dimension", *Approximation of functions by special classes of operators*, Interuniv. Collect. Sci. Works, Vologda, 1987, pp. 35–41 (in Russian).