

# ON STABLE RANK OF $H^\infty$ ON COVERINGS OF FINITE BORDERED RIEMANN SURFACES

ALEXANDER BRUDNYI

ABSTRACT. We prove that the Bass stable rank of the algebra of bounded holomorphic functions on an unbranched covering of a finite bordered Riemann surface is equal to one.

## 1. FORMULATION OF MAIN RESULTS

Let  $S'$  be a (not necessarily connected) unbranched covering of a finite bordered Riemann surface  $S$ . In this paper we continue the study initiated in [Br2] of the algebra  $H^\infty(S')$  of bounded holomorphic functions on  $S'$ . (We write  $H^\infty := H^\infty(\mathbb{D})$ , where  $\mathbb{D} \subset \mathbb{C}$  is the open unit disk.) It was shown in our previous work that algebras  $H^\infty(S')$  and  $H^\infty$  share many common properties (e.g., they are Hermite, their maximal ideal spaces are two-dimensional with vanishing second Čech cohomology groups, etc., see [Br2]–[Br4] for the corresponding results). The purpose of this paper is to prove that these algebras have also the same Bass stable rank. The latter notion is defined as follows.

Let  $A$  be an associative ring with unit. For a natural number  $n$  let  $U_n(A)$  denote the set of *unimodular* elements of  $A^n$ , i.e.,

$$U_n(A) = \{(a_1, \dots, a_n) \in A^n : Aa_1 + \dots + Aa_n = A\}.$$

An element  $(a_1, \dots, a_n) \in U_n(A)$  is called *reducible* if there exist  $c_1, \dots, c_{n-1} \in A$  such that  $(a_1 + c_1a_n, \dots, a_{n-1} + c_{n-1}a_n) \in U_{n-1}(A)$ . The *stable rank*  $\text{sr}(A)$  is the least  $n$  such that every element of  $U_{n+1}(A)$  is reducible. The concept of the stable rank introduced by Bass [B] plays an important role in some stabilization problems of algebraic  $K$ -theory. Following Vaserstein [V2] we call a ring of stable rank 1 a *B-ring*. (We refer to this paper for some examples and properties of *B-rings*.)

In [T] Treil proved the following result.

**Theorem A.** *Let  $f, g \in H^\infty$ ,  $\|f\|_{H^\infty} \leq 1$ ,  $\|g\|_{H^\infty} \leq 1$  and*

$$(1.1) \quad \inf_{z \in \mathbb{D}} (|f(z)| + |g(z)|) =: \delta > 0.$$

*Then there exists a function  $G \in H^\infty$  such that the function  $\Phi = f + gG$  is invertible in  $H^\infty$ , and moreover  $\|G\|_{H^\infty} \leq C$ ,  $\|\Phi^{-1}\|_{H^\infty} \leq C$ , where the constant  $C$  depends only on  $\delta$ .*

---

2010 *Mathematics Subject Classification.* Primary 30H50. Secondary 46J10.

*Key words and phrases.* Bass stable rank, *B-ring*, bounded holomorphic function, unbranched covering, elementary matrix.

Research is supported in part by NSERC.

(Here and below for a normed space  $B$  its norm is denoted by  $\|\cdot\|_B$ .)

By the Carleson corona theorem condition (1.1) is satisfied if and only if  $(f, g) \in U_2(H^\infty)$ . Hence, Treil's theorem implies that  $H^\infty$  is a  $B$ -ring.

Theorem A was used by Tolokonnikov [To] to prove that algebras  $H^\infty(U)$  are  $B$ -rings for finitely connected domains and some Behrens domains  $U$ . Until now no other classes of Riemann surfaces  $U$  for which  $H^\infty(U)$  are  $B$ -rings were known. In the present paper, we prove the following extension of Theorem A.

**Theorem 1.1.** *Let  $S'$  be an unbranched covering of a finite bordered Riemann surface  $S$ . Let  $f, g \in H^\infty(S')$ ,  $\|f\|_{H^\infty(S')} \leq 1$ ,  $\|g\|_{H^\infty(S')} \leq 1$  and*

$$(1.2) \quad \inf_{z \in S'} (|f(z)| + |g(z)|) =: \delta > 0.$$

*Then there exists a function  $G \in H^\infty(S')$  such that the function  $\Phi = f + gG$  is invertible in  $H^\infty(S')$ , and moreover  $\max\{\|G\|_{H^\infty(S')}, \|\Phi^{-1}\|_{H^\infty(S')}\} \leq C$ , where the constant  $C$  depends only on  $\delta$  and  $S$ .*

By the corona theorem for  $H^\infty(S')$  (see [Br2, Cor. 1.6]) condition (1.2) is satisfied if and only if  $(f, g) \in U_2(H^\infty(S'))$ . Hence, Theorem 1.1 implies

**Theorem 1.2.**  *$H^\infty(S')$  is a  $B$ -ring.*

**Remark 1.3.** It is known that every  $B$ -ring is *Hermite* (see, e.g., [V2, Thm. 2.7]), i.e., any finitely generated stably free right module over the ring is free (equivalently, any rectangular left-invertible matrix over the ring can be extended to an invertible matrix). Let  $J \subset H^\infty(S')$  be a closed ideal and  $H_J^\infty := \{c + f : c \in \mathbb{C}, f \in J\}$  be the unital closed subalgebra generated by  $J$ . Then Corollary 1.2 implies that  $H_J^\infty$  is a  $B$ -ring (see, e.g., [V1, Thm. 4]); hence, it is Hermite. This gives a generalization of [Br3, Thm. 1.1] proved by a different method.

Let  $M_n(H^\infty(S'))$  be the algebra of  $n \times n$  matrices with entries in  $H^\infty(S')$  regarded as the subspace of bounded linear operators on  $(H^\infty(S'))^n$  equipped with the operator norm. We use Theorem 1.1 to describe the structure of the group  $SL_n(H^\infty(S')) \subset M_n(H^\infty(S'))$  of matrices with determinant 1.

Recall that a matrix in  $SL_n(H^\infty(S'))$  is *elementary* if it differs from the identity matrix by at most one non-diagonal entry.

**Theorem 1.4.** *Every matrix in  $SL_n(H^\infty(S'))$  of norm  $\leq M$  is a product of at most  $(n-1)(\frac{3n}{2} + 1)$  elementary matrices whose norms are bounded from above by a constant depending only on  $M, n$  and  $S$ .*

This result is new even for matrices with entries in  $H^\infty$ .

The proof of Theorem 1.1 is based on Theorem A and some results of the author presented in [Br4] and [Br5] along with some topological results. In the next section we collect some results required for the proof of Theorem 1.1. The proof is given in Section 4.

## 2. AUXILIARY RESULTS

**2.1.** Let  $\mathfrak{M}(A)$  denote the maximal ideal space of a commutative complex unital Banach algebra  $A$ , i.e., the set of nonzero homomorphisms  $A \rightarrow \mathbb{C}$  equipped with the *Gelfand topology*. In this part we present some facts about the maximal ideal space  $\mathfrak{M}(H^\infty(S'))$ , where  $r : S' \rightarrow S$  is a (not necessarily connected but second-countable) unbranched covering of a bordered Riemann  $S$ , see [Br3, Sect. 2], [Br4, Sect. 4] for details.

Recall that  $H^\infty(S')$  separates points of  $S'$  and the map  $\iota : S' \rightarrow \mathfrak{M}(H^\infty(S'))$  sending  $x \in S'$  to the evaluation functional  $\delta_x \in (H^\infty(S'))^*$  at  $x$  embeds  $S'$  into  $\mathfrak{M}(H^\infty(S'))$  as an open dense subset (– the corona theorem for  $H^\infty(S')$ ).

The covering  $r : S' \rightarrow S$  can be viewed as a fiber bundle over  $S$  with a discrete (at most countable) fiber  $F$ . Let  $E(S, \beta F)$  be the space obtained from  $S'$  by taking the Stone-Ćech compactifications of fibres under  $r$ . It is a normal Hausdorff space and  $r$  extends to a continuous map  $r_E : E(S, \beta F) \rightarrow S$  such that  $(E(S, \beta F), S, r_E, \beta F)$  is a fibre bundle on  $S$  with fibre  $\beta F$  and  $S'$  is an open dense subbundle of  $E(S, \beta F)$ . Each  $f \in H^\infty(S')$  admits an extension  $\hat{f} \in C(E(S, \beta F))$  and the algebra formed by such extensions separates points of  $E(S, \beta F)$ . Thus  $\iota$  extends to a continuous injection  $\hat{\iota} : E(S, \beta F) \rightarrow \mathfrak{M}(H^\infty(S'))$ ,  $(\hat{\iota}(\xi))(f) := \hat{f}(\xi)$ .

In what follows, we identify  $E(S, \beta F)$  with its image under  $\hat{\iota}$ . Also, for  $K \subset S$  we set  $K' := r^{-1}(K)$ ,  $K_E := r_E^{-1}(K)$  and for a subset  $U$  of a topological space we denote by  $\overset{\circ}{U}$ ,  $\bar{U}$  and  $\partial U$  its interior, closure and boundary.

It is well known that  $S$  can be regarded as a domain in a compact Riemann surface  $R$  such that  $R \setminus \bar{S}$  is the finite disjoint union of open disks with analytic boundaries. Let  $A(S) \subset H^\infty(S)$  be the subalgebra of functions continuous up to the boundary. We denote by  $\hat{r} : \mathfrak{M}(H^\infty(S')) \rightarrow \bar{S}$  the continuous surjective map induced by the transpose of the homomorphism  $A(S) \rightarrow H^\infty(S')$ ,  $f \mapsto f \circ r$ . Then  $E(S, \beta F)$  coincides with the open set  $\hat{r}^{-1}(S)$  and  $\hat{r}|_{E(S, \beta F)} = r_E$ .

Let  $U \subset R$  be open such that  $V := U \cap \bar{S} \neq \emptyset$ . Then  $\hat{r}^{-1}(V)$  is an open subset of  $\mathfrak{M}(H^\infty(S'))$  and due to the corona theorem  $\overset{\circ}{V}' := r^{-1}(\overset{\circ}{V})$ ,  $\overset{\circ}{V} := U \cap S$ , is an open dense subset of  $\hat{r}^{-1}(V)$ .

**Proposition 2.1.** *Each  $f \in H^\infty(\overset{\circ}{V}')$  admits an extension  $\hat{f} \in C(\hat{r}^{-1}(V))$ .*

*Proof.* We reduce the statement to some known results proved earlier by the author.

We have to extend  $f$  continuously to each point  $\xi \in \hat{r}^{-1}(V)$ . The set  $\hat{r}^{-1}(V)$  is the disjoint union of the open set  $\overset{\circ}{V}_E = \hat{r}^{-1}(\overset{\circ}{V})$  and the set  $\hat{r}^{-1}(V \cap \partial S)$ . So we consider two cases.

(1)  $\xi \in \hat{r}^{-1}(\overset{\circ}{V})$ .

Let  $O \subset \overset{\circ}{V}$  be an open simply connected neighbourhood of  $\hat{r}(\xi)$ . By the definition of the bundle  $E(S, \beta F)$ , the set  $O_E = r_E^{-1}(O)$  is homeomorphic to  $O \times \beta F$  and this homeomorphism maps  $O' = r^{-1}(O)$  biholomorphically onto  $O \times F$ . Then Lemma 3.1 of

[Br1] implies that  $f|_{O'} \in H^\infty(O')$  admits an extension  $\hat{f} \in C(O_E)$  as required (because  $O_E$  is an open neighbourhood of  $\xi$ ).

(2)  $\xi \in \hat{r}^{-1}(V \cap \partial S)$ .

Let  $\hat{r}(\xi)$  belong to a connected component  $\gamma$  of  $\partial S$ . By the definition of  $S$ , there are a relatively open neighbourhood  $A_\gamma \subset \bar{S}$  of  $\gamma$  and a homeomorphic map  $A_\gamma \rightarrow A := \{z \in \mathbb{C} : c < |z| \leq 1\}$ ,  $c > 0$ , which maps  $\gamma$  onto the unit circle  $\mathbb{S} \subset \mathbb{C}$  and is holomorphic on  $\mathring{A}_\gamma$ . Without loss of generality we identify  $A_\gamma$  with  $A$  and  $\gamma$  with  $\mathbb{S}$ . Then since  $V \cap A \neq \emptyset$ , there is a relatively open subset  $\Pi \subset V \cap A$  which is a rectangle in polar coordinates with one side of the boundary on  $\mathbb{S}$  such that  $\hat{r}(\xi) \in \Pi$ . Repeating literally the arguments of the proof of [Br4, Prop. 4.2], we obtain that each function from  $H^\infty(\mathring{\Pi}')$  admits a continuous extension to  $\hat{r}^{-1}(\Pi)$ . Since the latter is an open neighbourhood of  $\xi$ , this gives the required extension of  $f$  to  $\xi$ . We leave the details to the readers.  $\square$

**Remark 2.2.** Since  $\mathring{V}'$  is dense in  $\hat{r}^{-1}(V)$ , the above extension preserves supremum norm. Then the transpose of the restriction homomorphism  $H^\infty(S') \rightarrow H^\infty(\mathring{V}')$ ,  $f \mapsto f|_{\mathring{V}'}$ , induces a continuous map  $s_V : \mathfrak{M}(H^\infty(\mathring{V}')) \rightarrow \mathfrak{M}(H^\infty(S'))$  with image  $\hat{r}^{-1}(\bar{V})$  one-to-one on  $s_V^{-1}(\hat{r}^{-1}(V))$ .

**2.2.** A compact subset  $K \subset \mathfrak{M}(H^\infty(S'))$  is said to be *holomorphically convex* (with respect to the algebra  $H^\infty(S')$ ) if for every  $\xi \notin K$  there is  $f \in H^\infty(S')$  such that

$$\max_K |\hat{f}| < |\hat{f}(\xi)|;$$

here  $\hat{f} \in C(\mathfrak{M}(H^\infty(S')))$  is the Gelfand transform of  $f$ .

A holomorphically convex subset  $Z \subset \mathfrak{M}(H^\infty(S'))$  is called a *hull* if there is a proper ideal  $I \subset H^\infty(S')$  such that

$$Z = \{\xi \in \mathfrak{M}(H^\infty(S')) : \hat{f}(\xi) = 0 \quad \forall f \in I\}.$$

The algebra  $H^\infty(S')$  is a  $B$ -ring if and only if for every hull  $Z \subset \mathfrak{M}(H^\infty(S'))$  the map  $C(\mathfrak{M}(H^\infty(S')), \mathbb{C}^*) \rightarrow C(Z, \mathbb{C}^*)$ ,  $\mathbb{C}^* := \mathbb{C} \setminus \{0\}$ , induced by restriction to  $Z$  is onto, see [CS].

In the next two lemmas,  $S = \mathbb{D}$  and  $S' = S \times \mathbb{N}$  (- the countable disjoint union of open unit disks).

**Lemma 2.3.** *If  $K \subset \mathfrak{M}(H^\infty(S'))$  is holomorphically convex, then for every  $g \in C(K, \mathbb{C}^*)$ , there exists  $\tilde{g} \in C(\mathfrak{M}(H^\infty(S')), \mathbb{C}^*)$  such that  $\tilde{g}|_K = g$ .*

*Proof.* According to [Br5, Lm. 5.3] the homomorphism of the Čech cohomology groups  $H^1(\mathfrak{M}(H^\infty(S')), \mathbb{Z}) \rightarrow H^1(K, \mathbb{Z})$  induced by the restriction map to  $K$  is surjective. In turn, by the Arens-Royden theorem  $H^1(K, \mathbb{Z})$  and  $H^1(\mathfrak{M}(H^\infty(S')), \mathbb{Z})$  are connected components of topological groups  $C(K, \mathbb{C}^*)$  and  $C(\mathfrak{M}(H^\infty(S')), \mathbb{C}^*)$ , respectively. Hence, for each  $g \in C(K, \mathbb{C}^*)$ , there is  $g_1 \in C(\mathfrak{M}(H^\infty(S')), \mathbb{C}^*)$  such that  $g \cdot (g_1^{-1})|_K = e^h$  for some  $h \in C(K)$ . Let  $\tilde{h} \in C(\mathfrak{M}(H^\infty(S')))$  be an extension of  $h$  (existing by the Titze-Urysohn theorem). Then  $\tilde{g} = g_1 e^{\tilde{h}}$  is the required extension of  $g$ .  $\square$

**Lemma 2.4.** *Suppose  $K \subset \mathfrak{M}(H^\infty(S'))$  is holomorphically convex and  $Z \subset \mathfrak{M}(H^\infty(S'))$  is a hull. Then  $K \cup Z \subset \mathfrak{M}(H^\infty(S'))$  is holomorphically convex.*

*Proof.* Let  $\xi \notin K \cup Z$ . By the hypothesis, there exist  $f, g \in H^\infty(S')$  such that  $\hat{f}(\xi) = \hat{g}(\xi) = 1$  and

$$\max_K |f| =: c < 1, \quad g|_Z = 0.$$

Let  $M := \max_K |g|$ . We choose  $n \in \mathbb{N}$  such that  $c^n M < 1$ . Then for  $h := f^n g \in H^\infty(S')$  we have

$$\max_K |\hat{h}| \leq c^n M < 1 = |\hat{h}(\xi)|$$

This shows that the set  $K \cup Z$  is holomorphically convex. □

**2.3.** For the proof of Theorem 1.1 we require the following topological result.

**Lemma 2.5.** *There is a finite cover  $(U_j)_{j=1}^m$  of  $\bar{S}$  by compact subsets homeomorphic to  $\bar{\mathbb{D}}$  such that each  $U_i$  is contained in an open simply connected set  $V_i \subset R$  with simply connected intersection  $V_i \cap S$ , each  $U_i$  intersects with at most two other sets of the family and each non-void  $U_i \cap U_j$  is homeomorphic to  $I := [0, 1]$ .*

*Proof.* Since  $\bar{S}$  is triangulable, we may regard it as a two dimensional polyhedral manifold. It follows from the Whitehead theorem [W, Thm. (3.5)] that there are a (finite) one-dimensional polyhedron  $L \subset \bar{S}$  with sets of edges  $E_L$  and vertices  $V_L$  and a piecewise linear strong deformation retraction  $F : \bar{S} \times I \rightarrow S$  of  $\bar{S}$  onto  $L$  such that

(a)  $F^{-1}(x, 1) \subset \bar{S}$  is a connected polyhedron homeomorphic to a *star tree* with internal vertex  $x$  of degree 2 if either  $x \in \mathring{e}$  for some  $e \in E_L$  or  $x \in V_L$  is of degree  $\leq 2$  and of degree  $> 2$  if  $x \in V_L$  is of degree  $> 2$ , and this homeomorphism maps  $F^{-1}(x, 1) \cap \partial S$  onto the set of external points of the tree.

(b) If  $e \in E_L$ , then  $F^{-1}(\mathring{e}, 1) \cap \partial S$  is the disjoint union of two sets homeomorphic to  $I$ . Let  $E_L := \{e_1, \dots, e_m\}$ . We define

$$U_i := \overline{F^{-1}(\mathring{e}_i, 1)}, \quad 1 \leq i \leq m.$$

Then every  $U_i$  is a polyhedral submanifold of  $\bar{S}$  homeomorphic to  $\bar{\mathbb{D}}$  with the boundary formed by some arcs in  $\partial S$  along with some subsets of  $F^{-1}(v_{i_j}, 1)$ ,  $j = 1, 2$ , homeomorphic to  $I$ ; here  $v_{i_1}, v_{i_2} \in V_L$  are endpoints of  $e_i$ . Clearly every non-void intersection  $U_i \cap U_j \subset F^{-1}(e_i \cap e_j, 1)$  is homeomorphic to  $I$ . Moreover, it is readily seen that each  $U_i$  is contained in an open simply connected subset  $V_i \subset R$  with simply connected intersection  $V_i \cap S$  because  $\bar{S}$  is the strong deformation retract of some of its open neighbourhoods in  $R$  (see, e.g., [W, Thm. (3.3)]). □

### 3. PROOF OF THEOREM 1.2

We retain notation of Lemma 2.5. We set

$$\partial U_i^\circ = \overline{\partial U_i} \setminus \partial S \quad \text{and} \quad W_i := V_i \cap S, \quad 1 \leq i \leq m.$$

Then  $\partial U_i^\circ$  consists of two connected components homeomorphic to  $I$  and  $W_i$  is an open simply connected subset of  $S$ . By the definition,  $\partial U_i^\circ \subset \overline{W_i}$ .

Let  $A(W_i) \subset H^\infty(W_i)$  be the subalgebra of functions continuous up to the boundary. We denote by  $\hat{r}_i : \mathfrak{M}(H^\infty(W'_i)) \rightarrow \overline{W}_i$  the continuous surjective map induced by the transpose of the homomorphism  $A(W_i) \rightarrow H^\infty(W'_i)$ ,  $f \mapsto f \circ r_i$ .

Let  $K$  be either  $\partial U_i^\circ$  or its connected component. We set

$$\tilde{K} := \hat{r}_i^{-1}(K).$$

**Lemma 3.1.** *The set  $\tilde{K} \subset \mathfrak{M}(H^\infty(W'_i))$  is holomorphically convex.*

*Proof.* By our construction the open set  $V_i \setminus K$  is connected. By the Riemann mapping theorem there is a biholomorphic map  $\psi_i$  of  $V_i$  onto  $\mathbb{D}$ . Then  $\mathbb{D} \setminus \psi_i(K)$  is a connected open subset of  $\mathbb{D}$ . This implies that the compact set  $\psi_i(K) \subset \mathbb{C}$  is polynomially convex. Hence,  $K \Subset V_i$  is holomorphically convex with respect to the algebra  $H^\infty(V_i)$  and so it is holomorphically convex in  $\overline{W}_i$  with respect to the algebra  $A(W_i)$ . Since  $\hat{r}_i$  is a surjection onto  $\overline{W}_i$  and  $\tilde{K} \subset \mathfrak{M}(H^\infty(W'_i))$  is the preimage of  $K$ , it is holomorphically convex.  $\square$

Due to Remark 2.2 the transpose of the restriction homomorphism  $H^\infty(S') \rightarrow H^\infty(W'_i)$  induces a continuous map  $s_i : \mathfrak{M}(H^\infty(W'_i)) \rightarrow \mathfrak{M}(H^\infty(S'))$  with image  $\hat{r}^{-1}(\overline{W}_i)$  one-to-one on  $s_i^{-1}(\hat{r}^{-1}(V_i \cap \bar{S}))$ .

Let  $Z \subset \mathfrak{M}(H^\infty(S'))$  be a hull and  $g \in C(Z, \mathbb{C}^*)$ . To prove the theorem we have to extend  $g$  to a function  $\tilde{g} \in C(\mathfrak{M}(H^\infty(S')), \mathbb{C}^*)$ , see Section 2.2 above.

Clearly  $Z_i := s_i^{-1}(Z)$  is a hull for the algebra  $H^\infty(W'_i)$  and  $s_i^*g \in C(Z_i, \mathbb{C}^*)$ . Since  $W'_i := r^{-1}(W_i)$  is biholomorphic to  $\mathbb{D} \times F$ , the Treil theorem implies that there is  $g_i \in C(\mathfrak{M}(H^\infty(W'_i)), \mathbb{C}^*)$  which extends  $s_i^*g$  ( $:= g \circ s_i$ ). Hence  $\tilde{g}_i := g_i \circ s_i^{-1} \in C(\hat{r}^{-1}(V_i \cap \bar{S}), \mathbb{C}^*)$  extends  $g|_{Z \cap \hat{r}^{-1}(U_i)}$  (because  $U_i \subset V_i \cap \bar{S}$ ). If  $Z \cap \hat{r}^{-1}(U_i) = \emptyset$ , we define  $\tilde{g}_i = 1$ .

Next, we order the sets of the cover  $(U_i)_{i=1}^m$  as follows. Choose some  $U_{i_1} \subset \{U_1, \dots, U_m\}$ . If  $U_{i_p}$  is already chosen, we choose  $U_{i_{p+1}}$  so that

$$U_{i_{p+1}} \cap (\cup_{j=1}^p U_{i_j}) \neq \emptyset.$$

This is always possible because  $\bar{S}$  is a connected set. We extend  $g$  by induction on the indices of the order.

For  $j = 1$  we set  $\tilde{g} = \tilde{g}_{i_1}$  on  $\hat{r}^{-1}(U_{i_1})$ . Suppose that  $\tilde{g}$  is already defined on  $\cup_{j=1}^p \hat{r}^{-1}(U_{i_j})$ . Let us define it on  $\cup_{j=1}^{p+1} \hat{r}^{-1}(U_{i_j})$ . To this end let

$$g_{p,p+1} := \tilde{g} \tilde{g}_{p+1}^{-1} \quad \text{on} \quad \hat{r}^{-1}(\cup_{j=1}^p U_{i_j}) \cap \hat{r}^{-1}(U_{i_{p+1}}).$$

By the definition, the above intersection, say  $X$ , is either the preimage of  $\partial U_{i_{p+1}}^\circ$  or its connected component under  $\hat{r}$ . Due to Lemma 3.1, the set  $s_{i_{p+1}}^{-1}(X) \cup Z_{i_{p+1}}$  is holomorphically convex with respect to  $H^\infty(W_{i_{p+1}})$ . Moreover,  $s_{i_{p+1}}^*(g_{p,p+1}) \in C(s_{i_{p+1}}^{-1}(X), \mathbb{C}^*)$  and equals 1 on  $s_{i_{p+1}}^{-1}(X) \cap Z_{i_{p+1}}$ . Hence, it can be extended to a function in  $C(s_{i_{p+1}}^{-1}(X) \cup Z_{i_{p+1}}, \mathbb{C}^*)$  attaining value 1 on  $Z_{i_{p+1}}$ . Due to Lemma 2.3 the extended function can further be extended to a function from  $C(s_{i_{p+1}}^{-1}(\hat{r}^{-1}(U_{i_{p+1}})), \mathbb{C}^*)$ . Composing this extension with  $s_{i_{p+1}}^{-1}$  we obtain an extension  $\tilde{g}_{p,p+1}$  of  $g_{p,p+1}$  equal to 1 on  $Z \cap \hat{r}^{-1}(U_{i_{p+1}})$ . Let us define

$$\tilde{g}|_{\hat{r}^{-1}(U_{i_{p+1}})} := \tilde{g}_{p+1} \tilde{g}_{p,p+1}.$$

Then  $\tilde{g}|_{\hat{r}^{-1}(U_{i_{p+1}})}$  extends  $g|_{Z \cap \hat{r}^{-1}(U_{i_{p+1}})}$  and

$$\tilde{g}|_{\hat{r}^{-1}(U_{i_{p+1}})} \cdot \tilde{g}^{-1}|_{\hat{r}^{-1}(\cup_{j=1}^p U_{i_j})} = \tilde{g}_{p+1} \tilde{g}_{p,p+1} \tilde{g}^{-1} = 1 \quad \text{on} \quad \hat{r}^{-1}(\cup_{j=1}^p U_{i_j}) \cap \hat{r}^{-1}(U_{i_{p+1}}),$$

i.e.,  $\tilde{g}|_{\hat{r}^{-1}(U_{i_{p+1}})}$  is the required extension of  $\tilde{g}|_{\hat{r}^{-1}(\cup_{j=1}^p U_{i_j})}$  to  $\cup_{j=1}^{p+1} U_{i_j}$ . This completes the proof of the induction step and hence of the theorem.

#### 4. PROOFS OF THEOREMS 1.1 AND 1.4

*Proof of Theorems 1.1.* Without loss of generality we may assume that  $S'$  is a connected unbranched covering of  $S$ . Let  $f, g \in H^\infty(S')$ ,  $\|f\|_{H^\infty(S')} \leq 1$ ,  $\|g\|_{H^\infty(S')} \leq 1$  and

$$(4.1) \quad \inf_{z \in S'} (|f(z)| + |g(z)|) =: \delta > 0.$$

Due to Theorem 1.2 there exists a function  $G \in H^\infty(S')$  such that the function  $f + gG$  is invertible in  $H^\infty(S')$ . By  $\mathcal{G}_{f,g,\delta,S'}$  we denote the class of such functions  $G$ . We have to prove that

$$(4.2) \quad C = C(\delta, S) := \sup_{f,g,S'} \inf_{G \in \mathcal{G}_{f,g,\delta,S'}} \max\{\|G\|_{H^\infty(S')}, \|(f + gG)^{-1}\|_{H^\infty(S')}\}$$

is finite. (Here the first supremum is taken over all functions  $f, g$  satisfying the above hypotheses and all connected unbranched coverings  $S'$  of  $S$ .)

Let  $\{S'_i\}_{i \in \mathbb{N}}$  and  $\{f_i\}_{i \in \mathbb{N}}, \{g_i\}_{i \in \mathbb{N}}, f_i, g_i \in H^\infty(S'_i)$ , be sequences satisfying assumptions of the theorem such that

$$(4.3) \quad C = \lim_{i \rightarrow \infty} \inf_{G \in \mathcal{G}_{f_i, g_i, \delta, S'_i}} \max\{\|G\|_{H^\infty(S'_i)}, \|(f_i + g_i G)^{-1}\|_{H^\infty(S'_i)}\}.$$

The disjoint union  $S' := \sqcup_{i \in \mathbb{N}} S'_i$  is clearly an unbranched covering of  $S$  and functions  $f, g \in H^\infty(S')$  defined by the formulas

$$f|_{S'_i} := f_i, \quad g|_{S'_i} := g_i, \quad i \in \mathbb{N},$$

are of norms  $\leq 1$  and satisfy condition (4.1) on  $S'$ . Then due to Theorem 1.2 there exists a function  $G \in H^\infty(S')$  such that the function  $f + gG$  is invertible in  $H^\infty(S')$ . We set

$$G_i := G|_{S'_i}, \quad i \in \mathbb{N}.$$

Then due to (4.3)

$$C \leq \sup_{i \in \mathbb{N}} \max\{\|G_i\|_{H^\infty(S'_i)}, \|(f_i + g_i G_i)^{-1}\|_{H^\infty(S'_i)}\} = \max\{\|G\|_{H^\infty(S')}, \|(f + gG)^{-1}\|_{H^\infty(S')}\}.$$

This completes the proof of the theorem.  $\square$

*Proof of Theorem 1.4.* According to [Br4, Thm. 1.3 (b)] the covering dimension of the maximal ideal space  $\mathfrak{M}(H^\infty(S'))$  is 2. In turn, due to the Browder theorem [Bro, Thm. 6.11] the second homotopy group  $\pi_2(SL_n(\mathbb{C})) = 0$  (here  $SL_n(\mathbb{C})$  is the group of  $n \times n$  complex matrices with determinant 1). These two facts and the Hu theorem [Hu, (11.4)] imply that the homotopy classes of the continuous mappings  $f : \mathfrak{M}(H^\infty(S')) \rightarrow SL_n(\mathbb{C})$  are in a one-to-one correspondence with the elements of the first Čech cohomology group

$H^1(\mathfrak{M}(H^\infty(S')), \pi_1(SL_n(\mathbb{C})))$ . This group is trivial because the space  $SL_n(\mathbb{C})$  is simply connected. Hence, each  $f \in C(\mathfrak{M}(H^\infty(S')), SL_n(\mathbb{C}))$  is homotopic to a constant map with value  $1_n$  (the unit of  $SL_n(\mathbb{C})$ ), i.e., the space  $C(\mathfrak{M}(H^\infty(S')), SL_n(\mathbb{C}))$  is connected. Next, by the Arens theorem [A] the Gelfand transform induces a bijection between the sets of connected components of the spaces  $SL_n(H^\infty(S'))$  and  $C(\mathfrak{M}(H^\infty(S')), SL_n(\mathbb{C}))$ . Therefore the group  $SL_n(H^\infty(S'))$  is connected as well. In particular, each matrix in  $SL_n(H^\infty(S'))$  can be presented as a finite product of elementary matrices. (This is a well-known fact; it can be deduced, e.g., from [BRS, Thm. 2.1].) Then since  $H^\infty(S')$  is a  $B$ -ring (by Theorem 1.2), Lemma 9 and Remark 10 of [DV] imply that each matrix  $F \in SL_n(H^\infty(S'))$  can be presented as a product of at most  $(n-1)(\frac{3n}{2}+1)$  elementary matrices. Let us show that if

$$(4.4) \quad \|F\|_{M_n(H^\infty(S'))} \leq M,$$

these matrices would be chosen so that their norms were bounded from above by a constant depending only on  $M$ ,  $n$  and  $S$ .

As before we may assume that  $S'$  is connected. Let  $\mathcal{F}_{M,S',n}$  be the class of matrices  $F \in SL_n(H^\infty(S'))$  satisfying (4.4). For every  $F \in \mathcal{F}_{M,S',n}$  by  $\Pi_{F,M,S',n}$  we denote the set of all possible products of  $F$  by at most  $(n-1)(\frac{3n}{2}+1)$  elementary matrices. By the above arguments the set  $\Pi_{F,M,S',n}$  is non-void. For each  $\pi \in \Pi_{F,M,S',n}$  by  $\|\pi\|$  we denote maximum of norms of elementary matrices in  $\pi$ . We have to prove that

$$(4.5) \quad C = C(S, M, n) := \sup_{S', F \in \mathcal{F}_{M,S',n}} \inf_{\pi \in \Pi_{F,M,S',n}} \|\pi\| < \infty;$$

here  $S'$  runs over all connected unbranched coverings of  $S$ .

Let  $S'_i$  and  $F_i \in \mathcal{F}_{M,S'_i,n}$ ,  $i \in \mathbb{N}$ , be such that

$$(4.6) \quad C = \lim_{i \rightarrow \infty} \inf_{\pi \in \Pi_{F_i, M, S'_i, n}} \|\pi\|.$$

It is clear that the disjoint union  $S' := \sqcup_{i \in \mathbb{N}} S'_i$  is an unbranched covering of  $S$  and the matrix  $F \in H^\infty(S')$  defined by the formula

$$F|_{S'_i} := F_i, \quad i \in \mathbb{N},$$

belongs to the class  $\mathcal{F}_{M,S',n}$ . Then there is  $\pi \in \Pi_{F,M,S',n}$ . By  $\pi_i$  we denote the product obtained by restriction of elementary matrices in  $\pi$  to  $S'_i$ . Then each  $\pi_i \in \Pi_{F_i, M, S'_i, n}$  and so due to (4.6)

$$C \leq \sup_{i \in \mathbb{N}} \|\pi_i\| = \|\pi\| < \infty.$$

This completes the proof of the theorem.  $\square$

## REFERENCES

- [A] R. Arens, To what extent does the space of maximal ideals determine the algebra?, in "Function Algebras" (Birtel, Ed.), Scott-Foresman, Chicago, 1966.
- [B] H. Bass, K-theory and stable algebra, Publ. Mat. I. H. E. S. **22** (1964), 5–60.
- [Bro] W. Brouder, Torsion in  $H$ -spaces, Ann. Math. **74** (1961), 24–51.
- [BRS] A. Brudnyi, L. Rodman and I. M. Spitkovsky, Projective free algebras of continuous functions on compact abelian groups, J. Funct. Anal. **259** (2010), 918–932.



- [Br1] A. Brudnyi, Topology of maximal ideal space of  $H^\infty$ , J. Funct. Anal. **189** (1) (2002) 21–52.
- [Br2] A. Brudnyi, Projections in the space  $H^\infty$  and the Corona Theorem for subdomains of coverings of finite bordered Riemann surfaces, Ark. Mat. **42** (2004), 31–59.
- [Br3] A. Brudnyi, Grauert- and Lax–Halmos-type theorems and extension of matrices with entries in  $H^\infty$ , J. Funct. Anal. **206** (2004), 87–108.
- [Br4] A. Brudnyi, Projective freeness of algebras of bounded holomorphic functions on infinitely connected domains, arXiv:1905.01532.
- [Br5] A. Brudnyi, Dense stable rank and Runge type approximation theorem for  $H^\infty$  maps, arXiv:2006.04179.
- [CS] G. Corach and F. D. Suarez, Extension problems and stable rank in commutative Banach algebras, Topology Appl. **21** (1985), 1–8.
- [DV] R. K. Dennis and L. N. Vaserstein, On a question of M. Newman on the number of commutators. J. Algebra **118** (1988), 150–161.
- [Hu] S.-T. Hu, Mappings of a normal space into an absolute neighborhood retract, Trans. Amer. Math. Soc. **64** (1948), 336–358.
- [To] V. Tolokonnikov, Stable rank of  $H^\infty$  in multiply connected domains, Proc. Amer. Math. Soc. **123** (10) (1995), 3151–3156.
- [T] S. Treil, Stable rank of  $H^\infty$  is equal to one, J. Funct. Anal. **109** (1992), 130–154.
- [V1] L.N. Vaserstein, Stable range of rings and dimension of topological spaces, Funk. Anal. Pril. **5** (2) (1971) 17–27=Funct. Anal. Appl. **5** (1971) 102–110.
- [V2] L. N. Vaserstein, The stable range for rings and the dimension of topological spaces, Functional Anal. Appl., vol. 5 (1971), 102–110.
- [W] J. H. C. Whitehead, The immersion of an open 3-manifold in Euclidean 3-space, Proc. London Math. Soc. (3) Vol. 11 (1961), 81–90.

DEPARTMENT OF MATHEMATICS AND STATISTICS  
 UNIVERSITY OF CALGARY  
 CALGARY, ALBERTA, CANADA  
 T2N 1N4  
*E-mail address:* [abrudnyi@ucalgary.ca](mailto:abrudnyi@ucalgary.ca)