MORE ON THE RINGS $B_1(X)$ AND $B_1^*(X)$

ATANU MONDAL AND A. DEB RAY

ABSTRACT. This paper focuses mainly on the ring of all bounded Baire one functions on a topological space. The uniform norm topology arises from the sup-norm defined on the collection $B_1^*(X)$ of all bounded Baire one functions. With respect to this topology, $B_1^*(X)$ is a topological ring. It is proved that under uniform norm topology, the set of all units forms an open set and as a consequence of it, every maximal ideal of $B_1^*(X)$ is closed in $B_1^*(X)$ with uniform norm topology. Since the natural extension of uniform norm topology on $B_1(X)$, when $B_1^*(X) \neq B_1(X)$, does not show up these features, a topology called m_B -topology is defined on $B_1(X)$ suitably to achieve these results on $B_1(X)$. It is proved that the relative m_B topology coincides with the uniform norm topology on $B_1^*(X)$ if and only if $B_1(X) = B_1^*(X)$. Moreover, $B_1(X)$ with m_B -topology is 1st countable if and only if $B_1(X) = B_1^*(X)$. The last part of the paper establishes a correspondence between the ideals of $B_1^*(X)$ and a special class of Z_B -filters, called e_B -filters on a normal topological space X. It is also observed that for normal spaces, the cardinality of the collection of all maximal ideals of $B_1(X)$ and those of $B_1^*(X)$ are the same.

1. Introduction and Preliminaries

In [1], we have initiated the study of the ring $B_1(X)$ of Baire one functions on a topological space X and have achieved several interesting results. The bounded Baire one functions, denoted by $B_1^*(X)$ is in general a subring of $B_1(X)$. It has been proved in [1] that for a completely Hausdorff space X, total disconnectedness of X is a necessary condition for $B_1^*(X) = B_1(X)$, however, the converse is not always true. It is therefore a question, when do we expect the converse to hold? In this paper, introducing a suitable topology on $B_1(X)$, called m_B -topology, we establish a couple of necessary and sufficient conditions for $B_1^*(X) = B_1(X)$. We observe that $(B_1(X), m_B)$ is a 1st countable space if and only if $B_1(X) = B_1^*(X)$, which is an analogue of the result proved in [4] for pseudocompact spaces.

Defining zero sets of Baire one functions in the usual way in [2], we have established the duality between the ideals of $B_1(X)$ with a collection of typical subfamilies of the zero sets of Baire one functions, called Z_B -filters on X. The study of such correspondence has resemblance to the duality between ideals of C(X), the ring of all real valued continuous functions on X and the z-filters on X [5]. It is of course a natural query whether such correspondence remains true if we confine the ideals within the class of ideals of $B_1^*(X)$. It is not hard to observe that for any Z_B -filter \mathscr{F} on X, $Z_B^{-1}[\mathscr{F}] \cap B_1^*(X)$ is an ideal in $B_1^*(X)$. However, Z_B fails to take an ideal of $B_1^*(X)$ to a Z_B -filter on X. Introducing a new type of Z_B -filter, called e_B -filter

1

²⁰¹⁰ Mathematics Subject Classification. 13A15, 26A21, 54A05, 54C30, 54C50, 54H11, 54H13. Key words and phrases. Uniform norm topology, u_B -topology, m_B -topology, Z_B -filter, Z_B -ultrafilter, e_B -ideal, e_B -filter, e_B -ultrafilter.

on X, we obtain in this paper a similar correspondence between the ideals of $B_1^*(X)$ and the e_B -filters on any normal topological space X.

We now state some existing definitions and results that are required for this paper. The zero set Z(f) of a function $f \in B_1(X)$ is defined by $Z(f) = \{x \in X : f(x) = 0\}$ and the collection of all zero sets in $B_1(X)$ is denoted by $Z(B_1(X))$.

It is evident that, Z(f) = Z(|f|), for all $f \in B_1(X)$.

For any real number $r, \mathbf{r} \in B_1(X)$ (or $B_1^*(X)$) will always indicate the real valued constant function defined on X whose value is r. So $Z(\mathbf{0}) = X$ and $Z(\mathbf{s}) = \emptyset$, for any $s \neq 0$.

If a Baire one function f on X is a unit in the ring $B_1(X)$ then $\{x \in X : f(x) = 0\} = \emptyset$. The following results provide some sufficient conditions to determine the units in $B_1(X)$, where X is any topological space..

Theorem 1.1. [1] Let X be a topological space and $f \in B_1(X)$ be such that f(x) > 0 (orf(x) < 0), $\forall x \in X$, then $\frac{1}{f}$ exists and belongs to $B_1(X)$.

Now, if $Z(f) = \{x \in X : f(x) = 0\} = \emptyset$, for some $f \in B_1(X)$, then $Z(f^2) = \emptyset$ and so by using Theorem 1.1, as $f^2 > 0$ there exists a $g \in B_1(X)$ such that $f(fg) = f^2g = 1$, i.e., f is a unit in $B_1(X)$.

We must note that this consequence has already been pointed out in [7]. Therefore, we have the following characterization:

Theorem 1.2. In any topological space X, $f \in B_1(X)$ is a unit in $B_1(X)$ if and only if $Z(f) = \emptyset$.

Clearly, if a bounded Baire one function f on X is bounded away from zero, i.e. there exists some $m \in \mathbb{R}$ such that $0 < m \le f(x), \forall x \in X$ then $\frac{1}{f}$ is also a bounded Baire one function on X.

Theorem 1.3. [1] Let f be any Baire one function on X and $g: \mathbb{R} \to \mathbb{R}$ be a continuous function. Then their composition $g \circ f$ is also a Baire one function.

Here we recall from [6] that Baire one functions are described in terms of pull-backs of open sets as follows:

Theorem 1.4. [6] (i) For any topological space X and any metric space Y, $B_1(X,Y) \subseteq \mathscr{F}_{\sigma}(X,Y)$, where $B_1(X,Y)$ denotes the collection of Baire one functions from X to Y and $\mathscr{F}_{\sigma}(X,Y) = \{f : X \to Y : f^{-1}(G) \text{ is an } F_{\sigma} \text{ set, for any open set } G \subseteq Y\}$. (ii) For a normal topological space X, $B_1(X,\mathbb{R}) = \mathscr{F}_{\sigma}(X,\mathbb{R})$.

Theorem 1.5. [1] For any $f \in B_1(X)$, Z(f) is a G_δ set.

Definition 1.6. A nonempty subcollection \mathscr{F} of $Z(B_1(X))$ is said to be a Z_B -filter on X, if it satisfies the following conditions:

- (1) $\emptyset \notin \mathscr{F}$
- (2) if $Z_1, Z_2 \in \mathscr{F}$, then $Z_1 \cap Z_2 \in \mathscr{F}$
- (3) if $Z \in \mathscr{F}$ and $Z' \in Z(B_1(X))$ is such that $Z \subseteq Z'$, then $Z' \in \mathscr{F}$.

Definition 1.7. A Z_B -filter is called Z_B -ultrafilter on X, if it is not properly contained in any other Z_B -filter on X.

Theorem 1.8. [2] If I is any ideal of $B_1(X)$, then $Z_B[I] = \{Z(f) : f \in I\}$ is a Z_B -filter on X.

Theorem 1.9. [2] For any Z_B -filter \mathscr{F} on X, $Z_B^{-1}[\mathscr{F}] = \{f \in B_1(X) : Z(f) \in \mathscr{F}\}$ is an ideal in $B_1(X)$.

Let \mathscr{I}_B be the collection of all ideals in $B_1(X)$ and $\mathscr{F}_B(X)$ be the collection of all Z_B -filters on X. The map $Z_B: \mathscr{I}_B \mapsto \mathscr{F}_B(X)$ defined by $Z_B(I) = Z_B[I], \forall I \in \mathscr{I}_B$ is surjective but not injective in general. Although, the restriction map $Z_B: \mathcal{M}(B_1(X)) \mapsto \Omega_B(X)$ is a bijection, where $\mathcal{M}(B_1(X))$ and $\Omega_B(X)$ are respectively the collection of all maximal ideals in $B_1(X)$ and the collection of all Z_B -ultrafilters on X [2].

A criterion to paste two Baire one functions to obtain a Baire one function on a bigger domain is proved for normal topological spaces in the following theorem:

Theorem 1.10. (Pasting Lemma) Let X be a normal topological space with $X = A \cup B$, where both A and B are G_{δ} sets in X. If $f : A \mapsto \mathbb{R}$ and $g : B \mapsto \mathbb{R}$ are Baire one functions so that f(x) = g(x), for all $x \in A \cap B$, then the map $h : X \mapsto \mathbb{R}$ defined by

$$h(x) = \begin{cases} f(x) & \text{if } x \in A \\ g(x) & \text{if } x \in B. \end{cases}$$

is also a Baire one function.

Proof. Let C be any closed set in \mathbb{R} . Then $h^{-1}(C) = f^{-1}(C) \bigcup g^{-1}(C)$. By Theorem 1.4, $f^{-1}(C)$ and $g^{-1}(C)$ are G_{δ} sets in A and B respectively. Since A and B are G_{δ} sets in X, $f^{-1}(C)$ and $g^{-1}(C)$ are G_{δ} sets in X. We know finite union of G_{δ} sets is G_{δ} , so $h^{-1}(C)$ is a G_{δ} set in X. Therefore, using Theorem 1.4, we have h is a Baire one function.

The result remains true if the G_{δ} sets are replaced by F_{σ} sets. The proof being exactly the same as that of Theorem 1.10, we only state the result as follows:

Theorem 1.11. Let X be a normal topological space with $X = A \cup B$, where both A and B are F_{σ} sets in X. If $f: A \mapsto \mathbb{R}$ and $g: B \mapsto \mathbb{R}$ are Baire one functions so that f(x) = g(x), for all $x \in A \cap B$, then the map $h: X \mapsto \mathbb{R}$ defined by

$$h(x) = \begin{cases} f(x) & \text{if } x \in A \\ g(x) & \text{if } x \in B. \end{cases}$$

is also a Baire one function.

2. m_B-topology - An extension of uniform norm topology

Let X be any topological space. Defining the 'sup'-norm as usual we get that $B_1^*(X)$ is a Banach space. The topology induced by the sup-norm is known as the uniform norm topology and it is not hard to see that $B_1^*(X)$ with uniform norm topology is a topological ring. For any $f \in B_1^*(X)$, the collection $\{B(f,\epsilon) : \epsilon > 0\}$, where $B(f,\epsilon) = \{g \in B_1^*(X) : ||f-g|| \le \epsilon\}$, forms a base for neighbourhood system at f in the uniform norm topology on $B_1^*(X)$.

Theorem 2.1. The set \mathscr{U}_{B^*} of all units of $B_1^*(X)$ is an open set in $B_1^*(X)$ with the uniform norm topology.

Proof. Let $f \in \mathcal{U}_{B^*}$. So, f is bounded away from 0, i.e., $\exists \lambda > 0$ such that $f(x) > \lambda$, for all $x \in X$. We observe that for any $g \in B(f, \frac{\lambda}{2})$, $|f(x) - g(x)| \leq \frac{\lambda}{2}$, for all $x \in X$. This implies that, $\lambda < |f(x)| \leq |f(x) - g(x)| + |g(x)| \leq \frac{\lambda}{2} + |g(x)|$, for all $x \in X$. Therefore, $|g(x)| > \frac{\lambda}{2}$, for all $x \in X$. Hence, $g \in \mathcal{U}_{B^*}$ and $B(f, \frac{\lambda}{2}) \subseteq \mathcal{U}_{B^*}$. This completes the proof that \mathcal{U}_{B^*} is an open set.

completes the proof that \mathscr{U}_{B^*} is an open set. \square **Theorem 2.2.** If I is a proper ideal in $B_1^*(X)$ then \overline{I} (Closure of I in the uniform

norm topology) is also a proper ideal in $B_1^*(X)$.

Proof. Since $B_1^*(X)$ is a topological ring, \overline{I} is an ideal. We show that \overline{I} is a proper ideal of $B_1^*(X)$. Since I does not contain any unit, $\mathscr{U}_{B^*} \cap I = \emptyset \implies I \subseteq B_1^*(X) \setminus U_{B^*}$. Now \mathscr{U}_{B^*} being an open set in uniform norm topology, $B_1^*(X) \setminus \mathscr{U}_{B^*}$ is a closed set containing I which implies $\overline{I} \subseteq B_1^*(X) \setminus \mathscr{U}_{B^*}$. Therefore, $\overline{I} \cap \mathscr{U}_{B^*} = \emptyset$. Consequently, $1 \notin \overline{I}$, proving \overline{I} a proper ideal of $B_1^*(X)$.

Corollary 2.3. Each maximal ideal of $B_1^*(X)$ is a closed set in the uniform norm topology.

The natural extension of uniform norm topology on $B_1(X)$ would be the one for which $\{U(f,\epsilon):\epsilon>0\}$ is a base for neighbourhood system at $f\in B_1(X)$, where $U(f,\epsilon)=\{g\in B_1(X):|f(x)-g(x)|\leq \epsilon$, for all $x\in X\}$. We call this topology the u_B -topology on $B_1(X)$. Clearly on $B_1^*(X)$, the subspace topology obtained from the u_B -topology coincides with the uniform norm topology, so in that case, $B_1(X)$ is a topological ring and the set of all units in $B_1(X)$ forms an open set in this topology. However, it is neither a topological ring nor the set of all units in $B_1(X)$ forms an open set in this topology, unless $B_1(X)=B_1^*(X)$.

Theorem 2.4. If $B_1(X) \neq B_1^*(X)$ then $B_1(X)$ is not a topological ring with respect to u_B -topology.

Proof. Let X be a topological space such that $B_1(X) \neq B_1^*(X)$. So, there exists an $f \in B_1(X) \setminus B_1^*(X)$ with $f(x) \geq 1$, $\forall x \in X$. We shall show that the multiplication operation is not continuous at the point $(\mathbf{0}, f)$, for any $f \in B_1(X)$. We choose a neighbourhood $U(\mathbf{0}, 1)$ of the constant function $\mathbf{0}$. Clearly each function in $U(\mathbf{0}, 1)$ is bounded on X. Let $U(\mathbf{0}, \epsilon)$ and $U(f, \delta)$ be arbitrary basic neighbourhoods of $\mathbf{0}$ and f in $B_1(X)$ with u_B -topology. We notice that, the constant function $\frac{\epsilon}{2} \in U(\mathbf{0}, \epsilon)$ and $f \in U(f, \delta)$ but $\frac{\epsilon}{2} f$ is an unbounded function. So, $\frac{\epsilon}{2} f \notin U(\mathbf{0}, 1)$. This proves that the multiplication is not continuous. Hence, $B_1(X)$ with u_B -topology is not a topological ring.

In the next theorem we show that the collection of all units of $B_1(X)$ is not an open set in u_B -topology, if $B_1(X) \neq B_1^*(X)$.

Theorem 2.5. If $B_1(X) \neq B_1^*(X)$ then \mathscr{U}_B , the collection of all units in $B_1(X)$ is not an open set in u_B -topology.

Proof. Let $f \in B_1(X) \setminus B_1^*(X)$ be an unbounded Baire one function with $f \geq 1$ on X. We define $g = \frac{1}{f}$ on X. Since f(x) > 0 for all $x \in X$, by Theorem 1.1 g is a Baire one function. So g is a positive unit in $B_1(X)$ which takes values arbitrarily close to 0. We show that g is not an interior point of \mathcal{U}_B . Indeed, for each $\epsilon > 0$,

 $U(g,\epsilon) \nsubseteq \mathscr{U}_B$. We can select a point $a \in X$, such that $0 < g(a) < \epsilon$. Taking h = g - g(a) we get that $h \in U(g,\epsilon)$ but h(a) = 0. So, h is not a unit in $B_1(X)$, i.e., $h \notin \mathscr{U}_B$.

We put together the outcome of the above discussion in the following theorem:

Theorem 2.6. For a topological space X, the following statements are equivalent:

- (i) $B_1(X) = B_1^*(X)$.
- (ii) $B_1(X)$ with u_B -topology is a topological ring.
- (iii) The set of all units in $B_1(X)$ forms an open set in u_B -topology.

Corollary 2.7. If a Completely Hausdorff space X is not totally disconnected then $B_1(X)$ with u_B -topology is not a topological ring. Moreover, the collection of all units does not form an open set in $B_1(X)$.

Proof. Follows from Theorem 3.4 of [1].

To overcome the inadequacy of u_B -topology, we define another topology on $B_1(X)$ as follows: Define $M(g,u) = \{f \in B_1(X) : |f(x) - g(x)| \le u(x), \text{ for every } x \in X\}$ and $\widetilde{M}(g,u) = \{f \in B_1(X) : |f(x) - g(x)| < u(x), \text{ for every } x \in X\}$, where u is any positive unit in $B_1(X)$. It is not hard to check that the collection $\mathscr{B} = \{\widetilde{M}(g,u) : g \in B_1(X) \text{ and } u \text{ is any positive unit in } B_1(X)\}$ is an open base for some topology on $B_1(X)$.

Theorem 2.8. \mathscr{B} forms an open base for some topology on $B_1(X)$.

Proof. (i) \mathscr{B} covers $B_1(X)$ follows from the construction of \mathscr{B} .

(ii) Let $\widetilde{M}(g_1, u_1) \cap \widetilde{M}(g_2, u_2) \neq \emptyset$ and $f \in \widetilde{M}(g_1, u_1) \cap \widetilde{M}(g_2, u_2)$. Since $f \in \widetilde{M}(g_1, u_1)$, it follows that, for all $x \in X$, $|f(x) - g(x)| < u_1(x)$.

Set $u_1^*(x) = u_1(x) - |f(x) - g_1(x)|$, for all $x \in X$. Then $u_1^* \in B_1(X)$ with $u_1^*(x) > 0$, for all $x \in X$. So, u_1^* is a positive unit in $B_1(X)$.

We claim that, $M(f, u_1^*) \subseteq M(g_1, u_1)$.

Choose $h \in \widetilde{M}(f, u_1^*)$, which implies that $|h(x) - f(x)| < u_1^*(x), \forall x \in X$.

Now, $|h(x) - g_1(x)| \le |h(x) - f(x)| + |f(x) - g_1(x)| < u_1^*(x) + u_1(x) - u_1^*(x) = u_1(x)$. So, $h \in \widetilde{M}(g_1, u_1)$.

Analogously, $M(f, u_2^*) \subseteq M(g_2, u_2)$, where u_2^* is a positive unit in $B_1(X)$.

Let $u^* = \min\{u_1^*, u_2^*\}$. Then u^* is a positive unit in $B_1(X)$ and $M(f, u^*) \subseteq \widetilde{M}(g_1, u_1) \cap \widetilde{M}(g_2, u_2)$. Therefore, \mathscr{B} is an open base for some topology on $B_1(X)$.

Definition 2.9. The topology on $B_1(X)$ for which \mathscr{B} forms a base for open sets is called the m_B -topology.

Remark 2.10. It is immediate that, for any $g \in B_1(X)$, the collection $\{M(g, u) : u \text{ is a positive unit in } B_1(X)\}$ is a neighbourhood base about g in the m_B -topology.

The following theorems show that $B_1(X)$ with m_B -topology overcomes the draw-backs of the u_B -topology.

Theorem 2.11. $B_1(X)$ with m_B -topology is a topological ring.

Proof. We shall show that both addition and multiplication operations are continuous in m_B -topology. For any $f, g \in B_1(X)$ we consider a neighbourhood M(f+g, u)

at the point f + g, where u is a positive unit in $B_1(X)$. Then it is clear that, $M(f,\frac{u}{2})+M(g,\frac{u}{2})\subseteq M(f+g,u)$. So, addition is a continuous map.

To show multiplication is continuous at (f, g), we select a positive unit u of $B_1(X)$. We want to produce a positive unit v of $B_1(X)$ with $v \leq 1$ such that $M(f, v).M(g, v) \subseteq M(fg, u).$

If v has to satisfy the above relation, then we should have, whenever, $|h_1 - f| \leq v$ and $|h_2 - g| \le v$ then $|fg - h_1h_2| \le u$.

Now if $|h_1 - f| \le v$ and $|h_2 - g| \le v$, then

 $|fg-h_1h_2|$

 $= |h_1(h_2 - g) + g(h_1 - f)|$

 $= |h_1(h_2 - g) + (g - h_2)(h_1 - f) + h_2(h_1 - f)|$ $\leq |h_1|v + v^2 + |h_2|v$

 $\leq v[|h_1| + 1 + |h_2|]$

 $\leq v[v + |f| + 1 + v + |g|]$

 $\leq v[|f|+|g|+3]$

Therefore, the unit v defined as $v = \left(\frac{u}{2(|f| + |g| + 3)}\right) \wedge 1$ will serve our purpose.

Hence, $B_1(X)$ with m_B -topology is a topological ring.

Theorem 2.12. \mathcal{U}_B , the collection of all units in $B_1(X)$ is open in the m_B topology.

Proof. Let $u \in \mathcal{U}_B$. If we show that $M(u, \frac{|u|}{2}) \subseteq \mathcal{U}_B$, then u is an interior point of \mathscr{U}_B . Indeed, for any $v \in M(u, \frac{|u|}{2}) \Longrightarrow |v(x) - u(x)| < \frac{|u(x)|}{2}, \forall x \in X \Longrightarrow v(x) \neq 0$, for all $x \in X$. So by Theorem 1.2, $v \in \mathscr{U}_B$. This completes the proof.

Theorem 2.13. The closure \overline{I} of a proper ideal I of $B_1(X)$ with m_B -topology is also a proper ideal.

Proof. $B_1(X)$ with m_B -topology being a topological ring, \overline{I} is an ideal of $B_1(X)$. It is enough to show that \overline{I} is a proper ideal of $B_1(X)$. Since \mathcal{U}_B is an open set, proceeding as in Theorem 2.2 we obtain the result.

Theorem 2.14. $B_1^*(X)$ is a closed subset of $B_1(X)$ with m_B -topology.

Proof. Let $f \in B_1(X) \setminus B_1^*(X)$. Then $M(f,1) \subseteq B_1(X) \setminus B_1^*(X)$ implies that f is not a limit point of $B_1^*(X)$ and this proves the desired result.

Theorem 2.15. The uniform norm topology on $B_1^*(X)$ is weaker than the relative m_B -topology on $B_1^*(X)$.

Proof. For any $f \in B_1^*(X)$ and $\epsilon > 0$ we get $B(f, \epsilon) = \{g \in B_1^*(X) : |f(x) - g(x)| \le 1 \}$ $\{\epsilon\} = M(f, \epsilon) \cap B_1^*(X)$ and this completes the proof.

Theorem 2.16. Each maximal ideal in $B_1^*(X)$ is closed in $B_1(X)$ with m_B topology.

Proof. Each maximal ideal M in $B_1^*(X)$ is closed in $B_1^*(X)$ with respect to relative m_B -topology, since the uniform norm topology on $B_1^*(X)$ is weaker than the relative m_B -topology on $B_1^*(X)$ and M is closed under uniform norm topology. Also $B_1^*(X)$ is closed in $B_1(X)$ with m_B -topology. Hence M is closed in $B_1(X)$.

Theorem 2.17. For a normal (T_4) topological space X, every closed ideal I in $B_1(X)$ with m_B -topology is a Z_B -ideal.

Proof. Suppose $f, g \in B_1(X)$ with $g \in I$ and Z(f) = Z(g). It is enough to show that $f \in I$. Since I is closed, we show that for any positive unit u in $B_1(X)$, $M(f, u) \cap I \neq \emptyset$. Consider the map $h : X \mapsto \mathbb{R}$ given by

$$h(x) = \begin{cases} 0 & \text{if } |f(x)| \le u(x) \\ \frac{f(x) - u(x)}{g(x)} & \text{if } f(x) \ge u(x) \\ \frac{f(x) + u(x)}{g(x)} & \text{if } f(x) \le -u(x) \end{cases}$$

h is well defined. Observe that the collections $\{x \in X: |f(x)| \le u(x)\}, \{x \in X: f(x) \ge u(x)\}, \{x \in X: f(x) \le -u(x)\}$ are G_{δ} sets in X and $0, \frac{f(x)-u(x)}{g(x)}, \frac{f(x)+u(x)}{g(x)}$ are Baire one functions on their respective domains. So by Theorem 1.10, h is a Baire one function on X.

For all $x \in X$, $|h(x)g(x) - f(x)| \le u(x)$ implies $hg \in M(f, u)$. But $g \in I$ implies $hg \in I$. Therefore $hg \in M(f, u) \cap I$ and f is a limit point of I. This completes the proof.

In Theorem 2.15 we have seen that the uniform norm topology is weaker than the relative m_B -topology on $B_1^*(X)$. The following theorem shows that it is always strictly weaker unless $B_1(X) = B_1^*(X)$. In other words, the equality of uniform norm topology with relative m_B -topology on $B_1^*(X)$ characterizes $B_1(X) = B_1^*(X)$.

Theorem 2.18. The uniform norm topology on $B_1^*(X)$ is same as the relative m_B -topology on $B_1^*(X)$ if and only if $B_1(X) = B_1^*(X)$.

Proof. If $B_1(X) = B_1^*(X)$ then for any $g \in B_1(X)$ and any positive unit u in $B_1(X)$ we get $M(g,u) \cap B_1^*(X) = M(g,u) = \{f \in B_1^*(X) : |f(x) - g(x)| \le u(x), \text{ for every } x \in X\}$. u is a unit in $B_1(X)$ ($= B_1^*(X)$) implies that u is bounded away from 0. So $u(x) \ge \epsilon$, for all $x \in X$ and for some $\epsilon > 0$. Hence $g \in B(g,\epsilon)$ ($= M(g,\epsilon)$) $\subseteq M(g,u)$. Therefore M(g,u) is a neighbourhood of g in $B_1^*(X)$ in the uniform norm topology. Hence this result along with Theorem 2.15 implies that uniform norm topology on $B_1^*(X)$ = the relative m_B -topology on $B_1^*(X)$.

For the converse, let $B_1(X) \neq B_1^*(X)$. Then there exists $f \in B_1(X)$ such that $Z(f) = \emptyset$, f(x) > 0 for all $x \in X$ and f takes arbitrarily small values near 0. So f is a positive unit in $B_1(X)$. Now, for any two real numbers r, s, it will never happen that $|\mathbf{r} - \mathbf{s}| \leq f$. This means that $M(\mathbf{r}, f) \cap \{\mathbf{t} : t \in \mathbb{R}\} = \{\mathbf{r}\}$. So the set $\{\mathbf{r} : r \in \mathbb{R}\}$ is a discrete subspace of $B_1^*(X)$ in the relative m_B -topology. From this it follows that, the scalar multiplication operation $\psi : \mathbb{R} \times B_1^*(X) \to B_1^*(X)$, defined by $\psi(\alpha, g) = \alpha.g$ is not continuous at (r, \mathbf{s}) , where $r, s \in \mathbb{R}$. Thus $B_1^*(X)$ with relative m_B -topology is not a topological vector space and hence the relative m_B -topology is not same as uniform norm topology on $B_1^*(X)$.

Theorem 2.19. $B_1(X)$ with m_B -topology is first countable if and only if $B_1(X) = B_1^*(X)$.

Proof. If $B_1(X) = B_1^*(X)$ then by Theorem 2.18, $B_1(X)$ with m_B -topology is first countable.

Conversely, let $B_1(X) \neq B_1^*(X)$. So there exists $f \in B_1(X) \setminus B_1^*(X)$. Then $g = f^2 + 1$ is a positive unit in $B_1(X)$. We can find a strictly increasing sequence $\{a_n\}$ of positive real numbers and a countable subset $\{p_n\}$ of X such that $g(p_n) = a_n$, for all $n \in \mathbb{N}$.

Consider any countable collection of positive units in $B_1(X)$, say $\{\pi_n\}$.

Let $b_n = \frac{1}{2} \min \left[\pi_1(p_n), \pi_2(p_n), ..., \pi_n(p_n) \right]$. Then there always exists a real valued continuous function $\sigma : \mathbb{R} \to \mathbb{R}$ such that $\sigma(x) > 0$, for all $x \in \mathbb{R}^+$ and $\sigma(a_n) = b_n^{-1}$, for all $n \in \mathbb{N}$.

Define $\psi(x) = \frac{1}{\sigma(g(x))}, \forall x \in X$. By Theorem 1.1 and Theorem 1.3, ψ is a positive unit in $B_1(X)$ and $\psi(p_n) = \frac{1}{\sigma(g(p_n))} = b_n \leq \frac{1}{2}\pi_n(p_n)$, for all $n \in \mathbb{N}$.

Clearly $\frac{2}{3}\pi_n \in M(0,\pi_n)$ but $\frac{2}{3}\pi_n \notin M(0,\psi)$, because at each $p_n, \psi(p_n) \leq \frac{1}{2}\pi_n(p_n) < \frac{2}{3}\pi_n(p_n)$. So the neighbourhood $M(0,\psi)$ at 0 contains no $M(0,\pi_n)$, n=1,2,3,..., which shows that at the point 0 in $B_1(X)$ there is no countable neighbourhood base.

Hence, $B_1(X)$ with m_B -topology is not first countable.

Corollary 2.20. $B_1(X)$ with m_B -topology is metrizable if and only if $B_1(X) = B_1^*(X)$.

3. e_B -IDEALS AND e_B -FILTERS IN $B_1^*(X)$

For any Z_B -filter \mathscr{F} on X, $Z_B^{-1}[\mathscr{F}] \cap B_1^*(X)$ is an ideal in $B_1^*(X)$. Although, for any ideal I of $B_1^*(X)$, $Z_B[I]$ is not in general a Z_B -filter and it is evident from the following example:

Consider the ring $B_1^*(\mathbb{N})$ and the ideal I as the collection of all sequences of real numbers converge to 0. The sequence $\{\frac{1}{n}\} \in I$ but $Z(\{\frac{1}{n}\}) = \emptyset$. $\emptyset \in Z_B[I]$ shows that it is not a Z_B -filter.

In this section, we introduce a special class of Z_B -filters, called e_B -filters. We locate a class of special ideals of $B_1^*(X)$, called e_B -ideals, which behave the same way as the Z_B -ideals in $B_1(X)$. The e_B -ideals and e_B -filters play the pivotal role to establish the desired correspondence.

For $f \in B_1^*(X)$ and $\epsilon > 0$, we define $E_B^{\epsilon}(f) = f^{-1}([-\epsilon, \epsilon]) = \{x \in X : |f(x)| \le \epsilon\}$. Every set of this form is a member of $Z(B_1(X))$, as $E_B^{\epsilon}(f) = Z((|f| - \epsilon) \vee 0)$. In fact, the converse is also true, i.e., every zero set Z(h), $h \in B^*(X)$ is of the form $E_B^{\epsilon}(|h| + \epsilon)$. For every non-empty set $I(\subseteq B_1^*(X))$ we define $E_B^{\epsilon}[I] = \{E_B^{\epsilon}(f) : f \in I\}$ and $E_B(I) = \bigcup_{\epsilon > 0} E_B^{\epsilon}[I] = \{E_B^{\epsilon}(f) : f \in I, \epsilon > 0\}$. For any collection of zero

sets
$$\mathscr{F}$$
, i.e. $\mathscr{F} \subseteq Z(B_1(X))$, we consider $E_B^{\epsilon^{-1}}[\mathscr{F}] = \{f \in B_1^*(X) : E_B^{\epsilon}(f) \in \mathscr{F}\}$ and define $E_B^{-1}(\mathscr{F}) = \bigcap_{\epsilon>0} E_B^{\epsilon^{-1}}[\mathscr{F}] = \{f \in B_1^*(X) : E_B^{\epsilon}(f) \in \mathscr{F}, \, \forall \epsilon>0\}.$

One may easily see that for any two ideals I, J and subcollections \mathscr{F} and \mathscr{G} of $Z(B_1(X))$

$$\begin{array}{ll} \text{(i)} & I \subseteq J \Rightarrow E_B(I) \subseteq E_B(J). \\ \text{(ii)} & \mathscr{F} \subseteq \mathscr{G} \Rightarrow E_B^{-1}(\mathscr{F}) \subseteq E_B^{-1}(\mathscr{G}). \end{array}$$

We record the following facts in the following couple of theorems:

Theorem 3.1. For any subset I of $B_1^*(X)$, $I \subseteq E_B^{-1}(E_B(I))$, where $E_B^{-1}(E_B(I)) = \{ f \in B_1^*(X) : E_B^{\epsilon}(f) \in E_B(I), \forall \epsilon > 0 \}.$

Proof. Let $f \in I$. Then $E_B^{\epsilon}(f) \in E_B^{\epsilon}[I]$, $\forall \epsilon > 0$. By definition $f \in E_B^{-1}(E_B(I))$. Hence, we have the required inclusion.

Theorem 3.2. For any subcollection \mathscr{F} of $Z(B_1(X))$, $E_B(E_B^{-1}(\mathscr{F})) \subseteq \mathscr{F}$, where $E_B(E_B^{-1}(\mathscr{F})) = \bigcup_{\epsilon > 0} \{E_B^{\epsilon}(f) : E_B^{\delta}(f) \in \mathscr{F}, \text{ for all } \delta > 0\}.$

Proof. The proof follows trivially from the definitions of E_B and E_B^{-1} .

It is interesting to note that, in Theorem 3.1, the inclusion may be a strict one even if we consider I as an ideal in $B_1^*(X)$.

For example, consider the ring $B_1^*(\mathbb{N})$ and the function $f(n) = \frac{1}{n}$ in $B_1^*(\mathbb{N})$. Suppose $I = \langle f^2 \rangle$, i.e., the ideal in $B_1^*(\mathbb{N})$ generated by f^2 . It is quite clear that, $f \notin I$. Now we will show that $f \in E_B^{-1}(E_B(I))$ or equivalently, $E_B^{\epsilon}(f) \in E_B(I)$, $\forall \epsilon > 0$. For any $\epsilon > 0$, $E_B^{\epsilon}(f) = \{n \in \mathbb{N} : |f(n)| \le \epsilon\} = \{n \in \mathbb{N} : f^2(n) \le \epsilon^2\} = E_B^{\epsilon^2}(f^2) \in E_B(I)$. So, $I \subsetneq E_B^{-1}(E_B(I))$.

In Theorem 3.2 too, the inclusion may be proper even when \mathscr{F} is a Z_B -filter on X. As an example in support of our claim, we consider the ring $B_1^*(\mathbb{R})$ and the Z_B -filter $\mathscr{F} = \{Z \in Z(B_1(\mathbb{R})) : 0 \in Z\}$ on \mathbb{R} . Now we consider a function $f : \mathbb{R} \mapsto \mathbb{R}$ defined by $f(x) = \frac{|x|}{|x|+1}$. Clearly, $\{0\} = Z(f) \in \mathscr{F}$. We show that $\{0\} \notin E_B(E_B^{-1}(\mathscr{F}))$. Let $g = f + \epsilon$, for any arbitrary $\epsilon > 0$. It is easy to observe that $E_B^{\epsilon}(g) = \{0\}$ but if we take any positive number $\delta < \epsilon$ then $E_B^{\delta}(g) = \emptyset$, which does not belong to \mathscr{F} . Hence, $\{0\} \notin E_B(E_B^{-1}(\mathscr{F}))$.

Definition 3.3. An ideal I in $B_1^*(X)$ is called an e_B -ideal if $I = E_B^{-1}(E_B(I))$. I is an e_B -ideal if and only if for all $\epsilon > 0$, $E_B^{\epsilon}(f) \in E_B(I)$ implies $f \in I$.

From the definition it is clear that the intersection of e_B -ideals is an e_B -ideal.

Definition 3.4. A Z_B -filter \mathscr{F} is said to be an e_B -filter if $\mathscr{F} = E_B\left(E_B^{-1}(\mathscr{F})\right)$. Equivalently, \mathscr{F} is an e_B -filter if and only if whenever $Z \in \mathscr{F}$ there exist $\epsilon > 0$ and $f \in B_1^*(X)$ such that $Z = E_B^{\epsilon}(f)$ and $E_B^{\delta}(f) \in \mathscr{F}$, $\forall \delta > 0$.

In what follows, we consider X to be always a normal (T_4) topological space.

Theorem 3.5. If I is any proper ideal in $B_1^*(X)$ then $E_B(I)$ is an e_B -filter.

Proof. We prove this in two steps. At first we show that $E_B(I)$ is a Z_B -filter and then we establish $E_B(I) = E_B(E_B^{-1}(E_B(I)))$. To show $E_B(I)$ is a Z_B -filter we need to check

- a) $\emptyset \notin E_B(I)$.
- b) $E_B(I)$ is closed under finite intersection.
- c) $E_B(I)$ is closed under superset.

We assert that $\emptyset \notin E_B(I)$. If our assertion is not true then $\emptyset = E_B^{\epsilon}(f)$, for some $f \in I$ and some $\epsilon > 0$. So $|f(x)| > \epsilon$, $\forall x \in X$, which implies that f is bounded away from zero and so f is a unit in $B_1^*(X)$ [1]. This contradicts that I is proper. For b), Let $E_B^{\epsilon}(f)$, $E_B^{\delta}(g) \in E_B(I)$. Then there exist some $f_1, g_1 \in I$ and $\epsilon_1, \delta_1 > 0$ such that $E_B^{\epsilon}(f) = E_B^{\epsilon_1}(f_1)$ and $E_B^{\delta}(g) = E_B^{\delta_1}(g_1)$. We can choose ϵ_1, δ_1 in such a way that $\delta_1 < \epsilon_1$. Then $E_B^{\delta_1^2}(f_1^2 + g_1^2) \subseteq E_B^{\epsilon_1}(f_1) \cap E_B^{\delta_1}(g_1) = E_B^{\epsilon}(f) \cap E_B^{\delta}(g)$. Since I is an ideal, $f_1^2 + g_1^2 \in I$ and $E_B^{\delta_1^2}(f_1^2 + g_1^2) \in E_B(I)$. Therefore, $E_B(I)$ will be closed under finite intersections if we can show (c), i.e., $E_B(I)$ is closed under supersets, which we shall prove now.

Suppose, $E_B^{\epsilon}(f) \in E_B(I)$ and Z(f') $(f' \in B_1^*(X))$ be any member in $Z(B_1(X))$ so that $E_B^{\epsilon}(f) \subseteq Z(f')$. We shall show that $Z(f') \in E_B(I)$. Since $E_B^{\epsilon}(f) = E_B^{\epsilon^2}(f^2)$ and Z(f') = Z(|f'|), we can start with $f, f' \ge 0$.

Let $P = \{x \in X : |f(x)| \ge \epsilon\}.$

We define a function $g: X \mapsto \mathbb{R}$ by

$$g(x) = \begin{cases} 1 & \text{if } x \in E_B^{\epsilon}(f) \\ f'(x) + \frac{\epsilon}{f(x)} & \text{if } x \in P. \end{cases}$$

We observe that $E_B^{\epsilon}(f) \cap P = \{x \in X : f(x) = \epsilon\}$ and $\forall x \in E_B^{\epsilon}(f) \cap P$, $f'(x) + \frac{\epsilon}{f(x)} = 0 + \frac{\epsilon}{\epsilon} = 1$. It is clear that both $E_B^{\epsilon}(f)$ and P are G_{δ} sets and the constant function 1 and $f'(x) + \frac{\epsilon}{f(x)}$ are Baire one functions on $E_B^{\epsilon}(f)$ and P respectively. Therefore by Theorem 1.10, g is a Baire one function on X, in fact $g \in B_1^*(X)$. Now consider the function

 $fg: X \mapsto \mathbb{R}$ given by

$$(fg)(x) = \begin{cases} f(x) & \text{if } x \in E_B^{\epsilon}(f) \\ (ff')(x) + \epsilon & \text{if } x \in P. \end{cases}$$

Since I is an ideal $fg \in I$ and it is easy to check that $Z(f') = E_B^e(fg)$. Hence, $Z(f') \in E_B(I)$. Therefore, $E_B(I)$ is a Z_B -filter. By Theorem 3.1, we get $I \subseteq E_B^{-1}(E_B(I))$. Since the map E_B preserves inclusion, we obtain $E_B(I) \subseteq E_B(E_B^{-1}(E_B(I)))$. Also $E_B(I)$ is a Z_B -ideal, so by Theorem 3.2 $E_B(E_B^{-1}(E_B(I))) \subseteq E_B(I)$. Combining these two we have $E_B(E_B^{-1}(E_B(I))) = E_B(I)$. Hence, $E_B(I)$ is a E_B -filter. \square

Theorem 3.6. For any Z_B -filter \mathscr{F} on X, $E_B^{-1}(\mathscr{F})$ is an e_B -ideal in $B_1^*(X)$.

Proof. We first show that $E_B^{-1}(\mathscr{F})$ is an ideal in $B_1^*(X)$. Let $f,g \in E_B^{-1}(\mathscr{F})$. Therefore for any arbitrary $\epsilon > 0$, $E_B^{\frac{\epsilon}{2}}(f)$, $E_B^{\frac{\epsilon}{2}}(g) \in \mathscr{F}$. \mathscr{F} being a Z_B -filter on X, $E_B^{\frac{\epsilon}{2}}(f) \cap E_B^{\frac{\epsilon}{2}}(g) \in \mathscr{F}$. Also, we know $E_B^{\frac{\epsilon}{2}}(f) \cap E_B^{\frac{\epsilon}{2}}(g) \subseteq E_B^{\epsilon}(f+g)$. Hence, $E_B^{\epsilon}(f+g) \in \mathscr{F}$, or equivalently $f+g \in E_B^{-1}(\mathscr{F})$.

Now consider $f \in E_B^{-1}(\mathscr{F})$ and h be any bounded Baire one function on X with an upper bound M > 0 and ϵ be any arbitrary positive real number. So $|h(x)| \leq M$, for all $x \in X$. For any point $x \in E_B^{\frac{\epsilon}{M}}(f) \Longrightarrow |f(x)| \leq \frac{\epsilon}{M} \Longrightarrow |Mf(x)| \leq \epsilon \Longrightarrow |f(x)h(x)| \leq \epsilon \Longrightarrow x \in E_B^{\epsilon}(fh)$. This implies $E_B^{\frac{\epsilon}{M}}(f) \subseteq E_B^{\epsilon}(fh)$. So $E_B^{\epsilon}(fh) \in \mathscr{F}$, for any arbitrary $\epsilon > 0$. Therefore by definition of $E_B^{-1}(\mathscr{F})$, $fh \in E_B^{-1}(\mathscr{F})$. Hence $E_B^{-1}(\mathscr{F})$ is an ideal in $B_1^*(X)$.

 $E_B^{-1}(\mathscr{F})$ is an ideal in $B_1^*(X)$. By Theorem 3.1, $E_B^{-1}(\mathscr{F}) \subseteq E_B^{-1}(E_B(E_B^{-1}(\mathscr{F})))$. Also by Theorem 3.2, $E_B(E_B^{-1}(\mathscr{F})) \subseteq \mathscr{F}$. Since E_B^{-1} preserves inclusion, $E_B^{-1}(E_B(E_B^{-1}(\mathscr{F}))) \subseteq E_B^{-1}(\mathscr{F})$. Hence $E_B^{-1}(\mathscr{F}) = E_B^{-1}(E_B(E_B^{-1}(\mathscr{F})))$ and so, $E_B^{-1}(\mathscr{F})$ is an e_B -ideal.

Corollary 3.7. The correspondence $I \mapsto E_B(I)$ is one-one from the set of all e_B -ideals in $B_1^*(X)$ onto the set of all e_B -filters on X.

Theorem 3.8. If I is an ideal in $B_1^*(X)$ then $E_B^{-1}(E_B(I))$ is the smallest e_B -ideal containing I.

Proof. It follows from Theorem 3.5 and Theorem 3.6 that $E_B^{-1}(E_B(I))$ is an e_{B} -ideal. Also from Theorem 3.1 we have $I \subseteq E_B^{-1}(E_B(I))$. If possible let \mathscr{J} be any e_B -ideal containing I. So $I \subseteq \mathscr{J}$ and since E_B and E_B^{-1} preserve inclusion, we can write $E_B^{-1}(E_B(I)) \subseteq E_B^{-1}(E_B(\mathscr{J})) = \mathscr{J}$ (since \mathscr{J} is an e_B -ideal). It completes the proof.

Corollary 3.9. Every maximal ideal in $B_1^*(X)$ is an e_B -ideal.

Proof. Follows immediately from the theorem. \Box Corollary 3.10. Intersection of maximal ideals in $B_1^*(X)$ is an e_B -ideal.

Proof. Straightforward. \Box

Theorem 3.11. For any Z_B -filter \mathscr{F} on X, $E_B(E_B^{-1}(\mathscr{F}))$ is the largest e_B -filter contained in \mathscr{F} .

Proof. Theorem 3.5, Theorem 3.6 and Theorem 3.2 show that $E_B(E_B^{-1}(\mathscr{F}))$ is an e_B -filter contained in \mathscr{F} . If possible, let \mathscr{E} be any e_B -filter contained in \mathscr{F} . So $\mathscr{E} \subseteq \mathscr{F} \implies E_B(E_B^{-1}(\mathscr{E})) \subseteq E_B(E_B^{-1}(\mathscr{F})) \implies \mathscr{E} \subseteq E_B(E_B^{-1}(\mathscr{F}))$. This completes the proof.

Lemma 3.12. Let I and J be two ideals in $B_1^*(X)$ with J be an e_B -ideal. Then $I \subseteq J$ if and only if $E_B(I) \subseteq E_B(J)$.

Proof. $I \subseteq J \implies E_B(I) \subseteq E_B(J)$ follows from the definition of $E_B(I)$. For the converse, let $f \in I$. To show that $f \in J$.

Suppose $\epsilon > 0$ be any arbitrary positive number. $f \in I \implies E_B^{\epsilon}(f) \in E_B(I) \implies E_B^{\epsilon}(f) \in E_B(J) \implies f \in J$ (since J is an e_B -ideal). Therefore, $I \subseteq J$.

Lemma 3.13. For any two Z_B -filters \mathscr{F}_1 and \mathscr{F}_2 on X, with \mathscr{F}_1 be an e_B -filter, $\mathscr{F}_1 \subseteq \mathscr{F}_2$ if and only if $E_B^{-1}(\mathscr{F}_1) \subseteq E_B^{-1}(\mathscr{F}_2)$

Proof. Straightforward.

Lemma 3.14. Let \mathscr{A} be any Z_B -ultrafilter. If a zero set Z meets every member of \mathscr{A} then $Z \in \mathscr{A}$.

Proof. Consider the collection $\mathscr{A} \cup \{Z\}$. By hypothesis, it has finite intersection property. So it can be extended to a Z_B -filter with Z as one of its member. As this Z_B -filter contains a maximal Z_B -filter, it must be \mathscr{A} . So $Z \in \mathscr{A}$.

Theorem 3.15. Let \mathscr{A} be any Z_B -ultrafilter. A zero set Z in $Z(B_1(X))$ belongs to \mathscr{A} if and only if Z meets every member of $E_B(E_B^{-1}(\mathscr{A}))$.

Proof. \mathscr{A} is a Z_B -ultrafilter, so $E_B\big(E_B^{-1}(\mathscr{A})\big)\subseteq\mathscr{A}$. If we assume $Z\in\mathscr{A}$ then Z meets every member of \mathscr{A} [2] and hence Z meets every member of $E_B\big(E_B^{-1}(\mathscr{A})\big)$. Conversely, suppose Z meets every member of $E_B\big(E_B^{-1}(\mathscr{A})\big)$. We claim that Z intersects every member of \mathscr{A} . If not, there exists $Z'\in\mathscr{A}$ for which $Z\cap Z'=\emptyset$. Therefore Z and Z' are completely separated in X by $B_1(X)$ [1] and there exists $f\in B_1^*(X)$ such that f(Z)=1 and f(Z')=0. Clearly for all $\epsilon>0$, $Z'\subseteq Z(f)\subseteq E_B^\epsilon(f)$. Now $Z'\in\mathscr{A}$ implies that $E_B^\epsilon(f)\in\mathscr{A}$, for all $\epsilon>0$. If we choose $\epsilon<\frac{1}{2}$, then $Z\cap E_B^\epsilon(f)=\emptyset$, which contradicts that Z meets every member of $E_B(E_B^{-1}(\mathscr{A}))$. Hence, Z meets every member of \mathscr{A} and therefore $Z\in\mathscr{A}$.

Theorem 3.16. If \mathscr{A} is any Z_B -ultrafilter on X then $E_B^{-1}(\mathscr{A})$ is a maximal ideal in $B_1^*(X)$.

Proof. We know from Theorem 3.6 that $E_B^{-1}(\mathscr{A})$ is an ideal in $B_1^*(X)$. Let M^* be a maximal ideal in $B_1^*(X)$ which contains $E_B^{-1}(\mathscr{A})$. By using Lemma 3.12 we can write $E_B(E_B^{-1}(\mathscr{A})) \subseteq E_B(M^*)$. Theorem 3.5 asserts that $E_B(M^*)$ is a Z_B -filter on X, therefore every member of $E_B(M^*)$ meets every member of $E_B(E_B^{-1}(\mathscr{A}))$. By Theorem 3.15 every member of $E_B(M^*)$ belongs to \mathscr{A} . Hence $E_B(M^*) \subseteq \mathscr{A}$.

Every maximal ideal is an e_B -ideal, so $M^* = E_B^{-1}(E_B(M^*)) \subseteq E_B^{-1}(\mathscr{A})$. This implies $M^* = E_B^{-1}(\mathscr{A})$ which completes the proof.

Corollary 3.17. For any Z_B -ultrafilter $\mathscr A$ on X, $E_B^{-1}(\mathscr A) = E_B^{-1}(E_B(E_B^{-1}(\mathscr A)))$.

Definition 3.18. An e_B -filter is called an e_B -ultrafilter if it is not contained in any other e_B -filter. In other words a maximal e_B -filter is called an e_B -ultrafilter.

Using Zorn's lemma one can establish that, every e_B -filter is contained in an e_B -ultrafilter.

Theorem 3.19. If M^* is a maximal ideal in $B_1^*(X)$ and \mathscr{F} is an e_B -ultrafilter on X then

- a) $E_B(M^*)$ is an e_B -ultrafilter on X.
- b) $E_B^{-1}(\mathscr{F})$ is a maximal ideal in $B_1^*(X)$.

Proof. a) Since M^* is a maximal ideal in $B_1^*(X)$ then by Corollary 3.9 we have $M^* = E_B^{-1}(E_B(M^*))$. Suppose the e_B -filter $E_B(M^*)$ is contained in an e_B -ultrafilter \mathscr{F}' . So $E_B(M^*) \subseteq \mathscr{F}' \implies E_B^{-1}(E_B(M^*)) = M^* \subseteq E_B^{-1}(\mathscr{F}') \implies M^* = E_B^{-1}(\mathscr{F}')$ (since M^* is maximal ideal). Therefore, $E_B(M^*) = E_B(E_B^{-1}(\mathscr{F}')) = \mathscr{F}'$. Hence, $E_B(M^*)$ is an e_B -ultrafilter.

b) Let M^* be any maximal extension of the ideal $E_B^{-1}(\mathscr{F})$ in $B_1^*(X)$. Now $E_B^{-1}(\mathscr{F}) \subseteq M^* \implies \mathscr{F} = E_B(E_B^{-1}(\mathscr{F})) \subseteq E_B(M^*)$. By part (a) we can conclude that $\mathscr{F} = E_B(M^*)$, which gives us $E_B^{-1}(\mathscr{F}) = M^*$. Hence we are done.

Corollary 3.20. Let M^* be an e_B -ideal, then M^* is maximal in $B_1^*(X)$ if and only if $E_B(M^*)$ is an e_B -ultrafilter.

Corollary 3.21. An e_B -filter \mathscr{F} is an e_B -ultrafilter if and only if $E_B^{-1}(\mathscr{F})$ is a maximal ideal in $B_1^*(X)$.

Corollary 3.22. The correspondence $M^* \mapsto E_B(M^*)$ is one-one from the set of all maximal ideals in $B_1^*(X)$ onto the set of all e_B -ultrafilters.

Theorem 3.23. If \mathscr{A} is a Z_B -ultrafilter then it is the unique Z_B -ultrafilter containing $E_B(E_B^{-1}(\mathscr{A}))$. In fact, $E_B(E_B^{-1}(\mathscr{A}))$ is the unique e_B -ultrafilter contained in \mathscr{A} .

Proof. Let \mathscr{A}^* be a Z_B -ultrafilter containing $E_B\left(E_B^{-1}(\mathscr{A})\right)$ and $Z \in \mathscr{A}^*$. Clearly Z meets every member of \mathscr{A}^* and so it meets every member of $E_B\left(E_B^{-1}(\mathscr{A})\right)$. By Theorem 3.15 $Z \in \mathscr{A}$ and $\mathscr{A}^* \subseteq \mathscr{A}$. Since both \mathscr{A} and \mathscr{A}^* are Z_B -ultrafilters, hence, we have $\mathscr{A} = \mathscr{A}^*$. So \mathscr{A} is unique one containing $E_B\left(E_B^{-1}(\mathscr{A})\right)$.

For the second part, $E_B(E_B^{-1}(\mathscr{A}))$ is e_B -ultrafilter follows from Theorem 3.16 and Theorem 3.19. To prove the uniqueness we suppose \mathscr{E} be any e_B -ultrafilter contained in \mathscr{A} . Then $\mathscr{E} \subseteq \mathscr{A} \implies E_B(E_B^{-1}(\mathscr{E})) = \mathscr{E} \subseteq E_B(E_B^{-1}(\mathscr{A})) \implies \mathscr{E} = E_B(E_B^{-1}(\mathscr{A}))$. This completes the proof.

Corollary 3.24. Every e_B -ultrafilter is contained in a unique Z_B -ultrafilter.

Proof. Let $\mathscr E$ be an e_B -ultrafilter which is contained in two Z_B -ultrafilters $\mathscr F_1$ and $\mathscr F_2$. Therefore, $\mathscr E=E_B\big(E_B^{-1}(\mathscr E)\big)\subseteq E_B\big(E_B^{-1}(\mathscr F_1)\big)\subseteq \mathscr F_1$. Now by Theorem 3.16, Theorem 3.19 (a) and Theorem 3.23 $E_B\big(E_B^{-1}(\mathscr F_1)\big)$ is the unique e_B -ultrafilter contained in $\mathscr F_1$. Hence, $\mathscr E=E_B\big(E_B^{-1}(\mathscr F_1)\big)$. Similarly, we have $\mathscr E=E_B\big(E_B^{-1}(\mathscr F_2)\big)$.

Therefore, $E_B(E_B^{-1}(\mathscr{F}_1)) = E_B(E_B^{-1}(\mathscr{F}_2))$ is contained in both \mathscr{F}_1 and \mathscr{F}_2 . Now let $Z \in \mathscr{F}_1$. Then \mathscr{F}_1 being a Z_B -ultrafilter, Z intersects every member of $E_B(E_B^{-1}(\mathscr{F}_2))$. So by Theorem 3.15, $Z \in \mathscr{F}_2$. This gives us $\mathscr{F}_1 \subseteq \mathscr{F}_2$. By a similar argument, we can show $\mathscr{F}_2 \subseteq \mathscr{F}_1$. Hence, $\mathscr{F}_1 = \mathscr{F}_2$

Corollary 3.25. There is a one to one correspondence between the collection of all Z_B -ultrafilters on X and the collection of all e_B -ultrafilters on X.

Proof. Let us define a map λ from the set of all Z_B -ultrafilters on X to the set of all e_B -ultrafilters on X by $\lambda(\mathscr{A}) = E_B\big(E_B^{-1}(\mathscr{A})\big)$. We shall show that the map is one-one and onto. To show the injectivity, suppose $\lambda(\mathscr{A}) = \lambda(\mathscr{B})$, which implies $E_B\big(E_B^{-1}(\mathscr{A})\big) = E_B\big(E_B^{-1}(\mathscr{B})\big)$. By Theorem 3.23 we can say $E_B\big(E_B^{-1}(\mathscr{A})\big) = E_B\big(E_B^{-1}(\mathscr{B})\big)$ is contained in both \mathscr{A} and \mathscr{B} . Now let $Z \in \mathscr{A}$. So it intersects every member of $E_B\big(E_B^{-1}(\mathscr{B})\big)$ and by Theorem 3.15 $Z \in \mathscr{B}$. This gives us $\mathscr{A} \subseteq \mathscr{B}$. By a similar argument we can show that $\mathscr{B} \subseteq \mathscr{A}$. Hence, $\mathscr{A} = \mathscr{B}$ and λ is one-one. Again, let \mathscr{E} be any e_B -ultrafilter. By Corollary 3.24 there is a unique Z_B -ultrafilter \mathscr{A} containing \mathscr{E} . Now, $\mathscr{E} \subseteq \mathscr{A} \implies E_B(E_B^{-1}(\mathscr{E})) \subseteq E_B(E_B^{-1}(\mathscr{A})) \implies \mathscr{E} \subseteq E_B(E_B^{-1}(\mathscr{A})) \subseteq \mathscr{A}$. Therefore by using Thorem 3.23, we get $\mathscr{E} = E_B\big(E_B^{-1}(\mathscr{A})\big)$ and this implies λ is onto with $\lambda(\mathscr{A}) = \mathscr{E}$.

In [2] we have shown that there is a bijection between the collection of all maximal ideals in $B_1(X)$ and the collection of all Z_B -ultrafilters on X. Using Corollary 3.22 and Corollary 3.25 we therefore obtain the following:

Theorem 3.26. For any normal (T_4) topological space X, $\mathcal{M}(B_1(X))$ and $\mathcal{M}(B_1^*(X))$ have the same cardinality, where $\mathcal{M}(B_1(X))$, $\mathcal{M}(B_1^*(X))$ are the collections of all maximal ideals in $B_1(X)$ and $B_1^*(X)$ respectively.

Proof. Let us define a map $\psi: \mathcal{M}\big(B_1(X)\big) \to \mathcal{M}\big(B_1^*(X)\big)$ by $\psi(M) = E_B^{-1}\big(Z_B[M]\big)$. From [2] we know that $Z_B[M]$ is a Z_B -ultrafilter on X and by Theorem 3.16 $E_B^{-1}\big(Z_B[M]\big)$ is a maximal ideal in $B_1^*(X)$. We claim that ψ is one-one and onto. Suppose $\psi(M) = \psi(N)$, for some $M, N \in \mathcal{M}\big(B_1(X)\big)$. Clearly, $E_B\big(E_B^{-1}\big(Z_B[M]\big)\big) = E_B\big(E_B^{-1}\big(Z_B[N]\big)\big)$ contained in $Z_B[M]$ as well as $Z_B[N]$. Since each member of $Z_B[M]$ intersects every member of $E_B\big(E_B^{-1}\big(Z_B[N]\big)\big)$, so by Theorem 3.15 every member of $Z_B[M]$ belongs to $Z_B[N]$. Therefore, $Z_B[M] \subseteq Z_B[N]$. Similarly, $Z_B[N] \subseteq Z_B[M]$. Hence, $Z_B[M] = Z_B[N]$ and by Theorem 2.11 of [2] we have M = N, i.e., ψ is one-one.

To prove that ψ is onto we let $M^* \in \mathcal{M}\big(B_1^*(X)\big)$. By Theorem 3.19 $E_B(M^*)$ is an e_B -ultrafilter. Let \mathscr{A} be the unique Z_B -ultrafilter containing $E_B(M^*)$. $Z_B^{-1}[\mathscr{A}]$ belongs to $\mathcal{M}\big(B_1(X)\big)$ [2]. We know $E_B\big(E_B^{-1}(\mathscr{A})\big)$ is the unique e_B -ultrafilter contained in \mathscr{A} . Therefore, $E_B\big(E_B^{-1}(\mathscr{A})\big) = E_B(M^*)$ and $E_B^{-1}(\mathscr{A}) = M^*$ (since both $E_B^{-1}(\mathscr{A})$ and M^* are maximal ideals in $B_1^*(X)$ and maximal ideals are e_B -ideals). Consider $M = Z_B^{-1}[\mathscr{A}]$, clearly $M \in \mathcal{M}\big(B_1(X)\big)$. Now $\psi(M) = E_B^{-1}\big(Z_B[M]\big) = E_B^{-1}\big(Z_B[Z_B^{-1}[\mathscr{A}]]\big) = E_B^{-1}(\mathscr{A}) = M^*$ ([2]), i.e., $Z_B^{-1}[\mathscr{A}]$ is the preimage of M^* . Hence, ψ is onto and it establishes a one to one correspondence between $\mathcal{M}\big(B_1(X)\big)$ and $\mathcal{M}\big(B_1^*(X)\big)$.

The following property characterizes maximal ideals of $B_1^*(X)$.

Theorem 3.27. An ideal M^* in $B_1^*(X)$ is a maximal ideal if and only if whenever $f \in B_1^*(X)$ and every $E_B^{\epsilon}(f)$ intersects every member of $E_B(M^*)$, then $f \in M^*$.

Proof. Suppose $f \in B_1^*(X)$ and every $E_B^{\epsilon}(f)$ meets every member of $E_B(M^*)$, where M^* is a maximal ideal in $B_1^*(X)$. We claim $f \in M^*$. If not, then the ideal $< M^*, f >$ generated by M^* and f must be equal to $B_1^*(X)$. Hence 1 = h + fg, where $g \in B_1^*(X)$ and $h \in M^*$. Let u be an upper bound of g and ϵ $(0 < \epsilon < \frac{1}{2})$ be any pre assigned positive number.

Then we have $\emptyset = E_B^{\epsilon}(1) = E_B^{\epsilon}(h+fg) \supseteq E_B^{\frac{\epsilon}{2}}(h) \cap E_B^{\frac{\epsilon}{2}}(fg) \supseteq E_B^{\frac{\epsilon}{2}}(h) \cap E_B^{\frac{\epsilon}{2u}}(f)$. This is a contradiction to our hypothesis as $E_B^{\frac{\epsilon}{2}}(h) \cap E_B^{\frac{\epsilon}{2u}}(f) = \emptyset$. Hence, $f \in M^*$. Conversely, suppose M^* is any ideal in $B_1^*(X)$ with the given property. Let M^{**} be a maximal ideal containing M^* in $B_1^*(X)$ and $f \in M^{**}$. As $E_B(M^{**})$ is an e_B -filter, $E_B^{\epsilon}(f)$ meets every member of $E_B(M^{**})$, for any $\epsilon > 0$. Since $E_B(M^*) \subseteq E_B(M^{**})$, $E_B^{\epsilon}(f)$ meets every member of $E_B(M^*)$ also. By our hypothesis $f \in M^*$. Therefore, $M^* = M^{**}$.

One may easily observe the following remarks.

Remark 3.28. Any e_B -ideal of $B_1^*(X)$ is closed in $B_1^*(X)$ with uniform norm topology.

Proof. Let I be any e_B -ideal in $B_1^*(X)$. By Theorem 2.2 \overline{I} (Closure of I in the uniform norm topology) is also a proper ideal in $B_1^*(X)$. Clearly, $I \subseteq \overline{I}$. Let us assume $g \in \overline{I}$ and $\epsilon > 0$ be any pre assigned positive real number. Therefore, there must exist an $f \in B(g, \frac{\epsilon}{2}) \cap I$. Now for all $x \in E_B^{\frac{\epsilon}{2}}(f)$, we have $|g(x)| \leq |g(x) - f(x)| + |f(x)| \leq \epsilon$. So, $E_B^{\frac{\epsilon}{2}}(f) \subseteq E_B(g)$. The Z_B -filter $E_B(I)$ contains $E_B^{\epsilon}(f)$, hence it contains $E_B^{\epsilon}(g)$, for all $\epsilon > 0$. Therefore, $g \in E_B^{-1}(E_B(I)) = I$ (as I is an I is an I is an I is closed. \square

Remark 3.29. Any e_B -ideal of $B_1^*(X)$ is closed in $B_1^*(X)$ with respect to the relative m_B -topology. Since $B_1^*(X)$ is closed in $B_1(X)$ with respect to m_B -topology, it then follows that every e_B -ideal of $B_1^*(X)$ is a closed set in $B_1(X)$ with m_B -topology.

References

- A. Deb Ray and Atanu Mondal, On Rings Of Baire One Functions, Appl. Gen. Topol., 20(1) (2019), 237-249.
- [2] A. Deb Ray and Atanu Mondal, Ideals In B₁(X) And Residue Class Rings Of B₁(X) Modulo An Ideal, Appl. Gen. Topol., 20(2) (2019), 379-393.
- [3] Amir Veisi, e_c-Filters and e_c-ideals in the functionally countable subalgebra of C*(X), Appl. Gen. Topol., 20(2) (2019), 395-405.
- [4] E. Hewitt, Rings of real-valued continuous functions. I, Trans. Amer. Math. Soc., 64 (1948), 45–99.
- [5] L. Gillman and M. Jerison, Rings of Continuous Functions. New York: Van Nostrand Rein- hold Co., 1960.
- [6] Libor Vesely, Characterization of Baire-One Functions Between Topological Spaces, Acta Universitatis Carolinae. Mathematica et Physica, 33 (2) (1992), 143-156.
- [7] M.R. Ahmadi Zand, Common properties of some function rings on a topological space, arXiv:2107.02110.

Department of Commerce (E), St. Xavier's college, Mother Teresa sarani, Kolkata - 700016, INDIA

Email address: atanu@sxccal.edu

DEPARTMENT OF PURE MATHEMATICS, UNIVERSITY OF CALCUTTA, 35, BALLYGUNGE CIRCULAR ROAD, KOLKATA - 700019, INDIA

Email address: debrayatasi@gmail.com