arXiv:2007.05602v3 [math.DS] 22 Feb 2022

QUANTITATIVE STATISTICAL PROPERTIES OF TWO-DIMENSIONAL PARTIALLY HYPERBOLIC SYSTEMS

ROBERTO CASTORRINI AND CARLANGELO LIVERANI

ABSTRACT. We study a class of two dimensional partially hyperbolic systems, not necessarily skew products, in an attempt to develop a general theory. As a main result, we provide explicit conditions for the existence of finitely many physical measures (and SRB) and prove exponential decay of correlations for mixing measures. In addition, we obtain precise information on the regularity of such measures (they are absolutely continuous with respect to Lebesgue with density in some Sobolev space). To illustrate the scope of the theory, we show that our results apply to the case of fast-slow partially hyperbolic systems, and for such systems we obtain more precise results on the structure of the SRB measures.

CONTENTS

1. Introduction	2
2. Partially hyperbolic systems	4
2.1. Strongly dominated vertical partially hyperbolic systems (SVPH)	4
2.2. Fast-slow systems	6
3. Main Results	7
3.1. SVPH systems	7
3.2. Fast-Slow systems	8
4. Transfer operators	8
4.1. Transversality of unstable cones	8
5. Preliminary estimates	11
5.1. SVPH [#] systems	11
5.2. C^r -norm	11
5.3. Admissible curves	12
5.4. Estimates for derivatives: SVPH case	14
5.5. Sharp estimates for derivatives: $SVPH^{\sharp}$ case	16
5.6. Iteration of curves: SVPH case	18
5.7. Iteration of curves: the SVPH ^{\sharp} case	19
5.8. Distortion: the SVPH case	20
5.9. Distortion: the SVPH ^{\sharp} case	24
6. A first Lasota-Yorke inequality	27

Roberto Castorrini, Laboratoire de Probabilités, Statistique et Modélisation (LPSM), Sorbonne Université, Université de Paris, 4 Place Jussieu, 75005 Paris, France

CARLANGELO LIVERANI, DIPARTIMENTO DI MATEMATICA, II UNIVERSITÀ DI ROMA (TOR VERGATA), VIA DELLA RICERCA SCIENTIFICA, 00133 ROMA, ITALY.

E-mail addresses: castorrini@lpsm.paris, liverani@mat.uniroma2.it.

This work was partially supported by the PRIN Grant "Regular and stochastic behaviour in dynamical systems" (PRIN 2017S35EHN) and by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement No 787304). C.L. acknowledges the MIUR Excellence Department Project awarded to the Department of Mathematics, University of Rome Tor Vergata, CUP E83C18000100006. R.C. acknowledges the Gran Sasso Science Institute (L'Aquila 67100, Italy), where most of the work was done. CL is member of the GNFM of INDAM. We thank Florestan Martin-Baillon who contributed to the very first stage of this project. It is also a pleasure to thank Viviane Baladi, Mark Demers, Peyman Eslami, Jacopo de Simoi, Dmitry Dolgopyat, Sébastien Gouëzel, Oliver Butterley and Masato Tsujii for many helpful discussions on related topics.

6.1. Differential Operators	28
6.2. Differential operators: the SVPH ^{\sharp} case	30
6.3. Conclusion of the first Lasota-Yorke inequality	32
7. A second Lasota-Yorke inequality: preliminaries	35
7.1. \mathcal{H}^s -norm of \mathcal{L}	36
7.2. Transversality	37
8. A second Lasota-Yorke inequality: Results	38
8.1. Partitions of unity	39
8.2. Fourier basic estimate: case SVPH	41
8.3. Decomposition of Fourier space	43
8.4. The case $\xi^{\perp} \in \mathbf{C}_u$ $(\xi \in \mathcal{Z}_u)$	44
8.5. Proof of Theorem 8.2	47
8.6. Fourier basic estimate: case $SVPH^{\sharp}$	47
8.7. Proof of Theorem 8.3	51
9. The final Lasota-Yorke Inequality	52
10. The map F_{ε}	54
10.1. The maps F_{ε} are SVPH [#]	54
10.2. A non-transversality argument	55
10.3. Proof of Theorem 4.5 .	58
10.4. Eigenfunctions regularity (quantitative)	59
10.5. Proof of Theorem 4.6	60
Appendix A. Proof of Lemma 2.6	62
Appendix B. Proof of Lemma 5.2	65
Appendix C. Black-box: iteration of curves	66
C.1. Proof of Lemma 5.10	66
C.2. Proof of Lemma 5.13	70
C.3. Proof of Lemma 5.14	70
C.4. Proof of Lemma 5.16	73
Appendix D. Proof of Lemma 6.6	76
Appendix E. The space \mathcal{H}^s	78
Appendix F. Vector Field regularity	78
Appendix G. Extension of curves	80
References	83

1. INTRODUCTION

One of the main challenges of the field of Dynamical Systems is to understand the ergodic properties of *partially hyperbolic* systems. Substantial progress has been made in the study of ergodicity starting with [33, 47, 53] until establishing very general results, e.g. [13], in the case of volume preserving diffeomorphisms. Nevertheless, if the invariant measure is not a priori known, then establishing the existence of SRB measures is a serious challenge in itself, see [12, 1, 50] for some important partial results. Moreover, it is well known, at least since the work of Krylov [42], that for many applications ergodicity does not suffice, and mixing (usually in the form of effective quantitative estimates on the decay of correlations) is of paramount importance. Some results on correlation decay exist in the case of mostly expanding central direction [2], and mostly contracting central direction [24, 16]. Such results, albeit important, are often not easy to apply since it is very difficult to estimate the central Lyapunov exponent.

For a central direction with zero Lyapunov exponents (or close to zero) there exist quantitative results on exponential decay of correlations only for group extensions of Anosov maps and Anosov flows [25, 17, 23, 43, 51], but none of them apply to an open class (with the notable exception of [15, 52]; also some form of rapid mixing is known to be typical for large classes of flows [30, 45]).

Hence, the problem of effectively studying the quantitative mixing properties of partially hyperbolic systems is wide open.

Recently, motivated by deep physical reasons [26, 9, 44], the second author has proposed the study of a simple class of partially hyperbolic systems with the goal of developing a theory applicable to a large class of fast-slow systems. Some encouraging results exist [19, 20, 22]. However, the amount of work needed to prove the above partial results has proven rather daunting, and to extend such an approach to more realistic systems seems extremely challenging. To attain substantial progress, it seems necessary to introduce new ideas supplementing the approaches developed so far.

In the last years, starting with [11, 32, 7], a powerful method to investigate the statistical properties of hyperbolic systems has evolved: the functional approach. It consists in the study of the spectral properties of the transfer operator on appropriate Banach spaces. Although the basic idea can be traced back, at least, to Von Neumann ergodic theorem, the new ingredient consists in the understanding that non standard functional spaces must be used, and in the insight of how to embed the key geometrical properties of the system in the topology of the Banach space. See [5] for a recent review of this approach, and [18] for an introduction.

This point of view has produced many important results, e.g. see [43, 40, 41, 31, 29, 27, 8] just to cite a few. It is then natural to investigate if the functional approach can be extended to partially hyperbolic systems. Some results that hint at this possibility already exist (e.g. [3, 28]), however, a general approach is totally missing. Nonetheless, the idea that some quantitative form of accessibility should play a fundamental role has slowly emerged, e.g. see [48, 46, 14].

In this paper we attempt to further the latter approach by combining ideas from [3] and [32]. We find checkable conditions that imply the existence of finitely many physical (and SRB) measures for a large class of two dimensional endomorphisms; in addition, such measures are exponentially mixing (see Theorem 3.1). This implies all the standard statistical results (CLT, Large deviations, deterministic stability, etc; see [4, 5] and references therein for details). Next, we show that the hypothesis of Theorem 3.1 are fulfilled for an open set of physically relevant systems (fast-slow systems), see Theorem 3.3. Moreover, for such systems, we obtain some precise quantitative information on the SRB measures (Theorem 3.4). In addition, we show how the results obtained here provide detailed information on the structure of the peripheral eigenfunctions of the transfer operator, see Theorem 4.6, which hopefully should allow further progress. Indeed, we believe that this approach can be further refined and extended to produce similar results in a much more general class of systems.

It is customary to think that the constants appearing in Lasota-Yorke type inequalities are largely irrelevant. This is certainly not the case for the fast-slow systems discussed in section 10 where the constant in front of the weak norm is highly non-uniform. This is reflected in the possible concentration of the invariant measures. Moreover, the possibility to consider the class of maps discussed there as a perturbation of a limiting case depends crucially on the size of such constants. It was then essential to try to push the estimates to their extreme in order to find out if perturbative ideas could be applied. It turns out that our estimates are not sharp enough to do so. However, we have identified precisely the obstructions to this approach, hence clarifying the focus of future research.

The paper is organized as follows: in the next section we describe the systems under consideration: we call them SVPH, for general partially hyperbolic systems. We will also study, in more detail, a special case: fast-slow systems. Since many results for fast-slow are obtained by refining some estimates already established for the SVPH systems, we made an effort to separate clearly the results for the two classes of systems. Hence, the reader that is not interested in the more technical part of the paper can easily skip it. In section 3 we state our main results on the invariant measures of the systems: Theorem 3.1 and 3.3, which are, respectively, direct consequences of the two technical results Theorems 4.4 and 4.5 on the transfer operators stated in section 4. Finally, the latter section contains two further results (Theorem 4.6 and 4.8) about the spectrum of the transfer operators for fast-slow systems. In section 5 we introduce the necessary notation and prove several facts needed to define the Banach spaces we are interested in. In particular, sections 5.6 and 5.8 contain most of the hard estimates needed in the following and are rather technical, so we postponed the proofs to Appendix C; which can be skipped in a first reading without losing the logic of the argument. In

section 6 we prove a first Lasota-Yorke inequality. Unfortunately, the spaces considered in this section do not embed compactly in each other, hence one cannot deduce the quasi-compactness of the operator from such inequalities. Sections 7 and 8 are the core of the paper where some inequalities relating the previous norms to the Sobolev norms \mathcal{H}^s are obtained. In section 9 we harvest the work done and prove Theorem 4.4 which implies Theorem 3.1. In section 10 we show that fast-slow systems satisfy the hypotheses of Theorem 4.4, hence our theory applies and Theorem 3.3 follows. Also, we take advantage of the peculiarities of the fast-slow systems to prove some sharper results on the SRB measure, and, more generally, the spectral projections (or resonances).

The paper also includes seven appendices that contain some necessary technical results which would have disrupted the flow of the argument if included in the main text.

Notation. As we would like to apply our results to open sets of maps F, all the constants appearing in the text are really functions of F. We will call a constant uniform if it depends continuously only on the C^r norm of the map F, on $(\lambda_- - \mu_+)^{-1}$, χ_c^{-1} , $(1 - \chi_u)^{-1}$, $(1 - \iota_*)^{-1}$ and C_* (see hypothesis (**H1**) for the definition of λ_- , μ_+ , χ_c , ι_* and C_*).¹

In order to make the reading more fluid, we will use the notation $f \leq g$ to mean that there exists a uniform constant $C_{\sharp} > 0$, such that $f \leq C_{\sharp}g$. The values of the constants C_{\sharp} can change from one occurrence to the next. Moreover, in the following we will use $C_{a,b,\ldots}, c_{a,b,\ldots}$ to designate constants that depend also on the quantities a, b, \ldots . When the quantities in the subscripts are fixed, these constants are uniform; hence, since no confusion can arise, we will call them uniform as well.

Note that $\chi_c, \chi_u \in (0, 1)$, which determine the size of the central and unstable cone, respectively, are not uniquely determined by the map. Given our convention, we must keep track of how the constants depend on χ_u^{-1} and we cannot hide such a dependency inside a constant C_{\sharp} . Indeed, in the next sections it will be apparent that it may be convenient to choose χ_u as small as possible while it is convenient to choose χ_c as large as possible.

Finally, to simplify notations, we use $\{a, b, ...\}^+$ and $\{a, b, ...\}^-$ to designate the maximum and minimum between the quantities a, b, ..., respectively.

2. Partially hyperbolic systems

In this section we introduce the class of systems we are interested in, the main assumptions and some definitions necessary to present the results. In this work \mathbb{T}^2 and \mathbb{T} represent the quotients $\mathbb{R}^2/\mathbb{Z}^2$ and \mathbb{R}/\mathbb{Z} respectively. For a local diffeomorphism $F : \mathbb{T}^2 \to \mathbb{T}^2$ we define the functions $\mathfrak{m}_F^*, \mathfrak{m}_F :$ $\mathbb{T}^2 \times \mathbb{R}^2 \setminus \{0\} \to \mathbb{R}_+$ as ²

(2.1)
$$\mathfrak{m}_F(z,v) = \frac{\|D_z Fv\|}{\|v\|} \quad ; \quad \mathfrak{m}_F^*(z,v) = \frac{\|(D_z F)^{-1}v\|}{\|v\|}.$$

2.1. Strongly dominated vertical partially hyperbolic systems (SVPH).

The systems we are interested in are defined in Definition 2.4, but before that we need to introduce some notation.

Let $r \geq 4$ and $F : \mathbb{T}^2 \mapsto \mathbb{T}^2$ be a surjective \mathcal{C}^r local diffeomorphism. We call F a partially hyperbolic system³ if there exist a continuous splitting, not necessarily invariant, of the tangent bundle into subspaces $\mathcal{T}\mathbb{T}^2 = E^c \oplus E^u$, $\sigma > 1$ and c > 0 such that for each $n \in \mathbb{N}$

(2.2)
$$\begin{aligned} \|DF_{|E^{u}}^{n}\| &> c\sigma^{n} \\ \|DF_{|E^{c}}^{n}\| &< c\sigma^{-n} \|DF_{|E^{u}}^{n}\|. \end{aligned}$$

¹The name is motivated by the fact that our results are thought for application to families of maps for which $(\lambda_{-} - \mu_{+})^{-1}, \chi_{c}^{-1}, (1 - \chi_{u})^{-1}, (1 - \iota_{\star})^{-1}$ and C_{\star} are uniformly bounded, hence for such families our constants will be uniform in the usual sense.

²By $\|\cdot\|$ we mean the Riemannian metric in \mathbb{T}^2 induced by the Euclidean norm in \mathbb{R}^2 .

³In the present case the term *partially expanding* would be more appropriate, as there is only an expanding direction which is dominant.

Notice that for non-invertible maps the unstable direction is not necessarily unique, nor backward invariant. It is then more convenient to work with cones instead than distributions. Indeed, it is well known (see e.g [37]) that the above conditions are equivalent to the existence of smooth invariant transversal cone fields $\mathbf{C}_u(z), \mathbf{C}_c(z)$, which satisfy conditions equivalent to (2.2).

To simplify the following arguments we will restrict ourselves to maps without critical points. To further simplify matters we restrict to orientation preserving maps (if not, one can always consider F^2)

(**H0**) for all $p \in \mathbb{T}^2$ we have $\det(D_p F) > 0$.

In addition, to simplify notations, we make the assumption that the cone fields can be chosen constant, since this hypothesis applies to all the examples we have in mind.⁴ Hence we require the following hyperbolicity hypothesis:

(H1) There exist
$$\chi_u \in (0,1), \chi_c \in (0,1]$$
 and $0 < \mu_- < 1 < \mu_+ < \lambda_- \leq \lambda_+$ such that, setting

(2.3)
$$\mathbf{C}_{u} := \{ (\xi, \eta) \in \mathcal{T}_{z} \mathbb{T}^{2} : |\eta| \leq \chi_{u} |\xi| \}$$
$$\mathbf{C}_{c} := \{ (\xi, \eta) \in \mathcal{T}_{z} \mathbb{T}^{2} : |\xi| \leq \chi_{c} |\eta| \},$$

defining (recall equations 2.1)

(2.4)
$$\begin{aligned} \lambda_n^-(z) &:= \inf_{v \in \mathbb{R}^2 \setminus \mathbf{C}_c} \mathfrak{m}_{F^n}(z, v) & \lambda_n^+(z) &:= \sup_{v \in \mathbb{R}^2 \setminus \mathbf{C}_c} \mathfrak{m}_{F^n}(z, v), \\ \mu_n^-(z) &:= \inf_{v \in \mathbf{C}_c \setminus \{0\}} \mathfrak{m}_{F^n}^*(z, v) & \mu_n^+(z) &:= \sup_{v \in \mathbf{C}_c \setminus \{0\}} \mathfrak{m}_{F^n}^*(z, v), \end{aligned}$$

and letting $\lambda_n^- = \inf_z \lambda_n^-(z)$ and $\lambda_n^+ = \sup_z \lambda_n^+(z)$ we assume the following: There exist uniform constants $C_* \ge 1$ and $\iota_* \in (0, 1)$ such that, for all $z \in \mathbb{T}^2$ and $n \in \mathbb{N}$,⁵

(2.5)
$$D_z F \mathbf{C}_u \subset \{(\xi, \eta) : |\eta| \le \iota_\star \chi_u |\xi|\} \Subset \mathbf{C}_u; \quad (D_z F)^{-1} \mathbf{C}_c \subset \{(\xi, \eta) : |\xi| \le \iota_\star \chi_c |\eta|\},$$

(2.6)
$$C_{\star}^{-1}\mu_{-}^{n} \leq \mu_{n}^{-}(z) \leq \mu_{n}^{+}(z) \leq C_{\star}\mu_{+}^{n}; \quad C_{\star}^{-1}\lambda_{-}^{n} \leq \lambda_{n}^{-}(z) \leq \lambda_{n}^{+}(z) \leq C_{\star}\lambda_{+}^{n}.$$

From now on we set $\mu := \{\mu_+, \mu_-^{-1}\}^+ > 1$. The above conditions imply, in particular, det $(DF) \neq 0$. Up to now we have just described a rather general two dimensional partially hyperbolic map. Next, we impose a constraint on the topology of the map

(H2) Let Υ be the family of closed curve $\gamma \in \mathcal{C}^r(\mathbb{T}, \mathbb{T}^2)$ such that ⁶

c0)
$$\gamma' \neq 0$$
,

- c1) γ has homotopy class (0, 1),
- c2) $\gamma'(t) \in \mathbf{C}_c$, for each $t \in \mathbb{T}$.

We assume that for each $\gamma \in \Upsilon$ there exist $\{\gamma_i\}_{i=1}^N \subset \Upsilon$ such that $F^{-1}(\Upsilon) = \{\gamma_i\}_{i=1}^N$.

We also need a *pinching* condition

(2.7)
$$\zeta_r := 6(r+1)!$$

We assume that F satisfies

(2.8)

Remark 2.1. Notice that condition (2.8) implies in particular that $\mu < \lambda_{-}$. The presence of the factorials in (2.7) is probably not optimal. This is a condition we did not try to optimize since it is irrelevant for our main application in which μ is close to 1.

 $\mu^{\zeta_r} < \lambda_-.$

⁴One can reduce to such a case by a change of variables.

 $^{{}^{5}}A \subseteq B$ means $\overline{A} \subset int(B) \cup \{0\}.$

⁶As usual we consider equivalent two curves that differ only by a C^r non-singular reparametrization. In the following we will mostly use curves that are parametrized by vertical length.

We will call a partially hyperbolic system satisfying (2.8) strongly dominated.

Remark 2.2. Note that, since F is a local diffeomorphism, then it can be lifted to a diffeomorphism \mathbb{F} of \mathbb{R}^2 with the projection π map being mod 1, so that $\pi(0) = 0$. Then we can define $\mathbb{G}(x,\theta) =$ $\mathbb{F}(x,\theta) - (0,\theta)$ and write $F \circ \pi(x,\theta) = \pi(\mathbb{G}(x,\theta) + (0,\theta))$. Thus in the following, with a slight abuse of notation, we will often confuse the map with his covering and write

(2.9)
$$F(x,\theta) = (f(x,\theta), \theta + \omega(x,\theta)).$$

In addition, note that if the map satisfies condition (H2) then for each $x \in \mathbb{R}^2$ the curve $\gamma_x(t) = (x, t)$, $t \in \mathbb{T}$ has a preimage $\nu \in \Upsilon$ homotopic to the curve $\bar{\gamma}_p(t) = p + (0,t), \ p \in \nu, \ F(p) = (x,0)$. This implies that $F(\bar{\gamma}_p(t))$ is a curve homotopic to γ_x . Thus for each $(x,\theta) \in \mathbb{R}^2$ the lift has the property $\mathbb{F}(x, \theta + 1) = \mathbb{F}(x, \theta) + (0, 1)$, which implies that \mathbb{G} , and hence ω , is periodic in the second variable.

In the following we will need some quantitative information on the Lipschitz constant of the graphs describing the "unstable manifolds." To simplify matters, we prove the needed results in Lemma F.1. We require then that our maps satisfy the hypotheses of such a Lemma. However, be aware that such hypotheses are not optimal and the following condition is used only in Lemma F.1. Hence, the next assumption becomes superfluous if in a given system one can prove Lemma F.1 independently. Also note that, in some cases, it is implied by (H3).

(H4) With the notation (2.9) we require, for each $p \in \mathbb{T}^2$,

$$\partial_x f(p) > \{2(1 + \|\partial_x \omega\|_\infty), |\partial_\theta f(p)|\}^+$$

Remark 2.3. Note that one can always have $\chi_c = 1$ by a linear change of variables, yet we prefer to keep track of χ_c since it may be useful in cases where assumption (H4) is not used.

Definition 2.4 (SVPH systems). We call a map F a strongly dominated vertical partially hyperbolic system (SVPH for simplicity) if it satisfies assumptions (H0),.., (H4).

Remark 2.5. Note that if F satisfies (H0), (H1) and (H2), then so does F^n , $n \in \mathbb{N}$. Thus it may be convenient to consider F^n , instead of F, to check (H3) and (H4).

From now on we will write a SVPH in the form (2.9) when convenient. Here we provide simple explicit conditions implying (H0),.., (H4). The proof is in Appendix A.

Lemma 2.6. Let $\lambda := \inf_{\mathbb{T}^2} \partial_x f, \Lambda := \sup_{\mathbb{T}^2} \partial_x f$ and suppose that:

- (1) $\partial_x f(p) > \{2(1 + \|\partial_x \omega\|_\infty), |\partial_\theta f(p)|\}^+, \quad \forall p \in \mathbb{T}^2,$
- (2) $\|\partial_x \omega\|_{\infty} + \|\partial_\theta \omega\|_{\infty} < \frac{1}{2}$,

- $\begin{array}{l} (2) \|\partial_{x}\omega\|_{\infty} + \|\partial_{\theta}\omega\|_{\infty} < \frac{1}{2}, \\ (3) \|\partial_{\theta}\omega\|_{\infty} < \frac{1+\|\partial_{x}\omega\|_{\infty}}{\lambda^{-1}}, \\ (4) \ 1+\|\partial_{\theta}f\|_{\infty} + \|\partial_{\theta}\omega\|_{\infty} + \|\partial_{x}\omega\|_{\infty} < \lambda, \\ (5) \ \|\partial_{\theta}f\|_{\infty} < \frac{1}{2} \left(-1 + \sqrt{1+2\lambda^{2}\Lambda^{-1}}\right), \end{array}$
- (6) $\chi_c \|\partial_x \omega\|_{\infty} + \|\partial_\theta \omega\|_{\infty} < \frac{\ln \lambda}{4\zeta_r},$

with ζ_r as in (2.7). Then F satisfies assumptions (H0),...,(H4) with χ_u given by (A.4), (A.11), χ_c given by (A.6), in particular $\chi_c = 1$ is allowed, and

(2.10)
$$\mu := \{ (1 - \chi_c \|\partial_x \omega\|_\infty - \|\partial_\theta \omega\|_\infty)^{-1}, e^{\chi_c \|\partial_x \omega\|_\infty + \|\partial_\theta \omega\|_\infty)} \}^+$$

In the next section we introduce an interesting example of SVPH. Despite of their simple form, the endomorphisms we are going to consider still include a large class of physically relevant systems.

2.2. Fast-slow systems.

We are specially interested in the following class of systems, introduced in [21] (and inspired by the more physically relevant model in [26]). We will call these systems fast-slow.

Definition 2.7 (Fast-Slow systems). Let $F_0(x,\theta) = (f(x,\theta),\theta)$ be $\mathcal{C}^r(\mathbb{T}^2,\mathbb{T}^2)$, for $r \geq 4$, such that $\inf_{(x,\theta)\in\mathbb{T}^2}\partial_x f(x,\theta) \geq \lambda > 2$. For any $\omega \in \mathcal{C}^r(\mathbb{R}^2,\mathbb{R})$, periodic of period one, and $\varepsilon > 0$, we define (2.11) $F_{\varepsilon}(x,\theta) = (f(x,\theta), \theta + \varepsilon \omega(x,\theta)).$

We call the maps F_{ε} fast-slow if they satisfy assumption (5) of Lemma 2.6.

In the following we will need also the next definition.

Definition 2.8. The function $\omega \in C^0(\mathbb{T}^2, \mathbb{R})$ is called x-constant with respect to F_0 if there exist $\theta \in \mathbb{T}$, $\Phi_\theta \in C^0(\mathbb{T}, \mathbb{R})$ and a constant $c \in \mathbb{R}$ such that, for each $x \in \mathbb{T}$,

$$\omega(x,\theta) = \Phi_{\theta}(f(x,\theta)) - \Phi_{\theta}(x) + c.$$

Note that, given a specific F_{ε} , one can often check that ω is not x-constant by looking at the shortest periodic orbits. Moreover, according to [20], being not x-constant is a generic condition. Our main results on F_{ε} hold under the assumption that ω is not x-constant.

3. Main Results

Here we detail our main results, first for general systems, then for fast-slow systems.

3.1. SVPH systems. A physical measure is an F-invariant probability measure μ_{ph} such that the set

$$B(\mu_{ph}) := \left\{ p \in \mathbb{T}^2 : \frac{1}{n} \sum_{k=0}^{n-1} \delta_{F^k(p)} \to \mu_{ph} \quad \text{weakly as} \quad n \to \infty \right\}$$

has positive Lebesgue measure.

To state our first result we introduce a quantity inspired by [48]. Given $y \in \mathbb{T}^2$ and a line L in \mathbb{R}^2 passing through the origin, define

(3.1)
$$\widetilde{\mathcal{N}}_F(n, y, L) := \sum_{\substack{z \in F^{-n}(y) \\ DF^n(z) \mathbf{C}_u \supset L}} |\det DF^n(z)|^{-1},$$

and we set $\widetilde{\mathcal{N}}_F(n) = \sup_{y \in \mathbb{T}^2} \sup_L \widetilde{\mathcal{N}}_F(n, y, L).$

In addition, for each integer $1 \le s \le r - 1$ we define⁷

(3.2)
$$\alpha = \frac{\log(\lambda_{-}\mu^{-2})}{\log(\lambda_{+})}$$
$$\alpha_s := 2(2+s-\alpha) \quad ; \quad \beta_s := 2(s+2).$$

Theorem 3.1. Let $F \in \mathcal{C}^r(\mathbb{T}^2, \mathbb{T}^2)$ be SVPH. We assume that there exist $n_1 \in \mathbb{N}$ and $\nu_0 \in (0,1)$ such that, for some $1 \leq s \leq r-1$,

(3.3)
$$\left\{\mu^{\zeta_s}\lambda_-^{-1}, \sqrt{\widetilde{\mathcal{N}}_F(\lceil \alpha n_1 \rceil)}\mu^{\alpha_s n_1 + \beta_s \overline{m}_{\chi_u}}\right\}^+ < \nu_0$$

where \overline{m}_{χ_u} is defined in (5.10). Then there exist finitely many ergodic physical measures and they are absolutely continuous with densities in the Hilbert space $\mathcal{H}^s(\mathbb{T}^2)$, hence they are SRB measures as well.⁸ Moreover, each mixing physical measure μ_{ph} enjoys exponential decay of correlations for Hölder observables ϕ, φ , namely there exist $\nu > 0$ such that,

$$\left|\mu_{ph}\left(\phi\cdot\varphi\circ F^{n}\right)-\mu_{ph}(\phi)\mu_{ph}(\varphi)\right|\leq C_{\phi,\varphi}e^{-\nu n}$$

The proof of Theorem 3.1 can be found at the end of subsection 4.1.2.

Remark 3.2. Under the assumption (**H3**), condition (3.3) is automatically satisfied if $\tilde{\mathcal{N}}_F$ grows subexponentially with n. According to [50], this latter fact holds generically for partially hyperbolic systems in two dimensions (for more details see Remark 8.1). Hence, the result and all the consequences of Theorem 3.1 hold generically. Unfortunately, this does no say much about a specific map F. However, given a map F, (3.3) is an explicit condition about some power αn_1 of F that one can try to check. If successful, then Theorem 3.1 applies to F.

⁷Note that in (3.2) $\alpha \in (0, 1)$, thanks to hypothesis (2.8).

⁸In general non invertible systems do not have an unstable manifold, but many of them, depending on the past history selected, so the SRB should be absolutely continuous when restricted to all such manifolds; this is the case since the physical measures are absolutely continuous.

3.2. Fast-Slow systems. Even though it is generic and checkable, condition (3.3) of Theorem 3.1, may be quite laborious to check and it may entail some computer assisted strategy. It is then interesting to consider less general examples in which such a condition can be easily verified. An important example is given by the fast-slow systems F_{ε} introduced in section 2.2. For fast-slow systems condition (3.3) is directly related to the condition of ω not being x-constant (see Definition 2.8).⁹

Theorem 3.3. There exists ε_* such that, if ω is not x-constant, then for each $1 \le s \le r-1$ and $\varepsilon \in (0, \varepsilon^*)$, F_{ε} has only finitely many physical measures. The physical measures are absolutely continuous, with densities in the Hilbert space $\mathcal{H}^s(\mathbb{T}^2)$, hence they are SRB measures as well. Moreover, there exist $\nu > 0$ such that each mixing physical measure μ_{ph}^{ε} enjoys exponential decay of correlations for Hölder observables ϕ, φ , namely,

$$\left|\mu_{ph}^{\varepsilon}\left(\phi\cdot\varphi\circ F_{\varepsilon}^{n}\right)-\mu_{ph}^{\varepsilon}(\phi)\mu_{ph}^{\varepsilon}(\varphi)\right|\leq C_{\phi,\varphi,\varepsilon}e^{-\nu n}.$$

The proof of Theorem 3.3 can be found at the end of subsection 4.1.2.

In this case, we can also establish some refined properties of SRB measures.

Theorem 3.4. Let $h_{\varepsilon,1} \in L^1$, $||h_{\varepsilon,1}||_{L^1} = 1$, be the density of a physical measure. Then, setting $\gamma_{\varepsilon}(\theta) = \int h_{\varepsilon,1}(y,\theta) dy$, we have

(3.4)
$$\operatorname{dist}_{W_1}(h_{\varepsilon,1} \cdot \operatorname{Leb}, h_* \cdot \gamma_{\varepsilon} \cdot \operatorname{Leb}) \leq C_{\sharp} \varepsilon (\ln \varepsilon^{-1})^2.$$

where $\operatorname{dist}_{W_1}$ is the Wasserstein distance and Leb is the Lebesgue measure on \mathbb{T}^2 , and $h_*(\cdot, \theta)$ is the unique invariant probability density of $f(\cdot, \theta)$. In addition, for all $\beta > \frac{13}{2}$, there exists $C_\beta > 0$:

$$\|h_{\varepsilon,1}\|_{\mathcal{H}^1} \leq C_\beta \varepsilon^{-\beta}.$$

The proof of Theorem 3.4 can be found at the end of subsection 4.1.2 where, in fact, some more precise results are stated.

4. TRANSFER OPERATORS

We will prove Theorems 3.1 and 3.3, 3.4 using a Transfer operator.

Definition 4.1. Given a map $F : \mathbb{T}^2 \to \mathbb{T}^2$, we define $\mathcal{L}_F : L^1(\mathbb{T}^2) \to L^1(\mathbb{T}^2)$, the transfer operator associated to F, as

(4.1)
$$\mathcal{L}_F u(z) = \sum_{y \in F^{-1}(z)} \frac{u(y)}{|\det(D_y F)|}$$

Iterating (4.1) yields, for all $n \in \mathbb{N}$,

(4.2)
$$\mathcal{L}_F^n u(z) = \sum_{y \in F^{-n}(z)} \frac{u(y)}{|\det(D_y F^n)|}$$

By a simple change of variables it follows that $\|\mathcal{L}_F u\|_{L^1} \leq \|u\|_{L^1}$.

4.1. Transversality of unstable cones.

Starting from [50] and following [28],[48], [14] and [54], a link between the mixing property of a partially hyperbolic system¹⁰ and a transversality condition of unstable cones clearly emerges. Let us recall the following notion of transversality introduced in [48] by Tsujii.

Definition 4.2. Given $n \in \mathbb{N}$, $y \in \mathbb{T}^2$ and $z_1, z_2 \in F^{-n}(y)$, we say that z_1 is transversal to z_2 (at time n) if $D_{z_1}F^n\mathbf{C}_u \cap D_{z_2}F^n\mathbf{C}_u = \{0\}$, and we write $z_1 \pitchfork z_2$.

⁹This relation was already remarked in [50] and [14] in the special case of skew-products.

 $^{^{10}\}mathrm{Although}$ restricted to cases in which the central direction is unidimensional.

To make the notion of transversality more quantitative, for each $y \in \mathbb{T}^2$ and $z_1 \in F^{-n}(y)$ we define

(4.3)
$$\mathcal{N}_F(n, y, z_1) := \sum_{\substack{z_2 \notin |z_1|\\z_2 \in F^{-n}(y)}} |\det D_{z_2} F^n|^{-1}$$

and set $\mathcal{N}_F(n) = \sup_{y \in \mathbb{T}^2} \sup_{z_1 \in F^{-n}(y)} \mathcal{N}_F(n, y, z_1).$

Remark 4.3. Note that if all the preimages are non-transversal, then the sum in (4.3) corresponds to the classical transfer operator applied to one, $\mathcal{L}_{F}^{n}1$ (see (4.2)).

In essence, $\mathcal{L}_{F}^{n}1-\mathcal{N}_{F}(n)$ provides a quantitative version of the notion of accessibility in our systems.

In Lemma 7.4 we explain the relation between \mathcal{N}_F and $\widetilde{\mathcal{N}}_F$ (defined in 3.1), while in section 7.2 we explore the properties of $\widetilde{\mathcal{N}}_F$.

4.1.1. *Partially hyperbolic systems.* We are now ready to state the main technical result for SVPH. The proof of the following Theorem is in Section 9.

Theorem 4.4. Let $F \in C^r(\mathbb{T}^2, \mathbb{T}^2)$ be SVPH that satisfies (3.3). Then there exists a Banach space $\mathcal{B}_{s,*}, C^{r-1}(\mathbb{T}^2) \subset \mathcal{B}_{s,*} \subset \mathcal{H}^s(\mathbb{T}^2)$ such that $\mathcal{L}_F(\mathcal{B}_{s,*}) \subset \mathcal{B}_{s,*}$.¹¹ The restriction of \mathcal{L}_F to $\mathcal{B}_{s,*}$ is a bounded quasi-compact operator, with spectral radius one and essential spectral radius smaller than ν_0 .

Proof of Theorem 3.1. Note that if $\mathcal{L}_F h = h$, then the measure h Leb is invariant for F. On the other hand if μ_{ph} is a physical measure, then there exists a set $K \subset \mathbb{T}^2$, Leb(K) > 0, $\mu_{ph}(K) = 1$ such that, for each $x \in K$, $\lim_{n\to\infty} \frac{1}{n} \sum_{k=1}^{n-1} \delta_{F^k(x)}$ converges weakly to μ_{ph} . By Lusin theorem and the density of \mathcal{C}^{∞} in \mathcal{C}^0 , for each $\varepsilon > 0$ there exists $g_{\varepsilon} \in \mathcal{C}^{\infty}$ such that $\int_{\mathbb{T}^2} |\mathbb{1}_K - g_{\varepsilon}| \leq \varepsilon$. We can approximate it weakly by measures g_{ε} Leb. Then, for each $\phi \in \mathcal{C}^r$,

$$\mu_{ph}(\phi) = \frac{1}{\operatorname{Leb}(K)} \int_{K} \lim_{n \to \infty} \frac{1}{n} \sum_{k=1}^{n-1} \mathbb{1}_{K} \phi \circ F^{k} = \lim_{n \to \infty} \frac{1}{n \operatorname{Leb}(K)} \sum_{k=1}^{n-1} \int_{K} \mathbb{1}_{K}(x) \phi \circ F^{k}(x)$$
$$= \lim_{n \to \infty} \frac{1}{n \operatorname{Leb}(K)} \sum_{k=1}^{n-1} \int_{\mathbb{T}^{2}} \mathcal{L}_{F}^{k} g_{\varepsilon} \phi + \mathcal{O}(\|\phi\|_{\infty} \varepsilon)$$

where the second equality follows by Lebesgue dominate convergence Theorem. Since Theorem 4.4 implies that $\lim_{n\to\infty} \frac{1}{n} \sum_{k=1}^{n-1} \mathcal{L}_F^k g_{\varepsilon}$ converges in \mathcal{H}^s to the projector Π which projects on the finite dimensional eigenspace associated to the eigenvalue 1, we have

$$\mu_{ph}(\phi) = \int_{\mathbb{T}^2} \Pi g_{\varepsilon} \cdot \phi.$$

Thus μ_{ph} is a convex combination of $\{h_i\}$ such that $\mathcal{L}_F h_i = h_i$. Hence, there are finitely many ergodic physical measure and they are absolutely continuous with density in \mathcal{H}^s . Obviously their supports are disjoint, hence if one is mixing with density h, we have that, if P_* is the projector on the eigenspace associated to the eigenvalues of modulus one, then $P_*(\phi h) = \mu_{ph}(\phi)h$ for each $\phi \in \mathcal{C}^r$. Hence,

$$\mu_{ph}(\phi \cdot \varphi \circ F^n) = \int_{\mathbb{T}^2} \mathcal{L}_F^n(\phi h) \varphi = \mu_{ph}(\phi) \mu_{ph}(\varphi) + \mathcal{O}(e^{-\nu n})$$

where $e^{-\nu}$ is larger than the modulus of the largest eigenvalue not on the unit circle.

4.1.2. **Fast-Slow systems.** As already remarked, the condition that ω is not x-constant is much easier to check than (3.3). The following theorem is proven in section 10.3.

Theorem 4.5. There exists ε_* such that the map F_{ε} is a SVPH (see Definition 2.4) for any $\varepsilon \in (0, \varepsilon_*)$. In addition, if ω is not x-constant, then there exists $\sigma_* \in (0,1)$ such that the transfer operator $\mathcal{L}_{F_{\varepsilon}}$ is quasi compact on the spaces $\mathcal{B}_{s,*}$, with spectral radius one and essential spectral radius bounded by σ_* for all $\varepsilon \in (0, \varepsilon_*)$.

¹¹By $\mathcal{H}^{s}(\mathbb{T}^{2})$ we mean the usual Sobolev space (see Appendix E).

Consider the operator $P: L^1(\mathbb{T}^2) \to L^1(\mathbb{T}^2)$ defined by

$$Ph(x,\theta) = h_*(x,\theta) \int_{\mathbb{T}} dy h(y,\theta).$$

The following Theorem is proved in section 10.5.

Theorem 4.6. In the hypothesis of Theorem 4.5, we have the decomposition $\mathcal{L}_{F_{\varepsilon}} = \Pi + Q$ where $\Pi Q = Q\Pi = 0$, Π is the finite rank projector on the eigenspace associated to the eigenvalues of modulus one, and Q has spectral radius strictly smaller than one. Moreover,

(4.4)
$$\|\Pi - P\Pi\|_{L^1 \to (\mathcal{C}^1)'} \le C_{\sharp} \varepsilon [\ln \varepsilon^{-1}]^2.$$

Finally, for each $\tau > 0$, let h_{ν} , $\|h_{\nu}\|_{L^1} = 1$, be an eigenfunction associated to the eigenvalue ν with $|\nu| \ge e^{-\varepsilon^{\tau}}$. Then, setting $\beta(\theta) = \int h_{\nu}(y, \theta) dy$, we have

(4.5)
$$\|h_{\nu} - h_*\beta\|_{(\mathcal{C}^1)'} \le C_{\sharp}\varepsilon^{\min\{1,\tau\}}(\ln\varepsilon^{-1})^2.$$

Remark 4.7. The above Theorem is much stronger than the results in [50] (where only the existence of the physical measure is discussed and the results hold only generically) or [12, 1] (where no information on the SRB measure is provided and its existence is obtained under an additional condition on the contraction or the expansion in the center foliation, even though for more general systems).

However, the papers [19, 20, 21, 22] show that, using the standard pair technology and investigating limit theorems, in some special cases it is possible to obtain considerably more detailed information on the system. Unfortunately, on the one hand the necessary arguments in [20] are rather involved and, on the other hand, the conclusions concerning the physical measure in [19] hold only for mostly contracting systems (contrary to the present case that holds in full generality). It is then very important to investigate if the present strategy can provide further information.

First of all we have an explicit bound on the regularity of the eigenfunctions. The reader can find the proof of the following theorem at the end of section 10.4.

Theorem 4.8. If ω is not x-constant, then there exists $c_* > 0$ such that, for each $\varepsilon > 0$ small enough, and $\mathbf{r} \in (0,1)$, if $\nu \in \sigma_{\mathcal{B}_{1,*}}(\mathcal{L}_{F_{\varepsilon}}) \cap \{z \in \mathbb{C} : 1 - \mathbf{r}c_*[\ln \varepsilon^{-1}]^{-1} \leq |z|\}$, and u is an eigenvector of $\mathcal{L}_{F_{\varepsilon}}$ with eigenvalue ν , and $||u||_{\mathcal{B}_0} = 1$,¹² then for all $\beta > \frac{13}{2}$,

$$\|u\|_{\mathcal{H}^1} \le C_\beta \varepsilon^{-(1-\mathbf{r})^{-1}\beta}.$$

Remark 4.9. It is not clear how sharp the above Theorem is. Certainly some form of blow-up is inevitable. For example: let $f_{\theta}(\cdot) = f(x,\theta)$ and call $h_*(\cdot,\theta)$ the unique invariant probability density of f_{θ} . Let $\bar{\omega}(\theta) = \int_{\mathbb{T}} \omega(x,\theta)h_*(x,\theta)dx$. If $\bar{\omega}$ has non degenerate zeroes $\{\theta_i\}_{i=1}^N$ such that $\bar{\omega}'(\theta_i) < 0$, then [22] (see also the discussion below) implies that there must exist an eigenfunction u essentially concentrated in the $\sqrt{\varepsilon}$ neighborhood of each θ_i . This implies that $||u||_{\mathcal{H}^1} \geq C_{\sharp}\varepsilon^{-\frac{1}{4}}$. However, there is a large gap between such a lower bound and the upper bound provided by Theorem 4.8. In particular, much more information on the spectrum could be obtained if one could establish an upper bound of the type $\varepsilon^{-\beta}$ with $\beta < 1$. We regard this as an open problem.

Finally, in the setting of Remark 4.9, let $\widehat{P}: L^1 \to (\mathcal{C}^0)'$ be the finite rank operator defined by: for all $\varphi \in \mathcal{C}^0$

$$\int_{\mathbb{T}^2} \varphi(x,\theta) [\widehat{P}h](dx,d\theta) = \sum_j \int_{\mathbb{T}} dx \varphi(x,\theta_j) h_*(x,\theta_j) \int_{\mathbb{T} \times U_j} dy \, dsh(y,s) + \sum_j \int_{\mathbb{T}^2} dy \,$$

where U_j is the basin of attraction of the stable equilibrium point θ_j of the averaged dynamics

(4.6)
$$\begin{aligned} \bar{\theta} &= \bar{\omega}(\bar{\theta}) \\ \bar{\theta}(0,\theta) &= \theta. \end{aligned}$$

¹²See Section 6 for the definition of the space \mathcal{B}_0 .

Then, an immediate consequence of Theorem 4.5 and [22, Proposition 4] is that the eigenfunctions h for the eigenvalue 1 satisfy, for $\gamma \in (0, \frac{1}{4})$,

(4.7)
$$\|h - \widehat{P}h\|_{L^1 \to (\mathcal{C}^1)'} \le \left(C_{\sharp} \varepsilon^{1/2 - 2\gamma} + C_{\sharp} \varepsilon \ln \varepsilon^{-1}\right).$$

Remark 4.10. Note that the results of [22] are conditional to the existence of the physical measure which has been previously proven only generically or in special cases, see Remark 4.7. On the contrary here the existence of the physical measures is ensured by Theorems 4.5, 4.6, regardless of the value of the central Lyapunov exponent.

Proof of Theorems 3.3 and 3.4. The proof follows from Theorems 4.5 by the same exact arguments used in the proof of Theorem 3.1. Theorem 3.4 follows trivially from Theorem 4.8. \Box

5. Preliminary estimates

In this section we provide several basic definitions and we prove many estimates that will be extensively used in the following. To deal with the fast-slow systems F_{ε} , and in particular to prove Theorem 4.8, we will need to know explicitly how the constants appearing in this section will behave depending on ε . This makes the computations much longer and tedious than in the general case, where it is not essential to explicitly bound so many constants. For this reason, we will keep the sharper estimates needed for F_{ε} as much as possible separated from general ones for the SVPH. The reader can thus skip them if not interested in the results, proven in section 10, about F_{ε} .

For convenience, and for possible future use, we will consider the fast-slow case as a special case of a larger class of systems, the "SVPH[#] systems".

5.1. **SVPH^{\ddagger}** systems.

The definition of SVPH^{\sharp} systems is a bit technical and is motivated by the need to prove Lemma 5.14. In particular, it depends on uniform constants \bar{c}_{\flat}, c_2^- introduced in Proposition 5.8 and equation (5.12) respectively, and the constants \bar{n}, C_{\flat} introduced in Lemma 5.10.

Definition 5.1 (SVPH[#] systems). A map F is called $SVPH^{\sharp}$ if it is a SVPH and it satisfies the following additional conditions: there exist $\hat{n} \in \mathbb{N}$ uniform with $\hat{n} > \bar{n}$ such that, given $B_{\hat{n}} := B_{\hat{n},\bar{c}_{\flat},c_{2}^{-},C_{\flat}}$ defined in (5.35) of Lemma 5.14, we have¹³

(5.1)
$$B_{\hat{n}}^{\frac{1}{n}} \mu^{30} \lambda_{-}^{-1} \le \frac{1}{2}$$

(5.2)
$$\hat{n} < c_2^{-} \ln \chi_u^{-1}.$$

We will see in section 10.1 that, for ε small enough, fast-slow systems are SVPH^{\ddagger}.

5.2. C^{r} -norm.

Since we will need to work with high order derivatives, it is convenient to choose a norm $\|\cdot\|_{\mathcal{C}^r}$ equivalent to the standard one, which ensures our spaces to be Banach Algebras. We thus define the weighted norm in $\mathcal{C}^r(\mathbb{T}^2, \mathcal{M}(m, n))$, where $\mathcal{M}(m, n)$ are the $m \times n$ matrices,¹⁴

(5.3)
$$\begin{aligned} \|\varphi\|_{\mathcal{C}^{0}} &= \sup_{x \in \mathbb{T}^{2}} \sup_{i \in \{1,..,n\}} \sum_{j=i}^{m} |\varphi_{i,j}(x)| \\ \|\varphi\|_{\mathcal{C}^{\rho+1}} &= 2^{\rho+1} \|\varphi\|_{\mathcal{C}^{0}} + \sup_{i} \|\partial_{x_{i}}\varphi\|_{\mathcal{C}^{\rho}}. \end{aligned}$$

¹³Remark that condition (5.1) can be satisfied thanks to hypothesis (H3).

¹⁴According with the previous notations we set $x_1 = x$ and $x_2 = \theta$.

where, for a multi-index $\alpha = (\alpha_1, ..., \alpha_k)$ with $\alpha_k \in \{1, 2\}$, and we will use the notation $|\alpha| = k$ and $\partial^{\alpha} = \partial_{x_{\alpha_1}} \cdots \partial_{x_{\alpha_k}}$.¹⁵ The above definition implies

(5.4)
$$\|\varphi\|_{\mathcal{C}^{\rho}} \leq \sum_{k=0}^{\rho} 2^{\rho-k} \sup_{|\alpha|=k} \|\partial^{\alpha}\varphi\|_{\mathcal{C}^{0}}.$$

We will often need to compute the \mathcal{C}^{ρ} norm of φ along a curve $\nu \in \mathcal{C}^{r}(\mathbb{T}, \mathbb{T}^{2})$. In this case we use the notation $\|\varphi\|_{\mathcal{C}^{\rho}_{\nu}} := \|\varphi \circ \nu\|_{\mathcal{C}^{\rho}}$.

The following Lemma is proven in Appendix B. Note that the estimates in the Lemma are not sharp, however they try to optimize the balance between simplicity and usefulness.¹⁶

Lemma 5.2. For every $\rho, n, m, s \in \mathbb{N}_0, \psi \in \mathcal{C}^{\rho}(\mathbb{T}^2, \mathcal{M}(n, m))$ and $\varphi \in \mathcal{C}^{\rho}(\mathbb{T}^2, \mathcal{M}(m, s))$ we have $\|\varphi\psi\|_{\mathcal{C}^{\rho}} \leq \|\varphi\|_{\mathcal{C}^{\rho}} \|\psi\|_{\mathcal{C}^{\rho}}.$

Also, there exists $C_j^{\star} > 0$, $j \in \mathbb{N}$, such that, if $\varphi \in \mathcal{C}^{\rho}(\mathbb{T}^2, \mathcal{M}(n, m))$ and $\psi \in \mathcal{C}^{\rho}(\mathbb{T}^2, \mathbb{T}^2)$,

(5.5)
$$\|\varphi \circ \psi\|_{\mathcal{C}^{\rho}} \le C^{\star}_{\rho} \sum_{s=0}^{\rho} \|\varphi\|_{\mathcal{C}^{s}} \sum_{k \in \mathcal{K}_{\rho,s}} \prod_{l=1}^{\rho} \|D\psi\|_{\mathcal{C}^{l-1}}^{k_{l}}$$

where $\mathcal{K}_{\rho,s} = \{k \in \mathbb{N}_0^{\rho} : \sum_{l=1}^{\rho} k_l = s, \sum_{l=1}^{\rho} lk_l \le \rho\}.$

Using the above Lemma it follows that there exists a constant $\Lambda > 1$ such that

(5.6) $\|DF^n\|_{\mathcal{C}^r} + \|(DF^n)^{-1}\|_{\mathcal{C}^r} \le \Lambda^n, \quad \forall n \in \mathbb{N}.$

5.3. Admissible curves.

In this section we introduce the notion of *admissible curve* in order to define important auxiliary spaces and norms in the next section. We start by fixing some notations and defining exactly what we mean by *inverse branch*.

Lemma 5.3. Let γ be a differentiable closed curve in the homotopy class (0,1) such that $\gamma'(t) \notin \mathbf{C}_u$ for each $t \in \mathbb{T}$ and $F^{-1}\gamma = \bigcup_{k=1}^d \nu_k$, where the ν_k are disjoint closed curves in the homotopy class (0,1). Then, there exist open sets $\Omega_{\gamma}, \Omega_{\nu_k}$, with $\overline{\Omega}_{\gamma} = \mathbb{T}^2$, and diffeomorphisms (the inverse branches) $\mathfrak{h}_{\nu_k}: \Omega_{\gamma} \to \Omega_{\nu_k}$ satisfying,

•
$$F \circ \mathfrak{h}_{\nu_h} = Id|_{\Omega_{\alpha_h}}$$

• If $\nu_k, \nu_j \in F^{-1}\gamma$, $k \neq j$, then $\Omega_{\nu_k} \cap \Omega_{\nu_j} = \emptyset$,

•
$$\bigcup_{\nu_k \in F^{-1}\gamma} \overline{\Omega}_{\nu_k} = \mathbb{T}^2.$$

Remark 5.4. If $\gamma \in \Upsilon$, then the hypotheses of the Lemma are satisfied thanks to hypothesis (H2).

Proof of Lemma 5.3. The circle $\mathbf{q} = \{(a,0)\}_{a \in \mathbb{T}}$ intersects each ν_k in only one point $p_k = \nu_k \cap \mathbf{q}$. Indeed, by the backward invariance of the complement of \mathbf{C}_u , ν_k is locally monotone so it can meet twice \mathbf{q} only if it wraps around the torus more than once, which cannot happen since ν_k belongs to the homotopy class (0, 1). We can then label the ν_k so that the map $k \to p_k$ is orientation preserving (mod d), let us call it positively oriented.¹⁷ Also, calling $\tilde{\gamma}$ the curve obtained by translating γ by $\frac{1}{2}$ in the horizontal direction, we consider $A := F^{-1}(\tilde{\gamma}) \cap \mathbf{q}$. Since F is a local diffeomorphism, if $\tilde{p} \in A$, in a neighborhood of \tilde{p} the set $F^{-1}(\tilde{\gamma})$ consists of a curve with derivative outside \mathbf{C}_u , hence transversal to \mathbf{q} . Accordingly A is a finite collection of points. Suppose that $\tilde{p}_k \in A$ is between p_k and p_{k+1} , then $\mathbb{T}^2 \setminus \nu_k$ is a cylinder and ν_{k+1} separates the cylinder in two disjoint regions (by Jordan curve theorem), thus \tilde{p}_k belongs to a cylinder defined by the curves ν_k, ν_{k+1} . We can then follow the curve in $F^{-1}\tilde{\gamma}$ starting from \tilde{p}_k , such a curve cannot exit the cylinder (since γ and $\tilde{\gamma}$ are disjoint). If it intersects again \mathbf{q} at a point p' then the image, under F, of the segment of \mathbf{q} between \tilde{p}_k and p' is an unstable

 $^{^{15}}$ Notice that this is at odd with the usual multi-index definition in PDE, however we prefer it for homogeneity with the case, treated later, of non-commutative vector fields.

¹⁶See [6, 36] for precise, but much more cumbersome, formulae.

¹⁷This definition is ambiguous if d = 2, but in such a case the ambiguity is irrelevant.

curve that starts and ends at $\tilde{\gamma}$, hence it must cross γ , contrary to the hypothesis. It follows that $p' = \tilde{p}_k$, that is $F^{-1}\tilde{\gamma} = \bigcup_{k=1}^d \tilde{\nu}_k$, where the $\tilde{\nu}_k$ are disjoint closed curves, of homotopy type (0, 1), and $\tilde{p}_k = \tilde{\nu}_k \cap \mathfrak{q}$. As before, we can label the curves so that the \tilde{p}_k are positively oriented and so are $\tilde{p}_{k-1}, p_k, \tilde{p}_k$, where the indexes are mod d. Next, for $i \in \{1, \dots, d\}$ and $q \in \nu_i$, we define the horizontal segment $\{\xi_q(t)\}_{t \in (-\delta_-(q), \delta_+(q))}$ where $\xi_q(t) = q + e_1 t$, $\xi_q(\delta_+(q)) \in \tilde{\nu}_i$ and $\xi_q(-\delta_-(q)) \in \tilde{\nu}_{i-1}$. We then define the regions

(5.7)
$$\Omega_{\nu_i} = \bigcup_{q \in \nu_i} \xi_q.$$

Clearly, $\Omega_{\nu_i} \cap \Omega_{\nu_j} = \emptyset$ if $i \neq j$, and $\bigcup_i \overline{\Omega}_{\nu_i} = \mathbb{T}^2$. Note that $F : \Omega_{\nu_i} \cup \tilde{\nu}_{i-1} \to \mathbb{T}^2$ is a bijection, although the inverse is not continuous. However, if we restrict the map to the set Ω_{ν_i} then it is a diffeomorphism between Ω_{ν_i} and $\Omega_{\gamma} = \mathbb{T}^2 \setminus \{\tilde{\gamma}\}$. Thus it is well defined the diffeomorphism $\mathfrak{h}_{\nu_i} : \Omega_{\gamma} \to \Omega_{\nu_i}$ such that $F \circ \mathfrak{h}_{\nu_i} = Id|_{\Omega_{\gamma}}$.

From now on we call \mathfrak{h}_{ν} the inverse branch of F associated to ν and simply \mathfrak{h} when the curve ν is clear from the context. We denote by \mathfrak{H} the set of inverse branches of F. Likewise, for each $n \in \mathbb{N}$ we denote with \mathfrak{H}_n the set of inverse branches of F^n . As usual, we wish to identify the elements of \mathfrak{H}_n as compositions of elements of \mathfrak{H} . Unfortunately, Lemma 5.3 tells us that each $\mathfrak{h} \in \mathfrak{H}$ is defined on a domain obtained by removing a curve in Υ from \mathbb{T}^2 . Therefore the composition of two inverse branches in \mathfrak{H} may not be well defined. We can however consider the following sets: denoting as $\mathcal{D}_{\mathfrak{h}}$ and $\mathcal{R}_{\mathfrak{h}}$ the domain and the range of \mathfrak{h} respectively. For a curve $\gamma \in \Upsilon$ and $n \in \mathbb{N}$ we define¹⁸

(5.8)
$$\begin{aligned} \mathfrak{H}_{\gamma,n} &:= \{ \mathfrak{h} \in \mathfrak{H}_n : \mathcal{D}_{\mathfrak{h}} = \mathbb{T}^2 \setminus \{ \tilde{\gamma} \} \}, \\ \mathfrak{H}_{*,\gamma}^n &:= \left\{ \mathfrak{h}_n = (\mathfrak{h}_1^*, \cdots, \mathfrak{h}_n^*) \in \mathfrak{H}^n : \mathcal{D}_{\mathfrak{h}_j^*} \subset \mathcal{R}_{\mathfrak{h}_{j-1}^*}, j \in \{2,..,n\}, \mathcal{D}_{\mathfrak{h}_1^*} = \mathbb{T}^2 \setminus \{ \tilde{\gamma} \} \right\}. \end{aligned}$$

In $\mathfrak{H}^n_{*,\gamma}$ there exists the obvious equivalence relation $\mathfrak{h}_n \sim \mathfrak{h}'_n$ if $\mathfrak{h}^*_n \circ \cdots \circ \mathfrak{h}^*_1 = \mathfrak{h}'^*_n \circ \cdots \circ \mathfrak{h}'^*_1$ and the quotient of $\mathfrak{H}^n_{*,\gamma}$ is naturally isomorphic to $\mathfrak{H}_{\gamma,n}$. In the following we will use the two notations interchangeably. Finally, we define

$$\mathfrak{H}_{\gamma}^{\infty} = \left\{ \mathfrak{h} = (\mathfrak{h}_{1}^{*}, \cdots) \in \mathfrak{H}^{\mathbb{N}} : \mathcal{D}_{\mathfrak{h}_{j+1}^{*}} \subset \mathcal{R}_{\mathfrak{h}_{j}^{*}}, j \in \mathbb{N} ; \mathcal{D}_{\mathfrak{h}_{1}^{*}} = \mathbb{T}^{2} \setminus \{ \tilde{\gamma} \} \right\}.$$

For $\mathfrak{h} \in \mathfrak{H}^{\infty}_{\gamma}$, the symbol \mathfrak{h}_n will denote the restriction of \mathfrak{h} to $\mathfrak{H}^n_{*,\gamma}$ and we will say that $\mathfrak{h} \sim \mathfrak{h}'$ iff their restrictions are equivalent for each $n \in \mathbb{N}$.¹⁹

In the following we will often suppress the subscripts γ, ν if it does not create confusion.

5.3.1. Some further notation. For technical reasons it is convenient to work with cones which are slightly smaller than \mathbf{C}_u and \mathbf{C}_c . Take $\epsilon \in (0, 1 - \iota_{\star})$ small and,²⁰ setting $\epsilon^* = 1 - \epsilon$, let us consider the cone

(5.9)
$$\mathbf{C}_{\epsilon,u} = \{(x,y) \in \mathbb{R}^2 : |y| \le \chi_u \epsilon^* |x|\}$$

which is strictly contained in \mathbf{C}_u . In the same way it is defined $\mathbf{C}_{\epsilon,c}$. For each $p \in \mathbb{T}^2$ let $\mathfrak{H}_p^n := \{\mathfrak{h} \in \mathfrak{H}^n : p \in \mathcal{D}_{\mathfrak{h}}\}$. By the expansion of the unstable cone under backward dynamics and the backward invariance of the central cone we can define $m_{\chi_u}(p,\mathfrak{h}): \mathbb{T}^2 \times \mathfrak{H}_p^\infty \to \mathbb{N}$ and $m_{\chi_u} \in \mathbb{N}$ as

(5.10)

$$m_{\chi_{u}}(p, \mathfrak{h}) = \min\{n \in \mathbb{N} : D_{p}\mathfrak{h}_{n}(\mathbb{R}^{2} \setminus \mathbf{C}_{\epsilon, u}) \subset \mathbf{C}_{\epsilon, c}\}$$

$$m_{\chi_{u}}(p) = \inf_{\mathfrak{h} \in \mathfrak{H}^{\infty}} m_{\chi_{u}}(p, \mathfrak{h}) ; \quad m_{\chi_{u}} = \inf_{p \in \mathbb{T}^{2}} m_{\chi_{u}}(p)$$

$$m_{\chi_{u}}^{+}(p) = \sup_{\mathfrak{h} \in \mathfrak{H}^{\infty}} m_{\chi_{u}}(p, \mathfrak{h}) ; \quad m_{\chi_{u}}^{+} = \sup_{p \in \mathbb{T}^{2}} m_{\chi_{u}}^{+}(p).$$

¹⁸Here we are using the notation $\mathfrak{H}^n = \underbrace{\mathfrak{H} \times \cdots \times \mathfrak{H}}_{n\text{-times}}$ and $\tilde{\gamma} = \gamma + (1/2, 0)$.

¹⁹As it is messy to define infinite compositions, we define the equivalence relation indirectly.

²⁰During the following sections ϵ will have to satisfy different conditions. However, it is important to note that, once the conditions are satisfied, the value of ϵ is fixed and it can be considered a uniform constant. Thus for the time being we will keep track of ϵ , but we will stop when there is no danger of confusion.

Note that, by (2.5), $\mathbf{C}_{\epsilon,c} \supset D_p \mathfrak{h} \mathbf{C}_c$, for each $\mathfrak{h} \in \mathfrak{H}_p^1$. By a direct computation (see Sub-Lemma C.1 for the details) equation (5.10) implies

(5.11)
$$\lambda^{-}_{m_{\chi_u}(p,\mathfrak{h})}(p)^{-1}\mu^{m_{\chi_u}} < \epsilon^* \chi_c \chi_u, \quad \forall p \in \mathbb{T}^2, \mathfrak{h} \in \mathfrak{H}^{\infty},$$

(5.12) $c_2^{-}\log\chi_u^{-1} \le m_{\chi_u} \le m_{\chi_u}^+ < c_2^+(\log\chi_u^{-1} + 1),$

for some uniform constants $c_2^-, c_2^+ > 0$. Next, consider a vector $v = (1, u_0) \in \mathbf{C}_u$, so that $|u_0| \in [-\chi_u, \chi_u]$. By forward invariance of the unstable cone, there exist continuous functions $\Upsilon_n, \Xi_n : \mathbb{N} \times \mathbb{T}^2 \times [-\chi_u, \chi_u] \to \mathbb{R}$ such that

$$D_p F^n v = \Upsilon_n(p, u_0)(1, \Xi_n(p, u_0)),$$

where $\|\Xi_n\|_{\infty} \leq \chi_u$. We are interested in the evolution of the slope field Ξ_n . For this purpose it is convenient to introduce the dynamics $\Phi(p, u_0) = (F(p), \Xi(p, u_0))$, for $p \in \mathbb{T}^2$, $u_0 \in [-\chi_u, \chi_u]$ and where we use the notation $\Xi = \Xi_1$. The map Φ describes how the slopes of the cones change while iterating F. Note that

(5.13)
$$\Phi^n(p, u_0) = (F^n(p), \Xi_n(p, u_0))$$

Finally, for $n \in \mathbb{N}$ and $\mathfrak{h} \in \mathfrak{H}^{\infty}$, let us define the function

(5.14)
$$u_{\mathfrak{h},n}(p,u_0) = \pi_2 \circ \Phi^n(\mathfrak{h}_n(p),u_0) : \mathbb{T}^2 \times [-\chi_u,\chi_u] \to [-\chi_u,\chi_u],$$

where π_2 is the projection on the second coordinate. By Lemma F.1, applied with $u = u' = u_0$ and $\varepsilon_0 = 1$, we see that $u_{\mathfrak{h},n}(p,u_0)$ is Lipschitz and the Lipschitz constant can be computed using (F.2).

5.3.2. Admissible central and unstable curves. In the following $\pi_k : \mathbb{T}^2 \to \mathbb{T}$ will denote the projection on the k^{th} component, for k = 1, 2. Also, for $\varphi \in \mathcal{C}^r(\mathbb{T}, \mathbb{C})$ we use the notation $(\varphi)^{(j)}(t) = \frac{d^j}{dt^j}\varphi(t)$ and φ' in the case j = 1.

Definition 5.5. Let c be a positive constant, then $\Gamma_j(c)$ is the set of the C^r closed curves $\gamma : \mathbb{T} \to \mathbb{T}^2$ which are parametrized by vertical length, i.e. $\gamma(t) = (\gamma_1(t), t)$, satisfy conditions c0), c1) and c2) of assumption (H2), and:

c3) for every $2 \le \ell \le j$: $\|\gamma^{(\ell)}(t)\| \le c^{(\ell-1)!}$.

Given c > 0 and $j \leq r$ we will call $\gamma \in \Gamma_j(c)$ a (j, c)-admissible central curve (or simply admissible curve if the context is clear). We will choose c in Corollary 5.11.

Similarly, a curve $\eta \in C^r(I, \mathbb{T}^2)$ of length δ defined on a compact interval $I = [0, \delta]$ of \mathbb{T} is called an admissible unstable curve if $\eta'(t) \in \mathbf{C}_u$, it is parametrized by horizontal length and its j-derivative is bounded by $c^{(j-1)!}$.

The basic objects used in the paper are integrals along admissible (or pre-admissible) curves. To estimate precisely such objects are necessary the technical estimates developed in the next subsections.

5.4. Estimates for derivatives: SVPH case.

We start with the following simple, but very helpful, proposition.

Proposition 5.6. There exists a uniform constant $C_* \geq 1$ such that, for every $z \in \mathbb{T}^2$, any $n \in \mathbb{N}$, any vectors $v^u \in \mathbf{C}_u$ and $v^c \in \mathbf{C}_c$ such that $(a, b) := D_z F^n v^c \notin \mathbf{C}_u$, we have :

$$C_*^{-1} \frac{\|D_z F^n v^u\|}{\|v^u\|} \frac{|b|}{\|v^c\|} \le |\det D_z F^n| \le C_* \frac{\|D_z F_z^n v^u\|}{\|v^u\|} \frac{|b|}{\|v^c\|}$$

Proof. Recall that for a matrix $D \in GL(2,\mathbb{R})$ and vectors $v_1, v_2 \in \mathbb{R}^2$ linearly independent²¹

(5.15)
$$|\det D| = \frac{|Dv_1 \wedge Dv_2|}{|v_1 \wedge v_2|} = \frac{||Dv_1||}{||v_1||} \frac{||Dv_2||}{||v_2||} \frac{\sin(\measuredangle(Dv_1, Dv_2))}{\sin(\measuredangle(v_1, v_2))}$$

²¹By $\measuredangle(v, w)$ we mean the absolute value of the angle between v and w, hence it has value in $[0, \pi]$.

Let $\theta = \measuredangle (DF^n v^u, DF^n v^c)$, $\theta_1 = \measuredangle (DF^n v^u, e_1)$, $\theta_2 = \measuredangle (DF^n v^c, e_1)$ and $\theta_u = \arctan \chi_u$. Since $D_z F^n v^u \in DF\mathbf{C}_u$ we have $|\theta_1| \le c \theta_u$, for some fixed $c \in (0, 1)$. On the other hand, by hypothesis, $|\theta_2| \ge \theta_u$. Thus

$$\frac{|\theta|}{|\theta_2|} \le \frac{|\theta_2| + |\theta_1|}{|\theta_2|} \le 1 + c$$
$$\frac{|\theta|}{|\theta_2|} \ge \frac{|\theta_2| - |\theta_1|}{|\theta_2|} \ge 1 - c.$$

The Lemma follows since $||DF^nv^c|| \sin \theta_2 = b$.

Next, we introduce the following quantities for each $n \in \mathbb{N}, p \in \mathbb{T}^2$ and some constants $C_{\sharp} > 0$:

(5.16)
$$C_{\mu,n} := C_{\sharp} \frac{1-\mu^{-n}}{\mu-1} \le C_{\sharp} \min\{n, (\mu-1)^{-1}\}; \quad C_{\mu,0} = 0.$$

Remark 5.7. Note that we can always estimate $C_{\mu,n}$ by $(\mu-1)^{-1}$, which is independent on n, and we will do so if we need estimates uniform in n. Nevertheless, it can be useful to keep track of constants which deteriorate when μ approaches one (as for $C_{\mu,n}$), in view of the fast-slow case.

Next, we provide sharp estimates of various quantities relevant in the next sections.

Proposition 5.8. There exist $\bar{c}_b \geq 1$ such that, for any $n \in \mathbb{N}$ and $p \in \mathbb{T}^2$, we have:

(5.17)
$$\begin{aligned} \lambda_n^+(p) &\leq \bar{c}_{\flat} \lambda_n^-(p) \\ \|(DF^n)^{-1}\|_{\mathcal{C}^0(\mathbb{T}^2)} &\leq C_{\sharp} \mu^n. \end{aligned}$$

Proof. Let $v^c \in \mathcal{T}_{F^n(p)} \mathbb{T}^2$ with $v^c \in \mathbf{C}_c$ unitary, and $w_u \in \mathbf{C}_u$. Define

$$\tilde{w}_u = \frac{D_p F^n w_u}{\|D_p F^n w_u\|} \in \mathbf{C}_u$$

For each $v \in \mathcal{T}_{F^n(p)} \mathbb{T}^2$ we can write $v = \alpha v^c + \beta \tilde{w}_u$, then

$$||(D_{F^n p} F^n)^{-1} v|| \le |\alpha| ||(D_{F^n p} F^n)^{-1} v^c|| + |\beta| ||(D_{F^n p} F^n)^{-1} \tilde{w}_u||$$

By (2.4) and (2.6) we have the following

- (1) $||(D_{F^n p} F^n)^{-1} \tilde{w}_u|| \leq C_\star \lambda_-^{-n},$
- (2) $||(D_{F^n p} F^n)^{-1} v^c|| \le C_\star \mu^n.$

Hence,

$$||(D_{F^n p} F^n)^{-1} v|| \le C_{\star} \mu^n |\alpha| + C_{\star} \lambda_{-}^{-n} |\beta|,$$

A direct computation shows

$$\{|\alpha|, |\beta|\}^+ \leq \frac{1 + |\langle v^c, \tilde{w}_u \rangle|}{1 - \langle v^c, \tilde{w}_u \rangle^2} \|v\| \leq \frac{1 + \cos \vartheta}{1 - (\cos \vartheta)^2} \|v\|$$

where

$$\cos\vartheta := \cos\left[\inf_{v \in \mathbf{C}_u, w \in \mathbf{C}_c} \{|\measuredangle(v, w)|\}\right] \le \frac{1 + \chi_u}{\sqrt{2(1 + \chi_u^2)}} < 1.$$

From the above the second statement of (5.17) follows. The strategy for proving the first of (5.17) is similar. We take $w_1, w_2 \notin \mathbf{C}_c$ unitary and $v^c = (0, 1) \in \mathbf{C}_c$, and we set $\tilde{v}_c = \frac{(D_{F^n p} F^n)^{-1} v^c}{\|(D_{F^n p} F^n)^{-1} v^c\|} \in \mathbf{C}_c$. Notice that $\|D_p F^n \tilde{v}^c\| \leq C \mu^n$. Let $w_2 = \alpha w_1 + \beta \tilde{v}^c$. By (2.5) it follows that there exists a minimal angle between $w_1 \notin \mathbf{C}_c$ and $\tilde{v}^c \in (DF)^{-1}\mathbf{C}_c$, thus $|\alpha| + |\beta| \leq C$ for some constant $C_{\sharp} > 0$. Hence,

$$\|D_p F^n w_1 - D_p F^n w_2\| \le |1 - \alpha| \|D_p F^n w_1\| + C_{\sharp} \mu^n \le (1 + C_{\sharp}) \|D_p F^n w_1\| + C_{\sharp} \mu^n.$$

Since $||D_p F^n w_1|| \ge C \lambda_n^-(p)$, it follows that

$$\left|1 - \frac{\|D_p F^n w_2\|}{\|D_p F^n w_1\|}\right| \le \left|\frac{D_p F^n w_1}{\|D_p F^n w_1\|} - \frac{D_p F^n w_2}{\|D_p F^n w_1\|}\right| \le (1 + C_{\sharp}) + C_{\sharp} \frac{\mu^n}{\lambda_n^-(p)}$$

Equation (5.17) follows by the arbitrariness of w_1, w_2 and since $\mu < \lambda_-$.

5.5. Sharp estimates for derivatives: \mathbf{SVPH}^{\sharp} case. In this section we provide sharper estimates of the first derivatives. To do so, it is convenient to introduce the following quantities for each $n \in \mathbb{N}, m \leq n, p \in \mathbb{T}^2$ and some constants $C_{\sharp} > 0$:

(5.18)
$$\bar{\varsigma}_{n,m}(p) := \{C_{\mu,n-m}, \lambda_m^+(F^{n-m}(p))\}^+$$

(5.19)
$$\varsigma_{n,m}(p) := \{1, C_{\mu,m}(\lambda_{n-m}^{-}(p))^{-1} + C_F \bar{\varsigma}_{n,m}(p)\}^+; \quad \varsigma_{n,n} := \varsigma_n$$

(5.20)
$$C_F := \chi_u + \|\omega\|_{\mathcal{C}^r},$$

and we will use the notation $\bar{\varsigma}_{n,m} := \|\bar{\varsigma}_{n,m}\|_{\infty}$ and $\varsigma_{n,m} = \|\varsigma_{n,m}\|_{\infty}$. We have the following improvement of Proposition 5.8.

Proposition 5.9. For each c > 0, $m \le n$ and $\nu \in \Gamma_2(c)$ such that $DF^{n-m}\nu' \in \mathbf{C}_c$,²²

(5.21)
$$\|DF^{n}\|_{\mathcal{C}^{0}_{\nu}} \leq C_{\sharp}\lambda_{n}^{+} \\ \|DF^{n}\|_{\mathcal{C}^{1}_{\nu}} \leq C_{\sharp}\lambda_{n}^{+}\bar{\varsigma}_{n,m}\mu^{n-m} \\ \|DF^{n}\|_{\mathcal{C}^{2}_{\nu}} \leq C_{\sharp}(\lambda_{n}^{+}\bar{\varsigma}_{n,m}\mu^{n-m})^{2} + C_{\sharp}\lambda_{n}^{+}\bar{\varsigma}_{n,m}\mu^{n-m}\mathfrak{c} \\ \|\frac{d}{dt}(D_{\nu(t)}F^{n})^{-1}\| \leq C_{\sharp}\mu^{2n-m}\varsigma_{n,m}\circ\nu(t) \\ \|\frac{d^{2}}{dt^{2}}(D_{\nu(t)}F^{n})^{-1}\| \leq \mu^{n}C_{\mu,n}(1+C_{F}\lambda_{n}^{+})\circ\nu(t)\left\{\mu^{2n}C_{\mu,n}\lambda_{m}^{+}\circ\nu(t)+\mathfrak{c}\right\}.$$

Proof. By (2.4), we have

$$(5.22) ||D_x F^k|| \le C_{\sharp} \lambda_k^+(x).$$

Moreover, for each $n, k \in \mathbb{N}$, we have

(5.23)
$$\frac{d}{dt}D_{\nu(t)}F^{n} = \sum_{s=1}^{2}\sum_{k=0}^{n-1}D_{F^{k+1}(\nu(t))}F^{n-k-1}\partial_{x_{s}}(D_{F^{k}(\nu(t))}F)D_{\nu(t)}F^{k}(D_{\nu(t)}F^{k}\nu')_{s}$$
$$\frac{d}{dt}(D_{\nu(t)}F^{n})^{-1} = \sum_{s=1}^{2}\sum_{k=0}^{n-1}(D_{\nu(t)}F^{k})^{-1}\left[\partial_{x_{s}}(DF)^{-1}(D_{F(\cdot)}F^{n-k-1})^{-1}\right] \circ F^{k}(\nu(t))$$
$$\cdot (D_{\nu(t)}F^{k}\nu')_{s}.$$

The above, also differentiating once more, implies that, for $k \leq n$,

(5.24)
$$\begin{aligned} \|\frac{d}{dt}(D_{\nu(t)}F^{k})\| &\leq \sum_{j=0}^{\kappa-1} \lambda_{k-j-1}^{+} \lambda_{k}^{+} \mu^{\{j,n-m\}^{-}} \lambda_{\{j-n+m,0\}^{+}}^{+} \\ &\leq C_{\sharp} \lambda_{k}^{+} \mu^{n-m} \{C_{\mu,n-m}, \lambda_{m}^{+}\}^{+} = C_{\sharp} \lambda_{k}^{+} \bar{\varsigma}_{n,m} \mu^{n-m}, \\ \|\frac{d^{2}}{dt^{2}} (D_{\nu(t)}F^{n})\| &= \|\sum_{\ell,s} (\partial_{x_{\ell}} \partial_{x_{s}} D_{x}F^{n}) \nu_{\ell}^{\prime} \nu_{s}^{\prime} + \sum_{s} \partial_{x_{s}} D_{x}F^{n} \nu_{s}^{\prime\prime} \| \\ &\leq C_{\sharp} (\lambda_{n}^{+} \mu^{n-m} \bar{\varsigma}_{n,m})^{2} + C_{\sharp} \lambda_{n}^{+} \bar{\varsigma}_{n,m} \mu^{n-m} \mathfrak{c}. \end{aligned}$$

1. 1

To estimate the second of (5.23), note that there exist $\xi, \tilde{\xi} \in \mathcal{C}^{r-1}(\mathbb{T}^2, \mathbb{R}^2)$, $\|\xi\|_{\mathcal{C}^{r-1}} \leq C_{\sharp}$, such that, for all $w \in \mathbb{R}^2$, and $0 < |\alpha| \leq r-1$,

(5.25)
$$\begin{aligned} \left\|\partial^{\alpha}(DF)^{-1}w - e_{1}\langle\partial^{\alpha}\xi, w\rangle\right\| &\leq C_{\sharp}\|w\|\|\omega\|_{\mathcal{C}^{|\alpha|+1}}\\ \left\|\partial^{\alpha}(DF)w - e_{1}\langle\partial^{\alpha}\tilde{\xi}, w\rangle\right\| &\leq C_{\sharp}\|w\|\|\omega\|_{\mathcal{C}^{|\alpha|+1}}. \end{aligned}$$

 $^{^{22}\}text{Recall Section 5.2}$ for the definition of $\|\cdot\|_{\mathcal{C}^r_\nu}$ and (5.19) for the notations used.

Thus, setting $\eta_k(p) = D_p F^k e_1 || D_p F^k e_1 ||^{-1}$, we have $||\eta_k - e_1|| \le C_\sharp \chi_u$ and, for all $w \in \mathbb{R}^2$,

(5.26)
$$\begin{aligned} \left\| (D_x F^k)^{-1} \partial_{x_i} (D_p F)^{-1} w \right\| &\leq \left\| (D_x F^k)^{-1} \eta_k(x) \langle \partial_{x_i} \xi, w \rangle \right\| \\ &+ \left\| (D_x F^k)^{-1} \left[\partial_{x_i} (D_p F)^{-1} w - \eta_k(x) \langle \partial_{x_i} \xi, w \rangle \right] \right\| \\ &\leq C_{\sharp} \frac{\|w\|}{\lambda_k^-(x)} + C_{\sharp} \mu^k \|w\| (\chi_u + \|\omega\|_{\mathcal{C}^2}), \end{aligned}$$

where $F^k(x) = p$. Hence, recalling (5.20) and using the above and (5.17), we have, for all $k \leq n$

(5.27)
$$\begin{aligned} \|\frac{d}{dt}((D_{\nu}F^{k})^{-1})\| &\leq C_{\sharp} \sum_{j=0}^{\{k,n-m\}^{-}-1} \mu^{k-1} \{(\lambda_{j}^{-} \circ \nu)^{-1} + C_{F}\mu^{j}\} \\ &+ C_{\sharp} \sum_{j=n-m}^{k-1} \mu^{k-j-1} \{(\lambda_{j}^{-} \circ \nu)^{-1} + C_{F}\mu^{j}\} \mu^{n-m} \lambda_{m-n+j}^{+} \circ F^{n-m} \circ \nu \\ &\leq C_{\sharp} \mu^{k+n-m} \left[1 + C_{\mu,m} (\lambda_{n-m}^{-} \circ \nu)^{-1} + C_{F} \{C_{\mu,n-m}, \lambda_{k-n+m}^{+} \circ F^{n-m} \circ \nu\}^{+}\right]. \end{aligned}$$

Therefore we have

(5.28)
$$\|\frac{d}{dt}((D_{\nu(t)}F^n)^{-1})\| \le C_{\sharp}\mu^{2n-m}\varsigma_{n,m}\circ\nu(t),$$

which yields the statement for the first derivative. Next, differentiating once more the second of (5.23),

$$\begin{aligned} \frac{d^2}{dt^2} (D_{\nu}F^n)^{-1} &= \sum_{s=1}^2 \sum_{k=0}^{n-1} \left[\frac{d}{dt} (D_{\nu(t)}F^k)^{-1} \right] \left[\partial_{x_s} (DF)^{-1} (D_{F(\cdot)}F^{n-k-1})^{-1} \right] \circ F^k(\nu) \\ &\cdot (D_{\nu(t)}F^k\nu')_s + \sum_{s,\ell=1}^2 \sum_{k=0}^{n-1} (D_{\nu}F^k)^{-1} \left\{ \partial_{x_\ell} \left[\partial_{x_s} (DF)^{-1} (D_{F(\cdot)}F^{n-k-1})^{-1} \right] \right\} \circ F^k(\nu) \\ &\cdot (D_{\nu}F^k\nu')_\ell (D_{\nu}F^k\nu')_s + \sum_{s=1}^2 \sum_{k=0}^{n-1} (D_{\nu}F^k)^{-1} \left[\partial_{x_s} (DF)^{-1} (D_{F(\cdot)}F^{n-k-1})^{-1} \right] \circ F^k(\nu) \\ &\cdot \left\{ \left[\frac{d}{dt} D_{\nu}F^k \right] \nu' + D_{\nu}F^k\nu'' \right\}. \end{aligned}$$

We estimate the three sums above separately. By (5.26) and (5.28), the first one is bounded by

$$C_{\sharp} \sum_{k=0}^{n-m-1} \mu^{k+2n-m-1} \varsigma_{n,m} \circ \nu + C_{\sharp} \sum_{k=n-m}^{n-1} \mu^{n+k-m} \mu^{k} \varsigma_{n,m} \circ \nu \mu^{n-k-1} \mu^{n-m} \lambda_{m-n+k}^{+} \circ F^{n-m} \nu$$

$$\leq C_{\sharp} \mu^{3n-2m} (\varsigma_{n,m} \bar{\varsigma}_{n,m}) \circ \nu.$$

The second one is equal to

$$\sum_{k=0}^{n-1} (D_{\nu}F^{k})^{-1} \left\{ \partial_{x_{\ell},x_{s}}^{2} (DF)^{-1} \cdot (D_{F(\cdot)}F^{n-k-1})^{-1} \right\} \circ F^{k}(\nu) \cdot (D_{\nu}F^{k}\nu')_{\ell} (D_{\nu}F^{k}\nu')_{s} + (D_{\nu}F^{k})^{-1} \left\{ \partial_{x_{s}} (DF)^{-1} \partial_{x_{\ell}} (D_{F(\cdot)}F^{n-k-1})^{-1} \right\} \circ F^{k}(\nu) \cdot (D_{\nu}F^{k}\nu')_{\ell} (D_{\nu}F^{k}\nu')_{s},$$

so we can use (5.23), (5.25), (5.26) to get the bound

$$C_{\sharp} \sum_{k=0}^{n-1} \left[(\lambda_{k}^{-} \circ \nu(t))^{-1} + C_{F} \mu^{k} \right] \mu^{n-k} \left\{ 1 + \left[C_{\mu,n-k} \mu^{n-k} + C_{F} \lambda_{n-k}^{+} \circ \nu \right] \right\} \mu^{2\{k,n-m\}^{-1}} \cdot (\lambda_{\{0,k-n+m\}^{+}}^{+} \circ F^{n-m} \circ \nu(t))^{2} \leq \mu^{5n-2m} C_{\mu,n}^{2} (1 + C_{F} \lambda_{m}^{+})^{2}.$$

For the last term we estimate as above and, recalling (5.24), we obtain the bound

$$\mu^n C_{\mu,n} \left\{ 1 + C_F \lambda_n^+(\nu(t)) \right\} (\bar{\varsigma}_{n,m} \mu^{n-m} + \mathfrak{c}).$$

Collecting the above estimates, the last of the (5.21) readily follows.

5.6. Iteration of curves: SVPH case.

A key point in the following arguments is to check how the central admissible curves behave under iteration. As they are just merely technical, we postpone the proofs, except for Corollary 5.11 and 5.15, to Appendix C to make the reading more fluent.

Lemma 5.10. Let F be SVPH. There exist uniform constants $\bar{n} \in \mathbb{N}$, $^{23}C_{\flat} > 1$ and $\eta < 1$ such that, if $\mathfrak{c} > \frac{9}{4}C_{\flat}^{3}\mu^{3\bar{n}}$, for each $c_{\star} > \mathfrak{c}/2$, $\gamma \in \Gamma_{\ell}(c_{\star})$, $1 \leq \ell \leq r$, and $n \geq \bar{n}$, setting $\nu_{n} \in F^{-n}\gamma$, there exist diffeomorphisms $h_{n,\nu} =: h_{n} \in \mathcal{C}^{r}(\mathbb{T})$ such that:

The curve $\hat{\nu}_n = \nu_n \circ h_n$ is in $\Gamma_\ell(\eta^n c_\star + c/2)$ and

(5.29)
$$\|h_n\|_{\mathcal{C}^{\ell}} \leq \begin{cases} C_{\flat}\mu^n & \text{if } \ell = 1\\ C_{\flat}^4(c_{\star}C_F + 1)C_{\mu,n}\mu^{2n} & \text{if } \ell = 2\\ C_{\flat}^{2\ell!} \left(c_{\star}^{\ell!}C_F + 1\right)C_{\mu,n}^{a_{\ell}}\mu^{\ell!n} & \text{if } \ell > 2 \end{cases}$$

where $a_{\ell} = (\ell - 1)! \sum_{k=0}^{\ell-1} \frac{1}{k!}$, and $C_{\mu,n}$ as in (5.16).

Lemma 5.10 implies immediately the following key result.

Corollary 5.11. Let $c = \chi_u^{-\varpi_{\bar{n},\chi_u}}$, for some $\varpi_{\bar{n},\chi_u} \ge 1$. Under the hypothesis of Lemma 5.10, for each n such that $\eta^n \le \frac{1}{2}$, we have the inclusion $F^{-n}\Gamma_\ell(c) \subset \Gamma_\ell(c)$.

Proof. The statement follows by Lemma 5.10 choosing $\varpi_{\bar{n},\chi_u}$ and c_{\star} such that

(5.30)
$$c_{\star} = \mathfrak{c} = \chi_{u}^{-\varpi_{\bar{n},\chi_{u}}} = \frac{9}{2} C_{\flat}^{3} \mu^{3\bar{n}}$$

The result then follows since $\eta^n \leq \frac{1}{2}$.

Remark 5.12. From now on we will use Γ to denote $\Gamma_r(\mathfrak{c})$ where \mathfrak{c} is defined in Corollary 5.11 and has thus the stated invariance property. As it clear from the proof of the above Corollary, the hypothesis of Lemma 5.10 suffice to guarantee the backward invariance of the central curves for a SVPH system with the choice $\mathfrak{c} = \chi_u^{-\varpi_{\bar{n},\chi_u}}$.

The above results tell us that the space of admissible central curves is stable under backward iteration of the map. Arguing as above, but forward in time (see [50, Lemma 3.2]), it can be proven that the space of admissible unstable curves is stable under the iteration of F^n , for n greater than \bar{n} . In particular, if $\eta : I \to \mathbb{T}^2$ is an admissible unstable curve, and η_n is the image of η under F^n , then there exists a diffeomorphism $p_{n,\eta} =: p_n$ such that

(5.31)
$$p'_{n}(t) = \frac{\|D_{\eta}F^{n} \cdot \eta'(t)\|}{\|\eta'(t)\|}$$

and $\eta_n \circ p_n = F^n \circ \eta$ is an admissible unstable curve. Moreover, as F acts as an expanding map along those curves, we have the following standard distortion estimate for each $n \ge 1$:

(5.32)
$$\frac{p'_n(t)}{p'_n(s)} \le C_{\sharp}, \quad \forall t, s \in I.$$

We will need to control the backward evolution also of curves not in the center cone. The last result of this section is the following Lemma, which proof can be found in Appendix C, Section C.2.

18

²³The constant \bar{n} is chosen in (C.10).

Lemma 5.13. Let F be a SVPH and $\Delta_{\gamma} \in L^{\infty}(\mathbb{T}, [1, +\infty])$ and consider any closed curve $\gamma \in C^r$, homotopic to (0, 1), such that $\|\gamma'(t)\| = 1$ and $\|\gamma^{(j+1)}(t)\| \leq \Delta_{\gamma}(t)^j$,²⁴ for all $j \in \{1, \ldots, r\}$ and $t \in \mathbb{T}$. For $\mathfrak{h} \in \mathfrak{H}^{\infty}$ let $n_0 \geq 0$ and m be the smallest integers such that, for all $t \in \mathbb{T}$,

(5.33)
$$D_{\gamma(t)}\mathfrak{h}_{n_0}\gamma'(t)\notin \mathbf{C}_u \quad and \quad D_{\gamma(t)}\mathfrak{h}_m\gamma'(t)\in \mathrm{Int}\left(\mathbf{C}_c\right),$$

and assume that $m > \{\bar{n}, n_0\}^+$.²⁵ Let $\nu_k = \mathfrak{h}_k(\gamma), k \in \mathbb{N}$. For $k \ge n_0$ then there exists a reparametrization h_k such that, setting $\hat{\nu}_k = \nu_k \circ h_k, \pi_2 \circ \hat{\nu}_k(t) = t$.

If, for some \mathfrak{h} , we have $m < \infty$, then:

For $\eta < 1$ given in Lemma 5.10,²⁶ Λ as in (5.6), $\overline{m} = \sigma m$, where

(5.34)
$$\sigma = \left\{ 1, \left\lceil \frac{\ln(\mu \| \Delta_{\gamma} \|_{\infty} \Lambda)}{\ln \eta^{-1}} \right\rceil \right\}^{+},$$

and c as in Corollary 5.11, we have $\hat{\nu}_{\overline{m}} \in \Gamma_j(c)$ for each $j \ge 1$, and the \mathcal{C}^j -norm of $h_{\overline{m}}$ satisfies (5.29) with $c_\star = \chi_u^{-2} \|\Delta_\gamma\|_{\infty} (\mu \Lambda)^m$.

5.7. Iteration of curves: the SVPH^{\sharp} case. In this sub-section we provide an improvement of Lemma 5.13, for lower derivatives, in the SVPH^{\sharp} case. Lemma 5.14 deals with the derivatives of $\hat{\nu}_n$, while Lemma 5.16 bounds the derivative of the reparametrization.

Lemma 5.14. In the hypothesis of Lemma 5.13, let F be a $SVPH^{\sharp}$ as in Definition 5.1 where²⁷

(5.35)
$$B_{\hat{n}} := \bar{c}_{\flat}^3 C_{\flat}^9 (1 + \chi_c^2)^{\frac{3}{2}} (1 + 6\varsigma_{\hat{n}} C_{\mu,\hat{n}}).$$

Then there exist $C_3, C_4 \ge 1$ uniform, such that for all $\gamma \in \Gamma_\ell(c_\star)$, $c_\star > 0$, and $n_\star \in \{\hat{n}, \ldots, c_2^{-} \ln \chi_u^{-1}\}$, setting $c_{n_\star} = c_\flat^{1/n_\star}$, $c_\flat = \bar{c}_\flat a_{n_\star}^{n_\star}$, $a_{n_\star} = ((1 + \chi_c^2)^{1/2} C_\flat^3)^{n_\star^{-1}}$,²⁸

(5.36)
$$b_{n_{\star}} := (C_4 C_{\flat}^5 \varsigma_{n_{\star}} C_{\mu,n_{\star}})^{\frac{1}{n_{\star}}}$$
$$\overline{\mathbf{s}}_{n_{\star}} = C_{\flat} \mu^{4n_{\star}} \varsigma_{n_{\star}}^2 C_{\mu,n_{\star}} + C_{\flat} \varsigma_{n_{\star}} C_{\mu,n_{\star}} \mu^{5n_{\star}} C_3 + C_{\mu,n_{\star}}^2 \mu^{6n_{\star}} C_3^2$$
$$\mathbf{s}_{n_{\star}} = \overline{\mathbf{s}}_{n_{\star}} + 6\varsigma_{n_{\star}} C_{\mu,n_{\star}}^3 \mu^{6n_{\star}} C_3^2 + b_{n_{\star}}^{2n_{\star}} C_{\mu,n_{\star}} \mu^{6n_{\star}} C_3,$$

we have, for all $n \in \mathbb{N}$,

(5.37)
$$\begin{aligned} \|\hat{\nu}_{n}''(t)\| &\leq c_{\flat}c_{n_{\star}}^{n}\mu^{2n}\lambda_{n}^{-}(\gamma\circ h_{n}(t))^{-1}c_{\star} + C_{\mu,n_{\star}}\mu^{3n_{\star}}C_{3} \\ \|\hat{\nu}_{n}'''\| &\leq c_{\flat}^{2}(1+6\varsigma_{n_{\star}}C_{\mu,n_{\star}})c_{n_{\star}}^{n}\mu^{3n}(\lambda_{n}^{-}(\gamma\circ h_{n}(t)))^{-1}c_{\star}^{2} \\ &+ c_{\flat}^{2}b_{n_{\star}}^{n,\star}\mu^{2n+2n_{\star}}\lambda_{n}^{-}(\gamma\circ h_{n}(t))^{-1}c_{\star} + \mathbf{s}_{n_{\star}}. \end{aligned}$$

The proof of Lemma 5.14 can be found in section C.3 of Appendix C. The above Lemma implies the following sharper version of Corollary 5.11.

Corollary 5.15. If $\ell \in \{2,3\}$ then, in the hypothesis of Lemma 5.14, for all $n \geq \hat{n}$, we have $F^{-n}\Gamma_{\ell}(\mathfrak{c}) \subset \Gamma_{\ell}(\mathfrak{c})$ with the sharper choice

²⁴We will apply this Lemma with $\Delta_{\gamma}(t)$ given by (G.1).

 $^{^{25}\}mathrm{The}$ other possibilities are already covered by Corollary 5.11.

²⁶See (C.10) for a precise definition of η .

²⁷Recall that condition (5.1) can be satisfied thanks to hypothesis (H3). Also note that for the present Lemma μ^3 would suffice, however we will need μ^{30} in (8.64).

²⁸Recall (5.19) for the definition of ς_n while \bar{c}_{\flat} is defined in Proposition 5.9.

Proof. We use Lemma 5.14. By (5.37) and (5.1) we have, for each $n \ge \hat{n}$,

$$\begin{split} \|\hat{\nu}_n''(t)\| &\leq \frac{1}{2}\mathfrak{c} + C_{\mu,n_\star}\mu^{3n_\star}C_3 \leq \mathfrak{c} \\ \|\hat{\nu}_n'''\| &\leq \frac{1}{2}\mathfrak{c}^2 + c_\flat^2 b_{n_\star}^n \mu^{4n} (\lambda_n^-(\gamma \circ h_n))^{-1}\mathfrak{c} + \mathfrak{s}_{n_\star} \\ &\leq \frac{1}{2}\mathfrak{c}^2 + \frac{\sqrt{\mathfrak{s}_{n_\star}}}{2}\mathfrak{c} + \mathfrak{s}_{n_\star} \leq \mathfrak{c}^2, \end{split}$$

where in the last line we used our choice of c.

To improve Lemma 5.13 and control the backward evolution of curves not in the center cone, we introduce a further quantity. Given a smooth curve γ such that $\pi_1 \circ \gamma'(t) \neq 0$ for each $t \in \mathbb{T}$, let

(5.39)
$$\vartheta_{\gamma}(t) = \left\{ \frac{|\pi_{2} \circ \gamma'(t)|}{|\pi_{1} \circ \gamma'(t)|}, \chi_{u} \right\}^{+}$$
$$\vartheta_{\gamma} = \inf_{t} \{\vartheta_{\gamma}(t)\}.$$

The last result of this section is the next Lemma, which proof is in Appendix C, section C.4.

Lemma 5.16. In the case $j \in \{1, 2\}$ we have the following sharper version of Lemma 5.13: If F is a SVPH^{\$\$\$}, for each $p \in \gamma$ and $n_{\star} \in \{\hat{n}, \dots, c_2^{-} \log \chi_u^{-1}\}$, let $\overline{m}(p, \mathfrak{h}, n_{\star}) \equiv \overline{m}$ be the minimum integer such that

$$(5.40) \qquad c_{\flat}c_{n_{\star}}^{\overline{m}}\mu^{2\overline{m}}\lambda_{\overline{m}-m}^{+}(\hat{\nu}_{m}\circ h_{\overline{m}-m}(t))^{-1}M_{m,n_{0}}(t) \leq C_{\mu,n_{\star}}\mu^{3n_{\star}}C_{3},$$

$$(5.40) \qquad c_{\flat}^{2}b_{n_{\star}}^{n_{\star}}\mu^{2\overline{m}+2n_{\star}}\lambda_{\overline{m}-m}^{-}(\gamma\circ h_{\overline{m}-m}(t))^{-1}M_{m,n_{0}}(t) \leq \frac{1}{2}\mathfrak{s}_{n_{\star}}$$

$$c_{\flat}^{2}(1+6\varsigma_{n_{\star}}C_{\mu,n_{\star}})c_{n_{\star}}^{\overline{m}}\mu^{3\overline{m}}(\lambda_{\overline{m}-m}^{-}(\gamma\circ h_{\overline{m}-m}(t)))^{-1}\overline{M}_{m,n_{0}}(t) \leq \frac{1}{2}\mathfrak{s}_{n_{\star}},$$

where

$$M_{m,n_0}(t) := C_{\sharp} \left[\Lambda^{2n_0} \mu^{2m} \Delta_{\gamma} \circ h_{n_0}(t) + C_{\mu,m} \mu^{3m} \vartheta_{\hat{\nu}_{n_0}}^{-1} \right] \left\{ 1 + \mu^{2m} \vartheta_{\hat{\nu}_{n_0}}^{-1} C_F \right\}$$

(5.41) $\overline{M}_{m,n_0}(t) := \mu^{8m} C^2_{\mu,m} (1 + C_F \mu^m \vartheta^{-1}_{\hat{\nu}_{n_0}})$

$$\times \left[1 + \Lambda^{2n_0} \Delta_{\gamma} \circ h_{n_0} \vartheta_{\hat{\nu}_{n_0}} + \Lambda^{3n_0} (\Delta_{\gamma} \circ h_{n_0})^2 \vartheta_{\hat{\nu}_{n_0}}\right] \vartheta_{\hat{\nu}_{n_0}}^{-2}$$

and $a_{n_{\star}}, b_{n_{\star}}, c_{n_{\star}}$, $\mathbf{s}_{n_{\star}}$ are defined in Lemma 5.14. Then $\hat{\nu}_{\overline{m}} \in \Gamma_3(\mathbf{c})$ and, setting $\bar{h}_{k-n_0} = h_{n_0}^{-1} \circ h_k$,

(5.42)
$$C_{\sharp}\Lambda^{-n_{0}}\vartheta_{\hat{\nu}_{n_{0}}}(\bar{h}_{\overline{m}-n_{0}}(t))^{-1}\mu^{-\overline{m}+n_{0}} \leq |h'_{\overline{m}}(t)| \leq C_{\sharp}\Lambda^{n_{0}}\vartheta_{\hat{\nu}_{n_{0}}}(\bar{h}_{\overline{m}-n_{0}}(t))^{-1}\mu^{\overline{m}-n_{0}} \\ |h''_{\overline{m}}| \leq C_{\sharp}\mu^{3\overline{m}}\vartheta_{\hat{\nu}_{n_{0}}}^{-1}\Lambda^{3n_{0}}\left\{\Delta_{\gamma}\vartheta_{\hat{\nu}_{n_{0}}}^{-1} + M_{m,n_{0}}(C_{F}C_{\mu,\overline{m}}+1) + C_{\mu,\overline{m}}\right\}.$$

5.8. Distortion: the SVPH case.

We conclude this section with some technical distortion results needed in the following.

Lemma 5.17. For all $n \in \mathbb{N}$, $\nu \in F^{-n}(\Gamma(\mathfrak{c}))$ and $x, y \in \nu$, we have

(5.43)
$$e^{-\mu^n C_{\mu,n} \|x-y\|} \le \frac{\lambda_n^+(x)}{\lambda_n^+(y)} \le e^{\mu^n C_{\mu,n} \|x-y\|}.$$

Proof. We prove it by induction. To start with, let $x = \nu(t_1), y = \nu(t_2)$ such that $||x - y|| \leq \tau_n$ for some τ_n to be chosen shortly. For n = 1 we have, for all unit vector $v \notin \mathbf{C}_c$,

$$\frac{\|D_x Fv\|}{\|D_y Fv\|} \le e^{\ln\left[1 + \frac{\|D_x Fv - D_y Fv\|}{\|D_y Fv\|}\right]} \le e^{\frac{\|D_x Fv - D_y Fv\|}{\|D_y Fv\|}}.$$

(5.44)
$$\|D_x Fv - D_y Fv\| \le \int_{t_1}^{t_2} \|\frac{d}{ds} D_{\nu(s)} Fv\| ds \le C_{\sharp} |t_2 - t_1| \le C_{\sharp} \|x - y\|,$$

the case n = 1 follows. Assume it is true for each k < n, then, by the triangular inequality

$$\begin{aligned} \|D_{y}F^{n}v - D_{x}F^{n}v\| &\leq \sum_{k=0}^{n-1} \|D_{F^{k+1}y}F^{n-k-1}(D_{F^{k}y}F - D_{F^{k}x}F)D_{x}F^{k}v\| \\ &\leq C_{\sharp}\sum_{k=0}^{n-1}\lambda_{n-k-1}^{+}(F^{k}y)\lambda_{k}^{+}(x)\|D_{F^{k}y}F - D_{F^{k}x}F\| \leq C_{\sharp}\sum_{k=0}^{n-1}\lambda_{n-k-1}^{+}(F^{k}y)\lambda_{k}^{+}(x)\mu^{k}\|x-y\| \end{aligned}$$

Since $\nu \in F^{-n}(\Gamma(c))$, $||D_{F^k y}F - D_{F^k x}F|| \leq C_{\sharp}\mu^k ||x - y||$. Also remark that (5.17) and the induction hypothesis imply

$$\lambda_{n-k}^{+}(F^{k}y)\lambda_{k}^{+}(x) \le e^{\mu^{k}C_{\mu,k}\|x-y\|}\lambda_{n-k}^{+}(F^{k}y)\lambda_{k}^{+}(y) \le C_{\sharp}\lambda_{n}^{-}(y),$$

provided we have chosen τ_n small enough. Accordingly, since $||D_y F^n v|| \ge \lambda_n^-(y)$,

$$\frac{\|D_x F^n v\|}{\|D_y F^n v\|} \le e^{\frac{\|D_x F^n v - D_y F^n v\|}{\|D_y F^n v\|}} \le e^{C_{\sharp} \sum_{k=0}^{n-1} \mu^k \|x - y\|}.$$

We can now choose v such that $||D_x F^n v|| = \lambda_n^+(x)$ so

$$\frac{\lambda_n^+(x)}{\lambda_n^+(y)} \le \frac{\|D_x F^n v\|}{\|D_y F^n v\|} \le e^{C_{\sharp} \sum_{k=0}^{n-1} \mu^k \|x-y\|} \le e^{C_{\mu,n} \mu^n \|x-y\|},$$

which proves the upper bound, for points close enough. Next, for all $x, y \in \nu$ we can consider close intermediate points $\{x_i\}_{i=0}^l$, $x_0 = x$, $x_l = y$, to which the above applies, hence

$$\frac{\lambda_n^+(x)}{\lambda_n^+(y)} \le \frac{\|D_x F^n v\|}{\|D_y F^n v\|} = \prod_{i=0}^{l-1} \frac{\|D_{x_i} F^n v\|}{\|D_{x_{i+1}} F^n v\|} \le e^{\mu^n C_{\mu,n} \sum_{i=0}^{l-1} \|x_{i+1} - x_i\|}.$$

Taking the limit for $l \to \infty$ we have the distance, along the curve, between x and y which is bounded by $C_{\sharp} ||x - y||$. This proves the upper bound. The lower bound is proven similarly.

Next, we prove other two distortion Lemmata, inspired by Lemma 6.2 in [32]. Even though the basic idea of the proof is the same, the presence of the central direction creates some difficulties.

Lemma 5.18. For each $c_{\star} \geq c$, $\gamma \in \Gamma(c_{\star})$, $n > \overline{n}$ and $0 \leq \rho \leq r - 1$, we have

(5.45)
$$\sum_{\nu_{n}\in F^{-n}\gamma} \left\| \frac{h'_{n}}{\det D_{\hat{\nu}_{n}}F^{n}} \right\|_{\mathcal{C}^{\rho}(\mathbb{T})} \leq C_{\sharp}c_{\star}^{\rho!-1}(1+C_{F}c_{\star}^{\beta_{\rho}})^{\{\rho+1,(\rho+1)\rho/2\}^{+}}C_{\mu,n}^{\tilde{a}_{\rho}}\mu^{\tilde{b}_{\rho}n} \\ \sum_{\nu_{n}\in F^{-n}\gamma} \left\| \frac{1}{\det D_{\hat{\nu}_{n}}F^{n}} \right\|_{\mathcal{C}^{\rho}(\mathbb{T})} \leq C_{\sharp}c_{\star}^{\rho!-1}(1+C_{F}c_{\star}^{\beta_{\rho}})^{\{\rho+1,(\rho+1)\rho/2\}^{+}}C_{\mu,n}^{\tilde{a}_{\rho}}\mu^{(\tilde{b}_{\rho}+1)n}$$

where $\beta_{\rho} = 1$ for $\rho \leq 2$ and $\beta_{\rho} = \rho!$ otherwise; $\tilde{a}_{\rho} = \rho + 1$, $\tilde{b}_{\rho} = \rho! - 1$, for $\rho \leq 2$, and $\tilde{a}_{\rho} = a_{\rho}\rho(\rho+1)/2 + 1$, $\tilde{b}_{\rho} = \rho!\rho(\rho+1)/2 + 1$, otherwise.

Proof. For every $\nu \in F^{-n}\gamma$ define

$$\Psi_{\nu_n}(t) = \frac{h'_n(t)}{\det D_{\hat{\nu}_n(t)}F^n}$$

and recall that in dimension one holds $\|\Psi_{\nu_n}\|_{\mathcal{C}^0} \leq \|\Psi_{\nu_n}\|_{L^1} + \|\Psi'_{\nu_n}\|_{L^1}$. We then first look for a bound of the $W^{1,1}(\mathbb{T})$ -norm of Ψ_{ν_n} . Since $e_1 = (1,0) \in \mathbb{C}_u$, $D_{\hat{\nu}_n} F^n \hat{\nu}'_n \notin \mathbb{C}_u$ and recalling that $F^n \hat{\nu}_n = \gamma \circ h_n$, we have

$$h'_n D_{\hat{\nu}_n} F^n e_1 \wedge \gamma' \circ h_n = D_{\hat{\nu}_n} F^n e_1 \wedge D_{\hat{\nu}_n} F^n \hat{\nu}'_n = \det(D_{\hat{\nu}_n} F^n) e_1 \wedge \hat{\nu}'_n.$$

Thus we have the equation

(5.46)
$$\frac{|h'_n(t)|}{|\det D_{\hat{\nu}_n(t)}F^n|} = \frac{\|e_1 \wedge \hat{\nu}'_n(t)\|}{\|D_{\hat{\nu}_n(t)}F^n e_1 \wedge \gamma' \circ h_n(t)\|}$$

 $^{^{29}\}text{Recall}$ the definition of a_{ρ} in Lemma 5.10

Since $\|\gamma'\| \ge 1$ we have, recalling definition (5.39),

(5.47)
$$\|D_{\hat{\nu}_n} F^n e_1 \wedge \gamma' \circ h_n\| \ge C_{\sharp} \vartheta_{\gamma} \circ h_n \|D_{\hat{\nu}_n} F^n e_1\|$$

Therefore, since $\|\hat{\nu}'_n\|^2 \leq 1 + \chi_c^2$, we have

(5.48)
$$\sum_{\nu_n \in F^{-n}\gamma} \|\Psi_{\nu_n}\|_{L^1} \lesssim \sum_{\nu_n \in F^{-n}\gamma} \left\|\frac{1}{\vartheta_{\gamma} \circ h_n \|D_{\hat{\nu}_n} F^n \cdot e_1\|}\right\|_{L^1}.$$

Recall that, by Lemma 5.3, for each $\hat{\nu}_n$ we have an inverse branch $\mathfrak{h}_{\hat{\nu}_n} : \Omega_{\gamma} \to \Omega_{\hat{\nu}_n}$ such that $F^n \circ \mathfrak{h}_{\hat{\nu}_n} = Id_{\Omega_{\gamma}}$. Note that the domain $\Omega_{\hat{\nu}_n}$ can be written as $\bigcup_{t \in \mathbb{T}} \xi_{t,\hat{\nu}_n}$, where $\xi_{t,\hat{\nu}_n}(s) = \hat{\nu}_n(t) + se_1$ are horizontal segments defined on an interval I_t of length $\delta_{\hat{\nu}_n(t)}$ whose images are unstable curves $\xi^{\sharp}_{t,\gamma}$ with length $(\xi^{\sharp}_{t,\gamma}) = \delta^{\sharp}_{t,\gamma} \geq 1$. Let $p_{n,\xi_{t,\nu_n}}$ be the diffeomorphism associated to ξ_{t,ν_n} , see formula (5.31). By equation (5.32) $p'_{n,\xi_{t,\nu_n}}(s) \leq p'_{n,\xi_{t,\nu_n}}(0) = \|D_{\hat{\nu}_n(t)}F^ne_1\|$. It follows

$$1 \le \delta_{t,\gamma}^{\sharp} = \int_{I_t} \left\| \frac{d}{ds} F^n(\xi_{t,\hat{\nu}_n}(s)) \right\| ds \le C_{\sharp} \delta_{\hat{\nu}_n(t)} p'_{n,\xi_{t,\nu_n}}(0) = C_{\sharp} \delta_{\hat{\nu}_n(t)} \| D_{\hat{\nu}_n(t)} F^n e_1 \|,$$

from which

(5.49)
$$\|D_{\hat{\nu}_n(t)}F^n e_1\| \gtrsim \frac{1}{\delta_{\hat{\nu}_n(t)}}$$

Since by Lemma 5.3 the Ω_{ν_n} are all disjoints and the ν_n are parametrized vertically, by (5.49) we have³⁰

(5.50)
$$\sum_{\nu_n \in F^n \gamma} \left\| \frac{1}{\|D_{\hat{\nu}_n(t)} F^n e_1\|} \right\|_{L^1} \lesssim \sum_{\hat{\nu}_n \in F^n \gamma} \int_{\mathbb{T}} \delta_{\hat{\nu}_n(t)} = \sum_{\nu_n \in F^n \gamma} m(\Omega_{\hat{\nu}_n}) = m(\mathbb{T}^2) = 1.$$

Using this in (5.48) yields

(5.51)
$$\sum_{\nu_n \in F^n \gamma} \|\Psi_{\nu_n}\|_{L^1} \le C_{\sharp} \vartheta_{\gamma}^{-1} \le C_{\sharp}$$

since $|\pi_1 \circ \gamma'(t)|^{-1} \ge \chi_c^{-1} > 1 > \chi_u$ implies $\vartheta_{\gamma}^{-1} \le 1$. To bound the L^1 norm of the derivative we can notice that:

(5.52)
$$\|\Psi_{\nu_n}'\|_{L^1} \le \left\|\frac{\Psi_{\nu_n}'}{\Psi_{\nu_n}}\right\|_{\mathcal{C}^0} \|\Psi_{\nu_n}\|_{L^1}$$

For each $0 \le i \le n$, let $\nu_{n-i} = F^i \nu_n$ and h_i be the diffeomorphism such that $\hat{\nu}_i = \nu_i \circ h_i$ is parametrized by vertical length. Define the diffeomorphisms h_i^* by

(5.53)
$$\hat{\nu}_i = F \circ \hat{\nu}_{i+1} \circ (h_{i+1}^*)^{-1},$$

where $\nu_0 = \gamma$ and $h_0^* = h_0 = Id$. Note that $h_i = h_1^* \circ \cdots \circ h_i^*$. We can then write

$$\Psi_{\nu_n}(t) = \frac{\frac{d}{dt}h_n(t)}{\det D_{\hat{\nu}_n(t)}F^n} = \frac{\prod_{i=1}^n (h_i^*)' \circ h_{i+1}^* \circ \cdots \circ h_n^*}{\prod_{i=1}^n (\det D_{\hat{\nu}_i}F) \circ h_{i+1}^* \circ \cdots \circ h_n^*} (t)$$
$$= \prod_{i=1}^n (\psi_i \circ h_{i+1}^* \circ \cdots \circ h_n^*)(t),$$

where $\psi_i(t) = (h_i^*)'(t) \cdot (\det D_{\hat{\nu}_i(t)}F)^{-1}$. Hence,

(5.54)
$$\left|\frac{\Psi_{\nu_n}'}{\Psi_{\nu_n}}\right| \le \sum_{i=1}^n \left| \left(\frac{\psi_i'}{\psi_i} \circ h_{i+1}^* \circ \cdots \circ h_n^*\right) (h_{i+1}^* \circ \cdots \circ h_n^*)' \right|$$

Differentiating twice (5.53), yields

$$\hat{\nu}_i' \circ h_{i+1}^* \cdot (h_{i+1}^*)' = D_{\hat{\nu}_{i+1}} F \hat{\nu}_{i+1}'.$$

³⁰Here m(A) is the Lebesgue measure of a set A.

Since $\hat{\nu}_n \in \Gamma(c_\star)$, multiplying by e_2 and remembering (2.9) we have

$$(h_{i+1}^*)' = \langle e_2, D_{\hat{\nu}_{i+1}}F\hat{\nu}'_{i+1} \rangle = (\partial_x\omega) \circ \hat{\nu}_{i+1} \cdot \langle e_1, \hat{\nu}'_{i+1} \rangle + 1 + (\partial_\theta\omega) \circ \hat{\nu}_{i+1}$$

Thus $\|\psi_i\|_{\mathcal{C}^0} \leq C_{\sharp}$ and, by (5.5), it follows that $\|\psi_i\|_{\mathcal{C}^1} \leq C_{\sharp}(1+C_Fc_{\star})$, while for $\rho \geq \ell > 1$, $\|\psi_i\|_{\mathcal{C}^\ell} \leq C_{\sharp}(c_{\star}^{(\ell-1)!} + C_Fc_{\star}^{\ell!})$. Thus, setting $h_{i,n} = h_{i+1}^* \circ \cdots \circ h_n^*$, by (5.5) we have

$$\begin{split} \left\| \frac{\Psi_{\nu_n}}{\Psi_{\nu_n}} \right\|_{\mathcal{C}^{\ell-1}} &\lesssim \sum_{i=0}^{n-1} \left\| (\log \psi_i \circ h_{i,n})' \right\|_{\mathcal{C}^{\ell-1}} \lesssim \sum_{i=0}^{n-1} \left\| \log \psi_i \circ h_{i,n} \right\|_{\mathcal{C}^{\ell}} \\ &\lesssim \sum_{i=0}^{n-1} \left\| \log \psi_i \right\|_{\mathcal{C}^{\ell}} \sum_{j=0}^{\ell-1} \left\| h_{i,n} \right\|_{\mathcal{C}^{\ell}}^j \end{split}$$

which, recalling (5.29), yields

(5.55)
$$\left\|\frac{\Psi_{\nu_n}'}{\Psi_{\nu_n}}\right\|_{\mathcal{C}^{\ell-1}} \lesssim \begin{cases} C_{\mu,n}(C_F c_{\star} + 1) & \text{for } \ell = 1\\ C_{\mu,n} c_{\star}(C_F c_{\star} + 1) & \text{for } \ell = 2\\ c_{\star}^{(\ell-1)!}(C_F c_{\star}^{\ell!} + 1)^{\ell} C_{\mu,n}^{\ell a_{\ell}} \mu^{n\ell \ell!} & \text{for } 2 < \ell \le \rho. \end{cases}$$

In particular, the above estimates in the case $\ell = 1$ and (5.52) gives

$$\sum_{\nu_n \in F^n \gamma} \|\Psi_{\nu_n}'\|_{L^1} \le C_{\sharp} C_{\mu,n} \mu^n (C_F c_{\star} + 1) \sum_{\nu_n \in F^n \gamma} \|\Psi_{\nu_n}\|_{L^1} \le C_{\sharp} C_{\mu,n} \mu^n (C_F c_{\star} + 1),$$

which gives

$$\sum_{\nu_n \in F^{-n}\gamma} \|\Psi_{\nu_n}\|_{\mathcal{C}^0} \lesssim C_{\mu,n} (1 + C_F c_\star).$$

Then

$$\sum_{\nu_n \in F^{-n}\gamma} \|\Psi_{\nu_n}\|_{\mathcal{C}^1} \lesssim \sum_{\nu_n \in F^{-n}\gamma} \|\Psi_{\nu_n}'\|_{\mathcal{C}^0} \lesssim \sum_{\nu_n \in F^{-n}\gamma} \left\|\frac{\Psi_{\nu_n}'}{\Psi_{\nu_n}}\right\|_{\mathcal{C}^0} \|\Psi_{\nu_n}\|_{\mathcal{C}^0} \lesssim C^2_{\mu,n} (1+C_F c_\star)^2$$
$$\sum_{\nu_n \in F^{-n}\gamma} \|\Psi_{\nu_n}\|_{\mathcal{C}^2} \lesssim \sum_{\nu_n \in F^{-n}\gamma} \left\|\frac{\Psi_{\nu_n}'}{\Psi_{\nu_n}}\right\|_{\mathcal{C}^1} \|\Psi_{\nu_n}\|_{\mathcal{C}^1} \lesssim C^3_{\mu,n} c_\star (1+C_F c_\star)^3.$$

To conclude we can obtain the general case $\rho \in \{3, \ldots, r-1\}$ by induction as follows:

$$\begin{split} \sum_{\nu_n \in F^{-n}\gamma} \|\Psi_{\nu_n}\|_{\mathcal{C}^{\rho}} &\lesssim \sum_{\nu_n \in F^{-n}\gamma} \|\Psi_{\nu_n}'\|_{\mathcal{C}^{\rho-1}} \lesssim \sum_{\nu_n \in F^{-n}\gamma} \left\|\frac{\Psi_{\nu_n}'}{\Psi_{\nu_n}}\right\|_{\mathcal{C}^{\rho-1}} \|\Psi_{\nu_n}\|_{\mathcal{C}^{\rho-1}} \\ &\lesssim c_{\star}^{(\rho-1)!} (C_F c_{\star}^{\rho!} + 1)^{\rho} C_{\mu,n}^{\rho a_{\rho}} \mu^{n\rho\rho!} \sum_{\nu_n \in F^{-n}\gamma} \|\Psi_{\nu_n}\|_{\mathcal{C}^{\rho-1}} \\ &\lesssim c_{\star}^{\rho!-2} (C_F c_{\star}^{\rho!} + 1)^{(\rho+1)\rho/2-3} C_{\mu,n}^{a_{\rho}} \sum_{k=0}^{\rho} k \mu^{n\rho!} \sum_{\nu_n \in F^{-n}\gamma} \|\Psi_{\nu_n}\|_{\mathcal{C}^2} \\ &\lesssim c_{\star}^{\rho!-1} (C_F c_{\star}^{\rho!} + 1)^{(\rho+1)\rho/2} C_{\mu,n}^{a_{\rho}} \frac{\rho^{(\rho+1)}}{2} + 1} \mu^{\rho! \frac{\rho(\rho+1)}{2} + 1)n}. \end{split}$$

The procedure to prove the second of (5.45) is analogous, with the difference that, by (5.46) and (C.8), the estimate for $\rho = 0$ gives another $C_{\sharp}\mu^n$, while the computation for $\rho \ge 1$ is exactly the same, but using $\psi_i = (\det D_{\hat{\nu}_i(t)}F)^{-1}$ instead.

The next result is a refinement of the previous Lemma in the more general case in which γ' is not contained neither in \mathbf{C}_c nor in \mathbf{C}_u , and it is parametrised by arc-length. Let h_0 be such that $\hat{\nu}_0 = \gamma \circ h_0$ is parametrised vertically. To state the result it is convenient to define the following quantities

(5.56)
$$\vartheta_{\hat{\nu}_0,m}(t) = \inf_{\|s-t\| \le c_{\sharp}m\| \omega\|_{\infty}} \vartheta_{\hat{\nu}_0}(s)$$

Lemma 5.19. In the same hypothesis of Lemma 5.13 with $n_0 = 0$, we have

(5.57)
$$\sum_{\nu_{\overline{m}}\in F^{-\overline{m}}\gamma} \left\| \frac{h'_{\overline{m}}}{\det D_{\hat{\nu}_{\overline{m}}}F^{\overline{m}}} \right\|_{\mathcal{C}^{\rho}(\mathbb{T})} \leq C_{\sharp}(\chi_{u}^{-1}\Delta_{\gamma})^{c_{\sharp}}\vartheta_{\gamma}^{-1}\Lambda^{c_{\sharp}\overline{m}}, \quad \rho \geq 2.$$

Proof. We use the same notations of the proof of Lemma 5.18. Note that (5.48) holds true for each $\gamma' \notin \mathbf{C}_u$, so we can proceed exactly as in the proof of above Lemma 5.18, but with the following differences: ϑ_{γ} in (5.51) is not bounded by a uniform constant C_{\sharp} , as γ is not an admissible central curve. Moreover, we use the estimates in (5.55) with $c_{\star} = \chi_u^{-2} ||\Delta_{\gamma}||_{\infty} (\mu \Lambda)^m$ provided by Lemma 5.13, so that the last of (5.57) follows.

By the above we can also obtain a key estimate of the L^{∞} norm of $\mathcal{L}^n 1$.

Corollary 5.20. Let $\mathcal{L} := \mathcal{L}_F$ be the transfer operator defined in (4.1). Then, for each $n \in \mathbb{N}$,

(5.58)
$$\|\mathcal{L}^n 1\|_{L^{\infty}(\mathbb{T}^2)} \le C_{\mu,n} (1 + C_F \mathfrak{c}) \mu^n.$$

Proof. For any $x \in \mathbb{T}^2$ we want to estimate the quantity

(5.59)
$$\mathcal{L}^n 1(x) = \sum_{y \in F^{-n}x} \frac{1}{|\det D_y F^n|}.$$

Recall the notation in Section 5.3 and take $y \in \gamma$, where $\gamma \in \Gamma(\mathfrak{c})$ is an admissible central curve. Then, for every $x \in F^{-n}(y)$, there exist $t \in \mathbb{T}$ and $\nu \in F^{-n}\gamma$ such that $x = \nu(h_n(t)) = \hat{\nu}(t)$. Hence

$$\sup_{y \in \gamma} \sum_{x \in F^{-n}(y)} \left| \frac{1}{\det D_x F^n} \right| \le \sum_{\nu \in F^{-n}\gamma} \left\| \frac{h'_{\nu,n}}{\det D_{\hat{\nu}} F^n} \right\|_{\mathcal{C}^0} \| (h'_{\nu,n})^{-1} \|_{\mathcal{C}^0}.$$

By equations (C.8) and (2.6) we know that $||(h'_n)^{-1}||_{\mathcal{C}^0} \leq C_{\sharp}\mu^n$, for every ν and n. Moreover, Lemma 5.18 gives the bound

$$\sum_{\nu \in F^{-n}\gamma} \left\| \frac{h'_{\nu}}{\det D_{\hat{\nu}}F^n} \right\|_{\mathcal{C}^0} \le C_{\mu,n}(1+C_F\mathfrak{c})\mu^n.$$

Remark 5.21. With some extra work the estimate (5.58) can be made sharper, however the above bound is good enough for our current purposes. We will need an improvement, provided in Lemma 10.5, in Section 10.

5.9. Distortion: the SVPH^{\ddagger} case. In the case of SVPH^{\ddagger} we need an improvement of Lemma 5.19 for the first derivatives. To state the result it is convenient to define the following quantities:

(5.60)

$$\begin{aligned}
\vartheta_{\hat{\nu}_{0},m}(t) &= \inf_{|s-t| \leq c_{\sharp}m \|\omega\|_{\infty}} \vartheta_{\hat{\nu}_{0}}(s) \\
\mathbb{J}_{\gamma,n} &= \int_{\mathbb{T}} \left[\vartheta_{\hat{\nu}_{0},n}(s) \right]^{-1} ds, \\
\mathbb{I}_{\gamma,m} &= C_{\mu,m} \mu^{4m} \left[1 + \mu^{m} (\vartheta_{\hat{\nu}_{0}} \circ \bar{h}_{m})^{-1} C_{F} \right] (\vartheta_{\hat{\nu}_{0}} \circ \bar{h}_{m})^{-1} \Delta_{\gamma}.
\end{aligned}$$

Lemma 5.22. In the same hypothesis of Lemma 5.16 with $n_0 = 0$, we have

$$(5.61) \qquad \sum_{\substack{\nu_{\overline{m}} \in F^{-\overline{m}}\gamma}} \left\| \frac{h'_{\overline{m}}}{\det D_{\hat{\nu}_{\overline{m}}}F^{\overline{m}}} \right\|_{\mathcal{C}^{0}(\mathbb{T})} \leq C_{\sharp}\mathbb{I}_{\gamma,\overline{m}}\mathbb{J}_{\gamma,\overline{m}}$$

$$(5.61) \qquad \sum_{\substack{\nu_{\overline{m}} \in F^{-\overline{m}}\gamma}} \left\| \frac{h'_{\overline{m}}}{\det D_{\hat{\nu}_{\overline{m}}}F^{\overline{m}}} \right\|_{\mathcal{C}^{1}(\mathbb{T})} \leq C_{\sharp}(\mathbb{I}_{\gamma,\overline{m}})^{2}\mathbb{J}_{\gamma,\overline{m}}$$

$$\sum_{\substack{\nu_{\overline{m}} \in F^{-\overline{m}}\gamma}} \left\| \frac{h'_{\overline{m}}}{\det D_{\hat{\nu}_{\overline{m}}}F^{\overline{m}}} \right\|_{\mathcal{C}^{2}(\mathbb{T})} \leq C_{\mu,\overline{m}}^{4}\mu^{5\overline{m}}(1+C_{F}\mu^{m}\vartheta_{\hat{\nu}_{0}}^{-1})O_{m}^{\star}\mathbb{I}_{\gamma,m}^{3}\mathbb{J}_{\gamma,\overline{m}}$$

where $O_m^{\star} = M_{m,0} (1 + C_F M_{m,0})^3$.³¹

³¹Where $M_{m,n}$ is defined in (5.41).

Proof. We use the same notations of the proof of Lemma 5.18. To clarify the argument we use the more precise notation $h_{\hat{\nu}_n}$ rather than h_n , since the dependence on the branch is relevant in the following. Thus, setting, for each $n \leq \overline{m}$, $\Psi_{\nu_n}(t) = \frac{h'_{\nu_n}(t)}{\det D_{\hat{\nu}_n(t)}F^n}$ and recalling (5.49), we can write

$$\sum_{\nu_n \in F^{-n}\gamma} \|\Psi_{\hat{\nu}_n}\|_{L^1} \le C_{\sharp} \sum_{\nu_n \in F^{-n}\gamma} \int_{\mathbb{T}} \frac{\delta_{\hat{\nu}_n}(s)}{\vartheta_{\gamma} \circ h_{\hat{\nu}_n}(s)} ds.$$

Setting $F^n(\hat{\nu}_n(t)) = \hat{\nu}_0 \circ \bar{h}_{\hat{\nu}_n}(t)$ we have $h_{\hat{\nu}_n}(t) = h_0 \circ \bar{h}_{\hat{\nu}_n}(t)$ and $\hat{\nu}_0 = \gamma \circ h_0$, $\pi_2(\hat{\nu}'_0) = 1$. Then, since $|\pi_2(F^n(x,\theta)) - \theta| \le n \|\omega\|_{\infty}$ it follows that,

$$|\bar{h}_{\hat{\nu}_m}(t) - t| = |\pi_2(F^m(\hat{\nu}_m(t))) - t))| \le m \|\omega\|_{\infty}$$

Accordingly,

$$\vartheta_{\gamma} \circ h_{\hat{\nu}_n}(s) = \vartheta_{\hat{\nu}_0} \circ \bar{h}_{\hat{\nu}_n}(s) \ge \inf_{|s-\tau| \le c_{\sharp}n \|\omega\|_{\infty}} \vartheta_{\hat{\nu}_0}(\tau) = \vartheta_{\hat{\nu}_0,n}(s).$$

Thus

(5.62)
$$\sum_{\hat{\nu}_n \in F^{-n}\gamma} \|\Psi_{\hat{\nu}_n}\|_{L^1} \le C_{\sharp} \int_{\mathbb{T}} \frac{\sum_{\hat{\nu}_n \in F^{-n}\gamma} \delta_{\hat{\nu}_n(s)}}{\vartheta_{\hat{\nu}_0,n}(s)} ds \le C_{\sharp} \int_{\mathbb{T}} [\vartheta_{\hat{\nu}_0,n}(s)]^{-1} ds.$$

Next, let $m_{\star}(t)$ be the smallest integer k for which $\hat{\nu}'_k \in \mathbf{C}_c$. Note that, setting $m_1 = \overline{m} - m_{\star}$ and $\tilde{\Psi}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m_{\star}}}(t) = \frac{\bar{h}'_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m_{\star}}}}{\det D_{\hat{\nu}_{\overline{m}}(t)}F^{m_1}}$, where $\bar{h}_{\hat{\nu}_n,\hat{\nu}_m}$ is such that $F^{n-m}\hat{\nu}_m = \hat{\nu}_n \circ \bar{h}_{\hat{\nu}_n,\hat{\nu}_m}$,

(5.63)
$$\sum_{\hat{\nu}_{\overline{m}}\in F^{-\overline{m}}\gamma} \Psi_{\hat{\nu}_{\overline{m}}}(t) = \sum_{F^{m_{\star}}\hat{\nu}_{m_{\star}}\in\gamma} \sum_{F^{m_{1}}\hat{\nu}_{\overline{m}}\in\hat{\nu}_{m_{\star}}} \frac{h'_{\hat{\nu}_{m_{\star}}} \circ \bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m_{\star}}} \cdot \bar{h}'_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m_{\star}}}}{\det D_{\hat{\nu}_{\overline{m}}}F^{m_{1}} \cdot \det D_{\hat{\nu}_{m_{\star}}} \circ \bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m_{\star}}}F^{m_{\star}}}$$
$$= \sum_{\hat{\nu}_{m_{\star}}\in F^{-m_{\star}}\gamma} \Psi_{\hat{\nu}_{m_{\star}}} \circ \bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m_{\star}}} \sum_{\hat{\nu}_{\overline{m}}\in F^{-m_{1}}\hat{\nu}_{m_{\star}}} \tilde{\Psi}_{\hat{\nu}_{\overline{m}}}.$$

Since $\hat{\nu}'_{m_{\star}} \in \mathbf{C}_c$, we apply Lemma 5.18, with $c_{\star} = \|\hat{\nu}''_{m_{\star}}\|$ and remember (C.45), we write

(5.64)
$$\begin{split} \sum_{\hat{\nu}_{\overline{m}}\in F^{-m_1}\hat{\nu}_{m_\star}} \left\|\tilde{\Psi}_{\hat{\nu}_{\overline{m}}}\right\|_{\mathcal{C}^0} &\leq C_{\mu,\overline{m}}(1+C_F M_{m_\star,0})\\ \sum_{\hat{\nu}_{\overline{m}}\in F^{-m_1}\hat{\nu}_{m_\star}} \left\|\tilde{\Psi}_{\hat{\nu}_{\overline{m}}}\right\|_{\mathcal{C}^1} &\leq C_{\mu,\overline{m}}^2(1+C_F M_{m_\star,0})^2\\ \sum_{\hat{\nu}_{\overline{m}}\in F^{-m_1}\hat{\nu}_{m_\star}} \left\|\tilde{\Psi}_{\hat{\nu}_{\overline{m}}}\right\|_{\mathcal{C}^2} &\leq C_{\mu,\overline{m}}^3 \mu^{\overline{m}} M_{m_\star,0}(1+C_F M_{m_\star,0})^3 \end{split}$$

Next, using (5.46),³²

(5.65)
$$\frac{\Psi'_{\hat{\nu}_{m_{\star}}}}{\Psi_{\hat{\nu}_{m_{\star}}}} = \frac{e_{1} \wedge \hat{\nu}'_{m_{\star}}}{e_{1} \wedge \hat{\nu}'_{m_{\star}}} \\ - \frac{\partial_{t} \left(D_{\hat{\nu}_{m_{\star}}} F^{m_{\star}} e_{1} \right) \wedge \gamma' \circ h_{m_{\star}} + D_{\hat{\nu}_{m_{\star}}} F^{m_{\star}} e_{1} \wedge \gamma'' \circ h_{m_{\star}} \cdot h'_{m_{\star}}}{D_{\hat{\nu}_{m_{\star}}} F^{m_{\star}} e_{1} \wedge \gamma' \circ h_{m_{\star}}} \\ = - \left(\left[(D_{\hat{\nu}_{m_{\star}}} F^{m_{\star}})^{-1} \partial_{t} \left(D_{\hat{\nu}_{m_{\star}}} F^{m_{\star}} \right) \right] e_{1} \right) \wedge \hat{\nu}'_{m_{\star}} \\ - e_{1} \wedge (D_{\hat{\nu}_{m_{\star}}} F^{m_{\star}})^{-1} \gamma'' \circ h_{m_{\star}} \cdot (h'_{m_{\star}})^{2}, \end{cases}$$

where we used that $e_1 \wedge \hat{\nu}''_{m_{\star}} = 0$, since the vectors are parallel, and $e_1 \wedge \hat{\nu}'_{m_{\star}} = 1$. Letting η , $\|\eta\| = 1$ such that $(DF^{m_{\star}})^{-1}\eta \wedge e_1 = 0$ we can write $e_1 = a\eta + be_2$, with $|b| \leq \chi_u$. Hence, using (5.23), (5.25),

 $^{^{32}}$ Remark that two forms are isomorphic to functions, and since here they are never zero we can disregard the norm.

(C.1), (5.42) and Sublemma C.1 we have

$$(5.66) \qquad \| \left[(D_{\hat{\nu}_{m_{\star}}} F^{m_{\star}})^{-1} \partial_{t} \left(D_{\hat{\nu}_{m_{\star}}} F^{m_{\star}} \right) \right] e_{1} \| \leq \\ \leq \sum_{k=0}^{m_{\star}-1} \| (D_{\hat{\nu}_{m_{\star}}} F^{k+1})^{-1} \left[\partial_{x_{i}} D_{F^{k}(\hat{\nu}_{m_{\star}})} F \right] D_{\hat{\nu}_{m_{\star}}} F^{k} e_{1} \| \cdot \| D_{\hat{\nu}_{m_{\star}}} F^{k} \hat{\nu}_{m_{\star}}' \| \\ \leq \sum_{k=0}^{m_{\star}-1} C_{\sharp} (1 + \mu^{k} C_{F} \lambda_{k}^{+}) \| (D_{F^{k} \hat{\nu}_{m_{\star}}} F^{m_{\star}-k})^{-1} D_{\hat{\nu}_{m_{\star}}} F^{m_{\star}} \hat{\nu}_{m_{\star}}' \| \\ \leq C_{\mu,m_{\star}} \mu^{m_{\star}} (1 + \lambda_{m_{\star}}^{+} (\hat{\nu}_{m_{\star}}) C_{F}) | h_{m_{\star}}' | \\ \leq C_{\mu,m_{\star}} \mu^{2m_{\star}} (1 + \mu^{m_{\star}} (\vartheta_{\hat{\nu}_{0}} \circ \bar{h}_{m_{\star}})^{-1} C_{F}) (\vartheta_{\hat{\nu}_{0}} \circ \bar{h}_{m_{\star}})^{-1}.$$

On the other hand, writing, as before, $\gamma'' = \eta + \alpha e_2$ with $\|\eta\| \le \|\gamma''\|$, $|\alpha| \le C_F \|\gamma''\|$ and $DF^{m_*}e_1 \land \eta = 0$, yields

$$|e_1 \wedge (D_{\hat{\nu}_{m_{\star}}}F^{m_{\star}})^{-1}\gamma''| = |\langle e_2, (D_{\hat{\nu}_{m_{\star}}}F^{m_{\star}})^{-1}\gamma''\rangle| \le C_{\sharp}\mu^{m_{\star}}(1+\mu^{m_{\star}}C_F\vartheta_{\hat{\nu}_0})\vartheta_{\hat{\nu}_0}\Delta_{\gamma}.$$

Thus, recalling (5.29),

(5.67)
$$\left| \frac{\Psi'_{\hat{\nu}_{m_{\star}}}}{\Psi_{\hat{\nu}_{m_{\star}}}} \circ \bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m_{\star}}} \cdot \bar{h}'_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m_{\star}}} \right| \le C_{\sharp} \mathbb{I}_{\gamma,m_{\star}}.$$

By (5.63), (5.67) and (5.62) it follow, for all $n \in \{m_{\star}, \ldots, \overline{m}\}$,

(5.68)
$$\sum_{\hat{\nu}_{n}\in F^{-n}\gamma} \|\Psi_{\hat{\nu}_{n}}\|_{\mathcal{C}^{0}} \leq \sum_{\hat{\nu}_{n}\in F^{-n}\gamma} \left(1 + \left\|\frac{\Psi_{\hat{\nu}_{n}}}{\Psi_{\hat{\nu}_{n}}}\right\|_{\mathcal{C}^{0}}\right) \|\Psi_{\hat{\nu}_{n}}\|_{L^{1}} \leq C_{\sharp} \mathbb{I}_{\gamma,n} \sum_{\hat{\nu}_{n}\in F^{-n}\gamma} \|\Psi_{\hat{\nu}_{n}}\|_{L^{1}} \leq C_{\sharp} \mathbb{I}_{\gamma,n} \int_{\mathbb{T}} \left[\vartheta_{\hat{\nu}_{0},n}(s)\right]^{-1} ds$$

which proves the first of (5.57). Next,

$$\|\Psi'_{\hat{\nu}\overline{m}}\|_{\mathcal{C}^0} \leq \left\|\frac{\Psi'_{\hat{\nu}\overline{m}}}{\Psi_{\hat{\nu}\overline{m}}}\right\|_{\mathcal{C}^0} \|\Psi_{\hat{\nu}\overline{m}}\|_{\mathcal{C}^0}.$$

leads immediately to the second of (5.61).

To conclude the lemma we must compute $\Psi''_{\hat{\nu}_{m_*}}$, which can be obtained by (5.65):

$$(5.69) \frac{\Psi_{\hat{\nu}_{m_{\star}}}'}{\Psi_{\hat{\nu}_{m_{\star}}}} = \left[\frac{\Psi_{\hat{\nu}_{m_{\star}}}'}{\Psi_{\hat{\nu}_{m_{\star}}}}\right]' + \left[\frac{\Psi_{\hat{\nu}_{m_{\star}}}'}{\Psi_{\hat{\nu}_{m_{\star}}}}\right]^{2} = -\left\{\partial_{t}\left[(D_{\hat{\nu}_{m_{\star}}}F^{m_{\star}})^{-1}\partial_{t}\left(D_{\hat{\nu}_{m_{\star}}}F^{m_{\star}}\right)\right]e_{1}\right\} \wedge \hat{\nu}_{m_{\star}}' - \left\{\left[(D_{\hat{\nu}_{m_{\star}}}F^{m_{\star}})^{-1}\partial_{t}\left(D_{\hat{\nu}_{m_{\star}}}F^{m_{\star}}\right)\right]e_{1}\right\} \wedge \hat{\nu}_{m_{\star}}'' - e_{1} \wedge \left[\partial_{t}(D_{\hat{\nu}_{m_{\star}}}F^{m_{\star}})^{-1}\right]\gamma'' \circ h_{m_{\star}} \cdot (h_{m_{\star}}')^{2} - e_{1} \wedge (D_{\hat{\nu}_{m_{\star}}}F^{m_{\star}})^{-1}\gamma''' \circ h_{m_{\star}} \cdot (h_{m_{\star}}')^{3} - 2e_{1} \wedge (D_{\hat{\nu}_{m_{\star}}}F^{m_{\star}})^{-1}\gamma'' \circ h_{m_{\star}} \cdot h_{m_{\star}}'h_{m_{\star}}'' + \left[\frac{\Psi_{\hat{\nu}_{m_{\star}}}'}{\Psi_{\hat{\nu}_{m_{\star}}}}\right]^{2}.$$

We estimate the lines of (5.69) one at a time. For the first line, by (5.23) we can write

$$(D_{\hat{\nu}_{m_{\star}}}F^{m_{\star}})^{-1}\partial_t(D_{\hat{\nu}_{m_{\star}}}F^{m_{\star}}) = \sum_{s=1}^2 \sum_{k=0}^{m_{\star}-1} (D_{\hat{\nu}_{m_{\star}}}F^{k+1})^{-1}\partial_{x_s}(D_{F^k(\hat{\nu}_{m_{\star}}(t))}F)D_{\hat{\nu}_{m_{\star}}(t)}F^k(D_{\hat{\nu}_{m_{\star}}(t)}F^k\hat{\nu}_{m_{\star}}')_s.$$

We can thus use the fourth of (5.21), with $n = m = m_{\star}$ and $c = M_{m_{\star},0}$, and argue as in (5.66) to bound the first line of (5.69) by

(5.70)
$$C_{\sharp}(1+\mu^{m_{\star}}C_{F}\vartheta_{\hat{\nu}_{0}}^{-1})\left[(\lambda_{m_{\star}}^{+}\circ\hat{\nu}_{m_{\star}})^{2}+(\lambda_{m_{\star}}^{+}\circ\hat{\nu}_{m_{\star}})^{2}+\lambda_{m_{\star}}^{+}\circ\hat{\nu}_{m_{\star}}M_{m_{\star},0}\right] \leq C_{\sharp}(1+\mu^{m_{\star}}C_{F}\vartheta_{\hat{\nu}_{0}}^{-1})^{2}\mu^{4m_{\star}}\vartheta_{\hat{\nu}_{0}}^{-2}(1+\Delta_{\gamma}\vartheta_{\hat{\nu}_{0}}).$$

To estimate the second line we use the second line of (5.23), arguing as above, and (5.42)

(5.71)
$$C_{\mu,m_{\star}}\mu^{4m_{\star}}(1+\mu^{m_{\star}}C_{F}\vartheta_{\hat{\nu}_{0}}^{-1})^{2}\Delta_{\gamma}\vartheta_{\hat{\nu}_{0}}^{-1}+\mu^{4m_{\star}}(1+\mu^{m_{\star}}C_{F}\vartheta_{\hat{\nu}_{0}}^{-1})\Delta_{\gamma}^{2}\vartheta_{\hat{\nu}_{0}}^{-2}$$

Finally, again by (5.42) and recalling (5.67), the last line is estimated by

(5.72)
$$\mu^{4m_{\star}}(1+\mu^{m_{\star}}C_F\vartheta_{\hat{\nu}_0}^{-1})M_{m_{\star},0}\vartheta_{\hat{\nu}_0}^{-2}+C_{\sharp}\mathbb{I}^2_{\gamma,m_{\star}}\mu^{-2m}$$

Thus

(5.73)
$$\left|\frac{\Psi_{\hat{\nu}_{m_{\star}}}'}{\Psi_{\hat{\nu}_{m_{\star}}}}\right| \leq \mu^{7m_{\star}} C_{\mu,m_{\star}}^2 (1 + \mu^{m_{\star}} C_F \vartheta_{\hat{\nu}_0}^{-1})^2 (1 + \Delta_{\gamma}^2 + \mu^{m_{\star}} \vartheta_{\hat{\nu}_0}^{-1}) \vartheta_{\hat{\nu}_0}^{-2}.$$

To conclude note that

(5.74)
$$\begin{aligned} \left| \frac{\left(\Psi_{\nu_{m\star}} \circ \bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m\star}}\right)''}{\Psi_{\nu_{m\star}} \circ \bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m\star}}} \right| &= \left| \frac{\Psi_{\nu_{\overline{m}}}'' \circ \bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m\star}}}{\Psi_{\nu_{\overline{m}}} \circ \bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m\star}}} \left(\bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m\star}} \right)^2 + \frac{\Psi_{\nu_{\overline{m}}}' \circ \bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m\star}}}{\Psi_{\nu_{\overline{m}}} \circ \bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m\star}}} \bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m\star}} \right| \\ &\leq \left| \frac{\Psi_{\nu_{\overline{m}}}''}{\Psi_{\nu_{\overline{m}}}} \right| \mu^{2\overline{m}} + C_{\mu,\overline{m}} \left| \frac{\Psi_{\nu_{\overline{m}}}'}{\Psi_{\nu_{\overline{m}}}} \right| (1 + C_F M_{m\star,0}) \mu^{2\overline{m}} \\ &\leq C_{\mu,m\star}^3 \mu^{10m\star} (1 + C_F \mu^{m\star} \vartheta_{\hat{\nu}_0}^{-1})^3 \Delta_{\gamma}^2 \vartheta_{\hat{\nu}_0}^{-2} \end{aligned}$$

$$\leq C_{\mu,m_{\star}}\mu^{2m_{\star}}(1+C_{F}\mu^{m_{\star}}\vartheta_{\hat{\nu}_{0}}^{-1})\mathbb{I}_{\gamma,m}^{2},$$

where, in the last lines we have used (5.29), with $c_{\star} = M_{m_{\star},0}$, and (5.73), (5.67). We finally have, by (5.64), (5.74), and (5.68),

$$\sum_{\nu_{m}\in F^{m}\gamma} \|\Psi_{\nu_{\overline{m}}}\|_{\mathcal{C}^{2}} \leq \sum_{\hat{\nu}_{m_{\star}}\in F^{-m_{\star}\gamma}} \|\Psi_{\hat{\nu}_{m_{\star}}}\circ\bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m_{\star}}}\|_{\mathcal{C}^{2}} \sum_{\hat{\nu}_{\overline{m}}\in F^{-m_{1}}\hat{\nu}_{m_{\star}}} \left\|\tilde{\Psi}_{\hat{\nu}_{\overline{m}}}\right\|_{\mathcal{C}^{2}}$$

$$\leq \sum_{\hat{\nu}_{m_{\star}}\in F^{-m_{\star}\gamma}} \left\{ \mathbb{I}_{\gamma,m_{\star}} + \left\|\frac{[\Psi_{\hat{\nu}_{m_{\star}}}\circ\bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m_{\star}}}]''}{\Psi_{\hat{\nu}_{m_{\star}}}\circ\bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m_{\star}}}}\right\|_{\mathcal{C}^{0}} \right\} \|\Psi_{\hat{\nu}_{m_{\star}}}\circ\bar{h}_{\hat{\nu}_{\overline{m}},\hat{\nu}_{m_{\star}}}\|_{\mathcal{C}^{0}}$$

$$\times C^{3}_{\mu,\overline{m}}\mu^{\overline{m}}M_{m_{\star},0}(1+C_{F}M_{m_{\star},0})^{3}$$

$$\leq C^{4}_{\mu,\overline{m}}\mu^{5\overline{m}}(1+C_{F}\mu^{m_{\star}}\vartheta_{\hat{\nu}_{0}}^{-1})M_{m_{\star},0}(1+C_{F}M_{m_{\star},0})^{3}\mathbb{I}^{3}_{\gamma,m}\mathbb{J}_{\gamma,\overline{m}}.$$

The Lemma follows since $m \leq m_{\star}$.

6. A FIRST LASOTA-YORKE INEQUALITY

We define a class of geometric norms inspired by [32] and [3]. Given $u \in \mathcal{C}^r(\mathbb{T}^2, \mathbb{R})$ and an integer $\rho < r$, we denote by \mathcal{B}_{ρ} the completion of $\mathcal{C}^r(\mathbb{T}^2, \mathbb{R})$ with respect to the norm:

(6.1)
$$\|u\|_{\rho} := \max_{|\alpha| \le \rho} \sup_{\substack{\gamma \in \Gamma(\mathbb{C}) \\ \|\phi\|_{\mathcal{C}}|\alpha| = 1}} \int_{\mathbb{T}} \phi(t) (\partial^{\alpha} u)(\gamma(t)) dt$$

This defines a decreasing sequence of Banach spaces continuously embedded in L^1 , namely

(6.2)
$$||u||_{L^1} \le C ||u||_{\rho_1} \le C ||u||_{\rho_2}, \quad \text{for every} \quad 0 \le \rho_1 \le \rho_2 \le r-1$$

To see this we observe that, since $\sigma_x(t) = (x, t) \in \Gamma$,

$$\begin{aligned} \|u\|_{L^{1}} &= \sup_{\|\phi\|_{C^{0}(\mathbb{T}^{2})} \leq 1} \int_{\mathbb{T}}^{\cdot} dx \int_{\mathbb{T}}^{\cdot} dy \phi(x, y) u(x, y) \leq \int_{\mathbb{T}}^{\cdot} dx \sup_{\|\phi\|_{C^{0}(\mathbb{T}^{2})} \leq 1} \int_{\mathbb{T}}^{\cdot} dy \phi(x, y) u(x, y) \\ &\leq \int_{\mathbb{T}}^{\cdot} dx \sup_{\|\phi\|_{C^{0}(\mathbb{T})} \leq 1} \int_{\mathbb{T}}^{\cdot} dt \ \phi(t) u(\sigma_{x}(t)) \leq \int_{\mathbb{T}}^{\cdot} dx \|u\|_{0} = \|u\|_{0}. \end{aligned}$$

The above proves the first inequality of (6.2), the others being trivial. Next, we prove a Lasota-Yorke type inequality between the spaces \mathcal{B}_{ρ} and $\mathcal{B}_{\rho-1}$.

Theorem 6.1. Let $F \in C^r(\mathbb{T}^2, \mathbb{T}^2)$ be a SVPH. Let $\mathcal{L} := \mathcal{L}_F$ be the transfer operator defined in (4.1), and \bar{n} be the integer given in Lemma 5.10. For each $\rho \in [1, r-1]$ and $n > \bar{n}$, there exists $C_{n,\rho}$ such that

(6.3)
$$\|\mathcal{L}^{n}u\|_{0} \leq (1+C_{F}\mathfrak{c})C_{\mu,n}\mu^{n}\|u\|_{0}$$

(6.4)
$$\|\mathcal{L}^{n}u\|_{\rho} \leq c_{\rho,F} \frac{C_{\mu,n}^{a_{\rho}} \mu^{b_{\rho}n}}{\lambda_{-}^{\rho_{n}}} \|u\|_{\rho} + C_{n,\rho} \|u\|_{\rho-1}$$

where $\bar{a}_{\rho} = 1 + 2a_{\rho}\rho^{2} + \rho$, $\bar{b}_{\rho} = (\rho! + 1)(\rho^{2} + 2)$ and $\mathfrak{c}_{\rho,F} := \mathfrak{c}^{\rho!(2\rho+1)}(1 + C_{F}\mathfrak{c}^{\rho!})^{\bar{c}_{\rho}}$ where $\bar{c}_{\rho} = 1 + \{\rho + 1, \rho(\rho+1)/2\}^{+}$.

We postpone the proof of Theorem 6.1 to section 6.3. First we need to develop several results on the commutators between differential operators and transfer operators.

6.1. Differential Operators.

For $s, \rho \in \mathbb{N}$ we denote by $P_{s,\rho}$ a differential operator of order at most ρ defined as a finite linear combination of compositions of at most ρ vector fields, and we write

(6.5)
$$P_{s,\rho}u = \sum_{j=0}^{s} \sum_{\alpha \in A \subset \mathbb{N}^{j}} v_{j,\alpha_{1}} \cdots v_{j,\alpha_{j}} u,$$

where A is a finite set and for every $i \leq j$, v_{j,α_i} are vector fields in $\mathcal{C}^{\rho+j-s}$, with the convention that $v_{j,\alpha_1} \cdots v_{j,\alpha_j} u = u$ if j = 0. We denote by $\Psi^{s,\rho}$ the set of differential operators $P_{s,\rho}$. For a function $u \in \mathcal{C}^r(\mathbb{T}^2, \mathbb{R})$ and a smooth vector field v, we denote $\partial_v u(x) = \langle \nabla_x u, v(x) \rangle$.

We start by studying the structure of the commutator between \mathcal{L} and the differential operators. Next, we will estimate the coefficients of the commutator.

Proposition 6.2. Given smooth vector fields $v_1, \dots, v_s \in C^{\rho}$, we have

$$\partial_{v_s} \cdots \partial_{v_1} \mathcal{L}^n = \mathcal{L}^n \partial_{F^n v_s} \cdots \partial_{F^n v_1} + \mathcal{L}^n P_{s-1,\rho}$$

where $F^*v(x) := (D_x F)^{-1}v(F(x))$ is the pullback of v by the map F and $P_{s-1,\rho} \in \Psi^{s-1,\rho}$ whose coefficients may depend on n.

Proof. Let us start with s = 1. Let $v_1 \in \mathcal{C}^{\rho}(\mathbb{T}^2, \mathbb{T}^2)$ and define

(6.6)
$$J_n(p) = (\det D_p F^n)^{-1}; \quad \phi_n(p) = \log |\det D_p F^n|.$$

For each $\mathfrak{h} \in \mathfrak{H}^n$ we have

$$\begin{split} \langle \nabla \left[J_n \circ \mathfrak{h} \cdot u \circ \mathfrak{h} \right], v_1 \rangle &= \langle J_n \circ \mathfrak{h}(D\mathfrak{h})^* \nabla u \circ \mathfrak{h}, v_1 \rangle - \langle (D\mathfrak{h})^* \nabla (\det DF^n) \circ \mathfrak{h} J_n^2 \circ \mathfrak{h} u \circ \mathfrak{h}, v_1 \rangle \\ &= J_n \circ \mathfrak{h} \langle (D\mathfrak{h})^* \nabla u \circ \mathfrak{h}, v_1 \rangle - J_n \circ \mathfrak{h} \langle (D\mathfrak{h})^* \nabla \phi_n \circ \mathfrak{h} u \circ \mathfrak{h}, v_1 \rangle. \end{split}$$

Then, since $DF^n \circ \mathfrak{h}D\mathfrak{h} = \mathrm{Id}_{\mathcal{R}_{\mathfrak{h}}}$, for each $\mathfrak{h} \in \mathfrak{H}^n$ and $x \in \mathcal{D}_{\mathfrak{h}}^{33}$

(6.7)
$$\langle \nabla \left[J_n \circ \mathfrak{h} \cdot u \circ \mathfrak{h} \right] (x), v_1(x) \rangle = J_n \circ \mathfrak{h}(x) \left[\partial_{F^{n*}v_1} u - \partial_{F^{n*}v_1} \phi_n u \right] \circ \mathfrak{h}(x).$$

Observing that

(6.8)
$$\mathcal{L}^{n}u = \sum_{\mathfrak{h}\in\mathfrak{H}^{n}} u\circ\mathfrak{h}J_{n}\circ\mathfrak{h}\mathbb{1}_{\mathcal{R}_{\mathfrak{h}}}\circ\mathfrak{h}.$$

it follows

(6.9)
$$\langle \nabla_x \mathcal{L}^n u, v_1(x) \rangle = \mathcal{L}^n \left(\partial_{F^{n^*} v_1} u \right) (x) - \mathcal{L}^n (\partial_{F^{n^*} v_1} \phi_n \cdot u) (x),$$

 $^{^{33}\}mathrm{Recall}$ that $\mathcal{D}_{\mathfrak{h}},\mathcal{R}_{\mathfrak{h}}$ indicate respectively the domain and the range of $\mathfrak{h}.$

which prove the result since the multiplication operator $P_{0,\rho} := -\partial_{F^{n^*}v_1} \phi_n \in \Psi^{0,\rho}$. Next, we argue by induction on s:

$$(6.10) \qquad \begin{aligned} \partial_{v_{s+1}} \cdots \partial_{v_1} \mathcal{L}^n u &= \partial_{v_{s+1}} \left[\mathcal{L}^n \partial_{F^{n^*} v_s} \cdots \partial_{F^{n^*} v_1} u + \mathcal{L}^n P_{s-1,\rho} u \right] \\ &= \mathcal{L}^n \partial_{F^{n^*} v_{s+1}} \cdots \partial_{F^{n^*} v_1} u + \mathcal{L}^n (\partial_{F^{n^*} v_{s+1}} \phi_n \cdot \partial_{F^{n^*} v_s} \cdots \partial_{F^{n^*} v_1} u) \\ &+ \mathcal{L}^n \partial_{F^{n^*} v_{s+1}} P_{s-1,\rho} u + \mathcal{L}^n (\partial_{F^{n^*} v_{s+1}} \phi_n \cdot P_{s-1,\rho} u), \end{aligned}$$

which yields the Lemma with

(6.11)
$$P_{s,\rho} = \partial_{F^{n^*}v_{s+1}} P_{s-1,\rho} + \partial_{F^{n^*}v_{s+1}} \phi_n \cdot \left[\partial_{F^{n^*}v_s} \cdots \partial_{F^{n^*}v_1} + P_{s-1,\rho}\right] \\ + \partial_{F^{n^*}v_{s+1}} P_{s-1,\rho}.$$

In the case $v_j \in \{e_1, e_2\}$ for each j, we have the following Corollary as an immediate iterative application of formulae (6.7) and (6.9).

Corollary 6.3. For each
$$t \ge 1$$
, $n \in \mathbb{N}$ $\alpha = (\alpha_1, ..., \alpha_t) \in \{1, 2\}^t$ and $\mathfrak{h} \in \mathfrak{H}^n$,
(6.12) $\partial^{\alpha}[J_n \circ \mathfrak{h} \cdot u \circ \mathfrak{h}] = J_n \circ \mathfrak{h} \cdot [P_{n,t}^{\alpha}u] \circ \mathfrak{h}$,

in particular

(6.13)
$$\partial^{\alpha} \mathcal{L}^n u = \mathcal{L}^n P_{n,t}^{\alpha} u,$$

the operators $P_{n,t}^{\alpha}$ being defined by the following relations, for each $u \in \mathcal{C}^t$,

(6.14)
$$\begin{cases} P_{n,0}^{\alpha}u = u, \\ P_{n,1}^{\alpha}u = A_{n}^{\alpha_{1}}u - A_{n}^{\alpha_{1}}\phi_{n} \cdot u, \\ P_{n,t}^{\alpha}u = A_{n,1}^{\alpha}u - \sum_{k=1}^{t}A_{n,k+1}^{\alpha}((A_{n}^{\alpha_{k}}\phi_{n}) \cdot P_{n,k-1}^{\alpha}u) & \text{for } t \ge 2, \end{cases}$$

where $A_n^{\alpha_i} = \partial_{F^n^* e_{\alpha_i}}$, $A_{n,k}^{\alpha} := A_n^{\alpha_t} \cdots A_n^{\alpha_k}$, $A_{n,t+1}^{\alpha} = Id$ and ϕ_n is defined in (6.6).

Proposition 6.4. For each $n \in \mathbb{N}$ let $P_{n,t}^{\alpha} \in \Psi^{t,t}$ given by (6.14). For any $0 \leq t < r$, $\psi \in \mathcal{C}^{r}(\mathbb{T}^{2},\mathbb{C})$ with $supp \psi \subset U = \mathring{U} \subset \mathbb{T}^{2}$, $\nu \in \Gamma(\mathfrak{c})$ such that $DF^{n-m}\nu' \in \mathbf{C}_{c}$, $\varphi \in \mathcal{C}^{t}(\mathbb{T},\mathbb{C})$ with $\|\varphi\|_{\mathcal{C}^{t}} \leq 1$, multi-index α , $|\alpha| = t$ and $u \in \mathcal{C}^{r}(\mathbb{T}^{2})$ we have

(6.15)
$$\int_{\mathbb{T}} \varphi(\tau) P_{n,t}^{\alpha}(\psi u)(\nu(\tau)) d\tau \leq \tilde{C}(t,n,m) \|\psi\|_{\mathcal{C}^{t}(U)} \|u\|_{t},$$

where $\tilde{C}(t, n, m) \leq C_{\sharp} \Lambda^{c_{\sharp} t n}$.

Proof. For simplicity we set $\partial_k = \partial_{x_k}$ for $k \in \{1, 2\}$. First of all notice that, if we set $d_{k,i} = \langle (DF^n)^{-1}e_k, e_i \rangle$, then $A_n^{\alpha_j} = \sum_{i=1}^2 d_{\alpha_j,i}\partial_{x_i}$. Furthermore, by formula (5.6), $\|d_{j,i}\|_{\mathcal{C}^t} \leq \|(DF^n)^{-1}\|_{\mathcal{C}^t} \leq \Lambda^n$, for each $1 \leq t \leq r$. We are going to prove (6.15) by induction on t. For t = 0 it is obvious, and for t = 1 it follows from $\|(DF^n)^{-1}\| \leq C_{\sharp}\mu^n$. Let us assume it for any $k \leq t - 1$. By (6.14) the integral in (6.15) splits into³⁴

(6.16)
$$\int \varphi(\tau) P_{n,t}^{\alpha}(\psi u)(\nu(\tau)) d\tau$$
$$= \int \varphi \left[A_n^{\alpha_t} \cdots A_n^{\alpha_1}(\psi u) \right] \circ \nu - \int \varphi \sum_{k=1}^t \left[A_{n,k+1}^{\alpha}((A_n^{\alpha_k}\phi_n) \cdot P_{n,k-1}^{\alpha}(\psi u)) \right] \circ \nu$$

The first integral is equal to

(6.17)
$$\sum_{\substack{i_1,\cdots,i_t\\i_l\in\{1,2\}}}\sum_{J,J_0,J_1,\dots,J_t}\int \varphi \cdot \big(\prod_{j\in J}\partial_j u\big)\big(\prod_{j\in J_0}\partial_j\psi\big)\big(\prod_{j\in J_1}\partial_j d_{\alpha_1,i_1}\big)\cdots\big(\prod_{j\in J_t}\partial_j d_{\alpha_t,i_t}\big),$$

 $^{^{34}}$ Unless differently specified, in the following all the integrals are on T.

where the second sum is made over all the partitions $J, J_0, J_1, ..., J_t$ of $\{1, ..., t\}$ such that $J_j \subset \{j + 1, ..., t\}, j \geq 1.^{35}$ Note that $\left\| (\prod_{k=1}^t \prod_{j \in J_k} \partial_j) d_{\alpha_k, i_k} \right\|_{\mathcal{C}^{\sharp J}_{\nu}} \leq \Lambda^{n\{\sharp J + \sum_{k=1}^t \sharp J_k\}}$ and $\|\prod_{j \in J_0} \partial_j \psi\|_{\mathcal{C}^{\sharp J}_{\nu}} \leq \|\psi\|_{\mathcal{C}^{\sharp J}}$. Consequently, from (6.17) and the definition (6.1), we have

(6.18)
$$\left| \int \varphi(\tau) A_{n,1}^{\alpha}(\psi u)(\nu(\tau)) d\tau \right| \leq C_{\sharp} \Lambda^{c_{\sharp} t n} \|\psi\|_{\mathcal{C}^{t}} \|u\|_{t}.$$

To bound the second integral in (6.16) we first note that

(6.19)
$$A_{n}^{\alpha_{k}}\phi_{n}(x) = \sum_{j=0}^{n-1} \langle (D_{x}F^{j})^{*}\nabla\phi_{1}\circ F^{j}(x), (D_{x}F^{n})^{-1}e_{\alpha_{k}}\rangle$$
$$= \sum_{j=0}^{n-1} \langle \nabla\phi_{1}, (DF^{n-j})^{-1}e_{\alpha_{k}}\rangle \circ F^{j}(x),$$

thus (5.5) implies

(6.20)
$$\|A_n^{\alpha_k}\phi_n\|_{\mathcal{C}^l} \le C_{\sharp} \sum_{j=0}^{n-1} \|(DF^{n-j})^{-1}\|_{\mathcal{C}^l} \Lambda^{nl} \le C_{\sharp} \sum_{j=0}^{n-1} \Lambda^{c_{\sharp}(n-j+l)} \le C_{\sharp} \Lambda^{c_{\sharp}n}.$$

We can then use (6.18) to estimate

(6.21)
$$\left| \int \varphi A_{n,k+1}^{\alpha} ((A_n^{\alpha_k} \phi_n) \cdot P_{n,k-1}^{\alpha}(\psi u)) \right| \leq C_{\sharp} \Lambda^{c_{\sharp}n} \|A_n^{\alpha_k} \phi_n\|_{\mathcal{C}^{t-k-1}_{\nu}} \|P_{n,k-1}^{\alpha}(\psi u))\|_{t-k-1} \leq C_{\sharp} \Lambda^{c_{\sharp}n} \|P_{n,k-1}^{\alpha}(\psi u))\|_{t-k-1}.$$

To bound the last term we take $\phi \in \mathcal{C}^{t-k-1}$, $\|\phi\|_{\mathcal{C}^{t-k-1}} = 1, \gamma \in \Gamma$, and we consider

$$\int \phi \partial^{t-k-1} [P_{n,k-1}^{\alpha}(\psi u)] \circ \gamma.$$

We can then split the integral as in (6.16), although this time $\alpha = (\alpha_1, \dots, \alpha_{k-1})$. For the first term we take t - k - 1 derivatives in (6.17) and, arguing as we did to prove (6.18), we have

$$\left|\int \phi(\tau)\partial^{t-k-1}A_{n,1}^{\alpha}(\psi u)(\gamma(\tau))d\tau\right| \leq C_{\sharp}\Lambda^{c_{\sharp}tn}\|\psi\|_{\mathcal{C}^{t}}\|u\|_{t}$$

The second term is estimated in the same way, using the inductive assumption. The statement of the Proposition then follows using this in (6.21).

6.2. Differential operators: the SVPH^{\ddagger} case. When treating SVPH^{\ddagger} systems we will need an improved estimate of the constant $\tilde{C}(t, n, m)$ appearing in Proposition 6.4 for low derivatives.

Proposition 6.5. If F is SVPH^{\sharp}, then the constants $\tilde{C}(t, n, m)$ of Proposition 6.4 satisfy

(6.22)
$$\tilde{C}(t,n,m) \leq \begin{cases} 1 & t = 0 \\ C_{\sharp}\mu^n & t = 1 \\ C_{\mu,n}^3\mu^{4n} \sup_{\zeta \in \operatorname{supp}(\varphi) \cap \nu(\mathbb{T})} (1 + C_F\lambda_n^+(\zeta))\{\lambda_m^+ \circ F^{n-m}(\zeta) + \mathfrak{c}\} & t = 2. \end{cases}$$

³⁵We use the conventions $\prod_{j \in \emptyset} \partial_j A = A$ and $\sharp B$ denote the cardinality of the set B.

Proof. We use the same notation of Proposition 6.4 and of its proof, where the cases t = 0 and t = 1have already been handled. The special case t = 2 corresponds to $\alpha = (\alpha_1, \alpha_2)^{36}$

$$\begin{aligned} P_{n,2}^{\alpha}(\psi u) &= A_{n,1}^{\alpha}(\psi u) - A_{n}^{\alpha_{2}}(\phi_{n})A_{n}^{\alpha_{1}}(\psi u) - A_{n,1}^{\alpha}(\phi_{n})\psi u - A_{n}^{\alpha_{1}}\phi_{n}A_{n}^{\alpha_{2}}(\psi u) \\ &- A_{n}^{\alpha_{1}}\phi_{n}A_{n}^{\alpha_{2}}\phi_{n} \cdot \psi u \\ &= \left\{A_{n,1}^{\alpha}\psi - A_{n}^{\alpha_{2}}\phi_{n}A_{n}^{\alpha_{1}}\psi - (A_{n}^{\alpha_{1}}\phi_{n}A_{n}^{\alpha_{2}}\phi_{n} + A_{n,1}^{\alpha}\phi_{n})\psi\right\}u \\ &- \left\{A_{n}^{\alpha_{2}}\psi + \psi A_{n}^{\alpha_{2}}\phi_{n}\right\}A_{n}^{\alpha_{1}}u - \left\{A_{n}^{\alpha_{1}}\psi + \psi A_{n}^{\alpha_{1}}\phi_{n}\right\}A_{n}^{\alpha_{2}}u + \psi A_{n,1}^{\alpha}u \\ &=: \Phi_{1} + \Phi_{2} + \Phi_{3} + \Phi_{4}. \end{aligned}$$

We then want to integrate the above terms along the curve ν against a test function $\varphi \in \mathcal{C}^2$. Recalling that the coefficients of the differential operators $A_n^{\alpha_j}$ have \mathcal{C}^r norm bounded by $\|(DF^n)^{-1}\|_{\mathcal{C}^r}$, we thus have

$$\int \varphi \Phi_1 \circ \nu \le C_{\sharp} \max_{i,j} \{ \|A_{n,1}^{\alpha}\psi\|_{\mathcal{C}^0_{\nu}}, \|\psi A_{n,1}^{\alpha}\phi_n\|_{\mathcal{C}^0_{\nu}}, \\ (1 + \|A_n^{\alpha_i}\phi_n\|_{\mathcal{C}^0_{\nu}})^2 \|\psi\|_{\mathcal{C}^0}, \|A_n^{\alpha_i}\phi_n A_n^{\alpha_j}\psi\|_{\mathcal{C}^0_{\nu}} \} \|u\|_0.$$

The bounds for Φ_2 and Φ_3 are similar:

$$\left| \int \varphi \Phi_2 \circ \nu \right| + \left| \int \varphi \Phi_3 \circ \nu \right| \le C_{\sharp} \| (DF^n)^{-1} \|_{\mathcal{C}^1_{\nu}} \max_i \{ \|A_n^{\alpha_i} \psi\|_{\mathcal{C}^1_{\nu}}, \|A_n^{\alpha_i} \phi_n\|_{\mathcal{C}^1_{\nu}} \|\psi\|_{\mathcal{C}^1} \} \|u\|_1.$$

Next, for any two vector $v, w \in \mathbb{R}^2$, $i, j \in \{1, 2\}$ and $x = (x_1, x_2) \in \mathbb{T}^2$,³⁷

$$\begin{aligned} \partial_{F^*v}(\partial_{F^*w}u) &= \partial_{F^*v}(\langle \nabla u, (DF)^{-1}w \rangle) = \langle \nabla(\langle \nabla u, (DF)^{-1}w \rangle), (DF)^{-1}v \rangle \\ &= \sum_{j,k} \partial_{x_j x_k}^2 u \cdot \left[(DF)^{-1}v \right]_k \left[(DF)^{-1}w \right]_j + \sum_{j,k} \partial_{x_k} u \, \partial_{x_j} \left[(DF)^{-1}w \right]_k \cdot \left[(DF)^{-1}v \right]_j. \end{aligned}$$

Thus,

$$\int \varphi \Phi_4 \circ \nu \le C_{\sharp} \{ \mu^{2n} \| \psi \|_{\mathcal{C}^2}, \mu^n \| (DF^n)^{-1} \|_{\mathcal{C}^2_{\nu}} \| \psi \|_{\mathcal{C}^1}, \| (DF^n)^{-1} \|_{\mathcal{C}^1_{\nu}}^2 \| \psi \|_{\mathcal{C}^1} \}^+ \| u \|_2.$$

It follows by the property of the C^r norm and (6.2), that

$$\int \varphi P_{2,n}^{\alpha}(\psi u) \circ \nu \leq C_{\sharp}\{1, \|A_{n,1}^{\alpha}\phi_n\|_{\mathcal{C}^0_{\nu}}, \max_{i\in\{1,2\}} \|A_n^{\alpha_i}\phi_n\|_{\mathcal{C}^0_{\nu}}^2, \\ \max_{i\in\{1,2\}} \|(DF^n)^{-1}\|_{\mathcal{C}^1_{\nu}} \|A_n^{\alpha_i}\phi_n\|_{\mathcal{C}^1_{\nu}}, \mu^n \|(DF^n)^{-1}\|_{\mathcal{C}^2_{\nu}}, \|(DF^n)^{-1}\|_{\mathcal{C}^1_{\nu}}^2\}^+ \|\psi\|_{\mathcal{C}^2_{\nu}} \|u\|_{2}.$$

We have thus proved that

$$\tilde{C}(2,n) = C_{\sharp} \{ 1, \|A_{n,1}^{\alpha}\phi_n\|_{\mathcal{C}^0_{\nu}}, \max_{i \in \{1,2\}} \|A_n^{\alpha_i}\phi_n\|_{\mathcal{C}^0_{\nu}}^2, \max_{i \in \{1,2\}} \|(DF^n)^{-1}\|_{\mathcal{C}^1_{\nu}} \|A_n^{\alpha_i}\phi_n\|_{\mathcal{C}^1_{\nu}}, \\ \mu^n \|(DF^n)^{-1}\|_{\mathcal{C}^2_{\nu}}, \|(DF^n)^{-1}\|_{\mathcal{C}^1_{\nu}}^2 \}^+.$$

To conclude we need a bound of the above quantity. It is enough to find estimates for $\|(A_{n,1}^{\alpha}\phi_n)\|_{\mathcal{C}^0_{\mu}}$ and $\|A_n^{\alpha_1}\phi_n\|_{\mathcal{C}^1_{\nu}} \cdot \|(DF^n)^{-1}\|_{\mathcal{C}^1_{\nu}}$, the other quantities being already estimated in Proposition 5.9. First, we can use formulae (6.19) and (5.21),

(6.23)
$$|\partial_{F^{n^*}e_{\ell}}\phi_n(x)| \le C_{\sharp} \sum_{j=0}^{n-1} \mu^{n-j} \le C_{\mu,n} \mu^n.$$

 $^{^{36}}$ We use the following notation: Φ_1 equals the third line from the bottom, the other Φ_i are, ordered, the terms in the second line from the bottom. ³⁷Here we denote $[(DF)^{-1}w]_k := \langle (DF)^{-1}w, e_k \rangle$.

In particular $||A_n^{\alpha_1}\phi_n||_{\mathcal{C}^0_{\nu}} \leq C_{\mu,n}\mu^n$. Next, we take another derivative of (6.19) in the direction of $F^{n^*}e_q$ and, setting $g_{\ell,n,j}(x) = \langle \nabla \phi_1, (DF^{n-j})^{-1}e_\ell \rangle(x)$, we have

(6.24)
$$\partial_{F^{n^*}e_q}(\partial_{F^{n^*}e_\ell}\phi_n(x)) = \sum_{j=0}^{n-1} \langle \nabla(g_{\ell,n,j} \circ F^j(x)), (D_x F^n)^{-1}e_q \rangle$$
$$= \sum_{j=0}^{n-1} \langle (D_x F^j)^* \nabla g_{\ell,n,j} \circ F^j(x), (D_x F^n)^{-1}e_q \rangle$$
$$= \sum_{j=0}^{n-1} \langle \nabla g_{\ell,n,j} \circ F^j(x), (D_x F^{n-j})^{-1}e_q \rangle.$$

By a direct computation we see that, recalling (5.23) and (5.25),

$$\|\nabla g_{\ell,n,j}\| \le C_{\sharp} \max_{i} \{ \|\partial_{x_{i}} (DF^{n-j})^{-1}\| \} \le C_{\mu,n-j} (1 + C_{F} \mu^{n-j} \lambda_{n-j}^{+}) \mu^{n-j}.$$

We use this in (6.24) obtaining

$$\|A_{n,1}^{\alpha}\phi_n\|_{\mathcal{C}^0_{\nu}} \le \sup_{q,\ell} \|\partial_{F^{n^*}e_q}(\partial_{F^{n^*}e_\ell}\phi_n)\|_{\mathcal{C}^0_{\nu}} \le C^2_{\mu,n}(1+C_F\mu^n\lambda_n^+)\mu^{2n}.$$

Finally, we use (6.19) to compute

$$\begin{aligned} \left| \frac{d}{dt} (A_n^{\alpha_1} \phi_n \circ \nu) \right| &\leq \sum_{j=0}^{n-1} \left| \langle D(\nabla \phi_1) \circ F^j(D_\nu F^j) \nu', (DF^{n-j})^{-1} \circ F^j \circ \nu e_{\alpha_1} \rangle \right| \\ &+ \left| \langle \nabla \phi_1 \circ (F^j \nu), [(DF^{n-j})^{-1} \circ (F^j \nu)]' e_{\alpha_1} \rangle \right| \\ &\leq C_{\mu,n} \lambda_m^+ \circ (F^{n-m} \nu) \mu^n + C_{\mu,n} \mu^n (1 + C_F \lambda_m^+ \circ (F^{n-m} \nu)) \\ &\leq C_{\mu,n} \lambda_m^+ \circ (F^{n-m} \nu) \mu^n, \end{aligned}$$

where we have used (5.21) and that, recalling (5.19), $\varsigma_{n,m} \leq C_{\mu,n}(1+C_F\lambda_m^+)$. We thus obtain

$$\|A_n^{\alpha_1}\phi_n\|_{\mathcal{C}^1_{\nu}} \cdot \|(DF^n)^{-1}\|_{\mathcal{C}^1_{\nu}} \le C_{\mu,n}\mu^{3n} \sup_{\zeta \in \operatorname{supp}(\varphi) \cap \nu(\mathbb{T})} [(1+C_F\lambda_m^+)\lambda_m^+] \circ F^{n-m}(\zeta).$$

The proposition follows collecting all the above estimates and recalling again (5.21) for the estimate of $||(DF^n)^{-1}||_{\mathcal{C}^1_{\nu}}, ||(DF^n)^{-1}||_{\mathcal{C}^2_{\nu}}$ and noticing that $\varsigma_{n,m} \leq C_{\mu,n}(1+C_F\lambda_m^+)$.

6.3. Conclusion of the first Lasota-Yorke inequality.

This section is devoted first to the proof of Theorem 6.1 and then is concluded by a useful Corollary.

Proof of Theorem 6.1. Given Lemma 5.18, the proof of Theorem 6.1 is almost exactly the same as in [32], hence we provide the full proof for $\rho = 0, 1$ and give a sketched proof for the case $1 < \rho \leq r - 1$. Let us prove (6.3) first, since it is an immediate consequence of Lemma 5.18 and Definition 6.1 in the case $\rho = 0$. Indeed, by changing the variables and recalling the notation of Section 5.3 and Lemma 5.18, we have,

$$\begin{split} &\int_{\mathbb{T}} \phi(t) \mathcal{L}^{n} u(\gamma(t)) dt = \sum_{\nu \in F^{-n} \gamma} \int_{\mathbb{T}} |\det D_{\nu(t)} F^{n}|^{-1} \cdot (u \circ \nu)(t) \cdot \phi(t) dt \\ &= \sum_{\nu \in F^{-n} \gamma} \int_{\mathbb{T}} |\det D_{\hat{\nu}} F^{n}|^{-1} \cdot (u \circ \hat{\nu})(t) \cdot (\phi \circ h_{n})(t) h_{n}'(t) dt \\ &\leq \sum_{\nu \in F^{-n} \gamma} \left\| h_{n}' \left| \det D_{\hat{\nu}} F^{n} \right|^{-1} \right\|_{\mathcal{C}^{0}} \|u\|_{0} \leq (1 + C_{F} \mathfrak{c}) C_{\mu,n} \|\phi\|_{\mathcal{C}^{0}(\mathbb{T})} \|u\|_{0}. \end{split}$$

Let us now proceed with the case $\rho = 1$, from which we deduce the general case by similar computations. We must bound the quantity, for all $\|\phi\|_{\mathcal{C}^1} \leq 1$,

$$\int_{\mathbb{T}} \phi(t)(\partial_{v}\mathcal{L}^{n}u)(\gamma(t))dt = \int_{\mathbb{T}} \phi(t)\langle \nabla(\mathcal{L}^{n}u)(\gamma(t)), v\rangle dt$$

where now $\phi \in \mathcal{C}^1(\mathbb{T})$ with norm one and v is a unitary \mathcal{C}^r vector field. From Proposition 6.2 the above quantity is equal to

(6.25)
$$\sum_{\nu \in F^{-n}\gamma} \int_{\mathbb{T}} \frac{1}{|\det D_{\nu}F^{n}|} \phi \cdot \partial_{F^{n^{*}}v} u(\nu) + \int_{\mathbb{T}} \mathcal{L}^{n}(Q_{n,0}u)\phi,$$

where $Q_{n,0}$ is an operator of multiplication by a \mathcal{C}^1 function.

By Proposition 6.4 applied with $\psi = 1$, plus the result for $\rho = 0$, the last term is then bounded by $C_n \|u\|_1.$

In order to bound the first term of (6.25) we need an analogous of Lemma 6.5 in [32]. The idea is to decompose the vector field v into a vector tangent to the central curve γ and a vector field approximately in the unstable direction so that the first one can be integrated by parts, while for the other we can exploit the expansion. The proof of the following Lemma follows that of the aforementioned paper, since the key point is the splitting of the tangent space in two directions, one of which is expanding. Once more, however, the presence of the central direction creates difficulties. We give the proof adapted to our case in Appendix D.

Lemma 6.6. Let \bar{n} be the integer provided by Lemma 5.10. For every $n > \bar{n}, \gamma \in \Gamma_{\rho}(c), \rho \leq r$, $\nu \in F^{-n}\gamma$, and any vector field $v \in \mathcal{C}^{\rho}$, with $\|v\|_{\mathcal{C}^{\rho}} \leq 1$, defined in some neighborhood $M(\gamma)$ of γ , there exist a neighborhood $M'(\gamma)$ of γ and a decomposition

$$(6.26) v = \hat{v}^c + \hat{v}^u,$$

where \hat{v}^c and \hat{v}^u are $\mathcal{C}^{\rho}(M'(\gamma))$ vector fields such that, setting $F^n(N(\nu)) = M'(\gamma)$,³⁸

- $\hat{v}^{c}(\gamma(t)) = g(t)\gamma'(t), \text{ where } g \in \mathcal{C}^{r} \text{ and } \|g\|_{\mathcal{C}^{\rho}} \leq C_{\sharp} \mathfrak{C}^{\rho!} \mathcal{C}^{a_{\rho}}_{\mu,n} \mu^{\rho!n},$ $\|(F^{n})^{*} \hat{v}^{u}\|_{\mathcal{C}^{\rho}(N(\nu))} \leq \lambda_{-}^{-n} \mathfrak{C}^{\rho!\rho} \mathcal{C}^{\rho a_{\rho}}_{\mu,n} \mu^{\rho!n},$ $\|(F^{n})^{*} \hat{v}^{c}\|_{\mathcal{C}^{\rho}(N(\nu))} \leq \mathfrak{C}^{\rho!\rho} \mathcal{C}^{2\rho a_{\rho}+1}_{\mu,n} \mu^{[(\rho+1)(\rho!+1)+\rho!]n},$ $\|\hat{v}^{u}\|_{\mathcal{C}^{\rho}(M'(\gamma))} + \|\hat{v}^{c}\|_{\mathcal{C}^{\rho}(M'(\gamma))} \leq C_{n}.$

By the above decomposition, the addends in the first term in (6.25) become

(6.27)
$$\int \frac{1}{|\det D_{\nu}F^n|} \phi \cdot \partial_{F^{n^*}\hat{v}^c} u(\nu) + \int \frac{1}{|\det D_{\nu}F^n|} \phi \cdot \partial_{F^{n^*}\hat{v}_u} u(\nu).$$

Since $\gamma(t) = F^n \nu(t)$ we have $g(t) D_{\nu(t)} F^n \cdot \nu'(t) = \hat{v}^c(F^n \nu(t))$, hence:

$$g(t)\nu'(t) = (D_{\nu(t)}F^n)^{-1} \cdot \hat{v}^c(F^n\nu(t)) = F^{n^*}\hat{v}^c(\nu(t)),$$

hence $|g'| \leq ||F^{n^*} \hat{v}^c||_{\mathcal{C}^1}$. Accordingly,

$$\begin{split} &\int \frac{\phi(t)}{|\det D_{\nu(t)}F^{n}|} \partial_{F^{n^{*}}\hat{v}^{c}} u(\nu(t)) dt = \int \frac{g(t)\phi(t)}{|\det D_{\nu(t)}F^{n}|} \frac{d}{dt} (u(\nu(t))) dt \\ &= \int \frac{g(t)\phi(t)}{|\det D_{\hat{\nu}\circ h_{n}^{-1}(t)}F^{n}|} \left[\frac{d}{dt} (u \circ \hat{\nu}) \right] \circ h_{n}^{-1}(t) \left[h_{n}^{-1}(t) \right]' dt \\ &= \int \frac{[g\phi] \circ h_{n}(t)}{|\det D_{\hat{\nu}(t)}F^{n}|} (u \circ \hat{\nu})'(t) dt = -\int \frac{d}{dt} \left(\frac{[g\phi] \circ h_{n}(t)}{|\det D_{\hat{\nu}(t)}F^{n}|} \right) u(\hat{\nu}(t)) dt \\ &\leq \left\| \frac{[g\phi] \circ h_{n}}{\det D_{\hat{\nu}}F^{n}} \right\|_{\mathcal{C}^{1}} \|u\|_{0}. \end{split}$$

 $^{^{38}\}text{The constants}~a_{\rho}$ are defined in Lemma 5.10.

Summing over $\nu \in F^{-n}\gamma$ and using Lemma 5.18 we obtain

(6.28)
$$\sum_{\nu \in F^{-n}\gamma} \int \phi(t) \partial_{F^{n^*}\hat{v}^c} u(\nu(t)) dt \lesssim C^5_{\mu,n} \mu^{7n} (1 + C_F \mathfrak{c})^2 \|u\|_{0}.$$

The second term of (6.27) is

(6.29)
$$\int \frac{\phi}{|\det D_{\nu}F^{n}|} \partial_{F^{n^{*}}\hat{v}^{u}} u(\nu) = \int \frac{\phi}{|\det D_{\nu}F^{n}|} \langle \nabla u, F^{n^{*}}\hat{v}^{u} \rangle \circ \nu$$
$$\leq C_{\sharp} \left\| \frac{\phi \circ h_{n}h_{n}'}{\det D_{\hat{\nu}}F^{n}} \right\|_{\mathcal{C}^{1}} \|F^{n^{*}}\hat{v}^{u} \circ \hat{\nu}\|_{\mathcal{C}^{1}} \|u\|_{1}$$
$$\leq C_{\sharp} \|h_{n}\|_{\mathcal{C}^{1}} \left\| \frac{h_{n}'}{\det D_{\hat{\nu}}F^{n}} \right\|_{\mathcal{C}^{1}} \varepsilon \lambda_{-}^{-n} C_{\mu,n} \mu^{n} \|u\|_{1},$$

where we made the usual change of variables $t = h_n(s)$ and used Lemma 6.6. Finally, using (6.28) and (6.29) in (6.27), and recalling (5.29), we have by Lemma 5.18, with $\rho = 1$,

(6.30)
$$\|\mathcal{L}^{n}u\|_{1} \leq \varepsilon \lambda_{-}^{-n} C_{\mu,n}^{3} \mu^{2n} (1+C_{F}\varepsilon) \|u\|_{1} + C_{n} \|u\|_{0}$$

For the general case $1 \le \rho \le r-1$ one has to control the term $\int_{\mathbb{T}} \phi(t) \partial_{v_s} \cdots \partial_{v_1} \mathcal{L}^n u(\nu(t)) dt$, for vector fields $v_j \in \mathcal{C}^{\rho}, j = 1, ..., s$ and $s \le \rho$. Using again Propositions 6.2 and 6.4, the latter is bounded by

(6.31)
$$\sum_{\nu \in F^{-n}\gamma} \int \frac{1}{|\det D_{\nu}F^{n}|} \phi \cdot \partial_{F^{n^{*}}v_{s}\cdots F^{n^{*}}v_{1}} u(\nu) + C_{n,\rho} ||u||_{\rho-1}.$$

Now the strategy is exactly the same as before. We use Lemma 6.6 to decompose each $v_j = \hat{v}_j^u + \hat{v}_j^c$. We take $\sigma \in \{u, c\}^s$, $k = \#\{i | \sigma_i = c\}$ and let π be a permutation of $\{1, \ldots, s\}$ such that $\pi\{1, \ldots, k\} = \{i | \sigma_i = c\}$. Using integration by parts, we can write the integral in (6.31) as

$$\int \frac{\phi}{\det D_{\nu}F^{n}} \partial_{F^{n^{*}}v_{s}} \dots \partial_{F^{n^{*}}v_{1}} u(\nu) = \sum_{\sigma \in \{u,c\}^{s}} \int \frac{\phi}{\det D_{\nu}F^{n}} \left(\prod_{s=1}^{1} \partial_{F^{n^{*}}\hat{v}_{i}^{\sigma_{i}}}\right) u(\nu)$$

$$= \sum_{\sigma \in \{u,c\}^{s}} \int \frac{\phi}{\det D_{\nu}F^{s}} \prod_{i=s}^{k} \partial_{F^{n^{*}}\hat{v}_{\pi(i)}^{c}} \prod_{i=k+1}^{1} \partial_{F^{n^{*}}\hat{v}_{\pi(i)}^{u}} u(\nu) + C_{n,\rho} \|u\|_{\rho-1}$$

$$= \sum_{\sigma \in \{u,c\}^{s}} (-1)^{k} \int \prod_{i=k+1}^{s} \partial_{F^{n^{*}}\hat{v}_{\pi(i)}^{u}} u(\nu) \prod_{i=k}^{1} \partial_{F^{n^{*}}\hat{v}_{\pi(i)}^{c}} \left(\frac{\phi}{\det D_{\nu(t)}F^{n}}\right) + C_{n,\rho} \|u\|_{\rho-1}$$

By Lemma 6.6, $\|F^{n^*}\hat{v}^c_{\pi(i)}\|_{\mathcal{C}^{\rho}(\nu)} \leq \mathbb{C}^{\rho!\rho}C^{2\rho a_{\rho}+1}_{\mu,n}\mu^{(\rho+1)(\rho!+1)+\rho!}$ while

$$\|\prod_{i=k+1}^{s} F^{n^{*}} \hat{v}_{\pi(i)}^{u}\|_{\mathcal{C}^{\rho}(\nu)} \leq \mathbb{C}^{\rho!\rho} C \lambda_{-}^{-(s-k)n} (C_{\mu,n}^{\rho a_{\rho}} \mu^{\rho \rho!n})^{s-k} d\mu^{\rho} d\mu^$$

It follows by Lemma 5.18, equation (5.29) and the fact that $\|\phi\|_{\mathcal{C}^r} \leq 1$, that³⁹

$$\sum_{\nu \in F^{-n}\gamma} \int \frac{\phi}{\det DF^{n}} \partial_{F^{n*}v_{1}} \dots \partial_{F^{n*}v_{\rho}} u \circ \nu$$

$$\leq \mathfrak{e}^{2\rho!\rho} \lambda_{-}^{-\rho n} C_{\mu,n}^{\rho^{2}a_{\rho}} \mu^{\rho^{2}\rho!n} \|h_{n}\|_{\mathcal{C}^{\rho}} \sum_{\nu \in F^{-n}\gamma} \left\| \frac{h'_{n}}{\det D_{\hat{\nu}}F^{n}} \right\|_{\mathcal{C}^{\rho}} \|u\|_{\rho} + C_{n,\rho} \|u\|_{\rho-1}$$

$$\leq \mathfrak{e}^{\rho!(2\rho+1)} (1 + C_{F} \mathfrak{e}^{\rho!})^{\bar{c}_{\rho}} \lambda_{-}^{-\rho n} C_{\mu,n}^{2\rho^{2}a_{\rho} + \tilde{a}_{\rho}} \mu^{[(\rho!+1)(\rho^{2}+1) + \tilde{b}_{\rho}]n} \|u\|_{\rho} + C_{n,\rho} \|u\|_{\rho-1}$$

hence, recalling the definitions of \tilde{a}, \tilde{b} in Lemma (5.18), we have (6.4) with $\bar{a}_{\rho} = 1 + 2a_{\rho}\rho^2 + \rho$, $\bar{b}_{\rho} = (\rho! + 1)(\rho^2 + 2)$ and $\bar{c}_{\rho} = 1 + \{\rho + 1, \rho(\rho + 1)/2\}^+$

³⁹Notice that the coefficient in front of the strong norm is obtained in the case $s = \rho$ and k = 0, while all the other terms are bounded again by $C_{n,\rho} ||u||_{\rho-1}$.

The last result of this section is the following Corollary of Theorem 6.1.

Corollary 6.7. Let us assume that, for every integer $0 \le \rho \le r - 1$,⁴⁰

(6.32)
$$\lambda_{-}^{-1}\mu^{6(\rho+1)!} < 1.$$

Let $\delta_* \in (\lambda_-^{-1}\mu^{6(\rho+1)!}, 1)$. Then, for each $n \in \mathbb{N}$,

(6.33)
$$\|\mathcal{L}^{n}u\|_{\rho} \leq C_{\sharp}\delta_{*}^{n}\|u\|_{\rho} + C_{\mu,n}\mu^{n}\|u\|_{0}$$

Proof. The strategy is to take the integer \bar{n} large enough such that the coefficient of the strong norm in (6.4) is smaller than 1, and then to iterate the estimates. First, recalling (5.30), we can estimate⁴¹ $c_{\rho,F} \leq C_{\sharp} \mu^{\frac{3}{2}\rho![\rho^2+5\rho+6]\bar{n}}$. Hence, a direct computation yields

$$\mathbb{C}_{\rho,F}\mu^{\bar{b}_{\rho}\bar{n}} \leq C_{\sharp}\mu^{[6(\rho+1)!\rho+13]\bar{n}}$$

Hence, by (6.32), we can choose $\delta \in (\lambda_{-}^{-1}\mu^{6(\rho+1)!}, 1)$. Accordingly, since $C_{\mu,n}$ grows only linearly in n, we can choose $\bar{n} \in \mathbb{N}$ large enough such that $\mathfrak{c}_{\rho,F}C_{\mu,\bar{n}}^{\bar{n}}\mu^{\bar{b}_{\rho}\bar{n}}\lambda_{-}^{-\rho\bar{n}} < \delta^{\rho\bar{n}}$ for every $\rho \in [1, r-1]$. Let us proceed by induction on ρ . For $\rho = 0$ the statement is implied by (6.3). Let us assume it true for each integer smaller then or equal to $\rho - 1$. By Theorem 6.1 and (6.32), we have

(6.34)
$$\|\mathcal{L}^{\bar{n}}u\|_{\rho} \le C_{\sharp} \delta^{\rho \bar{n}} \|u\|_{\rho} + C_{\bar{n}} \|u\|_{\rho-1}.$$

For every $m \in \mathbb{N}$ we write $m = \bar{n}q + r$, $0 \leq r < \bar{n}$, and iterate (6.34) to have

$$\|\mathcal{L}^{m}u\|_{\rho} = \|\mathcal{L}^{\bar{n}}(\mathcal{L}^{m-\bar{n}}u)\|_{\rho} \leq C_{\sharp}\delta^{\rho\bar{n}}\|\mathcal{L}^{m-\bar{n}}u\|_{\rho} + C_{\bar{n}}\|\mathcal{L}^{m-\bar{n}}u\|_{\rho-1} \leq \cdots$$

$$\cdots \leq C_{\sharp}\delta^{q\rho\bar{n}}\|\mathcal{L}^{r}u\|_{\rho} + C_{\sharp}\sum_{k=0}^{q-1}\delta^{k\rho\bar{n}}\|\mathcal{L}^{m-(k+1)\bar{n}}u\|_{\rho-1} \leq C_{\sharp}\delta^{\rho\bar{n}}\|u\|_{\rho} + C_{\mu,m}\mu^{m}\|u\|_{\rho-1},$$

where we used $\|\mathcal{L}^{m-(k+1)\bar{n}}u\|_{\rho-1} \leq C_{\mu,m}\mu^{m-(k+1)}\|u\|_{\rho-1}$ by the inductive assumption. We iterate the last inequality ρ times and obtain

$$\begin{aligned} \|\mathcal{L}^{\rho m}u\|_{\rho} &\leq C^{\rho-1}_{\mu,m}(\mu^{\rho-1}\delta^{\rho})^{m} \|u\|_{\rho} + C^{\rho}_{\mu,m}\mu^{\rho m} \|u\|_{0} \\ &\leq C^{\rho}_{\mu,m}(\mu^{\rho}\delta^{\rho})^{m} \|u\|_{\rho} + C^{\rho}_{\mu,m}\mu^{\rho m} \|u\|_{0}. \end{aligned}$$

We then consider the above inequality for m such that $\rho m = \bar{n}$, so that $C^{\rho}_{\mu,m}(\mu^{\rho}\delta^{\rho})^m < \tilde{\delta}^{\bar{n}}$, for some $\tilde{\delta} \in (\delta, 1)$. Hence,

(6.35)
$$\|\mathcal{L}^{\bar{n}}u\|_{\rho} \le \tilde{\delta}^{\bar{n}} \|u\|_{\rho} + C_{\mu,\bar{n}}\mu^{\bar{n}} \|u\|_{0}.$$

Finally, we iterate once again (6.35) and we obtain the result for some $\delta_* \in (\tilde{\delta}, 1)$.

Remark 6.8. Although Corollary 6.7 provides a Lasota-Yorke inequality, a fundamental ingredient is missing: the embedding of \mathcal{B}_{ρ} in \mathcal{B}_{0} is not compact.

7. A second Lasota-Yorke inequality: preliminaries

The main result of the following two sections is the second step towards the proof of Theorem 4.4, namely a Lasota-Yorke type inequality between the Hilbert space \mathcal{H}^s and \mathcal{B}_{ρ}^{42} . We will see in Corollary 9.2 that this solves the compactness problem mentioned in Remark 6.8. First we state some results on the \mathcal{H}^s -norm of the transfer operator.

 $^{^{40}}$ Note that the following is implied by **(H3)**.

 $^{^{41}\}mathrm{As}$ previously mentioned, all the estimates for the exponent of μ are far from being optimal.

⁴²See Appendix E for definitions and the needed properties of $\mathcal{H}^{s}(\mathbb{T}^{2})$.

7.1. \mathcal{H}^s -norm of \mathcal{L} .

Lemma 7.1. Let $F \in C^r(\mathbb{T}^2, \mathbb{T}^2)$ satisfying (H1). For each $n \in \mathbb{N}$ and $1 \leq s \leq r$, there exist $A_s, Q(n,s) > 0$ such that, for every $u \in \mathcal{H}^s(\mathbb{T}^2, \mathbb{R})$,

(7.1)
$$\|\mathcal{L}^n u\|_{L^2} \le \|\mathcal{L}^n 1\|_{\infty}^{\frac{1}{2}} \|u\|_{L^2}$$

(7.2)
$$\|\mathcal{L}^{n}u\|_{\mathcal{H}^{s}}^{2} \leq A_{s}\mu^{2sn}\|\mathcal{L}^{n}1\|_{\infty}\|u\|_{\mathcal{H}^{s}}^{2} + Q(n,s)\|u\|_{\mathcal{H}^{s-1}}^{2},$$

where

(7.3)
$$Q(n,1) \le C_{\mu,n}^3 \mu^{3n} (1 + C_F \mathfrak{c}).$$

Proof. First of all notice that

(7.4)
$$\|\mathcal{L}^{n}u\|_{L^{2}}^{2} \leq \|u\|_{L^{2}} \left(\int (\mathcal{L}^{n}u \circ F^{n})^{2}\right)^{\frac{1}{2}} \leq \|u\|_{L^{2}} \left(\int (\mathcal{L}^{n}u)^{2}\mathcal{L}^{n}1\right)^{\frac{1}{2}} \\ \leq \|u\|_{L^{2}} \|\mathcal{L}^{n}1\|_{\infty}^{\frac{1}{2}} \|\mathcal{L}^{n}u\|_{L^{2}},$$

hence (7.1). Next, by (6.13) and (6.14) we have, for each $v_i \in \{e_1, e_2\}$,

(7.5)
$$\begin{aligned} \|\partial_{v_s} \cdots \partial_{v_1} \mathcal{L}^n u\|_{L^2}^2 &\leq \|\mathcal{L}^n (\partial_{F^{n*}v_s} \cdots \partial_{F^{n*}v_1}) u\|_{L^2}^2 \\ &+ \sum_{s=1}^s \|\mathcal{L}^n (A^{\alpha}_{n,k} ((A^{\alpha}_n \phi_n) \cdot P^{\alpha}_{k-1} u))\|_{L^2}^2. \end{aligned}$$

$$+\sum_{k=1} \|\mathcal{L}(A_{n,k}((A_{n},\phi_{n}),F_{k-1}))\| \leq C_{k-1}$$

Let us analyze the first term above when s = 2. Notice that

$$\partial_{F^{n*}v_2}(\partial_{F^{n*}v_1}u) = \langle \nabla \left(\langle \nabla u, (DF^n)^{-1}v_1 \rangle \right), (DF^n)^{-1}v_2 \rangle = \langle (DF^n)^{-1}v_1 D^2 u, (DF^n)^{-1}v_2 \rangle + \langle D((DF^n)^{-1}v_1) \nabla u, (DF^n)^{-1}v_2 \rangle.$$

where $D^2 f$ indicates the Hessian of a function f and D(V) is the Jacobian of the vector field V. The term with higher derivatives of u has coefficients bounded by $||(DF^n)^{-1}||^2$, while the other term is a differential operator of order one applied to u. In the general case we have some $\overline{P}_{s-1,\rho}$ such that

(7.6)
$$\begin{aligned} |\mathcal{L}^{n}(\partial_{F^{n*}v_{s}}\cdots\partial_{F^{n*}v_{1}})u| &\leq \|(DF^{n})^{-1}\|^{s} \sup_{w_{1},\cdots,w_{s}\in\{e_{1},e_{2}\}} \mathcal{L}^{n}(|\partial_{w_{s}}\cdots\partial_{w_{1}}u|) \\ &+ |\mathcal{L}^{n}\overline{P}_{s-1,\rho}u|. \end{aligned}$$

Hence, by (7.1), (E.5) and (5.17), there exists a constant $C_1(n, s)$ such that

(7.7)
$$\|\mathcal{L}^{n}(\partial_{F^{n*}v_{s}}\cdots\partial_{F^{n*}v_{1}})u\|_{L^{2}}^{2} \leq C_{\sharp}\|\mathcal{L}^{n}1\|_{\infty}\mu^{2sn}\|u\|_{\mathcal{H}^{s}}^{2} + C_{1}(n,s)\|u\|_{\mathcal{H}^{s-1}}^{2}.$$

Similarly there exists $C_2(s, n)$ such that

(7.8)
$$\sum_{k=1}^{\iota} \|\mathcal{L}^n(A_{n,k}^{\alpha}((A_n^{\alpha_k}\phi_n) \cdot P_{n,k-1}^{\alpha}u))\|_{L^2}^2 \le C_2(n,s) \|u\|_{\mathcal{H}^{s-1}}^2.$$

By (7.5), (7.7) and (7.8) we obtain

$$\|\mathcal{L}^{n}(\partial_{F^{n*}v_{1}}\cdots\partial_{F^{n*}v_{s}})u\|_{L^{2}}^{2} \leq C_{\sharp}\|\mathcal{L}^{n}1\|_{\infty}\mu^{2sn}\|u\|_{\mathcal{H}^{s}}^{2} + Q(n,s)\|u\|_{\mathcal{H}^{s-1}}^{2}$$

It remains to prove that in the case s = 1 we have an explicit bound on Q(n, 1). Recall that by (6.9) and (7.1) we have, for any $v \in \{e_1, e_2\}$,

(7.9)
$$\|\langle \nabla \mathcal{L}^{n} u, v \rangle \|_{L^{2}} \leq \|\mathcal{L}^{n} \langle \nabla u, (DF^{n})^{-1} v \rangle \|_{L^{2}} + \|\mathcal{L}^{n} (\langle \nabla \phi_{n}, (DF^{n})^{-1} v \rangle u)\|_{L^{2}},$$
$$\leq \|\mathcal{L}^{n} 1\|_{\infty}^{\frac{1}{2}} \left(\|\langle \nabla u, (DF^{n})^{-1} v \rangle \|_{L^{2}} + \|\langle \nabla \phi_{n}, (DF^{n})^{-1} v \rangle u\|_{L^{2}} \right).$$

A bound for the first term is straightforward, since by (5.17)

(7.10)
$$\| \langle \nabla u, (DF^n)^{-1} v \rangle \|_{L^2} \le C_{\sharp} \mu^n \| \nabla u \|_{L^2}.$$
For the second term we use formula (6.19) and we have

(7.11)
$$\begin{aligned} \|\langle \nabla \phi_n, (DF^n)^{-1}v \rangle u\|_{L^2} &\leq \sum_{j=0}^{n-1} \|\langle \nabla \phi_1 \circ F^j(x), (D_{F^j x} F^{n-j})^{-1}v \rangle\|_{\infty} \|u\|_{L^2} \\ &\leq C_{\sharp} \sum_{j=0}^{n-1} \mu^{n-j} \|u\|_{L^2} \leq C_{\mu,n} \mu^n \|u\|_{L^2}, \end{aligned}$$

By (7.9), (7.10), (7.11) and (5.58) we obtain (7.3).

7.2. Transversality.

In this Section we investigate the quantities $\mathcal{N}_F, \widetilde{\mathcal{N}}_F$ defined in section 4.1. Recall that

(7.12)
$$\mathcal{N}_{F}(n) = \sup_{y \in \mathbb{T}^{2}} \sup_{z_{1} \in F^{-n}(y)} \mathcal{N}_{F}(n, y, z_{1})$$
$$\widetilde{\mathcal{N}}_{F}(n) = \sup_{y \in \mathbb{T}^{2}} \sup_{L} \widetilde{\mathcal{N}}_{F}(n, y, L).$$

Both \mathcal{N}_F and $\widetilde{\mathcal{N}}_F$ depend on the map F, however in the following we will often drop the F dependence to ease notation. An important advantage of $\widetilde{\mathcal{N}}$ over \mathcal{N} is the following

Proposition 7.2. $\widetilde{\mathcal{N}}(n)$ is sub-multiplicative, i.e $\widetilde{\mathcal{N}}(n+m) \leq \widetilde{\mathcal{N}}(n)\widetilde{\mathcal{N}}(m)$, for every $n, m \in \mathbb{N}$.

Proof. For any $y \in \mathbb{T}^2$, and line L we have

$$\widetilde{\mathcal{N}}(y,L,n+m) = \sum_{\substack{z \in F^{-n-m}(y) \\ DF^{n+m}(z)\mathbf{C}_u \supset L}} |\det DF^{n+m}(z)|^{-1}$$

$$= \sum_{\substack{\hat{z} \in F^{-n}(y) \\ DF^{n}(\hat{z})\mathbf{C}_u \supset L}} \sum_{\substack{z \in F^{-m}(\hat{z}) \\ DF^{m}(z)\mathbf{C}_u \supset (DF^{n}(\hat{z}))^{-1}L}} \frac{1}{|\det DF^{m}(z)\det DF^{n}(\hat{z})|}$$

$$\leq \sum_{\substack{\hat{z} \in F^{-n}(y) \\ DF^{n}(\hat{z})\mathbf{C}_u \supset L}} \frac{1}{|\det DF^{n}(\hat{z})|} \sup_{\hat{z} = L'} \sum_{\substack{z \in F^{-m}(\hat{z}) \\ DF^{m}(z)\mathbf{C}_u \supset L'}} \frac{1}{|\det DF^{m}(z)|}.$$

Taking the sup over $y \in \mathbb{T}^2$ and L we get the claim. \Box

Remark 7.3. The above Proposition, in spite of its simplicity, turns out to be pivotal. The submultiplicativity of the sequence $\widetilde{\mathcal{N}}(n)$ implies the existence of $\lim_{n\to\infty} \widetilde{\mathcal{N}}(n)^{\frac{1}{n}}$. Also, an estimate of $\widetilde{\mathcal{N}}(n_0)$ for some $n_0 \in \mathbb{N}$ yields an estimate for all $n \in \mathbb{N}$.

The result below, inspired by [14], provides the relation between \mathcal{N} and $\widetilde{\mathcal{N}}$.

Lemma 7.4. Let
$$\alpha = \frac{\log(\lambda - \mu^{-2})}{\log(\lambda_+)} \in (0, 1)$$
 and $m_0 = m_0(n) = \lceil \alpha n \rceil$ we have, for all $n \in \mathbb{N}$
$$\mathcal{N}(n)^{\frac{1}{n}} \leq \|\mathcal{L}^{n-m_0}1\|_{\infty}^{\frac{1}{n}} \left(\widetilde{\mathcal{N}}(m_0)^{\frac{1}{m_0}}\right)^{\alpha}.$$

Proof. Given $y \in \mathbb{T}^2$, we consider $z_1, z_2 \in F^{-n}(y)$ such that $D_{z_1}F^n\mathbf{C}_u \cap D_{z_2}F^n\mathbf{C}_u \neq \{0\}$ and the line $L := L(z_1) := D_{z_1}F^n(\mathbb{R} \times \{0\}).$

Let $v_{\pm} = (1, \pm \chi_u) \in \mathbf{C}_u$ and $\theta_n := \max_{\pm} \measuredangle (D_{z_1} F^n e_1, D_{z_1} F^n v_{\pm})$. Notice that, for n = 0, $|\cos \theta_0|^{-1} = |\cos(\arctan(\chi_u))|^{-1} = \sqrt{1 + \chi_u^2} =: a_0 \le 2$. In fact, by the invariance of the unstable cone $|\cos \theta_n|^{-1} \le a_0$, for all $n \in \mathbb{N}$. On the other hand, by formula (5.15), Proposition 5.6 and condition (2.6) we have

(7.13)
$$\begin{aligned} |\tan \theta_n| &\leq a_0 |\sin \theta_n| = a_0 |\sin \measuredangle (e_1, v_{\pm})| \frac{|\det D_{z_1} F^n| || v_{\pm} ||}{||D_{z_1} F^n e_1|| ||D_{z_1} F^n v_{\pm}|} \\ &\leq |\sin(\arctan \chi_u)| a_0 C_* C_*^2 \mu_+^n \lambda_-^{-n} \\ &= a_0 C_* C_*^2 \chi_u \mu_+^n \lambda_-^{-n}, \end{aligned}$$

where we have used that $\sin(\arctan x) = x(\sqrt{1+x^2})^{-1}$.

Next, note that in the projective space \mathbb{RP}^2 the cones $D_{z_1}F^n\mathbf{C}_u$ and $D_{z_2}F^n\mathbf{C}_u$ are canonically identified with two intervals $I_1 = [a_1, b_1]$ and $I_2 = [a_2, b_2]$, respectively. While the line is a point that we also denote by L. From the assumption on the cones, and (7.13), we have that the projective distance between L and each one of the extremal points of I_2 is bounded by

(7.14)
$$\min\{\operatorname{dist}(L, a_2), \operatorname{dist}(L, b_2)\} \le a_0 C_* C_*^2 \chi_u \lambda_-^{-n} \mu_+^n.$$

Let us now take m < n to be chosen later and, for $\tilde{z} = F^{n-m}(z_2)$, consider the cone $D_{\tilde{z}}F^m\mathbf{C}_u$ corresponding to the interval I_3 in the projective space. By the forward invariance of the unstable cone it is clear that $D_{\tilde{z}}F^m\mathbf{C}_u \supset D_{z_2}F^n\mathbf{C}_u$, meaning that $I_3 \supset I_2$. We are going to prove that $L \in I_3$. Let $w_{n,m} := D_{z_2}F^{n-m}v_{\pm}$. Arguing as before, but remembering also condition (2.5), we have

(7.15)
$$|\sin \measuredangle (D_{\tilde{z}}F^{m}w_{n,m}, D_{\tilde{z}}F^{m}v_{\pm})| = |\sin \measuredangle (w_{n,m}, v_{\pm})| \frac{|\det D_{\tilde{z}}F^{m}|||v_{\pm}||||w_{n,m}||}{||D_{\tilde{z}}F^{m}w_{n,m}||||D_{\tilde{z}}F^{m}v_{\pm}||} \\ \geq C_{*}^{-1}C_{*}^{-2}|\sin(\arctan(\iota_{*}\chi_{u}))|\lambda_{+}^{-m}\mu_{-}^{n} \\ = C_{*}^{-1}C_{*}^{-2}\frac{\iota_{*}\chi_{u}}{\sqrt{1+(\iota_{*}\chi_{u})^{2}}}\lambda_{+}^{-m}\mu_{-}^{n}.$$

It follows that, setting $B_{\iota_{\star}} := a_0 C_{\star}^2 C_{\star}^4 \iota_{\star}^{-1} \ge 1$, if $\lambda_+^{-m} \mu_-^n \ge B_{\iota_{\star}} \lambda_-^{-n} \mu_+^n$, then $L \in I_3$. By a direct computation, and recalling that $\mu := \{\mu_+, \mu_-^{-1}\}^+$, we see that the choice $m = \lceil \alpha n - \beta_{\iota_{\star}} \rceil$, with $\alpha := \frac{\log(\lambda_- \mu^{-2})}{\log(\lambda_+)} > 0$ and $\beta_{\iota_{\star}} := \frac{\log B_{\iota_{\star}}}{\log \lambda_+} \ge 0$ yields the wanted inequality. Also, note that $\alpha < 1$ since $\lambda_- < \lambda_+ \mu^2$.

The above computation shows that, given $z_1 \in F^{-n}(y)$, for every $z_2 \in F^{-n}(y)$ which is non-transversal to z_1 , the line L is contained in the cone $D_{\tilde{z}}F^m\mathbf{C}_u$, for $\tilde{z} = F^{n-m}(z_2)$. In particular, for every $y \in \mathbb{T}^2$, one has

$$\begin{split} \sup_{z_{1}\in F^{-n}(y)} & \sum_{\substack{z_{2}\in F^{-n}(y)\\z_{2}\notin z_{1}}} |\det D_{z_{2}}F^{n}|^{-1} \leq \sup_{z_{1}\in F^{-n}(y)} & \sum_{\substack{z_{2}\in F^{-n}(y)\\D_{F^{n-m}(z_{2})}F^{m}\mathbf{C}_{u}\supset D_{z_{1}}F^{n}(\mathbb{R}\times\{0\})} |\det D_{z_{2}}F^{n}|^{-1} \\ \leq & \sup_{L\subset \mathbb{R}^{p^{2}}} & \sum_{\substack{\tilde{z}\in F^{-m}(y)\\D\tilde{z}F^{m}\mathbf{C}_{u}\supset L}} |\det D_{\tilde{z}}F^{m}|^{-1} & \sum_{z_{2}\in F^{-n+m}(\tilde{z})} |\det D_{z_{2}}F^{n-m}|^{-1} \\ \leq & \mathcal{L}^{n-m}\mathbf{1}(y) \sup_{L\subset \mathbb{R}^{p^{2}}} & \sum_{\substack{z\in F^{-m}(y)\\D_{z}F^{m}\mathbf{C}_{u}\supset L}} |\det D_{z}F^{m}|^{-1}, \end{split}$$

where we have used (4.2). The above inequality implies

$$\mathcal{N}(n) \le \|\mathcal{L}^{n-m}1\|_{\infty} \widetilde{\mathcal{N}}(m) \le \|\mathcal{L}^{\lceil n(1-\alpha)\rceil}1\|_{\infty} \widetilde{\mathcal{N}}(\lceil \alpha n \rceil).$$

8. A SECOND LASOTA-YORKE INEQUALITY: RESULTS

To state the main result we need a few definitions. From Appendix E we recall that, for positive integers $N \in \mathbb{N}$ and $s \geq 1$, and for $u \in \mathcal{C}^r(\mathbb{T}^2)$,

(8.1)
$$\|\mathcal{L}^N u\|_{\mathcal{H}^s}^2 = \sum_{\xi \in \mathbb{Z}^2} |\langle \xi \rangle^s \mathcal{FL}^N u(\xi)|^2,$$

where $\langle \xi \rangle = \sqrt{1 + \|\xi\|^2}$. Since we will work in Fourier space, it is convenient to introduce the notion of the dual of a cone in \mathbb{R}^2 by:

(8.2)
$$\mathbf{C}^{\perp} = \{ v \in \mathbb{R}^2 : \exists u \in \mathbf{C} : \langle v, u \rangle = 0 \},$$

and if $\xi \in \mathbb{Z}^2 \setminus \{0\}$ we define $\xi^{\perp} := (\xi_1^{\perp}, \xi_2^{\perp}) = (-\xi_2, \xi_1) \|\xi\|^{-1}$. In addition, we define $\rho(\xi^{\perp}) = |\xi_2^{\perp}|/|\xi_1^{\perp}|$, for $\xi_1^{\perp} \neq 0$, $\rho(\pm e_2) = \infty$, and

(8.3)
$$\vartheta(\xi^{\perp}) := \{\rho(\xi^{\perp}), \chi_u\}^+.$$

Let us also define the sequence

$$\mathbb{L}_n := \|\mathcal{L}^n 1\|_{\infty}$$

Finally, to state the main result one last key assumption is needed. Let us define

(8.5)
$$n_0(F) := \min\{n \in \mathbb{N} : \forall p \in \mathbb{T}^2 \quad \exists z_1, z_2 \in F^{-n}p : z_1 \pitchfork z_2\}.$$

We will always assume that the map F satisfies

$$(8.6) n_0(F) < \infty.$$

For simplicity, in the following we will just use the notation n_0 instead of $n_0(F)$.

Remark 8.1. In [50] it is proven that assumption (8.6) is generic. More precisely, the author proves that for surface partially hyperbolic systems F, $\mathcal{N}_F(n)$ is generically strictly smaller than 1, for n large.

The goal of this Section is to prove the following Theorem.

Theorem 8.2. (SVPH) Let F be a SVPH. Let m_{χ_u} and n_0 be the integers given in (5.10) and (8.5) respectively. There exist uniform constants $C_1, c_{\sharp} > 0$, $\Lambda > 2$ and $\sigma > 1$ such that, for each $q_0 \ge n_0$ and any $1 \le s < r$, if $M \ge \sigma m_{\chi_u}$ and $N = M + q_0$,

(8.7)
$$\|\mathcal{L}^N u\|_{\mathcal{H}^s} \le C_1 \left(\sqrt{[\mathbb{L}_M \mathcal{N}(q_0)]^{\frac{1}{N}} \mu^{2s}} \right)^N \|u\|_{\mathcal{H}^s} + \Omega_{\chi_u}(M, s) \|u\|_{s+2}.$$

where $\Omega_{\chi_u}(M,s) \leq C_{\epsilon,q_0} \mu^{c_{\sharp}M} Q(M,s) C_{\mu,M}^{c_{\sharp}M} \mathbb{d}^{-4s} \mathbb{L}_M$ and Q(M,s) is given in Lemma 7.1.

In the case of SVPH^{\$\$} we have a sharper control on the constants as specified by the next theorem.

Theorem 8.3. (SVPH^{\sharp}) If the map F is a SVPH^{\sharp} (see Definition 5.1) and satisfies

(8.8)
$$\chi_u^{-1} \|\omega\|_{\mathcal{C}^r} \le C_5$$

for a uniform constant $C_5 > 0$, then there exist $\beta_3 \in \mathbb{R}^+$, depending only on C_5 , such that

(8.9)
$$\Omega_{\chi_u}(M,1) \le C_{\epsilon,q_0} C_{\mu,M}^{\beta_3} \mu^{\beta_3 M} (\ln \chi_u^{-1})^{\beta_3} \chi_u^{\frac{-13}{2} - c_\sharp \ln \mu^{-1}} \mathbb{L}_M^{\frac{1}{2}}.$$

We will prove Theorem 8.2 in Section 8.5 and Theorem 8.3 in Section 8.7, after several preliminary steps in which first we show how to localise the relevant objects and then we show how to estimate some of the Fourier coefficients using the $\|\cdot\|_s$ norms, for s large enough.

8.1. Partitions of unity.

We will use notations and definitions introduced in Section 5.3.1. First of all we want to decompose the transfer operator using suitable partitions of unity. For each point $z \in \mathbb{T}^2$, and $q_0 \ge n_0$, let us set $\delta_{q_0}(z) := \mu_{q_0}^-(z)\lambda_{q_0}^+(z)^{-1}$,⁴³ and define,⁴⁴

(8.10)
$$\mathcal{U}_{z,q_0} = \{ y \in \mathbb{T}^2 : \| y - z \| < \min\{1/2, \mathrm{d}\epsilon \delta_{q_0}(z) \} \},\$$

where 45

(8.11)
$$\mathbf{d} = \mathbf{d}(\chi_u) = C_{\mu, \mathrm{b}\ln\chi_u^{-1}}^{-1} \mu^{-\mathrm{b}\ln\chi_u^{-1}} L_\star(\chi_u, q_0)^{-1} C_0 \chi_u$$

for some uniform constants b, C_0 to be chosen later.⁴⁶

By Besicovitch covering theorem there exists a finite subset \mathcal{A} and points $\{z_{\alpha}\}_{\alpha \in \mathcal{A}}$ such that $\mathbb{T}^2 \subset \bigcup_{\alpha} \mathcal{U}_{\alpha}$ where $\mathcal{U}_{\alpha} = 5\mathcal{U}_{z_{\alpha},q_0}$, and such that the number of intersections is bounded by some fixed constant

⁴³The functions μ_n^- and λ_n^+ are defined in (2.4).

⁴⁴The constant ϵ is the one introduced just before equation (5.9).

 $^{{}^{45}}L_{\star}$ is the Lipschitz constant given in (G.2) with $v = \chi_u$.

⁴⁶The choice of is b, made in Proposition 8.6 and depends on the parameter t specifying the Sobolev space we are interested in. The choice of C_0 is made in Lemma 8.10.

 C_{\sharp} . We then define a family of smooth function $\{\psi_{\alpha}\}_{\alpha}$ supported on \mathcal{U}_{α} such that $\sum_{\alpha} \psi_{\alpha} = 1$. Note that we can choose the ψ_{α} such that

(8.12)
$$\|\psi_{\alpha}\|_{\mathcal{C}^t} \leq C_{\sharp} \mathrm{d}^{-t} \epsilon^{-t} \mu^{q_0 t} \lambda_+^{q_0 t}.$$

Next, we construct a refinement of the above partition using the inverse branches introduced in Section 5.3. For each $\alpha \in \mathcal{A}$ we pick a curve $\gamma_{\alpha} \in \Upsilon$, $\gamma'_{\alpha} = e_2$, such that $\mathcal{U}_{\alpha} \cap \gamma_{\alpha} = \{\emptyset\}$ and let $\tilde{\gamma}_{\alpha} = \gamma + (1/2, 0)$. Recalling, from equation (5.8), that $\mathfrak{H}_{\gamma_{\alpha},1} = \{\mathfrak{h} \in \mathfrak{H} : \mathcal{D}_{\mathfrak{h}} = \mathbb{T}^2 \setminus \{\tilde{\gamma}_{\alpha}\}\}$, for each $\mathfrak{h} \in \mathfrak{H}_{\gamma_{\alpha},1} \cup \mathfrak{H}_{\tilde{\gamma}_{\alpha},1}$ either $\mathfrak{h}(\mathbb{T}^2) \cap \gamma_{\alpha} = \emptyset$ or $\mathfrak{h}(\mathbb{T}^2) \cap \tilde{\gamma}_{\alpha} = \emptyset$. Note that the cardinality of $\mathfrak{H}_{\alpha,0} := \mathfrak{H}_{\gamma_{\alpha},1}$ and $\mathfrak{H}_{\tilde{\gamma}_{\alpha},1}$ is exactly d. We can then consider the set $\mathfrak{H}_{\alpha}^n = \{(\mathfrak{h}_1, \cdots, \mathfrak{h}_n) \in \mathfrak{H}^n : \mathfrak{h}_j \in \mathfrak{H}_{\gamma_{j-1},\alpha}, j \in \{1, \cdots, n\}\}$ where $\gamma_0 = \tilde{\gamma}_{\alpha}$ and, for j > 0, $\gamma_j = \gamma_{\alpha}$ if $\mathfrak{h}_j(\mathbb{T}^2) \cap \tilde{\gamma}_{\alpha} = \emptyset$ and $\gamma_j = \tilde{\gamma}_{\alpha}$ if $\mathfrak{h}_{j-1}(\mathbb{T}^2) \cap \tilde{\gamma}_{\alpha} \neq \emptyset$. Note that \mathfrak{H}_{α}^n has exactly one element for each equivalence class of $\mathfrak{H}_{*,\gamma_{\alpha}}^n$, defined in equation (5.8), hence it is isomorphic to $\mathfrak{H}_{\gamma_{\alpha,n},n}$ and has exactly d^n elements. To simplify notation, given $\alpha \in \mathcal{A}$ and $q_0 \in \mathbb{N}$, in the following we will denote $\mathfrak{H}^{q_0} := \mathfrak{H}_{*,\gamma_{\alpha}}^{q_0}$, which is then a set with cardinality d^{q_0} .

Next, let $\alpha, \alpha' \in \mathcal{A}$ and define

(8.13)
$$\psi_{\alpha\alpha',\mathfrak{h}}(z) = \psi_{\alpha} \circ F^{q_0}(z) \mathbb{1}_{\mathfrak{h},\alpha}(z) \psi_{\alpha'}(z), \ \forall \mathfrak{h} \in \mathfrak{H}^{q_0}, z \in \mathbb{T}^2,$$

where $\mathbb{1}_{\mathfrak{h},\alpha} := \mathbb{1}_{\mathcal{U}_{\alpha,\mathfrak{h}}}$. Note that (8.13) defines a \mathcal{C}^r partition of unity, supported on $\{\mathcal{U}_{\alpha,\alpha',\mathfrak{h}}\}_{\mathfrak{h}\in\mathfrak{H}^{q_0}}$, with $\mathcal{U}_{\alpha\alpha',\mathfrak{h}} := \mathfrak{h}(\mathcal{U}_{\alpha}) \cap \mathcal{U}_{\alpha'}$, and intersection multiplicity bounded by C_{\sharp} .

In order to keep the notation simple, we use the index $\bar{\alpha} = \alpha \alpha'$ to indicate quantities which depend both on α and α' and we write $\sum_{\bar{\alpha}}$ for $\sum_{\alpha \in \mathcal{A}} \sum_{\alpha' \in \mathcal{A}}$.

The following result is similar to [3, Lemma 9], but the proof is adapted to our case.⁴⁷

Lemma 8.4. For each $u \in C^r(\mathbb{T}^2)$

(8.14)
$$\|u\|_{\mathcal{H}^s}^2 \le C_{\sharp} \sum_{\alpha \in \mathcal{A}} \|u\psi_{\alpha}\|_{\mathcal{H}^s}^2$$

(8.15)
$$\sum_{\bar{\alpha}} \sum_{\mathfrak{h} \in \mathfrak{H}^{q_0}} \| u\psi_{\bar{\alpha},\mathfrak{h}} \|_{\mathcal{H}^s}^2 \le C_{\sharp} \| u \|_{\mathcal{H}^s}^2 + C_{\psi}(s) \| u \|_{L^1}^2,$$

where $C_{\psi}(s) = C_{\epsilon,q_0} \mathbf{d}^{-4s}$.

Proof. For the first inequality note that

$$\|u\|_{\mathcal{H}^s}^2 = \left\|\sum_{\alpha \in \mathcal{A}} u\psi_{\alpha}\right\|_{\mathcal{H}^s}^2 = \sum_{(\alpha, \alpha') \in \mathcal{A} \times \mathcal{A}} \langle \psi_{\alpha} u, \psi_{\alpha'} u \rangle_s.$$

By the definition of the $\langle \cdot, \cdot \rangle_s$ the above sum is zero if the supports of ψ_{α} and $\psi_{\alpha'}$ do not intersect. For the other terms, denoting with \mathcal{A}^* the set of elements in $\mathcal{A} \times \mathcal{A}$ for which the above supports intersect, we have:

$$\sum_{\mathcal{A}^*} \langle \psi_{\alpha} u, \psi_{\alpha'} u \rangle_s \le \sum_{\mathcal{A}^*} \frac{\|\psi_{\alpha} u\|_{\mathcal{H}^s}^2 + \|\psi_{\alpha'} u\|_{\mathcal{H}^s}^2}{2} \le C_{\sharp} \sum_{\alpha \in \mathcal{A}} \|\psi_{\alpha} u\|_{\mathcal{H}^s}^2.$$

We now prove (8.15). By formula (E.4) we have, recalling (8.12),

$$\begin{split} \sum_{\bar{\alpha},\mathfrak{h}} \|u\psi_{\bar{\alpha},\mathfrak{h}}\|_{\mathcal{H}^{s}}^{2} &= \sum_{\bar{\alpha},\mathfrak{h}} \sum_{|\beta| \leq s} C_{\beta} \langle \partial^{\beta}(u\psi_{\bar{\alpha},\mathfrak{h}}), \partial^{\beta}(u\psi_{\bar{\alpha},\mathfrak{h}}) \rangle_{L^{2}} \\ &\leq C_{\sharp} \sum_{\bar{\alpha},\mathfrak{h}} \sum_{|\beta| \leq s} \langle (\partial^{\beta}u)\psi_{\bar{\alpha},\mathfrak{h}}, (\partial^{\beta}u)\psi_{\bar{\alpha},\mathfrak{h}} \rangle_{L^{2}} + \langle (\partial^{\beta-1}u)\partial\psi_{\bar{\alpha},\mathfrak{h}}, (\partial^{\beta}u)\psi_{\bar{\alpha},\mathfrak{h}} \rangle_{L^{2}} \\ &+ C_{\epsilon,q_{0}} \mathrm{d}^{-2s} \|u\|_{\mathcal{H}^{s-1}}^{2} \end{split}$$

 $^{^{47}}$ Similar inequalities hold more generally for some anisotropic norms, as used originally in [7] and go under the name of *fragmentation-reconstitution* inequalities in [5].

where we have usded the fact that $\sum_{\bar{\alpha},\mathfrak{h}} |\partial^{\alpha}\psi_{\bar{\alpha},\mathfrak{h}}| |\partial^{\gamma}u| \leq C_{\sharp} \sup_{\bar{\alpha},\mathfrak{h}} ||\psi_{\bar{\alpha},\mathfrak{h}}||_{\mathcal{C}^{|\alpha|}} |\partial^{\gamma}u|$ due to the uniform finite intersection multiplicity of the partition of unity. Hence, since $\psi_{\bar{\alpha},\mathfrak{h}}^2 \leq \psi_{\bar{\alpha},\mathfrak{h}}$,

$$\begin{split} \sum_{\bar{\alpha},\mathfrak{h}} \|u\psi_{\bar{\alpha},\mathfrak{h}}\|_{\mathcal{H}^{s}}^{2} &\leq C_{\sharp} \|u\|_{\mathcal{H}^{s}}^{2} + C_{\epsilon,q_{0}} \mathrm{d}^{-1} \|u\|_{\mathcal{H}^{s}} \|u\|_{\mathcal{H}^{s-1}}^{2} + C_{\epsilon,q_{0}} \mathrm{d}^{-2s} \|u\|_{\mathcal{H}^{s-1}}^{2} \\ &\leq C_{\sharp} \|u\|_{\mathcal{H}^{s}}^{2} + C_{\epsilon,q_{0}} \mathrm{d}^{-2s} \|u\|_{\mathcal{H}^{s-1}}^{2} \leq C_{\sharp} \|u\|_{\mathcal{H}^{s}}^{2} + C_{\epsilon,q_{0}} \mathrm{d}^{-4s} \|u\|_{L^{1}}^{2}, \end{split}$$

where in the last line we used Lemma E.1.

Remark 8.5. For future purposes we note that, under condition (8.8), recalling equation (G.2), we have

$$L_{\star}(\chi_{u}, q_{0}) = C_{\sharp} C_{q_{0}} C_{\mu, \ln \chi_{u}^{-1}} \chi_{u}^{1-c_{\sharp} \ln \mu},$$

which implies that, by (8.11), $C_{\psi}(1) \leq \epsilon^{-2} C_{q_0} C_{\mu, c_{\sharp} \ln \chi_u^{-1}} \chi_u^{c_{\sharp} \ln \mu}$.

8.2. Fourier basic estimate: case SVPH.

The next Proposition is the main ingredient for the proof of Theorem 8.2.

Proposition 8.6. Let $n_0 \in \mathbb{N}$, $\xi \in \mathbb{Z}^2$, $q_0 \ge n_0$, $\alpha, \alpha' \in \mathcal{A}$, and $\mathfrak{h} \in \mathfrak{H}^{q_0}$ be such that $D_p \mathfrak{h}_{n_0} \xi^{\perp} \notin \mathbf{C}_u$, for each $p \in supp \ \psi_{\bar{\alpha},\mathfrak{h}}$, then, for each $t \ge 2$, there exists $M_{\xi} \equiv M_{\xi,t}$, such that

(8.16)
$$\langle \xi \rangle^t | \mathcal{FL}^{q_0}(\psi_{\bar{\alpha},\mathfrak{h}}\mathcal{L}^{M_{\xi}}u)(\xi)| \le K_1(t, M_{\xi}) \|u\|_t,$$

where $K_1(t, M_{\xi}) \leq C_{q_0} C_{\mu, M_{\xi}}^2 \mu^{c_{\sharp} M_{\xi}} \chi_u^{-c_{\sharp}} \Lambda^{c_{\sharp} M_{\xi}}$. Moreover, there exists $\sigma > 1$ such that, with m_{χ_u} as in (5.10),

(8.17)
$$\overline{m}_{\chi_u} := \sigma m_{\chi_u} \ge \sup_{\{\xi: \xi^\perp \notin \mathbf{C}_u\}} M_{\xi}.$$

Proof. Let $\xi = (\xi_1, \xi_2)$, chose $j \in \{1, 2\}$ such that $\|\xi\| \leq 2|\xi_j|$, and $M_{\xi} > 0$ to be chosen later. Since $\xi_j \mathcal{F} u = -i\mathcal{F}\partial_{x_j} u$, and $\|\mathcal{F} u\|_{\infty} \lesssim \|u\|_{L^1}$, using (6.1) we have, for each $t \geq 1$ and setting $u_{\bar{\alpha}, \bar{\beta}}^{M_{\xi}} = \psi_{\bar{\alpha}, \bar{\beta}} \mathcal{L}^{M_{\xi}} u$,

(8.18)
$$\langle \xi \rangle^t | \mathcal{FL}^{q_0}(\psi_{\bar{\alpha},\mathfrak{h}}\mathcal{L}^{M_{\xi}}u)(\xi)| \lesssim \|\mathcal{L}^{q_0}(u_{\bar{\alpha},\mathfrak{h}}^{M_{\xi}})\|_{L^1} + |\xi_j|^t | \mathcal{FL}^{q_0}(u_{\bar{\alpha},\mathfrak{h}}^{M_{\xi}})| \\ \lesssim \|u\|_0 + |\mathcal{F}\partial_{x,\iota}^t\mathcal{L}^{q_0}(u_{\bar{\alpha},\mathfrak{h}}^{M_{\xi}})|.$$

Letting $J_k(p) = (\det D_p F^k)^{-1}$ and recalling (8.13), we have

(8.19)
$$\begin{bmatrix} \mathcal{F}\partial_{x_{j}}^{t}\mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}}^{M_{\xi}}) \end{bmatrix}(\xi) = \int_{\mathbb{T}^{2}} dz \ e^{-2\pi i \langle z,\xi \rangle} \partial_{z_{j}}^{t} \left\{ \left[J_{q_{0}}\psi_{\bar{\alpha},\mathfrak{h}}\mathcal{L}^{M_{\xi}}u \right] \circ \mathfrak{h} \right\}(z) \\ = \sum_{|\eta_{1}|+|\eta_{2}|=t} C_{\eta_{1},\eta_{2}} \int_{\mathcal{U}_{\alpha}\cap F^{q_{0}}(\mathcal{U}_{\alpha'})} dz \ e^{-2\pi i \langle z,\xi \rangle} \partial^{\eta_{1}} \left[\psi_{\alpha}\psi_{\alpha'} \circ \mathfrak{h} \right](z) \cdot \partial^{\eta_{2}} \left\{ \left[J_{q_{0}}\mathcal{L}^{M_{\xi}}u \right] \circ \mathfrak{h} \right\}(z)$$

To simplify notation let $\bar{\psi}_{\bar{\alpha},\mathfrak{h}} := \psi_{\alpha}\psi_{\alpha'}\circ\mathfrak{h}$. Using the change of variables $\gamma_{\ell}(\tau) = z_{\alpha} + \ell\xi + \tau\xi^{\perp}$, where $\ell, \tau \in I_{q_0} = [-\mathrm{d}\epsilon\delta_{q_0}(z_{\alpha}), \mathrm{d}\epsilon\delta_{q_0}(z_{\alpha})]^{48}$ we have

(8.20)
$$\left| \mathcal{F}\partial_{x_{j}}^{t} \mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}}^{M_{\xi}}) \right| \leq C_{\sharp} \sup_{|\eta_{1}|+|\eta_{2}|=t} \int_{I_{q_{0}}} d\ell \left| \int_{I_{q_{0}}} d\tau \left\{ \partial^{\eta_{1}} \bar{\psi}_{\bar{\alpha},\mathfrak{h}} \cdot \partial^{\eta_{2}} \left[J_{q_{0}} \mathcal{L}^{M_{\xi}} u \right] \circ \mathfrak{h} \right\} (\gamma_{\ell}(\tau)) \right|.$$

For each $\tilde{\mathfrak{h}} \in \mathfrak{H}^{\infty}$ let $\bar{\mathfrak{h}} = \tilde{\mathfrak{h}} \circ \mathfrak{h}$, provided it is well-defined. Let $m := m(p, \mathfrak{h})$ be the smallest integer such that, for each $\bar{\mathfrak{h}}$ and $p \in \operatorname{supp} \psi_{\bar{\alpha},\mathfrak{h}}$, $D_p \bar{\mathfrak{h}}_m \xi^{\perp} \in \mathbf{C}_{\epsilon,c}$. Note that, by hypothesis and recalling (5.11), $m - n_0 \leq c_2^+ (\ln \chi_u^{-1} + 1)$.

⁴⁸By definition, the curves γ_{ℓ} cover $\mathcal{U}_{z_{\alpha},q_0}$, given in (8.10), which contain the support of $\psi_{\bar{\alpha}}$.

For $z \in \gamma_{\ell}$ let $\overline{m}_{\ell}(z, \tilde{\mathfrak{h}} \circ \mathfrak{h}) := \sigma m(z, \mathfrak{h})$, with σ as in (5.34), and set $\overline{m}_{\ell}(\tilde{\mathfrak{h}}) = \sup_{z \in \gamma_{\ell}} \overline{m}_{\ell}(z, \tilde{\mathfrak{h}} \circ \mathfrak{h})$.⁴⁹ We then define

(8.21)
$$M_{\xi} \equiv M_{\xi,t} = \sup_{\ell} \sup_{\tilde{\mathfrak{h}} \in \mathfrak{H}^{\infty}} \overline{m}_{\ell}(\tilde{\mathfrak{h}}).$$

Observe that, the assumption $D_p \bar{\mathfrak{h}}_{n_0} \xi^{\perp} \notin \mathbf{C}_u$ and condition (2.5) imply (8.17) for $t \geq 2$.

In the following we estimate the inner integrals of (8.20) for each fixed ℓ , since this does not create confusion we thus drop the ℓ subscript in \overline{m}_{ℓ} to ease notation.

Define $\hat{\mathfrak{H}}_{\alpha}^{*} = \{\mathfrak{h}_{\overline{m}(\overline{\mathfrak{h}})}\}_{\overline{\mathfrak{h}}\in\mathfrak{H}^{\infty}}, \, \hat{\mathfrak{H}}_{\alpha} = \{\hat{\mathfrak{h}} \, : \, \hat{\mathfrak{h}}\circ\mathfrak{h}\in\hat{\mathfrak{H}}_{\alpha}^{*}\}, \, v_{\overline{\alpha},\hat{\mathfrak{h}}} = \mathcal{L}^{M_{\xi}-\overline{m}(\hat{\mathfrak{h}})+q_{0}}u \text{ and write}$

(8.22)
$$\mathfrak{H}^{M_{\xi}} = \bigcup_{\hat{\mathfrak{h}} \in \hat{\mathfrak{H}}_{\alpha}} \left\{ \mathfrak{h}' \circ \hat{\mathfrak{h}} : \mathfrak{h}' \in \mathfrak{H}^{M_{\xi} - \overline{m}(\hat{\mathfrak{h}}) + q_{0}} \right\}.$$

This allows to define the decomposition

$$\mathcal{L}^{M_{\xi}} u = \sum_{\hat{\mathfrak{h}} \in \hat{\mathfrak{H}}_{\alpha}} J_{\overline{m}(\hat{\mathfrak{h}}) - q_{0}} \circ \hat{\mathfrak{h}} \cdot \left[\mathcal{L}^{M_{\xi} - \overline{m}(\hat{\mathfrak{h}}) + q_{0}} u \right] \circ \hat{\mathfrak{h}} = \sum_{\hat{\mathfrak{h}} \in \hat{\mathfrak{H}}_{\alpha}} J_{\overline{m}(\hat{\mathfrak{h}}) - q_{0}} \circ \hat{\mathfrak{h}} \cdot v_{\bar{\alpha}, \hat{\mathfrak{h}}} \circ \hat{\mathfrak{h}}.$$

Thus, recalling (6.12),

$$\begin{split} & \left| \mathcal{F}\partial_{x_{j}}^{t}\mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}}^{M_{\xi}}) \right| \\ & \leq C_{\sharp} \sup_{|\eta_{1}|+|\eta_{2}|=t} \sum_{\hat{\mathfrak{h}}\in\hat{\mathfrak{H}}_{\alpha}} \int_{I_{q_{0}}} d\ell \left| \int_{I_{q_{0}}} d\tau \left\{ \partial^{\eta_{1}} \bar{\psi}_{\bar{\alpha},\mathfrak{h}} \cdot J_{\overline{m}(\hat{\mathfrak{h}})} \circ \hat{\mathfrak{h}} \circ \mathfrak{h} \left[P_{\overline{m}(\hat{\mathfrak{h}}),|\eta_{2}|}^{\eta_{2}} v_{\bar{\alpha},\hat{\mathfrak{h}}} \right] \circ \hat{\mathfrak{h}} \circ \mathfrak{h} \right\} (\gamma_{\ell}(\tau)) \right| . \end{split}$$

Next, we apply Lemma G.1 to γ_{ℓ} with $\delta = d\epsilon \delta_{q_0}(z_{\alpha})$, note that the hypotheses of the Lemma are satisfied thanks to the assumptions on ξ . We thus obtain closed curves $\tilde{\gamma}_{\ell}$ with j + 1 derivative bounded by $C_{q_0,j}\Delta_{\tilde{\gamma}}^j$. It follows

$$\begin{split} & \left| \mathcal{F}\partial_{x_{j}}^{t} \mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}}^{M_{\xi}}) \right| \\ & \leq C_{\sharp} \sup_{|\eta_{1}|+|\eta_{2}|=t} \sum_{\hat{\mathfrak{h}}\in\hat{\mathfrak{H}}_{\alpha}} \int_{I_{q_{0}}} d\ell \left| \int_{\mathbb{T}} d\tau \left\{ \partial^{\eta_{1}} \bar{\psi}_{\bar{\alpha},\mathfrak{h}} \cdot J_{\overline{m}(\hat{\mathfrak{h}})} \circ \hat{\mathfrak{h}} \circ \mathfrak{h} \left[P_{\overline{m}(\hat{\mathfrak{h}}),|\eta_{2}|}^{\eta_{2}} v_{\bar{\alpha},\hat{\mathfrak{h}}} \right] \circ \hat{\mathfrak{h}} \circ \mathfrak{h} \right\} (\tilde{\gamma}_{\ell}(\tau)) \right|. \end{split}$$

Next, we apply, for each inverse branch $\hat{\mathfrak{h}} \circ \mathfrak{h}$, Lemma 5.13 to the curves $\tilde{\gamma}_{\ell}$ and obtain admissible central curves $\hat{\nu}_{\ell} = \nu_{\ell} \circ h_{\ell,\overline{m}}$.⁵⁰ Thus, we can rewrite the inner integrals in the right hand side of the above equation as follows

$$\begin{split} &\int_{\mathbb{T}} d\tau \left\{ \partial^{\eta_1} \bar{\psi}_{\bar{\alpha},\mathfrak{h}} \cdot J_{\overline{m}(\hat{\mathfrak{h}})} \circ \hat{\mathfrak{h}} \circ \mathfrak{h} \left[P^{\eta_2}_{\overline{m}(\hat{\mathfrak{h}}),|\eta_2|} v_{\bar{\alpha},\hat{\mathfrak{h}}} \right] \circ \hat{\mathfrak{h}} \circ \mathfrak{h} \right\} (\tilde{\gamma}_{\ell}(\tau)) \\ &= \int_{\mathbb{T}} d\tau \Psi_{\hat{\nu}_{\ell}}(\tau) \left\{ (\partial^{\eta_1} \bar{\psi}_{\bar{\alpha},\mathfrak{h}}) \circ F^{\overline{m}(\hat{\mathfrak{h}})} \cdot \left[P^{\eta_2}_{\overline{m}(\hat{\mathfrak{h}}),|\eta_2|} v_{\bar{\alpha},\hat{\mathfrak{h}}} \right] \right\} (\hat{\nu}_{\ell}(\tau)), \end{split}$$

where $\Psi_{\hat{\nu}_{\ell}}(\tau) = h'_{\ell,\overline{m}}[\det D_{\hat{\nu}_{\ell}(\tau)}F^{\overline{m}(\hat{\mathfrak{h}})}]^{-1}$. By Proposition 6.4 applied with $n = \overline{m}(\hat{\mathfrak{h}}), \varphi = \Psi_{\hat{\nu}_{\ell}}(\partial^{\eta_1}\bar{\psi}_{\bar{\alpha},\mathfrak{h}}) \circ F^{\overline{m}(\hat{\mathfrak{h}})} \|_{\mathcal{C}^{[\eta_2]}_{\hat{\nu}_{\epsilon}}}^{-1}, \psi = 1, u = v_{\bar{\alpha},\hat{\mathfrak{h}}} \text{ and } U = \mathbb{T}^2$, the above integral is bounded by

(8.23)
$$\tilde{C}(t,\overline{m}(\hat{\mathfrak{h}}),m)\|\Psi_{\hat{\nu}_{\ell}}(\partial^{\eta_1}\bar{\psi}_{\bar{\alpha},\mathfrak{h}})\circ F^{\overline{m}(\hat{\mathfrak{h}})}\|_{\mathcal{C}^{|\eta_2|}_{\hat{\nu}_{\ell}}}\|v_{\bar{\alpha},\hat{\mathfrak{h}}}\|_{|\eta_2|},$$

⁴⁹Notice that \overline{m} depends on ξ through γ_{ℓ} . Also, it would be more precise to call it $\overline{m}_{\ell}(\tilde{\mathfrak{h}} \circ \mathfrak{h})$, but we keep the notation as simple as possible.

⁵⁰Notice that $\nu_{\ell} = \hat{\mathfrak{h}} \circ \mathfrak{h} \circ \gamma_{\ell}$ depends on $\hat{\mathfrak{h}}$, but we drop this dependence for simplicity.

where $\tilde{C}(t, \overline{m}(\hat{\mathfrak{h}}), m) \leq C_{\sharp} \Lambda^{c_{\sharp} M_{\xi}}$. Accordingly,

(8.24)
$$\left| \mathcal{F}\partial_{x_{j}}^{t} \mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}}^{M_{\xi}}) \right| \leq C_{\sharp} \Lambda^{c_{\sharp}M_{\xi}} \sup_{|\eta_{1}|+|\eta_{2}|=t} \sum_{\mathfrak{\hat{h}}\in\mathfrak{\hat{h}}_{\alpha}} \|\Psi_{\hat{\nu}_{\ell}} \cdot (\partial^{\eta_{1}} \bar{\psi}_{\bar{\alpha},\mathfrak{h}}) \circ F^{\overline{m}(\mathfrak{\hat{h}})}\|_{\mathcal{C}^{|\eta_{2}|}_{\hat{\nu}_{\ell}}} \|v_{\bar{\alpha},\mathfrak{\hat{h}}}\|_{t} |I_{q_{0}}|,$$

where $|I_{q_0}| \leq 2d\epsilon \delta_{q_0}(z_{\alpha}) \leq 2d\epsilon \lambda_{-}^{-q_0} \mu^{q_0}$. Let us estimate the terms in the above sum, setting temporarily $\overline{m} = \overline{m}(\hat{\mathfrak{h}})$. By Corollary 6.7

(8.25)
$$\|v_{\bar{\alpha},\hat{\mathfrak{h}}}\|_{t} = \|\mathcal{L}^{M_{\xi}-\overline{m}(\mathfrak{h})+q_{0}}u\|_{t} \le C_{\mu,M_{\xi}}\mu^{M_{\xi}}C_{q_{0}}\|u\|_{t}$$

and, by Lemma 5.19, for each $\alpha \in \mathcal{A}$

(8.26)
$$\sum_{\hat{\mathfrak{h}}\in\hat{\mathfrak{H}}_{\alpha}} \|\Psi_{\hat{\nu}(\hat{\mathfrak{h}})}\|_{\mathcal{C}^{t}} \leq \sum_{\hat{\mathfrak{h}}\in\mathfrak{H}^{M_{\xi}}} \|\Psi_{\hat{\nu}_{\ell}(\hat{\mathfrak{h}})}\|_{\mathcal{C}^{t}} \leq A_{u}(t,\overline{m},m),$$

where,

(8.27)
$$A_u(t, M_{\xi}, m) := C_{\sharp} (\chi_u^{-1} \Delta_{\tilde{\gamma}})^{c_{\sharp}} \vartheta_{\tilde{\gamma}}^{-1} \Lambda^{c_{\sharp} M_{\xi}}, t \ge 2.$$

Lemma G.1 implies

(8.28)
$$\Delta_{\tilde{\gamma}} \leq C_{q_0,\epsilon} \vartheta(\xi^{\perp})^{-1} \mu^{5m} C_{\mu,m}(\|\omega\|_{\mathcal{C}^2} + \vartheta(\xi^{\perp})) \{1, \mu^{5m} C_{\mu,m}(\|\omega\|_{\mathcal{C}^2} + \vartheta(\xi^{\perp})\}^+ \mu^m, \\ \vartheta_{\tilde{\gamma}} \geq \vartheta_{\gamma} = \{\rho(\xi^{\perp}), \chi_u\}^+ =: \vartheta(\xi^{\perp}).$$

With the choice $\mathbf{b} = 1$, then by equations (8.13) and (8.12), we have the bound $\|(\partial^{\eta_1} \bar{\psi}_{\bar{\alpha},\mathfrak{h}}) \circ F^{\overline{m}(\hat{\mathfrak{h}})}\|_{\mathcal{C}^{|\eta_2|}_{\hat{\nu}_{\ell}}} \leq C_{\epsilon,q_0} \Lambda^{c_{\sharp}tM_{\xi}} \mu^{c_{\sharp}M_{\xi}} \chi_u^{-c_{\sharp}t}$, by (8.25), (8.26), (8.27), (8.28) and (8.24) we conclude.

8.3. Decomposition of Fourier space.

Let $\mathcal{Z}_u = \{\xi : \xi^{\perp} \in \mathbf{C}_u\}$ and $\mathcal{Z}_u^c = \mathbb{Z}^2 \setminus \mathcal{Z}_u$. Recalling that $\rho(\xi^{\perp}) = |\xi_2^{\perp}| |\xi_1^{\perp}|^{-1}, \rho(e_2) = \infty$, $\mathcal{Z}_u = \{\xi : \rho(\xi^{\perp}) \le \chi_u\} \quad ; \quad \mathcal{Z}_u^c = \{\xi : \rho(\xi^{\perp}) > \chi_u\}.$

Next, take $N = q_0 + M$, for some $M \in \mathbb{N}$ to be chosen shortly. For simplicity, it is convenient to introduce the following notation for $A \subset \mathbb{Z}^2, \mathfrak{h}, \mathfrak{h}' \in \mathfrak{H}^{q_0}$:

(8.29)
$$S_{q_0,M}^{\bar{\alpha}}(A,\mathfrak{h},\mathfrak{h}') = \sum_{\xi \in \mathbb{Z}^2} \mathbb{1}_A(\xi) \langle \xi \rangle^{2s} [\mathcal{FL}^{q_0}(u^M_{\bar{\alpha},\mathfrak{h}})](\xi) [\overline{\mathcal{FL}^{q_0}(u^M_{\bar{\alpha},\mathfrak{h}'})}](\xi),$$

where $u_{\bar{\alpha},\mathfrak{h}}^M = \psi_{\bar{\alpha},\mathfrak{h}} \mathcal{L}^M u$. Then, using equation (8.14) we have

$$(8.30) \qquad \begin{aligned} \|\mathcal{L}^{N}u\|_{\mathcal{H}^{s}}^{2} \leq C_{\sharp}\sum_{\bar{\alpha}} \|\psi_{\alpha}\mathcal{L}^{q_{0}}(\psi_{\alpha'}\mathcal{L}^{M}u)\|_{\mathcal{H}^{s}}^{2} \leq C_{\sharp}\sum_{\bar{\alpha}} \left\|\sum_{\mathfrak{h}\in\mathfrak{H}^{q_{0}}}\mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}}^{M})\right\|_{\mathcal{H}^{s}}^{2} \\ &= C_{\sharp}\sum_{\bar{\alpha}}\sum_{(\mathfrak{h},\mathfrak{h}')\in\mathfrak{H}^{q_{0}}\times\mathfrak{H}^{q_{0}}}\langle\mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}}^{M}),\mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}'}^{M})\rangle_{s} \\ &= C_{\sharp}\sum_{\bar{\alpha}}\sum_{(\mathfrak{h},\mathfrak{h}')\in\mathfrak{H}^{q_{0}}\times\mathfrak{H}^{q_{0}}}\sum_{\xi\in\mathbb{Z}^{2}}\langle\xi\rangle^{2s}[\mathcal{F}\mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}}^{M})](\xi)[\overline{\mathcal{F}\mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}'}^{M})}](\xi) \\ &= C_{\sharp}\sum_{\bar{\alpha}}\sum_{(\mathfrak{h},\mathfrak{h}')\in\mathfrak{H}^{q_{0}}\times\mathfrak{H}^{q_{0}}}S_{q_{0},M}^{\bar{\alpha}}(\mathcal{Z}_{u},\mathfrak{h},\mathfrak{h}') + C_{\sharp}\sum_{\bar{\alpha}}\sum_{(\mathfrak{h},\mathfrak{h}')\in\mathfrak{H}^{q_{0}}\times\mathfrak{H}^{q_{0}}}S_{q_{0},M}^{\bar{\alpha}}(\mathcal{Z}_{u}^{C},\mathfrak{h},\mathfrak{h}') \end{aligned}$$

We start with the second term in the above equation, next we will treat the term with $\xi \in \mathcal{Z}_u$.

Lemma 8.7 (Bound on \mathcal{Z}_{u}^{c}). Recall by (8.17) that $\overline{m}_{\chi_{u}} := \sup_{\xi \in \mathcal{Z}_{u}^{c}} M_{\xi}^{51}$ For each $M \geq \overline{m}_{\chi_{u}}$, $1 \leq s \leq r-1$, $\mathfrak{h} \in \mathfrak{H}^{q_{0}}$ and $N = q_{0} + M$,

(8.31)
$$\sum_{\xi \in \mathbb{Z}^2} |\langle \xi \rangle^s \mathbb{1}_{\mathcal{Z}^c_u}(\xi) [\mathcal{FL}^{q_0}(u^M_{\bar{\alpha},\mathfrak{h}})](\xi)|^2 \lesssim \Theta_s ||u||^2_{s+2},$$

where $\Theta_s = C_{q_0} C^6_{\mu,M} \mu^{c_{\sharp}M} \Lambda^{c_{\sharp}M}.$

 $^{^{51}}$ Recall that this is finite by Proposition 8.6.

Proof. Remark that $\Lambda^M \ge \mu^{-M} \chi_u^{-1}$. Since $\xi^{\perp} \notin \mathbf{C}_u$ we can apply Proposition 8.6 with $n_0 = 0$. For each $M \ge M_{\xi}$, by (8.16), we have

(8.32)
$$\sum_{\xi \in \mathbb{Z}^2} |\langle \xi \rangle^s \mathbb{1}_{\mathcal{Z}^c_u} \mathcal{FL}^{q_0}(u^M_{\bar{\alpha},\mathfrak{h}'})|^2 = \sum_{\xi \in \mathbb{Z}^2} \langle \xi \rangle^{-4} |\langle \xi \rangle^{s+2} \mathbb{1}_{\mathcal{Z}^c_u} \mathcal{FL}^{q_0}(\psi_{\bar{\alpha},\mathfrak{h}'} \mathcal{L}^{M_{\xi}}(\mathcal{L}^{M-M_{\xi}}u))|^2 \\ \lesssim C_{q_0} C^4_{\mu,M} \mu^{c_{\sharp}M} \Lambda^{c_{\sharp}M} \|\mathcal{L}^{M-M_{\xi}}u\|^2_{s+2}.$$

The statement (8.31) for s > 1 follows since, by Corollary 6.7,

(8.33)
$$\|\mathcal{L}^{M-M_{\xi}}u\|_{t}^{2} \leq C_{\mu,M}^{2}\mu^{2M}\|u\|_{t}^{2}, \quad \forall t \geq 1.$$

8.4. The case $\xi^{\perp} \in \mathbf{C}_u$ $(\xi \in \mathcal{Z}_u)$.

In this case we cannot apply Proposition 8.6 directly as we did in the previous section. The reason is that ξ^{\perp} could belong to the unstable direction, so its preimage never enters the central cone. Here transversality plays a major role.

Lemma 8.8 (Bound on \mathcal{Z}_u). If there exists $q_0 \in \mathbb{N}$ such that for each $\xi \in \mathbb{Z}$ the hypothesis of Proposition 8.6 are satisfied, then there exist C_{q_0} such that, for each $M \geq \overline{m}_{\chi_u}$ and each $\delta \in (0, 1)$,

$$\sum_{(\mathfrak{h},\mathfrak{h}')\in\mathfrak{H}^{q_{0}}\times\mathfrak{H}^{q_{0}}}\sum_{\xi\in\mathbb{Z}^{2}}\mathbb{1}_{\mathcal{Z}_{u}}(\xi)\langle\xi\rangle^{2s}[\mathcal{FL}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}}^{M})](\xi)[\overline{\mathcal{FL}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}'}^{M})}](\xi)$$

$$\leq \left(\mathcal{N}(q_{0})\mu^{2sq_{0}}+\delta\right)\sum_{\mathfrak{h}\in\mathfrak{H}^{q_{0}}}\|u_{\bar{\alpha},\mathfrak{h}}^{M}\|_{\mathcal{H}^{s}}^{2}+C_{q_{0},s}\delta^{-1}\sum_{\mathfrak{h}\in\mathfrak{H}^{q_{0}}}\|u_{\bar{\alpha},\mathfrak{h}}^{M}\|_{\mathcal{H}^{s-1}}^{2}$$

$$+C_{q_{0}}Q(M,s)\sqrt{\Theta_{s}}\|u\|_{s+2}\|u\|_{\mathcal{H}^{s}},$$

where Q(M,s) is given in (7.2) and Θ_s in Lemma 8.7.

The rest of this Section is devoted to the proof of the above Lemma. We argue in three Steps.

8.4.1. Step I (Local transversality). First we need a definition of transversality uniform on the elements of the partition of unity (8.13):

Definition 8.9. Given $n \in \mathbb{N}$ and $\mathfrak{h}, \mathfrak{h}' \in \mathfrak{H}^n$ we say that $\mathfrak{h} \uparrow_{\alpha}^n \mathfrak{h}'$ (\mathfrak{h} is transversal to \mathfrak{h}' on α at time n) if for every $z \in \mathfrak{h}(\mathcal{U}_{\alpha})$ and $w \in \mathfrak{h}'(\mathcal{U}_{\alpha})$ such that $F^n(z) = F^n(w) \in \mathcal{U}_{\alpha}$:

$$(8.34) D_z F^n \mathbf{C}_{\epsilon,u} \cap D_w F^n \mathbf{C}_{\epsilon,u} = \{0\}.$$

Next, we relate the (pointwise) Definition 4.2 to the (local) Definition 8.9.

Lemma 8.10. The constant C_0 in (8.11) can be chosen such that: for all $\alpha \in \mathcal{A}$, $p \in \mathcal{U}_{\alpha} \subset \mathbb{T}^2$ and $\mathfrak{h}, \mathfrak{h}' \in \mathfrak{H}^{q_0}$ if $z_1 = \mathfrak{h}(p)$ and $z_2 = \mathfrak{h}'(p)$, then $z_1 \pitchfork z_2$ implies $\mathfrak{h} \pitchfork^{q_0}_{\alpha} \mathfrak{h}'$.

Proof. Let $p \in \mathcal{U}_{\alpha} \subset \mathbb{T}^2$ and $\mathfrak{h}, \mathfrak{h}' \in \mathfrak{H}^{q_0}$, if $z_1 = \mathfrak{h}(p)$ and $z_2 = \mathfrak{h}'(p)$, recall that $z_1 \pitchfork z_2$ means

(8.35)
$$D_{z_1} F^{q_0} \mathbf{C}_u \cap D_{z_2} F^{q_0} \mathbf{C}_u = \{0\}.$$

As $\mathbf{C}_{u,\epsilon} \in \mathbf{C}_u$, clearly $D_{z_1} F^{q_0} \mathbf{C}_{u,\epsilon} \in D_{z_1} F^{q_0} \mathbf{C}_u$. So the above implies also

$$D_{z_1}F^{q_0}\mathbf{C}_{u,\epsilon}\cap D_{z_2}F^{q_0}\mathbf{C}_{u,\epsilon}=\{0\}.$$

Let $\tilde{p} \in \mathcal{U}_{\alpha}$, $\tilde{p} \neq p$, and define $\tilde{z}_1 = \mathfrak{h}(\tilde{p})$ and $\tilde{z}_1 = \mathfrak{h}'(\tilde{p})$. We claim that, for each $v \in \mathbf{C}_{u,\epsilon}$, the difference between $D_{z_1}F^{q_0}v$ and $D_{\tilde{z}_1}F^{q_0}v$ is smaller than the difference between $D_{z_1}F^{q_0}\mathbf{C}_u$ and $D_{z_1}F^{q_0}\mathbf{C}_{u,\epsilon}$, provided we choose \mathcal{U}_{α} small enough. This suffices to conclude the argument.

We compute a lower bound for the opening of the connected components of $D_{z_1}F^{q_0}\mathbf{C}_u \setminus D_{z_1}F^{q_0}\mathbf{C}_{u,\epsilon}$. By Proposition 5.6, and by formula (5.15), we deduce that, for each unitary vectors $v \in \mathbf{C}_{u,\epsilon}$ and $w \notin \mathbf{C}_u \cup \mathbf{C}_c$,

$$\measuredangle(D_{z_1}F^{q_0}v, D_{z_1}F^{q_0}w) = \frac{|\det D_{z_1}F^{q_0}|\measuredangle(v,w)}{\|D_{z_1}F^{q_0}v\|\|D_{z_1}F^{q_0}w\|} \ge \frac{C_*\chi_u\epsilon}{\mu_{q_0}^-(z)\lambda_{q_0}^+(z)} = C_*\chi_u\epsilon\delta_{q_0}(z_1).$$

On the other hand let us recall that $u_{\mathfrak{h},q_0}(p)$ defined in (5.14) gives the slope of the boundary of the cone $D_{\mathfrak{h}_{\alpha}(p)}F^{q_0}\mathbf{C}_u$, and it is a Lipschitz function of p. In particular, Lemma G.1 provides an estimate for the Lipschitz constant $L_{\star}(q_0)$ given in (G.2). Then, by the definition of \mathcal{U}_{z,q_0} in (8.10) and (8.11), we have the claim, since

$$\begin{aligned} \|D_{z_1}F^{q_0}v - D_{\tilde{z}_1}F^{q_0}v\| &\leq L_{\star}(q_0)\|z_1 - \tilde{z}_1\| \leq L_{\star}(q_0)L_{\star}(\chi_u, q_0)^{-1}C_0\chi_u\epsilon\delta_{q_0}(z_1)\\ &\leq C_{\sharp}C_0\chi_u\epsilon\delta_{q_0}(z_1). \end{aligned}$$

Clearly the same is true replacing $z_1, \tilde{z}_1, \mathfrak{h}$ with $z_2, \tilde{z}_2, \mathfrak{h}'$, and the result follows.

It is thus natural to work with the local transversality. Using Definition 8.9, and recalling notation (8.29), we introduce the following decomposition into transversal and non transversal terms, which we will estimate separately,

(8.36)
$$\sum_{(\mathfrak{h},\mathfrak{h}')\in\mathfrak{H}^{q_0}\times\mathfrak{H}^{q_0}} S_{q_0,M}^{\bar{\alpha}}(\mathcal{Z}_u,\mathfrak{h},\mathfrak{h}') = \sum_{\mathfrak{h}\mathfrak{h}_{\alpha}^{q_0}\mathfrak{h}'} S_{q_0,M}^{\bar{\alpha}}(\mathcal{Z}_u,\mathfrak{h},\mathfrak{h}') + \sum_{\mathfrak{h}\mathfrak{H}_{\alpha}^{q_0}\mathfrak{h}'} S_{q_0,M}^{\bar{\alpha}}(\mathcal{Z}_u,\mathfrak{h},\mathfrak{h}').$$

Step II (Estimate of transversal terms). In this step we will prove that

(8.37)
$$\sum_{\mathfrak{h}\mathfrak{h}_{\alpha}^{q_{0}}\mathfrak{h}'} S_{q_{0},M}^{\bar{\alpha}}(\mathcal{Z}_{u},\mathfrak{h},\mathfrak{h}') \leq C_{q_{0}}Q(M,s)\sqrt{\Theta_{s}}\|u\|_{s+2}\|u\|_{\mathcal{H}^{s}}$$

where Θ_s is given in Lemma 8.7.

If $\mathfrak{h} \uparrow_{\alpha}^{q_0} \mathfrak{h}'$, then for any $\xi \in \mathbb{Z}^2 \setminus \{0\}$, either $\xi^{\perp} \notin D_{\mathfrak{h}(p)} F^{q_0} \mathbf{C}_{\epsilon,u}$ or $\xi^{\perp} \notin D_{\mathfrak{h}'(p)} F^{q_0} \mathbf{C}_{\epsilon,u}$, for all $p \in \operatorname{supp}(\psi_{\bar{\alpha},\mathfrak{h}})$. We can then decompose $\mathcal{Z}_u = Z(\mathfrak{h}) \cup Z(\mathfrak{h}')$, where

(8.38)
$$Z(\mathfrak{h}) = \{\xi \in \mathcal{Z}_u : \xi^{\perp} \notin D_{\mathfrak{h}(p)} F^{q_0} \mathbf{C}_{\epsilon, u} \quad \forall p \in \mathrm{supp} \psi_{\bar{\alpha}, \mathfrak{h}} \},$$

and we write, recalling (8.29),

(8.39)
$$S_{q_0,M}^{\bar{\alpha}}(\mathcal{Z}_u,\mathfrak{h},\mathfrak{h}') = S_{q_0,M}^{\bar{\alpha}}(\mathcal{Z}_u \cap Z(\mathfrak{h}),\mathfrak{h},\mathfrak{h}') + S_{q_0,M}^{\bar{\alpha}}(\mathcal{Z}_u \cap Z(\mathfrak{h}'),\mathfrak{h},\mathfrak{h}').$$

It is enough to estimate the first addend, the second being analogous. By the Cauchy-Schwarz inequality we have

(8.40)
$$|S_{q_0,M}^{\bar{\alpha}}(\mathcal{Z}_u \cap Z(\mathfrak{h}),\mathfrak{h},\mathfrak{h}')| \leq \left(\sum_{\xi \in \mathbb{Z}^2} |\langle \xi \rangle^s \mathbb{1}_{\mathcal{Z}_u \cap Z(\mathfrak{h})} \mathcal{FL}^{q_0}(u_{\bar{\alpha},\mathfrak{h}}^M)|^2\right)^{\frac{1}{2}} \|\mathcal{L}^{q_0}(u_{\bar{\alpha},\mathfrak{h}'}^M)\|_{\mathcal{H}^s}$$

Moreover, by (7.2), $\|\mathcal{L}^{q_0}(u_{\bar{\alpha},\mathfrak{h}'}^M)\|_{\mathcal{H}^s} \leq C_{\sharp}Q(M,s)\|u\|_{\mathcal{H}^s}$. We can bound the sum inside the square root following exactly the same argument of the proof of Lemma 8.7, since the key condition $\xi \in \mathcal{Z}_u^c$ is now replaced by $\xi \in Z(\mathfrak{h})$, with the difference that this time $\vartheta(\xi^{\perp}) = \chi_u$, since $\xi \in \mathcal{Z}_u$. We thus have

$$|S_{q_0,M}^{\bar{\alpha}}(\mathcal{Z}_u \cap Z(\mathfrak{h}),\mathfrak{h},\mathfrak{h}')| \lesssim Q(M,s)\sqrt{\Theta_s} \|u\|_{s+2} \|u\|_{\mathcal{H}^s}.$$

Summing over $\mathfrak{h} \uparrow_{\alpha}^{q_0} \mathfrak{h}'$, we conclude the proof of (8.37).

Step III (Estimate of non-transversal terms). We now want to estimate the $\mathfrak{h} \not{\!\!\!/}_{\alpha}^{q_0} \mathfrak{h}'$ terms in (8.36). Our aim is to prove that

(8.41)
$$\begin{split} \sum_{\mathfrak{h}\notin_{\alpha}^{q_{0}}\mathfrak{h}'} \left| \langle \mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}}^{M}), \mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}'}^{M}) \rangle_{s} \right| \lesssim \mathcal{N}(q_{0}) \mu^{2sq_{0}} \sum_{\mathfrak{h}\in\mathfrak{H}^{q_{0}}} \|u_{\bar{\alpha},\mathfrak{h}}^{M}\|_{\mathcal{H}^{s}}^{2} \\ + C_{q_{0}} \sum_{\mathfrak{h}\in\mathfrak{H}^{q_{0}}} \|u_{\bar{\alpha},\mathfrak{h}}^{M}\|_{\mathcal{H}^{s-1}}^{2}. \end{split}$$

Keeping the same notation used previously, we write

(8.42)
$$\sum_{\mathfrak{h}\notin_{\alpha}^{q_{0}}\mathfrak{h}'} \langle \mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}}^{M}), \mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}'}^{M}) \rangle_{s} = \sum_{\mathfrak{h}\in\mathfrak{H}^{q_{0}}} \sum_{\mathfrak{h}':\mathfrak{h}'\notin_{\alpha}^{q_{0}}\mathfrak{h}} \langle \mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}}^{M}), \mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}'}^{M}) \rangle_{s}.$$

By equation (E.4), there are $C_{\gamma,\beta}$ such that

(8.43)
$$\langle \mathcal{L}^{q_0} u^M_{\bar{\alpha},\mathfrak{h}}, \mathcal{L}^{q_0}(u^M_{\bar{\alpha},\mathfrak{h}'}) \rangle_s = \sum_{\gamma+\beta \leq s} C_{\gamma,\beta} \langle \partial^{\gamma}_{x_1} \partial^{\beta}_{x_2} \mathcal{L}^{q_0}(u^M_{\bar{\alpha},\mathfrak{h}}), \partial^{\gamma}_{x_1} \partial^{\beta}_{x_2} \mathcal{L}^{q_0}(u^M_{\bar{\alpha},\mathfrak{h}'}) \rangle_{L^2}.$$

We then use equation (7.6) and we have, for every γ, β such that $\gamma + \beta \leq s$

$$|\partial_{x_1}^{\gamma}\partial_{x_2}^{\beta}(\mathcal{L}^{q_0}u^M_{\bar{\alpha},\mathfrak{h}})| \leq \|(DF^{q_0})^{-1}\|_{\infty}^s \mathcal{L}^{q_0}(|\partial_{x_1}^{\gamma}\partial_{x_2}^{\beta}u^M_{\bar{\alpha},\mathfrak{h}}|) + \mathcal{L}^{q_0}(P^{q_0}_{s-1}u^M_{\bar{\alpha},\mathfrak{h}})$$

where $P_{s-1}^{q_0}$ is a differential operator of order s-1. By (5.17) $||(DF^{q_0})^{-1}||_{\infty}^s \leq C\mu^{sq_0}$. Clearly the same inequality holds for \mathfrak{h}' and we use this in (8.43) to obtain

$$\begin{split} &\sum_{\gamma+\beta\leq s} C_{\gamma,\beta} \left| \langle \partial_{x_{1}}^{\gamma} \partial_{x_{2}}^{\beta} \mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}}^{M}), \partial_{x_{1}}^{\gamma} \partial_{x_{2}}^{\beta} \mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}'}^{M}) \rangle_{L^{2}} \right| \\ &\lesssim \mu^{2sq_{0}} \sum_{\gamma+\beta=s} C_{\gamma,\beta} \langle \mathcal{L}^{q_{0}}(|\partial_{x_{1}}^{\gamma} \partial_{x_{2}}^{\beta} u_{\bar{\alpha},\mathfrak{h}}^{M}|), \mathcal{L}^{q_{0}}(|\partial_{x_{1}}^{\gamma} \partial_{x_{2}}^{\beta} u_{\bar{\alpha},\mathfrak{h}'}^{M}|) \rangle_{L^{2}} \\ &+ C_{q_{0}} \sum_{\gamma+\beta=s} \| \mathcal{L}^{q_{0}}(|\partial_{x_{1}}^{\gamma} \partial_{x_{2}}^{\beta} u_{\bar{\alpha},\mathfrak{h}}^{M}|) \|_{L^{2}} \| u_{\bar{\alpha},\mathfrak{h}'}^{M} \|_{\mathcal{H}^{s-1}} \\ &+ C_{q_{0}} \sum_{\gamma+\beta=s} \| \mathcal{L}^{q_{0}}(|\partial_{x_{1}}^{\gamma} \partial_{x_{2}}^{\beta} u_{\bar{\alpha},\mathfrak{h}'}^{M}|) \|_{L^{2}} \| u_{\bar{\alpha},\mathfrak{h}}^{M} \|_{\mathcal{H}^{s-1}} + C_{q_{0}} \| u_{\bar{\alpha},\mathfrak{h}}^{M} \|_{\mathcal{H}^{s-1}} \| u_{\bar{\alpha},\mathfrak{h}'}^{M} \|_{\mathcal{H}^{s-1}} \end{split}$$

It follows that, for each $\delta > 0$,

$$(8.44) \qquad \sum_{\gamma+\beta\leq s} C_{\gamma,\beta} \left| \langle \partial_{x_{1}}^{\gamma} \partial_{x_{2}}^{\beta} (\mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}}^{M}), \partial_{x_{1}}^{\gamma} \partial_{x_{2}}^{\beta} \mathcal{L}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}'}^{M}) \rangle_{L^{2}} \right| \leq \mu^{2sq_{0}} \sum_{\gamma+\beta=s} C_{\gamma,\beta} \langle \mathcal{L}^{q_{0}}(|\partial_{x_{1}}^{\gamma} \partial_{x_{2}}^{\beta} u_{\bar{\alpha},\mathfrak{h}}^{M}|), \mathcal{L}^{q_{0}}(|\partial_{x_{1}}^{\gamma} \partial_{x_{2}}^{\beta} u_{\bar{\alpha},\mathfrak{h}'}^{M}|) \rangle_{L^{2}} + \delta \left[\|u_{\bar{\alpha},\mathfrak{h}}^{M}\|_{\mathcal{H}^{s}}^{2} + \|u_{\bar{\alpha},\mathfrak{h}'}^{M}\|_{\mathcal{H}^{s}}^{2} \right] + C_{q_{0},s} \delta^{-1} \left[\|u_{\bar{\alpha},\mathfrak{h}}^{M}\|_{\mathcal{H}^{s-1}}^{2} + \|u_{\bar{\alpha},\mathfrak{h}'}^{M}\|_{\mathcal{H}^{s-1}}^{2} \right]$$

Since $u^{M}_{\bar{\alpha},\mathfrak{h}}$ and $u^{M}_{\bar{\alpha},\mathfrak{h}'}$ are supported on invertibility domains of F^{q_0} ,

(8.45)
$$\mathcal{L}^{q_0}|(\partial_{x_1}^{\gamma}\partial_{x_2}^{\beta}u^M_{\alpha,\tau}|) = \frac{|\partial_{x_1}^{\gamma}\partial_{x_2}^{\beta}u^M_{\alpha,\tau}| \circ \tau}{|\det DF^{q_0}| \circ \tau}, \quad \tau \in \{\mathfrak{h}, \mathfrak{h}'\}.$$

We define $\chi_{\tau} := |\partial_{x_1}^{\gamma} \partial_{x_2}^{\beta} u^M_{\alpha,\tau}| \circ \tau$ and $g_{\tau} := |\det DF^N| \circ \tau$ and we have

(8.46)
$$\langle \mathcal{L}^{q_0}(|\partial_{x_1}^{\gamma}\partial_{x_2}^{\beta}u_{\bar{\alpha},\mathfrak{h}}^{M}|), \mathcal{L}^{q_0}(|\partial_{x_1}^{\gamma}\partial_{x_2}^{\beta}u_{\bar{\alpha},\mathfrak{h}'}^{M}|)\rangle_{L^2} = \int_{\mathbb{T}^2} \frac{\chi_{\mathfrak{h}}\chi_{\mathfrak{h}'}}{\sqrt{g_{\mathfrak{h}}g_{\mathfrak{h}'}}\sqrt{g_{\mathfrak{h}}g_{\mathfrak{h}'}}} \\ \leq \frac{1}{2} \int_{\mathbb{T}^2} \frac{\chi_{\mathfrak{h}}^2}{g_{\mathfrak{h}'}g_{\mathfrak{h}}} + \frac{1}{2} \int_{\mathbb{T}^2} \frac{\chi_{\mathfrak{h}'}^2}{g_{\mathfrak{h}'}g_{\mathfrak{h}}},$$

where we used the elementary inequality $ab \leq \frac{1}{2}(a^2 + b^2)$ with $a = \frac{\chi_{\mathfrak{h}}}{\sqrt{g_{\mathfrak{h}}g_{\mathfrak{h}'}}}, b = \frac{\chi_{\mathfrak{h}'}}{\sqrt{g_{\mathfrak{h}}g_{\mathfrak{h}'}}}$. In order to obtain (8.41), we need to sum equation (8.44) over $\mathfrak{h} \in \mathfrak{H}^{q_0}$ and $\mathfrak{h}' \not/ \mathfrak{h}^{q_0}_{\alpha} \mathfrak{h}$. Let us begin with the first term. Consider one of the integrals in (8.46), for example the first one. Recalling $\mathcal{N}(q_0)$ as in Definition 4.2 and equation (7.12), Lemma 8.10 implies

(8.47)
$$\sum_{\mathfrak{h}} \sum_{\mathfrak{h}':\mathfrak{h}'\mathfrak{H}^{q_{0}}_{\alpha}\mathfrak{h}} \int_{\mathbb{T}^{2}} \frac{\chi^{2}_{\mathfrak{h}}}{g_{\mathfrak{h}'}g_{\mathfrak{h}}} \leq \sum_{\mathfrak{h}} \int_{\mathbb{T}^{2}} \frac{\chi^{2}_{\mathfrak{h}}}{g_{\mathfrak{h}}} \sum_{\mathfrak{h}':\mathfrak{h}'\mathfrak{H}^{M}_{\alpha}\mathfrak{h}} \frac{1}{g_{\mathfrak{h}'}} \leq \mathcal{N}(q_{0}) \sum_{\mathfrak{h}} \int_{\mathbb{T}^{2}} \frac{|\partial^{\gamma}_{x_{1}}\partial^{\beta}_{x_{2}}u^{M}_{\overline{\alpha},\mathfrak{h}}|^{2} \circ \mathfrak{h}}{|\det DF^{N}| \circ \mathfrak{h}}$$
$$= \mathcal{N}(q_{0}) \sum_{\mathfrak{h}} \|\mathcal{L}^{q_{0}}|\partial^{\gamma}_{x_{1}}\partial^{\beta}_{x_{2}}u^{M}_{\overline{\alpha},\mathfrak{h}}|^{2}\|_{L^{1}} = \mathcal{N}(q_{0}) \sum_{\mathfrak{h}} \|\partial^{\gamma}_{x_{1}}\partial^{\beta}_{x_{2}}u^{M}_{\overline{\alpha},\mathfrak{h}}\|^{2}_{L^{2}}.$$

By symmetry we have

(8.48)
$$\mu^{2sq_0} \sum_{\mathfrak{h}\mathfrak{H}^{q_0}_{\alpha}\mathfrak{h}'} \sum_{\gamma+\beta=s} C_{\gamma,\beta} \left| \langle \mathcal{L}^{q_0}(|\partial_{x_1}^{\gamma}\partial_{x_2}^{\beta}u_{\bar{\alpha},\mathfrak{h}}^M|), \mathcal{L}^{q_0}(|\partial_{x_1}^{\gamma}\partial_{x_2}^{\beta}u_{\bar{\alpha},\mathfrak{h}'}^M|) \rangle_{L^2} \right|$$
$$\leq \mu^{2sq_0} \mathcal{N}(q_0) \sum_{\mathfrak{h}\in\mathfrak{H}^{q_0}} \sum_{\gamma+\beta=s} \|\partial_{x_1}^{\gamma}\partial_{x_2}^{\beta}u_{\bar{\alpha},\mathfrak{h}}^M\|_{L^2}^2 \leq \mu^{2sq_0} \mathcal{N}(q_0) \sum_{\mathfrak{h}\in\mathfrak{H}^{q_0}} \|u_{\bar{\alpha},\mathfrak{h}}^M\|_{\mathcal{H}^s}^2,$$

which corresponds to the first addend of the r.h.s. of (8.41).

Summing the other terms of (8.44) over \mathfrak{h} yields (8.41) which, together with (8.37), conclude the proof of Lemma 8.8.

8.5. Proof of Theorem 8.2.

By (8.30) and Lemmata 8.7 and 8.8, we have

(8.49)
$$\begin{aligned} \|\mathcal{L}^{N}u\|_{\mathcal{H}^{s}}^{2} &\leq C_{q_{0}}\Theta_{s}\|u\|_{s+2}^{2} + C_{q_{0}}Q(M,s)\sqrt{\Theta_{s}}\|u\|_{s+2}\|u\|_{\mathcal{H}^{s}} \\ &+ (\mathcal{N}(q_{0})\mu^{2sq_{0}} + \delta)\sum_{\bar{\alpha}}\sum_{\mathfrak{h}\in\mathfrak{H}^{q_{0}}}\|u_{\bar{\alpha},\mathfrak{h}}^{M}\|_{\mathcal{H}^{s}}^{2} + C_{q_{0},s}\delta^{-1}\sum_{\bar{\alpha}}\sum_{\mathfrak{h}\in\mathfrak{H}^{q_{0}}}\|u_{\bar{\alpha},\mathfrak{h}}^{M}\|_{\mathcal{H}^{s-1}}^{2}. \end{aligned}$$

Recalling that $u_{\bar{\alpha},\mathfrak{h}}^M = \psi_{\bar{\alpha},\mathfrak{h}} \mathcal{L}^M u$, we can use equations (8.15) and (7.2) to write,⁵²

(8.50)
$$\sum_{\bar{\alpha}} \sum_{\mathfrak{h} \in \mathfrak{H}^{q_0}} \|u_{\bar{\alpha},\mathfrak{h}}^M\|_{\mathcal{H}^s}^2 = \sum_{\bar{\alpha}} \sum_{\mathfrak{h} \in \mathfrak{H}^{q_0}} \|\psi_{\bar{\alpha},\mathfrak{h}}\mathcal{L}^M u\|_{\mathcal{H}^s}^2 \le C_{\sharp} \|\mathcal{L}^M u\|_{\mathcal{H}^s}^2 + C_{\psi}(s)\|\mathcal{L}^M u\|_{L^1}^2 \le C_{\mathfrak{h}} \|\mathcal{L}^M u\|_{\mathcal{H}^s}^2 + C_{\psi}(s)\|\mathcal{L}^M u\|_{L^1}^2 \le C_{\mathfrak{h}} \|\mathcal{L}^M u\|_{\mathcal{H}^s}^2 + C_{\psi}(s)\|u\|_{L^1}^2 \le C_{\mathfrak{h}} \|\mathcal{L}^M u\|_{\mathcal{H}^s}^2 \le C_{\mathfrak{h}} \|\mathcal{L}^M u\|_{\mathcal{H}^s}^2 + C_{\psi}(s)\|u\|_{L^1}^2 \le C_{\mathfrak{h}} \|\mathcal{L}^M u\|_{\mathcal{H}^s}^2 \le C_{\mathfrak{h}} \|\mathcal{L}^M u\|_{\mathcal{H}^s}^2 + C_{\psi}(s)\|u\|_{L^1}^2 \le C_{\mathfrak{h}} \|\mathcal{L}^M u\|_{\mathcal{H}^s}^2 \le C_{\mathfrak{h}} \|\mathcal{L}^M u\|_{\mathcal{H}^s}^2 + C_{\psi}(s)\|u\|_{\mathcal{H}^s}^2 \le C_{\mathfrak{h}} \|\mathcal{L}^M u\|_{\mathcal{H}^s}^2 \le C_{\mathfrak{h}} \|\mathcal{$$

and

(8.51)
$$\sum_{\bar{\alpha}} \sum_{\mathfrak{h} \in \mathfrak{H}^{q_0}} \|u_{\bar{\alpha},\mathfrak{h}}^M\|_{\mathcal{H}^{s-1}}^2 \le C_{\sharp} \|\mathcal{L}^M u\|_{\mathcal{H}^{s-1}}^2 + C_{\psi}(s-1)\|\mathcal{L}^M u\|_{L^1}^2 \le \|\mathcal{L}^M u\|_{\mathcal{H}^{s-1}}^2 + C_{\psi}(s-1)\|u\|_{L^1}^2 \le \|\mathcal{L}^M u\|_{\mathcal{H}^{s-1}}^2 \le C_{\sharp} \|\mathcal{L}^M u\|_{\mathcal{H}^{$$

$$\leq \|\mathcal{L}^{M}1\|_{\infty}Q(M,s-1)\|u\|_{\mathcal{H}^{s-1}}^{2} + C_{\psi}(s-1)\|u\|_{L^{1}}^{2}.$$

Next, choosing $\delta = \mathcal{N}(q_0)$, using Lemma E.1 with $\varsigma = \mathcal{N}(q_0)^2 \mu^{2sq_0} Q(M, s)^{-1} C_{\epsilon,q_0}^{-1}$, substituting equations (8.50) and (8.51) in (8.49), setting

(8.52)
$$\overline{Q}(M,s) = \{\{1, \mathcal{N}(q_0)^{-3}\}^+ \mu^{2sq_0} Q(M,s)^2 C_{\epsilon,q_0} \mathbb{L}_M, C_{\psi}(s)\}^+,$$

and recalling (8.4) for the definition of \mathbb{L}_M , we obtain

$$(8.53) \qquad \begin{aligned} \|\mathcal{L}^{N}u\|_{\mathcal{H}^{s}}^{2} &\leq C_{\sharp}\mathbb{L}_{M}\mathcal{N}(q_{0})\mu^{2sN}\|u\|_{\mathcal{H}^{s}}^{2} + C_{q_{0}}\Theta_{s}\|u\|_{s+2}^{2} \\ &+ C_{q_{0}}Q(M,s)\sqrt{\Theta_{s}}\|u\|_{s+2}\|u\|_{\mathcal{H}^{s}} + \overline{Q}(M,s)\|u\|_{L^{1}}^{2} \\ &\leq C_{\sharp}\mathbb{L}_{M}\mathcal{N}(q_{0})\mu^{2sN}\|u\|_{\mathcal{H}^{s}}^{2} + C_{\epsilon,q_{0}}\left[Q(M,s)^{2}\mathbb{L}_{M}\Theta_{s}\mathcal{N}(q_{0})^{-1} + \overline{Q}(M,s)\right]\|u\|_{s+2}^{2} \end{aligned}$$

where, in the last line, we used (6.2) to estimate $||u||_{L^1} \leq C_{\sharp} ||u||_{s+2}$. Recalling (8.52), it follows

$$\|\mathcal{L}^{N}u\|_{\mathcal{H}^{s}} \leq C_{\sharp} \left(\sqrt{[\mathbb{L}_{M}\mathcal{N}(q_{0})]^{\frac{1}{N}}\mu^{2s}}\right)^{N} \|u\|_{\mathcal{H}^{s}} + C_{\epsilon,q_{0}}\sqrt{\overline{Q}(M,s)\Theta_{s}}\|u\|_{s+2},$$

from which, by equations (8.31), (8.15) and Lemma 7.1, we obtain (8.7) in the case s > 1.

8.6. Fourier basic estimate: case SVPH^{\ddagger}. To prove Theorem 8.3 we must improve the constant $K_1(s, M_{\xi})$ in Proposition 8.6 for t = 2. This is done in the following proposition.

Proposition 8.11. Under the assumption of Proposition 8.6 if, in addition, the map F is a $SVPH^{\sharp}$ and satisfies condition (8.8) for a uniform constant $C_5 > 0$, then there exist $C_{n_0,q_0,\epsilon}, \beta_1, \beta_2 > 0$ such that the conclusions of Proposition 8.6 hold with

(8.54)
$$K_1(2, M_{\xi}) \le C_{n_0, q_0, \epsilon} (\ln \chi_u^{-1})^{\beta_1} \mu^{\beta_2 \ln \chi_u^{-1}} \vartheta(\xi^{\perp})^{-7}.$$

⁵²We also use repeatedly $\|\mathcal{L}^n u\|_{L^1} \le \|u\|_{L^1}$.

Proof. We follow the proof of Proposition 8.6 word by word until the definition of \overline{m}_{ℓ} which we define here differently in the following way.

Let $\overline{m}_{\ell}(z, \tilde{\mathfrak{h}} \circ \mathfrak{h}, n_{\star})$ be the smallest integer satisfying (5.40), for $z \in \gamma_{\ell}$, with $n_{\star} = \{\hat{n}, \min\{c_2^{-}\log \chi_u^{-1}, m\}\}^+$, where m was defined right after (8.20). In addition, set $\overline{m}_{\ell}(\tilde{\mathfrak{h}}) = \sup_{z \in \gamma_{\ell}} \overline{m}_{\ell}(z, \tilde{\mathfrak{h}} \circ \mathfrak{h}, n_{\star})$ and define

(8.55)
$$M_{\xi} \equiv M_{\xi,2} = \sup_{\ell} \sup_{\tilde{\mathfrak{h}} \in \mathfrak{H}^{\infty}} \overline{m}_{\ell}(\tilde{\mathfrak{h}}).$$

We can follow the proof of Proposition 8.6 literally and we see that the main task is to estimate $\sum_{\hat{\mathfrak{h}}\in\hat{\mathfrak{H}}_{\alpha}} \|\Psi_{\hat{\nu}_{\ell}}(\partial^{\eta_1}\psi_{\alpha}) \circ F^{\overline{m}(\hat{\mathfrak{h}})}\|_{\mathcal{C}^2_{\hat{\nu}_{\ell}}}$ in (8.26), and so, to have a sharper bound of $A_u(t, M_{\xi}, m)$ for $\tau = 2$. To this goal, we use the improvements Proposition 6.5 and Lemma 5.22 instead of Proposition 6.4 and Lemma 5.19 respectively to get, by (6.22), setting $I_{\hat{\nu}_{\ell}} = h_{\ell \bar{m}}(I_{q_0})$,

(8.56)
$$\tilde{C}(2,\overline{m},m) \le C^3_{\mu,\overline{m}} \mu^{4\overline{m}} \sup_{\zeta \in \hat{\nu}_{\ell}(I_{\hat{\nu}_{\ell}})} (1 + C_F \lambda^+_{\overline{m}}(\zeta)) \{\lambda^+_m(F^{\overline{m}-m}(\zeta)) + C_{q_0} \Delta_{\tilde{\gamma}}\},$$

and, accordingly, we have equation (8.26) with

(8.57)
$$A_{u}(\tau, M_{\xi}, m) := \begin{cases} C_{\sharp} \mathbb{I}_{\tilde{\gamma}, M_{\xi}} \mathbb{J}_{\tilde{\gamma}, M_{\xi}} & \tau = 0\\ C_{\sharp} (\mathbb{I}_{\tilde{\gamma}, M_{\xi}})^{2} \mathbb{J}_{\tilde{\gamma}, M_{\xi}} & \tau = 1\\ C_{\mu, M_{\xi}}^{4} \mu^{5M_{\xi}} (1 + C_{F} \mu^{m} \vartheta_{\hat{\nu}_{0}}^{-1}) O_{m}^{\star} \mathbb{I}_{\tilde{\gamma}, m}^{3} \mathbb{J}_{\tilde{\gamma}, M_{\xi}} & \tau = 2 \end{cases}$$

First note that, by (8.12), (8.11) and (G.2) we can choose b large enough so that $b \ln \chi_u^{-1} > m$ and hence

(8.58)
$$\|\bar{\psi}_{\bar{\alpha},\mathfrak{h}}\|_{\mathcal{C}^{s}} \leq C_{\epsilon,q_{0}} C_{\mu,\mathrm{b}\ln\chi_{u}^{-1}}^{4s} \mu^{4s\mathrm{b}\ln\chi_{u}^{-1}}.$$

Next, let $g_{\alpha} = \partial^{\eta_1} \bar{\psi}_{\bar{\alpha},\mathfrak{h}}$, and $G_{\alpha}(s) = g_{\alpha} \circ \tilde{\gamma}_{\ell} \circ h_{\ell,\overline{m}}(s)$. Then, since $F^{\overline{m}} \hat{\nu}_{\ell} = \tilde{\gamma}_{\ell} \circ h_{\ell,\overline{m}}$, we have $g_{\alpha} \circ F^{\overline{m}} \circ \hat{\nu}_{\ell} = G_{\alpha}$ and if we choose

$$\begin{aligned} G'_{\alpha} &= \langle \nabla g_{\alpha} \circ \tilde{\gamma}_{\ell} \circ h_{\ell,\overline{m}}, \tilde{\gamma}'_{\ell} \circ h_{\ell,\overline{m}} h'_{\ell,\overline{m}} \rangle \\ G''_{\alpha} &= \langle (D^2 g_{\alpha}) \circ \tilde{\gamma}'_{\ell} \circ h_{\ell,\overline{m}} \cdot \tilde{\gamma}'_{\ell} \circ h_{\ell,\overline{m}}, \tilde{\gamma}'_{\ell} \circ h_{\ell,\overline{m}} \rangle (h'_{\ell,\overline{m}})^2 \\ &+ \langle \nabla g_{\alpha} \circ \tilde{\gamma}_{\ell} \circ h_{\ell,\overline{m}}, \tilde{\gamma}''_{\ell} \circ h_{\ell,\overline{m}} (h'_{\ell,\overline{m}})^2 + \tilde{\gamma}'_{\ell} \circ h_{\ell,\overline{m}} h''_{\ell,\overline{m}} \rangle. \end{aligned}$$

Note that by (G.1) and the first of (8.28) we have

$$\|\tilde{\gamma}_{\ell}^{\prime\prime}\|_{\infty} \leq C_{q_0,\epsilon} \mu^{11m} C_{\mu,m}^2.$$

Moreover, (8.8) and (5.20), imply $C_F \leq C_{\sharp}\chi_u$. We can use this to compute, using (8.28), (5.41),⁵³

(8.59)
$$\begin{aligned} \Delta_{\tilde{\gamma}} &\leq C_{q_0,\epsilon} \mu^{11m} C_{\mu,m}^2 \\ M_{m,q_0} &\leq C_{q_0,\epsilon} C_{\mu,m}^2 \mu^{15m} \vartheta_{\tilde{\gamma}}^{-1} \\ \overline{M}_{m,q_0} &\leq C_{q_0,\epsilon} C_{\mu,m}^4 \mu^{32m} \vartheta_{\tilde{\gamma}}^{-2} \end{aligned}$$

Accordingly, using also (5.42) (with $h_{\overline{m}} = h_{\overline{m},\ell}$), (8.28), (8.58) and (8.8), we have

$$\begin{aligned} \| (\partial^{2} \bar{\psi}_{\bar{\alpha}, \mathfrak{h}}) \circ F^{\overline{m}(\hat{\mathfrak{h}})} \|_{\mathcal{C}^{0}_{\bar{\nu}_{\ell}}} &\leq C_{\epsilon, q_{0}} C^{8}_{\mu, \mathrm{b} \ln \chi^{-1}_{u}} \mu^{8\mathrm{b} \ln \chi^{-1}_{u}} \\ \| (\partial \bar{\psi}_{\bar{\alpha}, \mathfrak{h}}) \circ F^{\overline{m}(\hat{\mathfrak{h}})} \|_{\mathcal{C}^{1}_{\bar{\nu}_{\ell}}} &\leq C_{\epsilon, q_{0}} C^{4}_{\mu, \mathrm{b} \ln \chi^{-1}_{u}} \vartheta^{-1}_{\bar{\gamma}} \mu^{M_{\xi} + 4s\mathrm{b} \ln \chi^{-1}_{u}} \\ (8.60) \qquad \| \bar{\psi}_{\bar{\alpha}, \mathfrak{h}} \circ F^{\overline{m}(\hat{\mathfrak{h}})} \|_{\mathcal{C}^{2}_{\bar{\nu}_{\ell}}} &\leq C_{\epsilon, q_{0}} C^{8}_{\mu, \mathrm{b} \ln \chi^{-1}_{u}} \mu^{3M_{\xi} + 24\mathrm{b} \ln \chi^{-1}_{u}} \vartheta^{-1}_{\bar{\gamma}} \left\{ \vartheta^{-1}_{\bar{\gamma}} + M_{m, n_{0}} \right\} \\ &\leq C_{\epsilon, q_{0}} C^{8}_{\mu, \mathrm{b} \ln \chi^{-1}_{u}} \mu^{3M_{\xi} + 25\mathrm{b} \ln \chi^{-1}_{u}} \vartheta^{-1}_{\bar{\gamma}} \left\{ C^{2}_{\mu, n} \mu^{15m} + C_{\mu, m} \mu^{6m} \vartheta^{-1}_{\bar{\gamma}} \right\} \\ &\leq C_{\epsilon, q_{0}} C^{10}_{\mu, \mathrm{b} \ln \chi^{-1}_{u}} \mu^{3M_{\xi} + 40\mathrm{b} \ln \chi^{-1}_{u}} \vartheta^{-2}_{\bar{\gamma}}. \end{aligned}$$

⁵³Recall that $q_0 \ge n_0$.

Since $(\Psi_{\hat{\nu}_{\ell}}G_{\alpha})'' = \Psi_{\hat{\nu}_{\ell}}''G_{\alpha} + 2\Psi_{\hat{\nu}_{\ell}}'G_{\alpha}' + \Psi_{\hat{\nu}_{\ell}}G_{\alpha}''$, by equations (8.26), (8.57), (5.60) and (8.60)

(8.61)
$$\sum_{\hat{\mathfrak{h}}\in\hat{\mathfrak{H}}_{\alpha}} \|\Psi_{\hat{\nu}_{\ell}}(\partial^{\eta_{1}}\psi_{\alpha})\circ F^{\overline{m}(\hat{\mathfrak{h}})}\|_{\mathcal{C}^{2}_{\hat{\nu}_{\ell}}} \leq C_{\epsilon,q_{0}}C^{10}_{\mu,\operatorname{b}\ln\chi_{u}^{-1}}\mu^{3M_{\xi}+40\operatorname{b}\ln\chi_{u}^{-1}}\Big\{A_{u}(2,M_{\xi},m) + A_{u}(1,M_{\xi},m)\vartheta_{\tilde{\gamma}}^{-1} + A_{u}(0,M_{\xi},m)\vartheta_{\tilde{\gamma}}^{-2}\Big\}.$$

To conclude, we need to relate all the quantities to $\vartheta_{\tilde{\gamma}}$. First we notice that, by (8.28) and recalling (C.43), it follows that for each $\zeta \in \mathbb{T}$.

(8.62)
$$\lambda_m^+ \circ F^{\overline{m}-m} \circ \hat{\nu}_{\ell}(\zeta) \le C_{\sharp} \vartheta_{\tilde{\gamma}}^{-1} \mu^m.$$

Notice that the expansion is almost constant for the points of interest. Indeed, by Lemma 5.17, for any $p, p_* \in \overline{U} = \mathcal{U}_{\alpha,\mathfrak{h}} \cap \operatorname{Im}(\hat{\nu}_{\ell})$ and $n \leq M_{\xi}$,

$$\lambda_{n}^{+}(p) \leq \lambda_{n}^{+}(p_{*})e^{\mu^{M_{\xi}}C_{\mu,M_{\xi}}\|p-p_{*}\|} \leq \lambda_{n}^{+}(p_{*})e^{\mu^{M_{\xi}}C_{\mu,M_{\xi}}\mathrm{d}\epsilon\delta_{q_{0}}(p_{*})}.$$

If we choose b so that $b \ln \chi_u^{-1} \ge M_{\xi}$, then (8.11) and (G.2) imply

(8.63)
$$\lambda_n^+(p) \le C_{\sharp} \lambda_n^+(p_*).$$

Next, recall the choice $n_{\star} = \{\hat{n}, \min\{c_2^{-}\log\chi_u^{-1}, m\}\}^+$ in Lemma 5.14. To continue we need to check the conditions (5.40). Recalling (8.59), the first of the (5.40) is implied by

$$\lambda_{\overline{m}-m}^{-}(\hat{\nu}_{m} \circ h_{\overline{m}-m}(t)) \ge c_{\flat}c_{n_{\star}}^{\overline{m}}\mu^{2\overline{m}+15m-3n_{\star}}C_{q_{0},\epsilon}C_{\mu,m}\vartheta_{\tilde{\gamma}}^{-1}$$

The second is implied by

$$c_{\flat}^{2}b_{n_{\star}}^{n_{\star}}\mu^{2\overline{m}+2n_{\star}}\lambda_{\overline{m}-m}^{-}(\gamma \circ h_{\overline{m}-m}(t))^{-1}C_{q_{0},\epsilon}C_{\mu,m}\mu^{15m}\vartheta_{\bar{\gamma}}^{-1} \leq \frac{1}{2}b_{n_{\star}}^{2n_{\star}}\mu^{6n_{\star}}C_{3}$$

which follows by

$$\lambda_{\overline{m}-m}^{-}(\gamma \circ h_{\overline{m}-m}(t)) \ge 2c_{\flat}^{2}\mu^{2\overline{m}+15m-4n_{\star}}C_{q_{0},\epsilon}C_{\mu,m}\vartheta_{\tilde{\gamma}}^{-1}.$$

And the third of (5.40) is implied by

$$c_{\flat}^{2} c_{n_{\star}}^{\overline{m}} \mu^{3\overline{m}} (\lambda_{\overline{m}-m}^{-}(\gamma \circ h_{\overline{m}-m}(t)))^{-1} C_{q_{0},\epsilon} C_{\mu,m}^{4} \mu^{32m} \vartheta_{\tilde{\gamma}}^{-2} \leq \frac{1}{2} C_{\mu,n_{\star}}^{2} \mu^{6n_{\star}} C_{3}^{2}$$

which follows by

$$\lambda_{\overline{m}-m}^{-}(\gamma \circ h_{\overline{m}-m}(t)) \geq 2c_{\flat}^{2}c_{n_{\star}}^{\overline{m}}\mu^{3\overline{m}+32m-6n_{\star}}C_{q_{0},\epsilon}C_{\mu,m}^{2}\vartheta_{\tilde{\gamma}}^{-2}.$$

The above, due to (8.62) and recalling (5.16), are implied by

$$\begin{split} \lambda^{\overline{m}}_{-} &\geq C_{\sharp} m c^{\overline{m}}_{n_{\star}} \mu^{15\overline{m}} \vartheta^{-2}_{\tilde{\gamma}} \\ \lambda^{\overline{m}}_{-} &\geq C_{\sharp} m \mu^{14\overline{m}} \vartheta^{-2}_{\tilde{\gamma}} \\ \lambda^{\overline{m}}_{-} &\geq C_{\sharp} m^2 c^{\overline{m}}_{n_{\star}} \mu^{30\overline{m}} \vartheta^{-3}_{\tilde{\gamma}}, \end{split}$$

which, in turn, are satisfied if

(8.64)
$$\left(\frac{\lambda_{-}}{\mu^{30}c_{n_{\star}}}\right)^{\overline{m}} \ge C_{\sharp}m^{2}\vartheta_{\tilde{\gamma}}^{-3}$$

Recalling (5.1) and noting that $B_n \geq c_n^3$, the above is implied by $2^{\overline{m}} \geq C_{\sharp} m^2 \vartheta_{\tilde{\gamma}}^{-3}$ which implies $\overline{m} \leq C_6 \ln \vartheta(\xi^{\perp}) \leq C_6 \ln \chi_u^{-1}$ for some constant C_6 , hence

(8.65)
$$M_{\xi} \le C_6 \ln \vartheta(\xi^{\perp})^{-1}.$$

Note that this implies $\sup_{\xi^{\perp} \notin \mathbf{C}_u} M_{\xi} \leq C_6 \ln \chi_u^{-1} < \infty$, which proves (8.17) for t = 2.

The above allows us to make the choice $\mathbb{b} = C_6$. In addition, since $c_{n_\star} \leq e^{c_\sharp / \ln \vartheta_{\tilde{\gamma}}^{-1}}$ we have $c_{n_\star}^{\overline{m}} \leq C_\sharp$. Hence, we can choose \overline{m} so that

(8.66)
$$\lambda_{\overline{m}}^+ \circ \hat{\nu}_{\ell}(s) \le C_{\mu,\overline{m}}^2 \mu^{30\overline{m}} \vartheta(\xi^{\perp})^{-3}.$$

By the above, (8.56), (8.62) and (8.59), we then have the estimate of the first term in the sum (8.24),

(8.67)
$$\tilde{C}(2,\overline{m}(\mathfrak{h}),m) \le C_{q_0,\epsilon} C_{\mu,M_{\xi}}^7 \mu^{45M_{\xi}} \vartheta(\xi^{\perp})^{-3}$$

We now proceed with the bound of the other terms in (8.24). We want to use Lemma 5.19, hence we need to estimate the quantities in (5.60). Our first task is to study $\vartheta_{\hat{\nu}_0,m}(s)$. By definition $\hat{\nu}_{n_0}(s) = (\sigma(s), s)$ for some function σ and (5.39) implies that $\vartheta_{\hat{\nu}_0,m}(s) = |\sigma'(s)|^{-1}$. Since, using the notation of Lemma 5.13, $F^{n_0}\hat{\nu}_{n_0} = \hat{\nu}_0 \circ \bar{h}_{n_0}$, we have

$$C_{n_0}|\pi_2(\hat{\nu}'_0 \circ \bar{h}_{n_0}(s))| \ge |\pi_2([D_{\hat{\nu}_{n_0}(s)}F^{n_0}]^{-1}\bar{\nu}'_0 \circ \bar{h}_{n_0}(s)\bar{h}'_{n_0}(s))| = |\sigma'(s)| \ge C_{n_0}|\pi_2(\bar{\nu}'_0 \circ \bar{h}_{n_0}(s))|.$$

By the proof of Lemma G.1, see (G.5), we have that $\pi_2(\tilde{\gamma}'_0) = \pi_2(\nu'_0)$ is monotone, hence so is $\pi_2(\hat{\nu}'_0(s)) = \pi_2(\nu'_0 \circ \bar{h}_{n_0}(s))$. Thus, if $|t| \ge |s|$, we have

(8.68)
$$|\sigma'(s)| \le C_{n_0} |\pi_2(\hat{\nu}'_0 \circ \bar{h}_{n_0}(s))| \le C_{n_0} |\pi_2(\hat{\nu}'_0 \circ \bar{h}_{n_0}(t))| \le C_{n_0} |\sigma'(t)|.$$

Also the proof of Lemma G.1 implies that, for $|s| \ge \frac{1}{4}$, $|\sigma'(s)| \le c_{\sharp}$. Accordingly,

$$\mathbb{J}_{\tilde{\gamma},M_{\xi}} = \int_{\mathbb{T}} \left[\vartheta_{\hat{\nu}_{n_{0}},M_{\xi}}(s) \right]^{-1} ds \leq C_{\sharp} + \int_{-\frac{1}{4}}^{\frac{1}{4}} \sup_{|s-t| \leq c_{\sharp}\chi_{u} \ln \chi_{u}^{-1}} |\sigma'(t)| dt \\
\leq C_{\sharp} + \int_{-c_{\sharp}\chi_{u} \ln \chi_{u}^{-1}}^{c_{\sharp}\chi_{u} \ln \chi_{u}^{-1}} \vartheta(\xi^{\perp})^{-1} + C_{n_{0}} \int_{-\frac{1}{4}}^{-c_{\sharp}\chi_{u} \ln \chi_{u}^{-1}} |\sigma'(s+c_{\sharp}\chi_{u} \ln \chi_{u}^{-1})| ds \\
+ C_{n_{0}} \int_{c_{\sharp}\chi_{u} \ln \chi_{u}^{-1}}^{-\frac{1}{4}} |\sigma'(s-c_{\sharp}\chi_{u} \ln \chi_{u}^{-1})| ds \\
\leq C_{n_{0}} \left[1 + \vartheta(\xi^{\perp})^{-1}\chi_{u} \ln \chi_{u}^{-1} \right] \leq C_{n_{0}} \ln \chi_{u}^{-1},$$

since, $\int_{\mathbb{T}} |\sigma'| = \left| \int_{\mathbb{T}} \sigma' \right| \leq 1$, since $\hat{\nu}'_{n_0} \notin \mathbf{C}_c$ and the curve does not wrap around the torus horizontally. By the proof of Lemma G.1 we have that $\vartheta_{\hat{\nu}_0}(s)$ is monotone for $|s| \leq \frac{1}{4}$.

Next, by (8.59) we have

$$\begin{split} \mathbb{I}_{\tilde{\gamma},M_{\xi}} &\leq C^{3}_{\mu,M_{\xi}} \mu^{16M_{\xi}} \vartheta_{\tilde{\gamma}}^{-1} \\ O^{\star}_{M_{\xi}} &\leq C_{q_{0},\epsilon} C^{8}_{\mu,M_{\xi}} \mu^{60M_{\xi}} \vartheta_{\tilde{\gamma}}^{-1}. \end{split}$$

Using the above estimates in (8.57) it follows

(8.70)
$$\begin{aligned} A_{u}(0, M_{\xi}, m) &\leq C_{n_{0}} C_{\mu, M_{\xi}}^{3} \mu^{16M_{\xi}} \vartheta_{\tilde{\gamma}}^{-1} \ln \chi_{u}^{-1}, \\ A_{u}(1, M_{\xi}, m) &\leq C_{n_{0}} C_{\mu, M_{\xi}}^{6} \mu^{32M_{\xi}} \vartheta_{\tilde{\gamma}}^{-2} \ln \chi_{u}^{-1} \\ A_{u}(2, M_{\xi}, m) &\leq C_{n_{0}, q_{0}, \epsilon} C_{\mu, M_{\xi}}^{21} \mu^{114M_{\xi}} \vartheta_{\tilde{\gamma}}^{-4} \ln \chi_{u}^{-1}. \end{aligned}$$

Substituting the above in (8.61) yields

(8.71)
$$\sum_{\hat{\mathfrak{h}}\in\hat{\mathfrak{H}}_{\alpha}} \|\Psi_{\hat{\nu}_{\ell}}(\partial^{\eta_{1}}\psi_{\alpha})\circ F^{\overline{m}(\hat{\mathfrak{h}})}\|_{\mathcal{C}^{2}_{\hat{\nu}_{\ell}}} \leq C_{n_{0},q_{0},\epsilon}C^{10}_{\mu,\mathfrak{h}\ln\chi_{u}^{-1}}C^{21}_{\mu,M_{\xi}} \times \mu^{114M_{\xi}+4\mathfrak{h}\ln\chi_{u}^{-1}}\vartheta(\xi^{\perp})^{-4}\ln\chi_{u}^{-1}.$$

By (8.24), (8.25), (8.67), (8.71), ams since $|I_0| \leq C_{q_0,\epsilon} C_{\mu,b \ln \chi_u^{-1}}^{-1} \mu^{-b \ln \chi_u^{-1}}$, there exist $\beta_1, \beta_2 > 0$ such that

$$\left| \mathcal{F} \partial_{x_j}^2 \mathcal{L}^{q_0}(u_{\bar{\alpha},\mathfrak{h}}^{M_{\xi}}) \right| \le C_{n_0,q_0,\epsilon} C_{\mu,\mathrm{b}\ln\chi_u^{-1}}^9 C_{\mu,M_{\xi}}^{\beta_1} \mu^{\beta_2 M_{\xi} + 3\mathrm{b}\ln\chi_u^{-1}} \vartheta(\xi^{\perp})^{-7} \ln\chi_u^{-1} \|u\|_2,$$

which concludes the proof of (8.54), recalling equations (8.18), (8.65).

Thanks to the previous result, we can now conclude the proof of Theorem 8.3.

8.7. Proof of Theorem 8.3.

Under the additional assumption of F being a SVPH^{\ddagger} and satisfying condition (8.8), we are going to prove (8.9) following word by word the SVPH case with the following three additions: 1) By Remark 8.5 we have a sharper estimate of $C_{\psi}(1)$ given in Lemma 8.4:

$$C_{\psi}(1) \le C_{\epsilon,q_0} C_{\mu,c_{\sharp} \ln \chi_u^{-1}} \chi_u^{c_{\sharp} \ln \mu}.$$

2) Here we improve the constant Θ_1 of Lemma 8.7 for s = 1. In the proof of Lemma 8.7, after (8.33), we add the following computation for s = 1: For any R > 0 let $B_R = \{\xi \in \mathbb{Z}^2 : ||\xi|| \leq R\}$ and $B_R^c = \mathbb{Z}^2 \setminus B_R$. Then

$$\sum_{\xi \in \mathbb{Z}^2} |\langle \xi \rangle \mathbb{1}_{\mathcal{Z}_u^c} \mathcal{FL}^{q_0}(u_{\bar{\alpha}, \mathfrak{h}'}^M)|^2 = \sum_{\xi \in \mathcal{Z}_u^c \cap B_R} \langle \xi \rangle^{-2} |\langle \xi \rangle^2 \mathcal{FL}^{q_0}(u_{\bar{\alpha}, \mathfrak{h}'}^M)|^2 + \sum_{\xi \in \mathcal{Z}_u^c \cap B_R^c} \langle \xi \rangle^{-3} |\langle \xi \rangle^2 \mathcal{FL}^{q_0}(u_{\bar{\alpha}, \mathfrak{h}'}^M)| |\langle \xi \rangle^3 \mathcal{FL}^{q_0}(u_{\bar{\alpha}, \mathfrak{h}'}^M)|.$$

Arguing as in the proof of Proposition 8.6 and by (8.33) yield

(8.72)
$$\sum_{\xi \in \mathcal{Z}_u^c \cap B_R} |\langle \xi \rangle \mathcal{FL}^{q_0}(u_{\bar{\alpha},\mathfrak{h}'}^M)|^2 \le C_{\mu,M}^2 \mu^{2M} \|u\|_2^2 \sum_{\xi \in \mathcal{Z}_u^c \cap B_R} \langle \xi \rangle^{-2} K_1(2, M_\xi)^2$$

and

(8.73)
$$\sum_{\xi \in \mathcal{Z}_{u}^{c} \cap B_{R}^{c}} |\langle \xi \rangle \mathcal{FL}^{q_{0}}(u_{\bar{\alpha},\mathfrak{h}'}^{M})|^{2} \leq C_{\mu,M}^{2} \mu^{2M} ||u||_{2} ||u||_{3} \sum_{\xi \in \mathcal{Z}_{u}^{c} \cap B_{R}^{c}} \langle \xi \rangle^{-3} K_{1}(2, M_{\xi}) K_{1}(3, M_{\xi}).$$

We use the estimate of $K_1(2, M_{\xi})$ in (8.54) for the sum in (8.72), with $\vartheta(\xi^{\perp}) = \rho(\xi^{\perp})$, since $\xi \in \mathcal{Z}_u^c$,

(8.74)
$$\sum_{\xi \in \mathcal{Z}_{u}^{c} \cap B_{R}} \langle \xi \rangle^{-2} K_{1}(2, M_{\xi})^{2} \leq C_{n_{0}, q_{0}, \epsilon} (\ln \chi_{u}^{-1})^{c_{\sharp}} \mu^{c_{\sharp} \ln \chi_{u}^{-1}} \sum_{\xi \in \mathcal{Z}_{u}^{c} \cap B_{R}} \langle \xi \rangle^{-2} \rho(\xi^{\perp})^{-14} \leq C_{n_{0}, q_{0}, \epsilon} (\ln \chi_{u}^{-1})^{c_{\sharp}} \mu^{c_{\sharp} \ln \chi_{u}^{-1}} \chi_{u}^{-13} \log R,$$

since

(8.75)
$$\sum_{\xi \in \mathcal{Z}_u^c \cap B_R} \langle \xi \rangle^{-2} \rho(\xi^{\perp})^{-14} \le \int_0^R \int_{\{\tan \theta > \chi_u\}} \frac{1}{1 + \rho^2} \frac{1}{(\tan \theta)^{14}} \rho d\rho d\theta \lesssim \chi_u^{-13} \log R.$$

Similarly, for the sum in (8.73), we have

(8.76)

$$\sum_{\xi \in \mathcal{Z}_{u}^{c} \cap B_{R}^{c}} \langle \xi \rangle^{-3} K_{1}(2, M_{\xi}) K_{1}(3, M_{\xi})$$

$$\leq C_{n_{0},q_{0},\epsilon} \mu^{c_{\sharp} \ln \chi_{u}^{-1}} \chi_{u}^{-c_{\sharp}} \sum_{\xi \in \mathcal{Z}_{u}^{c} \cap B_{R}^{c}} \langle \xi \rangle^{-3} \rho(\xi^{\perp})^{-7} \Lambda^{c_{\sharp} M_{\xi}}$$

$$\leq C_{n_{0},q_{0},\epsilon} \mu^{c_{\sharp} \ln \chi_{u}^{-1}} \chi_{u}^{-c_{\sharp}} R^{-1}.$$

Choosing $R = \chi_u^{-c_{\sharp}} \Lambda^{c_{\sharp}M_{\xi}}$ by (8.72) and (8.73) we have the following estimate:

(8.77)
$$\sum_{\xi \in \mathbb{Z}^2} |\langle \xi \rangle \mathbb{1}_{\mathcal{Z}_u^c} \mathcal{FL}^{q_0}(u^M_{\bar{\alpha}, \mathfrak{h}'})|^2 \le C_{n_0, q_0, \epsilon} C^2_{\mu, M} \mu^{2M} (\ln \chi_u^{-1})^{c_\sharp} \chi_u^{-13 - c_\sharp \ln \mu} \|u\|_3^2.$$

In conclusion, under the assumption that the map is a SVPH[#] and satisfies condition (8.8), the constant Θ_1 of Lemma 8.7 for s = 1 becomes

(8.78)
$$\Theta_1 = C_{n_0,q_0,\epsilon} C_{\mu,M}^2 \mu^{2M} (\ln \chi_u^{-1})^{c_\sharp} \chi_u^{-13-c_\sharp \ln \mu}.$$

3) To estimate the right hand side of (8.40) in the case s = 1 we use

$$\sum_{\xi \in \mathcal{Z}_u \cap Z(\mathfrak{h}) \cap B_R} \langle \xi \rangle^{-2} \chi_u^{-14} \le C_{\sharp} \chi_u^{-13} \log R$$

instead of (8.75) where $Z(\mathfrak{h})$ is defined in (8.38). By Lemma 7.1, $Q(M,1) \leq C^3_{\mu,M} \mu^{3M}$. Finally, using (8.52), (8.78), we can find $\beta_3 > 0$ such that

$$\sqrt{Q}(M,1)\Theta_1 \le C_{\epsilon,q_0} C_{\mu,M}^{\beta_3} \mu^{\beta_3 M} (\ln \chi_u^{-1})^{\beta_3} \chi_u^{\frac{-13}{2} - c_{\sharp} \ln \mu^{-1}} \mathbb{L}_M^{\frac{1}{2}},$$

which concludes the proof of Theorem 8.3.

9. The final Lasota-Yorke Inequality

We state and prove our main technical Theorem which implies the Theorems stated in section 4. For each integer $1 \le s \le r - 1$ we define the following norm

$$\|\cdot\|_{s,*} := \|\cdot\|_{\mathcal{H}^s} + \|\cdot\|_{s+2}.$$

Theorem 9.1. Let $F \in C^r(\mathbb{T}^2, \mathbb{T}^2)$ be a SVPH and $\alpha = \frac{\log(\lambda - \mu^{-2})}{\log(\lambda_+)}$. Let \overline{m}_{χ_u} be as in (8.17), n_0 as in (8.5) and $C_1 > 0$ provided in Theorem 8.2. We assume that there exist: constants $c, K > 0, \nu_0 \in (0, 1)$, integer $n_1 \ge n_0$, and uniform constants $\tau_0 \ge 1, \kappa_1 \ge \kappa_0 \in \mathbb{N}$ such that, for some $1 \le s \le r - 3$,⁵⁴

(9.1)
$$\sup_{m \le n} \|\mathcal{L}^m 1\|_{\infty} \le K \mu^{cn^{\tau_0}}, \qquad \forall n < \kappa_1 n_1 + \overline{m}_{\chi_u},$$

(9.2)
$$\left\{\mu^{\zeta_s}\lambda_-^{-1}, \sqrt{\widetilde{\mathcal{N}}_F(\lceil \alpha n_1 \rceil)}\mu^{\alpha_s n_1^{\tau_0}\kappa_1^{\tau_0-1} + \beta_s \overline{m}_{\chi_u}^{\tau_0}\kappa_0^{-1}}\right\}^+ \le \nu_0 < 1$$

(9.3)
$$(C_1 K)^{\frac{1}{\kappa_0 n_1 + \overline{m}_{\chi_u}}} \nu_0^{\frac{1}{n_1 \kappa_0 + \overline{m}_{\chi_u}}} < 1$$

where $\widetilde{\mathcal{N}}_F$ is given in (3.1), $\alpha_s = c[(1-\alpha)^{\tau_0}+1]+2s$, $\beta_s = 2(s+c)$ and ζ_s given in (3.2). Moreover, for $\kappa \in (\kappa_0, \kappa_1)$, choose

(9.4)
$$\sigma_{\kappa} \in \left(\left\{\mu^{\zeta_s}\lambda_{-}^{-1}, (C_1K)^{\frac{1}{\kappa n_1 + \overline{m}_{\chi_u}}}\nu_0^{\frac{n}{n_1\kappa + \overline{m}_{\chi_u}}}\right\}^+, 1\right).$$

Then, for each $n \in \mathbb{N}$ and $\bar{\sigma}_{\kappa} \in (\sigma_{\kappa}, 1)$ we have, Ω_{χ_u} being as in Theorem 8.2,

(9.5)
$$\|\mathcal{L}^{n}u\|_{s,*} \leq C_{\sharp}\Omega_{\chi_{u}}(\kappa n_{1} + \overline{m}_{\chi_{u}}, s) \left\{\sigma_{\kappa}^{n}\|u\|_{s,*} + (1 - \sigma_{\kappa})^{-1}C_{\mu,n}\mu^{n}\|u\|_{0}\right\}$$

(9.6)
$$\|\mathcal{L}^n u\|_{s,*} \le C_{\sharp} \Omega_{\chi_u} (\kappa n_1 + \overline{m}_{\chi_u}, s) \overline{\sigma}_{\kappa}^n \|u\|_{s,*}$$

$$+ C_{\bar{\sigma}_{\kappa}} \Omega_{\chi_u} (\kappa n_1 + \overline{m}_{\chi_u}, s)^3 C^3_{\mu, n} \mu^{3n} \|u\|_{L^1}$$

In particular, $\Omega_{\chi_u}(\kappa n_1 + \overline{m}_{\chi_u}, 1)$ is given by (8.9) if F is also a SVPH[#] satisfying (8.8).

Proof. We use Theorem 8.2 with $q_0 = \kappa n_1 \ge n_0$ and $\kappa \in (\kappa_0, \kappa_1)$. First, by conditions (9.1) and (9.2) and Lemma 7.4, we observe that, setting $N = q_0 + M$,

(9.7)

$$\begin{bmatrix} \mathbb{L}_{\overline{m}_{\chi_{u}}} \mathcal{N}(q_{0}) \end{bmatrix}^{\frac{1}{N}} \mu^{2s} \leq \begin{bmatrix} K \mu^{c\overline{m}_{\chi_{u}}^{\tau_{0}}} \mathcal{N}(q_{0}) \end{bmatrix}^{\frac{1}{N}} \mu^{2s}$$

$$\leq \left(K \mu^{c\overline{m}_{\chi_{u}}^{\tau_{0}}} \mathbb{L}_{q_{0} - \lceil \alpha q_{0} \rceil} \widetilde{\mathcal{N}}(\lceil \alpha q_{0} \rceil) \right)^{\frac{1}{N}} \mu^{2s}$$

$$\leq (K^{2} \widetilde{\mathcal{N}}(\lceil \alpha q_{0} \rceil) \mu^{q_{0}^{\tau_{0}} \alpha_{s} + \beta_{s} \overline{m}_{\chi_{u}}^{\tau_{0}}})^{\frac{1}{N}}.$$

Therefore, by equation (8.7),

(9.8)
$$\|\mathcal{L}^N u\|_{\mathcal{H}^s} \le C_1 K \left(\widetilde{\mathcal{N}}(\lceil \alpha q_0 \rceil) \mu^{q_0^{\tau_0} \alpha_s + \beta_s \overline{m}_{\chi u}^{\tau_0}} \right)^{\frac{1}{2}} \|u\|_{\mathcal{H}^s} + \Omega_{\chi_u}(M,s) \|u\|_{s+2}.$$

Moreover, by the sub-multiplicativity of $\hat{\mathcal{N}}$

$$\widetilde{\mathcal{N}}(\lceil \alpha q_0 \rceil) = \widetilde{\mathcal{N}}(\lceil \alpha \kappa n_1 \rceil) \le \widetilde{\mathcal{N}}(\lceil \alpha n_1 \rceil)^{\kappa}$$

⁵⁴Recall ζ_s defined in (2.7), while (9.1), (9.2) and (9.3) constrain κ_0 , κ_1 and n_1 .

It follows by the definition of ν_0 that

$$\sqrt{\widetilde{\mathcal{N}}(\lceil \alpha q_0 \rceil)} \mu^{q_0^{\tau_0} \alpha_s + \beta_s \overline{m}_{\chi_u}^{\tau_0}} \leq \sqrt{\left[\widetilde{\mathcal{N}}(\lceil \alpha n_1 \rceil) \mu^{\frac{\alpha_s q_0^{\tau_0} + \beta_s \overline{m}_{\chi_u}^{\tau_0}}{\kappa}\right]^{\kappa}} \leq \nu_0^{\kappa}.$$

Accordingly

(9.9)
$$\|\mathcal{L}^N u\|_{\mathcal{H}^s} \le \sigma_{\kappa}^N \|u\|_{\mathcal{H}^s} + \Omega_{\chi_u}(M,s) \|u\|_{s+2}.$$

On the other hand, the assumption $\mu^{\zeta_s} \lambda_-^{-1} \leq \nu_0$ implies (6.32), so that we can choose δ_* in (6.33) such that, for all $n \in \mathbb{N}$,

(9.10)
$$\|\mathcal{L}^{n}u\|_{s+2} \leq C_{\sharp}\sigma_{\kappa}^{n}\|u\|_{s+2} + C_{\mu,n}\mu^{n}\|u\|_{0}.$$

Iterating (9.9) by multiple of N and using (9.10) yields (9.5).

Next, we want to compare the norm $\|\cdot\|_0$ with the L^1 -norm. Let us fix $\ell > 0$. Take an admissible central curve γ and notice that, for any $\phi \in \mathcal{C}^0(\mathbb{T})$ with $\|\phi\|_{\infty} = 1$, we have

$$\left|\int_{\mathbb{T}}\phi(t)(u)(\gamma(t)+\ell e_1)dt-\int_{\mathbb{T}}\phi(t)u(\gamma(t))dt\right|=\left|\int_0^\ell ds\int_{\mathbb{T}}\phi(t)\partial_z u(\gamma(t)+se_1)dt\right|.$$

Writing $\gamma(t) = (\sigma(t), t)$ we can make the change of variables $\psi(s, t) = \gamma(t) + se_1 = (\sigma(t) + s, t)$. Since $\det(D\psi) = -1$ and setting $D_{\ell} = \{\psi(s, t) ; t \in \mathbb{T}, s \in [0, \ell]\}$, we have

$$\left| \int_{\mathbb{T}} \phi(t) u(\gamma(t) + \ell e_1) dt - \int_{\mathbb{T}} \phi(t) u(\gamma(t)) dt \right| = \left| \int_{D_\ell} \phi(z) \partial_z u(x, z) dx dz \right| \\ \leq \|\phi\|_{L^{\infty}} \sqrt{\ell} \|u\|_{\mathcal{H}^1}.$$

Hence

$$\left|\int_{\mathbb{T}}\phi(t)u(\gamma(t)+se_1)dt\right| \geq \left|\int_{\mathbb{T}}\phi(t)u(\gamma(t))dt\right| - \sqrt{s}\|u\|_{\mathcal{H}^1}.$$

Integrating in $s \in [0, \ell]$ and taking the sup on γ and ϕ yields

(9.11)
$$\|u\|_{0} \leq \ell^{-1} \|u\|_{L^{1}} + \frac{2\ell^{\frac{1}{2}}}{3} \|u\|_{\mathcal{H}^{1}}$$

Applying (9.11) to (9.5) with the choice $\ell^{-1} = C_{\sigma_{\kappa}}^2 C_{\mu,n}^2 \sigma_{\kappa}^{-2n} \mu^{2n}$, where $C_{\sigma_{\kappa}} = C_{\sharp} (1 - \sigma_{\kappa})^{-1}$, yields

$$\|\mathcal{L}^n u\|_{s,*} \le C_{\sharp} \Omega_{\chi_u}(\kappa n_1 + \overline{m}_{\chi_u}, s) \left\{ \sigma_{\kappa}^n \|u\|_{s,*} + C^3_{\sigma_{\kappa}} C^3_{\mu,n} \mu^{3n} \sigma_{\kappa}^{-2n} \|u\|_0 \right\}.$$

Next, for each $\bar{\sigma}_{\kappa} \in (\sigma_{\kappa}, 1)$, let n_{κ} be the smallest integer such that

$$C_{\sharp}\Omega_{\chi_u}(\kappa n_1 + \overline{m}_{\chi_u}, s)\sigma_{\kappa}^{n_{\kappa}} \le \bar{\sigma}_{\kappa}^{n_{\kappa}}.$$

For each $n \in \mathbb{N}$, write $n = kn_{\kappa} + m$ with $m < \kappa$, then iterating the above equation yields

$$\begin{aligned} \|\mathcal{L}^{n}u\|_{s,*} &\leq \bar{\sigma}_{\kappa}^{kn_{\kappa}} \|\mathcal{L}^{m}u\|_{s,*} + C_{\sharp}\Omega_{\chi_{u}}(\kappa n_{1} + \overline{m}_{\chi_{u}}, s)\mu^{3kn_{\kappa}}C_{\sigma_{\kappa}}^{3}C_{\mu,n_{\kappa}}^{3}\sigma_{\kappa}^{-2n_{\kappa}}\sum_{j=0}^{k-1}\bar{\sigma}_{\kappa}^{n_{\kappa}}\|u\|_{L^{1}} \\ &\leq C_{\sharp}\Omega_{\chi_{u}}(\kappa n_{1} + \overline{m}_{\chi_{u}}, s)\bar{\sigma}_{\kappa}^{n}\|u\|_{s,*} + C_{\sharp}\frac{\Omega_{\chi_{u}}(\kappa n_{1} + \overline{m}_{\chi_{u}}, s)^{3}\mu^{3n}C_{\sigma_{\kappa}}^{3}C_{\mu,n}^{3}}{\bar{\sigma}_{\kappa}^{2n_{\kappa}}(1 - \bar{\sigma}_{\kappa}^{n_{\kappa}})}\|u\|_{L^{1}} \end{aligned}$$

which implies (9.6).

Corollary 9.2. Under the assumptions of Theorem 9.1 there exists a Banach space $\mathcal{B}_{s,*}$ such that $\mathcal{C}^{r-1}(\mathbb{T}^2) \subset \mathcal{B}_{s,*} \subset \mathcal{H}^s(\mathbb{T}^2)$ on which the operator $\mathcal{L}_F : \mathcal{B}_{s,*} \to \mathcal{B}_{s,*}$ has spectral radius one and is quasi compact with essential spectral radius bounded by σ_{κ} .

Proof. Let $\mathcal{B}_{s,*}$ be the completion of $\mathcal{C}^{r-1}(\mathbb{T}^2)$ with respect to $\|\cdot\|_{s,*}$, then $\mathcal{C}^{r-1}(\mathbb{T}^2) \subset \mathcal{B}_{s,*} \subset \mathcal{H}^s(\mathbb{T}^2)$. Iterating (9.6), and since \mathcal{L}_F is a L^1 contraction, implies that the spectral radius is bounded by one, but since the adjoint of \mathcal{L}_F has eigenvalue one, so does \mathcal{L} , hence the spectral radius is one.

To bound the essential spectral radius note that the immersion $\mathcal{B}_{s,*} \hookrightarrow \mathcal{H}^s$ is continuous by definition of the norm. Moreover the immersion $\mathcal{H}^s \hookrightarrow L^1$ is compact for every s by Sobolev embeddings theorems, hence $\mathcal{B}_{s,*} \hookrightarrow L^1$ is compact. Thus, by (9.6) and Hennion theorem [35] follows that the essential spectral radius is bounded by $\bar{\sigma}_{\kappa}$ and hence the claim by the arbitrariness of $\bar{\sigma}_{\kappa}$. \Box

Proof of Theorem 4.4. According to Corollary 9.2, it is enough to check the conditions of Theorem 9.1. Since $\mu > 1$, Corollary 5.20 implies $\sup_{k \le n} \|\mathcal{L}^k 1\|_{\infty} \le K_{\mu}\mu^n$ for each $n \in \mathbb{N}$, for some constant K_{μ} depending on μ . Hence (9.1) is satisfied with c = 1 and $\tau_0 = 1$ and arbitrary $\kappa_1 \in \mathbb{N}$. Next, $\mu^{\zeta_s} \lambda_{-}^{-1} < 1$ is implied by hypothesis (H3). Therefore, condition (3.3) coincides with (9.2) with $\alpha_s, \beta_s, \zeta_s$ given in (3.2). Finally, choosing any κ_0 such that

(9.12)
$$\kappa_0 > \frac{\ln(C_1 K_{\mu}^{\frac{1}{2}})}{\ln \nu_0^{-1}}$$

we have also (9.3), whereby we conclude.

10. The map F_{ε}

Here we apply Theorem 9.1 to the maps F_{ε} , see (2.11), and we prove Theorems 4.5, 4.6 and 4.8.

10.1. The maps F_{ε} are SVPH^{\$}.

We use Lemma 2.6, applied with ω replaced by $\varepsilon\omega$, to check that the maps given in (2.11) are SVPH for ε small enough. Condition (5) is part of the assumptions on F_{ε} and implies $\lambda > 2 ||\partial_{\theta}f||_{\infty}$, which implies (4) and, since $\lambda > 2$, (1) for ε small enough. Finally, conditions (2), (3) and (6) are also immediate for ε small. Hence, there exists $\varepsilon_0 > 0$ such that, for all $\varepsilon < \varepsilon_0$, F_{ε} satisfies the hypothesis of Lemma 2.6 and hence it is a SVPH.

In particular, we choose $\mathbf{C}^{u}_{\varepsilon} = \{(\xi,\eta) \in \mathbb{R}^{2} : |\eta| \leq \varepsilon u_{\star}|\xi|\}, {}^{55}$ and $\mathbf{C}^{c} = \{(\xi,\eta) \in \mathbb{R}^{2} : |\xi| \leq \eta|\}$ with (10.1) $u_{\star} = 2 \|\partial_{x}\omega\|_{\infty} =: \varepsilon^{-1}\chi_{u}.$

Let $(1, \varepsilon u) \in \mathbf{C}^u_{\varepsilon}$, for $p = (x, \theta) \in \mathbb{T}^2$. In this case equation (A.1) yields

(10.2)
$$D_p F_{\varepsilon}(1, \varepsilon u) = (\partial_x f + \varepsilon u \partial_{\theta} f)(1, \varepsilon \Xi_{\varepsilon}(p, u)),$$

where

(10.3)
$$\Xi_{\varepsilon}(p,u) = \frac{\partial_x \omega + \varepsilon u \partial_{\theta} \omega + u}{\partial_x f + \varepsilon u \partial_{\theta} f}$$

We have also a more explicit formula for iteration of the map Ξ_{ε} . For any $k \ge 0$ and $p \in \mathbb{T}^2$, let us denote $p_k = F_{\varepsilon}^k(p)$. Then we have the recursive formula:

(10.4)
$$\Xi_{\varepsilon}^{(n)}(p,u) = \Xi_{\varepsilon}(p_{n-1},\Xi^{(n-1)}(p,u)).$$

On the other hand, recalling (A.2):

(10.5)
$$\partial_u \Xi_{\varepsilon}(p,u) = \frac{\partial_x f + \varepsilon (\partial_\theta \omega \partial_x f - \partial_\theta f \partial_x \omega)}{(\partial_x f + \varepsilon u \partial_\theta f)^2}.$$

In fact, with the above choice of u_{\star} and using (10.3), for ε small enough,

$$D_p F_{\varepsilon}(\mathbf{C}^u_{\varepsilon}) \subset \{(\eta, \xi) : |\xi| \le \frac{3}{2} \lambda^{-1} \varepsilon \chi_u |\eta|\},\$$

hence condition (2.5) holds with $\iota_{\star} = \frac{3}{2}\lambda^{-1} \leq \frac{3}{4} < 1.^{56}$

Moreover, in (A.7) it is shown that for some $\bar{c} > 0$, $\mu_{\pm} = e^{\pm \bar{c}\varepsilon}$. Finally, by (A.9), it follows that condition (2.6) holds with $C_{\star} = 1$, $\lambda_{+} = 2 \sup_{\mathbb{T}^2} \partial_x f$ and $\lambda_{-} = 2\lambda/3$.

⁵⁵Observe that in this special case $\chi_u(\varepsilon) = \varepsilon u_{\star}$, thus we have an unstable cone of size ε .

⁵⁶Recall that $\lambda = \inf \partial_x f > 2$.

The above discussion shows that all the quantities χ_c , ι_{\star} and C_{\star} are independent of ε , and μ_+ and $(\lambda_- - \mu_+)^{-1}$, $(1 - \chi_u)^{-1}$ are bounded uniformly in ε . Therefore, the uniform constant in the sense of the Notation at the end of Section 1 are, in fact, uniform in the usual sense, hence the name. Finally, it is immediate to check that, possibly taking ε_0 smaller, the extra conditions (5.2) and (5.1)

for the invariance of the curves are also satisfied. Indeed, choosing $\hat{n} = c_3^{-1} \log \varepsilon^{-1}$, with $c_3^{-1} < c_2^{-1}$ (which implies (5.2)), and recalling (5.16)-(5.20), we have $C_{\mu,\hat{n}} \lesssim \log \varepsilon^{-1}$ and $\sigma_{\hat{n}} \lesssim \log \varepsilon^{-1} + e^{c_{\sharp}\varepsilon}$, which implies (5.1) if ε is small enough. Thus, F_{ε} is a SVPH[#] for each $\varepsilon \leq \varepsilon_0$.⁵⁷

To conclude the preliminaries, it is useful to note that, if we set $\psi(p) = \langle \nabla \omega, (-\frac{\partial_{\theta} f}{\partial_x f}, 1) \rangle(p)$, for every $p \in \mathbb{T}^2$ and $n \in \mathbb{N}$ we have

$$\det D_p F_{\varepsilon}^n = \prod_{k=0}^{n-1} \det D_{F_{\varepsilon}^k p} F_{\varepsilon} = \prod_{k=0}^{n-1} \left[\partial_x f(F_{\varepsilon}^k p) (1 + \varepsilon \psi(F_{\varepsilon}^k p)) \right],$$

hence

(10.6) $e^{-\bar{c}\varepsilon n}\lambda^n \leq \det D_p F_{\varepsilon}^n \leq e^{\bar{c}\varepsilon n}\Lambda^n, \quad \forall p \in \mathbb{T}^2, \forall n \in \mathbb{N}.$

10.2. A non-transversality argument.

Here the aim is to prove Proposition 10.2 which guarantees that, after some fix time which does not depend on ε , for each point we have at least a couple of pre-images with transversal unstable cones, provided ω is not x-constant. This will imply the existence of the integer n_1 required by Theorem 9.1.

In the following we denote as $\mathfrak{H}_{\varepsilon}$ the set of the inverse branches of F_{ε} .⁵⁸ Moreover, $\mathfrak{H}_{\varepsilon}^{n}$ will be the set of elements of the form $\mathfrak{h}_{1} \circ \cdots \circ \mathfrak{h}_{n}$, for $\mathfrak{h}_{j} \in \mathfrak{H}_{\varepsilon}$ and $\mathfrak{H}_{\varepsilon}^{\infty} := \mathfrak{H}_{\varepsilon}^{\mathbb{N}}$. For $\mathfrak{h} \in \mathfrak{H}_{\varepsilon}^{\infty}$ the symbol \mathfrak{h}_{n} will denote the restriction of \mathfrak{h} on $\mathfrak{H}_{\varepsilon}^{n}$.

Remark 10.1. Since F_0 and F_{ε} are homotopic coverings they are isomorphic, that is there exist $I_{\varepsilon} : \mathbb{T}^2 \to \mathbb{T}^2$ such that $F_{\varepsilon} = F_0 \circ I_{\varepsilon}$. This induces an isomorphism $\mathcal{I}_{\varepsilon} : \mathfrak{H}_0 \to \mathfrak{H}_{\varepsilon}$ defined by $\mathcal{I}_{\varepsilon}\mathfrak{h} = I_{\varepsilon}^{-1} \circ \mathfrak{h}$. Hence, the same is true for the sets $\mathfrak{H}_{\varepsilon}^n = \mathfrak{H}^n$ and $\mathfrak{H}_{\varepsilon}^\infty$. In the following we will then identify inverse branches of F_{ε}^n and F_0^n by these isomorphisms, and drop the script ε from the notation when it is not necessary.

Proposition 10.2. If ω is not x-constant with respect to F_0 (see Definition 2.8), then there exist $\varepsilon_0 > 0$ and $n_0 \in \mathbb{N}$ such that, for every $\varepsilon \leq \varepsilon_0$, $p \in \mathbb{T}^2$ and vector $v \in \mathbb{R}^2$, there exists $q \in F_{\varepsilon}^{-n_0}(p)$ such that $v \notin D_q F_{\varepsilon}^{n_0} \mathbf{C}_{\varepsilon}^u$.

Proof. We argue by contradiction. Suppose that for every $\varepsilon_0 > 0$ and $\ell \in \mathbb{N}$ there exist $\varepsilon_\ell \in [0, \varepsilon_0]$, $p_\ell \in \mathbb{T}^2$ and $v_\ell = (1, \varepsilon_\ell u_\ell)$ with $|u_\ell| \leq u_\star$ such that⁵⁹

(10.7)
$$D_q F^{\ell}_{\varepsilon_{\ell}} \mathbf{C}^u_{\varepsilon_{\ell}} \supset v_{\ell}, \quad \forall q \in F^{-\ell}_{\varepsilon_{\ell}}(p_{\ell}),$$

namely, all the above cones have a common direction. Since the sequence $\{p_{\ell}, u_{\ell}\} \subset \mathbb{T}^2 \times [-u_{\star}, u_{\star}]$, it has an accumulation point (p_*, u_*) . In analogy with (5.13), for $p \in \mathbb{T}^2$ and $u \in [-u_{\star}, u_{\star}]$ we define

(10.8)
$$\Phi_{\varepsilon}^{n}(p,u) = \left(F_{\varepsilon}^{n}(p), \Xi_{\varepsilon}^{(n)}(p,u)\right),$$

where $\Xi_{\varepsilon}^{(n)}$ is given by formula (10.4). Condition (10.7) in terms of this dynamics says that the slope u_{ℓ} is contained in the interval $\Xi_{\varepsilon_{\ell}}^{(\ell)}(q, [-u_{\star}, u_{\star}])$ for every $\ell \in \mathbb{N}$ and $q \in F_{\varepsilon_{\ell}}(p_{\ell})$. Hence, it can be written as:

(10.9)
$$\forall \ell \in \mathbb{N}, \quad \exists (p_\ell, u_\ell) : \quad \pi_2 \circ \Phi_{\varepsilon_\ell}^\ell \left(q, [-u_\star, u_\star] \right) \supset \{u_\ell\}, \quad \forall q \in F_{\varepsilon_\ell}^{-\ell}(p_\ell),$$

 $^{^{57}}$ See Remark 5.12 and Definition 5.1 for details.

⁵⁸Accordingly \mathfrak{H}_0 is the set of inverse branches of F_0 .

⁵⁹We use the notation with subscript ℓ for a generic object that depends on ℓ through ε_{ℓ} , but we keep the notation as simple as possible when there is no need to specify.

where $\pi_2 : \mathbb{T}^2 \times [-u_\star, u_\star] \to [-u_\star, u_\star]$ is the projection on the second coordinate. Now, for $m \in \mathbb{N}$, $\varepsilon \in [0, \varepsilon_0], u_0 \in [-u_\star, u_\star]$ and $\mathfrak{h} \in \mathfrak{H}^\infty$, let us define

(10.10)
$$u_{\mathfrak{h},m}^{\varepsilon}(p) = \pi_2 \circ \Phi_{\varepsilon}^m(\mathfrak{h}_m(p), u_0) : \mathbb{T}^2 \to [-u_\star, u_\star]$$

Next, we prove the following result, which will allow us to conclude the proof.

Lemma 10.3. The sequence of functions defined in (10.10) satisfies:

- (i) For every $\varepsilon \in [0, \varepsilon_0]$ and $\mathfrak{h} \in \mathfrak{H}^{\infty}$, there exists $u_{\mathfrak{h},\infty}^{\varepsilon}(\tilde{p}) := \lim_{m \to \infty} u_{\mathfrak{h},m}^{\varepsilon}(\tilde{p})$, and the limit is uniform in $\tilde{p} \in \mathbb{T}^2$.
- (ii) For every $\mathfrak{h} \in \mathcal{H}^{\infty}$, the sequence $\{u_{\mathfrak{h},\infty}^{\varepsilon}\}_{\varepsilon}$ converges to $\overline{u}_{\mathfrak{h},\infty}$ uniformly for $\varepsilon \to 0$.
- (iii) The functions $\overline{u}_{\mathfrak{h},\infty}$ are independent of \mathfrak{h} , we call them \tilde{u} . In addition, \tilde{u} satisfies

(10.11)
$$\tilde{u}(F_0(q)) = \Xi_0(q, \tilde{u}(q)), \quad \forall q \in \mathbb{T} \times \{\theta_*\}.$$

Proof. Applying Lemma F.1 with $u = u' \equiv u_0 \in [-u_\star, u_\star]$, $\varepsilon_0 = 1$, $A = 2\chi_c u_\star$ and B = 0 we have that there exists $\nu \in (0, 1)$ such that, for each $\mathfrak{h} \in \mathfrak{H}^{\infty}$, $q \in \mathbb{T}^2$, $\varepsilon, \varepsilon' \in [0, 1)$, $m \in \mathbb{N}$ and n > m,⁶⁰

(10.12)
$$\begin{aligned} |u_{\mathfrak{h},m}^{\varepsilon}(q) - u_{\mathfrak{h},m}^{\varepsilon'}(q)| &\leq C_{\sharp}\mu^{3m}|\varepsilon - \varepsilon' \\ |u_{\mathfrak{h},m}^{\varepsilon}(q) - u_{\mathfrak{h},m}^{\varepsilon}(q)| &\leq C_{\sharp}\nu^{m}. \end{aligned}$$

It follows that there exists $u_{\mathfrak{h},\infty}^{\varepsilon}(q) := \lim_{m \to \infty} u_{\mathfrak{h},m}^{\varepsilon}(q)$, and the limit is uniform in q. Next, for each $\delta > 0$, we choose ε_* and m such that $C_{\sharp}\mu^{3m}\varepsilon_* \leq \frac{\delta}{4}$ and $\nu^m \leq \frac{\delta}{4}$, then, for each $\varepsilon, \varepsilon' \leq \varepsilon_*$ and $q \in \mathbb{T}^2$

$$\begin{aligned} |u^{\varepsilon}_{\mathfrak{h},\infty}(q) - u^{\varepsilon'}_{\mathfrak{h},\infty}(q)| &\leq |u^{\varepsilon}_{\mathfrak{h},\infty}(q) - u^{\varepsilon'}_{\mathfrak{h},m}(q)| + |u^{\varepsilon}_{\mathfrak{h},m}(q) - u^{\varepsilon'}_{\mathfrak{h},m}(q)| + |u^{\varepsilon'}_{\mathfrak{h},m}(q) - u^{\varepsilon'}_{\mathfrak{h},\infty}(q)| \\ &\leq 2\nu^m + C_{\sharp}\mu^{3n}|\varepsilon - \varepsilon'| \leq \delta. \end{aligned}$$

The above proves the first two items. Let us proceed with the third one.

First we claim that, for $q \in \mathbb{T}^2$, if \mathfrak{h}_q is such that $q = \mathfrak{h}_q(F_{\varepsilon}(q))$, then

(10.13)
$$u_{\mathfrak{h}\circ\mathfrak{h}_q,\infty}^{\varepsilon}(F_{\varepsilon}(q)) = \Xi_{\varepsilon}(q, u_{\mathfrak{h},\infty}^{\varepsilon}(q)), \quad \forall q \in \mathbb{T}^2.$$

Indeed, since $u_{\mathfrak{h},\infty}^{\varepsilon}$ belongs to the unstable cone, by (10.10), for every $\mathfrak{h} \in \mathfrak{H}^{\infty}$ and $q \in \mathbb{T}^2$,

$$\left(F_{\varepsilon}(q), \Xi_{\varepsilon}(q, u_{\mathfrak{h},\infty}^{\varepsilon}(q))\right) = \Phi_{\varepsilon}(q, u_{\mathfrak{h},\infty}^{\varepsilon}(q)) = \left(F_{\varepsilon}(q), u_{\mathfrak{h}\circ\mathfrak{h}_{q},\infty}^{\varepsilon}(F_{\varepsilon}(q))\right)$$

which implies the claim taking the projection on the second coordinate.

For every $\ell \in \mathbb{N}$, let us now consider ε_{ℓ} , p_{ℓ} and u_{ℓ} as given in (10.9) and let ℓ_j so that (p_{ℓ_j}, u_{ℓ_j}) is a convergent sequence. Equation (10.7) implies

(10.14)
$$|u_{\ell_j} - u_{\mathfrak{h}, n_{\ell_j}}^{\varepsilon_{\ell_j}}(p_{\ell_j})| \le C_{\sharp} \nu^{n_{\ell_j}}.$$

Taking the limit for $j \to \infty$ in the above inequality yields⁶¹

(10.15)
$$u_* = \lim_{j \to \infty} u_{\ell_j} = \lim_{j \to \infty} u_{\mathfrak{h}, n_{\ell_j}}^{\varepsilon_{\ell_j}}(p_{\ell_j}) = \overline{u}_{\mathfrak{h}, \infty}(p_*),$$

regardless of the choice of the inverse branch $\mathfrak{h} \in \mathfrak{H}^{\infty}$. Let \mathfrak{h}_q be the inverse branch such that $q = \mathfrak{h}_q(F_{\varepsilon}(q))$, and set $q_{\ell} = \mathfrak{h}_q(p_{\ell})$ in equation (10.13) to obtain:

(10.16)
$$u_{\mathfrak{h}\circ\mathfrak{h}_{q},\infty}^{\varepsilon_{\ell}}(p_{\ell}) = \Xi_{\varepsilon_{\ell}}(q_{\ell}, u_{\mathfrak{h},\infty}^{\varepsilon_{\ell}}(q_{\ell})).$$

By item (ii) above, and by the continuity of the map F_{ε} , we can take the limit as $\ell_j \to \infty$ in the last equation and obtain

$$\overline{u}_{\mathfrak{h}\circ\mathfrak{h}_{q},\infty}(p_{*})=\Xi_{0}(q_{*},\overline{u}_{h,\infty}(q_{*})),$$

where q_* is such that $F_0(q_*) = p_*$. By (10.15), the above equation becomes $u_* = \Xi_0(q_*, \overline{u}_{\mathfrak{h},\infty}(q_*))$, and, since $\Xi_0(q_*, \cdot)$ is invertible, this implies that there exists $u_*(q_*)$ independent of $\mathfrak{h} \in \mathcal{H}^\infty$ such that

$$u_*(q_*) = \bar{u}_{\mathfrak{h}}(q_*) = \lim_{j \to \infty} u_{\mathfrak{h},\infty}^{\varepsilon_{\ell_j}}(q_{\ell_j})$$

⁶⁰The second equation of (10.12) is a direct consequence of (F.5) which implies that $\Xi_{\varepsilon}(p, \cdot)$ is a contraction.

⁶¹Recall that (p_*, u_*) is an accumulation point of the sequence (p_ℓ, u_ℓ) given in (10.9)

Hence, by induction, $\overline{u}_{\mathfrak{h},\infty}(q)$ is independent on \mathfrak{h} for each $q \in \bigcup_{k \in \mathbb{N}} F_0^{-k}(p_*) =: \Lambda_{\theta_*}$, let us call it $u_*(q)$. Taking the limit in equation (10.13) we have, for each $q \in \Lambda_{\theta_*}$,

(10.17)
$$u_*(F_0(q)) = \Xi_0(q, u_*(q)).$$

Note that the $\overline{u}_{\mathfrak{h},\infty}$ are uniform limits of continuous functions and hence are continuous functions such that $\overline{u}_{\mathfrak{h},\infty}|_{\Lambda_{\theta_*}} = u_*$. Since Λ_{θ_*} is dense in $\mathbb{T} \times \{\theta_*\}$.⁶² It follows that the $\overline{u}_{\mathfrak{h},\infty}$ equal some continuous function \tilde{u} defined on $\mathbb{T} \times \{\theta_*\}$ and independently of \mathfrak{h} . In addition, \tilde{u} satisfies (10.11).⁶³

We can now conclude the proof of Proposition 10.2. By Lemma 10.3 we can find a function $\tilde{u}: \mathbb{T}^2 \to \mathbb{R}$ and $\theta_* \in \mathbb{T}^1$ such that (10.11) holds, namely:

(10.18)
$$\tilde{u}(F_0(q)) = \frac{\partial_x \omega(q) + \tilde{u}(q)}{\partial_x f(q)}, \quad q \in \mathbb{T}^1 \times \{\theta_*\}$$

Let us use the notation $g_{\theta}(x)$ for a function $g(x, \theta)$ and observe that, integrating (10.18) and recalling that ω is periodic by hypothesis, we have

$$\int_{0}^{1} \tilde{u}_{\theta_{*}}(x) dx = \int_{0}^{1} f_{\theta_{*}}'(x) \tilde{u}_{\theta_{*}}(f_{\theta_{*}}(x)) dx - \int_{0}^{1} \partial_{x} \omega(x,\theta_{*}) dx = \sum_{i=0}^{d-1} \int_{U_{i}} f_{\theta_{*}}'(x) \tilde{u}_{\theta_{*}}(f_{\theta_{*}}(x)) dx$$
$$= d \int_{0}^{1} \tilde{u}_{\theta_{*}}(t) dt,$$

where U_i are the invertibility domains of f_{θ_*} , and d > 1 its topological degree. Hence $\int_{\mathbb{T}} \tilde{u}_{\theta_*}(x) dx = 0$. So there is a potential given by $\Psi_{\theta_*}(x) = \int_0^x \tilde{u}_{\theta_*}(z) dz$. Finally, integrating equation (10.18) from 0 to x, there exists c > 0 such that

$$\omega_{\theta_*}(x) = \Psi_{\theta_*}(f_{\theta_*}(x)) - \Psi_{\theta_*}(x) + c,$$

which contradicts the assumption on ω whereby proving the Proposition.

For reasons which will be clear shortly, we introduce a new quantity related to $\mathcal{N}_{F_{\varepsilon}}$ and $\mathcal{N}_{F_{\varepsilon}}$ which can be interpreted as a kind of normalization of the latter one. The definition is inspired by [14].

Definition 10.4. For each $p = (x, \theta) \in \mathbb{T}^2$, $v \in \mathbb{R}^2$, $n \in \mathbb{N}$ and $\varepsilon > 0$ we define

(10.19)
$$\widetilde{\mathfrak{N}}(x,\theta,v,n) := \frac{1}{h_*(x,\theta)} \sum_{\substack{(y,\eta) \in F_{\varepsilon}^{-n}(x,\theta) \\ DF_{\varepsilon}^n(y,\eta) \mathbf{C}_{\varepsilon}^u \supset v}} \frac{h_*(y,\theta)}{|\det DF_{\varepsilon}^n(y,\eta)|},$$

where, for every $\theta \in \mathbb{T}$, $h_*(\cdot, \theta) =: h_{*\theta}(\cdot)$ is the density of the unique invariant measure of $f(\cdot, \theta)$. As before we will denote $\widetilde{\mathfrak{N}}(n) := \sup_p \sup_v \widetilde{\mathfrak{N}}(p, v, n)$.

The motivation to introduce this quantity is twofold. One reason lies in Lemma 10.5 below in which, using a shadowing argument similar to [20, Appendix B], we exploit the following fact: for each $\theta \in \mathbb{T}$, setting $f_{\theta}(\cdot) = f(\cdot, \theta)$, we have

(10.20)
$$\frac{1}{h_{*\theta}(x)} \sum_{y \in f_{\theta}(x)} \frac{h_{*\theta}(y)}{(f_{\theta}^n)'(y)} = 1, \quad \forall x \in \mathbb{T}.$$

 $^{^{62}}$ It follows from the expansivity of $f(\cdot, \theta_*)$ that the preimmages of any point form a dense set.

⁶³Just approximate any point with a sequence $\{q_j\} \subset \Lambda_{\theta_*}$ and take the limit in (10.17).

On the other hand it is easy to see that \mathfrak{N} has the same properties of $\widetilde{\mathcal{N}}_{F_{\varepsilon}}$. In particular, arguing exactly in the same way as in Proposition 7.2 and Lemma 7.4, one can show that

(10.21)
$$\mathfrak{N}(n)$$
 is submultiplicative,

(10.22)
$$\mathcal{N}_{F_{\varepsilon}}(n)^{\frac{1}{n}} \leq C_{\sharp} \|\mathcal{L}_{F_{\varepsilon}}^{n\lfloor 1-\alpha \rfloor} 1\|_{\infty}^{\frac{1}{n}} \left(\widetilde{\mathfrak{N}}(\lfloor \alpha n \rfloor)^{\frac{1}{\lfloor \alpha n \rfloor}}\right)^{\alpha}, \quad \text{for some } \alpha \in (0,1)$$

(10.23)
$$\widetilde{\mathfrak{N}}(n) \le \sup_{(x,\theta)\in\mathbb{T}^2} \frac{1}{h_*(x,\theta)} (\mathcal{L}_{F_{\varepsilon}}^n h_*)(x,\theta)$$

This implies that we can check condition (9.2) of Theorem (9.1) with $\widetilde{\mathcal{N}}$ replaced by $\widetilde{\mathfrak{N}}$. To ease notation in the following we set $\mathcal{L}_{F_{\varepsilon}} =: \mathcal{L}_{\varepsilon}$.

Lemma 10.5. There are constants $C, c_* > 0$ such that, for each $n < C\varepsilon^{-\frac{1}{2}}$,

(10.24)
$$\sup_{(x,\theta)\in\mathbb{T}^2}\frac{1}{h_*(x,\theta)}(\mathcal{L}^n_{\varepsilon}h_*)(x,\theta) \le e^{c_*n^2\varepsilon}.$$

Proof. Let $F_{\varepsilon}^{n}(q) = (x, \theta)$ and define $q_{k} = (x_{k}, \theta_{k}) = F_{\varepsilon}^{k}(q)$, for every $0 \leq k \leq n$. Then,

(10.25)
$$|\theta - \theta_k| \le \sum_{j=k}^{n-1} \varepsilon ||\omega||_{\infty} \le C_{\sharp} (n-k) \varepsilon.$$

Let us set $f_{\theta}(y) = f(y, \theta)$. Since f_{θ} is homotopic to f_{θ_k} , for each k, there is a correspondence between inverse branches, hence there exists x_* such that $|f_{\theta}^n(x_*) - x_n| = 0$. Moreover, let $\xi_k = f_{\theta}^k(x_*) - x_k$. Since f is expanding, by the mean value theorem and (10.25), there is $(\bar{x}, \bar{\theta})$ such that

$$|\xi_{k+1}| = |\langle \nabla f(\bar{x}, \bar{\theta}), (\xi_k, \theta_k - \theta) \rangle| \ge \lambda |\xi_k| - C_{\sharp} n\varepsilon.$$

Since $\xi_n = 0$, we find by induction $|\xi_k| \leq \sum_{j=k}^{n-1} \lambda^{-j+k} C_{\sharp} \varepsilon n \leq C_{\sharp} \varepsilon n$. Moreover, since h_* is differentiable⁶⁴ we also have

$$h_*(x_k,\theta_k) - h_{*\theta}(f^k_\theta(x_*))| \le C_\sharp \varepsilon n$$

Next, since $|\det D_q F_{\varepsilon} - \partial_x f(q)| \le C_{\sharp} \varepsilon$,

$$\frac{(f_{\theta}^{n})'(x_{*})}{\det DF_{\varepsilon}^{n}(x_{0},\theta_{0})} = \prod_{k=0}^{n-1} \frac{f_{\theta}'(f_{\theta}^{k}(x_{*}))}{\det DF_{\varepsilon}(x_{k},\theta_{k})} \leq \prod_{k=0}^{n-1} \frac{f_{\theta}'(f_{\theta}^{k}(x_{*}))}{\det DF_{\varepsilon}(f_{\theta}^{k}(x_{*}),\theta)} \left[1 + C_{\sharp}n\varepsilon\right] \leq e^{c_{\sharp}n^{2}\varepsilon}.$$

It follows that,

$$\frac{1}{h_*(x,\theta)} \sum_{(y,\vartheta)\in F_{\varepsilon}^{-n}(x,\theta)} \left(\frac{h_*(y,\vartheta)}{|\det DF_{\varepsilon}^n(y,\vartheta)|} \right) \leq \frac{e^{c_{\sharp}n^2\varepsilon}}{h_{*\theta}(x)} \sum_{x_*\in f_{\theta}^{-n}(x)} \frac{h_{*\theta}(x_*)}{(f_{\theta}^n)'(x_*)} = e^{c_{\sharp}n^2\varepsilon},$$

where we have used (10.20).

10.3. Proof of Theorem 4.5.

By section 10.1 F_{ε} is SVPH[#] for $\varepsilon \leq \varepsilon_0$. We check the other hypotheses of Theorem 9.1 for F_{ε} , under the assumption that ω is not *x*-constant. In this case the existence of n_0 independent of ε is guaranteed by Proposition 10.2. Notice that $\chi_u = u_{\star}\varepsilon$, i.e the unstable cone $\mathbf{C}_{\varepsilon}^u$ is of order ε while the center cone \mathbf{C}^c is of order one. Hence, by (5.11), there exists $c_0 > 0$ such that $\overline{m}_{\chi_u} \leq \lfloor c_0 \log \varepsilon^{-1} \rfloor$. ⁶⁵ Provided $n_1\kappa_1 = c_1 \log \varepsilon^{-1}$, for some $c_1 > 0$ and, by Lemma 10.5, we have

$$\sup_{m \le n} \|\mathcal{L}_{\varepsilon}^m 1\|_{\infty} \le \frac{|\sup h_*|}{|\inf h_*|} \sup_{m \le n} \left\| \frac{1}{h_*} \mathcal{L}_{\varepsilon}^m h_* \right\|_{\infty} \le \frac{|\sup h_*|}{|\inf h_*|} e^{c_* \varepsilon n^2}, \quad \forall n \le \{c_0 + c_1\} \log \varepsilon^{-1},$$

 $^{^{64}\}mathrm{See}$ [19] for the details.

 $^{^{65}}$ For simplicity in the following we drop the $\lfloor \cdot \rfloor$ notation.

hence condition (9.1) with $K = \frac{|\sup h_*|}{|\inf h_*|}$, $\tau_0 = 2$ and $c = c_* \varepsilon$. Next, we prove that there exists a uniform constant ν_0 such that

(10.26)
$$\left\{ e^{\bar{c}\varepsilon\zeta_s}\lambda^{-\frac{1}{2}}, \widetilde{\mathfrak{N}}(\lceil\alpha n_1\rceil)e^{c_{\sharp}\varepsilon[n_1^2\kappa_1+m_{\chi_u}^2]} \right\}^+ \le \nu_0 < 1,$$

i.e condition (9.1) with $\widetilde{\mathcal{N}}_F$ replaced by $\widetilde{\mathfrak{N}}$ which, as we already observed, is an alternative condition under which Theorem 9.1 holds for F_{ε} . Let $\varepsilon_1 \in (0, \varepsilon_0)$, ε_0 given in Proposition 10.2, be such that, for each $1 \leq s \leq r-1$

(10.27)
$$\mu^{\zeta_s} \lambda^{-1} = e^{\bar{c}\varepsilon\zeta_s} \lambda^{-1} < 1, \quad \forall \varepsilon \in (0, \varepsilon_1)$$

Accordingly, for every $p = (x, \theta) \in \mathbb{T}^2$ and $v \in \mathbb{R}^2$, there exists $q_* \in F^{-n_0}(p)$, n_0 provided by Proposition 10.2, such that

$$\frac{1}{h_*(x,\theta)} \sum_{\substack{(y,\theta)\in F_{\varepsilon}^{-n_0}(p)\\DF_{\varepsilon}^{n_0}(y,\eta)\mathbf{C}_{\varepsilon}^{u}\supset v}} \frac{h_*(y,\theta)}{|\det D_q F_{\varepsilon}^{n_0}|} \leq \frac{1}{h_*(x,\theta)} (\mathcal{L}_{\varepsilon}^{n_0}h_*)(x,\theta) - \frac{\mathbb{k}}{|\det D_{q_*} F_{\varepsilon}^{n_0}|},$$

where $\mathbf{k} = \frac{\inf h_*}{\sup h_*}$. By Lemma 10.5 and equation (10.6), the last expression is bounded by $e^{c_* n_0^2 \varepsilon} - \frac{\mathbf{k}}{\Lambda^{n_0} e^{\overline{\varepsilon}\varepsilon}}$. Choosing $\varepsilon_2 < \min\left(\varepsilon_1, \frac{1}{cn_0^2}\log(1 + \frac{\mathbf{k}}{\Lambda^{n_0} e^{\overline{\varepsilon}\varepsilon}}\right)$, we have that $\widetilde{\mathfrak{N}}(n_0) \leq \overline{\sigma} < 1$ for every $\varepsilon \in [0, \varepsilon_2]$. Consequently, choosing $n_1 = \alpha^{-1}n_0$, for each $\nu_0 > \overline{\sigma}$, there exists $\varepsilon_* \in (0, \varepsilon_2)$ such that

(10.28)
$$\widetilde{\mathfrak{N}}(\lceil \alpha n_1 \rceil) e^{c_{\sharp} \varepsilon [n_1^2 \kappa_1 + (\ln \varepsilon^{-1})^2]} \le \bar{\sigma} e^{c_{\sharp} \varepsilon (\ln \varepsilon^{-1})^2} \le \nu_0 < 1, \quad \forall \varepsilon \in (0, \varepsilon_*).$$

By (10.27) and (10.28) we obtain (10.26). Finally, condition (9.3) is satisfied choosing

(10.29)
$$\kappa_0 \ge \frac{\ln C_1 K}{\ln \nu_0^{-1}}$$

Thus Theorem 9.1 applies and Theorem 4.5 follows by Corollary 9.2 choosing $\kappa = \kappa_1$.

10.4. Eigenfunctions regularity (quantitative).

As we have already seen in Corollary 9.2, the main consequence of Theorem 9.1 is that there exists a Banach space $\mathcal{B}_{s,*} \subset \mathcal{H}^s$ on which the transfer operator $\mathcal{L}_{\varepsilon}$ is quasi compact for each $\varepsilon < \varepsilon_*$ with a uniform essential spectral radius.

Using inequality (9.5), we can say much more about the constants. Indeed, by the previous section Theorem 9.1 applies for κ_0 as in (10.29) and $\kappa_1 = n_0^{-1} \alpha c_1 \log \varepsilon^{-1}$. Hence, for each $n, \kappa \in \mathbb{N}, \kappa \in (\kappa_0, \kappa_1)$,

$$\left\|\mathcal{L}^{n}_{\varepsilon}u\right\|_{s,*} \leq \Omega_{\chi_{u}}(\kappa n_{1} + \overline{m}_{\chi_{u}}, s) \left\{\sigma^{n}_{\kappa}\|u\|_{s,*} + (1 - \sigma_{\kappa})^{-1}C_{\mu,n}\mu^{n}\|u\|_{0}\right\},$$

where $m_{\chi_u} = c_0 \log \varepsilon^{-1}$ and σ_{κ} is given in (9.4). The choice $\kappa = C_{\sharp} \log \varepsilon^{-1}$, in the proof of Theorem 4.5, yielded a spectral radius uniform in ε , but it provides no control on the constant $\Omega_{\chi_u}(\kappa n_1 + \overline{m}_{\chi_u}, s)$. On the contrary, the choice $\kappa = 2\kappa_0 \in \mathbb{N}$ (independent of ε) implies, for some $c_{\star} > 0$,

$$\sigma_{\kappa_0} \in (e^{-c_\star \left(\log \varepsilon^{-1}\right)^{-1}}, 1),$$

hence a weaker contraction, but it allows a better control of the constants. Indeed, observe that by (5.16)

$$C_{\mu,n_1+m_{\chi_u}} \le C_{\sharp} \min\{\log \varepsilon^{-1}, \varepsilon^{-1}\} = C_{\sharp} \log \varepsilon^{-1}$$

Hence, it follows by (8.9) that we can find $\beta_3, \beta_3, C_{\sharp} > 0$ and $\epsilon > 0$ such that

(10.30)
$$\Omega_{\chi_u}(2\kappa_0 n_1 + \overline{m}_{\chi_u}, 1) \le C_{\sharp} \varepsilon^{-\frac{13}{2}} (\log \varepsilon^{-1})^{\beta_1} e^{\beta_2 \varepsilon \log \varepsilon^{-1}}$$

Thus, for each $\beta > \frac{13}{2}$, we have, for all $n \in \mathbb{N}$,

(10.31)
$$\begin{aligned} \|\mathcal{L}_{\varepsilon}u\|_{0} &\leq Ce^{\bar{c}n\varepsilon}\|u\|_{0}\\ \|\mathcal{L}_{\varepsilon}^{n}u\|_{1,*} &\leq C_{\alpha}\varepsilon^{-\beta}e^{-\frac{c_{*}n}{\ln\varepsilon^{-1}}}\|u\|_{1,*} + C_{\beta}\varepsilon^{-\beta}\|u\|_{0}. \end{aligned}$$

Proof of Theorem 4.8. Suppose $\mathcal{L}_{\varepsilon} u = \nu u$ with $\nu^n > e^{-r \frac{c \star n}{\ln \varepsilon^{-1}}}$, r < 1, then

$$\|u\|_{1,*} = \nu^{-n} \|\mathcal{L}_{\varepsilon}^{n} u\|_{1,*} \le C_{\beta} \varepsilon^{-\beta} \nu^{-n} e^{-\frac{c*n}{\ln \varepsilon^{-1}}} \|u\|_{1,*} + C_{\beta} \nu^{-n} \varepsilon^{-\beta} \|u\|_{0}$$

We choose n to be the smallest integer such that $C_{\beta}\varepsilon^{-\beta}e^{-\frac{(1-r)c_*n}{\ln \varepsilon^{-1}}} \leq \frac{1}{2}$, which yields

$$\|u\|_{\mathcal{H}^1} \le \|u\|_{1,*} \le C_\beta \varepsilon^{-(1-r)^{-1}\beta} \|u\|_0$$

which concludes the proof.

10.5. Proof of Theorem 4.6.

Let $\sigma_{ph}(\mathcal{L}_{F_{\varepsilon}}) = \{z \in \mathbb{C} : |z| = 1\}$ be the peripheral spectrum. By Theorem 4.5 the cardinality of $\sigma_{ph}(\mathcal{L}_{F_{\varepsilon}})$ is finite and if $e^{i\vartheta} \in \sigma_{ph}(\mathcal{L}_{F_{\varepsilon}})$, then the corresponding eigenspace is finite dimensional. In addition, since the operator is power bounded, there cannot exists Jordan blocks, thus the algebraic and geometric multiplicity are equal.

Hence, there exist $N \in \mathbb{N}$ and $\{\vartheta_j, h_j, \ell_j\}_{j=1}^N$ such that $\vartheta_0 = 1$, $\ell_0(\varphi) = \int_{\mathbb{T}^2} \varphi$, $\vartheta_j \in [0, 2\pi)$, $h_j \in \mathcal{B}_{*,s}$, $\ell_j \in \mathcal{B}'_{*,s}$ and $\mathcal{L}_{F_{\varepsilon}}h_j = e^{i\vartheta_j}h_j$, $\ell_j(\mathcal{L}_{F_{\varepsilon}}\varphi) = e^{i\vartheta_j}\ell_j(\varphi)$ for all $\varphi \in \mathcal{B}_{*,s}$. It follows that we have the following spectral decomposition

(10.32)
$$\mathcal{L}_{F_{\varepsilon}} = \sum_{j} e^{i\vartheta_{j}} \Pi_{j} + Q$$

where $\Pi_j h = h_j \ell_j(h)$, $\Pi_j \Pi_k = \delta_{jk} \Pi_j$ and Q has spectral radius strictly smaller than one. Moreover, see [10, Section 5] for a proof which applies verbatim to the present context, the eigenvectors associated to the eigenvalue one are the physical measures and since they are absolutely continuous with respect to Lebesgue they are SRB as well. Also note that

(10.33)
$$\lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} e^{-i\vartheta_j k} \mathcal{L}_{F_{\varepsilon}}^k = \Pi_j,$$

where the limit is meant in the $\mathcal{B}_{s,*}$ topology. By Lemma 9.6 it follows the Π_j are bounded operators from $L^1 \to \mathcal{B}_{*,s}$.

Also, choosing $\varphi \in \mathcal{C}^{\infty}$ is such that $\alpha := \int_{\mathbb{T}^2} \varphi h_j > 0$,

$$\begin{aligned} |\ell_j(h)| &= \alpha^{-1} \left| \int_{\mathbb{T}^2} \varphi \Pi_j h \right| = \alpha^{-1} \left| \lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} e^{-i\vartheta_j k} \int_{\mathbb{T}^2} \varphi \mathcal{L}_{F_\varepsilon}^k h \right| \\ &\leq \alpha^{-1} \lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} \int_{\mathbb{T}^2} |\varphi \circ F_\varepsilon^k| \, |h| \leq \alpha^{-1} \|\varphi_j\|_\infty \|h\|_{L^1}. \end{aligned}$$

Which implies that there exists $\tilde{\ell}_j \in L^{\infty}$ such that

$$\ell_j(h) = \int_{\mathbb{T}^2} \tilde{\ell}_j h$$

Note that the above also implies $\tilde{\ell}_j \circ F_{\varepsilon} = e^{i\vartheta_j}\tilde{\ell}_j$. The above means that, for all $l \in \mathbb{N}$,

$$\int_{\mathbb{T}^2} \tilde{\ell}_j^l \mathcal{L}_{F_{\varepsilon}} h = \int_{\mathbb{T}^2} \tilde{\ell}_j^l \circ F_{\varepsilon} h = e^{i\vartheta_j l} \int_{\mathbb{T}^2} \tilde{\ell}_j^l h.$$

This implies that $e^{i\vartheta_j l}$ belongs to the spectrum of $(\mathcal{L}_{F_{\varepsilon}})'$, hence of $\mathcal{L}_{F_{\varepsilon}}$. Since there can be only finitely many elements of $\sigma_{ph}(\mathcal{L}_{F_{\varepsilon}})$, it must be $\vartheta_j = \frac{2\pi p}{q}$ for some $p, q \in \mathbb{N}$, that is the $\{\vartheta_j\}$ form a collection of finite groups.

Next we would like to better understand the structure of the peripheral spectrum, and prove equations (4.4) and (4.5).

Let $(x_k, \theta_k) = F_{\varepsilon}^k(x, \theta)$ and $f_{\theta}(x') = f(x', \theta)$. By [20, Lemma 4.2], for each $\theta \in \mathbb{T}$, there exists $Y_{\theta,n}$ such that $\pi_2(F_{\varepsilon}^n(x, \theta)) = f_{\theta}^n(Y_{\theta,n}(x))$ and, for all $k \leq n$,

$$\begin{aligned} \|x_k - f_{\theta}^k \circ Y_{\theta,n}\|_{\infty} &\leq C_{\sharp} \varepsilon k \\ |\theta_k - \theta| &\leq C_{\sharp} k \varepsilon \\ \|1 - \partial_x Y_{\theta,n}\|_{\infty} &\leq C_{\sharp} \varepsilon n^2. \end{aligned}$$

For each $n \in \{C_{\sharp} \ln \varepsilon^{-1}, \dots, C_{\sharp} \varepsilon^{-\frac{1}{2}}\}$ we have

$$\begin{split} \int_{\mathbb{T}^2} \varphi \mathcal{L}_{F_{\varepsilon}}^n h &= \int_{\mathbb{T}^2} \varphi \circ F_{\varepsilon}^n h = \int_{\mathbb{T}^2} \varphi(f_{\theta}^n(Y_{\theta,n}(x)), \theta) h(x, \theta) + \mathcal{O}(\varepsilon n \| \partial_{\theta} \varphi \|_{\mathcal{C}^0} \| h \|_{L^1}) \\ &= \int_{\mathbb{T}^2} \varphi(f_{\theta}^n(x, \theta) h(Y_{\theta,n}^{-1}(x), \theta) + \mathcal{O}(\varepsilon n^2 \| \partial_{\theta} \varphi \|_{\mathcal{C}^0} \| h \|_{L^1}) \\ &= \int_{\mathbb{T}^2} \varphi(x, \theta) [\mathcal{L}_{\theta}^n(h \circ Y_{\theta,n}^{-1})](x, \theta) + \mathcal{O}(\varepsilon n^2 \| \partial_{\theta} \varphi \|_{\mathcal{C}^0} \| h \|_{L^1}). \end{split}$$

Let \mathcal{L}_{θ} be the transfer operator associated to f_{θ} and let $h_*(\cdot, \theta)$ be the associated unique invariant density. Then, for each $\theta \in \mathbb{T}$, \mathcal{L}_{θ} has a uniform spectral gap $\sigma \in (0, 1)$ on the Sobolev space $\mathcal{H}^1(\mathbb{T})$. Thus

$$\int_{\mathbb{T}^2} \left| \left[\mathcal{L}^n_{\theta}(h \circ Y_{\theta,n}^{-1}) \right](x,\theta) - h_*(x,\theta) \int_{\mathbb{T}} (h \circ Y_{\theta,n}^{-1})(y,\theta) dy \right| dx d\theta \le C_{\sharp} \sigma^n \|h\|_{\mathcal{H}^1},$$

since $\|h \circ Y_n^{-1}(\cdot, \theta)\|_{\mathcal{H}^1} \le C_{\sharp} \|h(\cdot, \theta)\|_{\mathcal{H}^1}$. Hence,

(10.34)
$$\int_{\mathbb{T}^2} \varphi \mathcal{L}_{F_{\varepsilon}}^n h = \int_{\mathbb{T}^2} dx \varphi(x,\theta) h_*(x,\theta) \int_{\mathbb{T}} dy h(y,\theta) + \mathcal{O}(\varepsilon n^2 \|\partial_{\theta}\varphi\|_{\mathcal{C}^0} \|h\|_{L^1}) \\ + \mathcal{O}(\sigma^n \|\varphi\|_{\mathcal{C}^0} \|h\|_{\mathcal{H}^1}).$$

We can then choose $n = c \ln \varepsilon^{-1}$, for c large enough, and obtain

(10.35)
$$\|\mathcal{L}_{F_{\varepsilon}}^{c_{\sharp}\ln\varepsilon^{-1}}h - Ph\|_{(\mathcal{C}^{1})'} \leq C_{\sharp}\varepsilon[\ln\varepsilon^{-1}]^{2}\|h\|_{L^{1}} + C_{\sharp}\varepsilon^{3\alpha}\|h\|_{\mathcal{H}^{1}}.$$

Equation (9.6) yields

(10.36)
$$\|\mathcal{L}_{F_{\varepsilon}}^{k}h\|_{\mathcal{H}^{1}} \leq C_{\alpha}\varepsilon^{-\alpha}e^{-\frac{c_{\sharp}k}{\ln\varepsilon^{-1}}}\|h\|_{\mathcal{B}_{1,\ast}} + C_{\alpha}\varepsilon^{-3\alpha}\ln\varepsilon^{-1}\|h\|_{L^{1}}.$$

Hence, by equation (10.35) we have that, for each $\varphi \in \mathcal{C}^1$ and $h \in \mathcal{B}_{1,*}$,

(10.37)
$$\int_{\mathbb{T}^{2}} \varphi \Pi_{0} h = \lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-1} \int_{\mathbb{T}^{2}} \varphi \mathcal{L}_{F_{\varepsilon}}^{k} h$$
$$= \lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-c_{\sharp} \ln \varepsilon^{-1}} \int_{\mathbb{T}^{2}} \varphi \mathcal{L}_{F_{\varepsilon}}^{c_{\sharp} \ln \varepsilon^{-1}} \mathcal{L}_{F_{\varepsilon}}^{k} h$$
$$= \lim_{n \to \infty} \frac{1}{n} \sum_{k=0}^{n-c_{\sharp} \ln \varepsilon^{-1}} \int_{\mathbb{T}^{2}} \varphi P \mathcal{L}_{F_{\varepsilon}}^{k} h + \mathcal{O}(\varepsilon [\ln \varepsilon^{-1}]^{2} \|\varphi\|_{\mathcal{C}^{1}} \|h\|_{L^{1}})$$
$$= \int_{\mathbb{T}^{2}} \varphi P \Pi_{0} h + \mathcal{O}(\varepsilon [\ln \varepsilon^{-1}]^{2} \|\varphi\|_{\mathcal{C}^{1}}).$$

Hence, by the density of $\mathcal{B}_{1,*}$ in L^1 and since Π_0 extends naturally to a bounded operator on L^1 , we have

(10.38)
$$\|\Pi_0 - P\Pi_0\|_{L^1 \to (\mathcal{C}^1)'} \le C_{\sharp} \varepsilon [\ln \varepsilon^{-1}]^2.$$

The same argument proves a similar result for the projectors Π_j yielding (4.4).

It remains to prove equation (4.5). For each $\tau > 0$ consider $h \in \mathcal{B}_{*,1}$ such that $\mathcal{L}_{F_{\varepsilon}}h = \nu h$ with $|\nu| \ge e^{-\varepsilon^{\tau}}$. Then, for all $\varphi \in \mathcal{C}^1$ and $n \in \mathbb{N}$, we have

$$\begin{split} \int_{\mathbb{T}^2} \varphi h &= \nu^{-n} \int_{\mathbb{T}^2} \varphi \mathcal{L}_{F_{\varepsilon}}^n h = \nu^{-n} \int_{\mathbb{T}^2} \varphi \circ F_{\varepsilon}^n h \\ &= \nu^{-n} \int_{\mathbb{T}^2} \varphi (f_{\theta}^n \circ Y_{\theta,n}(x), \theta) h(x, \theta) + \mathcal{O} \left(\nu^{-n} n \varepsilon \|\varphi\|_{\mathcal{C}^1} \|h\|_{L^1} \right) \\ &= \nu^{-n} \int_{\mathbb{T}^2} \varphi (f_{\theta}^n(x), \theta) h(Y_{\theta,n}^{-1}(x), \theta) + \mathcal{O} \left(\nu^{-n} n^2 \varepsilon \|\varphi\|_{\mathcal{C}^1} \|h\|_{L^1} \right) \\ &= \nu^{-n} \int_{\mathbb{T}^2} \varphi (x, \theta) (\mathcal{L}_{\theta}^n [h_{\theta} \circ Y_{\theta,n}^{-1}])(x) + \mathcal{O} \left(\nu^{-n} n^2 \varepsilon \|\varphi\|_{\mathcal{C}^1} \|h\|_{L^1} \right) \end{split}$$

where $h_{\theta}(x) = h(x, \theta)$. Note that, by inequality (10.36), we have $\|h\|_{\mathcal{H}^1} \leq C_{\sharp} \varepsilon^{-3\alpha} \ln \varepsilon^{-1} \|h\|_{L^1}$. Thus

$$\int_{\mathbb{T}^2} \varphi(x,\theta) (\mathcal{L}^n_{\theta}[h_{\theta} \circ Y_{\theta,n}^{-1}])(x) = \int_{\mathbb{T}^2} dx d\theta \varphi(x,\theta) h_*(x,\theta) \int_{\mathbb{T}} dy h(Y_{\theta,n}^{-1}(y),\theta) + \mathcal{O}(\sigma^n \|\varphi\|_{\mathcal{C}^0} \|h\|_{\mathcal{H}^1})$$
$$= \int_{\mathbb{T}^2} \varphi Ph + \mathcal{O}(\left[\sigma^n \varepsilon^{-3\alpha} \ln \varepsilon^{-1} + n^2 \varepsilon\right] \|\varphi\|_{\mathcal{C}^0} \|h\|_{L^1}).$$

To conclude we choose $n = c \ln \varepsilon^{-1}$, with c large enough, and obtain

$$\int_{\mathbb{T}^2} \varphi h = \nu^{-n} \int_{\mathbb{T}^2} \varphi P h + \nu^{-n} \mathcal{O}\left(\varepsilon(\ln \varepsilon^{-1})^2 \|\varphi\|_{\mathcal{C}^1} \|h\|_{L^1}\right)$$
$$= \int_{\mathbb{T}^2} \varphi P h + \mathcal{O}\left(\varepsilon^{\min\{1,\tau\}}(\ln \varepsilon^{-1})^2 \|\varphi\|_{\mathcal{C}^1} \|h\|_{L^1}\right).$$

It follows that there exists a $\beta_h \in \mathcal{H}^1(\mathbb{T}), \ \beta_h(\theta) = \int_{\mathbb{T}} dy h(y,\theta)$, such that

$$\|h - h_* \beta_h\|_{(\mathcal{C}^1)'} \le C_{\sharp} \varepsilon^{\min\{1,\tau\}} (\ln \varepsilon^{-1})^2 \|h\|_{L^1}$$

Appendix A. Proof of Lemma 2.6

To start, note that (1) coincides with the first part of (H4), which implies in particular that $\lambda > 2$. We have thus to prove only (H0) up to (H3) and the second part of (H4).

We start with **(H0)**. It suffices to show that $\partial_x f(p) > [\partial_\theta f \partial_x \omega - \partial_x f \partial_\theta \omega](p)$, for each $p \in \mathbb{T}^2$. The latter, by (2) and (4), is implied by $\lambda(1 - |\partial_\theta \omega|) > \lambda - 1 - ||\partial_\theta \omega||_{\infty} - ||\partial_x \omega||_{\infty}$ which, in turn, is implied by (3).

Next we prove (H1). Following [19] we start by proving that $D_p F(\mathbf{C}_u) \in \mathbf{C}_u$ and $D_p F^{-1}(\mathbf{C}_c) \in \mathbf{C}_c$. We consider a vector $(1, u) \in \mathbf{C}_u$ and we write a formula for the unstable slope field

(A.1)
$$D_p F(1, u) = (\partial_x f + u \partial_\theta f)(1, \Xi(p, u)), \qquad \Xi(p, u) = \frac{\partial_x \omega(p) + u \partial_\theta \omega(p) + u}{\partial_x f(p) + u \partial_\theta f(p)}$$

Notice that

(A.2)
$$\frac{d}{du}\Xi(p,\cdot) = \frac{\partial_x f + (\partial_\theta \omega \partial_x f - \partial_\theta f \partial_x \omega)}{(\partial_x f + u \partial_\theta f)^2} = \frac{\det DF(x,\theta)}{(\partial_x f + u \partial_\theta f)^2} > 0,$$

since det DF > 0 by (H0). Hence, checking the invariance of \mathbf{C}_u under DF is equivalent to showing that, for each $p \in \mathbb{T}^2$, $|\Xi(p, \pm \chi_u)| \leq \chi_u$. That is

(A.3)
$$\|\partial_{\theta}f\|_{\infty}\chi_{u}^{2} - (\lambda - \|\partial_{\theta}\omega\|_{\infty} - 1)\chi_{u} + \|\partial_{x}\omega\| \le 0.$$

Setting $\phi = \lambda - \|\partial_{\theta}\omega\|_{\infty} - 1$, inequality (A.3) has positive solutions since $\phi > 0$ by (4), which also implies

$$\frac{\phi^2 - 4\|\partial_\theta f\|_{\infty}\|\partial_x \omega\|_{\infty}}{2} \ge (\|\partial_\theta f\|_{\infty} - \|\partial_x \omega\|_{\infty})^2 > 0$$

Setting $\Phi_{\pm} = \phi \pm \sqrt{\phi^2 - 4 \|\partial_{\theta} f\|_{\infty} \|\partial_x \omega\|_{\infty}}$, we can then choose

(A.4)
$$\chi_u \in \left(\frac{\Phi_-}{2\|\partial_\theta f\|_{\infty}}, 1\right).$$

Note that the interval it is not empty due to (4).

On the other hand, if $(c, 1) \in \mathbf{C}_c$ we consider the center slope field

(A.5)
$$\Xi^{-}(p,c) = \frac{(1+\partial_{\theta}\omega(p))c - \partial_{\theta}f(p)}{\partial_{x}f(p) - \partial_{x}\omega(p)c}$$

and by an analogous computation we obtain $|\Xi^{-}(p, \pm \chi_{c})| < \chi_{c}$ if

(A.6)
$$\chi_c \in \left(\frac{\Phi_-}{2\|\partial_x \omega\|_{\infty}}, 1\right]$$

This also proves the second part of (H4).

Again, the interval it is not empty due to (4), we have thus proved (2.5). Next, by the invariance of the cones we can define real quantities λ_n, μ_n, u_n and c_n such that, for each $p \in \mathbb{T}^2$,⁶⁶

$$D_p F^n(1,0) = \lambda_n(p) (1, u_n(p)) ; \quad D_p F^n (c_n(p), 1) = \mu_n(p)(0, 1),$$

with $||u_n||_{\infty} \leq \chi_u$, $||c_n||_{\infty} \leq \chi_c$. Moreover, by definition

$$D_p F(c_n(p), 1) = \frac{\mu_n(p)}{\mu_{n-1}(F(p))} (c_{n-1}(F(p)), 1),$$

from which it follows, by (2.9),

$$\mu_n(p) = \mu_{n-1} \left(F(p) \right) \left(1 + \partial_\theta \omega(p) + c_n(p) \partial_x \omega(p) \right)$$

We set $b := \|\partial_{\theta}\omega\|_{\infty} + \chi_c \|\partial_x \omega\|_{\infty}$. Since $\|c_n\|_{\infty} \leq \chi_c$, we have that the condition (2) of the Lemma implies b < 1, and

(A.7)

$$(1-b)^n \le \mu_n(p) \le (1+b)^n$$

Similarly,

$$\begin{split} \lambda_n(p) &= \lambda_{n-1}(F(p))(\partial_x f(p) + \partial_\theta f(p) u_n(p)) \\ &= \prod_{k=0}^{n-1} \partial_x f(F^k p) \left(\partial_x f(F^k p) + \frac{\partial_\theta f(F^k)}{\partial_x f(F^k p)} u_{n-k}(F^k p) \right) \end{split}$$

which, setting $a := \chi_u \| \frac{\partial_{\theta} f}{\partial_x f} \|_{\infty}$, implies

(A.8)
$$(1-a)^n \prod_{k=0}^{n-1} \partial_x f(F^k(p)) \le \lambda_n(p) \le (1+a)^n \prod_{k=0}^{n-1} \partial_x f(F^k(p)),$$

which yields the second of (2.6) with $C_{\star} = 1$ and

(A.9)
$$\lambda_{+} = (1+a)\lambda \text{ and } \lambda_{-} = (1-a)\lambda_{+}$$

since, by the definition of χ_u in (A.4), we can check that $\lambda_- > 1$. To conclude, we need to check that $\frac{(1+b)}{(1-a)\lambda} < 1$, form which we deduce (H1). This is implied by

$$1 + \|\partial_{\theta}\omega\|_{\infty} + \|\partial_{x}\omega\|_{\infty} + \|\partial_{\theta}f\|_{\infty} < \lambda$$

which correspond to equation (4).

By (A.7), we can make the choice (2.10) which, by (2), implies $\mu \leq e^{2\chi_c \|\partial_x \omega\|_{\infty} + 2\|\partial_\theta \omega\|_{\infty}}$ and hence we obtain (H3) by (6).

It remains to prove (**H2**). Since $\lambda > 2$, F has rank at least two at each point, hence it is a covering map and each point has the same number of preimages, says d. Let then $\gamma : [0,1] \to \mathbb{T}^2$ be a smooth closed curve $\gamma(t) = (c(t), t)$ such that $\gamma' \in \mathbf{C}_c$ with homotopy class (0,1). If $p = (x,\theta) \in \gamma(t)$ then $F^{-1}(p) = \{q_1, \dots, q_d\}$. Note that, by the implicit function theorem, locally $F^{-1}\gamma$ is a curve, also, due to the above discussion, it belongs to the central cone. If we call η the local curve in $F^{-1}\gamma$ such that $\eta(0) = q_i$ we can extend it uniquely to a curve $\nu : [0,1] \to \mathbb{T}^2$. We will prove that $\nu(1) = q_i = \nu(0)$.

⁶⁶Note that the definition of λ_n differs from the one of λ_n^{\pm} in (2.4), since we are considering iteration of vectors inside the unstable cone. Nevertheless, they are related since there exists an integer m such that $F^m(\mathbb{R}^2 \setminus \mathbf{C}_c) \in \mathbf{C}_u$.

In turn this implies that $F^{-1}\gamma$ is the union of d closed curves ν_1, \dots, ν_d with $\nu'_i \in \mathbf{C}_c$, each one with homotopy class (0, 1), by the lifting property of covering maps (see [34, Proposition 1.30]). We argue by contradiction: assume that $\nu(1) = q_j \neq q_i$. Let $q_k = (x_k, \theta_k), k \in \{1, \dots, d\}$, then

$$\theta_i + \omega(x_i, \theta_i) = \theta_j + \omega(x_j, \theta_j)$$

implies

(A.10)
$$|\theta_i - \theta_j| \le \frac{\|\partial_x \omega\|_{\infty}}{1 - \|\partial_\theta \omega\|_{\infty}} |x_i - x_j|.$$

Hence the segment joining q_i and q_j belongs to the unstable cone if

(A.11)
$$\chi_u \ge \frac{\|\partial_x \omega\|_{\infty}}{1 - \|\partial_\theta \omega\|_{\infty}}$$

which is possible since (2) implies that this condition is compatible with (A.4). It follows that the image of the segment $\ell = \{tq_i + (1-t)q_j\}$ is an unstable curve and hence it cannot join p to itself without wrapping around the torus. In particular, if $q_i \neq q_j$, then the horizontal length of $F(\ell)$ must be larger than one. Then, setting $\delta = |x_i - x_j|$,

(A.12)
$$1 \le \int_0^1 \left| \langle e_1, D_{\ell(t)} F \ell'(t) \rangle \right| \le \|\partial_x f\|_\infty \left(1 + \chi_u \frac{\|\partial_\theta f\|_\infty}{\|\partial_x f\|_\infty} \right) |x_i - x_j| \le (1+a)\Lambda\delta.$$

To conclude we must show that ν cannot move horizontally by δ , whereby obtaining the wanted contradiction. Let $\nu(t) = (\alpha(t), \beta(t))$, then

$$\binom{c'(t)}{1} = \gamma'(t) = DF\nu' = \binom{\alpha'\partial_x f + \beta'\partial_\theta f}{\alpha'\partial_x\omega + (1+\partial_\theta\omega)\beta'}.$$

Since we know that $|c'| \leq \chi_c$ and $|\alpha'| \leq \chi_c |\beta'|$ we have

$$\begin{aligned} |\beta'| &\leq (1 - \chi_c \|\partial_x \omega\|_{\infty} - \|\partial_\theta \omega\|_{\infty})^{-1} \\ |\alpha'| &\leq \frac{\chi_c}{\lambda} + \frac{\|\partial_\theta f\|_{\infty}}{\lambda(1 - \chi_c \|\partial_x \omega\|_{\infty} - \|\partial_\theta \omega\|_{\infty})} \end{aligned}$$

I follows that it must be

$$\frac{1}{(1+a)\Lambda} \le \delta \le \int_0^1 |\alpha'(t)| dt \le \frac{\chi_c}{\lambda} + \frac{\|\partial_\theta f\|_\infty}{\lambda(1-\chi_c\|\partial_x \omega\|_\infty - \|\partial_\theta \omega\|_\infty)}$$

We thus have a contradiction if we can choose χ_c such that

$$\left(1 + \frac{\|\partial_{\theta}f\|_{\infty}}{\lambda}\right) \Lambda \left[\frac{\chi_c}{\lambda} + \frac{\|\partial_{\theta}f\|_{\infty}}{\lambda(1 - \|\partial_x\omega\|_{\infty} - \|\partial_{\theta}\omega\|_{\infty})}\right] < 1$$

which, by (A.6), is possible only if

$$\frac{\Phi_{-}}{2\|\partial_{x}\omega\|_{\infty}} < \left(1 + \frac{\|\partial_{\theta}f\|_{\infty}}{\lambda}\right)^{-1} \frac{\lambda}{\Lambda} - \frac{\|\partial_{\theta}f\|_{\infty}}{1 - \|\partial_{x}\omega\|_{\infty} - \|\partial_{\theta}\omega\|_{\infty}} =: A$$

Note that if $A \ge 1$, then the inequality is trivially satisfied. We must then consider only the case A < 1. A direct computation shows that the above inequality is implied by

(A.13)
$$\|\partial_{\theta}f\|_{\infty} < A[\phi - A\|\partial_{x}\omega\|_{\infty}] = A[\lambda - \|\partial_{\theta}\omega\|_{\infty} - 1 - A\|\partial_{x}\omega\|_{\infty}]$$

Let us set for simplicity $\varpi := \|\partial_{\theta}\omega\|_{\infty} + \|\partial_{x}\omega\|_{\infty}$. Since A < 1 the above equation is in turn implied by the following inequality

(A.14)
$$\|\partial_{\theta}f\|_{\infty} < \left[\left(1 + \frac{\|\partial_{\theta}f\|_{\infty}}{\lambda} \right)^{-1} \frac{\lambda}{\Lambda} - \frac{\|\partial_{\theta}f\|_{\infty}}{1 - \varpi} \right] \left(\lambda - (1 + \varpi) \right).$$

By elementary algebra (A.14) is equivalent to

(A.15)
$$\|\partial_{\theta}f\|_{\infty}(\|\partial_{\theta}f\|_{\infty}+1) < \frac{\lambda^2}{\Lambda} \left(1 - \frac{1}{\lambda + \varpi}\right).$$

Since $\lambda > 2$, (A.15) is implied by $\|\partial_{\theta} f\|_{\infty}(\|\partial_{\theta} f\|_{\infty} + 1) < \frac{1}{2}\lambda^{2}\Lambda^{-1}$, which is true if $\|\partial_{\theta} f\|_{\infty} < \frac{1}{2}\left(-1 + \sqrt{1 + 2\lambda^{2}\Lambda^{-1}}\right)$. Hence the conclusion by condition (5).

Appendix B. Proof of Lemma 5.2

We start considering $\varphi, \psi \in \mathcal{C}^{\rho}(\mathbb{T}^2, \mathbb{R})$. First we prove, by induction on ρ , that

(B.1)
$$\sup_{|\alpha|=\rho} \|\partial^{\alpha}(\varphi\psi)\|_{\mathcal{C}^{0}} \leq \sum_{k=0}^{\rho} \binom{\rho}{k} \sup_{|\beta|=\rho-k} \|\partial^{\beta}\varphi\|_{\mathcal{C}^{0}} \sup_{|\gamma|=k} \|\partial^{\gamma}\psi\|_{\mathcal{C}^{0}}$$

Indeed, it is trivial for $\rho = 0$ and

$$\begin{split} \|\partial_{x_{i}}\partial^{\alpha}(\varphi\psi)\|_{\mathcal{C}^{0}} &= \|\partial^{\alpha}(\psi\partial_{x_{i}}\varphi + \varphi\partial_{x_{i}}\psi)\| \\ &\leq \sum_{k=0}^{\rho} \binom{\rho}{k} \sup_{|\beta|=\rho-k} \|\partial^{\beta}\partial_{x_{i}}\varphi\|_{\mathcal{C}^{0}} \sup_{|\gamma|=k} \|\partial^{\gamma}\psi\|_{\mathcal{C}^{0}} + \sum_{k=0}^{\rho} \binom{\rho}{k} \sup_{|\beta|=\rho-k} \|\partial^{\beta}\partial_{x_{i}}\psi\|_{\mathcal{C}^{0}} \sup_{|\gamma|=k} \|\partial^{\gamma}\psi\|_{\mathcal{C}^{0}} \\ &\leq \sum_{k=0}^{\rho} \binom{\rho}{k} \sup_{|\beta|=\rho-k} \|\partial^{\beta}\varphi\|_{\mathcal{C}^{0}} \sup_{|\gamma|=k} \|\partial^{\gamma}\psi\|_{\mathcal{C}^{0}} + \sum_{k=0}^{\rho+1} \binom{\rho}{\rho+1-k} \sup_{|\beta|=\rho-k} \|\partial^{\beta}\varphi\|_{\mathcal{C}^{0}} \sup_{|\gamma|=k} \|\partial^{\gamma}\psi\|_{\mathcal{C}^{0}}, \end{split}$$

from which (B.1) follows taking the sup on α , *i* and since $\binom{\rho}{k} + \binom{\rho}{\rho+1-k} = \binom{\rho+1}{k}$. We then have the first statement of the Lemma, indeed

$$\begin{aligned} \|\varphi\psi\|_{\mathcal{C}^{\rho}} &\leq \sum_{k=0}^{\rho} 2^{\rho-k} \sum_{j=0}^{k} \binom{k}{j} \sup_{|\beta|=k-j} \|\partial^{\beta}\varphi\|_{\mathcal{C}^{0}} \sup_{|\gamma|=j} \|\partial^{\gamma}\psi\|_{\mathcal{C}^{0}} \\ &\leq \sum_{j=0}^{\rho} \sum_{l=0}^{\rho-j} \binom{\rho}{j} 2^{\rho-j-l} \sup_{|\beta|=l} \|\partial^{\beta}\varphi\|_{\mathcal{C}^{0}} \sup_{|\gamma|=j} \|\partial^{\gamma}\psi\|_{\mathcal{C}^{0}} \leq \|\varphi\|_{\mathcal{C}^{\rho}} \|\psi\|_{\mathcal{C}^{\rho}} \end{aligned}$$

since $\binom{\rho}{j} \leq 2^{\rho}$. The extension to function with values in the matrices follows trivially since we have chosen a norm in which the matrices form a norm algebra.

To prove the second statement we proceed again by induction on ρ . The case $\rho = 0$ is immediate since $\mathcal{K}_{0,0}$ contains only the zero string. Let us assume that the statement is true for every $k \leq \rho$ and prove it for $\rho + 1$. By equation (B.1) and the inductive hypothesis (5.5), we have, for each $|\alpha| = \rho + 1$,

$$\begin{aligned} \partial^{\alpha}(\varphi \circ \psi) &| \leq C_{\sharp} \sup_{|\beta|=\rho} \sup_{|\tau_{1}|,|\tau_{2}|=1} \left| \partial^{\beta} \left[(\partial^{\tau_{1}} \varphi) \circ \psi \cdot \partial^{\tau_{2}} \psi \right] \right| \\ &\leq C_{\rho} \sup_{|\tau_{1}|,|\tau_{2}|=1} \sup_{|\alpha_{0}|+|\alpha_{1}|=\rho} \left\| \partial^{\alpha_{0}} \left[(\partial^{\tau_{1}} \varphi) \circ \psi \right] \right\|_{\mathcal{C}^{0}} \left\| \partial^{\alpha_{1}} \partial^{\tau_{2}} \psi \right\|_{\mathcal{C}^{0}} \\ &\leq C_{\rho} \sup_{|\tau_{1}|=1} \sup_{\alpha_{0} \leq \rho} \left\| (\partial^{\tau_{1}} \varphi) \circ \psi \right\|_{\mathcal{C}^{\alpha_{0}}} \left\| D\psi \right\|_{\mathcal{C}^{\rho-\alpha_{0}}} \\ &\leq C_{\rho} C_{\rho}^{\star} \sup_{\alpha_{0} \leq \rho} \sum_{s=0}^{\alpha_{0}} \left\| \varphi \right\|_{\mathcal{C}^{s+1}} \sum_{k \in \mathcal{K}_{\alpha_{0},s}} \prod_{l=1}^{\alpha_{0}} \left\| D\psi \right\|_{\mathcal{C}^{l-1}}^{k_{l}} \cdot \left\| D\psi \right\|_{\mathcal{C}^{\rho-\alpha_{0}}} \\ &\leq C_{\rho} C_{\rho}^{\star} \sup_{\alpha_{0} \leq \rho} \sum_{s=0}^{\alpha_{0}} \left\| \varphi \right\|_{\mathcal{C}^{s+1}} \sum_{k \in \mathcal{K}_{\rho,s+1}} \prod_{l=1}^{\rho+1} \left\| D\psi \right\|_{\mathcal{C}^{l-1}}^{k_{l}} \\ &\leq C_{\rho} C_{\rho}^{\star} \sum_{s=0}^{\rho+1} \left\| \varphi \right\|_{\mathcal{C}^{s}} \sum_{k \in \mathcal{K}_{\rho+1,s}} \prod_{l=1}^{\rho+1} \left\| D\psi \right\|_{\mathcal{C}^{l-1}}^{k_{l}}. \end{aligned}$$

The result follows by choosing $C^{\star}_{\rho+1}$ large enough. \Box

APPENDIX C. BLACK-BOX: ITERATION OF CURVES

C.1. **Proof of Lemma 5.10.** Fix $\gamma \in \Gamma_j(c_*)$ and $n \in \mathbb{N}$. Let ν_n be a pre-image of γ under F^n and consider $\mathfrak{h} \in \mathfrak{H}^{\infty}$ such that $\nu_n = \mathfrak{h}_n \circ \gamma$. Let $h_n : \mathbb{T} \to \mathbb{T}$ be the diffeomorphism such that $\hat{\nu}_n = \nu_n \circ h_n$ is parametrized by vertical length. We then want to check properties c1, ..., c3) for $\hat{\nu}_n$. The first two follow immediately by assumption (**H2**), thus we only have to check property c3) (from Definition 5.5). By definition we have

(C.1)
$$F^n \hat{\nu}_n = \gamma \circ h_n.$$

Differentiating equation (C.1) twice we obtain

(C.2)
$$(\partial_t D_{\hat{\nu}_n} F^n) \hat{\nu}'_n + D_{\hat{\nu}_n} F^n \hat{\nu}''_n = \gamma'' \circ h_n (h'_n)^2 + \gamma' \circ h_n h''_n.$$

Similarly, if we differentiate equation (C.1) *j*-th times, we obtain

(C.3)
$$R_{j}(F^{n},\hat{\nu}_{n}) + D_{\hat{\nu}_{n}}F^{n}\hat{\nu}_{n}^{(j)} = \gamma^{(j)} \circ h_{n}(h_{n}')^{j} + Q_{j}(h_{n},\gamma) + \gamma' \circ h_{n} \cdot h_{n}^{(j)},$$

where, by (5.25), $R_j(F^n, \hat{\nu}_n) = A_{j,n} + B_{j,n}$ with $A_{j,n} \in \mathbf{C}_u$ and $\|B_{j,n}\| \leq \|\omega\|_{\mathcal{C}^r} \|\tilde{B}_{j,n}\|$. Moreover, $A_{j,n}, \tilde{B}_{j,n}$ are the sum of monomials, with coefficients depending only of $(\partial^{\alpha} F^n) \circ \hat{\nu}_n$ with $|\alpha| \leq j$, in the variables $\hat{\nu}_n^{(s)}$, $s \in \{0, \ldots, j-1\}$, where, if k_s is the degree of $\hat{\nu}^{(s)}$, then we have $\sum_{s=1}^{j-1} sk_s = j$.

the variables $\hat{\nu}_{n}^{(s)}$, $s \in \{0, \dots, j-1\}$, where, if k_s is the degree of $\hat{\nu}^{(s)}$, then we have $\sum_{s=1}^{j-1} sk_s = j$. Likewise the Q_j are the sum of monomials that are linear in $\gamma^{(\sigma)}$, $\sigma \in \{2, \dots, j-1\}$, and of degree p_s in $h_n^{(s)}$, $s \in \{1, \dots, j-\sigma+1\}$, such that $\sum_{s=1}^{j-\sigma+1} s p_s = j$.⁶⁷ In order to obtain an estimate for $\|\hat{\nu}_n^{(j)}\|$ it is convenient to introduce the vectors $\eta_{n,j} = D_{\hat{\nu}_n} F^n \hat{\nu}_n^{(j)}$. We then define the unitary vectors $\eta_{n,j}^{\perp}, \hat{\eta}_{n,j}$ such that $\langle \eta_{n,j}^{\perp}, \eta_{n,j} \rangle = 0$ and $\hat{\eta}_{n,j} = \frac{\eta_{n,j}}{\|\eta_{n,j}\|}$. Multiplying equation (C.3) by $\eta_{n,j}^{\perp}$ and $\hat{\eta}_{n,j}$ respectively, we obtain the system of equations

(C.4)
$$\langle \eta_{n,j}^{\perp}, R_j(F^n, \hat{\nu}_n) \rangle = \left\langle \eta_{n,j}^{\perp}, \gamma^{(j)} \circ h_n(h'_n)^j + Q_j(h_n, \gamma) + \gamma' \circ h_n \cdot h_n^{(j)} \right\rangle$$
$$\langle \hat{\eta}_{n,j}, R_j(F^n, \hat{\nu}_n) \rangle + \|\eta_{n,j}\| = \left\langle \hat{\eta}_{n,j}, \gamma^{(j)} \circ h_n(h'_n)^j + Q_j(h_n, \gamma) + \gamma' \circ h_n \cdot h_n^{(j)} \right\rangle.$$

Notice that, since $\hat{\nu}_n^{(j)}$ and $\gamma^{(j)}$, j > 1, are horizontal vectors, by the invariance of the unstable cone $\eta_{n,j} \in \mathbf{C}_u$. Moreover $\gamma' \in \mathbf{C}_c$ by assumption and $\|\eta_{n,j}^{\perp}\| = 1$, thus there exists $\vartheta \in (0,1)$ such that

(C.5)
$$|\langle \eta_{n,j}^{\perp}, \gamma' \circ h_n \rangle| \ge \vartheta \|\gamma' \circ h_n\| \ge \vartheta.$$

Using (C.5), setting $\tilde{R}_{j,n} := |\langle \eta_{n,j}^{\perp}, R_j(F^n, \hat{\nu}_n) \rangle| + ||Q_j(h_n, \gamma)||$, and $R_{j,n} := ||R_j(F^n, \hat{\nu}_n)|| + ||Q_j(h_n, \gamma)||$, equation (C.4) yields

(C.6)
$$|h_{n}^{(j)}| \leq \frac{|h_{n}'|^{j} \|\gamma^{(j)} \circ h_{n}\| \chi_{u} + \tilde{R}_{j,n}}{\vartheta \|\gamma' \circ h_{n}\|}, \\ \|\eta_{n,j}\| \leq \|\gamma^{(j)} \circ h_{n}\| \|h_{n}'|^{j} + \|\gamma' \circ h_{n}\| \|h_{n}^{(j)}\| + R_{j,n}$$

By equation (C.1) it follows that

(C.7)
$$\|\hat{\nu}'_n\| = |h'_n| \| (D_{\hat{\nu}_n} F^n)^{-1} \gamma' \circ h_n \|,$$

which yields, by (2.6) and the fact that $\hat{\nu}'_n = ((\pi_1 \circ \hat{\nu}_n)', 1) \in \mathbf{C}_c$,

(C.8)
$$\frac{\mu^{-n}}{C_{\star}\sqrt{1+\chi_c^2}} \le |h'_n| \le \frac{C_{\star}\mu^n \|\hat{\nu}'_n\|}{\|\gamma' \circ h_n\|} \le \sqrt{1+\chi_c^2} C_{\star}\mu^n =: \bar{C}_{\star}\mu^n.$$

⁶⁷The reader can check this by induction (equation (C.2) gives the case j = 2). E.g., if a term T in R_j has the form $T = \prod_{s=0}^{j-1} \alpha_s(\hat{\nu}_n^{(s)})$ where $\alpha_s(x)$ is homogeneous of degree k_s in x, then $\partial_t T$ will be a sum of terms of the same type with homogeneity degrees k'_s . Let us compute such homogeneity degrees: if the derivative does not hit a $\hat{\nu}_n^{(s)}$, s > 0, then, by the chain rule, we will get a monomial with $k'_1 = k_1 + 1$ while all the other homogeneity degree are unchanged: $k'_s = k_s$ for s > 1. Hence, $\sum_{s=0}^{j} k'_s = j + 1$. If the derivative hits one $\hat{\nu}_n^{(i)}$, then it produces a monomial with $k'_s = k_s$ for $s \notin \{i, i+1\}$ while $k'_i = k_i - 1$ and $k'_{i+1} = k_{i+1} + 1$. Then $\sum_{s=0}^{j} k'_s = j - ik_i - (i+1)k_{i+1} + i(k_i - 1) + (i+1)(k_{i+1} + 1) = j + 1$.

Using this in (C.6) and observing that $\|\eta_{n,j}\| = \|D_{\hat{\nu}_n} F^n \hat{\nu}_n^{(j)}\| \ge \lambda_n^- \|\hat{\nu}_n^{(j)}\|$, we obtain

(C.9)
$$\|\hat{\nu}_{n}^{(j)}\| \leq \|\gamma^{(j)} \circ h_{n}\| (\lambda_{n}^{-})^{-1} (\bar{C}_{\star} \mu^{\bar{n}})^{j} A + R_{j,n}^{\star},$$

where $A = (1 + \vartheta^{-1})$ and $R_{j,n}^{\star} = (\lambda_n^-)^{-1} A R_{j,n}$. Next we choose \bar{n} . The choice depends on a uniform constant $C_{\flat} \geq \{3\bar{C}_{\star}, \frac{2\bar{C}_{\star}}{\vartheta}, C_{\sharp}r\}^+$ that will be chosen, large enough, in (C.24):⁶⁸

(C.10)
$$A(\bar{C}_{\star}\mu^{\bar{n}})^{r+1}(\lambda_{\bar{n}})^{-1}\left[1+C_{\flat}^{(r+1)^{2}}\mu^{\bar{n}(r+1)!}\right] < 1, \text{ and set}$$
$$\eta := \left(A(\bar{C}_{\star}\mu^{\bar{n}})^{r+1}(\lambda_{\bar{n}})^{-1}\left[1+C_{\flat}^{(r+1)^{2}}\mu^{\bar{n}(r+1)!}\right]\right)^{\frac{1}{2\bar{n}(r+1)!}} < 1$$

Note in particular that, as \bar{C}_{\star} and ϑ are uniform constants, so are both \bar{n} and η .

We are ready to conclude. For j = 1 the Lemma is trivial since $\|\hat{\nu}_{\bar{n}}\| \leq \sqrt{1 + \chi_c^2}$ and $h_{\bar{n}}$ can be bounded by (C.8), provided $C_{\flat} \geq \bar{C}_{\star}$. Equation (C.2) implies that $\|R_2(F^{2\bar{n}}, \hat{\nu}_{2\bar{n}})\| \leq C_{\bar{n}}$ while $\|\tilde{R}_2(F^{2\bar{n}}, \hat{\nu}_{2\bar{n}})\| \leq C_{\bar{n}}C_F$ and $Q_2 = 0$,⁶⁹ thus $R_{2,2\bar{n}} \leq C_{\bar{n}}$. Then the first of (C.6), remembering equations (C.8) and (C.9) imply

(C.11)
$$\begin{aligned} \|h_n^{(2)}\| &\leq C_{\sharp} C_F(\bar{C}_*^2 c_\star \mu^n + C_{\bar{n}}) \quad \forall n \leq 2\bar{n} \\ \|\hat{\nu}_n^{(2)}\| &\leq A(\lambda_n^-)^{-1} \left\{ c_\star (\bar{C}_\star \mu^n)^2 + C_{\bar{n}} \right\}. \end{aligned}$$

Next, we proceed by induction on $j < \ell$ to prove that for each $\bar{n} \leq n \leq 2\bar{n}$

(C.12)
$$\begin{aligned} \|h_n^{(j)}\| &\leq C_\flat^{j^2} C_F\left(c_\star^{(j-1)!} \mu^{j!n} + C_{\bar{n}}\right) \\ \|\hat{\nu}_n^{(j)}\| &\leq (\eta^n c_\star + c/2)^{(j-1)!}. \end{aligned}$$

By (C.11) we have the case j = 2, let us assume it for all $2 \le s \le j$. Recalling the structure of R_j, \tilde{R}_j, Q_j , see equation (C.3) and comments thereafter, and provided $c_* \ge C_{\bar{n}}$ we have⁷⁰

$$R_{j+1,n} \leq C_{\sharp} \left\{ \sum_{k} c_{\star}^{\sum_{s=1}^{j} (s-1)!k_{s}} + C_{\flat}^{(j+1)j} \sum_{\sigma=2}^{j} \sum_{p} c_{\star}^{(\sigma-1)! + \sum_{s=2}^{j+2-\sigma} p_{s}(s-1)!} \mu^{n \sum_{s=1}^{j+2-\sigma} p_{s}(s-1)!} + C_{\bar{n}} \right\}$$
$$\tilde{R}_{j+1,n} \leq C_{\sharp} C_{F} \left\{ \sum_{k} c_{\star}^{\sum_{s=1}^{j} (s-1)!k_{s}} + C_{\flat}^{(j+1)j} \sum_{\sigma=2}^{j} \sum_{p} c_{\star}^{(\sigma-1)! + \sum_{s=2}^{j+2-\sigma} p_{s}(s-1)!} \mu^{n \sum_{s=1}^{j+2-\sigma} p_{s}(s-1)!} + C_{\bar{n}} \right\}.$$

It is convenient to define

$$\tau_j = \begin{cases} 2 & \text{if } j = 2\\ 6 & \text{if } j = 3\\ j! - 1 & \text{if } j > 3 \end{cases}$$

Note if j = 2, then $\sum_{s=1}^{2} (s-1)! k_s = 2 = \tau_2$, $\sigma = 2$ and $1 + \sum_{s=2}^{2} p_s(s-1)! = 2 = \tau_2$. If j > 2 note that $k_j \leq 1$, otherwise $\sum_{s=1}^{j} sk_s > j+1$, hence

$$\sum_{s=1}^{j} (s-1)! k_s \le (j-3)! \sum_{s=1}^{j-1} sk_s + (j-1)! k_j = (j-3)! (j+1-jk_j) + (j-1)! k_j$$
$$\le (j-3)! \{j+1, j^2 - 3j + 3\}^+ \le \tau_j.$$

If $\sigma = j$, then

$$(\sigma - 1)! + \sum_{s=1}^{j+2-\sigma} p_s(s-1)! = (j-1)! + j + 1 \le \tau_j.$$

 $^{^{68}}$ Note that this is possible due to hypothesis (H3), see (2.7).

⁶⁹Recall that $C_F = \chi_u + \|\omega\|_{\mathcal{C}^r}$, as defined just after (5.26).

⁷⁰The sum in the exponent of c_{\star} starts from 2 since bound of h' does not contains c_{\star} .

If $\sigma = j - 1$, then

$$(\sigma-1)! + \sum_{s=1}^{j+2-\sigma} p_s(s-1)! = (j-2)! + j + 1 \le (j-1)! + j + 1 \le \tau_j.$$

On the other hand if $\sigma < j - 1$, then we have

$$(\sigma - 1)! + \sum_{s=1}^{j+2-\sigma} p_s(s-1)! \le (j-3)! + (j-\sigma)!(j+1) \le (j-3)![1+(j-2)(j+1)] \le (j-3)!(j^2-j-1) \le \tau_j.$$

Accordingly, since the sums in k and p have at most j^j terms, and $j \leq r$, since $C_{\flat} \geq \{\frac{2\bar{C}_*}{\vartheta}, C_{\sharp}r\}^+$,

(C.13)

$$R_{j+1,n} \leq C_{\flat}^{(j+1)^{2}} \left\{ c_{\star}^{\tau_{j}} \mu^{n(j-1)!(j+1)} + C_{\bar{n}} \right\}$$

$$\tilde{R}_{j+1,n} \leq \frac{1}{2} C_{\flat}^{(j+1)^{2}} C_{F} \left\{ c_{\star}^{\tau_{j}} \mu^{n(j-1)!(j+1)} + C_{\bar{n}} \right\}$$

$$R_{j+1,n}^{\star} \leq A(\lambda_{n}^{-})^{-1} C_{\flat}^{(j+1)^{2}} \left\{ c_{\star}^{\tau_{j}} \mu^{n(j-1)!(j+1)} + C_{\bar{n}} \right\}.$$

Let us show the first of (C.12). Substituting the above in the first of (C.6) and using (C.8) we have

$$\|h_n^{(j+1)}\| \le \frac{(\bar{C}_\star \mu^n)^{j+1}}{\vartheta} c_\star^{j!} C_F + \frac{1}{2} C_F C_\flat^{(j+1)^2} (c_\star^{j!} \mu^{n(j-1)!(j+1)} + C_{\bar{n}}).$$

We can then write

$$||h_n^{(j+1)}|| \le C_b^{(j+1)^2} C_F\left(c_\star^{j!} \mu^{n(j+1)!} + C_{\bar{n}}\right).$$

which is the first of (C.12) for $n \leq 2\bar{n}$.

Next, we substitute (C.13) in (C.9), using (C.10), and choosing $2c > C_{\bar{n}}$ we write

$$\|\hat{\nu}_{n}^{(j+1)}\| \leq \eta^{nj!} \left\{ c_{\star}^{j!} + C_{\bar{n}} \right\} \leq \left\{ \eta^{n} c_{\star} + \varepsilon/2 \right\}^{j!}.$$

Hence also the second of (C.12) is satisfied for $n \leq 2\bar{n}$.

In particular $\hat{\nu}_n \in \Gamma_{\ell}(\eta^n c_{\star} + c/2)$ for each $\ell \leq r$ and $\bar{n} \leq n \leq 2\bar{n}$. Next, let us set $c_{\star,1} = \eta^{\bar{n}} c_{\star} + c/2 \leq c_{\star}$, and, for each integer $k \geq 2$,

$$\frac{\mathbb{c}}{2} \le c_{\star,k} := \eta^{\bar{n}} c_{\star,k-1} + \frac{\mathbb{c}}{2} \le \eta^{\bar{n}k} c_{\star} + \frac{\mathbb{c}}{2(1-\eta^{\bar{n}})}$$

It follows that $\hat{\nu}_{k\bar{n}} \in \Gamma_{\ell}(c_{\star,k})$ and, for all $m \in \{\bar{n}, \dots 2\bar{n}\},^{71}$

$$\hat{\nu}_{k\bar{n}+m} = \mathfrak{h}_{k\bar{n}+m-1}^* \circ \cdots \circ \mathfrak{h}_{k\bar{n}+1}^* \circ \hat{\nu}_{k\bar{n}} \circ h_{m,k+1}^*,$$

and, by (C.12),

(C.14)
$$\|h_{m,(k+1)}^*\|_{\mathcal{C}^j} \le C_{\flat}^{j^2} C_F\left(c_{\star,k}^{(j-1)!} \mu^{j!m} + C_{\bar{n}}\right).$$

Hence, applying iteratively the above argument to $\hat{\nu}_n$ for $k\bar{n} \leq n \leq (k+1)\bar{n}$, we obtain the second of (C.12) for each $n \geq \bar{n}$.

It remains to prove the estimate for h_n , $n \ge 2\bar{n}$. We write $n = m + k\bar{n}$, $m \in \{\bar{n}, \ldots, 2\bar{n}\}$ and

(C.15)
$$h_n = h_{m,k+1}^* \circ h_{\bar{n},k}^* \circ \dots \circ h_{\bar{n},1}^* = h_{m,k+1}^* \circ h_{k\bar{n}}.$$

Note that (C.8) yields $||h_n||_{C^1} \leq C_{\flat} \mu^n$, since $C_{\flat} \geq 3\bar{C}_{\star}$. It is then natural to start by investigating the second derivative. In fact, it turns out to be more convenient to study the following ratio

(C.16)
$$\frac{h_n''}{h_n'} = \left(\log[(h_{m,k+1}^*)' \circ h_{k\bar{n}}]\right)' + \frac{h_{k\bar{n}}''}{h_{k\bar{n}}'} =: Q_1 + Q_2.$$

To estimate the norm of such a ratio we start estimating the sup norm, then we will proceed by induction.

⁷¹Recall the definition of \mathfrak{h}_n^* in (5.8).

Since (5.17) and (C.7) imply $|h'_{\bar{n},i}| \ge c_0 \mu^{-n}$ for each *i*, for some uniform constant c_0 , formula (5.5) and (C.14) yield

(C.17)
$$\|(\log h_{m,k+1}^{*\prime})'\|_{\mathcal{C}^{\ell}} \le C_{\sharp} C_F C_{\flat}^{(\ell+2)^2} c_{\star,k}^{(\ell+1)!} \mu^{(\ell+2)!m}.$$

It follows

$$\|Q_1\|_{\mathcal{C}^0} \le C_{\sharp} C_F C_{\flat}^4 c_{\star,k} \mu^{2m} \mu^{k\bar{n}} \le C_{\sharp} C_F C_{\flat}^4 c_{\star} \mu^n.$$

To estimate $||Q_2||_{\mathcal{C}^0}$ we write

(C.18)
$$Q_2 = \frac{h_{k\bar{n}}''}{h_{k\bar{n}}'} = \frac{\left(\prod_{i=1}^k h_{\bar{n},i}^{*\prime} \circ h_{i\bar{n}}\right)'}{\prod_{i=1}^k h_{\bar{n},i}^{*\prime} \circ h_{i\bar{n}}} = \left(\log\prod_{i=1}^k h_{\bar{n},i}^{*\prime} \circ h_{i\bar{n}}\right)' = \sum_{i=1}^k \left(\log h_{\bar{n},i}^{*\prime} \circ h_{i\bar{n}}\right)'.$$

Using formulae (C.8), (C.17) and (C.16) we have, since $\bar{n} \leq m$,

(C.19)
$$\begin{aligned} \|Q_2\|_{\mathcal{C}^0} &\leq \sum_{i=1}^k \|(\log h_{\bar{n},i}^{*\prime})'\|_{\mathcal{C}^0} \|h_{i\bar{n}}'\|_{\mathcal{C}^0} \leq C_{\sharp} C_{\flat}^4 C_F c_{\star,1} \mu^{2\bar{n}} \sum_{i=1}^k \mu^{i\bar{n}} \\ &\leq C_{\sharp} C_{\flat}^4 C_F c_{\star,1} \mu^{2\bar{n}} \frac{\mu^{k\bar{n}} - 1}{\mu - 1} \leq \mu^{2\bar{n}} C_{\flat}^4 C_F c_{\star,1} C_{\mu,k\bar{n}} \mu^{k\bar{n}} \leq C_{\flat}^4 C_F c_{\star,1} C_{\mu,n} \mu^n. \end{aligned}$$

Hence, using the above and (C.8), it follows by (C.18)

(C.20)
$$\|h_n''\|_{\mathcal{C}^0} \le C_{\sharp} \left\|\frac{h_n''}{h_n'}\right\|_{\mathcal{C}^0} \|h_n'\|_{\mathcal{C}^0} \le C_{\sharp}\mu^n \left\|\frac{h_n''}{h_n'}\right\|_{\mathcal{C}^0} \le C_{\flat}^4 C_F c_{\star} C_{\mu,n}\mu^{2n}.$$

This proves the second of (5.29). We can now prove the general case by induction on $j \leq \ell$. Assume it true for all i < j. Using again (5.5), by the inductive assumption and C.17 we have

(C.21)
$$\begin{aligned} \|Q_1\|_{\mathcal{C}^{j-1}} &= \|(\log[(h_{m,k+1}^*)'])' \circ h_{k\bar{n}}\|_{\mathcal{C}^{j-1}} \|h_{k\bar{n}}'\|_{\mathcal{C}^{j-1}} \\ &\leq C_{\flat}^{4(j-2)!} (c_{\star,1}^{j!} C_{\mu,n}^{a_j} + 1) \mu^{nj!}. \end{aligned}$$

On the other hand, by formulae (C.18), (5.5), (C.17) and the inductive assumption

$$\begin{aligned} \|Q_2\|_{\mathcal{C}^{j-1}} &\leq C_{\sharp} \sum_{i=1}^k \|(\log[(h_{m,k+1}^*)'])' \circ h_{k\bar{n}}\|_{\mathcal{C}^{j-1}} \|h_{i\bar{n}}'\|_{\mathcal{C}^{j-1}} \\ &\leq C_{\sharp} \sum_{i=1}^k \|(\log[(h_{m,k+1}^*)'])'\|_{\mathcal{C}^{j-1}} \|h_{i\bar{n}}'\|_{\mathcal{C}^{j-2}}^{j-1} \|h_{i\bar{n}}'\|_{\mathcal{C}^{j-1}} \\ &\leq C_{\sharp} C_{\flat}^{(j+1)^2 + 2 \cdot (j-1)! + 2 \cdot j!} C_F c_{\star,k}^{j!} \mu^{(j+1)!m} \sum_{i=1}^k \sum_{q=0}^{j-1} ([c_{\star}^{(j-1)!} C_F + 1] C_{\mu,i\bar{n}}^{a_j} \mu^{j!i\bar{n}})^q \\ &\leq C_{\sharp} C_{\flat}^{3\cdot j!} [c_{\star}^{2\cdot j!} C_F + 1] \mu^{(j+1)!m} \sum_{i=1}^k \sum_{q=0}^{j-1} (C_{\mu,i\bar{n}}^{a_j} \mu^{j!i\bar{n}})^q. \end{aligned}$$
To estimate the last sum notice that by definition

To estimate the last sum, notice that by definition

$$\begin{array}{ll} \mathrm{i} \ 1 \leq \! C_{\mu,\bar{n}i} \leq C_{\mu,\bar{n}k}, & \forall i \leq k, \\ \mathrm{ii} \ C_{\mu^a,n} \leq C_{\mu,n}, & \forall a > 1, \end{array}$$

Hence,

$$\sum_{i=1}^{k} \sum_{q=0}^{j-1} (C^{a_j}_{\mu,\bar{n}i} \mu^{j!i\bar{n}})^q \le C^{a_j j}_{\mu,\bar{n}k} \sum_{i=1}^{k} \mu^{j!(j-1)\bar{n}i} \le C^{a_j(j-1)+1}_{\mu,\bar{n}k} \mu^{j!(j-1)\bar{n}k}.$$

Using this in (C.22) we obtain

(C.23)
$$\|Q_2\|_{\mathcal{C}^{j-1}} \le C_{\sharp} C_{\flat}^{3 \cdot j!} [C_F c_{\star,i}^{2 \cdot j!} + 1] \mu^{(j+1)!m} C_{\mu,\bar{n}k}^{a_j(j-1)+1} \mu^{j!(j-1)\bar{n}k} .$$

Therefore, by the inductive assumption, equations (C.21), (C.23) and (C.16), and provided we choose C_{\flat} large enough, we finally have

(C.24)
$$\begin{aligned} \|h_n''\|_{\mathcal{C}^{j-1}} &\leq \left\|\frac{h_n''}{h_n'}\right\|_{\mathcal{C}^{j-1}} \|h_n\|_{\mathcal{C}^j} \\ &\leq C_\flat^{2(j+1)!} [C_F c_{\star,1}^{(j+1)!} + 1] C_{\mu,n}^{a_{j+1}} \mu^{(j+1)!n} \end{aligned}$$

C.2. **Proof of Lemma 5.13.** Let $\mathfrak{h} \in \mathfrak{H}^{\infty}$ be such that $\nu_m = \mathfrak{h}_m \gamma$. Recalling (5.6), we can apply (5.5) and we have for each $j \leq r$

(C.25)
$$\|\nu_m\|_{\mathcal{C}^{j+1}} = \|\mathfrak{h}_m \circ \gamma\|_{\mathcal{C}^{j+1}} \le C_{\sharp} (\|\Delta_{\gamma}\|_{\infty} \Lambda^m)^j.$$

We set $\phi(t) := (\pi_2 \circ \nu_m)(t)$. By (5.33) there exists $c_{u,\gamma} \ge \chi_u \mu^{-m}$ such that we have $|\phi'| > c_{u,\gamma} > 0$, so it is well defined the diffeomorphism $h_m(t) = \phi^{-1}(t)$, and $\hat{\nu}_m = \nu_m \circ h_m$ is parametrized by vertical length. We want to estimate the higher order derivatives of h_m using a formula for inverse functions given in [39]. For the reader convenience we write it down here for our case:

(C.26)
$$h_m^{(j+1)}(t) = \frac{d^{j+1}\phi^{-1}(t)}{dt^{j+1}}$$
$$= \sum_{k=0}^{j} [\phi' \circ \phi^{-1}(t)]^{-j-k-1} \sum_{\substack{b_1 + \dots + b_k = j+k \\ b_l \ge 2}} B_{j,k,\{b_l\}_{l=1}^k} \prod_{l=1}^k \phi^{(b_l)} \circ \phi^{-1}(t),$$

where $B_{j,k,\{b_l\}_{l=1}^k} = \frac{(j+k)!}{k!b_1!\cdots b_k!}$. It follows by (C.25) and (C.26) that for each t

(C.27)
$$|h_m^{(j+1)}(t)| \le C_{\sharp} \left(c_{u,\gamma}^{-2} \| \Delta_{\gamma} \|_{\infty} \Lambda^m \right)^j$$

By (C.25), (C.27) and formula (5.5) for the composition,

(C.28)
$$\begin{aligned} \|\hat{\nu}_{m}\|_{\mathcal{C}^{j+1}} &= \|\nu_{m} \circ h_{m}\|_{\mathcal{C}^{j+1}} \leq C_{\sharp} \sum_{s=0}^{j+1} \|\hat{\nu}_{m}\|_{\mathcal{C}^{s}} \sum_{k \in \mathcal{K}_{\rho,s}} \prod_{l \in \mathbb{N}} \|h_{m}\|_{\mathcal{C}^{l}}^{k_{l}} \\ &\leq C_{\sharp} (\bar{c}_{u,\gamma} \|\Delta_{\gamma}\|_{\infty} \Lambda^{c_{\sharp}m})^{2j} \leq (\bar{c}_{u,\gamma} \|\Delta_{\gamma}\|_{\infty} \Lambda^{c_{\sharp}m})^{(j+1)!}, \end{aligned}$$

where $\bar{c}_{u,\gamma} = \{c_{u,\gamma}^{-2}, 1\}^+$. Hence, setting $c_{\star}(m) = \bar{c}_{u,\gamma} \|\Delta_{\gamma}\|_{\infty} \Lambda^{c_{\sharp}m}$ we have that $\hat{\nu}_m \in \Gamma_j(c_{\star}(m))$. Since $\overline{m} > m > \overline{n}$ we can apply Lemma 5.10 and we have that the curve $\hat{\nu}_{\overline{m}} = \nu_m \circ h_{\overline{m}}$ belongs to $\Gamma_j(\eta^{\overline{m}}c_{\star}(m) + \frac{c}{2})$. By definition, $c_{\star}(m) \leq \chi_u^{-2} \|\Delta_{\gamma}\|_{\infty} (\mu\Lambda)^m$ and by Corollary 5.11, $c \geq \overline{c}_{u,\gamma}$ (since $j \geq 3$), having chosen ϖ large enough. The statement then follows choosing $\overline{m} = \sigma m$, with σ defined in (5.34).

C.3. **Proof of Lemma 5.14.** We use the notation of (C.1). To prove the first of (5.37) it is convenient to go back to equation (C.2) and, recalling (5.23), (5.26), for each $v \in \mathbb{R}^2$, ||v|| = 1, we have

(C.29)

$$\begin{aligned} \left| \langle v, \hat{\nu}_{n}'' \rangle - \langle v, \hat{\nu}_{n}' \rangle \frac{h_{n}''}{h_{n}'} \right| &\leq |\langle v, (D_{\hat{\nu}_{n}}F^{n})^{-1}\gamma'' \circ h_{n}(h_{n}')^{2} \rangle| \\ &+ \sum_{k=0}^{n-1} \sum_{i=1}^{2} \left| \langle v, (D_{\hat{\nu}_{n}}F^{k+1})^{-1} \left[\partial_{x_{i}}D_{F^{k}(\hat{\nu}_{n})}F \right] D_{\hat{\nu}_{n}}F^{k}\hat{\nu}_{n}' \rangle \right| \, \| (D_{\hat{\nu}_{n}}F^{k})\hat{\nu}_{n}' \| \\ &\leq |\langle v, (D_{\hat{\nu}_{n}}F^{n})^{-1}\gamma'' \circ h_{n}(h_{n}')^{2} \rangle| \\ &+ C_{\sharp} \sum_{k=0}^{n-1} \left(\frac{1}{\lambda_{k+1}^{-1}(\hat{\nu}_{n}(t))} + C_{F}\mu^{k+1} \right) \| (D_{\hat{\nu}_{n}}F^{k})\hat{\nu}_{n}' \|^{2}. \end{aligned}$$

Note that, recalling (5.12), for each $n \leq n_{\star} \leq c_2^{-} \log \chi_u^{-1}$ we have $(D_{\hat{\nu}_n(t)}F^n)^{-1}e_1 \notin \mathbf{C}_c$. Consequently

(C.30)
$$|\langle v, (D_{\hat{\nu}_n} F^n)^{-1} \gamma'' \circ h_n \rangle| \le (\lambda_n^-(\hat{\nu}_n(t)))^{-1} \|\gamma'' \circ h_n(t)\|, \quad \forall n \le n_\star.$$

Next, if v is perpendicular to $\hat{\nu}'_n$, then it must be $|v_2| \leq \chi_c |v_1|$, hence

(C.31)
$$|\langle v, \hat{\nu}_n'' \rangle| = |v_1| \|\hat{\nu}_n''\| \ge (1 + \chi_c^2)^{-\frac{1}{2}} \|\hat{\nu}_n''\|$$

On the other hand, if v is perpendicular to $\hat{\nu}''_n$, then $v = e_2$ and $|\langle v, \hat{\nu}'_n \rangle| = 1$. Accordingly, using (C.29) first with v orthonormal to $\hat{\nu}'_n$ and then with $v = e_2$, and recalling Proposition 5.9 and equations (2.6), (C.8) we have for $n \le n_{\star}$,⁷²

(C.32)
$$\|\hat{\nu}_{n}''(t)\| \leq (1+\chi_{c}^{2})^{\frac{1}{2}} (\lambda_{n}^{-}(\hat{\nu}_{n}(t)))^{-1} C_{\flat}^{2} \mu^{2n} \|\gamma'' \circ h_{n}(t)\| + \sum_{k=0}^{n-1} C_{\star}^{2} \mu^{2k} C_{F,k},$$

$$\|h_n''/h_n'\| \le (\lambda_n^-(\hat{\nu}_n(t)))^{-1} C_{\flat}^2 \mu^{2n} \|\gamma'' \circ h_n(t)\| + \sum_{k=0} C_{\star}^2 \mu^{2k} C_{F,k}$$

where $C_{F,k} = C_{\sharp} \left(\frac{1}{\lambda_{k+1}^{-}(\hat{\nu}_{n}(t))} + C_{F} \mu^{k+1} \right)$. Setting $\bar{c}_{n_{\star}} = \left[(1 + \chi_{c}^{2})^{\frac{1}{2}} C_{\flat}^{2} \right]^{\frac{1}{n_{\star}}}$ we obtain

(C.33)
$$\|\hat{\nu}_{n_{\star}}''(t)\| \leq \bar{c}_{n_{\star}}^{n_{\star}} \mu^{2n_{\star}} (\lambda_{n_{\star}}^{-}(\hat{\nu}_{n_{\star}}(t)))^{-1} \|\gamma'' \circ h_{n_{\star}}(t)\| + \sum_{k=0}^{n_{\star}-1} C_{\star}^{2} \mu^{2k} C_{F,k}.$$

Condition (5.1) implies $\bar{c}_{\hat{n}}^{\hat{n}}\mu^{2\hat{n}}\lambda_{-}^{-\hat{n}} \leq \frac{1}{2}$, so we can proceed by induction since, setting $h_{l,m}^* = h_{ln_{\star}+m} \circ h_{ln_{\star}}^{-1}$, if $n = ln_{\star} + m$, $m \leq n_{\star}$, then

(C.34)
$$\begin{aligned} \|\hat{\nu}_{n}''(t)\| &\leq \bar{c}_{n_{\star}}^{n_{\star}} \mu^{2m} (\lambda_{m}^{-}(\hat{\nu}_{n}(t)))^{-1} \|\hat{\nu}_{ln_{\star}}'' \circ h_{l,m}^{*}(t)\| + \sum_{k=0}^{n_{\star}-1} C_{\star}^{2} \mu^{2k} C_{F,k} \\ &\leq \bar{c}_{n_{\star}}^{n+n_{\star}} \mu^{2n} (\lambda_{m}^{-}(\hat{\nu}_{n}(t)))^{-1} (\lambda_{n_{\star}}^{-}(\hat{\nu}_{ln_{\star}} \circ h_{l,m}^{*}(t)))^{-1} \dots (\lambda_{n_{\star}}^{-}(\gamma \circ h_{n}(t)))^{-1} c_{\star} \\ &+ \sum_{s=1}^{l} \bar{c}_{n_{\star}}^{sn_{\star}} \mu^{2sn_{\star}} \lambda_{-}^{-sn_{\star}} \sum_{k=0}^{n_{\star}-1} C_{\star}^{2} \mu^{2k} C_{F,k} \\ &\leq \bar{c}_{n_{\star}}^{n+n_{\star}} \mu^{2n} \bar{c}_{\flat} \bar{c}_{\flat}^{\frac{n}{n_{\star}}} (\lambda_{n}^{+}(\gamma \circ h_{n}(t)))^{-1} c_{\star} + C_{\mu,n_{\star}} \mu^{3n_{\star}} C_{3}, \end{aligned}$$

for some appropriate C_3 . This implies the first of the (5.37).

It remains to bound the third derivative of $\hat{\nu}_n$. The strategy is basically the same. Recalling that $\hat{\nu}'_n = (D_{\hat{\nu}_n} F^n)^{-1} \gamma' \circ h_n h'_n$, we differentiate this expression twice and multiply by a unitary vector v orthogonal to $\hat{\nu}'_n$:

(C.35)
$$\langle \hat{\nu}_{n}^{\prime\prime\prime}, v \rangle = \left\langle [(D_{\hat{\nu}_{n}}F^{n})^{-1}]^{\prime\prime}\gamma^{\prime} \circ h_{n}h_{n}^{\prime} + 2[(D_{\hat{\nu}_{n}}F^{n})^{-1}]^{\prime}(\gamma^{\prime\prime} \circ h_{n}(h_{n}^{\prime})^{2} + \gamma^{\prime} \circ h_{n}h_{n}^{\prime\prime}) + [(D_{\hat{\nu}_{n}}F^{n})^{-1}](\gamma^{\prime\prime\prime} \circ h_{n}(h_{n}^{\prime})^{3} + 3\gamma^{\prime\prime} \circ h_{n}h_{n}^{\prime}h_{n}^{\prime\prime}), v \right\rangle.$$

We will estimate the norms of the terms in the first line of the above equation one at a time, for each $n \leq n_{\star}$. First, using (5.21) with m = 0 and $c = \|\hat{\nu}_n''\|$ (that we have estimated in the first of the (5.37)), and (5.29) we have, for some uniform $A_1 > 0$

$$\| [(D_{\hat{\nu}_n} F^n)^{-1}]'' \gamma' \circ h_n h'_n \| \leq C_{\flat} \mu^{4n} \varsigma_n^2 C_{\mu,n} + A_1 c_{\flat} C_{\flat} \varsigma_n c_{n_\star}^{n_\star} \mu^{4n} (\lambda_n^-(\hat{\nu}_n(t)))^{-1} \| \gamma'' \circ h_n(t) \|$$

+ $C_{\flat} \varsigma_n \mu^{2n} C_{\mu,n_\star} \mu^{3n_\star} C_3.$

Next, notice that $(D_{\hat{\nu}_{n_{\star}}}F^{n_{\star}})^{-1}\gamma'' \notin \mathbf{C}_{c}$, hence by the second of (5.23) and subsequent, there is uniform $A_{2} > 0$ such that

(C.36)
$$\|[(D_{\hat{\nu}_n}F^n)^{-1}]'\gamma'' \circ h_n(h'_n)^2\| \le A_2 C_b^2 \mu^{3n} (\lambda_n^-(\hat{\nu}_n(t)))^{-1} \varsigma_n \|\gamma'' \circ h_n(t)\|.$$

⁷²The constant C_{\flat} is introduced just before (C.10).

It is convenient to write the third term as

$$[D_{\hat{\nu}_n}F^n)^{-1}]'\gamma' \circ h_n h_n'' = \frac{h_n''}{h_n'} [D_{\hat{\nu}_n}F^n)^{-1}]'\gamma' \circ h_n h_n'$$

= $\frac{h_n''}{h_n'} \left(\hat{\nu}_n'' - [(D_{\hat{\nu}_n}F^n)^{-1}]\gamma'' \circ h_n (h_n')^2 - \hat{\nu}_n' \frac{h_n''}{h_n'} \right).$

The last term vanishes when we multiplied by v; hence, by (C.30) and (C.32), we have⁷³

$$\begin{split} \left| \langle [D_{\hat{\nu}_n} F^n)^{-1}]' \gamma' \circ h_n h_n'', v \rangle \right| &\leq \left| \frac{h_n''}{h_n'} \right| \left\{ \| \hat{\nu}_n'' \| + \| [D_{\hat{\nu}_n} F^n)^{-1}] \gamma'' \circ h_n (h_n')^2 \| \right\} \\ &\leq c_{n_\star}^{n_\star} C_{\mathfrak{h}}^{\mathfrak{h}} \mu^{4n} (\lambda_n^- (\hat{\nu}_n(t)))^{-2} \| \gamma'' \circ h_n(t) \|^2 \\ &+ 2 c_{n_\star}^{n_\star} C_{\mu,n} C_{\mathfrak{h}}^2 C_3 \mu^{5n} (\lambda_n^- (\hat{\nu}_n(t)))^{-1} \| \gamma'' \circ h_n(t) \| + C_{\mu,n}^2 \mu^{6n} C_3^2 . \end{split}$$

For the two terms in the second line of (C.35), when the matrix hits γ'' or γ''' , we can use (C.30) for $n \leq n_{\star}$ and (5.29) with $\|\gamma'' \circ h_n(t)\|$ instead of c_{\star} . Collecting all the above estimates in (C.35) we finally have, recalling also (C.31),

$$\begin{aligned} (1+\chi_{c}^{2})^{-\frac{1}{2}} \| \hat{\nu}_{n}^{\prime\prime\prime} \| &\leq C_{\flat} \mu^{4n} \varsigma_{n}^{2} C_{\mu,n} + A_{1} c_{\flat} C_{\flat} \varsigma_{n} c_{n\star}^{n} \mu^{4n} (\lambda_{n}^{-}(\hat{\nu}_{n}(t)))^{-1} \| \gamma^{\prime\prime} \circ h_{n}(t) \| \\ &+ C_{\flat} \varsigma_{n} \mu^{2n} C_{\mu,n\star} \mu^{3n\star} C_{3} \\ &+ A_{2} C_{\flat}^{2} \mu^{3n} \varsigma_{n} (\lambda_{n}^{-}(\hat{\nu}_{n}(t)))^{-1} \| \gamma^{\prime\prime} \circ h_{n}(t) \| \\ &+ c_{n\star}^{n\star} C_{\flat}^{4} \mu^{4n} (\lambda_{n}^{-}(\hat{\nu}_{n}(t)))^{-2} \| \gamma^{\prime\prime} \circ h_{n}(t) \|^{2} \\ &+ 2 c_{n\star}^{n\star} C_{\mu,n} C_{\flat}^{2} C_{3} \mu^{5n} (\lambda_{n}^{-}(\hat{\nu}_{n}(t)))^{-1} \| \gamma^{\prime\prime} \circ h_{n}(t) \| + C_{\mu,n}^{2} \mu^{6n} C_{3}^{2} \\ &+ C_{\flat}^{3} \mu^{3n} (\lambda_{n}^{-}(\hat{\nu}_{n}(t)))^{-1} \| \gamma^{\prime\prime\prime} \circ h_{n}(t) \| \\ &+ 3 (\lambda_{n}^{-}(\hat{\nu}_{n}(t)))^{-1} C_{\flat}^{5} \mu^{3n} C_{\mu,n} \| \gamma^{\prime\prime} \circ h_{n}(t) \| (\| \gamma^{\prime\prime} \circ h_{n}(t) \| C_{F} + 1). \end{aligned}$$

Setting $C_4 = \{A_1c_{\flat} + A_2 + 3 + 2C_3\}^+ (1 + \chi_c^2)^{1/2}$, and recalling the second of (5.36), yields

$$\begin{aligned} \|\hat{\nu}_{n}^{\prime\prime\prime}\| &\leq a_{n_{\star}}^{n_{\star}} \mu^{3n} (\lambda_{n}^{-}(\hat{\nu}_{n}(t)))^{-1} \|\gamma^{\prime\prime\prime} \circ h_{n}(t)\| + 6c_{n_{\star}}^{n_{\star}} \mu^{4n} C_{\flat}^{5} (\lambda_{n}^{-}(\hat{\nu}_{n}(t)))^{-2} \varsigma_{n} \|\gamma^{\prime\prime} \circ h_{n}(t)\|^{2} C_{\mu,n} \\ &+ b_{n_{\star}}^{n_{\star}} (\lambda_{n}^{-}(\hat{\nu}_{n}))^{-1} \mu^{5n} \|\gamma^{\prime\prime} \circ h_{n}(t)\| + \overline{\mathbf{s}}_{n_{\star}}. \end{aligned}$$

Using the first of (5.37), we can write

$$\begin{split} \|\hat{\nu}_{kn_{\star}}^{\prime\prime\prime}\| &\leq a_{n_{\star}}^{n_{\star}} \mu^{3n_{\star}} (\lambda_{n_{\star}}^{-} (\hat{\nu}_{(k-1)n_{\star}}(t)))^{-1} \|\nu_{(k-1)n_{\star}}^{\prime\prime\prime} \circ h_{n_{\star}}^{*}(t)\| \\ &+ 6c_{n_{\star}}^{n_{\star}} \mu^{4n_{\star}} C_{\flat}^{5} (\lambda_{n_{\star}}^{-} (\hat{\nu}_{(k-1)n_{\star}}(t)))^{-2} \varsigma_{n_{\star}} C_{\mu,n_{\star}} \left[c_{\flat} c_{n_{\star}}^{(k-1)n_{\star}} \mu^{2(k-1)n_{\star}} \lambda_{(k-1)n_{\star}}^{-} (\gamma \circ h_{(k-1)n_{\star}}(t))^{-1} c_{\star} \right]^{2} \\ &+ b_{n_{\star}}^{n_{\star}} (\lambda_{n_{\star}}^{-} (\hat{\nu}_{(k-1)n_{\star}}))^{-1} \mu^{5n_{\star}} \left[c_{\flat} c_{n_{\star}}^{(k-1)n_{\star}} \mu^{2(k-1)n_{\star}} \lambda_{(k-1)n_{\star}}^{-} (\gamma \circ h_{(k-1)n_{\star}}(t))^{-1} c_{\star} \right] \\ &+ \overline{\mathbf{s}}_{n_{\star}} + 6c_{n_{\star}}^{n_{\star}} \mu^{4n_{\star}} C_{\flat}^{5} (\lambda_{n_{\star}}^{-} (\hat{\nu}_{(k-1)n_{\star}}(t)))^{-2} \varsigma_{n_{\star}} C_{\mu,n_{\star}}^{3} \mu^{6n_{\star}} C_{3}^{2} \\ &+ b_{n_{\star}}^{n_{\star}} (\lambda_{n_{\star}}^{-} (\hat{\nu}_{(k-1)n_{\star}}))^{-1} \mu^{5n_{\star}} C_{\mu,n_{\star}} \mu^{3n_{\star}} C_{3}. \end{split}$$

We can then proceed by induction as in (C.34), and since condition (5.1) implies both $a_{\hat{n}}^{\hat{n}}\mu^{3\hat{n}}\lambda_{-}^{-\hat{n}} \leq \frac{1}{2}$ and $\bar{c}_{n_{\star}}^{n_{\star}}\mu^{2n_{\star}}\lambda_{-}^{-n_{\star}} \leq \frac{1}{2}$, we have

$$\begin{aligned} \|\hat{\nu}_{kn_{\star}}^{\prime\prime\prime}\| \leq & c_{\flat}^{2} (1 + 6\varsigma_{n_{\star}} C_{\mu,n_{\star}}) c_{n_{\star}}^{kn_{\star}} \mu^{3kn_{\star}} (\lambda_{kn_{\star}}^{-} (\gamma \circ h_{kn_{\star}}(t)))^{-1} c_{\star}^{2} \\ &+ c_{\flat}^{2} b_{n_{\star}}^{n_{\star}} \mu^{2kn_{\star} + 2n_{\star}} \lambda_{kn_{\star}}^{-} (\gamma \circ h_{kn_{\star}}(t))^{-1} c_{\star} + \mathbf{s}_{n_{\star}}, \end{aligned}$$

which implies the second of (5.37).

⁷³Recall also the lower bound for $|h'_n|$ in (C.8).
C.4. Proof of Lemma 5.16. Let $\nu_n = \mathfrak{h}_n \circ \gamma$ for each $n \in \mathbb{N}$. Then, $C_{\sharp} \vartheta_{\gamma}(t) |\pi_1 \circ \nu'_{n_0}(t)| \ge |\pi_2 \circ \nu'_{n_0}(t)| \ge \vartheta_{\gamma}(t) |\pi_1 \circ \nu'_{n_0}(t)| > 0$, and we can reparametrize ν_n , $n \ge n_0$, by vertical length $\hat{\nu}_n(t) = \nu_n(h_n(t))$. Note that

(C.37)
$$C_{\sharp}\vartheta_{\gamma}(t)^{-1} \le \|\hat{\nu}_{n_0}'(t)\| \le C_{\sharp}\vartheta_{\gamma}(t)^{-1}.$$

If
$$n_0 = 0$$
, then $C_{\sharp}\vartheta_{\hat{\nu}_0}(t)^{-1} = C_{\sharp}\vartheta_{\gamma} \circ h_0(t)^{-1} \leq |h'_0(t)| \leq C_{\sharp}\vartheta_{\hat{\nu}_0}(t)^{-1}$ and (C.29) yields⁷⁴

$$|h_{n_0}''(t)| \leq \frac{\|\gamma'' \circ h_{n_0}(t)\| \|h_{n_0}'(t)\|^3}{|\langle e_1, \hat{\nu}_{n_0}'(t)\rangle|} \leq C_{\sharp} \Delta_{\gamma} \circ h_{n_0}(t) \vartheta_{\hat{\nu}_{n_0}}(t)^{-2} \\ \|\hat{\nu}_{n_0}''(t)\| \leq C_{\sharp} \Delta_{\gamma} \circ h_{n_0}(t) \vartheta_{\hat{\nu}_{n_0}}(t)^{-1}.$$

If $n_0 > 0$, then

(C.38)
$$C_{\sharp}\Lambda^{-n_{0}}\vartheta_{\hat{\nu}_{n_{0}}}(t)^{-1} \leq |h'_{n_{0}}(t)| \leq C_{\sharp}\Lambda^{n_{0}}\vartheta_{\hat{\nu}_{n_{0}}}(t)^{-1} \\ \|\nu''_{n_{0}}(t)\| \leq C_{\sharp}\Lambda^{2n_{0}}\|\gamma''(t)\|$$

and

(C.39)
$$\begin{aligned} |h_{n_0}''(t)| &\leq C_{\sharp} \Lambda^{3n_0} \Delta_{\gamma} \circ h_{n_0}(t) \vartheta_{\hat{\nu}_{n_0}}(t)^{-2} \\ \|\hat{\nu}_{n_0}''(t)\| &\leq C_{\sharp} \Lambda^{2n_0} \Delta_{\gamma} \circ h_{n_0}(t) \vartheta_{\hat{\nu}_{n_0}}(t)^{-1} \\ \|\hat{\nu}_{n_0}'''(t)\| &\leq C_{\sharp} \Lambda^{3n_0} \Delta_{\gamma}^2 \circ h_{n_0}(t) \vartheta_{\hat{\nu}_{n_0}}(t)^{-2}. \end{aligned}$$

Remark that

(C.40)

$$\|(D_{\hat{\nu}_m(t)}F^k)\hat{\nu}'_m(t)\| \le \sqrt{1+\chi_c^2}C_{\star}\lambda_k^+(\hat{\nu}_m(t))$$

Let $m_{\star}(t)$ be the largest integer for which $\hat{\nu}'_{m_{\star}}(t) \notin \mathbf{C}_{c}$, then, recalling (2.4) and setting $F^{m-n_{0}}\hat{\nu}_{m} = \hat{\nu}_{n_{0}} \circ \bar{h}_{m-n_{0}}$, we have

(C.41)

$$\begin{aligned} |\bar{h}'_{m-n_{0}}(t)| &= |\langle e_{2}, D_{\hat{\nu}_{m}(t)}F^{m-n_{0}}\hat{\nu}'_{m}(t)\rangle| \\
&\leq C_{\sharp}\mu^{m-m_{\star}}\lambda^{+}_{m_{\star}-n_{0}}(\hat{\nu}_{m_{\star}}(t))\vartheta_{\hat{\nu}_{n_{0}}}(\bar{h}_{m-n_{0}}(t)) \\
&|\bar{h}'_{m-n_{0}}(t)| \geq C_{\sharp}\mu^{m_{\star}-m}\lambda^{-}_{m_{\star}-n_{0}}(\hat{\nu}_{m_{\star}}(t))\vartheta_{\hat{\nu}_{n_{0}}}(\bar{h}_{m-n_{0}}(t)).\end{aligned}$$

Next, we want to use equation (C.29), with γ replaced by $\hat{\nu}_{n_0}$, n by $m - n_0$ and $\hat{\nu}_k$ by $\hat{\nu}_{k+n_0}$ and. Setting $\eta_m = DF_{\hat{\nu}_{m-n_0}}e_1$, $\hat{\eta}_m = \eta_m \|\eta_m\|^{-1}$, if we write $\nu''_0 = a\eta_m + be_2$, then $|a| \leq C_{\sharp} \|\nu''_0\|$ and $|b| \leq C_{\sharp} \chi_u \|\nu''_0\|$. Accordingly, setting $m_0 = m - n_0$, equations (C.39) and (C.41), yield,

$$\begin{aligned} \|\hat{\nu}_{m}''(t)\| &\leq C_{\sharp} \left\{ \frac{1}{\lambda_{m_{0}}^{-}(\hat{\nu}_{m_{0}}(t))} + \mu^{m_{0}}C_{F} \right\} \\ &\times \Lambda^{2n_{0}} \Delta_{\gamma} \circ h_{n_{0}}(t) \vartheta_{\hat{\nu}_{n_{0}}}(\bar{h}_{m_{0}}(t))\lambda_{m_{\star}-n_{0}}^{+}(\hat{\nu}_{m_{\star}}(t))^{2}\mu^{2(m-m_{\star})} \\ &+ \sum_{k=0}^{m_{0}-1} C_{\sharp} \left\{ 1 + \mu^{k} \lambda_{m_{\star}-n_{0}}^{+}(\hat{\nu}_{m}(t))C_{F} \right\} \lambda_{m_{\star}-n_{0}}^{+}(\hat{\nu}_{m}(t))^{-1} \\ &\times \mu^{2\{k,m-m_{\star}\}^{-}} \lambda_{\{0,k-m+m_{\star}\}^{+}}^{+}(\hat{\nu}_{m_{\star}}(t))^{2} \\ |\bar{h}_{m_{0}}''(t)| &\leq C_{\sharp} \left\{ \frac{1}{\lambda_{m_{0}}(\hat{\nu}_{m_{0}}(t))} + \mu^{m_{0}}C_{F} \right\} \\ &\times \Lambda^{2n_{0}} \Delta_{\gamma} \circ h_{n_{0}}(t) \vartheta_{\hat{\nu}_{n_{0}}}(\bar{h}_{m_{0}}(t))^{2} \lambda_{m_{\star}-n_{0}}^{+}(\hat{\nu}_{m_{\star}}(t))^{3} \mu^{3(m-m_{\star})} \\ &+ \sum_{k=0}^{m_{0}-1} C_{\sharp} \left\{ 1 + \mu^{k} \lambda_{m_{\star}-n_{0}}^{+}(\hat{\nu}_{m}(t))C_{F} \right\} \mu^{2\{k,m-m_{\star}\}^{-}} \\ &\times \lambda_{\{0,k-m+m_{\star}\}^{+}}^{+}(\hat{\nu}_{m_{\star}}(t))^{2} \vartheta_{\hat{\nu}_{n_{0}}}(\bar{h}_{m_{0}}(t)) \mu^{m-m_{\star}}. \end{aligned}$$

To continue we need the following

⁷⁴Note that (C.29) holds also if γ is not parametrized vertically.

Sublemma C.1. We have

(C.43)
$$C_{\sharp}\chi_{c}^{-1}\mu^{-m_{\star}+n_{0}} \leq \vartheta_{\hat{\nu}_{n_{0}}}(\bar{h}_{m_{0}}(t))\lambda_{m_{\star}-n_{0}}^{+}(\hat{\nu}_{m_{\star}}(t)) \leq C_{\sharp}\chi_{c}^{-1}\mu^{m_{\star}-n_{0}}, \quad \forall t \in \mathbb{T}.$$

Proof. Let w, ||w|| = 1, such that $DF^{m_{\star}}w = ||DF^{m_{\star}}w||e_2$. Also, let $v \in \mathbb{R}^2$, ||v|| = 1 such that $DF^{m_{\star}-n_0}v = \frac{||DF^{m_{\star}-n_0}v||}{(1+\vartheta_{\hat{\nu}n_0}^2)^{1/2}}(q(\hat{\nu}'_{n_0}), \vartheta_{\hat{\nu}n_0})$, where, for $z = (z_1, z_2) \in \mathbb{R}^2$, $q(z) = \operatorname{sign}(z_1) \cdot \operatorname{sign}(z_2)$. Note that $v \notin \mathbf{C}_c$, otherwise we would have $\hat{\nu}'_{m_{\star}} \in \mathbf{C}_c$, contrary to the hypothesis. We can then write $v = ae_1 + bw$. Note that $w \in (DF)^{-1}\mathbf{C}_c$, moreover by (2.4) it follows $|w_1| \leq \iota_{\star}\chi_c|w_2|$ while $C_{\sharp}\chi_c|v_2| \geq |v_1| \geq \chi_c|v_2|$. In addition, $v_2 = bw_2$ and

$$\begin{aligned} a| &\geq \chi_c |v_2| - |bw_1| \geq \chi_c (1 - \iota_\star) |b| |w_2| \geq \chi_c (1 - \iota_\star) (1 + \chi_c^2 \iota_\star^2)^{-\frac{1}{2}} |b| \\ a| &\leq C_{\sharp} \chi_c |v_2| + |bw_1| \leq C_{\sharp} \chi_c |b| |w_2| \leq C_{\sharp} \chi_c |b| \end{aligned}$$

which implies $C_{\sharp}\chi_c^{-1} \leq \frac{|b|}{|a|} \leq C_{\sharp}\chi_c^{-1}$. Finally, by equations (2.4) and (5.17), we can write

$$\begin{split} \vartheta_{\hat{\nu}_{n_{0}}} &= \frac{|\langle e_{2}, DF^{m_{\star}-n_{0}}v\rangle|}{|\langle e_{1}, DF^{m_{\star}-n_{0}}v\rangle|} \leq \frac{|b|\mu^{m_{\star}} + |a| |\langle e_{2}, DF^{m_{\star}-n_{0}}e_{1}\rangle|}{|a| |\langle e_{1}, DF^{m_{\star}-n_{0}}e_{1}\rangle|} \\ &\leq C_{\sharp}\frac{|b|}{|a|}\mu^{m_{\star}-n_{0}}(\lambda_{m_{\star}-n_{0}}^{+}\circ\hat{\nu}_{m_{\star}})^{-1} + \iota_{\star}\chi_{u} \leq C_{\sharp}\frac{|b|}{|a|}\mu^{m_{\star}-n_{0}}(\lambda_{m_{\star}-n_{0}}^{+}\circ\hat{\nu}_{m_{\star}})^{-1} + \iota_{\star}\vartheta_{\hat{\nu}_{n_{0}}} \\ \vartheta_{\hat{\nu}_{n_{0}}} \geq \frac{|b|\mu^{-m_{\star}+n_{0}} - |a| |\langle e_{2}, DF^{m_{\star}-n_{0}}e_{1}\rangle|}{|a| |\langle e_{1}, DF^{m_{\star}-n_{0}}e_{1}\rangle|} \geq C_{\sharp}\chi_{c}^{-1}\mu^{-m_{\star}+n_{0}}(\lambda_{m_{\star}-n_{0}}^{+}\circ\hat{\nu}_{m_{\star}})^{-1} - \vartheta_{\hat{\nu}_{n_{0}}} \end{split}$$

that is (C.43).

Note that equation (C.41) and Sublemma C.1 imply

(C.44)
$$|\bar{h}'_{m_0}| \le C_{\sharp} \mu^{m_0}$$

Thus, recalling (C.42) and (C.41), and the definition of M_{m,n_0} in (5.41),

(C.45)
$$\begin{aligned} \|\hat{\nu}_{m}''(t)\| &\leq C_{\sharp} \left[\Lambda^{2n_{0}} \mu^{2m} \Delta_{\gamma} \circ h_{n_{0}}(t) + C_{\mu,m} \mu^{3m} \vartheta_{\hat{\nu}_{n_{0}}}^{-1} \right] \left\{ 1 + \mu^{2m} \vartheta_{\hat{\nu}_{n_{0}}}^{-1} C_{F} \right\} \\ &\leq M_{m,n_{0}}(t) \\ |\bar{h}_{m_{0}}''(t)| \leq C_{\sharp} \left[\Lambda^{2n_{0}} \mu^{3m} \Delta_{\gamma} \circ h_{n_{0}}(t) + C_{\mu,m} \mu^{4m} \vartheta_{\hat{\nu}_{n_{0}}}^{-1} \right] \left\{ 1 + \mu^{2m} \vartheta_{\hat{\nu}_{n_{0}}}^{-1} C_{F} \right\} \\ &\leq \mu^{m} M_{m,n_{0}}(t). \end{aligned}$$

Our next task is to estimate $\hat{\nu}_{m}^{\prime\prime\prime}$. To this end we first estimate $\hat{\nu}_{m_{\star}}^{\prime\prime\prime}$, to do so we use (C.35) where $\hat{\nu}_{n}, \gamma, h_{n}$ are replaced by $\hat{\nu}_{m_{\star}}, \hat{\nu}_{n_{0}}$ and $\bar{h}_{m_{0}^{\star}}, m_{0}^{\star} = m_{\star} - n_{0}$, respectively. In this case $\hat{\nu}_{n_{0}}^{\prime} \notin \mathbf{C}_{c}$, and so is $\mathfrak{h}_{k}(\hat{\nu}_{n_{0}})^{\prime}$ for each $k < m_{\star}, \mathfrak{h}_{k} \in \mathfrak{H}^{k}$. We will estimate the terms in (C.35) one by one. We will use Proposition 5.9 for the first one and (5.23) with the equivalent of (5.26) for the second.⁷⁵ Recalling Sublemma C.1 we have the following estimates

$$\begin{split} \left\| \left[(D_{\hat{\nu}_{m_{\star}}} F^{m_{0}^{\star}})^{-1} \right]^{\prime\prime} \hat{\nu}_{n_{0}}^{\prime} \bar{h}_{m_{0}^{\star}}^{\prime} \right\| &\leq C_{\sharp} \mu^{m_{0}^{\star}} C_{\mu,m_{0}^{\star}} (1 + C_{F} \lambda_{m_{0}^{\star}}^{+}) \Big\{ \mu^{2m_{0}^{\star}} C_{\mu,m_{0}^{\star}} \lambda_{m_{0}^{\star}}^{+} + \| \hat{\nu}_{m_{\star}}^{\prime\prime} \| \Big\} \| \hat{\nu}_{n_{0}}^{\prime} \| \| \bar{h}_{m_{0}^{\star}}^{\prime} \| \\ &\leq C_{\mu,m_{0}^{\star}} \mu^{5m_{0}^{\star}} (1 + C_{F} \mu^{m_{0}^{\star}} \vartheta_{\hat{\nu}_{n_{0}}}^{-1})^{2} \left[C_{\mu,m_{0}^{\star}} \mu^{m_{0}^{\star}} + \Lambda^{2n_{0}} \Delta_{\gamma} \circ h_{n_{0}}(t) \vartheta_{\hat{\nu}_{n_{0}}} \right] \vartheta_{\hat{\nu}_{n_{0}}}^{-2} \\ & \left\| \left[(D_{\hat{\nu}_{m_{\star}}} F^{m_{0}^{\star}})^{-1} \right]^{\prime} \hat{\nu}_{n_{0}}^{\prime} \bar{h}_{m_{0}^{\star}}^{\prime\prime} \Big\| \leq C_{\sharp} \mu^{m_{\star}} \left(1 + \lambda_{m_{0}^{\star}}^{+} (\hat{\nu}_{m_{\star}}) C_{F} \right) \vartheta_{\hat{\nu}_{n_{0}}}^{-1} | \bar{h}_{m_{0}^{\star}}^{\prime\prime} | \\ & \leq \mu^{5m_{0}^{\star}} C_{\mu,m_{0}^{\star}} (1 + C_{F} \mu^{m_{0}^{\star}} \vartheta_{\hat{\nu}_{n_{0}}}^{-1})^{2} \left[C_{\mu,m_{0}^{\star}} \mu^{m_{0}^{\star}} + \Lambda^{2n_{0}} \Delta_{\gamma} \circ h_{n_{0}}(t) \vartheta_{\hat{\nu}_{n_{0}}} \right] \vartheta_{\hat{\nu}_{n_{0}}}^{-2}, \end{split}$$

⁷⁵We use Proposition 5.9, with $\|\hat{\nu}_{m_{\star}}^{\prime\prime}\|$ instead of \mathfrak{c} , $\hat{\nu}_{m_{\star}}$ for ν , n and m replaced by m_{0}^{\star} .

where in the second and last inequality we have used equations (C.44), (C.45), (C.37). Analogously, recalling (5.23), (5.26), (C.39), (C.41), (C.45) and Sublemma C.1 we have⁷⁶

$$\begin{split} \left\| \left[(D_{\hat{\nu}_{m_{\star}}} F^{m_{0}^{\star}})^{-1} \right]' \hat{\nu}_{n_{0}}'' (\bar{h}_{m_{0}^{\star}}')^{2} \right\| &\leq C_{\mu,m_{\star}} \mu^{m_{\star}} \left(1 + \lambda_{m_{0}^{\star}}^{+} (\hat{\nu}_{m_{\star}}) C_{F} \right) \| \hat{\nu}_{n_{0}}'' \| |\bar{h}_{m_{0}^{\star}}'|^{2}, \\ &\leq \mu^{3m_{\star}} C_{\mu,m_{\star}} (1 + C_{F} \mu^{m_{0}^{\star}} \vartheta_{\hat{\nu}_{n_{0}}}^{-1}) \Lambda^{2n_{0}} \Delta_{\gamma} \circ h_{n_{0}} \vartheta_{\hat{\nu}_{n_{0}}}^{-1} \\ \left\| \left[(D_{\hat{\nu}_{m_{\star}}} F^{m_{0}^{\star}})^{-1} \right] \hat{\nu}_{n_{0}}''' (\bar{h}_{m_{0}^{\star}}')^{3} \right\| \leq C_{\mu,m_{\star}} \mu^{m_{\star}} \left(1 + \lambda_{m_{0}^{\star}}^{+} (\hat{\nu}_{m_{\star}}) C_{F} \right) \lambda_{m_{0}^{\star}}^{-1} (\hat{\nu}_{m_{\star}})^{-1} \| \hat{\nu}_{n_{0}}''' \| |\bar{h}_{m_{0}^{\star}}'|^{3} \\ &\leq \mu^{5m_{\star}} (1 + C_{F} \mu^{m_{0}^{\star}} \vartheta_{\hat{\nu}_{n_{0}}}^{-1}) \Lambda^{3n_{0}} \Delta_{\gamma}^{2} \vartheta_{\hat{\nu}_{n_{0}}}^{-1} \\ \left\| \left[(D_{\hat{\nu}_{m_{\star}}} F^{m_{0}^{\star}})^{-1} \right] \hat{\nu}_{n_{0}}'' \bar{h}_{m_{0}^{\star}}' \bar{h}_{m_{0}^{\star}}'' \right\| \leq C_{\mu,m_{\star}} \mu^{m_{\star}} \left(1 + \lambda_{m_{0}^{\star}}^{+} (\hat{\nu}_{m_{\star}}) C_{F} \right) \lambda_{m_{0}^{\star}}^{-1} (\hat{\nu}_{m_{\star}})^{-1} \| \hat{\nu}_{n_{0}}'' \| |\bar{h}_{m_{0}^{\star}}' \| \bar{h}_{m_{0}^{\star}}' \| \\ &\leq C_{\mu,m_{\star}} \mu^{7m_{\star}} (1 + C_{F} \mu^{m_{0}^{\star}} \vartheta_{\hat{\nu}_{n_{0}}}^{-1})^{2} \Lambda^{2n_{0}} \Delta_{\gamma} \circ h_{n_{0}} \left[C_{\mu,m_{0}^{\star}} \mu^{m_{0}^{\star}} + \Lambda^{2n_{0}} \Delta_{\gamma} \circ h_{n_{0}}(t) \vartheta_{\hat{\nu}_{n_{0}}} \right] \vartheta_{\hat{\nu}_{n_{0}}}^{-1}. \end{split}$$

Using the above estimates in (C.35), and Sublemma C.1 again, we conclude

$$\begin{aligned} \|\hat{\nu}_{m_{\star}}^{\prime\prime\prime\prime}\| &\leq \mu^{8m_{\star}} C_{\mu,m_{\star}}^{2} (1 + C_{F} \mu^{m_{0}^{\star}} \vartheta_{\hat{\nu}_{n_{0}}}^{-1})^{2} \left[1 + \Lambda^{2n_{0}} \Delta_{\gamma} \circ h_{n_{0}} \vartheta_{\hat{\nu}_{n_{0}}} + \Lambda^{3n_{0}} (\Delta_{\gamma} \circ h_{n_{0}})^{2} \vartheta_{\hat{\nu}_{n_{0}}} \right] \vartheta_{\hat{\nu}_{n_{0}}}^{-2} \\ &= \overline{M}_{m_{\star},n_{0}}(t). \end{aligned}$$

Next, for each $\overline{m} > m_{\star}$ let $F^{\overline{m}-m_{\star}}\hat{\nu}_{\overline{m}} = \hat{\nu}_{m_{\star}} \circ \overline{h}_{\overline{m}-m_{\star}}$. Then,

(C.46)
$$\begin{aligned} |\bar{h}'_{\overline{m}-m_{\star}}(t)| &= |\langle e_{2}, D_{\hat{\nu}\overline{m}(t)}F^{\overline{m}-m_{\star}}\hat{\nu}'_{\overline{m}}(t)\rangle| \leq C_{\sharp}\chi_{c}^{-1}\mu^{\overline{m}-m_{\star}}\\ |\bar{h}'_{\overline{m}-m_{\star}}(t)| \geq C_{\sharp}\chi_{c}^{-1}\mu^{-\overline{m}+m_{\star}}. \end{aligned}$$

We can now apply Lemmata 5.10 and 5.14, in particular (5.37), to $\hat{\nu}_{\overline{m}}$ and $\hat{h}_{\overline{m}-m_{\star}}$ with γ replaced by $\hat{\nu}_{m_{\star}}$, n by $\overline{m} - m_{\star}$, and c_{\star} and c_{\star}^2 replaced by $M_{m,n_0}(t)$ and $\overline{M}_{m,n_0}(t)$ respectively. We thus obtain

(C.47)
$$\begin{aligned} \|\hat{\nu}_{\overline{m}}^{\prime\prime}\| &\leq c_{\flat}c_{n_{\star}}^{\overline{m}-m_{\star}}\mu^{2(\overline{m}-m_{\star})}\lambda_{\overline{m}-m_{\star}}^{+}(\hat{\nu}_{m}\circ\bar{h}_{\overline{m}-m_{\star}})^{-1}M_{m,n_{0}} + C_{\mu,n_{\star}}\mu^{3n_{\star}}C_{3}, \\ \|\hat{\nu}_{\overline{m}}^{\prime\prime\prime}\| &\leq c_{\flat}^{2}(1+6\varsigma_{n_{\star}}C_{\mu,n_{\star}})c_{n_{\star}}^{\overline{m}-m_{\star}}\mu^{3(\overline{m}-m_{\star})}(\lambda_{\overline{m}-m_{\star}}^{-}(\gamma\circ h_{\overline{m}-m_{\star}}(t)))^{-1}\overline{M}_{m,n_{0}} \\ &+ c_{\flat}^{2}b_{n_{\star}}^{n_{\star}}\mu^{2(\overline{m}-m_{\star})+2n_{\star}}\lambda_{\overline{m}-m_{\star}}^{-}(\gamma\circ h_{\overline{m}-m_{\star}}(t))^{-1}M_{m,n_{0}} + \mathbf{s}_{n_{\star}}, \\ &|\bar{h}_{\overline{m}-m_{\star}}^{\prime\prime}| \leq C_{\sharp}(M_{m,n_{0}}C_{F}+1)\mu^{2(\overline{m}-m_{\star})}C_{\mu,\overline{m}-m_{\star}}. \end{aligned}$$

We are ready to conclude. Recalling Corollary 5.11, the first two of the above equations plus condition (5.40) give $\hat{\nu}_{\overline{m}} \in \Gamma_3(\mathfrak{c})$. Next we set $m_1 = \overline{m} - m$. If $F^{\overline{m}}\hat{\nu}_{\overline{m}} = \gamma \circ h_{\overline{m}}$, by definition we have

(C.48)
$$h_{\overline{m}} = h_{n_0} \circ \bar{h}_{m_0} \circ \bar{h}_{m_1}.$$

Hence, differentiating (C.48) and recalling (C.44), (C.46) and (C.38) we have the first of (5.42). Taking two derivatives of (C.48) and using the second lines of (C.39), (C.45) and the third of (C.47), we have⁷⁷

$$\begin{split} |h_{\overline{m}}''| &\leq |h_{n_0}'' \circ \bar{h}_{m_0} \circ \bar{h}_{m_1} \cdot \left[\bar{h}_{m_0}' \circ h_{m_1} \cdot \bar{h}_{m_1}'\right]^2 | \\ &+ |h_{n_0}' \circ \bar{h}_{m_0} \circ \bar{h}_{m_1} \left[h_{m_0}'' \circ \bar{h}_{m_1} \cdot (\bar{h}_{m_1}')^2 + \bar{h}_{m_1}'' \cdot \bar{h}_{m_0}' \circ \bar{h}_{m_1}\right] | \\ &\leq C_{\sharp} \mu^{3\overline{m}} \vartheta_{\hat{\nu}_{n_0}}^{-1} \Lambda^{3n_0} \left\{ \Delta_{\gamma} \vartheta_{\hat{\nu}_{n_0}}^{-1} + M_{m,n_0} (C_F C_{\mu,\overline{m}} + 1) + C_{\mu,\overline{m}} \right\}, \end{split}$$

form which the second of (5.42) follows and the Lemma is proven.

⁷⁶We write $\hat{\nu}_{n_0}'' = av + be_2$ where $v = DF^{m_0^*}e_1 ||DF^{m_0^*}e_1||^{-1}$, thus $|a| \le C_{\sharp} ||\hat{\nu}_{n_0}''||$ and $|b| \le C_{\sharp} \chi_u ||\hat{\nu}_{n_0}''||$ and then we use (5.25). Also, we treat $\hat{\nu}_{n_0}'''$ in the same manner. ⁷⁷Here we drop the dependence on t to ease notations.

Appendix D. Proof of Lemma 6.6

This appendix is devoted to the proof of Lemma 6.6.

As before we use the notation $F^k \hat{\nu}_k = \gamma \circ h_k$, $F^k \nu_k = \gamma$. As the computation is local it suffices to consider $p_n \in \hat{\nu}_n$ and $p_0 \in \gamma$ such that $F^n(p_n) = p_0$. Let $p_k = F^{n-k}p_n$. To ease notation we use a translation to reparaetrize the curves so that $\nu_k(0) = \hat{\nu}_k(0) = p_k$, note that $h_k(0) = 0$. Before discussing the splitting of the vector field we need some notations and few estimates.

It is convenient to perform the changes of variables $\phi_k^{-1}(x,y) = (x,0) + \hat{\nu}_k(y)$ and set

$$\widetilde{F}^k = \phi_0 \circ F^k \circ \phi_k^{-1} ; \quad \widetilde{F}_k = \phi_{k-1} \circ F \circ \phi_k^{-1}$$

Note that $\widetilde{F}^k = \widetilde{F}_k \circ \cdots \circ \widetilde{F}_1$ and $\widetilde{F}^n(0, y) = \phi_0 \circ F^n(\hat{\nu}_n(y)) = \phi_0(\gamma \circ h_n(y)) = (0, h_n(y))$, this implies that

$$D_{(0,y)}\tilde{F}^n = \begin{pmatrix} a^n(y) & 0\\ c^n(y) & d^n(y) \end{pmatrix}; \quad D_{(0,y)}\tilde{F}_k = \begin{pmatrix} a_k(y) & 0\\ c_k(y) & d_k(y) \end{pmatrix}; \quad D\phi_k^{-1} = \begin{pmatrix} 1 & (\hat{\nu}'_k)_1\\ 0 & 1 \end{pmatrix}$$

with $d^n(y) = h'_n(y)$ and $d_k(y) = h^*_k(y)$. Thus, we have the estimates on the \mathcal{C}^{ρ} norms of d^k by Lemma 5.10, also the changes of coordinates ϕ_k have uniformly bounded \mathcal{C}^{ρ} norms. From the above we easily get the formulae:

(D.1)
$$a^{k+1}(y) = a^k(y)a_{k+1}(h_k(y))$$

(D.2)
$$d^{k+1}(y) = d_{k+1}(h_k(y))d^k(y)$$

1.

(D.3)
$$c^{k}(y) = \sum_{j=1}^{\kappa} d_{k}(h_{k-1}(y)) \cdots d_{j+1}(h_{j}(y))c_{j}(h_{j-1}(y))a_{j-1}(h_{j-2}(y)) \cdots a_{1}(y).$$

Moreover,

$$DF^{k} = \begin{pmatrix} a^{k} + (\hat{\nu}'_{0})_{1}c^{k} & (\hat{\nu}'_{0})_{1}d^{k} - (\hat{\nu}'_{k})_{1} \begin{bmatrix} a^{k} + (\hat{\nu}'_{0})_{1}c^{k} \end{bmatrix} \\ c^{k} & d^{k} - (\hat{\nu}'_{k})_{1}c^{k} \end{pmatrix}$$

which, setting $y_k = h_k(y)$, yields the alternative representations and estimates

(D.4)

$$c_{k}(y_{k-1}) = \langle e_{2}, D_{(0,y_{k-1})}Fe_{1} \rangle$$

$$a_{k}(y_{k-1}) = \langle e_{1}, D_{(0,y_{k-1})}Fe_{1} \rangle - \nu_{k-1}'(y_{k-1})_{1} \langle e_{2}, D_{(0,y_{k-1})}Fe_{1} \rangle$$

$$|c^{k}(y)| = |\langle e_{2}, D_{(0,y)}F^{k}e_{1} \rangle| \leq \lambda_{k}^{+} \chi_{u}$$

$$\frac{\lambda_{k}^{-}}{\sqrt{1+\chi_{u}^{2}}} - \chi_{c}\chi_{u}\lambda_{k}^{+} \leq |a^{k}(y)| \leq \lambda_{k}^{+} + \chi_{c}\chi_{u}\lambda_{k}^{+}$$

Also, for further use,

(D.5)
$$\left(D\tilde{F}^k \right)^{-1} = \begin{pmatrix} a^k(y)^{-1} & 0\\ -d^k(y)^{-1}a^k(y)^{-1}c^k(y) & d^k(y)^{-1} \end{pmatrix} .$$

We are now ready to describe the splitting of the vector field. We do it in the new coordinates. Consider the subspace $E_n(y) = \{(\eta, u^n(y)\eta)\}_{\eta \in \mathbb{R}}$, where $u_n(y) = a^n(y)^{-1}c^n(y)$, which is a \mathcal{C}^r approximation of the unstable direction. Given a vector $v \in \mathbb{R}^2$ let us call $\tilde{v} = D\phi_0 v$ the vector in the new coordinates. Next, we decompose a vector \tilde{v} as

$$\tilde{v} = (1, u_n \circ \mathfrak{h}_n)\tilde{v}_1 + (\tilde{v}_2 - \tilde{v}_1 u_n \circ \mathfrak{h}_n)e_2$$

where $\mathfrak{h}_n \circ \widetilde{F}^n(0, y) = (0, y)$. Thus, setting $V(t) = v_1(\gamma(t)) - \gamma'(t)_1 v_2(\gamma(t))$, we have the decomposition (6.26), restricted to γ , with

(D.6)
$$v^{u}(\gamma(t)) = V(t)(1 + \gamma'(t)_{1}u_{n} \circ \mathfrak{h}_{n}(0, t), u_{n} \circ \mathfrak{h}_{n}(0, t))$$
$$v^{c}(\gamma(t)) = [v_{2}(\gamma(t)) - u_{n} \circ \mathfrak{h}_{n}(0, t)V(t)]\gamma'(t),$$

which gives, in particular, $v^c(\gamma(t)) = g(t)\gamma'(t)$ with $g(t) = v_2(\gamma(t)) - u_n \circ \mathfrak{h}_n(0,t)V(t)$. To extend the above decomposition in a neighborhood of γ we will proceed as in [32, Lemma 6.5].⁷⁸ First, we compute the derivatives along the curve, to this end note that in the new coordinates $t = y_n$. Differentiating (D.1) we have

(D.7)
$$\partial_y a^k(y)^{-1} = \left[\partial_y a^{k-1}(y)^{-1}\right] a_k(y_{k-1})^{-1} + a^{k-1}(y)^{-1} \partial_{y_{k-1}} a_k(y_{k-1})^{-1} \tilde{d}^{k-1},$$

and, by (D.4) and and Lemma 5.10,

$$\begin{aligned} |\partial_{y_{k-1}}a_k(y_{k-1})| &\leq C_{\sharp}(1+\|\nu_{k-1}''\|) \leq C_{\sharp}(1+\varepsilon) \\ \|\partial_y a_k\|_{\mathcal{C}^{\rho}} &\leq C_{\sharp}\|\nu_{k-1}\|_{\mathcal{C}^{\rho+1}} \leq C_{\sharp}\varepsilon^{\rho!}. \end{aligned}$$

Next, using (D.7), we can prove by induction that $||(a^n)^{-1}||_{\mathcal{C}^{\rho}} \leq C_{\sharp} \lambda_{-}^{-n} \mathfrak{C}^{\rho\rho!} C_{\mu,n}^{\rho_{\alpha_{\rho}}} \mu^{\rho\rho!n}$:⁷⁹

(D.8)
$$\begin{aligned} \|[a^{n}]^{-1}\|_{\mathcal{C}^{\rho}} &\leq C_{\sharp}\lambda_{-}^{-n} + \lambda^{-1}\|[a^{n-1}]^{-1}\|_{\mathcal{C}^{\rho}} + C_{\sharp}\|[a^{n-1}]^{-1}\|_{\mathcal{C}^{\rho-1}}\mathbb{C}^{\rho_{j}}C^{a_{\rho}}_{\mu,n-1}\mu^{\rho!(n-1)} \\ &\leq C_{\sharp}\lambda_{-}^{-n} + \mathbb{C}^{\rho!}C_{\sharp}\sum_{j=0}^{n-1}\lambda_{-}^{j-n}\|[a^{j}]^{-1}\|_{\mathcal{C}^{\rho-1}}C^{a_{\rho}}_{\mu,j}\mu^{\rho!j} \\ &\leq C_{\sharp}\lambda_{-}^{-n}\mathbb{C}^{\rho\rho!}C^{\rho a_{\rho}}_{\mu,n}\mu^{\rho\rho!n}. \end{aligned}$$

To compute $||(d^n)^{-1}||_{\mathcal{C}^{\rho}}$ we can use formula (5.5) and recall (5.29) and (C.8):

(D.9) $\|(d^n)^{-1}\|_{\mathcal{C}^{\rho}} = \|(h'_n)^{-1}\|_{\mathcal{C}^{\rho}} \le C_{\sharp} \mathfrak{C}^{\rho!} \mu^{(\rho+1)n} C^{a_{\rho+1}}_{\mu,n} \mu^{(\rho+1)!n} = C_{\sharp} \mathfrak{C}^{\rho!} C^{a_{\rho+1}}_{\mu,n} \mu^{(\rho+1)(\rho!+1)n}.$ Next, by (D.1), (D.2) and (D.3) we have

$$[a^{n}(y)]^{-1}c^{n}(y) = \sum_{j=1}^{n} d_{n}(h_{n-1}(y)) \cdots d_{j+1}(h_{j}(y))c_{j}(h_{j-1}(y))[a_{n}(h_{n-1}(y)) \cdots a_{j}(h_{j-1}(y))]^{-1},$$

$$[d^{n}(y)a^{n}(y)]^{-1}c^{n}(y) = \sum_{j=1}^{n} [d_{j-1}(h_{j-2}(y)) \cdots d_{1}(y))]^{-1}c_{j}(h_{j-1}(y))[a_{n}(h_{n-1}(y)) \cdots a_{j}(h_{j-1}(y))]^{-1}.$$

Hence, by (D.8), (D.9) and the first of (D.4), we obtain, using (B.1),

(D.10)
$$\begin{aligned} \|[d^{n}a^{n}]^{-1}c^{n}\|_{\mathcal{C}^{\rho}} &\leq C_{\sharp} \mathbb{C}^{\rho\rho!} C^{2\rho a_{\rho}+1}_{\mu,n} \mu^{(\rho+1)(\rho!+1)+\rho!} \\ \|[a^{n}]^{-1}c^{n}\|_{\mathcal{C}^{\rho}} &\leq C_{\sharp} \mathbb{C}^{\rho!} C^{a_{\rho}}_{\mu,n} \mu^{\rho!n}. \end{aligned}$$

We are ready to conclude. Since

$$(D_{\hat{\nu}_n(y)}F^n)^{-1} = D_{(0,y)}\phi_n^{-1}(D_{(0,y)}\widetilde{F}^n)^{-1}D_{\gamma \circ h_n(y)}\phi_0,$$

by (D.6) and (D.5) it follows

$$(D_{\hat{\nu}_n(y)}F^n)^{-1}v^u(\gamma \circ h_n(y)) = V(h_n(y)) \left(a^n(y)^{-1}, 0\right), (D_{\hat{\nu}_n(y)}F^n)^{-1}v^c(\gamma \circ h_n(y)) = d^n(y)^{-1} \cdot [v_2 - u_nv_1] \circ \gamma(h_n(y)) \left((\hat{\nu}'_n)_1(y), 1\right).$$

Recalling that $u_n(y) = a^n(y)^{-1}c^n(y)$, by (D.8), (D.9), (D.10), and since $\gamma \in \Gamma(c)$ and $||v||_{\mathcal{C}^r} \leq 1$, we have the result for the vector field along the curve. Finally, we extend v^u to a neighborhood of γ . It turns out the be more convenient to define first the extension

$$w(x,y) = F^n v^u(\hat{\nu}_n(y))$$

then $\hat{v}^u = \mathfrak{h}_n^* w$ and $F^{n^*} \hat{v}^u = w$. By these definitions it follows

$$\|F^{n^*}\hat{v}^u\|_{\mathcal{C}^{\rho}(N(\nu))} = \|F^{n^*}v^u\|_{\mathcal{C}^{\rho}_{\nu}} \le \lambda_{-}^{-n} \mathfrak{C}^{\rho!\rho} C^{\rho a_{\rho}}_{\mu,n} \mu^{\rho\rho!n} \\ \|\hat{v}^u\|_{\mathcal{C}^{\rho}(M'(\gamma))} \le C_n.$$

The definition of \hat{v}^c and relative estimates are analogous.

⁷⁸In the mentioned paper the authors need more regularity for the extended vector field. Here it is enough to obtain a vector field which is C^{ρ} .

 $^{^{79}\}text{Here}~a_{\rho}$ is the one given by Lemma 5.10.

QUANTITATIVE STATISTICAL PROPERTIES

Appendix E. The space \mathcal{H}^s

Let $u \in \mathcal{C}^{\infty}(\mathbb{T}^2)$. The Fourier Transform of u and its inverse are

(E.1)
$$\mathcal{F}u(\xi) = \int_{\mathbb{T}^2} e^{-i2\pi x\xi} u(x) dx, \quad \xi \in \mathbb{Z}^2,$$

(E.2)
$$u(x) = \sum_{\xi \in \mathbb{Z}^2} \mathcal{F}u(\xi) e^{i2\pi x\xi}, \quad x \in \mathbb{T}^2.$$

Then \mathcal{H}^s is the completion of $\mathcal{C}^{\infty}(\mathbb{T}^2)$ with respect to the inner product

(E.3)
$$\langle u, v \rangle_s = \sum_{\xi \in \mathbb{Z}^2} \langle \xi \rangle^{2s} \mathcal{F}u(\xi) \overline{\mathcal{F}v(\xi)}, \quad \langle \xi \rangle := \sqrt{1 + \|\xi\|^2}.$$

Notice that, arguing like in the proof of formula [38, (7.9.2)], we have

(E.4)
$$\langle u, v \rangle_s = \sum_{\gamma+\beta \le s} C_{\gamma,\beta} \langle \partial_{x_1}^{\gamma} \partial_{x_2}^{\beta} u, \partial_{x_1}^{\gamma} \partial_{x_2}^{\beta} v \rangle_0$$

Hence, there are constants $C, C_{\gamma,\beta} > 0$ such that

(E.5)
$$C^{-1} \sum_{\gamma+\beta=s} C_{\gamma,\beta} \|\partial_{x_1}^{\gamma} \partial_{x_2}^{\beta} u\|_{L^2}^2 \le \|u\|_{\mathcal{H}^s}^2 \le C \sum_{\gamma+\beta=s} C_{\gamma,\beta} \|\partial_{x_1}^{\gamma} \partial_{x_2}^{\beta} u\|_{L^2}^2.$$

Lemma E.1. For every $\varsigma \in (0,1)$ and $1 \leq s < r$ there exists constants C_s such that

$$\|u\|_{\mathcal{H}^{s-1}}^2 \leq \varsigma \|u\|_{\mathcal{H}^s}^2 + \frac{C_s}{\varsigma} \|u\|_{L^1}^2, \quad u \in \mathcal{C}^r(\mathbb{T}^2).$$

Proof. By definition of the norm we have, for all $\tau \in (1,2)$,

(E.6)
$$||u||_{\mathcal{H}^{s-1}}^2 = \sum_{\xi \in \mathbb{Z}^2} |\mathcal{F}u(\xi)|^2 \langle \xi \rangle^{2(s-1)} = \sum_{\xi \in \mathbb{Z}^2} |\mathcal{F}u(\xi)|^2 \langle \xi \rangle^{2s-2+\tau} \langle \xi \rangle^{-\tau}$$

By Young inequality $ab \leq \frac{\varsigma a^p}{p} + \frac{\varsigma^{-\frac{p}{q}} b^q}{p}$, for every $\varsigma > 0$ and $\frac{1}{p} + \frac{1}{q} = 1$. We apply this with $a = \langle \xi \rangle^{2s-2+\tau}$, $b = \langle \xi \rangle^{-\tau}$ and $p = \frac{2s}{(2s-2+\tau)}$, $q = \frac{2s}{2-\tau}$ to obtain:

$$\langle \xi \rangle^{2s-2+\tau} \langle \xi \rangle^{-\tau} \le \left(1 - \frac{2-\tau}{2s}\right) \varsigma \langle \xi \rangle^{2s} + \varsigma^{-1} \frac{(2-\tau) \langle \xi \rangle^{-\frac{2s\tau}{2-\tau}}}{2s}$$

Using this fact in (E.6) and recalling that $\|\mathcal{F}u\|_{\infty} \leq C \|u\|_{L^1}$, we get

$$\|u\|_{\mathcal{H}^{s-1}}^2 \leq \varsigma \sum_{\xi \in \mathbb{Z}^2} |\mathcal{F}u(\xi)|^2 \langle \xi \rangle^{2s} + \frac{C_s}{\varsigma} \|\mathcal{F}u\|_{\infty}^2 \leq \varsigma \|u\|_{\mathcal{H}^s}^2 + \frac{C_s}{\varsigma} \|u\|_{L^1}^2.$$

APPENDIX F. VECTOR FIELD REGULARITY

This appendix is devoted to proving the following regularity results on the iteration of a vector field. Note that the hypotheses (F.1) are implied by assumption (H4).

Lemma F.1. Let $\varepsilon_0 \in (0, 1]$, $A \in [0, 1/2]$, B > 0 and $u, u' \in \mathcal{C}^1(\mathbb{T}^2, \mathbb{R})$ such that $||u||_{\infty}, ||u'||_{\infty} \leq A\varepsilon_0^{-1}$ and $||\nabla u||_{\infty}, ||\nabla u'||_{\infty} \leq B\varepsilon_0^{-1}$. Consider a family of vertically partially hyperbolic map $F_{\varepsilon}, \varepsilon \leq \varepsilon_0$ such that

(F.1)
$$\begin{aligned} \left\| \frac{\partial_{\theta} f(p)}{\partial_{x} f(p)} \right\|_{\infty} &\leq 1\\ \partial_{x} f(p) \left[1 - A \left\| \frac{\partial_{\theta} f(p)}{\partial_{x} f(p)} \right\|_{\infty} \right] &\geq 2(1 + \varepsilon_{0} \| \partial_{x} \omega \|_{\infty}). \end{aligned}$$

For each $\mathfrak{h} \in \mathfrak{H}^{\infty}$ and $k \leq n \in \mathbb{N}$, we define the sequence of functions⁸⁰

$$\bar{u}_0(p,\varepsilon) = u(\mathfrak{h}_n(p))$$
$$\bar{u}_k(p,\varepsilon) = \pi_2 \circ \Phi_{\varepsilon}^k(\mathfrak{h}_n(p), u(\mathfrak{h}_n(p))).$$

and similarly for \bar{u}'_k . Then, for each $p, p' \in \mathbb{T}^2$ and $\varepsilon, \varepsilon' < \varepsilon_0$,

(F.2)

$$\begin{aligned} & |\bar{u}_{n}(p,\varepsilon) - \bar{u}_{n}'(p',\varepsilon')| \leq C_{\sharp} e^{4A} \mu^{3n} \bigg\{ \lambda_{n}^{+}(\mathfrak{h}_{n}(p))^{-1} ||u - u'||_{\infty} \\ & + (||\omega||_{\mathcal{C}^{2}} + \mu^{2n} \lambda_{n}^{+}(\mathfrak{h}_{n}(p))^{-1} C_{\mu,n}|u'|) ||p - p'|| + \big[1 + \lambda_{n}^{+}(\mathfrak{h}_{n}(p))^{-1} |u'|^{2} \big] |\varepsilon - \varepsilon'| \bigg\}. \end{aligned}$$

Proof. Let $p_k(p,\varepsilon) = \mathfrak{h}_k(p)$, for $\mathfrak{h} \in \mathcal{H}^{\infty}$, $p \in \mathbb{T}^2$. By (5.21) (or see [19] for details) we have

(F.3)
$$\|\partial_p p_k\| \le \|(D_{\mathfrak{h}_k(p)} F_{\varepsilon}^k)^{-1}\| \le C_{\sharp} \mu^k \le C_{\sharp} e^{c_{\sharp} \varepsilon k}.$$

For each u > 0 and for $k \le n$ let

(F.4)
$$\lambda(p,\varepsilon) = \frac{|\partial_x f(p)|}{1+\varepsilon \left(\|\partial_\theta \omega\|_\infty + \|\partial_x \omega\|_\infty\right)} \ge |\partial_x f(p)|\mu^{-1}$$
$$u_k(p,\varepsilon,u) = \Xi_\varepsilon (p_{n-k+1}(p,\varepsilon), u_{k-1}(p,\varepsilon,u)),$$

where in the first line we have used equation (A.7), which shows that this bound does not depend on the hypotheses of Lemma 2.6, and Ξ_{ε} is defined in (10.3). Note that $\bar{u}_n(p,\varepsilon) = u_n(p,\varepsilon,u_0(p,\varepsilon))$. Using (10.3) and (10.5) a direct computation yields, for $|u| \leq A\varepsilon_0^{-1}$,

(F.5)

$$\begin{aligned} |\Xi_{\varepsilon}(p,u)| &\leq \frac{|u| (1+\varepsilon \|\partial_{\theta}\omega\|_{\infty})}{|\partial_{x}f(p)| \left[1-\varepsilon |u| \|\frac{\partial_{\theta}f(p)}{\partial_{x}f(p)}\|_{\infty}\right]} + \frac{\|\partial_{x}\omega\|_{\infty}}{\partial_{x}f(p) \left[1-A\|\frac{\partial_{\theta}f(p)}{\partial_{x}f(p)}\|_{\infty}\right]} \\ &\leq \frac{1}{|\partial_{x}f(p)|} \mu e^{2\varepsilon_{0}|u|} |u| + \frac{1}{2} \|\partial_{x}\omega\|_{\infty} \\ |\partial_{u}\Xi_{\varepsilon}(p,u)| &\leq \frac{1}{\lambda(p,\varepsilon)} \left[1-\varepsilon \left|u\frac{\partial_{\theta}f(p)}{\partial_{x}f(p)}\right|\right]^{-2} \\ &\|\partial_{p}\Xi_{\varepsilon}(p,u)\| \leq C_{\sharp}(\|\omega\|_{\mathcal{C}^{2}} + |u|) \\ &\|\partial_{\varepsilon}\Xi_{\varepsilon}(p,u)\| \leq C_{\sharp}(1+|u|)|u|. \end{aligned}$$

The first line of the (F.5) and the second of (F.1) imply

$$|u_k(p,\varepsilon,u)| \le 2^{-k} |u| + ||\partial_x \omega||_{\infty}.$$

We can get a sharper bound defining

$$\begin{split} \Lambda_{k,j}(p) &:= \prod_{i=k+1}^{j} \lambda(p_i,\varepsilon) ; \quad \overline{\Lambda}_{k,j}(p) := \prod_{i=k+1}^{j} |\partial_x f(p_i)| \\ \Delta &:= \|\partial_x \omega\|_{\infty} \left\| \frac{\partial_{\theta} f(p)}{\partial_x f(p)} \right\|_{\infty}, \end{split}$$

then

(F.6)
$$|u_k(p,\varepsilon,u)| \le \Lambda_{n-k,n}(p)^{-1}|u| + ||\partial_x \omega||_{\infty}$$

⁸⁰See (10.8) for the definition of Φ_{ε}^{n} .

Moreover, setting $u_j = u_j(p, u, \varepsilon)$, $u'_j = u_j(p', u', \varepsilon')$, with $|u|, |u'| \leq \frac{A}{\varepsilon_0}$, and recalling (F.3), (F.4), (F.6), we have

$$|u_{k+1}(p,\varepsilon,u) - u_{k+1}(p',\varepsilon',u')| = |\Xi_{\varepsilon}(p_{n-k},u_k) - \Xi_{\varepsilon'}(p'_{n-k},u'_k)| \\ \leq C_{\sharp}(||\omega||_{\mathcal{C}^2} + |u'_k|)||p_{n-k} - p'_{n-k}|| + C_{\sharp}(1 + |u'_k|)|u'_k||\varepsilon - \varepsilon'| \\ + \Lambda_{n-k-1,n-k}(p)^{-1}e^{2^{-k+1}A+2\varepsilon_0\Delta}|u_k - u'_k| \\ \leq C_{\sharp}(||\omega||_{\mathcal{C}^2} + \overline{\Lambda}_{n-k,n}(p')^{-1}\mu^k|u'|)[\mu^{n-k}||p - p'|| \\ + (1 + \overline{\Lambda}_{n-k,n}(p')^{-1}\mu^k|u'|)|\varepsilon - \varepsilon'|] + \Lambda_{n-k-1,n-k}(p)e^{2^{-k+1}A+2\varepsilon_0\Delta}|u_k - u'_k|.$$

We can then iterate the above equation and obtain

$$\begin{split} u_{n}(p,\varepsilon,u) &- u_{n}(p',\varepsilon',u')| = \overline{\Lambda}_{0,n}(p)^{-1}\mu^{n}e^{4A+2n\varepsilon_{0}\Delta}|u-u'| \\ &+ C_{\sharp}\sum_{k=0}^{n-1}\overline{\Lambda}_{0,n-k}(p)^{-1}\mu^{n-k}e^{4A+2\varepsilon_{0}(n-k)\Delta}(||\omega||_{\mathcal{C}^{2}} + \overline{\Lambda}_{n-k,n}(p')^{-1}\mu^{k}|u'|)\mu^{n-k}||p-p'|| \\ &+ C_{\sharp}\sum_{k=0}^{n-1}\overline{\Lambda}_{0,n-k}(p)^{-1}\mu^{n-k}e^{4A+2\varepsilon_{0}(n-k)\Delta}(1+\overline{\Lambda}_{n-k,n}(p')^{-2}\mu^{k}|u'|^{2})|\varepsilon-\varepsilon'|. \end{split}$$

In addition equations (10.2) and (5.17) imply

$$\begin{split} \Lambda_{j,n}(p) &\geq C_{\sharp} \lambda_{n-j}^{+}(p_{n}) \\ |\partial_{p} \overline{\Lambda}_{j,n}(p)| &\leq \sum_{j=l+1}^{n} \left| \overline{\Lambda}_{l,n}(p) \frac{\partial_{x}^{2} f(p_{l})}{[\partial_{x} f(p_{l})]^{2}} \overline{\Lambda}_{j,l-1}(p) \right| \left\| \partial_{p} p_{l} \right\| &\leq C_{\sharp} C_{\mu,n} \mu^{n} \overline{\Lambda}_{j,n}(p). \end{split}$$

Thus,

$$\begin{aligned} |u_n(p,\varepsilon,u) - u_n(p',\varepsilon',u')| &\leq C_{\sharp} e^{4A + 2n\varepsilon_0 \Delta} \mu^n \bigg\{ \lambda_n^+(p_n)^{-1} |u - u'| \\ &+ (\|\omega\|_{\mathcal{C}^2} + \mu^{2n} \lambda_n^+(p_n)^{-1} C_{\mu,n} |u'|) \|p - p'\| + \big[1 + \lambda_n^+(p_n)^{-1} |u'|^2 \big] |\varepsilon - \varepsilon'| \bigg\}. \end{aligned}$$

The Lemma follows recalling that equation (2.10) and our hypotheses imply $e^{\varepsilon_0 \Delta} \leq \mu$.

APPENDIX G. EXTENSION OF CURVES

In this section we explain how to extend a segment to a close curve of homotopy class (0, 1) with precise dynamical properties and explicit bounds on the derivatives.

Lemma G.1. There exist constants δ_0 , $C_{n_0,j} > 0$, $j, n_0 \in \mathbb{N} \cup \{0\}$, and $L_{\star} \geq 1$ such that for each line segment $\gamma(t) = \gamma(0) + (1, v)t$, $t \in [-\delta, \delta]$ with $\delta \leq \delta_0$, such that, for each $\mathfrak{h} \in \mathfrak{H}^{\infty}_{\gamma_-}$, if $\gamma'(t) \notin D_{\mathfrak{h}_{n_0}(\gamma(t))}F^{n_0}\mathbf{C}_u$, then we can extend γ to a closed curve $\tilde{\gamma}$, parametrized by arclength, of homotopy class (0, 1) with the following properties:

- let γ_(t) = γ(0) + ½e₁ + e₂t, then for each k ∈ N we have γ̃ ∈ Dom(𝔥_k) and 𝔥_k ∘ γ̃ is a closed curve in the homotopy class (0, 1).
- $\vartheta_{\gamma} \leq \vartheta_{\tilde{\gamma}}^{81}$
- Let $m \ge n_0$ be the smallest integer such that $D_{\gamma(t)}\mathfrak{h}_m\gamma'(t) \in \mathbf{C}_{\epsilon,c}$, for all $t \in [-\delta, \delta]$, then $D_p\mathfrak{h}_{n_0}\tilde{\gamma}' \notin \mathbf{C}_{C_{\sharp}\epsilon,u}$ and $D_p\mathfrak{h}_m\tilde{\gamma}' \in \mathbf{C}_c$.
- For each $j \in \{1, \ldots, r-1\}$ and $t \in \mathbb{R}$,

(G.1)
$$\|\tilde{\gamma}^{(j+1)}(t)\| \leq C_{n_0,j} \frac{(L_{\star}\{L_{\star},1\}^+ \epsilon^{-1} \mu^m)^j}{(\chi_u + |\pi_2(\tilde{\gamma}'(t))|)^j} := C_{n_0,j} \Delta_{\tilde{\gamma}}^j$$

⁸¹Recall the definition of ϑ_{γ} in (5.39).

Moreover, if the conditions of Lemma F.1 are satisfied, then (G.1) holds true with

(G.2)
$$L_{\star}(n_{0}) = \sup_{|v| \leq 1} L_{\star}(v, n_{0}),$$
$$L_{\star}(v, n_{0}) = C_{\sharp} \mu^{5m} C_{\mu,m}(\|\omega\|_{\mathcal{C}^{2}} + \bar{\kappa}); \quad \bar{\kappa} = |v| + \chi_{u}; \quad m - n_{0} \leq C_{\sharp} \ln \bar{\kappa}^{-1}$$

Proof. By an isometric change of variables we can assume, without loss of generality, that $\gamma(0) = 0$. Hence $\gamma(t) = (1, v)t$ for $t \in [-\delta, \delta]$ and $\gamma'(t) = (1, v) =: \overline{v}$. Note that we can assume $|v| \leq 1$ since otherwise the Lemma is trivial.

Before getting to the extension per se, we need some results on the dynamics of the tangent vectors seen as elements of a projective space. We write a vector outside the central cone as $(1, \zeta)$, so ζ can be interpreted as a projective coordinate. Then, in analogy with (A.1), we have, for each $p \in \mathbb{T}^2$ and $\zeta \in \mathbb{R}$,

$$D_p F(1,\zeta) = (\partial_x F_1 + \partial_\theta F_1 \zeta)(1, \Xi(p,\zeta))$$
$$\Xi(p,\zeta) = \frac{\partial_x F_2 + \partial_\theta F_2 \zeta}{\partial_x F_1 + \partial_\theta F_1 \zeta}.$$

Also, computing as in (A.2),

$$\partial_{\zeta} \Xi(p,\zeta) = \frac{\det(D_p F)}{\left(\partial_x F_1 + \partial_\theta F_1 \zeta\right)^2}$$

Next, for each $q \in \mathbb{T}^2$, $j \in \mathbb{N}$, let $q_j = F^j(q)$, $z_0(\zeta) = \zeta$, $z_1(q,\zeta) = \Xi(q,z_0(\zeta))$ and, for $j \ge 1$, $z_{j+1}(q_j,\zeta) = \Xi(q_j,z_j(q_{j-1},\zeta))$. In particular, if $p \in \mathbb{T}^2$ and $\Gamma_j(p) = D_{\mathfrak{h}_j(p)}F^j\mathbf{C}_c$, then $\Gamma_j(p) = \{(1,\bar{z}_j(p,\zeta)) : |\zeta| \le \chi_c\}$ where $\bar{z}_j(p,\zeta) := z_j(\mathfrak{h}_j(p),\zeta)$. Note that if $\bar{z}_j(p,\chi_c) \notin \mathbf{C}_u$, then we have

(G.3)
$$|\bar{z}_j(p,\chi_c)| \le C_{\sharp} \lambda_j^-(\mathfrak{h}_j(p))^{-1} \mu^j \chi_c.$$

Moreover,

$$\partial_{\zeta} \bar{z}_j(p,\zeta) = \partial_{\bar{z}} \Xi(\mathfrak{h}_j(p), \bar{z}_{j-1}(p,\zeta)) \partial_{\zeta} \bar{z}_{j-1}(p,\zeta).$$

Iterating the above identities and recalling Propositions 5.6, 5.8 we have

$$C_{\sharp}\frac{\mu^{-j}}{\lambda_j^-(\mathfrak{h}_j(p))} \le |\partial_{\zeta}\bar{z}_j(p,\zeta)| \le C_{\sharp}\frac{\lambda_j^+(\mathfrak{h}_j(p))\mu^j}{\lambda_j^-(\mathfrak{h}_j(p))^2} \le C_{\sharp}\frac{\mu^{-j}}{\lambda_j^-(\mathfrak{h}_j(p))}.$$

It follows, by (G.3), that

$$|\bar{z}_j(p,\pm\chi_c)-\bar{z}_j(p,\pm\chi_c(1-\epsilon))| \ge C_{\sharp}\epsilon\mu^{-2j}|\bar{z}_j(p,\pm\chi_c)|.$$

Let $\mathfrak{a}_{j}(t,\mathfrak{h}) = \frac{\pi_{2}(D_{\gamma(t)}\mathfrak{h}_{j}\bar{v})}{\pi_{1}(D_{\gamma(t)}\mathfrak{h}_{j}\bar{v})}$, and $m(\mathfrak{h},t)$ be the smallest integer k such that $D_{\gamma(t)}\mathfrak{h}_{k}\bar{v} \in \mathbf{C}_{\epsilon,c}$.

In the following we consider only the case $\mathfrak{a}_{m(\mathfrak{h},t)}(t,\mathfrak{h}) \geq \chi_c$. The case $\mathfrak{a}_{m(\mathfrak{h},t)}(t,\mathfrak{h}) \leq -\chi_c$ is totally analogous. Also, we consider only t > 0 since the construction for t < 0 is exactly the same.

It follows $C_{\sharp}\bar{z}_{m(\mathfrak{h},t)}(\mathfrak{h}_{m(\mathfrak{h},t)}(\gamma(t)),\chi_c) \geq v \geq \bar{z}_{m(\mathfrak{h},t)}(\mathfrak{h}_{m(\mathfrak{h},t)}(\gamma(t)),\chi_c(1-\epsilon))$. Thus, setting $m'(\mathfrak{h},t) = m(\mathfrak{h},t) - n_0$, we have

(G.4)
$$\mathfrak{a}_{n_0}(\mathfrak{h},t) - \bar{z}_{m'(\mathfrak{h},t)}(\mathfrak{h}_{n_0}(\gamma(t)),\chi_c) \ge C_{\sharp}\epsilon\mu^{-2m}\bar{z}_{m'(\mathfrak{h},t)}(\mathfrak{h}_{n_0}(\gamma(t)),\chi_c) \ge C_{\sharp}\epsilon\mu^{-2m}v.$$

We are finally ready to extend our segment. For $\varphi \in \mathbb{R}$, let $w(\varphi) = (\cos \varphi, \sin \varphi)$, $\theta = \tan v$ and $a = \sqrt{1 + v^2}$. Then $\bar{v} = aw(\theta)$. We start by extending the curve to the interval $(\delta, \delta + A)$, with $A = \frac{C_{n_0}}{2a} \epsilon \mu^{-2m} \bar{\kappa} L_{\star}^{-1} < 1$. Next, let L_{\star} be the maximal Lipschitz constant of the $\bar{z}_j(\cdot, \pm \chi_c)$ for $j \leq m - n_0$.

Let $b \in \mathcal{C}^{\infty}(\mathbb{R}, [0, 1])$ be a bump function with b(t) = 0 for $t \leq 0$ and b(t) = 1 for $t \geq 1$. Also, let $B = \{aL_{\star}, 16\theta_c, \Bbbk\}^+$, for some $\Bbbk \geq 1$ to be chosen later, and where $\theta_c := \arctan(2\chi_c^{-1})$, and define, for $t \in [\delta, T]$, where T will be chosen shortly,

(G.5)
$$\hat{\gamma}'(t) = aw(\theta + b((t-\delta)A^{-1})B(t-\delta)) =: aw(\tilde{\theta}(t)).$$

Note that, by construction, $\tilde{\theta}(t) \geq \theta$. Moreover, for $t \in [\delta, \delta + A]$, we have

$$\|\hat{\gamma}(t) - \hat{\gamma}(\delta)\| \le \int_{\delta}^{\delta + A} \|aw(\tilde{\theta}(ts))\| ds \le Aa.$$

Thus, recalling (G.4), for $t \in [\delta, \delta + A]$,

$$\operatorname{rctan} \bar{z}_{m(\mathfrak{h},\gamma(\delta))}(\hat{\gamma}(t),\chi_c) \leq \arctan \bar{z}_{m(\mathfrak{h},\gamma(\delta))}(\hat{\gamma}(\delta),\chi_c) + L_{\star}aA$$
$$\leq \theta - C_{n_0}\epsilon\bar{\kappa}\mu^{-2m} + L_{\star}Aa < \theta \leq \tilde{\theta}(t),$$

which implies that $D_{\tilde{\gamma}(t)}\mathfrak{h}_m\hat{\gamma}'(t)\in \mathbf{C}_c$. In addition, for $t\geq \delta+A$, we have

$$\left|\frac{d}{dt}\tan\tilde{\theta}(t)\right| \ge B \ge aL_{\star} \ge \left|\frac{d}{dt}\bar{z}_{m}(\tilde{\gamma}(t))\right|,$$

thus $D_{\tilde{\gamma}(t)}\mathfrak{h}_m \hat{\gamma}'(t) \in \mathbf{C}_c$ also for $t \ge \delta + A$.

We now choose T > 0 be such that $\tilde{\theta}(T) = \theta_c$ so that $\hat{\gamma}'(T)$ is well inside the central cone. This implies $T \leq \delta + \theta_c B^{-1}$ and

$$|\pi_1(\hat{\gamma}(T))| \le C_{\sharp}T \le C_{\sharp}(\delta + B^{-1}) \le C_{\sharp}(\delta_0 + \mathbb{k}^{-1}) < 1/4,$$

provided δ_0 and \mathbb{k}^{-1} are small enough. It is then a simple exercise to define $\hat{\gamma} : [T, S] \to \mathbb{T}^2$ such that $\|\hat{\gamma}'\| = a$; $\hat{\gamma}'(t) \in \mathbf{C}_c$, for all $t \in [T, S]$, and $\hat{\gamma}(S) = (0, 1/2)$, $|\pi_1(\hat{\gamma})| \leq C_{\sharp}(\delta_0 + \mathbb{k}^{-1})$, $\hat{\gamma}'(S) = (-\chi_c/2, 1)$, $\hat{\gamma}^{(j)}(S) = 0$ for all j > 1 and $\sup_{t \in [T,S]} \|\hat{\gamma}^{(j)}(t)\| \leq C_{\sharp}$. By symmetry we have a closed curve $\hat{\gamma}$ of homotopy class (0, 1).

Note that $\hat{\gamma} \in \text{Dom}(\mathfrak{h}_k)$ for each $\mathfrak{h} \in \mathfrak{H}_{\gamma_-}^{\infty}$ and $k \in \mathbb{N}$. Then Lemma 5.3 implies that there exists inverse branches $\{\mathfrak{h}_{k,i}\}_{i=1}^{d^k}$, where d is the degree of F, such that $F^{-k}\hat{\gamma} = \bigcup_{i=1}^{d^k} \mathfrak{h}_{k,i} \circ \hat{\gamma}$. Since $\mathfrak{h}_{k,i}$ is a diffeomorphism, $\mathfrak{h}_{k,i} \circ \hat{\gamma}$ is a closed curve. In addition it must be of homotopy type (0,1), otherwise it would intersect an horizontal segment in more than one point and the image, under F^k , of the interval between two intersection points would be an unstable curve going from $\hat{\gamma}$ to itself. Since such a curve would be transversal to $\hat{\gamma}$ by hypothesis, it follows that it would have to wrap around the torus horizontally an hence intersect γ_- contradicting the fact that it is in the domain of $\mathfrak{h}_{k,i}$.

Recalling (G.5), formula (5.5) gives, for all $j \ge 2$,⁸²

$$\begin{aligned} \|\hat{\gamma}\|_{\mathcal{C}^{j+1}} &\leq C_{\sharp} \|w \circ \tilde{\theta}\|_{\mathcal{C}^{j}} \leq C_{\sharp} \sum_{s=0}^{j} \|w\|_{\mathcal{C}^{s}} \sum_{k \in \mathcal{K}_{j,s}} \prod_{l \in \mathbb{N}}^{j} \|\tilde{\theta}\|_{\mathcal{C}^{l}}^{k_{l}} \\ &\leq C_{\sharp} \sum_{s=0}^{j} \sum_{k \in \mathcal{K}_{j,s}} \prod_{l \in \mathbb{N}} \left(A^{-l}B\right)^{k_{l}} \leq A^{-j} \sum_{s=0}^{j} B^{s}. \end{aligned}$$

Thus, since $\|\hat{\gamma}'\| = a$, we can reparametrize the curve by arc-length. Calling $\tilde{\gamma}$ the reparametrized curve we obtain

$$\|\tilde{\gamma}^{(j)}(t)\| \leq \begin{cases} 0 & \text{if } |t| \leq \delta\\ C_{\sharp} A^{-j+1} B^{j-1} & \text{if } \delta \leq |t| \leq \delta + A\\ C_{\sharp} B^{j-1} & \text{if } |t| \geq \delta + A, \end{cases}$$

which yields (G.1) since

$$|\pi_2(\tilde{\gamma}'(t))| \ge \begin{cases} |v| & \text{if } |t| \le \delta + A\\ C_{\sharp}(|v| + B(t - \delta)) & \text{if } |t| \ge \delta + A. \end{cases}$$

⁸²Notice that, as $\|\tilde{\theta}\|_{\mathcal{C}^l} \leq C_{\sharp} A^{-l} B$, recalling the definition of $\mathcal{K}_{j,s}$ we have

$$\sum_{k \in \mathcal{K}_{j,s}} \prod_{l \in \mathbb{N}} (A^{-l}B)^{k_l} \lesssim \sum_{k \in \mathcal{K}_{j,s}} A^{-\sum_{l=1}^{\infty} lk_l} B^{\sum_{l=1}^{\infty} k_l} \lesssim A^{-j}B^s.$$

To conclude we estimate L_{\star} when the hypotheses of Lemma F.1 are satisfied. In a finite number of steps n_2 (depending only on the derivatives of F) we can have $\bar{z}_{n_2} \leq 1/2$, we can thus apply Lemma F.1 $\varepsilon_0 = \varepsilon = \varepsilon' = 1$, A = 1/2, $B \leq C_{\sharp}$ and $u = u' = \bar{z}_{n_2}(p)$, we have

$$\begin{aligned} |\bar{z}_{m-n_0}(p,\chi_c) - \bar{z}_{m-n_0}(p',\chi_c)| &\leq L_{m-n_0} \|p - p'\| \\ L_j &= C_{\sharp} \mu^{3j}(\|\omega\|_{\mathcal{C}^2} + \mu^{2j} \lambda_j^+(p)^{-1} C_{\mu,j}/2). \end{aligned}$$

Next, note that, by usual distortion arguments, equation (G.4) implies $\lambda_{m-n_0}^+ \ge C_{\sharp} \mu^{-n_0} \chi_c(\bar{\kappa})^{-1}$ and $m-n_0 \le C_{\sharp} \ln \bar{\kappa}^{-1}$, thus

$$L_{m-n_0} \le C_{\sharp} \mu^{5m} C_{\mu,m}(\|\omega\|_{\mathcal{C}^2} + \bar{\kappa}) = L_{\star}(v, n_0)$$

References

- Alves, J; Bonatti, C.; Viana, M. SRB measures for partially hyperbolic systems whose central direction is mostly expanding. Invent. Math. 140 (2000), no. 2, 351–398.
- [2] Alves, J; Luzzatto, S; Pinheiro, V Markov structures and decay of correlations for non-uniformly expanding dynamical systems. Ann. Inst. H. Poincaré Anal. Non Linéaire, 22(6):817–839, 2005.
- [3] Avila, A; Gouëzel, S; Tsujii, M. Smoothness of solenoidal attractors, Discrete and Continuous Dynamical Systems 15 (2006), no. 1, 21-35.
- Baladi, V. Positive transfer operators and decay of correlations. Advanced Series in Nonlinear Dynamics, 16. World Scientific Publishing Co., Inc., River Edge, NJ, 2000.
- [5] Baladi, V. Dynamical zeta functions and dynamical determinants for hyperbolic maps. A functional approach. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], 68. Springer, Cham, 2018.
- [6] F. deBruno Note sur une nouvelle formule de calcul différentiel Quart. J. Math., 1 (1856), pp. 359-360.
- Baladi, Viviane; Tsujii, Masato Anisotropic Hölder and Sobolev spaces for hyperbolic diffeomorphisms. Ann. Inst. Fourier (Grenoble) 57 (2007), no. 1, 127–154.
- [8] Baladi, Viviane; Demers, Mark F.; Liverani, Carlangelo Exponential decay of correlations for finite horizon Sinai billiard flows. Invent. Math. 211 (2018), no. 1, 39–177.
- Bernardin, Cédric; Huveneers, François; Lebowitz, Joel L.; Liverani, Carlangelo; Olla, Stefano Green-Kubo formula for weakly coupled systems with noise. Comm. Math. Phys. 334 (2015), no. 3, 1377–1412.
- [10] Wael Bahsoun, Carlangelo Liverani Anosov diffeomorphisms, anisotropic BV spaces and regularity of foliations. Preprint arXiv:2003.06811.
- Blank, Michael; Keller, Gerhard; Liverani, Carlangelo Ruelle-Perron-Frobenius spectrum for Anosov maps. Nonlinearity 15 (2002), no. 6, 1905–1973.
- [12] Bonatti C.; Viana, M. SRB measures for partially hyperbolic systems whose central direction is mostly contracting. Israel J. Math. 115 (2000), 157–193.
- [13] Burns, Keith; Wilkinson, Amie. On the ergodicity of partially hyperbolic systems. Ann. of Math. (2) 171 (2010), no. 1, 451–489.
- [14] Butterley, Oliver; Eslami, Peyman. Exponential Mixing for Skew Products with Discontinuities. Amer. Math. Soc., 369:783803, (2017).
- [15] Butterley, O; War, K. Open Sets of Exponentially Mixing Anosov Flows To appear in J. Eur. Math. Soc.
- [16] Castro Júnior A. A Backward inducing and exponential decay of correlations for partially hyperbolic attractors. Israel J. Math., 130:29–75, 2002.
- [17] Chernov, N. I Markov approximations and decay of correlations for Anosov flows. Ann. of Math. (2), 147(2):269– 324, 1998.
- [18] Mark Demers, Niloofar Kiamari, Carlangelo Liverani. Transfer operators in Hyperbolic Dynamics An introduction.
 33 Colloquio Brasilero de Matematica. Editora do IMPA. pp.252 (2021).
- [19] De Simoi, Jacopo; Liverani, Carlangelo, Statistical properties of mostly contracting fast-slow partially hyperbolic systems. Invent. Math. 206 (2016), no. 1, 147–227.
- [20] De Simoi, Jacopo; Liverani, Carlangelo Limit theorems for fast-slow partially hyperbolic systems. Invent. Math. 213 (2018), no. 3, 811–1016.
- [21] De Simoi, Jacopo; Liverani, Carlangelo The martingale approach after Varadhan and Dolgopyat. Hyperbolic dynamics, fluctuations and large deviations, 311339, Proc. Sympos. Pure Math., 89, Amer. Math. Soc., Providence, RI, 2015.
- [22] Jacopo De Simoi, Carlangelo Liverani, Christophe Poquet, and Denis Volk. Fast-Slow Partially Hyperbolic Systems Versus Freidlin Wentzell Random Systems. J. Stat. Phys., 166(3-4):650-679, 2017.
- [23] D. Dolgopyat. On decay of correlations in Anosov flows. Ann. of Math. (2), 147(2):357-390, 1998.

- [24] Dolgopyat, D. On dynamics of mostly contracting diffeomorphisms. Comm. Math. Phys., 213(1):181-201, 2000.
- [25] Dolgopyat, D. On mixing properties of compact group extensions of hyperbolic systems. Israel J. Math., 130:157–205, 2002.
- [26] Dolgopyat, Dmitry; Liverani, Carlangelo Energy transfer in a fast-slow Hamiltonian system. Comm. Math. Phys. 308 (2011), no. 1, 201–225.
- [27] Dyatlov, Semyon; Zworski, Maciej Ruelle zeta function at zero for surfaces. Invent. Math. 210 (2017), no. 1, 211–229.
- [28] F. Faure. Semiclassical origin of the spectral gap for transfer operators of a partially expanding map. Nonlinearity 24 (2011), no. 5, 1473–1498.
- [29] F. Faure; M. Tsujii. The semiclassical zeta function for geodesic flows on negatively curved manifolds. Invent. Math. 208 (2017), no. 3, 851–998.
- [30] Field, M.; Melbourne, I.; Török, A. Stability of mixing and rapid mixing for hyperbolic flows. Ann. of Math. (2), 166(1):269–291, 2007.
- [31] Giulietti, P.; Liverani, C.; Pollicott, M. Anosov flows and dynamical zeta functions. Ann. of Math. (2) 178 (2013), no. 2, 687–773.
- [32] S. Gouëzel, C. Liverani, Banach spaces adapted to Anosov systems, Ergodic Theory and Dynamical Systems 26 (2006), 189-217.
- [33] Grayson, M.; Pugh, C.; Shub, M. Stably ergodic diffeomorphisms. Ann. of Math. (2), 140(2):295–329, 1994.
- [34] Hatcher Allen, Algebraic Topology, Cornell University, New York, 2002.
- [35] H.Hennion, Sur un theoreme spectral et son application aux noyaux Lipchitziens, Proceedings of the American Mathematical Society, 118 (1993), 627–634.
- [36] Hernndéz Encinas, L.H.; Munoz Masqué, J. A short proof of the generalized Faà di Bruno's formula. Applied mathematics lectures 16 (2003), 975-979.
- [37] Hirsch, M.; Pugh, C.; Shub, M. Invariant manifolds. Volume 583 of Led. Notes in Math. Springer Verlag, 1977.
- [38] Hörmander, L. The analysis of linear partial differential operators. I. 2nd edition, Springer-Verlag, Berlin, 1990.
- [39] W.P. Johnson, Combinatorics of Higher Derivatives of Inverses. The American Mathematical Monthly Vol. 109, No. 3, pp. 273-277, (2002).
- [40] Keller, G.; Liverani, C. A spectral gap for a one-dimensional lattice of coupled piecewise expanding interval maps. Dynamics of coupled map lattices and of related spatially extended systems, 115–151, Lecture Notes in Phys., 671, Springer, Berlin, 2005.
- [41] Keller, Gerhard; Liverani, Carlangelo Uniqueness of the SRB measure for piecewise expanding weakly coupled map lattices in any dimension. Comm. Math. Phys. 262 (2006), no. 1, 33–50.
- [42] Krylov, N. S. Works on the foundations of statistical physics. Princeton University Press, Princeton, N.J., 1979. Translated from the Russian by A. B. Migdal, Ya. G. Sinai [Ja. G. Sinaĭ] and Yu. L. Zeeman [Ju. L. Zeeman], With a preface by A. S. Wightman, With a biography of Krylov by V. A. Fock [V. A. Fok], With an introductory article "The views of N. S. Krylov on the foundations of statistical physics" by Migdal and Fok, With a supplementary article "Development of Krylov's ideas" by Sinaĭ, Princeton Series in Physics.
- [43] Liverani, C. On contact Anosov flows. Ann. of Math. (2), 159(3):1275–1312, 2004.
- [44] Liverani, Carlangelo Transport in partially hyperbolic fast-slow systems. Proceedings of the International Congress of Mathematicians—Rio de Janeiro 2018. Vol. III. Invited lectures, 2643–2667, World Sci. Publ., Hackensack, NJ, 2018.
- [45] Melbourne, I. Rapid decay of correlations for nonuniformly hyperbolic flows. Trans. Amer. Math. Soc., 359(5):2421– 2441 (electronic), 2007.
- [46] Nakano, Yushi; Tsujii, Masato; Wittsten, Jens The partial captivity condition for U(1) extensions of expanding maps on the circle. Nonlinearity 29 (2016), no. 7, 1917–1925.
- [47] Pugh, C.; Shub, M. Stably ergodic dynamical systems and partial hyperbolicity. J. Complexity, 13(1):125–179, 1997.
- [48] Tsujii, M. Decay of correlations in suspension semi-flows of angle-multiplying maps, Ergodic Theory and Dynamical Systems 28 (01), 291-317, 2005.
- [49] Tsujii, M. Fat solenoidal attractors. Nonlinearity, 14, 1011–1027 (2001).
- [50] Tsujii, M. Physical measures for partially hyperbolic surface endomorphisms, Acta Math. 194 (2005), 37-132.
- [51] Tsujii, M. Quasi-compactness of transfer operators for contact Anosov flows. Nonlinearity, 23(7):1495–1545, 2010.
- [52] Masato Tsujii, Zhiyuan Zhang, Smooth mixing Anosov flows in dimension three are exponential mixing, Preprint arXiv:2006.04293.
- [53] Wilkinson, Amie. Stable ergodicity of the time-one map of a geodesic flow. Ergodic Theory Dynam. Systems 18 (1998), no. 6, 1545–1587.
- [54] Zhang, Zhiyuan, On the smooth dependence of SRB measures for partially hyperbolic systems. Comm. Math. Phys. 358 (2018), no. 1, 45–79.