Polar coordinates view on KM-arcs

Kanat Abdukhalikov and Duy Ho UAE University, PO Box 15551, Al Ain, UAE abdukhalik@uaeu.ac.ae, duyho92@gmail.com

Abstract

We study presentations of KM-arcs in polar coordinates. New characterizations on the point set of KM-arcs are obtained in terms of power sums and bilinear forms. Therefore, we provide purely algebraic criteria for the existence of KM-arcs. We also describe some constructions of KM-arcs which include known examples and describe their automorphism groups in this presentation.

Keywords: Arcs, ovals, hyperovals, power sums, Vandermonde sets.

1 Introduction

In [13], Korchmáros and Mazzocca initiated the study of (q + t, t)-arc of type (0, 2, t) in the projective plane PG(2, q). These objects are sets of q + t points meeting every line in 0, 2 or t points. They are also called *KM*-arcs of type t in honour of the authors of [13], as mentioned in [9]. KM-arcs were studied also in [8, 11, 16, 17].

KM-arcs have been investigated exclusively in homogeneous coordinates. This presentation has its own advantages with the established algebraic methods over finite fields. On the other hand, the drawback of this approach is that there are points of KM-arcs presented on the infinite line. Descriptions for structural results of KM-arcs are then required to be split between points in the affine plane and infinite points, as seen in the constructions by Gács and Weiner in [11].

In this paper, we describe KM-arcs in polar coordinates. This approach was initiated by investigations in [1, 2, 3, 4]. With this approach, KM-arcs can be described solely with points in the affine plane. As a consequence, structural results of KM-arcs can be obtained uniformly on the point set. To demonstrate this point of view, we provide new characterizations of KM-arcs in terms of bilinear forms and power sums. We also describe some examples of KM-arcs in this presentation.

The paper is organized as follows. In Section 2, we recall preliminary results on projective planes, affine planes and KM-arcs. In Section 3, we describe a configuration in Definition 1 that we use to describe KM-arcs in polar coordinates. Also in this section we obtain characterizations of KM-arcs in terms of bilinear forms and power sums, in Theorem 1 and Theorem 3, respectively. In Section 4 and Section 5, we describe constructions of KM-arcs in PG(2,q) using hyperovals and KM-arcs in PG(2,r), where q and r are powers of 2 with $q = r^l$. We also show that the KM-arcs constructed in Section 5 are translation KM-arcs and describe their automorphisms.

2 Preliminaries

In this section we recall some definitions and notation.

2.1 Polar representation

In this paper we consider finite fields only in characteristics 2. Let $F = \mathbb{F}_{2^m}$ be a finite field of order $q = 2^m$. Consider F as a subfield of $K = \mathbb{F}_{2^n}$, where n = 2m, so K is a two dimensional vector space over F.

The *conjugate* of $x \in K$ over F is

$$\bar{x} = x^q$$
.

Then the *trace* and the *norm* maps from K to F are

$$T(x) = Tr_{K/F}(x) = x + \bar{x} = x + x^q,$$

 $N(x) = N_{K/F}(x) = x\bar{x} = x^{1+q}.$

The *unit circle* of K is the set of elements of norm 1:

$$S = \{ u \in K \mid u\bar{u} = 1 \}.$$

Therefore, S is the multiplicative group of (q+1)st roots of unity in K. Since $F \cap S = \{1\}$, each non-zero element of K has a unique polar coordinate representation $x = \lambda u$ with $\lambda \in F^*$ and $u \in S$. For any $x \in K^*$ we have $\lambda = \sqrt{x\bar{x}}$ and $u = \sqrt{x/\bar{x}}$.

One can define a nondegenerate bilinear form $\langle \cdot, \cdot \rangle : K \times K \to F$ by

$$\langle x, y \rangle = T(x\bar{y}) = x\bar{y} + \bar{x}y.$$

Then the form $\langle \cdot, \cdot \rangle$ is alternating and symmetric, that is, $\langle a, a \rangle = 0$ and $\langle a, b \rangle = \langle b, a \rangle$.

Following [10], consider an element $\mathbf{i} \in K$ with property $T(\mathbf{i}) = \mathbf{i} + \mathbf{i}^q = 1$. Then $K = F(\mathbf{i})$ and \mathbf{i} is a root of a quadratic equation

$$z^2 + z + \delta = 0,$$

where $\delta = N(\mathbf{i}) \in F$. Any element $z \in K$ can be represented as $z = x + y\mathbf{i}$, where $x, y \in F$. For $z = x + y\mathbf{i}$ we have $x = \langle \mathbf{i}, z \rangle$, and $y = \langle 1, z \rangle$.

2.2 Affine and projective planes

Consider points of a projective plane PG(2,q) in homogeneous coordinates [12] as triples (x : y : z), where $x, y, z \in F$, $(x, y, z) \neq (0, 0, 0)$, and we identify (x : y : z) with $(\lambda x : \lambda y : \lambda z), \lambda \in F^*$. Then the points of PG(2,q) are

$$\{(x:y:1) \mid x \in F, y \in F\} \cup \{(x:1:0) \mid x \in F\} \cup \{(1:0:0)\}.$$

For $a, b, c \in F$, $(a, b, c) \neq (0, 0, 0)$, the line [a : b : c] in PG(2, q) is defined as

$$[a:b:c] = \{(x:y:z) \in PG(2,q) \mid ax + by + cz = 0\}.$$

Triples [a:b:c] and $[\lambda a:\lambda b:\lambda c]$ with $\lambda \in F^*$ define the same lines. The point (x:y:z) is incident with the line [a:b:c] if and only if ax + by + cz = 0. We shall call points of the form (x:y:0) the points at infinity. Then [0:0:1] indicates the line at infinity.

We define an affine plane $AG(2,q) = PG(2,q) \setminus [0:0:1]$, so points of this affine plane AG(2,q) are $\{(x:y:1) \mid x, y \in F\}$. Associating (x:y:1) with (x,y) we can identify points of the affine plane AG(2,q) with elements of the vector space V(2,q) = $\{(x,y) \mid x, y \in F\}$, and we will write AG(2,q) = V(2,q). Lines in AG(2,q) = V(2,q) are $\{(c,y) \mid y \in F\}$ and $\{(x,xb+a) \mid x \in F\}$, $a, b, c \in F$. These lines can be described by equations x = c and y = xb + a.

We introduce now another representation of PG(2, q) using the field K. Consider pairs (x : z), where $x \in K, z \in F, x \neq 0$ or $z \neq 0$, and we identify (x : z) with $(\lambda x : \lambda z), \lambda \in F^*$. Then points of PG(2, q) are

$$\{(x:1) \mid x \in K\} \cup \{(u:0) \mid u \in S\}.$$

For $\alpha \in K$, $\beta \in F$, $(\alpha, \beta) \neq (0, 0)$, we define the lines $[\alpha : \beta]$ in PG(2, q) as

$$[\alpha:\beta] = \{ (x:z) \in PG(2,q) \mid \langle \alpha, x \rangle + \beta z = 0 \}.$$

Pairs $[\alpha : \beta]$ and $[\lambda \alpha : \lambda \beta]$ with $\lambda \in F^*$ define the same lines. The point (x : z) is incident with the line $[\alpha : \beta]$ if and only if $\langle \alpha, x \rangle + \beta z = 0$. The element $u_{\infty} = (u : 0), u \in S$, will be referred to as the point at infinity in the direction of u. So [0:1] indicates the line at infinity.

We define an affine plane $AG(2,q) = PG(2,q) \setminus [0:1]$, so points of this affine plane AG(2,q) are $\{(x:1) \mid x \in K\}$. Associating (x:1) with $x \in K$ we can identify points of the affine plane AG(2,q) with elements of the field K, and we write AG(2,q) = K. Lines of AG(2,q) = K are of the form

$$L(u,\mu) = \{ x \in K \mid \langle u, x \rangle + \mu = 0 \},\$$

where $u \in S$ and $\mu \in F$ (cp. [6, subsection 2.1]).

Throughout the paper, we will consider these two representations of the projective plane PG(2,q), and for each of such projective planes we consider a fixed affine plane AG(2,q) described above. They will be written as AG(2,q) = V(2,q) and AG(2,q) = K.

2.3 KM-arcs and basic facts

In the projective plane PG(2,q), a KM-arc of type t (also known as a (q+t,t)-arc of type (0,2,t)) is a set H of q+t points meeting every line in 0, 2 or t points. If H is a KM-arc of type t in PG(2,q), 2 < t < q, then

- 1. q is even and t is a divisor of q;
- 2. each point of H is on exactly one *t*-secant and every other line through this point is a 2-secant of H, cp. [13];
- 3. there are q/t + 1 different t-secants to H, and they are concurrent at a unique point called the t-nucleus of H, cp. [11];
- 4. all other lines contain 0 or 2 points of H, cp. [17].

The case t = q is classified as H can be shown to be symmetric difference of two lines. The case t = 1 is a degenerate case when H is an *oval* of q + 1 points and therefore has a unique nucleus, which we call the 1-nucleus of H. When t = 2, the set H is called a *hyperoval*. In this case, if p is a point in PG(2,q) not on H, then there are q/2 + 1different 2-secants of H going through p. Thus, any point not on H can be considered as a 2-nucleus of H.

2.4 Vandermonde sets

We recall sets with the following property first considered by Gács and Weiner [11] (and subsequently by other authors in [7, 15]). Let 1 < t < q. A set $T = \{y_1, \dots, y_t\} \subseteq GF(q)$ is called a *Vandermonde set* if

$$\pi_k(T) := \sum_{y \in T} y^k = 0,$$

for all $1 \leq k \leq t - 2$.

Lemma 1 (cp. [15], [5]). For a set of t elements the Vandermonde property is invariant under transformations $y \rightarrow ay + b, a \neq 0$, if and only if t is even or b = 0.

Some examples of Vandermonde sets can be found in [15, Proposition 1.8]. In Section 4, we will use the fact that any additive subgroup of GF(q) is a Vandermonde set.

3 KM-arcs in polar presentation

In this section we obtain characterizations of KM-arcs in terms of bilinear forms and power sums.

3.1 The configuration of a KM-arc in polar presentation

Lemma 2. Any KM-arc in PG(2,q) is equivalent to a KM-arc in AG(2,q) with t-nucleus at 0.

Proof. Assume H has a point on infinity line L_{∞} . Let L be a line which does not intersect H. Consider a collineation γ interchanging the lines L and L_{∞} . Then $\gamma(H)$ has no points at infinity and a suitable affine translation of $\gamma(H)$ is a KM-arc with t-nucleus at 0.

Lemma 3. For $t \ge 2$, let H be a set of q + t points in K = AG(2, q). Assume that for every point p of H there exists a line L going through p such that $|L \cap H| \ge t$. Then His a KM-arc of type t if and only if every line intersects H at an even number of points.

Proof. If H is a KM-arc of type t then every line intersects H at 0, 2 or t points.

Assume every line intersects H at an even number of points. Let p be a point on H and let L be a line going through p such that $|L \cap H| \ge t$. Let \mathcal{L} be the set of q + 1 lines going through p. Since there are at most q points in $H \setminus (H \cap L)$, each line in $\mathcal{L} \setminus L$ contains exactly two points of H (including p). This also implies $|L \cap H| = t$. In particular, every line intersects H at 0, 2 or t points and so H is a KM-arc of type t.

We now describe a configuration H of points that is used throughout the paper.

Definition 1. Let $t \ge 2$. A set H of q + t points in K = AG(2, q) is called a *star-set* if all points of H belong to a union of $\frac{q}{t} + 1$ lines concurrent at 0, and each line contains t points of H.

By Lemma 3, a star-set H is a KM-arc of type $t \ge 2$ (with t-nucleus at 0) if and only if every line intersects H at an even number of points.

Theorem 1. Let H be a star-set. Then H is a KM-arc of type $t \ge 2$ with t-nucleus at 0 if and only if for each $v \in S$ and $1 \le k \le q - 2$,

$$\sum_{y \in H} \langle v, y \rangle^k = 0.$$
 (1)

Proof. The case t = 2 is proved in [5]. In the remainder of the proof, we assume t > 2. Assume H is a KM-arc of type t with t-nucleus at 0. Fix $v \in S$. For each $\mu \in F$, the line $\langle v, x \rangle = \mu$ intersects H at an even number of points. In particular, the equation

$$\langle v, x \rangle = \mu$$

has an even number of solutions in H. This implies condition (1) is true for each $v \in S$ and $1 \leq k \leq q-2$.

Conversely, assume condition (1) is true for each $v \in S$ and $1 \leq k \leq q-2$. We will show that every line $\langle v, x \rangle = \mu$ intersects H at an even number of points. Fix $v \in S$. For $\mu = 0$, the intersection H_v of H and the line $\langle v, x \rangle = 0$ is of size 0 or t. For each $\mu \in F^*$, let

$$Y_{\mu} := \{ y \in H \mid \langle v, y \rangle = \mu \} .$$

We note that, for each $1 \le k \le q-2$,

$$\sum_{y \in Y_{\mu}} \langle v, y \rangle^{k} = \begin{cases} \mu^{k} & \text{if } |Y_{\mu}| \text{ is odd,} \\ 0 & \text{if } |Y_{\mu}| \text{ is even.} \end{cases}$$

Let $\Omega := \{ \mu \in F^* \mid |Y_\mu| \text{ is odd} \}$. Assume that $\Omega \neq \emptyset$. Then

$$0 = \sum_{y \in H} \langle v, y \rangle^k = \sum_{\mu \in \Omega} \sum_{y \in Y_\mu} \langle v, y \rangle^k = \sum_{\mu \in \Omega} \mu^k.$$

Also, the sets Y_{μ} partition the set $H \setminus H_v$ of even size (either q or q+t), so that $|\Omega|$ is even. In particular, the sum of the vectors $(1, \mu, \mu^2, \cdots, \mu^{q-2}), \mu \in \Omega$, is the zero vector. On the other hand, the Vandermonde's determinant implies that these vectors are linearly independent. Hence $\Omega = \emptyset$.

We have shown that every line $\langle v, x \rangle = \mu$ intersects H at an even number of points. By Lemma 3, H is a KM-arc of type t with t-nucleus at 0.

3.2 KM-arcs and power sums

We recall and extend [5, Lemma 9] to the following.

Lemma 4. The power of the bilinear form $\langle \cdot, \cdot \rangle$ is given by

$$\langle a,b\rangle^k = \sum_{i=0}^{\lfloor (k-1)/2 \rfloor} \binom{k}{i} \langle a^{iq+k-i}, b^{iq+k-i}\rangle = \sum_{i=\lceil (k+1)/2 \rceil}^k \binom{k}{i} \langle a^{iq+k-i}, b^{iq+k-i}\rangle.$$

Proof. It was shown in [5, Lemma 9] that

$$\langle a,b\rangle^k = \sum_{i=0}^{\lfloor (k-1)/2 \rfloor} \binom{k}{i} \langle a^{iq+k-i}, b^{iq+k-i} \rangle.$$

For the remaining equality, when k is odd we have

$$\begin{split} \langle a,b\rangle^{k} &= (a^{q}b + ab^{q})^{k} = \sum_{i=0}^{k} \binom{k}{i} a^{iq}b^{i}a^{k-i}b^{(k-i)q} = \sum_{i=0}^{k} \binom{k}{i} a^{iq+k-i}b^{i+(k-i)q} \\ &= \sum_{i=0}^{(k-1)/2} \binom{k}{i} a^{iq+k-i}b^{i+(k-i)q} + \sum_{i=(k+1)/2}^{k} \binom{k}{i} a^{iq+k-i}b^{i+(k-i)q} \\ &= \sum_{i=(k+1)/2}^{k} \binom{k}{i} a^{i+(k-i)q}b^{iq+k-i} + \sum_{i=(k+1)/2}^{k} \binom{k}{i} a^{iq+k-i}b^{i+(k-i)q} \\ &= \sum_{i=(k+1)/2}^{k} \binom{k}{i} \left(a^{iq+k-i}b^{i+(k-i)q} + a^{i+(k-i)q}b^{iq+k-i} \right) \\ &= \sum_{i=(k+1)/2}^{k} \binom{k}{i} \langle a^{iq+k-i}, b^{iq+k-i} \rangle. \end{split}$$

Similar to the above, when k is even, we have

$$\begin{split} \langle a,b\rangle^k &= \sum_{i=k/2+1}^k \binom{k}{i} \langle a^{iq+k-i}, b^{iq+k-i} \rangle + \binom{k}{k/2} \langle a^{iq+k-i}, b^{iq+k-i} \rangle \\ &= \sum_{i=k/2+1}^k \binom{k}{i} \langle a^{iq+k-i}, b^{iq+k-i} \rangle, \end{split}$$

since $\binom{k}{k/2}$ is even.

We recall the following partial ordering \leq on the set of nonnegative integers. If

$$x = \sum_{i=0}^{m} x_i 2^i$$
 and $y = \sum_{i=0}^{m} y_i 2^i$

(where each x_i and each y_i is either 0 or 1), then $x \leq y$ if and only if $x_i \leq y_i$ for all *i*. In other words, $x \leq y$ if and only if all nonzero terms appearing in the binary expansion of x also appear in the binary expansion of y.

Let

$$D := \{iq + k - i \mid 1 \le k \le q - 2, 0 \le i \le \lfloor (k - 1)/2 \rfloor, i \preceq k\}$$

Theorem 2. Let H be a star-set. Then H is a KM-arc of type t with t-nucleus at 0 if and only if

$$\pi_d(H) := \sum_{y \in H} y^d = 0,$$

for all $d \in D$.

Proof. Let $H = \{y_1, \dots, y_{q+t}\}$. Fix $1 \le k \le q-2$. For each $v \in S$, we have

$$\begin{split} \sum_{j=1}^{q+t} \left\langle v, y_j \right\rangle^k &= \sum_{j=1}^{q+t} \sum_{i=0}^{\lfloor (k-1)/2 \rfloor} \binom{k}{i} \left\langle v^{iq+k-i}, y_j^{iq+k-i} \right\rangle \\ &= \sum_{i=0}^{\lfloor (k-1)/2 \rfloor} \sum_{j=1}^{q+t} \binom{k}{i} \left\langle v^{iq+k-i}, y_j^{iq+k-i} \right\rangle \\ &= \sum_{i=0}^{\lfloor (k-1)/2 \rfloor} \binom{k}{i} \left\langle v^{iq+k-i}, \pi_{iq+k-i}(H) \right\rangle \\ &= \sum_{0 \le i \le \lfloor (k-1)/2 \rfloor} \left\langle v^{iq+k-i}, \pi_{iq+k-i}(H) \right\rangle = P_k(v), \end{split}$$

where P_k is the polynomial obtained from expanding the terms in the last sum. We can assume P_k has degree at most q, as $v^{q+1} = 1$.

The power of v in $P_k(v)$ is either of the form k - 2i or q + 1 + 2i - k. If $i \neq i'$, then $k - 2i \neq k - 2i'$, and $k - 2i \neq 2i' - k + q + 1$. Hence $P_k(v)$ has the form $P_k(v) = \sum \pi_l v^s$.

By Theorem 1, H is a KM-arc of type t with t-nucleus at 0 if and only if $P_k(v) = 0$ has q+1 roots $v \in S$ for each $1 \leq k \leq q-2$. Equivalently, the coefficients of P_k are zeros, that is, $\pi_d(H) = 0$ for all $d \in D$.

Now we introduce an analog of the set D, which contains more elements but looks more symmetric. Let

$$D' := \{k + (q-1)i \mid 1 \le k \le q-1, i \le k\},\$$
$$E := \left\{\sum_{j=0}^{m-1} 2^j x_j > 0 \mid x_j \in \{0, 1, q\}\right\}.$$

An element $x \in E$ can also be described in binary form as

$$x = \sum_{j=0}^{2m-1} 2^j x_j > 0,$$

where $x_j \in \{0, 1\}$ for $0 \le j \le 2m - 1$, and $(x_j, x_{m+j}) \ne (1, 1)$, for $0 \le j \le m - 1$. Lemma 5. D' = E.

Proof. We rewrite $k = \sum_{j=0}^{m-1} 2^j k_j$, where $k_j \in \{0,1\}$. For $i \leq k$, we rewrite

$$i = \sum_{j=0}^{m-1} 2^j i_j$$

where $i_j \leq k_j$. Let $d := k + (q-1)i \in D'$. Then

$$d = \sum_{j=0}^{m-1} 2^j (k_j + (q-1)i_j).$$

For each j, there are three possible cases for $k_j + (q-1)i_j$, as follows.

- 1. $k_j = 0, i_j = 0$. Then $k_j + (q-1)i_j = 0$.
- 2. $k_j = 1, i_j = 0$. Then $k_j + (q-1)i_j = 1$.
- 3. $k_i = 1, i_j = 1$. Then $k_i + (q-1)i_j = q$.

This implies $d \in E$ so that $D' \subseteq E$. Conversely, for each j, each element of $\{0, 1, q\}$ can be rewritten in the form of $k_j + (q-1)i_j$ as above, so that $E \subseteq D'$ and consequently D' = E.

Theorem 3. Let H be a star-set. Then H is a KM-arc of type t with t-nucleus at 0 if and only if $\pi_e(H) = 0$ for all $e \in E$.

Proof. We note that $D \subset D' = E$. By Theorem 2, if $\pi_e(H) = 0$ for all $e \in E$, then H is a KM-arc of type t with t-nucleus at 0.

Conversely, assume H is a KM-arc of type t with t-nucleus at 0. By Theorem 1, for each $v \in S$ and $1 \le k \le q - 2$,

$$\sum_{y \in H} \langle v, y \rangle^k = 0.$$
 (2)

For k = q - 1, it can be checked that condition (2) also holds. Using Lemma 4, for $1 \le k \le q - 1$,

$$\sum_{y \in H} \langle v, y \rangle^k = \sum_{\substack{i \leq k \\ 0 \leq i \leq \lfloor (k-1)/2 \rfloor}} \langle v^{iq+k-i}, \pi_{iq+k-i}(H) \rangle$$
$$= \sum_{\substack{i \leq k \\ \lceil (k+1)/2 \rceil \leq i \leq k}} \langle v^{iq+k-i}, \pi_{iq+k-i}(H) \rangle = 0.$$

Similar to the proof of Theorem 2, it follows that $\pi_d(H) = 0$ for all $d \in D'$. By Lemma 5, $\pi_e(H) = 0$ for all $e \in E$.

In homogeneous coordinates, KM-arcs are considered with the line at infinity L_{∞} as a *t*-secant. Under this setting, Gács and Weiner [11] proved that the set of points on L_{∞} of a KM-arc is a Vandermonde set. In the following, we extend this result in polar coordinates.

Proposition 1. For $t \ge 2$, let $\{y_i \mid 1 \le i \le t\} \subset K$ be the set of points on a t-secant of a KM-arc H with t-nucleus at 0. Then the set

$$\{1/y_i \mid 1 \le i \le t\}$$

is a Vandermonde set.

Proof. We identify each point $z = z_1 + z_2 \mathbf{i} \in K$ as $(z_1 : z_2 : 1)$ in homogeneous coordinates. We note that the Vandermonde property is invariant under affine transformations by Lemma 1. Under a suitable map $x \mapsto \alpha x, \alpha \in K$, we can assume that the x-axis is a t-secant of H. This means we can choose the set $\{y_i \mid 1 \leq i \leq t\}$ on the x-axis, so that $y_i \in F$ and each y_i is identified as $(y_i : 0 : 1)$.

Assume the line $x = \mu y$ is a t-secant of H. Let φ be the collineation defined by

$$\varphi((x:y:z)) = (x + \mu y:z:y).$$

Under φ , each point $(y_i : 0 : 1)$ is mapped to $(1 : 1/y_i : 0)$ and the *t*-nucleus (0 : 0 : 1) is mapped to (0 : 1 : 0). Also, the *t*-secant $x = \mu y$ is mapped to the *y*-axis. By [11, Proposition 2.4], the set $\{1/y_1, \dots, 1/y_t\}$ is a Vandermonde set.

4 KM-arcs from hyperovals and KM-arcs

In this section we construct KM-arcs starting from KM-arcs in smaller dimensions. Let $r = 2^h$, where $h \mid m$. Let $F' = \mathbb{F}_r$, and $K' = \mathbb{F}_{r^2}$. We denote the group of (r+1)-st roots of unity in K' by S'.

We recall the relative trace map $Tr_{F/F'}: F \to F'$, where

$$Tr_{F/F'}(x) = x^{q/r} + x^{q/r^2} + \dots + x^r + x.$$

Let $V_1 := \{x \in F \mid Tr_{F/F'}(x) = 1\}$. The main result of this section is the following.

Theorem 4. Assume m/h is odd. Let $H' \subset K'$ be a (r + s, s)-arc of type $s \ge 1$ with s-nucleus at 0. Let $V_c := cV_1$ for some $c \in K^*$. Then $H := \{\lambda u \mid 1/\lambda \in V_c, u \in H'\}$ is a (q + sq/r, sq/r)-arc in K of type t = sq/r with t-nucleus at 0.

In particular,

1) if H' is an oval with 1-nucleus at 0, then H is a (q + q/r, q/r)-arc of type t = q/r with t-nucleus at 0.

2) if H' is a hyperoval not containing 0, then H is a (q+2q/r, 2q/r)-arc of type t = 2q/r with t-nucleus at 0.

The proof of Theorem 4 is presented at the end of this section. In Lemmas 6 and 7 we examine special power sums of the sets H' and V_1 .

Let

$$\mathcal{A} := \left\{ \frac{q}{r^{j}} \mid 1 \le j \le m/h \right\} \cup \{0\} = \left\{ 1, r, r^{2}, \cdots, \frac{q}{r} \right\} \cup \{0\}.$$

Lemma 6. Let $k = \sum_{j=0}^{h-1} 2^j a_j > 0$, where $a_j \in \mathcal{A}$. Let d = k + (q-1)i, for some $i \leq k$. Then $\pi_d(H') = 0$.

Proof. Let

$$D'_r := \{ l + (r-1)i \mid 1 \le l \le r-1, i \le l \}$$

As in Lemma 5, D'_r can also be expressed as

$$E_r := \left\{ \sum_{j=0}^{h-1} 2^j x_j > 0 \mid x_j \in \{0, 1, r\} \right\}.$$

Let $i \leq k$. We rewrite $i = \sum_{j=0}^{h-1} 2^j i_j$, where $i_j \in \{0, a_j\}$. Then

$$d = k + (q-1)i = \sum_{j=0}^{h-1} 2^j \left(a_j + (q-1)i_j \right).$$

For each j, we note that $a_j + (q-1)i_j \in \{a_j, qa_j\}$. Since $q \pmod{r^2 - 1} \in \{1, r\}$ and $a_j \pmod{r^2 - 1} \in \{0, 1, r\}$, it follows that

$$a_j + (q-1)i_j \pmod{r^2 - 1} \in \{0, 1, r\}.$$

Then, for $u \in K'$, there exists $d' \in D'_r = E_r$ such that

$$u^d = u^{k+(q-1)i} = u^{d'}.$$

From Theorem 3,

$$\pi_d(H') = \pi_{d'}(H') = 0.$$

Lemma 7. Let $1 \le k \le q - 2$. If $\pi_k(V_1) \ne 0$, then

$$k = q - 1 - \sum_{j=0}^{h-1} 2^j a_j,$$

where $a_j \in \mathcal{A}$ for each $0 \leq j \leq h - 1$. In this case, $\pi_k(V_1) = 1$.

Proof. Rewrite k in the form

$$k = \frac{q}{r}l + s,$$

where $0 \le l \le r-1$, and $0 \le s \le q/r-1$. We consider three cases depending on l.

1. l = 0. Then $1 \le k \le q/r - 1$. Let $V_0 := \{x \in F \mid Tr_{F/\mathbb{F}_r}(x) = 0\}$, which is an additive subgroup of F and hence a Vandermonde set of size q/r. Since V_1 is a translation of V_0 , by Lemma 1, it is also a Vandermonde set of size q/r.

It follows that $\pi_k(V_1) \neq 0$ if and only if

$$k = \frac{q}{r} - 1 = q - 1 - \sum_{j=0}^{h-1} 2^j \frac{q}{r}.$$

For $x \in V_0 \setminus \{0\}$,

$$x^{q/r-1} = (x^{q/r^2} + \dots + x^r + x)x^{-1} = x^{q/r^2 - 1} + \dots + x^{r-1} + 1,$$

and so $\pi_{q/r-1}(V_0) = 1$. Note that the set $V_0 \cup V_1$ is a Vandermonde set of size 2q/r, so that $\pi_{q/r-1}(V_0 \cup V_1) = 0$. Therefore, $\pi_{q/r-1}(V_1) = \pi_{q/r-1}(V_0) = 1$.

2. l = 1. Then k = q/r + s, and for $x \in V_1$, we have

$$x^{k} = x^{q/r}x^{s} = \left(x^{q/r^{2}} + \dots + x + 1\right)x^{s}.$$

To evaluate the sum $\sum_{x \in V_1} x^k$, we consider three cases of each sum $\sum_{x \in V_1} x^{q/r^i+s}$, depending on $2 \le i \le m/h$.

(a) $q/r^i + s < q/r - 1$. Then $\sum_{x \in V_1} x^{q/r^i + s} = 0$.

(b)
$$q/r^{i} + s = q/r - 1$$
. Then $\sum_{x \in V_{1}} x^{q/r^{i} + s} = 1$.
(c) $q/r^{i} + s > q/r - 1$. Rewrite $\frac{q}{r^{i}} + s = \frac{q}{r} + s_{0}$, where $0 \le s_{0} \le \frac{q}{r^{i}} - 1$. Then $x^{q/r^{i} + s} = x^{q/r + s_{0}} = \left(x^{q/r^{2}} + \dots + x + 1\right) x^{s_{0}}$,

and since for each $2 \leq i' \leq m/h$,

$$\frac{q}{r^{i'}} + s_0 \le \frac{q}{r^{i'}} + \frac{q}{r^i} - 1 < \frac{q}{r} - 1,$$

it follows that

$$\sum_{x \in V_1} x^{q/r^i + s} = 0.$$

Therefore $\pi_k(V_1) \neq 0$ if and only if $\pi_k(V_1) = 1$. This occurs when

$$s = \frac{q}{r} - \frac{q}{r^i} - 1,$$

if and only if

$$k = 2\frac{q}{r} - \frac{q}{r^{i}} - 1 = q - 1 - \left(\sum_{j=1}^{h-1} 2^{j}\frac{q}{r} + \frac{q}{r^{i}}\right).$$

3. l > 1. For $x \in V_1$, we have

$$x^{k} = x^{lq/r}x^{s} = \left(x^{q/r^{2}} + \dots + 1\right)^{l}x^{s}.$$

Let x^c be a term of the expansion of the above. Rewrite l as the binary expansion $l = \sum_{j=0}^{h-1} 2^j l_j$, where $l_j \in \{0, 1\}$, for $0 \le j \le h-1$. Then

$$c = \sum_{j=0}^{h-1} 2^{j} l_{j} a_{j} + s_{j}$$

where $a_j \in \mathcal{A} \setminus \left\{ \frac{q}{r} \right\}$. From the conditions on l and s, we have

$$c \le \frac{q}{r^2}l + s \le \frac{q}{r^2}(r-1) + \frac{q}{r} - 1 = 2\frac{q}{r} - \frac{q}{r^2} - 1.$$

Furthermore, $c \neq 2\frac{q}{r} - \frac{q}{r^2} - 1$, otherwise l = r - 1, $s = \frac{q}{r} - 1$ and k = q - 1, which is excluded from our assumption. From part 1) and part 2), $\pi_c(V_1) \neq 0$ if and only if $\pi_c(V_1) = 1$. This occurs when c = q/r - 1, if and only if

$$s = \frac{q}{r} - 1 - \sum_{j=0}^{h-1} 2^j l_j a_j,$$

if and only if

$$\begin{split} k &= \frac{q}{r}(l+1) - 1 - \sum_{j=0}^{h-1} 2^{j} l_{j} a_{j} \\ &= \frac{q}{r} - 1 + \sum_{j=0}^{h-1} 2^{j} l_{j} \frac{q}{r} - \sum_{j=0}^{h-1} 2^{j} l_{j} a_{j} \\ &= \frac{q}{r} - 1 + \sum_{j=0}^{h-1} 2^{j} \frac{q}{r} - \sum_{j=0}^{h-1} 2^{j} (1 - l_{j}) \frac{q}{r} - \sum_{j=0}^{h-1} 2^{j} l_{j} a_{j} \\ &= q - 1 - \sum_{j=0}^{h-1} 2^{j} \left[(1 - l_{j}) \frac{q}{r} + l_{j} a_{j} \right] \\ &= q - 1 - \sum_{j=0}^{h-1} 2^{j} a_{j}', \end{split}$$

where $a'_i \in \mathcal{A}$.

The proof now follows.

Proof of Theorem 4. We will assume that c = 1 since multiplication by $c \in K^*$ is a collineation of PG(2,q).

1. Let x, y be two points from H' such that the lines xF' and yF' are distinct in K'. This implies $x/y \notin F'$. But since m/h is odd, we have $F \cap K' = F'$. Hence $x/y \notin F$, and so the lines xF and yF are distinct in K.

Assume $s \geq 2$. For $x, y \in F', x \neq y$, we consider two distinct points $x\alpha, y\alpha$ from the same s-secant $\alpha F'$ of H' in K'. Let $\lambda_1, \lambda_2 \in V_1$. Since $Tr_{F/F'}(\lambda_2 x) = x, Tr_{F/F'}(\lambda_1 y) = y$, it follows that $\lambda_2 x \neq \lambda_1 y$, and so

$$\frac{x\alpha}{\lambda_1} \neq \frac{y\alpha}{\lambda_2}$$

This implies the line αF in K contains t = sq/r distinct points of H. Hence, H is a star-set.

2. Let $d = k + (q-1)i \in D'$. For each $\lambda \in F$, we have

$$1/\lambda^d = \lambda^{q-1-d} = \lambda^{q-1-k},$$

so that

$$\pi_d(H) = \pi_d(H') \sum_{1/\lambda \in V_1} \lambda^d = \pi_{q-1-k}(V_1)\pi_d(H').$$

From Lemma 7, if $\pi_{q-1-k}(V_1) \neq 0$, then

$$k = \sum_{i=0}^{h-1} 2^i a_i,$$

where $a_i \in \mathcal{A}$ for each $0 \leq i \leq h-1$. By Lemma 6, $\pi_d(H') = 0$. This implies $\pi_d(H) = 0$, and by Theorem 3, H is a KM-arc of type t with t-nucleus at 0.

Now we show that our construction is equivalent to the construction of Gács and Weiner [11]. Recall that they constructed a family of KM-arcs as follows. Let I be a direct complement of F' in the additive group of F. Let H_0 be a KM-arc of type $s \ge 1$ with affine part $\{(x_k : y_k : 1) \mid x_k, y_k \in F'\}$. Construct the following point set

$$J := \{ (x_k : y_k + i : 1) \mid (x_k : y_k : 1) \in H_0, i \in I \}.$$

Then J can be uniquely extended to a KM-arc of type t = sq/r in PG(2,q).

Now we consider a collineation γ defined by

$$\gamma((a:b:c)) = (a:c:a+b+cd),$$

where $d \in F'$ is chosen such that the line x + y + zd = 0 does not intersect H_0 , so $x_k + y_k + d \neq 0$. Under γ , the point $(x_k : y_k : 1)$ of H_0 is mapped to $\left(\frac{x_k}{x_k + y_k + d} : \frac{1}{x_k + y_k + d} : 1\right)$, which corresponds to the element

$$p_k = \frac{x_k}{x_k + y_k + d} + \frac{1}{x_k + y_k + d} \mathbf{i} \in K.$$

Also, the point $(x_k : y_k + i : 1)$ of J is mapped to

$$\frac{x_k}{x_k + y_k + d + i} + \frac{1}{x_k + y_k + d + i}\mathbf{i} = \frac{p_k}{1 + i(x_k + y_k + d)^{-1}} \in \frac{p_k}{1 + I}$$

Therefore,

$$\gamma(J) = \left\{\frac{p_k}{1+I}\right\}$$

On the other hand, if we consider F as a vector space over F' then I is a subspace of codimension 1, not containing F'. Hence, 1 + I is a coset, such that $(1 + I) \cap F' = \{1\}$. Further, we can choose $b \in K^*$ such that bV_1 is parallel to I (note that $V_0 := \{x \in F \mid Tr_{F/F'}(x) = 0\}$ is a subspace in F of codimension 1, and bV_0 , $b \in K^*$, runs the set of all subspaces of codimension 1 in F). Let $bV_1 \cap F' = a$. Then $a^{-1}bV_1 \cap F' = 1$ and $cV_1 \cap F' = 1$ for $c = a^{-1}b$. Hence $cV_1 = 1 + I$. It shows the equivalence of our construction to the construction of Gács and Weiner.

5 A family of KM-arcs type t = q/r

We present now some other examples of KM-arcs.

5.1 The construction

Let $r = 2^h$, where $h \mid m$. Let $F' = \mathbb{F}_r$, and $K' = \mathbb{F}_{r^2}$. We denote the group of (r + 1)-st roots of unity in K' by S'. Let \mathbf{u} be a generator of S' whose conjugate over F' is $\tilde{\mathbf{u}} = \mathbf{u}^r$. Let $b = \mathbf{u} + \tilde{\mathbf{u}}$ be the trace of \mathbf{u} over F' and (b_n) be the sequence in F' defined by the recurrence relation $b_{n+1} = bb_n + b_{n-1}$, with initial terms $b_0 = 1, b_1 = 0$. Explicitly, for each n,

$$b_n = \frac{\mathbf{u}^{n-1} + \tilde{\mathbf{u}}^{n-1}}{\mathbf{u} + \tilde{\mathbf{u}}}$$

Let $U := \{u_i \mid 0 \le i \le r\}$, where $u_i = b_i + b_{i+1}\mathbf{i}$, for each *i*. Alternatively, if we identify each point $z = x + y\mathbf{i}$ as $[x \ y]^T$, then elements of *U* are obtained recursively as

$$u_{i+1} = \begin{bmatrix} 0 & 1\\ 1 & b \end{bmatrix} u_i$$

with the initial term $u_0 = 1 = \begin{bmatrix} 1 & 0 \end{bmatrix}^T$. We also recall the relative trace map

$$Tr_{F/\mathbb{F}_r}(x) = x^{q/r} + x^{q/r^2} + \dots + x^r + x.$$

Define $V_1 := \{x \in F \mid Tr_{F/\mathbb{F}_r}(x) = 1\}$ and $V_c := cV_1$ for $c \in K^*$. In this subsection we prove the following theorem.

Theorem 5. Let $c \in K^*$. The set $H_r := \{\lambda u \mid 1/\lambda \in V_c, u \in U\}$ is a KM-arc of type t = q/r with t-nucleus at 0 in K.

The proof of Theorem 5 is presented at the end of the subsection.

Lemma 8. The set U has size r + 1.

Proof. It is sufficient to prove that the sequence (b_n) has period r+1. We have

$$b_{r+1} = \frac{\mathbf{u}^r + \tilde{\mathbf{u}}^r}{\mathbf{u} + \tilde{\mathbf{u}}} = 1,$$
$$b_{r+2} = \frac{\mathbf{u}^{r+1} + \tilde{\mathbf{u}}^{r+1}}{\mathbf{u} + \tilde{\mathbf{u}}} = 0$$

so that the period of (b_n) divides r + 1. On the other hand, $b_n = 0$ if and only if $\mathbf{u}^{n-1} = \tilde{\mathbf{u}}^{n-1}$ if and only if $\mathbf{u}^{2(n-1)} = 1$. This occurs if and only if r + 1 divides n - 1, which implies b_{r+2} is the smallest zero term of the sequence excluding b_1 . Therefore (b_n) has period r + 1 and consequently |U| = r + 1.

Let $U' := \{u_i \mid 0 \le i \le r\}$, where $u_i = b_i + b_{i+1}\mathbf{u}$, for each *i*. It can be checked that elements of U' are pairwise distinct and of order r + 1, so that U' = S'. This implies the following.

Lemma 9. The set U can be written as

$$U = \{x + y\mathbf{i} \mid x, y \in F', x^2 + y^2 + bxy = 1\}.$$

We also recall from Section 4 the set

$$\mathcal{A} = \left\{ \frac{q}{r^j} \mid 1 \le j \le m/h \right\} \cup \{0\}.$$

Lemma 10. Let $k = \sum_{j=0}^{h-1} 2^j a_j \in \{1, \cdots, q-2\}$, where $a_j \in \mathcal{A}$. Then for each $v \in S$,

$$\sum_{u \in U} \langle v, u \rangle^k = 0$$

Proof. 1. Let

$$\mathcal{B} := \left\{ \sum_{j=0}^{h-1} 2^j x_j \mid x_j \in \{0, a_j\} \right\}.$$

We claim that for each $x \in \mathcal{B}$,

$$\sum_{i=0}^{r} b_i^x b_{i+1}^{k-x} = 0.$$

Fix $x \in \mathcal{B}$. Let

$$y := k - x = \sum_{j=0}^{h-1} 2^j y_j,$$

where $y_j := a_j - x_j$ for each $0 \le j \le h - 1$. We also introduce the sets

$$\mathcal{B}_x := \left\{ \sum_{j=0}^{h-1} \pm 2^j x_j \right\}, \text{ and } \mathcal{B}_y := \left\{ \sum_{j=0}^{h-1} \pm 2^j y_j \right\}.$$

For each $0 \leq i \leq r$, we have

$$b_i^x = \frac{(\mathbf{u}^{i-1} + \tilde{\mathbf{u}}^{i-1})^x}{(\mathbf{u} + \tilde{\mathbf{u}})^x} = \frac{\prod_{j=0}^{h-1} (\mathbf{u}^{(i-1)2^j x_j} + \tilde{\mathbf{u}}^{(i-1)2^j x_j})}{(\mathbf{u} + \tilde{\mathbf{u}})^x} = \frac{\sum_{x' \in \mathcal{B}_x} \mathbf{u}^{(i-1)x'}}{(\mathbf{u} + \tilde{\mathbf{u}})^x}.$$

Similarly,

$$b_{i+1}^y = \frac{(\mathbf{u}^i + \tilde{\mathbf{u}}^i)^y}{(\mathbf{u} + \tilde{\mathbf{u}})^y} = \frac{\prod_{j=0}^{h-1} (\mathbf{u}^{i2^j y_j} + \tilde{\mathbf{u}}^{i2^j y_j})}{(\mathbf{u} + \tilde{\mathbf{u}})^y} = \frac{\sum_{y' \in \mathcal{B}_y} \mathbf{u}^{iy'}}{(\mathbf{u} + \tilde{\mathbf{u}})^y}.$$

Then

$$\sum_{i=0}^{r} b_i^x b_{i+1}^y = 0 \iff \sum_{i=0}^{r} \left(\sum_{x' \in \mathcal{B}_x} \mathbf{u}^{(i-1)x'} \right) \left(\sum_{y' \in \mathcal{B}_y} \mathbf{u}^{iy'} \right) = 0$$
$$\iff \sum_{i=0}^{r} \sum_{\substack{x' \in \mathcal{B}_x \\ y' \in \mathcal{B}_y}} \mathbf{u}^{i(x'+y')-x'} = \sum_{\substack{x' \in \mathcal{B}_x \\ y' \in \mathcal{B}_y}} \sum_{i=0}^{r} \mathbf{u}^{i(x'+y')-x'} = 0$$

We now consider the sum

$$x' + y' = \sum_{j=0}^{h-1} 2^j (\pm x_j \pm y_j)$$

For each j, we have $x_j + y_j = a_j$ and $x_j - y_j = 2x_j - a_j \in \{a_j, -a_j\}$, so that

$$\pm x_j \pm y_j \in \{a_j, -a_j\}$$

We note that $a_j \pmod{r+1} \in \{-1, 0, 1\}$. Also, x' + y' = 0 implies $a_j = 0$ for any j, which is impossible. It follows that $x' + y' \neq 0 \pmod{r+1}$. Therefore $\mathbf{u}^{x'+y'}$ generates a subgroup of S' of order $\frac{r+1}{\gcd(x'+y',r+1)} \neq 1$. Consequently,

$$\sum_{i=0}^{r} \mathbf{u}^{i(x'+y')-x'} = \mathbf{u}^{-x'} \sum_{i=0}^{r} \mathbf{u}^{i(x'+y')} = 0.$$

This proves $\sum_{i=0}^{r} b_i^x b_{i+1}^{k-x} = 0.$

2. To simplify the notation, let $A = \langle v, 1 \rangle, B = \langle v, \mathbf{i} \rangle$. For fixed $0 \le i \le r$, we have

$$\langle v, u_i \rangle^k = \langle v, b_i + b_{i+1} \mathbf{i} \rangle^k = (b_i A + b_{i+1} B)^{\sum_j 2^j a_j}$$
$$= \prod_{j=0}^{h-1} (b_i^{2^j a_j} A^{2^j a_j} + b_{i+1}^{2^j a_j} B^{2^j a_j}) = \sum_{x \in \mathcal{B}} b_i^x b_{i+1}^{k-x} A^x B^{k-x}$$

From part 1),

$$\sum_{u \in U} \langle v, u \rangle^k = \sum_{i=0}^r \sum_{x \in \mathcal{B}} b_i^x b_{i+1}^{k-x} A^x B^{k-x} = \sum_{x \in \mathcal{B}} \left(\sum_{i=0}^r b_i^x b_{i+1}^{k-x} \right) A^x B^{k-x} = 0.$$

Proof of Theorem 5. We will assume that c = 1 since multiplication by $c \in K^*$ is a collineation of PG(2,q).

- 1. Let $x = b_i + b_{i+1}\mathbf{i}$, $y = b_j + b_{j+1}\mathbf{i} \in U$, with $0 \le i, j \le r 1$, $i \ne j$. Suppose that the lines xF and yF are the same. Since the matrix $\begin{bmatrix} 0 & 1 \\ 1 & b \end{bmatrix}$ permutes elements of U cyclically, we may assume that x = 1. Then $b_{j+1} = 0$ and y = 1, a contradiction. This implies for $x, y \in U$, the lines xF and yF intersect at 0 only. Hence, H_r is a star-set.
- 2. Let $d = k + (q-1)i \in D'$. We have

$$\pi_d(H) = \pi_d(U) \sum_{1/\lambda \in V_1} \lambda^d = \pi_{q-1-k}(V_1)\pi_d(U).$$

From Lemma 7, if $\pi_{q-1-k}(V_1) \neq 0$, then

$$k = \sum_{i=0}^{h-1} 2^i a_i$$

where $a_i \in \mathcal{A}$ for each $0 \leq i \leq h-1$. From Lemma 10 and by the method used in the proof of Theorem 3, we have $\pi_d(U) = 0$. This implies $\pi_d(H_r) = 0$, and by Theorem 3, H_r is a KM-arc of type t = q/r with t-nucleus at 0.

5.2 Automorphisms of H_r

In this section, we discuss some properties of the automorphism group of KM-arcs H_r and show that they are translation KM-arcs.

An *elation* with axis the line l and centre the point R on l is a collineation which fixes the points of l and stabilises the lines through the centre R. The set of all elations with a fixed centre and a fixed axis form a subgroup of $P\Gamma L(3, q)$.

Let H be a KM-arc of type t > 2 with t-nucleus at N. Then H is an *elation KM-arc* with elation line l_0 if for every t-secant $l \neq l_0$ to H, the group of elations with axis l_0 that stabilise H acts transitively on the points of $l \cap H$. Furthermore, H is a *translation* KM-arc with translation line l_0 if the group of all elations with axis l_0 that stabilise H acts transitively on the points of $l \cap H$. Furthermore, and h arc stabilise H acts transitively on the group of all elations with axis l_0 that stabilise H acts transitively on the points of $H \setminus l_0$. In particular, every translation KM-arc is an elation KM-arc.

For $x, y \in F$, let $z = x + y\mathbf{i} \in K$. Let l_0 be the line $\langle 1, z \rangle = 0$. We consider several maps from the automorphism group $P\Gamma L(3,q)$ of PG(2,q).

1. We define θ as a map induced by the matrix

0	1	0	
1	b	0	;
0	0	1	

which generates the cyclic group C_{r+1} .

2. The map σ' is defined by

$$\sigma' := \begin{bmatrix} 1 & \delta + 1 \\ 0 & b \end{bmatrix} \sigma,$$

where $\sigma: z \mapsto z^2$ is a generator of $Gal(K/\mathbb{F}_2)$. Explicitly,

$$\sigma'(z) = \sigma'(x+y\mathbf{i}) = x^2 + y^2 + by^2\mathbf{i}.$$

Note that $(\sigma')^m \in GL(2,q)$.

3. Let $E_{a,b}$ be the map induced by the matrix

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ a & b & 1 \end{bmatrix},$$

where $a, b \in F$. Explicitly,

$$E_{a,b}(z) = \frac{z}{ax + by + 1} = \frac{z}{\langle z, b + a\mathbf{i} \rangle + 1}.$$

We note that $E_{a,b}$ is an elation with centre 0 and axis $\langle b + a\mathbf{i}, z \rangle = 0$. The group

$$\mathcal{E} = \{ E_{a,b} \mid Tr_{F/F'}(a) = Tr_{F/F'}(b) = 0 \}$$

is a group of the automorphisms of H_r that fixes every t-secant and acts transitively on the points of each t-secant of H_r . Then the group $\{E_{0,b} \mid Tr_{F/F'}(b) = 0\}$ fixes all points of l_0 and acts transitively on the points $H_r \cap l$ on any t-secant $l \neq l_0$. This shows that H_r is an elation KM-arc with elation line l_0 .

4. Let ψ be the map induced by the matrix

$$\begin{bmatrix} 1 & b & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Explicitly, $\psi(z) := x + by + y\mathbf{i}$. We note that ψ is an elation with centre at infinity and axis l_0 . Also, ψ has order 2.

5. Let $\rho_{\gamma,s,t}$ be the map induced by the matrix

$$\begin{bmatrix} \gamma & 0 & 0 \\ 0 & \gamma^{-1} & 0 \\ s & t & 1 \end{bmatrix},$$

where $\gamma \in F'$, $s, t \in F$, $Tr_{F/F'}(s) = \gamma + 1$, $Tr_{F/F'}(t) = \gamma^{-1} + 1$. We note that

$$\begin{bmatrix} \gamma & 0 & 0 \\ 0 & \gamma^{-1} & 0 \\ s & t & 1 \end{bmatrix}^k = \begin{bmatrix} \gamma^k & 0 & 0 \\ 0 & \gamma^{-k} & 0 \\ s_k & t_k & 1 \end{bmatrix},$$

where $s_k = (\gamma^{k-1} + \cdots 1)s$, and $t_k = (\gamma^{-(k-1)} + \cdots 1)t$. And so, if γ is a primitive element of F', then $\rho_{\gamma,s,t}$ has order r-1.

Theorem 6. H_r is a translation KM-arc with translation line l_0 .

Proof. 1. We first show that the map ψ is an automorphism of H_r . Since ψ is linear on F, it is sufficient to show that ψ fixes the set U. We recall from Lemma 9 that

$$U = \{x + y\mathbf{i} \mid x, y \in F', x^2 + y^2 + bxy = 1\}.$$

Let $u = x + y\mathbf{i} \in U$ with $\psi(u) = x' + y'\mathbf{i} = x + by + y\mathbf{i}$. Then

$$x'^{2} + y'^{2} + bx'y' = (x + by)^{2} + y^{2} + b(x + by)y$$

= $x^{2} + y^{2} + bxy = 1$,

which shows that $\psi(u) \in U$, and the claim follows. The action of ψ on H_r can then be described by its action on U. Since ψ has order 2, it has one fixed point (which is 1) and r/2 orbits of size 2.

2. We now show that the map $\rho_{\gamma,s,t}$ is an automorphism of H_r . Let $p = \lambda u$ be a point on H_r , where $u = x + y\mathbf{i} \in U$, and $\lambda \in F$ with $1/\lambda \in V_1$. We have

$$\rho_{\gamma,s,t}(\lambda u) = \frac{\lambda}{B}\gamma x + \frac{\lambda}{B}\gamma^{-1}y\mathbf{i},$$

where $B = s\lambda x + t\lambda y + 1$. We rewrite

$$\rho_{\gamma,s,t}(\lambda u) = \frac{\lambda}{AB} (A\gamma x) + \frac{\lambda}{AB} (A\gamma^{-1} y) \mathbf{i}$$
$$= \frac{\lambda}{AB} (x' + y' \mathbf{i}),$$

where $x' = A\gamma x, y' = A\gamma^{-1}y$ and A satisfies

$$\frac{1}{A^2} = \gamma^2 x^2 + \gamma^{-2} y^2 + bxy.$$

With this choice of A, we have

$$x'^2 + y'^2 + bx'y' = 1,$$

so that $x' + y'\mathbf{i} \in U$. Note that $A \in F'$, and so

$$Tr_{F/F'}\left(\frac{AB}{\lambda}\right) = A^2 Tr_{F/F'}\left(\frac{B^2}{\lambda^2}\right)$$

= $A^2 Tr_{F/F'}\left(\frac{s^2\lambda^2 x^2 + t^2\lambda^2 y^2 + 1}{\lambda^2}\right)$
= $A^2 Tr_{F/F'}\left(s^2 x^2 + t^2 y^2 + \frac{1}{\lambda^2}\right)$
= $A^2 \left(x^2 Tr_{F/F'}(s^2) + y^2 Tr_{F/F'}(t^2) + 1\right)$
= $A^2 (x^2 (\gamma^2 + 1) + y^2 (\gamma^{-2} + 1) + 1)$
= $A^2 (\gamma^2 x^2 + \gamma^{-2} y^2 + bxy) = 1,$

which shows that $\rho_{\gamma,s,t}(p) \in H_r$.

3. We consider a map $\rho := \rho_{\gamma,s,t}$, where γ is a primitive element of F'. In the following, we describe how ρ permutes the *t*-secants of H_r . Using the calculations above, we see that for $u = x + y\mathbf{i} \in U$, the *t*-secant uF is mapped to u'F, where $u' = x' + y'\mathbf{i} = A\gamma x + A\gamma^{-1}y\mathbf{i}$. It follows that ρ^k maps uF to vF, where

$$v = A^k \gamma^k x + A^k \gamma^{-k} y \mathbf{i}.$$

This implies that ρ^k fixes the *t*-secant uF if and only if $u \in \{1, i\}$ or $A^k \gamma^k = A^k \gamma^{-k} = 1$. The latter case implies that $\gamma^{2k} = 1$, which occurs if and only if k = r - 1. Hence, ρ fixes the *t*-secants l_0 and $\mathbf{i}F$, and $\langle \rho \rangle$ is transitive on the set of (r-1) remaining *t*-secants of H_r .

4. Let l be a t-secant of H_r different from $\mathbf{i}F$ and l_0 . For a suitable $k \in \{1, \ldots, r-1\}$, the map $\rho^k \psi \rho^{-k}$ is an elation that fixes l_0 and maps the line $\mathbf{i}F$ to l. Since $\rho^k \psi \rho^{-k}$ has order 2, it maps l to the line $\mathbf{i}F$. It follows that the group of automorphisms

$$\langle \rho^k \psi \rho^{-k} \mid k \in \{1, \dots, r-1\} \rangle$$

is a group of elations with axis l_0 which is transitive on the set of t-secants $H \setminus l_0$. Since H_r is an elation KM-arc with elation line l_0 , it follows that H_r is a translation KM-arc with translation line l_0 .

We do not know if our general construction of KM-arcs H_r is different from the one constructed by Korchmaros-Mazzocca, although their parameters look the same. Anyway, we see that some of the known KM-arcs can be easily described by polar coordinates.

5.3 Examples

We describe some examples of KM-arcs from the construction in Subsection 5.1. As before, let $F = \mathbb{F}_q$ and $K = \mathbb{F}_{q^2}$. We will also make use of the maps introduced in Subsection 5.2 and describe the full automorphism group of these examples.

Example 1 (KM-arcs of type q/2). Let r = 2, $V_1 = \{x \in F \mid Tr_{F/\mathbb{F}_2}(x) = 1\}$, and $U = \{1, \mathbf{i}, \mathbf{i} + 1\}$. Then $H_2 := \{\lambda u \mid 1/\lambda \in V_1, u \in U\}$ is a KM-arc of type q/2. According to [16, Corollary 4.4], all KM-arcs of type q/2 are projectively equivalent and they are also known as *projective triads*. The automorphism group of H_2 is

$$G_2 = (\mathcal{E}.\langle \theta, \tau \rangle).\langle \sigma' \rangle \cong (\mathcal{E}.S_3).\langle \sigma' \rangle,$$

where $\tau(x) = \bar{x}$ is the conjugation in K and S_3 is the symmetric group.

Example 2 (KM-arcs of type q/4). Let r = 4 and $V_1 = \{x \in F \mid Tr_{F/\mathbb{F}_4}(x) = 1\}$. Let $\omega \neq 1$ be an element in F such that $\omega^3 = 1$. Define

$$U := \{1, \mathbf{i}, 1 + \omega \mathbf{i}, \omega + \mathbf{i}, \omega + \omega \mathbf{i}\}.$$

Then $H_4 := \{\lambda u \mid 1/\lambda \in V_1, u \in U\}$ is a KM-arc of type q/4. In [9], it was proved that elation KM-arcs of type q/4 are translation KM-arcs and the full classification was provided. By [9, Theorem 3.12], H_4 is projectively equivalent to the one with parameters described therein. The automorphism group of H_4 is

$$G_4 = (\mathcal{E}.\langle \theta, \tau \rangle).\langle \sigma' \rangle \cong (\mathcal{E}.A_5).\langle \sigma' \rangle,$$

where the map τ induced by the matrix

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & t & 1 \end{bmatrix},$$

with $t \in F$ such that $Tr_{F/F'}(t) = \omega$, and A_5 is the alternating group.

Example 3 (KM-arcs of type q/8). Let r = 8 and $V_1 = \{x \in F \mid Tr_{F/\mathbb{F}_8}(x) = 1\}$. Let η be an element of \mathbb{F}_8 satisfying $\eta^3 + \eta + 1 = 0$. Define

$$U := \{1, \mathbf{i}, 1 + \eta \mathbf{i}, \eta + \eta^6 \mathbf{i}, \eta^6 + \eta^3 \mathbf{i}, \eta^3 + \eta^3 \mathbf{i}, \eta^3 + \eta^6 \mathbf{i}, \eta^6 + \eta \mathbf{i}, \eta + \mathbf{i}\}$$

Then $H_8 := \{\lambda u \mid 1/\lambda \in V_1, u \in U\}$ is a KM-arc of type q/8. Using GAP, we calculated that the automorphism group of H_8 is

$$G_8 = (\mathcal{E}.\langle \theta, \tau \rangle).\langle \sigma' \rangle,$$

where the map τ is induced by the matrix

$$\begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 0 \\ 0 & t & 1 \end{bmatrix},$$

with $t \in F$ such that $Tr_{F/F'}(t) = \eta^5$, and the group $(\mathcal{E} \cdot \langle \theta, \tau \rangle)/\mathcal{E}$ has order 504. Note that the automorphism τ is the conjugation in K corrected by some element $E_{0,b}$.

Acknowledgments

This research was supported by UAEU grant 31S366.

References

- K. Abdukhalikov, Bent functions and line ovals, Finite Fields Appl. 47, 94–124 (2017).
- [2] K. Abdukhalikov, Hyperovals and bent functions. European J. Combin. 79, 123–139 (2019).

- [3] K. Abdukhalikov, Short description of the Lunelli-Sce hyperoval and its automorphism group, Journal of Geometry (2019) 110:54.
- [4] K. Abdukhalikov, Equivalence classes of Niho bent functions. Des. Codes Cryptogr. (2021). https://doi.org/10.1007/s10623-021-00885-5
- [5] K. Abdukhalikov and Duy Ho, Vandermonde sets and hyperovals, arXiv:1911.10798.
- [6] S. Ball, Polynomials in finite geometries, Surveys in combinatorics, 1999 (Canterbury), 17–35, in: London Math. Soc. Lecture Note Ser., 267, Cambridge Univ. Press, Cambridge (1999).
- [7] A. Blokhuis, G. Marino, F. Mazzocca and O. Polverino, On almost small and almost large super-Vandermonde sets in GF(q), Des. Codes Cryptogr. 84, 197–201 (2017).
- [8] M. De Boeck and G. Van de Voorde, A linear set view on KM-arcs, J. Algebraic Combin. 44, 131–164 (2016).
- [9] M. De Boeck and G. Van de Voorde, Elation KM-arcs, Combinatorica 39, 501–544 (2019).
- [10] J. C. Fisher and B. Schmidt, Finite Fourier series and ovals in $PG(2, 2^h)$, J. Aust. Math. Soc. 81, no. 1, 21–34 (2006).
- [11] A. Gács and Z. Weiner, On (q + t, t)-arcs of type (0, 2, t), Des. Codes Cryptogr. 29, no 1-3, 131–139 (2003).
- [12] J. W. P. Hirschfeld, Projective geometries over finite fields. Second edition. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York (1998).
- [13] G. Korchmáros and F. Mazzocca, On (q + t)-arcs of type (0, 2, t) in a Desarguesian plane of order q, Math. Proc. Cambridge Philos. Soc. 108, no 3, 445–459 (1990).
- [14] R. Lidl and H. Niederreiter, Finite Fields. Cambridge University Press (1997).
- [15] P. Sziklai and M. Takáts, Vandermonde sets and super-Vandermonde sets, Finite Fields Appl. 14, 1056–1067 (2008).
- [16] P. Vandendriessche, Codes of Desarguesian projective planes of even order, projective triads and (q + t, t)-arcs of type (0, 2, t), Finite Fields Appl. 17, 521–531 (2011).
- [17] P. Vandendriessche, On KM-arcs in small Desarguesian planes, Electron. J. Combin. 24, no 1, Paper 1.51, 11 (2017).