

A note on the action of Hecke groups on subsets of quadratic fields

Mircea Cimpoeaş

Abstract

We study the action of the groups $H(\lambda)$ generated by the linear fractional transformations $x : z \mapsto -\frac{1}{z}$ and $w : z \mapsto z + \lambda$, where λ is a positive integer, on the subsets $\mathbb{Q}^*(\sqrt{n}) = \left\{ \frac{a+\sqrt{n}}{c} \mid a, b = \frac{a^2-n}{c}, c \in \mathbb{Z} \right\}$, where n is a square-free integer. We prove that this action has a finite number of orbits if and only if $\lambda = 1$ or $\lambda = 2$, and we give an upper bound for the number of orbits for $\lambda = 2$.

Keywords: Quadratic field, Hecke groups, Orbit

2020 Mathematics Subject Classification: 05A18; 05E18; 11A25; 11R11; 20F05

Introduction

E. Hecke introduced in [2] the groups $H(\lambda)$ generated by two linear fractional transformations $x : z \mapsto -\frac{1}{z}$ and $w : z \mapsto z + \lambda$, where λ is positive real number. He further showed that $H(\lambda)$ is discrete if and only if $\lambda = \lambda_k = 2 \cos\left(\frac{\pi}{k}\right)$, $k \geq 3$ an integer, or $\lambda \geq 2$. If $\lambda = \lambda_3 = 1$, then $H(1) = PSL(2, \mathbb{Z})$ is the modular group which consists in all the transformations $z \mapsto \frac{az+b}{cz+d}$ with $a, b, c, d \in \mathbb{Z}$ and $ac - bd = 1$.

The actions of the modular group on many discrete and non-discrete structures play significant roles in different branches of mathematics. Among these structures upon which the modular group acts are subsets of quadratic number fields. In [4], Q. Mushtaq studied the action of $PSL(2, \mathbb{Z})$ on the subset

$$\mathbb{Q}^*(\sqrt{n}) = \left\{ \frac{a + \sqrt{n}}{c} \mid a, b = \frac{a^2 - n}{c}, c \in \mathbb{Z} \right\},$$

where $n \geq 2$ is a square-free integer. Subsequent works by several authors considered properties emerging from this action; see for instance [5] and [3]. Recently, M. Aslam and A. Deajim [1] studied the action of $PSL(2, \mathbb{Z})$ on $\mathbb{Q}^*(\sqrt{n})$, when n is a negative square-free integer, and give a formula for the number of orbits of this action.

The aim of our paper is to study the action of the groups $H(\lambda)$, where $\lambda \geq 1$ is an integer, on the subsets $\mathbb{Q}^*(\sqrt{n})$, where n is a square-free integer. We prove that the action of $H(\lambda)$ on $\mathbb{Q}^*(\sqrt{n})$ has a finite number of orbits if and only if $\lambda \in \{1, 2\}$; see Theorem 1.4. For $\lambda = 2$ we give an upper bound for the number of orbits; see Theorem 1.5.

1 Main results

We fix n a square-free integer and we consider the sets:

$$\mathbb{Q}^*(\sqrt{n}) = \left\{ \alpha = \frac{a + \sqrt{n}}{c} : a, b = \frac{a^2 - n}{c}, c \in \mathbb{Z} \right\} \subset \mathbb{Q}(\sqrt{n})$$

$$A(n) = \{(a, b, c) \in \mathbb{Z}^3 : bc = a^2 - n\}.$$

The map $\frac{a + \sqrt{n}}{c} \mapsto (a, b, c)$ gives an 1-to-1 correspondence between $\mathbb{Q}^*(\sqrt{n})$ and $A(n)$.

Proposition 1.1. *For any integer $\lambda \geq 1$, the group $H(\lambda)$ acts on $\mathbb{Q}^*(\sqrt{n})$, hence on $A(n)$.*

Proof. Since $H(\lambda)$ is generated by $x : z \mapsto -\frac{1}{z}$ and $w_\lambda : z \mapsto z + \lambda$, it is enough to show that for any $\alpha = \frac{a + \sqrt{n}}{c} \in \mathbb{Q}^*(\sqrt{n})$, it holds that $x(\alpha), w_\lambda(\alpha) \in \mathbb{Q}^*(\sqrt{n})$. Indeed, we have

$$x(\alpha) = -\frac{1}{\alpha} = -\frac{c}{a + \sqrt{n}} = -\frac{c(a - \sqrt{n})}{a^2 - n} = \frac{-a + \sqrt{n}}{b}. \quad (1.1)$$

On the other hand

$$\frac{(-a)^2 - n}{b} = \frac{a^2 - n}{b} = c \in \mathbb{Z}. \quad (1.2)$$

From (1.1) and (1.2) it follows that $x(\alpha) \in \mathbb{Q}^*(\sqrt{n})$.

Also, we have that

$$w_\lambda(\alpha) = \alpha + \lambda = \frac{a + \lambda c + \sqrt{n}}{c}. \quad (1.3)$$

On the other hand

$$\frac{(a + \lambda c)^2 - n}{c} = \frac{a^2 + 2\lambda ac + \lambda^2 c^2 - n}{c} = \frac{bc + 2\lambda ac + \lambda^2 c^2}{c} = 2\lambda a + b + \lambda^2 c \in \mathbb{Z}. \quad (1.4)$$

From (1.3) and (1.4) it follows that $w_\lambda(\alpha) \in \mathbb{Q}^*(\sqrt{n})$. \square

From the proof of Proposition 1.1, it follows that

$$x(a, b, c) = (-a, c, b) \text{ and } y(a, b, c) = (-a - \lambda c, c, 2\lambda a + b + \lambda^2 c). \quad (1.5)$$

Lemma 1.2. *Let $\varepsilon := \frac{5 - \sqrt{7}}{3} \approx 0.78$. Let $(a, b, c) \in A(n)$ such that $c \geq \varepsilon|a|$ and $a^2 - n > 0$. If k is an integer with $|k| \geq 3$ and $(a', b', c') := (xw_k)(a, b, c)$ then $|a'| > |a|$ and $|c'| > \varepsilon|a'|$.*

Proof. Since $c > 0$ and $bc = a^2 - n > 0$, we have $b > 0$. From (1.5), it follows that

$$a' = -a - kc, \quad b' = c, \quad c' = b + 2ka + k^2c.$$

Since $|kc| = |k|c \geq 3\varepsilon|a| > |a|$, it follows that

$$|a'| \geq |k|c - |a| \geq (\varepsilon|k| - 1)|a| > |a|.$$

In order to prove $|c'| > \varepsilon|a'|$, we consider two cases:

(i) $ak > 0$. We have $|a'| = |a| + |k|c$. Hence:

$$c' = b + 2ka + k^2c > 2|k||a| + k^2c = |k|(2a + |k|c) > |k|(|a| + |k|c) = |k||a'| > \varepsilon|a'|.$$

(ii) $ak < 0$. We have $|a'| = -|a| + |k|c$. On the other hand

$$c' = b + 2ka + k^2c > 2ka + k^2c = -2|k||a| + |k|^2c.$$

It follows that

$$\begin{aligned} c' - \varepsilon|a'| &> -2|k|a + |k|^2c + \varepsilon|a| + \varepsilon|k|c = (|k|^2 - \varepsilon|k|)c + (-2|k| + \varepsilon)|a| \geq \\ &(\varepsilon|k|^2 - \varepsilon^2|k| - 2|k| + \varepsilon)|a| = \varepsilon(|k|^2 - \frac{2+\varepsilon}{\varepsilon}|k| + 1)|a| = \varepsilon|a|(k-3)(k-\frac{1}{3}). \end{aligned}$$

Since $|k| \geq 3$, we get $c' - \varepsilon|a'| > 0$, thus $c' > \varepsilon|a'|$. (We used the identity $\frac{2+\varepsilon}{\varepsilon} = \frac{10}{3}$)

□

Given $\alpha := (a, b, c) \in A(n)$, we denote $\|\alpha\| := |a|$. Note that $\|x(\alpha)\| = \|\alpha\| = |a|$.

Lemma 1.3. *The set $B(n) := \{(a, b, c) \in A(n) \mid |a| \leq |b|, |a| \leq |c|\}$ is finite.*

Proof. We consider two cases:

(i) $n > 0$. Since $a^2 - n = bc$, it follows that $|bc| = |a^2 - n| \geq |a|^2$, hence $a^2 \leq n$ and moreover $n - a^2 \geq a^2$, that is $|a| \leq \sqrt{\frac{n}{2}}$. It follows that

$$|b||c| = |a^2 - n| \leq |\frac{n}{2} - n| = \frac{n}{2},$$

hence $|b| \leq \frac{n}{2}$ and $|c| \leq \frac{n}{2}$.

(ii) $n < 0$. Since $bc = a^2 - n > a^2$, $|b| > |a|$ and $|c| > |a|$ it follows that

$$a^2 - n = |b||c| \geq (|a| + 1)|a| = a^2 + |a|,$$

hence $|a| \leq -n$. Therefore, $|b||c| \leq n^2 - n$, so $|b|, |c| \leq n^2 - n$.

□

Theorem 1.4. *Let $\lambda \geq 1$ be an integer. The action of $H(\lambda)$ on $\mathbb{Q}^*(n)$ has a finite number of orbits if and only if $\lambda \in \{1, 2\}$.*

Proof. Assume $\lambda \geq 3$. For any integer $s \geq 1$ with $(s-1)^2 > n$, we let

$$a_s := s^2 - s - n, \quad b_s := (s-1)^2 - n, \quad c_s := s^2 - n, \quad \alpha_s := (a_s, b_s, c_s).$$

Note that $a_s^2 - n = b_s c_s$, hence $\alpha_s \in A(n)$. Since

$$\lim_{s \rightarrow \infty} \frac{a_s}{c_s} = \lim_{s \rightarrow \infty} \frac{a_s}{b_s} = 1,$$

it follows that there exists an integer $s_0 \geq 1$ with $(s_0 - 1)^2 > n$, such that, for any $s \geq s_0$, we have $b_s > \varepsilon a_s$ and $c_s > \varepsilon a_s$, where $\varepsilon = \frac{5-\sqrt{7}}{3}$.

Let $s > t \geq s_0$ be two integers such that $(t - 1)^2 > n$.

We claim that $H(\lambda)\alpha_s \neq H(\lambda)\alpha_t$. In order to do that, we show that $\alpha_t \notin H(\lambda)\alpha_s$. An element β of the orbit $H(\lambda)\alpha_s$ is of the form $g\alpha_s$ with $g \in H(\lambda)$. Since $H(\lambda)$ is generated by x and w_λ , g has one of the forms

$$\begin{aligned} (i)g &= xw_{k_1}xw_{k_2}x \cdots xw_{k_j} \\ (ii)g &= xw_{k_1}xw_{k_2}x \cdots xw_{k_j}x \\ (iii)g &= w_{k_1}xw_{k_2}x \cdots xw_{k_j} \\ (iv)g &= w_{k_1}xw_{k_2}x \cdots xw_{k_j}x, \end{aligned}$$

where k_i 's are nonzero integers, multiples of λ , and $j \geq 0$ is an integer.

Since $\|x(\alpha)\| = \|\alpha\|$ for any $\alpha \in A(n)$, it is enough to tackle the cases (i) and (ii). Note that α_s and $x\alpha_s$ satisfy the conditions of Lemma 1.2. Hence, by applying Lemma 1.2, it follows that

$$\begin{aligned} \|(xw_{k_1}xw_{k_2}x \cdots xw_{k_j})(\alpha_s)\| &\geq a_s \\ \|(xw_{k_1}xw_{k_2}x \cdots xw_{k_j}x)(\alpha_s)\| &\geq a_s. \end{aligned}$$

Therefore, in all cases (i-iv), we have $\|g(\alpha_s)\| \geq \|\alpha_s\|$. Hence $\alpha_t \notin H(\lambda)\alpha_s$, as required.

From the above argument, it follows that the orbits $H(\lambda)\alpha_{s_0}, H(\lambda)\alpha_{s_0+1}, H(\lambda)\alpha_{s_0+2}, \dots$ are distinct, hence we have an infinite number of orbits.

In order to complete the proof, as $H(2)$ is a subgroup of $H(1)$, it is enough to show that the action of $H(2)$ on $A(n)$ has a finite number of orbits. Let $\alpha = (a, b, c) \in A(n)$. We claim that there exists $g \in H(2)$ such that $g(\alpha) \in B(n)$, where $B(n)$ was defined in the statement of Lemma 1.3. We use the following algorithm:

1. We let $g = 1$.
2. If $\alpha \in B(n)$, then we stop.
3. If $|a| > |c|$ then, according to the remainder theorem, there exists an integer $k \neq 0$ such that $-|c| < a + 2kc \leq |c|$, that is $|a + 2kc| \leq |c|$. We let

$$\alpha' := w_{2k}(\alpha) = (a', b', c'), \text{ where } a' = a + 2kc, b' = b + 2ka + 4k^2c \text{ and } c' = c.$$

We have $|a'| = |a + 2kc| \leq |c| < |a|$ and we replace α with α' and g with $w_{2k}g$. We return to 2.

4. If $|a| > |b|$ then, we let $\alpha' = x(\alpha) = (-a, c, b)$. We replace α with α' and g with xg . We return to 3.

Since, in the step 3., $\|\alpha'\| = |a'| < |a| = \|\alpha\|$, the above procedure eventually stop, and we obtain $g \in H(2)$ with $g(\alpha) \in B(n)$, as required. Since, according to Lemma 1.3, the set $B(n)$ is finite, it follows that the set of orbits of $H(2)$ is also finite. \square

Theorem 1.5. *The number $o(n)$ of orbits of the action of $H(2)$ on $\mathbb{Q}^*(n)$ is:*

$$o(n) \leq \begin{cases} \sum_{j=0}^{\lceil \sqrt{\frac{n}{2}} - 1 \rceil} d(n - j^2), & n < 0 \\ \sum_{j=1}^n d(j^2 - n), & n > 0 \end{cases},$$

where $d(n) =$ the number of integer divisors of n .

Proof. We consider the sets:

$$\begin{aligned} B^+(n) &= \{(a, b, c) \in B(n) \mid a > 0\} \\ B^0(n) &= \{(a, b, c) \in B(n) \mid a = 0\} \\ B^-(n) &= \{(a, b, c) \in B(n) \mid a < 0\} \end{aligned}$$

Obviously, $|B^+(n)| = |B^-(n)|$ and $|B^0(n)| = |\{(0, b, c) \mid bc = -n\}| = d(n)$. According to the proof of Theorem 1.4, for any $\alpha = (a, b, c) \in A(n)$, there exists a $g \in H(2)$ such that $g(\alpha) \in B(n)$. Note that, if $g(\alpha) \in B^-(n)$, then $xg(\alpha) \in B^+(n)$. It follows that

$$o(n) \leq |B^+(n)| + d(n). \quad (1.6)$$

Let $(a, b, c) \in B^+(n)$. We consider two cases:

- (i) $n < 0$. As in the proof of Lemma 1.3, we have $a \leq -n$. Since $bc = a^2 - n$, it follows that b is a divisor of $a^2 - n$, hence b can have $d(a^2 - n)$ possible values. Also, c is uniquely determined by a and b . It follows that

$$|B^+(n)| = \sum_{j=1}^n d(j^2 - n). \quad (1.7)$$

- (ii) $n > 0$. As in the proof of Lemma 1.3, we have $a \leq \sqrt{\frac{n}{2}}$. In fact $a < \sqrt{\frac{n}{2}}$, otherwise $bc = 0$ hence $b = 0$ or $c = 0$, a contradiction. Since $bc = a^2 - n$, it follows that b is a divisor of $n - a^2$, hence b can have $d(n - a^2)$ possible values. Also, c is uniquely determined by a and b . It follows that

$$|B^+(n)| = \sum_{j=1}^{\lceil \sqrt{\frac{n}{2}} - 1 \rceil} d(n - j^2). \quad (1.8)$$

The conclusion follows from (1.6), (1.7) and (1.8). \square

Remark 1.6. The inequality in Theorem 1.5 is strict. Let $\alpha := (1, -3, 2) \in B^+(7) \subseteq A(7)$. We have $w_{-2}(\alpha) = (-3, 1, 2)$, $xw_{-2}(\alpha) = (3, 2, 1)$ and $w_{-2}xw_{-2}(\alpha) = (1, -6, 1) \in B^+(7)$.

Let $\alpha := (2, 3, 2) \in B^+(-2) \subset A(-2)$. We have $w_{-2}(\alpha) = (-2, 3, 2)$ and therefore $xw_{-2}(\alpha) = (2, 2, 3) \in B^+(-2)$.

References

- [1] M. Aslam, *On the number of orbits arising from the action of $PSL(2, \mathbb{Z})$ on imaginary quadratic number fields*, <https://arxiv.org/pdf/1909.10448>, to appear in U.P.B. Sci. Bull. Series A (2020).
- [2] E. Hecke, *Über die bestimmung dirichletscher reihen durch ihre funktionalgleichung*, Math. Ann. 112 (1936), 664–699.
- [3] M. Malik, M. Riaz, *Orbits of $\mathbb{Q}(\sqrt{k^2m})$ under the action of the modular group $PSL(2, \mathbb{Z})$* , U.P.B. Sci. Bull. Series A **74** (2012), 109–116.
- [4] Q. Mushtaq, *Modular Group acting on Real Quadratic Fields*, Bull. Austral. Math. Soc. **Vol. 37** (1988), 303–309.
- [5] Q. Mushtaq, *On word structure of the modular group over finite and real quadratic fields*, Disc. Math., **178** (1998), 155–164.

Mircea Cimpoeaş, Simion Stoilow Institute of Mathematics, Research unit 5, P.O.Box 1-764, Bucharest 014700, Romania and University Politehnica of Bucharest, Faculty of Applied Sciences, Department of Mathematical Methods and Models, Bucharest, 060042, Romania.
E-mail: mircea.cimpoeas@imar.ro