

ON THE POSITIVE FIXED POINTS OF QUARTIC OPERATORS

Eshkabilov Yu. Kh. ¹ Nodirov Sh. D. ²

О положительных неподвижных точках кватрических операторов

В статье рассмотрен кватрик оператор с положительными коэффициентами на \mathbb{R}^2 . Исследованы положительные неподвижные точки кватрического оператора. Доказаны теоремы о количестве положительных неподвижных точек кватрического оператора.

Ключевые слова: Неподвижная точка; кватрический оператор; метод Феррари; правила Декарта; трансляционно-инвариантные меры Гиббса; дерево Кэли; модел; потенциал.

MSC 2010: 47H10, 65J15.

Keywords: Fixed point; quartic operator; Ferrari's method; Descartes rule; translation-invariant Gibbs measure; Cayley tree; model; hamiltonian; potential.

Introduction

It is well known that fixed point theory is a very important tool for solving problems in Nonlinear Functional Analysis and as well as to various theoretical and applied fields such as variational and linear inequalities, the dynamic systems theory, nonlinear analysis [8]. Nonlinear operators are connected with problems in statistical physics, biology, thermodynamics, statistical mechanics and so on [5], [9],[10]. One of the central problem in statistical physics is the description of the set of Gibbs measures for given models on Cayley trees. Translation-invariant Gibbs measures for models with a continuum set of spin values is reduced to find positive fixed points of nonlinear integral operators [5]-[6]. Shown that in [7],[11] to find positive fixed point of nonlinear integral operators require to find positive fixed points of nonlinear operators on the finite dimensional space \mathbb{R}^2 .

In [7] number of the positive fixed points of quadratic operator on \mathbb{R}^2 was studied. A number of positive fixed points of cubic operator on \mathbb{R}^2 were studied in [11]. In [7],[11] proven theorems, that quadratic and cubic operators can have one, two and three positive fixed points on \mathbb{R}^2 .

In this paper, we consider the operator of degree fourth (quartic) on \mathbb{R}^2 . We show that the existence of positive fixed point of quartic operators and they can have up to five positive fixed points. Besides, obtained the theorems for the number of positive fixed points of quartic operators. Furthermore, gives sufficient conditions for the number of positive fixed points of quartic operators (table 2) and at the end gives examples to satisfy conditions. Investigated quartic operators had arisen when examining translation-invariant Gibbs measures for models in [7] on the Cayley tree of order $k = 4$. In the paragraph 5 given model, which the study translation-invariant Gibbs measure reduced the study the fixed points of a quartic operator.

1 Quartic operators on \mathbb{R}_+^2

We introduce

$$\mathbb{R}_+^2 = \{(x, y) \in \mathbb{R}^2 : x \geq 0, y \geq 0\}, \quad \mathbb{R}_>^2 = \{(x, y) \in \mathbb{R}^2 : x > 0, y > 0\}.$$

We consider the following operator \mathcal{Q} on the cone \mathbb{R}_+^2 :

$$\mathcal{Q}(x, y) = \left(\sum_{i=0}^4 C_4^i a_i x^{4-i} y^i, \sum_{i=0}^4 C_4^i b_i x^{4-i} y^i \right), \quad (1)$$

where $C_4^i = \frac{4!}{(4-i)!i!}$ (binomial coefficient), $a_i > 0$ and $b_i > 0$ for all $i \in \{0, 1, 2, 3, 4\}$. This operator is composed of two polynomials of the fourth degree in two variables, therefore we call the operator \mathcal{Q} (1) a quartic. Clearly, an arbitrary non-trivial positive fixed point $(x_0, y_0) \in \mathbb{R}_+^2$ of the quartic operator (QO) \mathcal{Q} is a strictly positive, i.e. $(x_0, y_0) \in \mathbb{R}_>^2$. Denote by $N_{>}^{fix}(\mathcal{Q})$ the number of fixed points of the QO \mathcal{Q} that belongs to $\mathbb{R}_>^2$.

¹Karshi state university, Karshi, Uzbekistan. E-mail: yusup62@mail.ru

²Karshi state university, Karshi, Uzbekistan. E-mail: shoh0809@mail.ru

Lemma 1. If $\omega = (x_0, y_0) \in \mathbb{R}_>^2$ is a fixed point of the QO \mathcal{Q} , then $\xi_0 = \frac{y_0}{x_0}$ is root of the algebraic equation

$$a_4\xi^5 + (4a_3 - b_4)\xi^4 + (6a_2 - 4b_3)\xi^3 + (4a_1 - 6b_2)\xi^2 - (a_0 - 4b_1)\xi - b_0 = 0. \quad (2)$$

Proof. Let the point $\omega = (x_0, y_0) \in \mathbb{R}_>^2$ be a fixed point of QO \mathcal{Q} . Then

$$\sum_{i=0}^4 C_4^i a_i x_0^{4-i} y_0^i = x_0, \quad \sum_{i=0}^4 C_4^i b_i x_0^{4-i} y_0^i = y_0.$$

From the notation $\frac{y_0}{x_0} = \xi_0$ we obtain

$$x_0^4 \left(\sum_{i=0}^4 C_4^i a_i \xi_0^i \right) = x_0, \quad x_0^4 \left(\sum_{i=0}^4 C_4^i b_i \xi_0^i \right) = \xi_0 x_0.$$

Hence, we have

$$\frac{1}{\xi_0} = \frac{\sum_{i=0}^4 C_4^i a_i \xi_0^i}{\sum_{i=0}^4 C_4^i b_i \xi_0^i}.$$

By the last equality we get

$$a_4\xi_0^5 + (4a_3 - b_4)\xi_0^4 + (6a_2 - 4b_3)\xi_0^3 + (4a_1 - 6b_2)\xi_0^2 - (a_0 - 4b_1)\xi_0 - b_0 = 0.$$

Lemma 1 is proved. \square

Lemma 2. If ξ_0 is a positive root of the algebraic equation (2), then the point $\omega_0 = (x_0, \xi_0 x_0) \in \mathbb{R}_>^2$ and ξ_0 is a fixed point of the QO \mathcal{Q} , where

$$x_0 = \frac{1}{\sqrt[3]{\sum_{i=0}^4 C_4^i a_i \xi_0^i}}. \quad (3)$$

Proof. Let $\xi_0 > 0$ and ξ_0 is a root of the equation (2). Put $y_0 = \xi_0 x_0$, where x_0 is a given by the equality (3) and $\omega_0 = (x_0, \xi_0 x_0)$. From the equality $y_0 = \xi_0 x_0$ we have

$$\sum_{i=0}^4 C_4^i a_i x_0^{4-i} y_0^i = \sum_{i=0}^4 C_4^i a_i x_0^{4-i} (\xi_0 x_0)^i = x_0^4 \left(\sum_{i=0}^4 C_4^i a_i \xi_0^i \right) = \frac{1}{\sqrt[3]{\sum_{i=0}^4 C_4^i a_i \xi_0^i}},$$

i.e.

$$\sum_{i=0}^4 C_4^i a_i x_0^{4-i} y_0^i = x_0.$$

On the other hand

$$a_4\xi_0^5 + (4a_3 - b_4)\xi_0^4 + (6a_2 - 4b_3)\xi_0^3 + (4a_1 - 6b_2)\xi_0^2 - (a_0 - 4b_1)\xi_0 - b_0 = 0.$$

Then we get

$$\begin{aligned} b_0 + 4b_1\xi_0 + 6b_2\xi_0^2 + 4b_3\xi_0^3 + b_4\xi_0^4 &= a_0\xi_0 + 4a_1\xi_0^2 + 6a_2\xi_0^3 + 4a_3\xi_0^4 + a_4\xi_0^5 = \\ &= \xi_0 (a_0 + 4a_1\xi_0 + 6a_2\xi_0^2 + 4a_3\xi_0^3 + a_4\xi_0^4) \end{aligned}$$

From the last equality we have

$$\frac{\xi_0}{\sqrt[3]{\sum_{i=0}^4 C_4^i a_i \xi_0^i}} = \frac{\sum_{i=0}^4 C_4^i b_i \xi_0^i}{\left(\sqrt[3]{\sum_{i=0}^4 C_4^i a_i \xi_0^i} \right)^4} = x_0^4 \left(\sum_{i=0}^4 C_4^i b_i \xi_0^i \right) = \sum_{i=0}^4 C_4^i b_i x_0^{4-i} y_0^i = y_0.$$

□

We put

$$\mu_0 = a_4, \quad \mu_1 = 4a_3 - b_4, \quad \mu_2 = 6a_2 - 4b_3, \quad \mu_3 = 4a_1 - 6b_2, \quad \mu_4 = a_0 - 4b_1, \quad \mu_5 = b_0$$

and define the polynomial $P_5(\xi)$ of degree 5:

$$P_5(\xi) = \mu_0 \xi^5 + \mu_1 \xi^4 + \mu_2 \xi^3 + \mu_3 \xi^2 + \mu_4 \xi - \mu_5. \quad (4)$$

By the Lemmas 1, 2 the following corollary is correct.

Corollary 1. *The number of positive fixed points of the QO \mathcal{Q} equal to the number of positive roots of the polynomial $P_5(\xi)$.*

Lemma 3. *The polynomial $P_5(\xi)$ has at least one positive root.*

Proof. Clearly, that $P_5(0) = -\mu_5 < 0$ and $P_5(+\infty) = +\infty$. It means that there exists $c > 0$ such that $P_5(c) = 0$. □

Proposition 1. *The polynomial $P_5(\xi)$ can have up to five positive roots.*

Proof. We have the following table for the number of sign changes in the sequence of the coefficients of the polynomial $P_5(\xi)$ (Tab. 1). By using this table and the Descartes rule, we can conclude that the polynomial $P_5(\xi)$ can have up to five the positive roots (see [1], pp. 27-29).

$P_5(\xi)$	μ_0	μ_1	μ_2	μ_3	μ_4	μ_5	the number of sign changes
1.	+	-	+	-	+	-	5

Table 1.

□

2 Sufficient conditions for number of positive fixed points of the QO \mathcal{Q}

In this section we study a number of positive fixed points of the quartic operator \mathcal{Q} .

We use the following designations

$$Q = \left(\frac{a}{3}\right)^3 + \left(\frac{b}{2}\right)^2,$$

where

$$a = -\frac{p^2}{12} - r, \quad b = -\frac{p^3}{108} + \frac{pr}{3} - \frac{q^2}{8},$$

and

$$p = \frac{15\mu_0\mu_2 - 6\mu_1^2}{25\mu_0^2}, \quad q = \frac{50\mu_0^2\mu_3 + 8\mu_1^3 - 30\mu_0\mu_1\mu_2}{125\mu_0^3},$$

$$r = \frac{15\mu_0\mu_1^2\mu_2 - 50\mu_0^2\mu_1\mu_3 - 3\mu_1^4 - 125\mu_0^3\mu_4}{625\mu_0^4}.$$

When $Q < 0$, we define following number

$$z_0 = 2\sqrt{-\frac{a}{3}} \cos\left(\frac{2\pi}{3} + \frac{\alpha}{3}\right) - \frac{p}{3},$$

where

$$\cos \alpha = -\frac{b}{2} \left(-\frac{3}{a}\right)^{\frac{3}{2}}, \quad \alpha \in [0, \pi].$$

We also introduce

$$\xi_{1,2}^{ext} = \frac{1}{2} \left(\sqrt{2z_0} \pm \sqrt{2z_0 - 4 \left(\frac{p}{2} + z_0 + \frac{q}{2\sqrt{2z_0}} \right)} \right) - \frac{\mu_1}{5\mu_0},$$

$$\xi_{3,4}^{ext} = \frac{1}{2} \left(-\sqrt{2z_0} \pm \sqrt{2z_0 - 4 \left(\frac{p}{2} + z_0 - \frac{q}{2\sqrt{2z_0}} \right)} \right) - \frac{\mu_1}{5\mu_0}.$$

Let us note that if $z_0 > 0$ then the numbers $\xi_1^{ext}, \xi_2^{ext}, \xi_3^{ext}, \xi_4^{ext}$ are real.

For $z_0 > 0$, denote

$$\xi_{min} = \min\{\xi_1^{ext}, \xi_2^{ext}, \xi_3^{ext}, \xi_4^{ext}\}, \quad \xi_{max} = \max\{\xi_1^{ext}, \xi_2^{ext}, \xi_3^{ext}, \xi_4^{ext}\}.$$

Theorem 1. *Let $Q < 0, z_0 > 0, \xi_{max} < 0$. Then the QOQ has a unique fixed point in $\mathbb{R}_>^2$, i.e., $N_{>}^{fix}(Q) = 1$.*

Proof. By the Corollary 1 quite enough to find the positive roots of the polynomial $P_5(\xi)$.

Let $Q < 0$. Then by the Cardano's formula cubic equation $\eta^3 + a\eta + b = 0$ has three real roots. They have the following form [2]:

$$\eta_k = 2\sqrt{-\frac{a}{3}} \cos\left(\frac{\alpha + 2\pi(k-2)}{3}\right), \quad k = \overline{1, 2, 3},$$

where

$$\cos \alpha = -\frac{b}{2} \left(-\frac{3}{a}\right)^{\frac{3}{2}}, \quad \alpha \in [0, \pi].$$

We have $\eta_3 < \eta_1 < \eta_2$. The number $z_0 = \eta_3 - \frac{r}{3} (z_0 > 0)$ is the least root of the following cubic equation:

$$z^3 + pz^2 + \frac{p^2 - 4r}{4}z - \frac{q^2}{8} = 0. \quad (5)$$

On the other hand if all roots are strictly positive of the equation (5), then all roots $\omega_j, (j = \overline{1, \dots, 4})$ of the equation

$$\omega^4 + p\omega^2 + q\omega + r = 0$$

are real [3]. By the Ferrari's method the numbers $\xi_j^{ext} = \omega_j - \frac{\mu_1}{5\mu_0} (j = \overline{1, \dots, 4})$ are roots of the equation $P_5'(\xi) = 0$, i.e.

$$5\mu_0\xi^4 + 4\mu_1\xi^3 + 3\mu_2\xi^2 + 2\mu_3\xi + \mu_4 = 0. \quad (6)$$

Thus, the numbers $\xi_j^{ext} (j = \overline{1, \dots, 4})$ are the local extreme points of the function $P_5(\xi)$. Besides $\xi_{max} = \max_{j \in \{1, \dots, 4\}} \xi_j^{ext} < 0$ and $P_5(+\infty) = +\infty$. Therefore the function $P_5(\xi)$ is an increasing on the set $(\xi_{max}, +\infty)$. Consequently, by the inequality $P_5(0) < 0$ the polynomial $P_5(\xi)$ has a unique positive root. \square

Theorem 2. *Let $Q < 0, z_0 > 0, \xi_{min} > 0$. If the polynomial $P_5(\xi)$ satisfies one of the following conditions*

(a) $P_5(\xi_{min}) = 0$,

(b) $P_5(\xi_{max}) = 0$,

then the QOQ has at least two fixed points in $\mathbb{R}_>^2$, i.e., $N_{>}^{fix}(Q) \geq 2$.

Proof. Let $Q < 0, z_0 > 0, \xi_{min} > 0$. Then all four roots of the equation (6) is real and they strictly positive. And so the function $P_5(\xi)$ is an increasing on the set $(-\infty, \xi_{min}) \cup (\xi_{max}, +\infty)$.

(a) $P_5(\xi_{min}) = 0$. Then $\max_{\xi \in (-\infty, \xi_{min}]} P_5(\xi) = P_5(\xi_{min}) = 0$ and the number $\xi_1 = \xi_{min}$ is the root of the polynomial $P_5(\xi)$. Besides we know, that the function $P_5(\xi)$ is an increasing on the set $(\xi_{max}, +\infty)$. Therefore exist $\xi_2 \in (\xi_{max}, +\infty)$, where $P_5(\xi_2) = 0$. And so the polynomial $P_5(\xi)$ has at least two strictly positive roots.

(b) $P_5(\xi_{max}) = 0$. Then $\min_{\xi \in [\xi_{max}, +\infty)} P_5(\xi) = P_5(\xi_{max}) = 0$ and the number $\xi_1 = \xi_{max}$ is the root of the polynomial $P_5(\xi)$. The function $P_5(\xi)$ is an increasing on the set $[\xi_{max}, +\infty)$, so exist $\xi_{k_0} \neq \xi_{max}$ which the function $P_5(\xi)$ is decreasing on the set (ξ_{k_0}, ξ_{max}) . Also we have $P_5(0) < 0$ and $\max_{\xi \in [\xi_{k_0}, \xi_{max}]} P_5(\xi) = P_5(\xi_{k_0}) > 0$. Therefore the polynomial $P_5(\xi)$ has another unique positive root ξ_2 on the set $(0, \xi_{k_0})$. \square

Theorem 3. *Let $Q < 0, z_0 > 0, \xi_{min} > 0$. If for the polynomial $P_5(\xi)$ following condition*

(c) $P_5(\xi_{min}) > 0, P_5(\xi_{max}) < 0$

is satisfies, then the QO \mathcal{Q} has at least three fixed points in $\mathbb{R}_{>}^2$, i.e., $N_{>}^{fix}(\mathcal{Q}) \geq 3$.

Proof. Let $Q < 0, z_0 > 0, \xi_{min} > 0$. Then all four roots of the equation (6) is real and they strictly positive. And so the function $P_5(\xi)$ is an increasing on the set $(-\infty, \xi_{min}) \cup (\xi_{max}, +\infty)$.

(c) Let $P_5(\xi_{min}) > 0, P_5(\xi_{max}) < 0$. We have known that $P_5(0) < 0$ and $P_5(+\infty) = +\infty$. It indicates that the polynomial $P_5(\xi)$ has at least three positive roots, such that $\xi_1 \in (0, \xi_{min}), \xi_2 \in (\xi_{min}, \xi_{max})$ and $\xi_3 \in (\xi_{max}, +\infty)$. □

In the next, we assume that $Q < 0, z_0 > 0$. Then we can define following set of real numbers (set of extremal points of the function $P_5(\xi)$):

$$\mathfrak{A} = \{\xi_1^{ext}, \xi_2^{ext}, \xi_3^{ext}, \xi_4^{ext}\}.$$

Points of the set of \mathfrak{A} are real and they are different. Therefore we can have denote following

$$\lambda_1 = \xi_{min}, \lambda_2 = \min(\mathfrak{A} \setminus \{\xi_{min}\}), \lambda_3 = \max(\mathfrak{A} \setminus \{\xi_{max}\}), \lambda_4 = \xi_{max}.$$

Consequently, we have the following relation

$$\lambda_1 < \lambda_2 < \lambda_3 < \lambda_4.$$

It follows the function $P_5(\xi)$ is an increasing (decreasing) on the set $(-\infty, \lambda_1) \cup (\lambda_2, \lambda_3) \cup (\lambda_4, +\infty)$ ($(\lambda_1, \lambda_2) \cup (\lambda_3, \lambda_4)$). The function $P_5(\xi)$ has a local maximum value at the points λ_1, λ_3 and local minimum values at the points λ_2 and λ_4 . Using these property and by the corollary 1, we can give the following table:

On the number of the positive fixed points of the quartic operators, when $Q < 0, z_0 > 0, \lambda_1 > 0$.		
	Sufficient condition:	$N_{>}^{fix}(\mathcal{Q})$
1.	$P_5(\lambda_1) < 0, P_5(\lambda_3) < 0$	1
2.	$P_5(\lambda_1) = 0, P_5(\lambda_3) < 0$	2
3.	$P_5(\lambda_1) < 0, P_5(\lambda_3) = 0$	
4.	$P_5(\lambda_2) > 0, P_5(\lambda_4) = 0$	
5.	$P_5(\lambda_2) = 0, P_5(\lambda_4) > 0$	
6.	$P_5(\lambda_2) > 0, P_5(\lambda_4) = 0$	
7.	$P_5(\lambda_1) = 0, P_5(\lambda_4) = 0$	3
8.	$P_5(\lambda_1) = 0, P_5(\lambda_3) = 0$	
9.	$P_5(\lambda_2) = 0, P_5(\lambda_4) = 0$	
10.	$P_5(\lambda_2) > 0, P_5(\lambda_4) < 0$	
11.	$P_5(\lambda_1) < 0, P_5(\lambda_3) > 0, P_5(\lambda_4) < 0$	4
12.	$P_5(\lambda_1) > 0, P_5(\lambda_2) < 0, P_5(\lambda_4) > 0$	
13.	$P_5(\lambda_1) = 0, P_5(\lambda_3) > 0, P_5(\lambda_4) < 0$	
14.	$P_5(\lambda_1) > 0, P_5(\lambda_3) = 0$	
15.	$P_5(\lambda_2) = 0, P_5(\lambda_4) < 0$	
16.	$P_5(\lambda_1) > 0, P_5(\lambda_2) < 0, P_5(\lambda_4) = 0$	
17.	$P_5(\lambda_1) > 0, P_5(\lambda_2) < 0, P_5(\lambda_3) > 0, P_5(\lambda_4) < 0$	

Table 2.

3 Examples

In this section we give the concrete examples, which the QO \mathcal{Q} has the three and the five positive fixed points.

EXAMPLE 1. We consider an example for QO \mathcal{Q} , which satisfies the 7th condition in the table 2. We define QO \mathcal{Q} by the equality:

$$\mathcal{Q}(x, y) = (52x^4 + 8x^3y + 42x^2y^2 + 4xy^3 + y^4, 16x^4 + 4x^3y + 72x^2y^2 + 5xy^3 + 14y^4). \tag{7}$$

Then for the QO \mathcal{Q} we have

$$a_0 = 52, \quad a_1 = 2, \quad a_2 = 7, \quad a_3 = 1, \quad a_4 = 1, \quad b_0 = 16, \quad b_1 = 1, \quad b_2 = 12, \quad b_3 = \frac{5}{4}, \quad b_4 = 1.$$

Therefore we get

$$\mu_0 = a_4 = 1, \quad \mu_1 = 4a_3 - b_4 = -10, \quad \mu_2 = 6a_2 - 4b_3 = 37, \quad \mu_3 = 4a_1 - 6b_2 = -64,$$

$$\mu_4 = a_0 - 4b_1 = 52, \quad \mu_5 = b_0 = 16.$$

For the QO \mathcal{Q} (7) the polynomial $P_5(\xi)$ has the following form

$$P_5(\xi) = \xi^5 - 10\xi^4 + 37\xi^3 - 64\xi^2 + 52\xi - 16.$$

Then we have

$$Q = -\frac{7}{12500}, \quad z_0 = \frac{1}{20} (13 - \sqrt{105}),$$

$$\xi_1^{ext} = 1, \quad \xi_2^{ext} = 2, \quad \xi_3^{ext} = \frac{1}{10} (25 - \sqrt{105}), \quad \xi_4^{ext} = \frac{1}{10} (25 + \sqrt{105}).$$

By the denotes we have

$$\lambda_1 = 1, \quad \lambda_2 = \frac{1}{10} (25 - \sqrt{105}), \quad \lambda_3 = 2, \quad \lambda_4 = \frac{1}{10} (25 + \sqrt{105}).$$

So $Q < 0$, $z_0 > 0$, $\lambda_1 > 0$, $P_5(\lambda_1) = P_5(1) = 0$, $P_5(\lambda_3) = P_5(2) = 0$. By the 7th condition in the table 2, the QO \mathcal{Q} has three positive fixed points.

It is easy to verify that the polynomial $P_5(\xi)$ has three positive roots and they have the following form:

$$\xi_{1,2} = 1, \quad \xi_{3,4} = 2, \quad \xi_5 = 4.$$

By the Corollary 1 the QO \mathcal{Q} (7) has three positive fixed points.

EXAMPLE 2. Now, we consider the following QO \mathcal{Q} :

$$\mathcal{Q}(x, y) = \left(31x^4 + \frac{1}{2}x^3y + \frac{43}{3}x^2y^2 + \frac{1}{4}xy^3 + \frac{1}{5}y^4, 10x^4 + x^3y + 31x^2y^2 + \frac{2}{3}xy^3 + 3y^4 \right). \quad (8)$$

Then

$$P_5(\xi) = \frac{1}{5}\xi^5 - \frac{11}{40}\xi^4 + \frac{41}{3}\xi^3 - \frac{61}{2}\xi^2 + 30\xi - 106.$$

Also

$$a = -\frac{7}{3}, \quad b = -\frac{20}{27}, \quad Q = -\frac{1}{3},$$

and

$$z_0 = \frac{1}{8}.$$

Consequently,

$$\lambda_1 = 1, \quad \lambda_2 = 2, \quad \lambda_3 = 3, \quad \lambda_4 = 5.$$

The following relations are true :

$$Q = -\frac{1}{3} < 0, \quad z_0 = \frac{1}{8} > 0, \quad \lambda_1 = 1 > 0,$$

$$P_5(\lambda_1) = P_5(1) = \frac{37}{60} > 0, \quad P_5(\lambda_2) = P_5(2) = -\frac{4}{15} < 0,$$

$$P_5(\lambda_3) = P_5(3) = \frac{7}{20} > 0, \quad P_5(\lambda_4) = P_5(5) = -\frac{95}{12} < 0,$$

By the 17th condition in the table 2, the QO \mathcal{Q} (8) has five positive fixed points.

4 Application

In this section, we show that the number of fixed points of the operator \mathcal{Q} (1) corresponds to a non-trivial positive fixed point of the Hammerstein integral operator, which plays an important role in the theory of Gibbs measures.

A Cayley tree $\Gamma^k = (V, L)$ of order $k \geq 1$ is an infinite homogeneous tree, i.e. a graph without cycles, with exactly $k + 1$ edges incident to each vertices. Here V is the set of vertices and L that of edges.

Consider models where the spin takes values in the set $[0, 1]$, and is assigned to the vertices of the tree. For $A \subset V$ a configuration σ_A on A is an arbitrary function $\sigma_A : A \rightarrow [0, 1]$. Denote $\Omega_A = [0, 1]^A$ is the set of all configurations on A .

We consider the Hamiltonian of the model in [11]:

$$H(\sigma) = -J \sum_{\langle x, y \rangle \in L} \xi_{\sigma(x), \sigma(y)}, \quad \sigma \in \Omega_V \quad (9)$$

where $J \in \mathbb{R} \setminus \{0\}$ and $\beta = \frac{1}{T}$, $T > 0$ is temperature, $\xi : (u, v) \in [0, 1]^2 \rightarrow \xi_{uv} \in \mathbb{R}$ is a given bounded, measurable function. As usually, $\langle x, y \rangle$ stands for the nearest neighbor vertices on the Cayley tree of order four.

We introduce

$$C_+[0, 1] = \{f \in C[0, 1] : f(x) \geq 0\}, \quad C_+^0[0, 1] = C_+[0, 1] \setminus \{\theta \equiv 0\}, \quad C_{>}[0, 1] = \{f \in C[0, 1] : f(x) > 0\}.$$

Consider the following operator [5] R_k on $C_+[0, 1]$ defined by:

$$(R_k f)(t) = \left(\frac{\int_0^1 K(t, u) f(u) du}{\int_0^1 K(0, u) f(u) du} \right)^k, \quad k \in \mathbb{N}$$

where $K(t, u) = \exp(J\beta\xi_{tu})$, $f(t) > 0$, $t, u \in [0, 1]$.

In [4]-[6] was shown that a translation-invariant Gibbs measure of the model H (9) corresponds to a solution $f(t) \in C_+^0[0, 1]$ of the following equation:

$$(R_k f)(t) = f(t). \quad (10)$$

For every $k \geq 2$ we define the Hammerstein integral operator H_k acting in $C_+[0, 1]$ as following:

$$(H_k f)(t) = \int_0^1 K(t, u) f^k(u) du.$$

Lemma 4. *Let $k \geq 2$. The equation (10) has a non-trivial positive solution iff the Hammerstein integral operator has a non-trivial positive fixed point and*

$$N_+^{fix}(R_k) = N_+^{fix}(H_k),$$

where $N_+^{fix}(T)$ is a number of non-trivial positive fixed points of the operator T .

Proof. At first, we notice that the equation (10) has at least one solution in $C_{>}[0, 1]$ (see Theorem 3.5 in [6]).

Necessity. Let $k \geq 2$ and $f(t) \in C_+^0[0, 1]$ be a solution of the (10), then $f(0) = 1$. Define the following function

$$g(t) = \frac{\sqrt[k]{f(t)}}{\sqrt[k-1]{\int_0^1 K(0, u) f(u) du}}, \quad (11)$$

where $\int_0^1 K(0, u) f(u) du > 0$. From $f(0) = 1$, g corresponds to exactly one function f . Then we have

$$(H_k g)(t) = g(t).$$

Sufficiency. Let $k \geq 2$ and $g = g(t) \in C_+^0[0, 1]$ be a fixed point of the operator H_k . It is clear that $\int_0^1 K(0, u)g^k(u)du = g(0) > 0$. Define non-trivial positive continuous function

$$f(t) = \left(\frac{g(t)}{g(0)} \right)^k. \tag{12}$$

Because of g is a fixed point of the operator H_k , the last equality is well-defined, i.e., each function f corresponds to exactly one function g .

Then

$$(R_k f)(t) = \left(\frac{\int_0^1 K(t, u)f(u)du}{\int_0^1 K(0, u)f(u)du} \right)^k = \left(\frac{\int_0^1 K(t, u)g^k(u)du}{\int_0^1 K(0, u)g^k(u)du} \right)^k = \left(\frac{g(t)}{g(0)} \right)^k = f(t).$$

It is easy to check that the relations (11) and (12) establish a one-to-one correspondence between

$$F_\alpha = \{f(t) \in C_+^0[0, 1] : R_k f = f\} \quad \text{and} \quad G_\alpha = \{g(t) \in C_+^0[0, 1] : H_k g = g\}.$$

Consequently $N_+^{fix}(R_k) = N_+^{fix}(H_k)$.

This completes the proof. □

From Lemma 4 we can conclude that the number of the non-trivial positive fixed points of the operator H_k is equal to the number of translation-invariant Gibbs measures of the model H (9).

Let functions $\varphi_1(t)$, $\varphi_2(t)$ and $\phi_1(t)$, $\phi_2(t)$ belong to $C_{>}[0, 1]$. We consider the Hamiltonian H (9) on the Cayley tree of order $k = 4$ with the function of potential

$$\xi_{t,u} = \frac{1}{J\beta} \ln \left(\phi_1(t)\varphi_1(u) + \phi_2(t)\varphi_2(u) \right). \tag{13}$$

In a similar manner, we can prove that for the number non-trivial positive fixed points of the integral operator H_4 the following equality holds, as in the [11] (see Lemma 7):

$$N_+^{fix}(H_4) = N_{>}^{fix}(\mathcal{Q}).$$

By the last equality we can conclude that number of nontrivial positive fixed points of the quartic operator \mathcal{Q} (1) on the $\mathbb{R}_{>}^2$ is equal to number of translation-invariants Gibbs measures for the model H (9) on the Cayley tree of order four.

Open problem. It has not been constructed yet the exact function of potential $\xi_{t,u}$ (13) for satisfy the conditions of the Theorems 2-3. This requires constructing non-trivial positive continuous functions $\varphi_1(t)$, $\varphi_2(t)$ and $\phi_1(t)$, $\phi_2(t)$ in such a way that the corresponding coefficients

$$a_i = \int_0^1 \varphi_1(u) \phi_1^{k-i}(u) \phi_2^i(u) du, \quad b_i = \int_0^1 \varphi_2(u) \phi_1^{k-i}(u) \phi_2^i(u) du, \quad i = \overline{0, \dots, 4}$$

of the nonlinear operator \mathcal{Q} need satisfy the conditions of the Theorems 2-3 (as in the above examples 1-2).

Acknowledgements. We would like to thank reviewers for their insightful comments on the paper, as these comments led us to an improvement of the work.

References

1. Victor V. Prasolov. Polynomials. *Algorithms and Computation in Mathematics. Volume 11.* **316** (2000).
2. Nickalls R. W. D. Vieta, Descartes and the cubic equation. *Mathematical Gazette.* **90** (July 2006), 203-208.
3. Nickalls R. W. D. The quartic equation: invariants and Euler’s solution revealed. *The Mathematical Gazette.* vol. 93 (March; No. 526),(2009), pp. 66-75.

4. Rozikov, U. A. , Eshkabilov, Yu. Kh. On models with uncountable set of spin values on a Cayley tree: integral equations. *Math. Phys. Anal. Geom.* **13**, 275-286 (2010).
5. Eshkabilov Yu. Kh. , Haydarov F. H. , Rozikov U. A. Non-uniqueness of Gibbs Measure for Models with Uncountable Set of Spin Values on a Cayley Tree. *J. Stat. Phys.* **147**(2012),779-794.
6. Eshkabilov Yu. Kh. , Haydarov F. H. , Rozikov U. A. Uniqueness of Gibbs Measure for Models With Uncountable Set of Spin Values on a Cayley Tree. *Math. Phys. Anal. Geom.* **16** (2013), 1-17.
7. Eshkabilov Yu. Kh. , Nodirov Sh. D. , Haydarov F. H. Positive fixed points of quadratic operators and Gibbs measures. *Positivity*, **20** No.4, (2016), 929-943.
8. Amelia Bucur. About application of the fixed point theory. *Scientific Bulletin*. Vol. XXII. No 1(43),(2017), 13-17.
9. Juan Martinez-Moreno, Dhananjay Gopal, Vijay Gupta, Edixon Rojas, and Satish Shukla. Nonlinear Operator Theory and Its Applications. *Hindawi Journal of Function Spaces*, Article ID 9713872, Volume 2018, 2 pages.
10. Jamilov U. U. , Khamraev A. Yu. , Ladra M. On a Volterra Cubic Stochastic Operator. *Bull Math Biol*, **80**:2, (2018), 319-334.
11. Eshkabilov Yu. Kh. , Nodirov Sh. D. Positive Fixed Points of Cubic Operators on \mathbb{R}^2 and Gibbs Measures. *Journal of Siberian Federal University. Mathematics Physics*. 12(6),(2019), 663-673.