Technical Note - Exact simulation of the first passage time of Brownian motion to a symmetric linear boundary

Jong Mun Lee, Taeho Lee

July 14, 2020

Abstract

We state an exact simulation scheme for the first passage time of a Brownian motion to a symmetric linear boundary. This note summarizes a part of Chapter 4 in [1] which is the unpublished doctoral dissertation of Jong Mun Lee.

Keywords: Brownian motion, hitting time simulation, exact simulation, linear boundary

1 Theoretical Results

Let $\tau_{a,b}$ be the first hitting time of Brownian motion to a symmetric and linear boundary $\pm(a + bt)$ with given nonnegative coefficients a, b. The following theorem from [2] provides an explicit expression for the distribution of $\tau_{a,b}$ including the probability $\mathbb{P}[\tau_{a,b} = \infty]$.

Theorem 1.1 (Abundo [2]) For any a, b > 0, the distribution of $\tau_{a,b}$ is given by

$$\mathbb{P}[\tau_{a,b} \le t] = 1 - \sum_{k \in \mathbb{Z}} (-1)^k e^{-2k^2 ab} \int_{(-(a+bt)+2ak)/\sqrt{t}}^{(a+bt+2ak)/\sqrt{t}} \frac{e^{-x^2/2}}{\sqrt{2\pi}} dx,$$
$$\mathbb{P}[\tau_{a,b} < \infty] = 2 \sum_{k=1}^{\infty} (-1)^{k+1} e^{-2k^2 ab}.$$

We suggest an algorithm based on the Acceptance & Rejection (AR) method motivated by [3] which solves the same problem under the restriction b = 0. We first calculate the conditional density function $f_{a,b}(t)$ under the condition $[\tau_{a,b} < \infty].$

$$f_{a,b}(t) = \frac{d}{dt} \mathbb{P}[\tau_{a,b} \le t \mid \tau_{a,b} < \infty]$$

$$= -\frac{1}{C_{a,b}} \sum_{k \in \mathbb{Z}} \frac{(-1)^k e^{-2k^2 ab}}{\sqrt{8\pi t^3}} \left((-a + bt - 2ak) e^{-\frac{(a+bt+2ak)^2}{2t}} - (a - bt - 2ak) e^{-\frac{(-a-bt+2ak)^2}{2t}} \right)$$

$$= \frac{1}{C_{a,b}} \left[\frac{a - bt}{\sqrt{2\pi t^3}} e^{-\frac{(a+bt)^2}{2t}} + \sum_{k=1}^{\infty} \frac{(-1)^k e^{-2k^2 ab}}{\sqrt{2\pi t^3}} \left((-a + bt - 2ak) e^{-\frac{(a+bt+2ak)^2}{2t}} - (a - bt - 2ak) e^{-\frac{(-a-bt+2ak)^2}{2t}} \right) \right],$$
(1)

where $C_{a,b} = \mathbb{P}[\tau_{a,b} < \infty] = 2 \sum_{k=1}^{\infty} (-1)^{k+1} e^{-2k^2 a b}.$

To apply the AR method, we need a density whose scale-up dominates the true density $f_{a,b}$. The following theorem shows that the gamma distribution can be used.

Theorem 1.2 For any $\alpha \geq \frac{1}{2}$, there exist a constant M such that

$$f_{a,b}(t) \le M \cdot g(t;\alpha,b^2/2) \tag{2}$$

for all $t \ge 0$, where $g(t : \alpha, \lambda)$ is the density function of gamma distribution given by

$$g(t;\alpha,\lambda) = \frac{\lambda^{\alpha} t^{\alpha-1e^{-\lambda t}}}{\Gamma(\alpha)}$$
(3)

Proof To prove Theorem 1.2, we analyze the tail asymptotic of $f_{a,b}(t)$. The following two lemmas analyze the left tail asymptotic and the right tail asymptotic, respectively.

Lemma 1.1 (Left tail asymptotic) For any $n \ge 1$

$$f_{a,b}(t) = o(t^n)$$
 as $t \to 0$.

Proof Taking the positive terms and using the fact that $e^{-x} \le m!/x^m$ for any $m \ge 0$ and $x \ge 0$, we have for $0 \le t < a/(2b)$ and $m \ge 2$

$$\begin{split} C_{a,b} \cdot f_{a,b}(t) &\leq \frac{a - bt}{\sqrt{2\pi}\sqrt{t^3}} e^{-(a+bt)^2/(2t)} \\ &+ \sum_{k:odd} e^{-2k^2ab} \frac{1}{\sqrt{2\pi}\sqrt{t^3}} (-a + bt + 2ak) e^{-(-(a+bt) + 2ak)^2/(2t)} \\ &+ \sum_{k:even} e^{-2k^2ab} \frac{1}{\sqrt{2\pi}\sqrt{t^3}} (a - bt + 2ak) e^{-(a+bt + 2ak)^2/(2t)} \\ &\leq \frac{e^{-(a-bt)^2/(2t)}}{\sqrt{2\pi}\sqrt{t^3}} \sum_{k\geq 0} e^{-2k^2ab} a(2k+1) \\ &\leq \frac{m!2^m t^m}{(a-bt)^{2m}} \frac{1}{\sqrt{2\pi}\sqrt{t^3}} \sum_{k\geq 0} e^{-2k^2ab} a(2k+1) = c_m t^{m-3/2}, \end{split}$$

where c_m is some constant depends on a, b, m only.

Lemma 1.2 (Right tail asymptotic)

$$f_{a,b}(t) = O(t^{-\frac{1}{2}}e^{-\frac{b^2}{2}t}) \text{ as } t \to \infty.$$

Proof For large enough t

$$\begin{aligned} C_{a,b} \cdot f_{a,b}(t) &\leq \sum_{k=1}^{\infty} e^{-2k^2 a b} \frac{2}{\sqrt{2\pi}\sqrt{t^3}} \Big((a+bt+2ak)e^{-(-(a+bt)+2ak)^2/(2t)} \Big). \\ &= \frac{2e^{-(a+bt)^2/(2t)}}{\sqrt{2\pi}\sqrt{t^3}} \sum_{k=1}^{\infty} (a+bt+2ak) \exp\big(-\frac{4a^2k^2-4(a+bt)ak}{2t}-2k^2ab\big) \\ &= \frac{2e^{-(a+bt)^2/(2t)}}{\sqrt{2\pi}\sqrt{t^3}} \sum_{k=1}^{\infty} (a+bt+2ak) \exp\big(-2ab(k^2-k)\big) \\ &= O(t^{-\frac{1}{2}}e^{-\frac{b^2}{2}t}). \end{aligned}$$

From Lemma 1.1, we have $\lim_{t\to 0} \frac{f_{a,b}(t)}{g(t;\alpha,b^2/2)} = 0$ for any $\alpha, b > 0$. And from Lemma 1.2, we have $\limsup_{t\to\infty} \frac{f_{a,b}(t)}{g(t;\alpha,b^2/2)} < \infty$ for $\alpha \ge 1/2$ and thus Theorem 1.2 is proved.

Now we propose a prototype of our algorithm to sample $\tau_{a,b}$. We call Algorithm 1 a prototype since the evaluation of

Algorithm 1 Sampling of $\tau_{a,b}$ (Sketch)

1: Generate $U \sim U(0, 1)$ 2: if $U < C_{a,b}$ then Resolved = FALSE 3: repeat 4: Generate $V \sim \text{Gamma}(\alpha, b^2/2)$ 5: if $f_{a,b}(V) < M \cdot g(V; \alpha, b^2/2)$ then 6: 7: T = VResolved = TRUE 8: end if 9: until Resolved 10: 11: else $T = \infty$ 12: 13: end if 14: return T

 $C_{a,b}$ and $f_{a,b}(t)$ cannot be done in finite time. Though we cannot evaluate $C_{a,b}$ and $f_{a,b}(t)$ exactly, it is possible to determine the corresponding inequalities exactly in finite time. We explore some special feature of the quantities $C_{a,b}$ and $f_{a,b}(t)$ first.

Definition 1.1 (Oscillating sequence and series) A sequence of numbers $\{a_n\}_{n=1}^{\infty}$ is called an oscillating sequence

if there exists a positive integer N *such that for* $k \ge N$ *,*

$$-1 < \frac{a_{k+1} - a_k}{a_k - a_{k-1}} < 0.$$

And we call an infinite series $\sum_{n=1}^{\infty} a_n$ is an oscillating series if its partial sum $s_n = \sum_{k=1}^{n} a_k$ is an oscillating sequence.

Lemma 1.3 shows that the inequality [S > s] can be determined exactly in finite time for an oscillating series $S = \sum_{n=1}^{\infty} a_n$ and a constant $s \in \mathbb{R}$. The same technique is used in [4] which suggests an exact simulation scheme for a solution of SDE.

Lemma 1.3 Let $S = \sum_{n=1}^{\infty} a_n$ be an oscillating series where its oscillating property holds for $k \ge N$ and S_n be its partial sum. We have S > s if and only if

$$S_k \wedge S_{k+1} > s \tag{4}$$

for some $k \ge N$. Conversely, We have S < s if and only if

$$S_k \wedge S_{k+1} < s \tag{5}$$

for some $k \geq N$.

The following two lemmas shows that the series $C_{a,b}$ and $f_{a,b}(t)$ used in Algorithm 1 are oscillating series.

Lemma 1.4 $C_{a,b} = 2 \sum_{k=1}^{\infty} (-1)^{k+1} e^{-2k^2 ab}$ is an oscillating series.

Proof Let $a_k = (-1)^{k+1} e^{-2k^2 ab}$. Then we have

$$0 < -\frac{a^{(n+1)} - a^{(n)}}{a^{(n)} - p^{(n-1)}} = \frac{e^{-2(n+1)^2ab}}{e^{-2n^2ab}} = e^{-2(2n+1)ab} < 1.$$

Lemma 1.5 For any t > 0, $f_{a,b}(t)$ is an oscillating series with $N(t) = \max\left(\log \frac{6}{4ab}, \frac{bt}{2a}, 1\right) + 1$

Proof Let

$$h_n(t) := e^{-2n^2ab} \left((-a+bt+2an)e^{-(-(a+bt)+2an)^2/2t} - (a-bt+2an)e^{-(a+bt+2an)^2/2t} \right).$$

Then

$$-\frac{q^{(n+1)}(t) - q^{(n)}(t)}{q^{(n)}(t) - q^{(n-1)}(t)} = \frac{h_{n+1}(t)}{h_n(t)}.$$

First, we have

$$h_n(t) > 0 \Leftrightarrow (-a + bt + 2an)e^{-(-(a+bt)+2an)^2/2t} > (a - bt + 2an)e^{-(a+bt+2an)^2/2t}$$
$$\Leftrightarrow e^{4a^2n/t+4abn} > \frac{(a - bt + 2an)}{(-a + bt + 2an)}$$
$$\Leftrightarrow e^{4abn} > \frac{(a + 2an)}{(-a + 2an)}.$$

For n > 1, (a + 2an)/(-a + 2an) is strictly bounded above by 3. Thus, $h_n(t)$ is strictly positive when $n > \max\{\log 3/4ab, 1\}$. Next, we have that for $n > \max\{\log 3/4ab, 1\} \lor (b/2a)t$,

$$\begin{aligned} \frac{h_{n+1}(t)}{h_n(t)} < 1 \Leftrightarrow e^{-2(n+1)^2 a b} e^{-((a+bt)^2 + 4a^2(n+1)^2)/2t} \\ & \times \left((-a+bt+2a(n+1))e^{4(a+bt)a(n+1)/2t} - (a-bt+2a(n+1))e^{-4(a+bt)a(n+1)/2t} \right) \\ < e^{-2n^2 a b} e^{-((a+bt)^2 + 4a^2 n^2)/2t} \left((-a+bt+2an)e^{4(a+bt)an/2t} - (a-bt+2an)e^{-4(a+bt)a(n+1)/2t} \right) \\ \Leftrightarrow (-a+bt+2a(n+1))e^{4(a+bt)a(n+1)/2t} - (a-bt+2a(n+1))e^{-4(a+bt)a(n+1)/2t} \\ < (-a+bt+2an)e^{4(a+bt)a(3n+1)/2t} - (a-bt+2an)e^{4(a+bt)a(n+1)/2t} \\ \leqslant 2a(2n+1)e^{4(a+bt)a(n+1)/2t} < (-a+bt+2an)e^{4(a+bt)a(3n+1)/2t} \\ \Leftrightarrow \frac{2a(2n+1)}{(-a+bt+2an)} < e^{4(a+bt)an/t} \\ \Leftrightarrow \frac{2(a+2an)}{(-a+2an)} < e^{4abn}. \end{aligned}$$

The left-hand side of the last inequality is bounded above by 6. And to conclude, we have that

$$0 < -\frac{q^{(n+1)}(t) - q^{(n)}(t)}{q^{(n)}(t) - q^{(n-1)}(t)} = \frac{h_{n+1}(t)}{h_n(t)} < 1,$$

for $n \ge N(t) := \max\{\frac{\log 6}{4ab}, \frac{bt}{2a}, 1\} + 1.$

Since $C_{a,b}$, $f_{a,b}(t)$ are oscillating, from lemma 1.3, we can determine the inequalities in Algorithm 1 in finite time.

2 Algorithm

Now we write down our algorithm to sample $\tau_{a,b}$. We divide our algorithm into two phases. In Algorithm 2, we decide whether $\tau_{a,b}$ is finite or not. And in Algorithm 3, we sample $\tau_{a,b}$ under the condition $[\tau_{a,b} < \infty]$. In Algorithm 2 and Algorithm 3, we write $C^{(N)}$, $F^{(N)}$ to denote the partial sums of $C_{a,b}$ and $f_{a,b}(t)$, respectively. Algorithm 2 To determine whether $\tau_{a,b}$ is finite

1: Generate $U \sim U(0, 1)$ 2: N = 13: Resolved = FALSE 4: repeat if $C^{(N)} \vee C^{(N+1)} < U$ then 5: Resolved = TRUE 6: Finite = FALSE 7: else if $C^{(N)} \wedge C^{(N+1)} > U$ then 8: Resolved = TRUE 9: 10: Finite = TRUEend if 11: N = N + 112: 13: until Resolved

14: **return** Finite

Algorithm 3 To generate $\tau_{a,b}$ given that $\tau_{a,b} < \infty$

1: repeat Generate $U \sim U(0, 1)$ and $V \sim \text{Gamma}(\alpha, b^2/2)$ 2: $\triangleright N(V) = \max\{\log 6/4ab, (b/2a)V, 1\} + 1$ $N = \lceil N(V) \rceil$ 3: Resolved = FALSE 4: repeat 5: if $F^{(N)}(V) \vee F^{(N+1)}(V) < C_{a,b}Mg(V; \alpha, b^2/2)U$ then 6: Resolved = TRUE 7: Accept = FALSE 8: else if $F^{(N)}(V) \wedge F^{(N+1)}(V) > C_{a,b}Mg(V; \alpha, b^2/2)U$ then 9: Resolved = TRUE 10: Accept = TRUE 11: end if 12: N = N + 113: until Resolved 14. 15: until Accept 16: return V

References

- [1] Jong Mun Lee. Efficient simulation for greeks and bridges under diffusion models. PhD thesis, KAIST, 2017.
- [2] Mario Abundo. Some conditional crossing results of brownian motion over a piecewise-linear boundary. *Statistics* & *probability letters*, 58(2):131–145, 2002.
- [3] Zaeem A Burq and Owen D Jones. Simulation of brownian motion at first-passage times. *Mathematics and Computers in Simulation*, 77(1):64–71, 2008.
- [4] Nan Chen and Zhengyu Huang. Localization and exact simulation of brownian motion-driven stochastic differential equations. *Mathematics of Operations Research*, 38(3):591–616, 2013.