

ROTA-BAXTER SYSTEMS ON HOM-ASSOCIATIVE ALGEBRAS AND COVARIANT HOM-BIALGEBRAS.

APURBA DAS

ABSTRACT. Rota-Baxter systems were introduced by Brzeziński as a generalization of Rota-Baxter operators that are related to dendriform structures, associative Yang-Baxter pairs and covariant bialgebras. In this paper, we define Rota-Baxter systems on Hom-associative algebras and show how they induce Hom-dendriform structures and weak pseudotwistors on the underlying Hom-associative algebra. We introduce Hom-Yang-Baxter pairs and a notion of covariant Hom-bialgebra. Given a Hom-Yang-Baxter pair, we construct a twisted Rota-Baxter system and a (quasitriangular) covariant Hom-bialgebra. Finally, we consider perturbations of the coproduct in a covariant Hom-bialgebra.

CONTENTS

1. Introduction	1
2. Hom-associative algebras and infinitesimal Hom-bialgebras	2
3. Rota-Baxter systems on Hom-associative algebras	3
4. Hom-Yang-Baxter pairs	5
5. Covariant Hom-bialgebras	7
References	14

1. INTRODUCTION

Rota-Baxter operators are algebraic abstraction of the integral operator. A Rota-Baxter operator on an algebra A is a linear map $R : A \rightarrow A$ that satisfies

$$R(a) \cdot R(b) = R(R(a) \cdot b + a \cdot R(b)), \quad \text{for } a, b \in A.$$

They were first appeared in the fluctuation theory in probability by G. Baxter [4] and further studied by G.-C. Rota [24] and P. Cartier [6] in relationship with combinatorics. An important application of Rota-Baxter operators were found in the Connes-Kreimer's algebraic approach of the renormalization of quantum field theory [7]. In [3] M. Aguiar relates Rota-Baxter operators with dendriform algebras introduced by J.-L. Loday [21]. See [10,14–16] for some recent study on Rota-Baxter operators.

On the other hand, the notion of infinitesimal bialgebras was introduced by Joni and Rota [18] to give an algebraic framework for the calculus of Newton divided differences and further studied by Aguiar [2]. Infinitesimal bialgebras are the associative analog of Lie bialgebras and related to Rota-Baxter operators. Motivated from the classical Yang-Baxter equation in Lie algebra context, in [2], Aguiar introduced an associative analog of Yang-Baxter equation. A solution of the associative Yang-Baxter equation induces a Rota-Baxter operator and an infinitesimal bialgebra, called quasitriangular infinitesimal bialgebra.

In [5] Brzeziński introduced Rota-Baxter systems as a generalization of Rota-Baxter operator that are also related to dendriform structures. Brzeziński also considered a notion of associative Yang-Baxter pair (generalization of associative Yang-Baxter solution), a notion of covariant bialgebra (generalization of infinitesimal bialgebra) and find various relations between them. These relations are generalization of Aguiar's results that can be mentioned as follows:

Key words and phrases. Hom-associative algebras, Rota-Baxter systems, Hom-Yang-Baxter pairs, Covariant Hom-bialgebras.
2020 MSC. 17B38, 17B61, 16T10

infinitesimal bialgebra \longleftrightarrow associative Yang-Baxter solution \longleftrightarrow Rota-Baxter operator \longleftrightarrow
 Dendriform algebra
 covariant bialgebra \longleftrightarrow associative Yang-Baxter pair \longleftrightarrow Rota-Baxter system \longleftrightarrow Dendriform algebra.

Recently, Hom-type algebras are widely studied in the literature by many authors. In these algebras, the identities defining the structures are twisted by linear maps [23]. The notion of Hom-Lie algebras was introduced by Hartwig, Larsson and Silvestrov that appeared in examples of q -deformations of the Witt and Virasoro algebras [17]. Other types of algebras (e.g. associative, Leibniz, dendriform, Poisson etc.) twisted by linear maps have also been studied extensively [1,8,11,22,23]. Here we mainly put our attention to Hom-associative algebras. Various aspects of Hom-associative algebras were studied in [1,8,9]. In [25] Yau defined associative Hom-Yang-Baxter equation, infinitesimal Hom-bialgebras and generalize some results of Aguiar [2]. Rota-Baxter operators on Hom-associative algebras was defined by Makhlouf in [22] and relations with Hom-dendriform algebras and Hom-preLie algebras was obtained.

In this paper, we define Rota-Baxter system on Hom-associative algebras generalizing Rota-Baxter Hom-associative algebras. We show that a Rota-Baxter system induces a Hom-dendriform structure and therefore a Hom-preLie algebra structure. We show that a Rota-Baxter system on a Hom-associative algebra induces a weak pseudotwistor on the underlying Hom-associative algebra in the sense of [20].

Next, we introduce an associative Hom-Yang-Baxter pair which are Hom-analog of Yang-Baxter pairs of Brzeziński [5]. Associative Hom-Yang-Baxter pairs generalize associative Hom-Yang-Baxter solutions of [25]. We show that a Hom-Yang-Baxter pair induces a twisted version of Rota-Baxter system on the underlying Hom-associative algebra. Hence, by previous results, a Hom-Yang-Baxter pair induces a Hom-dendriform algebra and a Hom-preLie algebra.

Finally, we define a notion of covariant Hom-bialgebra. They are generalization of covariant bialgebras and infinitesimal Hom-bialgebras. Given a covariant bialgebra and a morphism, one can construct a covariant Hom-bialgebra, called ‘induced by composition’. We show that a Hom-Yang-Baxter pair induces a covariant Hom-bialgebra, called quasitriangular covariant Hom-bialgebra. We give some characterizations of covariant Hom-bialgebras. Finally, we consider perturbations of the coproduct in a covariant Hom-bialgebra generalizing the perturbation theory of Drinfel’d for quasi-Hopf algebras [12,13].

All vector spaces, linear maps and tensor products are over any field \mathbb{K} .

2. HOM-ASSOCIATIVE ALGEBRAS AND INFINITESIMAL HOM-BIALGEBRAS

In this section, we recall Hom-associative algebras and infinitesimal Hom-bialgebras. Our main references are [23,25].

A Hom-vector space is a pair (A, α) in which A is a vector space and $\alpha : A \rightarrow A$ is a linear map. A morphism between Hom-vector spaces (A, α) and (B, β) is given by a linear map $f : A \rightarrow B$ satisfying $\beta \circ f = f \circ \alpha$. Hom-vector spaces and morphisms between them forms a category.

2.1. Definition. A Hom-associative algebra consists of a Hom-vector space (A, α) together with a bilinear map $\mu : A \otimes A \rightarrow A$, $a \otimes b \mapsto a \cdot b$ satisfying

$$(a \cdot b) \cdot \alpha(c) = \alpha(a) \cdot (b \cdot c), \quad \text{for } a, b, c \in A.$$

A Hom-associative algebra as above may be denoted by (A, α, μ) or simply by A .

A Hom-associative algebra (A, α, μ) is said to be multiplicative if α preserves the multiplication, i.e. $\alpha(a \cdot b) = \alpha(a) \cdot \alpha(b)$, for $a, b \in A$. From now onward, by a Hom-associative algebra, we shall always mean a multiplicative Hom-associative algebra. A Hom-associative algebra (A, α, μ) with $\alpha = \text{id}$ is simply an associative algebra. Therefore, the class of associative algebras is a subclass of the class of Hom-associative algebras.

2.2. Definition. Let (A, μ) be an associative algebra and $\alpha : A \rightarrow A$ be an associative algebra morphism. Then $(A, \alpha, \mu_\alpha = \alpha \circ \mu)$ is a Hom-associative algebra, called ‘induced by composition’.

The dual notion of Hom-associative algebras is given by Hom-associative coalgebras.

2.3. Definition. A Hom-coassociative coalgebra is a Hom-vector space (C, α) together with a linear map $\Delta : C \rightarrow C \otimes C$ satisfying

$$(1) \quad (\Delta \otimes \alpha) \circ \Delta = (\alpha \otimes \Delta) \circ \Delta.$$

A Hom-coassociative coalgebra as above may be denoted by (C, α, Δ) or simply by C . A Hom-coassociative coalgebra (C, α, Δ) is called multiplicative if $(\alpha \otimes \alpha) \circ \Delta = \Delta \circ \alpha$.

In the rest of the paper, by a Hom-coassociative coalgebra, we shall always mean a multiplicative Hom-coassociative coalgebra.

2.4. Definition. An infinitesimal Hom-bialgebra is a quadruple (A, α, μ, Δ) in which (A, α, μ) is a Hom-associative algebra, (A, α, Δ) is a Hom-coassociative coalgebra and satisfying the following compatibility

$$(2) \quad \Delta \circ \mu = (\mu \otimes \alpha) \circ (\alpha \otimes \Delta) + (\alpha \otimes \mu) \circ (\Delta \otimes \alpha).$$

Using Sweedler's notation $\Delta(a) = a_{(1)} \otimes a_{(2)}$ (note that we do not use the summation sign in the Sweedler's notation which is hidden in the picture), the compatibility (2) can be expressed as

$$\Delta(ab) = \alpha(a) \cdot b_{(1)} \otimes \alpha(b_{(2)}) + \alpha(a_{(1)}) \otimes a_{(2)} \cdot \alpha(b), \quad \text{for } a, b \in A.$$

Infinitesimal Hom-bialgebras are Hom-analog of infinitesimal bialgebras introduced by Joni and Rota [18], further studied by Aguiar [2]. More precisely, an infinitesimal Hom-bialgebra (A, α, μ, Δ) with $\alpha = \text{id}$ is an infinitesimal bialgebra and a morphism between infinitesimal bialgebras gives rise to an infinitesimal Hom-bialgebra induced by composition [25].

3. ROTA-BAXTER SYSTEMS ON HOM-ASSOCIATIVE ALGEBRAS

In this section, we introduce Rota-Baxter systems on Hom-associative algebras and prove that they induce Hom-dendriform algebras. A Rota-Baxter system on a Hom-associative algebra gives rise to a weak pseudotwistor on the underlying Hom-associative algebra in the sense of [20].

3.1. Definition. Let (A, α, μ) be a Hom-associative algebra. A Rota-Baxter system on A consists of a pair (R, S) of linear maps on A satisfying $\alpha \circ R = R \circ \alpha$, $\alpha \circ S = S \circ \alpha$ and

$$(3) \quad R(a) \cdot R(b) = R(R(a) \cdot b + a \cdot S(b)),$$

$$(4) \quad S(a) \cdot S(b) = S(R(a) \cdot b + a \cdot S(b)).$$

If the Hom-associative algebra is given by an associative algebra (i.e. $\alpha = \text{id}$), one gets Rota-Baxter system on associative algebra introduced by Brzeziński [5].

In [22] the author introduced Rota-Baxter operators on Hom-associative algebras. A linear map $R : A \rightarrow A$ is a Rota-Baxter operator of weight $\lambda \in \mathbb{K}$ on a Hom-associative algebra (A, α, μ) if $\alpha \circ R = R \circ \alpha$ and satisfying

$$R(a) \cdot R(b) = R(R(a) \cdot b + a \cdot R(b) + \lambda a \cdot b), \quad \text{for } a, b \in A.$$

It follows that a Rota-Baxter operator R of weight 0 is a Rota-Baxter system (R, R) . More generally, if R is a Rota-Baxter operator of weight λ , then $(R, S = R + \lambda \text{id})$ is a Rota-Baxter system. First observe that (3) holds obviously for $R = R$ and $S = R + \lambda \text{id}$. To prove (4), we see that

$$\begin{aligned} S(R(a) \cdot b + a \cdot S(b)) &= (R + \lambda \text{id})(R(a) \cdot b + a \cdot R(b) + \lambda a \cdot b) \\ &= R(R(a) \cdot b + a \cdot R(b) + \lambda a \cdot b) + \lambda R(a) \cdot b + \lambda a \cdot R(b) + \lambda^2 a \cdot b \\ &= R(a) \cdot R(b) + \lambda R(a) \cdot b + \lambda a \cdot R(b) + \lambda^2 a \cdot b \\ &= (R(a) + \lambda a) \cdot (R(b) + \lambda b) = S(a) \cdot S(b). \end{aligned}$$

3.2. Definition. [11,22] A Hom-dendriform algebra is a Hom-vector space (D, α) together with bilinear maps $\prec, \succ : D \otimes D \rightarrow D$ satisfying $\alpha(a \prec b) = \alpha(a) \prec \alpha(b)$, $\alpha(a \succ b) = \alpha(a) \succ \alpha(b)$ and the following

identities are hold

$$\begin{aligned}(a \prec b) \prec \alpha(c) &= \alpha(a) \prec (b \prec c + b \succ c), \\ (a \succ b) \prec \alpha(c) &= \alpha(a) \succ (b \prec c), \\ (a \prec b + a \succ b) \succ \alpha(c) &= \alpha(a) \succ (b \succ c).\end{aligned}$$

A Hom-dendriform algebra as above may be denoted by $(D, \alpha, \prec, \succ)$. Hom-dendriform algebras Hom-analog of Loday's dendriform algebras [21].

If $(D, \alpha, \prec, \succ)$ is a Hom-dendriform algebra then it follows that $(D, \alpha, *)$ is a Hom-associative algebra, where $a * b = a \prec b + a \succ b$, for $a, b \in D$.

3.3. Definition. [22] A Hom-preLie algebra is a Hom-vector space (A, α) together with a bilinear map $\diamond : A \otimes A \rightarrow A$ that satisfies $\alpha(a \diamond b) = \alpha(a) \diamond \alpha(b)$ and the following identity

$$(a \diamond b) \diamond \alpha(c) - \alpha(a) \diamond (b \diamond c) = (b \diamond a) \diamond \alpha(c) - \alpha(b) \diamond (a \diamond c), \quad \text{for } a, b, c \in A.$$

The following result can be found in [22].

3.4. Proposition. Let $(D, \alpha, \prec, \succ)$ be a Hom-dendriform algebra. Then the triple (D, α, Δ) is a Hom-preLie algebra, where $a \diamond b = a \succ b - a \prec b$, for $a, b \in D$.

3.5. Proposition. Let (A, α, μ) be a Hom-associative algebra and (R, S) be a Rota-Baxter system. Then $(A, \alpha, \prec, \succ)$ is a Hom-dendriform algebra, where $a \prec b = a \cdot S(b)$ and $a \succ b = R(a) \cdot b$, for $a, b \in A$.

Proof. First observe that

$$\begin{aligned}\alpha(a \prec b) &= \alpha(a \cdot S(b)) = \alpha(a) \cdot S(\alpha(b)) = \alpha(a) \prec \alpha(b), \\ \alpha(a \succ b) &= \alpha(R(a) \cdot b) = R(\alpha(a)) \cdot \alpha(b) = \alpha(a) \succ \alpha(b).\end{aligned}$$

We have

$$\begin{aligned}(a \prec b) \prec \alpha(c) &= (a \cdot S(b)) \cdot \alpha(S(c)) = \alpha(a) \cdot (S(b) \cdot S(c)) \\ &= \alpha(a) \cdot S(b \cdot S(c) + R(b) \cdot c) \\ &= \alpha(a) \prec (b \prec c + b \succ c).\end{aligned}$$

Similarly,

$$(a \succ b) \prec \alpha(c) = (R(a) \cdot b) \cdot \alpha(S(c)) = R(\alpha(a)) \cdot (b \cdot S(c)) = \alpha(a) \succ (b \prec c),$$

and

$$\begin{aligned}(a \prec b + a \succ b) \succ \alpha(c) &= R(a \cdot S(b) + R(a) \cdot b) \cdot \alpha(c) = (R(a) \cdot R(b)) \cdot \alpha(c) \\ &= R(\alpha(a)) \cdot (R(b) \cdot c) \\ &= \alpha(a) \succ (b \succ c).\end{aligned}$$

Hence the proof. □

3.6. Remark. (i) A Rota-Baxter system (R, S) on a Hom-associative algebra (A, α, μ) induces a new Hom-associative structure on (A, α) with multiplication

$$(5) \quad a * b = a \cdot S(b) + R(a) \cdot b, \quad \text{for } a, b \in A.$$

(ii) It follows from the above Proposition and Proposition 3.4 that a Rota-Baxter system (R, S) on a Hom-associative algebra (A, α, μ) induces a Hom-preLie algebra (A, α, \diamond) , where $a \diamond b = R(a) \cdot b - a \cdot S(b)$, for $a, b \in A$.

3.7. Definition. [20] Let (A, α, μ) be a Hom-associative algebra. A linear map $T : A^{\otimes 2} \rightarrow A^{\otimes 2}$ is called a weak pseudotwistor if $\alpha^{\otimes 2} \circ T = T \circ \alpha^{\otimes 2}$ and there exists a linear map $\tau : A^{\otimes 3} \rightarrow A^{\otimes 3}$ (called a weak

companion of T) with $\alpha^{\otimes 3} \circ \tau = \tau \circ \alpha^{\otimes 3}$ that makes the following diagram commutative

$$(6) \quad \begin{array}{ccccc} & & A \otimes A \otimes A & \xrightarrow{\alpha \otimes \mu} & A \otimes A & \xleftarrow{\mu \otimes \alpha} & A \otimes A \otimes A & & \\ & \nearrow \text{id} \otimes T & & & \downarrow T & & & \nwarrow T \otimes \text{id} & \\ A \otimes A \otimes A & & & & & & & & A \otimes A \\ & \searrow \tau & & & & & & \swarrow \tau & \\ & & A \otimes A \otimes A & \xrightarrow{\alpha \otimes \mu} & A \otimes A & \xleftarrow{\mu \otimes \alpha} & A \otimes A \otimes A & & \end{array}$$

In this case, it can be seen that $(A, \alpha, \mu \circ T)$ is a new Hom-associative algebra [20].

3.8. Proposition. *Let (R, S) be a Rota-Baxter system on a Hom-associative algebra (A, α, μ) . Then the map $T : A^{\otimes 2} \rightarrow A^{\otimes 2}$ defined by*

$$T(a \otimes b) = R(a) \otimes b + a \otimes S(b)$$

is a weak pseudotwistor.

Proof. Let us define a map $\tau : A^{\otimes 3} \rightarrow A^{\otimes 3}$ by

$$\tau(a \otimes b \otimes c) = R(a) \otimes R(b) \otimes c + R(a) \otimes b \otimes S(c) + a \otimes S(b) \otimes S(c).$$

We show that T is a weak pseudotwistor with a weak companion τ . First observe that $\alpha^{\otimes 2} \circ T = T \circ \alpha^{\otimes 2}$ and $\alpha^{\otimes 3} \circ \tau = \tau \circ \alpha^{\otimes 3}$. Next we have

$$(7) \quad \begin{aligned} ((\mu \otimes \alpha) \circ \tau)(a \otimes b \otimes c) &= R(a) \cdot R(b) \otimes \alpha(c) + R(a) \cdot b \otimes S(\alpha(c)) + a \cdot S(b) \otimes S(\alpha(c)) \\ &= R(R(a) \cdot b) \otimes \alpha(c) + R(a \cdot S(b)) \otimes \alpha(c) + R(a) \cdot b \otimes S(\alpha(c)) + a \cdot S(b) \otimes S(\alpha(c)). \end{aligned}$$

On the other hand,

$$(8) \quad \begin{aligned} (T \circ (\mu \otimes \alpha) \circ (T \otimes \text{id}))(a \otimes b \otimes c) &= (T \circ (\mu \otimes \alpha))(R(a) \otimes b \otimes c + a \otimes S(b) \otimes c) \\ &= T(R(a) \cdot b \otimes \alpha(c) + a \cdot S(b) \otimes \alpha(c)) \\ &= R(R(a) \cdot b) \otimes \alpha(c) + R(a) \cdot b \otimes S(\alpha(c)) \\ &\quad + R(a \cdot S(b)) \otimes \alpha(c) + a \cdot S(b) \otimes S(\alpha(c)). \end{aligned}$$

It follows from (7) and (8) that the right hand side pentagon of (6) is commutative. Similarly,

$$\begin{aligned} ((\alpha \otimes \mu) \circ \tau)(a \otimes b \otimes c) &= R(\alpha(c)) \otimes R(b) \cdot c + R(\alpha(a)) \otimes b \cdot S(c) + \alpha(a) \otimes S(R(b) \cdot c) + \alpha(a) \otimes S(b \cdot S(c)) \\ &= (T \circ (\alpha \otimes \mu) \circ (\text{id} \otimes T))(a \otimes b \otimes c). \end{aligned}$$

This implies that the left-hand side pentagon of (6) is also commutative. Hence the proof. \square

The corresponding Hom-associative algebra structure on the Hom-vector space (A, α) induced from the above weak pseudotwistor coincides with (5).

4. HOM-YANG-BAXTER PAIRS

Let (A, α, μ) be a Hom-associative algebra. An element $r = r_{(1)} \otimes r_{(2)} \in A^{\otimes 2}$ is called α -invariant if $(\alpha \otimes \alpha)(r) = r$, or equivalently, $\alpha(r_{(1)}) \otimes \alpha(r_{(2)}) = r_{(1)} \otimes r_{(2)}$. As before, we omit the summation here. Let r, s be two α -invariant elements of $A^{\otimes 2}$. Define

$$\begin{aligned} r_{13}s_{12} &= r_{(1)} \cdot s_{(1)} \otimes \alpha(s_{(2)}) \otimes \alpha(r_{(2)}), \\ r_{12}s_{23} &= \alpha(r_{(1)}) \otimes r_{(2)} \cdot s_{(1)} \otimes \alpha(s_{(2)}), \\ r_{23}s_{13} &= \alpha(s_{(1)}) \otimes \alpha(r_{(1)}) \otimes r_{(2)} \cdot s_{(2)}. \end{aligned}$$

4.1. Definition. Let (A, α, μ) be a Hom-associative algebra. A pair (r, s) of two α -invariant elements of $A^{\otimes 2}$ is said to be a Hom-Yang-Baxter pair if the following conditions are hold:

$$\begin{cases} r_{13}r_{12} - r_{12}r_{23} + s_{23}r_{13} = 0, \\ s_{13}r_{12} - s_{12}s_{23} + s_{23}s_{13} = 0. \end{cases}$$

When A is an associative algebra (i.e. $\alpha = \text{id}$), one recover the notion of Yang-Baxter pairs introduced by Brzeziński [5]. On the other hand, if $r = s$, one gets the Hom-Yang-Baxter equation introduced by Yau [25].

Let (A, μ) be an associative algebra and (r, s) be an associative Yang-Baxter pair. If $\alpha : A \rightarrow A$ is an algebra morphism such that both r, s are α -invariant, then (r, s) is a Hom-Yang-Baxter pair on the Hom-associative algebra $(A, \alpha, \mu_\alpha = \alpha \circ \mu)$.

4.2. Example. Let (A, μ) be an associative algebra and $a, b \in A$ be two elements such that $a^2 = b^2 = ab = ba = 0$. Then $(r = a \otimes a, s = b \otimes b)$ is an associative Yang-Baxter pair. If $\alpha : A \rightarrow A$ is an algebra map fixing a and b (i.e. $\alpha(a) = a$ and $\alpha(b) = b$), then (r, s) is a Hom-Yang-Baxter pair on the Hom-associative algebra (A, α, μ_α) .

Let (A, μ) be an unital associative algebra (with unit 1) and $a, b \in A$ be such that $b^2 = ba = 0$. Then $(r = 1 \otimes a, s = 1 \otimes b)$ is an associative Yang-Baxter pair. Thus, if $\alpha : A \rightarrow A$ is an unital algebra morphism fixing a and b , then (r, s) is a Hom-Yang-Baxter pair on the Hom-associative algebra (A, α, μ_α) .

In [5] the author constructs a Rota-Baxter system from a Yang-Baxter pair in an associative algebra. In the case of Hom-associative algebras, we do not get a Rota-Baxter system on the underlying Hom-associative algebra. Rather, we get a certain twisted version of Rota-Baxter system.

4.3. Definition. Let (A, α, μ) be a Hom-associative algebra. A pair (R, S) of linear maps on A is said to be an (α^n) -Rota-Baxter system if $\alpha \circ R = R \circ \alpha$, $\alpha \circ S = S \circ \alpha$ and satisfies

$$\begin{aligned} R(\alpha^n(a)) \cdot R(\alpha^n(b)) &= R(R(a) \cdot \alpha^n(b) + \alpha^n(a) \cdot S(b)), \\ S(\alpha^n(a)) \cdot S(\alpha^n(b)) &= S(R(a) \cdot \alpha^n(b) + \alpha^n(a) \cdot S(b)). \end{aligned}$$

Thus, an (α^0) -Rota-Baxter system is simply a Rota-Baxter system (Definition 3.1). The notion of (α^n) -Rota-Baxter system is a generalization of (α^n) -Rota-Baxter operator of [19].

In similar to Proposition 3.5, one can show that if (R, S) is an (α^n) -Rota-Baxter system on a Hom-associative algebra (A, α, μ) then $(A, \alpha^{n+1}, \prec, \succ)$ is a Hom-dendriform algebra, where $a \prec b = \alpha^n(a) \cdot S(b)$ and $a \succ b = R(a) \cdot \alpha^n(b)$, for $a, b \in A$. As a consequence, $(A, \alpha^{n+1}, *)$ is a Hom-associative algebra and $(A, \alpha^{n+1}, \diamond)$ is a Hom-preLie algebra, where

$$a * b = R(a) \cdot \alpha^n(b) + \alpha^n(a) \cdot S(b) \quad \text{and} \quad a \diamond b = R(a) \cdot \alpha^n(b) - \alpha^n(a) \cdot S(b).$$

4.4. Theorem. Let (A, α, μ) be a Hom-associative algebra and (r, s) be a Hom-Yang-Baxter pair. Define $R, S : A \rightarrow A$ by

$$\begin{aligned} R(a) &:= (r_{(1)} \cdot a) \cdot \alpha(r_{(2)}) = \alpha(r_{(1)}) \cdot (a \cdot r_{(2)}), \\ S(a) &:= (s_{(1)} \cdot a) \cdot \alpha(s_{(2)}) = \alpha(s_{(1)}) \cdot (a \cdot s_{(2)}). \end{aligned}$$

Then (R, S) is a (α^2) -Rota-Baxter system on the Hom-associative algebra (A, α, μ) .

Proof. First we have

$$R(\alpha(a)) = (r_{(1)} \cdot \alpha(a)) \cdot \alpha(r_{(2)}) = (\alpha(r_{(1)}) \cdot \alpha(a)) \cdot \alpha^2(r_{(2)}) = \alpha(R(a)).$$

Similarly, we have $S(\alpha(a)) = \alpha(S(a))$. Next, we note that the condition $r_{13}r_{12} - r_{12}r_{23} + s_{23}r_{13} = 0$, or equivalently,

$$r_{(1)} \cdot \tilde{r}_{(1)} \otimes \alpha(\tilde{r}_{(2)}) \otimes \alpha(r_{(2)}) - \alpha(r_{(1)}) \otimes r_{(2)} \cdot \tilde{r}_{(1)} \otimes \alpha(\tilde{r}_{(2)}) + \alpha(r_{(1)}) \otimes \alpha(s_{(1)}) \otimes s_{(2)} \cdot r_{(2)} = 0$$

implies that

$$(9) \quad \begin{aligned} &(((r_{(1)} \cdot \tilde{r}_{(1)}) \cdot \alpha^2(a)) \cdot \alpha^2(\tilde{r}_{(2)})) \cdot (\alpha^3(b) \cdot \alpha^2(r_{(2)})) - ((\alpha(r_{(1)}) \cdot \alpha^2(a)) \cdot \alpha(r_{(2)} \cdot \tilde{r}_{(1)})) \cdot (\alpha^3(b) \cdot \alpha^2(\tilde{r}_{(2)})) \\ &\quad + ((\alpha(r_{(1)}) \cdot \alpha^2(a)) \cdot \alpha^2(s_{(1)})) \cdot (\alpha^3(b) \cdot \alpha(s_{(2)} \cdot r_{(2)})) = 0. \end{aligned}$$

We have

$$\begin{aligned} R(\alpha^2(a)) \cdot R(\alpha^2(b)) &= ((r_{(1)} \cdot \alpha^2(a)) \cdot \alpha(r_{(2)})) \cdot (\alpha(\tilde{r}_{(1)}) \cdot (\alpha^2(b) \cdot \tilde{r}_{(2)})) \\ &= (((\alpha(r_{(1)}) \cdot \alpha^2(a)) \cdot \alpha^2(r_{(2)})) \cdot (\alpha(\tilde{r}_{(1)}) \cdot (\alpha^2(b) \cdot \tilde{r}_{(2)}))) \\ &= (((r_{(1)} \cdot \alpha(a)) \cdot \alpha(r_{(2)})) \cdot \alpha(\tilde{r}_{(1)})) \cdot (\alpha^3(b) \cdot \alpha(\tilde{r}_{(2)})) \\ &= (((\alpha(r_{(1)}) \cdot \alpha^2(a)) \cdot (\alpha(r_{(2)}) \cdot \tilde{r}_{(1)})) \cdot (\alpha^3(b) \cdot \alpha(\tilde{r}_{(2)}))) \\ &= (((\alpha(r_{(1)}) \cdot \alpha^2(a)) \cdot \alpha(r_{(2)} \cdot \tilde{r}_{(1)})) \cdot (\alpha^3(b) \cdot \alpha^2(\tilde{r}_{(2)}))). \end{aligned}$$

Similarly,

$$\begin{aligned} R(R(a) \cdot \alpha^2(b)) &= \alpha(r_{(1)}) \cdot (((\tilde{r}_{(1)} \cdot a) \cdot \alpha(\tilde{r}_{(2)})) \cdot \alpha^2(b)) \cdot r_{(2)} \\ &= \alpha^2(r_{(1)}) \cdot (((\tilde{r}_{(1)} \cdot a) \cdot \alpha(\tilde{r}_{(2)})) \cdot \alpha^2(b)) \cdot \alpha(r_{(2)}) \\ &= \alpha^2(r_{(1)}) \cdot (((\alpha(\tilde{r}_{(1)}) \cdot \alpha(a)) \cdot \alpha^2(\tilde{r}_{(2)})) \cdot (\alpha^2(b) \cdot r_{(2)})) \\ &= (\alpha(r_{(1)}) \cdot ((\alpha(\tilde{r}_{(1)}) \cdot \alpha(a)) \cdot \alpha^2(\tilde{r}_{(2)}))) \cdot (\alpha^3(b) \cdot \alpha(r_{(2)})) \\ &= ((r_{(1)} \cdot (\alpha(\tilde{r}_{(1)}) \cdot \alpha(a))) \cdot \alpha^3(\tilde{r}_{(2)})) \cdot (\alpha^3(b) \cdot \alpha(r_{(2)})) \\ &= (((r_{(1)} \cdot \alpha(\tilde{r}_{(1)})) \cdot \alpha^2(a)) \cdot \alpha^3(\tilde{r}_{(2)})) \cdot (\alpha^3(b) \cdot \alpha^2(r_{(2)})) \\ &= (((r_{(1)} \cdot \tilde{r}_{(1)}) \cdot \alpha^2(a)) \cdot \alpha^2(\tilde{r}_{(2)})) \cdot (\alpha^3(b) \cdot \alpha^2(r_{(2)})) \end{aligned}$$

and

$$\begin{aligned} R(\alpha^2(a) \cdot S(b)) &= \alpha(r_{(1)}) \cdot ((\alpha^2(a) \cdot (\alpha(s_{(1)}) \cdot (b \cdot s_{(2)}))) \cdot r_{(2)}) \\ &= \alpha(r_{(1)}) \cdot (((\alpha(a) \cdot \alpha(s_{(1)})) \cdot \alpha(b \cdot s_{(2)})) \cdot r_{(2)}) \\ &= \alpha^2(r_{(1)}) \cdot (((\alpha(a) \cdot \alpha(s_{(1)})) \cdot \alpha(b \cdot s_{(2)})) \cdot \alpha(r_{(2)})) \\ &= \alpha^2(r_{(1)}) \cdot ((\alpha^2(a) \cdot \alpha^2(s_{(1)})) \cdot (\alpha(b \cdot s_{(2)}) \cdot r_{(2)})) \\ &= (\alpha(r_{(1)}) \cdot (\alpha^2(a) \cdot \alpha^2(s_{(1)}))) \cdot (\alpha^2(b \cdot s_{(2)}) \cdot \alpha(r_{(2)})) \\ &= ((r_{(1)} \cdot \alpha^2(a)) \cdot \alpha^3(s_{(1)})) \cdot (\alpha^3(b) \cdot (\alpha^2(s_{(2)}) \cdot r_{(2)})) \\ &= ((\alpha(r_{(1)}) \cdot \alpha^2(a)) \cdot \alpha^2(s_{(1)})) \cdot (\alpha^3(b) \cdot \alpha(s_{(2)} \cdot r_{(2)})). \end{aligned}$$

Thus, it follows from (9) that

$$R(\alpha^2(a)) \cdot R(\alpha^2(b)) = R(R(a) \cdot \alpha^2(b) + \alpha^2(a) \cdot S(b))$$

holds. In a similar way, by using the condition $s_{13}r_{12} - s_{12}s_{23} + s_{23}s_{13} = 0$, we can deduce that

$$S(\alpha^2(a)) \cdot S(\alpha^2(b)) = S(R(a) \cdot \alpha^2(b) + \alpha^2(a) \cdot S(b)).$$

Hence the pair (R, S) is an (α^2) -Rota-Baxter system on the Hom-associative algebra (A, α, μ) . \square

It follows from the above theorem that a Hom-Yang-Baxter pair (r, s) induces a Hom-dendriform structure $(A, \alpha^3, \prec, \succ)$, where $a \prec b = \alpha^2(a) \cdot (\alpha(s_{(1)}) \cdot (b \cdot s_{(2)}))$ and $a \succ b = ((r_{(1)} \cdot a) \cdot \alpha(r_{(2)})) \cdot \alpha^2(b)$. The associated Hom-associative and Hom-preLie algebras are respectively given by $(A, \alpha^3, *)$ and (A, α^3, \diamond) , where

$$\begin{aligned} a * b &= ((r_{(1)} \cdot a) \cdot \alpha(r_{(2)})) \cdot \alpha^2(b) + \alpha^2(a) \cdot (\alpha(s_{(1)}) \cdot (b \cdot s_{(2)})), \\ a \diamond b &= ((r_{(1)} \cdot a) \cdot \alpha(r_{(2)})) \cdot \alpha^2(b) - \alpha^2(a) \cdot (\alpha(s_{(1)}) \cdot (b \cdot s_{(2)})). \end{aligned}$$

5. COVARIANT HOM-BIALGEBRAS

In this section, we introduce a notion of covariant Hom-bialgebra. This structure generalizes covariant bialgebras of Brzeziński [5] and infinitesimal Hom-bialgebras of Yau [25]. Given a Hom-Yang-Baxter pair, we construct a covariant Hom-bialgebra, called quasitriangular covariant Hom-bialgebra. Some characterizations

of quasitriangular covariant Hom-bialgebra are given. Finally, we consider the perturbations of the coproduct in a covariant Hom-bialgebra generalizing perturbations of quasi-Hopf algebra à la Drinfel'd [12,13].

Let (A, α, μ) be a Hom-associative algebra. A bimodule consists of a Hom-vector space (M, β) together with action maps $l : A \otimes M \rightarrow M$, $(a, m) \mapsto a \bullet m$ and $r : M \otimes A \rightarrow M$, $(m, a) \mapsto m \bullet a$ satisfying $\beta(a \bullet m) = \alpha(a) \bullet \beta(m)$, $\beta(m \bullet a) = \beta(m) \bullet \alpha(a)$ and the following bimodule identities

$$(a \cdot b) \bullet \beta(m) = \alpha(a) \bullet (b \bullet m), \quad (a \bullet m) \bullet \alpha(b) = \alpha(a) \bullet (m \bullet b), \quad (m \bullet a) \bullet \alpha(b) = \beta(m) \bullet (a \cdot b),$$

for $a, b \in A$ and $m \in M$. See [9] for details. A bimodule is often denoted by (M, β) or simply by M when the action maps are understood. Any Hom-associative algebra is a bimodule over itself, called adjoint bimodule. The tensor product $M = A \otimes A$ can also be given an A -bimodule structure with linear map $\beta = \alpha \otimes \alpha : A \otimes A \rightarrow A \otimes A$ and action maps

$$(10) \quad a \bullet (b \otimes c) = \alpha(a) \cdot b \otimes \alpha(c), \quad (b \otimes c) \bullet a = \alpha(b) \otimes c \cdot \alpha(a), \quad \text{for } a \in A, b \otimes c \in A \otimes A.$$

5.1. Definition. Let (A, α, μ) be a Hom-associative algebra and (M, β) be a bimodule over it. A linear map $d : A \rightarrow M$ is said to be a derivation on A with values in the bimodule M if $\beta \circ d = d \circ \alpha$ and satisfying

$$d(a \cdot b) = a \bullet d(b) + d(a) \bullet b, \quad \text{for } a, b \in A.$$

Hence a linear map $d : A \rightarrow A \otimes A$ is a derivation on A with values in the bimodule $A \otimes A$ if and only if $\alpha^{\otimes 2} \circ d = d \circ \alpha$ and D satisfies

$$d \circ \mu = (\mu \otimes \alpha) \circ (\alpha \otimes d) + (\alpha \otimes \mu) \circ (d \otimes \alpha).$$

Let (A, α, μ) be a Hom-associative algebra and $\delta_1, \delta_2 : A \rightarrow A \otimes A$ be two derivations on A with values in the A -bimodule $A \otimes A$.

5.2. Definition. A linear map $D : A \rightarrow A \otimes A$ is said to be a covariant derivation with respect to (δ_1, δ_2) if D satisfies $\alpha^{\otimes 2} \circ D = D \circ \alpha$ and

$$\begin{aligned} D \circ \mu &= (\mu \otimes \alpha) \circ (\alpha \otimes \delta_1) + (\alpha \otimes \mu) \circ (D \otimes \alpha), \\ D \circ \mu &= (\mu \otimes \alpha) \circ (\alpha \otimes D) + (\alpha \otimes \mu) \circ (\delta_2 \otimes \alpha). \end{aligned}$$

It follows that any derivation d is a covariant derivation with respect to (d, d) . We are now in a position to define covariant Hom-bialgebras.

5.3. Definition. A covariant Hom-bialgebra consists of a tuple $(A, \alpha, \mu, \Delta, \delta_1, \delta_2)$ in which

- (i) (A, α, μ) is a Hom-associative algebra;
- (ii) (A, α, Δ) is a Hom-coassociative coalgebra;
- (iii) the linear maps $\delta_1, \delta_2 : A \rightarrow A \otimes A$ are derivations on A with values in the A -bimodule $A \otimes A$;
- (iv) the map $\Delta : A \rightarrow A \otimes A$ is a covariant derivation with respect to (δ_1, δ_2) .

If $\alpha = \text{id}$, one recover the notion of covariant bialgebras introduced by Brzeziński [5]. On the other hand, if $\Delta = \delta_1 = \delta_2$, one recover infinitesimal Hom-bialgebras defined in [25]. Further, if $\alpha = \text{id}$, we get infinitesimal bialgebras [2,18].

5.4. Proposition. Let $(A, \mu, \Delta, \delta_1, \delta_2)$ be a covariant bialgebra and $\alpha : A \rightarrow A$ be a morphism of covariant bialgebras. Then $(A, \alpha, \mu_\alpha, \Delta_\alpha, (\delta_1)_\alpha, (\delta_2)_\alpha)$ is a covariant Hom-bialgebra, where $\Delta_\alpha = \Delta \circ \alpha$, $(\delta_1)_\alpha = \delta_1 \circ \alpha$ and $(\delta_2)_\alpha = \delta_2 \circ \alpha$.

Proof. Since $\alpha : (A, \mu) \rightarrow (A, \mu)$ is an algebra morphism and $\alpha : (A, \Delta) \rightarrow (A, \Delta)$ is a coassociative coalgebra morphism, it follows that (A, α, μ_α) is a Hom-associative algebra and $(A, \alpha, \Delta_\alpha)$ is a Hom-coassociative

coalgebra. Moreover, for $i = 1, 2$, we have

$$\begin{aligned}
 (\delta_i)_\alpha \circ \mu_\alpha &= (\alpha^2)^{\otimes 2} \circ \delta_i \circ \mu \\
 &= (\alpha^2)^{\otimes 2} \circ (\text{id} \otimes \mu) \circ (\delta_i \otimes \text{id}) + (\alpha^2)^{\otimes 2} \circ (\mu \otimes \text{id}) \circ (\text{id} \otimes \delta_i) \\
 &= (\alpha^2 \otimes \mu_\alpha \circ \alpha^{\otimes 2}) \circ (\delta_i \otimes \text{id}) + (\mu_\alpha \circ \alpha^{\otimes 2} \otimes \alpha^2) \circ (\text{id} \otimes \delta_i) \\
 &= (\alpha \otimes \mu_\alpha) \circ \alpha^{\otimes 3} \circ (\delta_i \otimes \text{id}) + (\mu_\alpha \otimes \alpha) \circ \alpha^{\otimes 3} \circ (\text{id} \otimes \delta_i) \\
 &= (\mu_\alpha \otimes \alpha) \circ (\alpha \otimes (\delta_i)_\alpha) + (\alpha \otimes \mu_\alpha) \circ ((\delta_i)_\alpha \otimes \alpha).
 \end{aligned}$$

This shows that $(\delta_1)_\alpha$ and $(\delta_2)_\alpha$ are both derivations. Finally,

$$\begin{aligned}
 \Delta_\alpha \circ \mu_\alpha &= (\alpha^2)^{\otimes 2} \circ \Delta \circ \mu \\
 &= (\alpha^2)^{\otimes 2} \circ (\mu \otimes \text{id}) \circ (\text{id} \otimes \delta_1) + (\alpha^2)^{\otimes 2} \circ (\text{id} \otimes \mu) \circ (\Delta \otimes \text{id}) \\
 &= (\mu_\alpha \circ \alpha^{\otimes 2} \otimes \alpha^2) \circ (\text{id} \otimes \delta_1) + (\alpha^2 \otimes \mu_\alpha \circ \alpha^{\otimes 2}) \circ (\Delta \otimes \text{id}) \\
 &= (\mu_\alpha \otimes \alpha) \circ \alpha^{\otimes 3} \circ (\text{id} \otimes \delta_1) + (\alpha \otimes \mu_\alpha) \circ \alpha^{\otimes 3} \circ (\Delta \otimes \text{id}) \\
 &= (\mu_\alpha \otimes \alpha) \circ (\alpha \otimes (\delta_1)_\alpha) + (\alpha \otimes \mu_\alpha) \circ ((\Delta)_\alpha \otimes \alpha).
 \end{aligned}$$

Similarly, we can show that $\Delta_\alpha \circ \mu_\alpha = (\mu_\alpha \otimes \alpha) \circ (\alpha \otimes \Delta_\alpha) + (\alpha \otimes \mu_\alpha) \circ ((\delta_2)_\alpha \otimes \alpha)$. Therefore, Δ_α is a covariant derivation with respect to $((\delta_1)_\alpha, (\delta_2)_\alpha)$. Hence $(A, \alpha, \mu_\alpha, \Delta_\alpha, (\delta_1)_\alpha, (\delta_2)_\alpha)$ is a covariant Hom-bialgebra. \square

The covariant Hom-bialgebra $(A, \alpha, \mu_\alpha, \Delta_\alpha, (\delta_1)_\alpha, (\delta_2)_\alpha)$ obtained in Proposition 5.4 is called ‘induced by composition’.

In the following, we introduce another type of Hom-bialgebras dual to covariant Hom-bialgebras. Before that, we require some more terminologies.

Let (C, α, Δ) be a Hom-coassociative coalgebra. A bicomodule over it consists of a Hom-vector space (M, β) together with coaction maps $\Delta^l : M \rightarrow C \otimes M$ and $\Delta^r : M \rightarrow M \otimes C$ satisfying $(\alpha \otimes \beta) \circ \Delta^l = \Delta^l \circ \beta$, $(\beta \otimes \alpha) \circ \Delta^r = \Delta^r \circ \beta$ and

$$(\alpha \otimes \Delta^l) \circ \Delta^l = (\Delta \otimes \beta) \circ \Delta^l, \quad (\alpha \otimes \Delta^r) \circ \Delta^l = (\Delta^l \otimes \alpha) \circ \Delta^r, \quad (\beta \otimes \Delta) \circ \Delta^r = (\Delta^r \otimes \alpha) \circ \Delta^r.$$

It follows that any Hom-coassociative coalgebra is a bicomodule over itself. The tensor product $M = C \otimes C$ with the linear map $\beta = \alpha \otimes \alpha : C \otimes C \rightarrow C \otimes C$ and coaction maps $\Delta^l : C \otimes C \rightarrow (C \otimes C) \otimes C$, $\Delta^r : C \otimes C \rightarrow (C \otimes C) \otimes C$ defined by

$$\Delta^l = ((\alpha \otimes \text{id}) \circ \Delta) \otimes \alpha, \quad \Delta^r = \alpha \otimes ((\text{id} \otimes \alpha) \circ \Delta)$$

is a bicomodule over C .

Given a Hom-coassociative coalgebra (C, α, Δ) and a bicomodule (M, β) , a linear map $D : M \rightarrow C$ is a coderivation on C with coefficients in M if $\alpha \circ D = D \circ \beta$ and satisfies

$$\Delta \circ D = (\text{id}_C \otimes D) \circ \Delta^l + (D \otimes \text{id}_C) \circ \Delta^r.$$

It follows that a linear map $\partial : C \otimes C \rightarrow C$ is a coderivation on C with coefficients in the bicomodule $C \otimes C$ if $\alpha \otimes \partial = \partial \circ (\alpha \otimes \alpha)$ and

$$\Delta \circ \partial = (\alpha \otimes \partial) \circ (\Delta \otimes \alpha) + (\partial \otimes \alpha) \circ (\alpha \otimes \Delta).$$

Let ∂_1, ∂_2 be two coderivations on C with coefficients in $C \otimes C$. Then a linear map $\mu : C \otimes C \rightarrow C$ is called a covariant coderivation with respect to (∂_1, ∂_2) if

$$\begin{aligned}
 \Delta \circ \mu &= (\alpha \otimes \partial_1) \circ (\Delta \otimes \alpha) + (\mu \otimes \alpha) \circ (\alpha \otimes \Delta), \\
 \Delta \circ \mu &= (\alpha \otimes \mu) \circ (\Delta \otimes \alpha) + (\partial_2 \otimes \alpha) \circ (\alpha \otimes \Delta).
 \end{aligned}$$

5.5. Definition. A dual covariant Hom-bialgebra is a tuple $(A, \alpha, \mu, \Delta, \partial_1, \partial_2)$ in which (A, α, μ) is a Hom-associative algebra, (A, α, Δ) is a Hom-coassociative coalgebra, the maps $\partial_1, \partial_2 : A \otimes A \rightarrow A$ are coderivations on A with coefficients in the bicomodule $A \otimes A$ such that $\mu : A \otimes A \rightarrow A$ is a covariant coderivation with respect to (∂_1, ∂_2) .

5.6. Proposition. *Let $(A, \alpha, \mu, \Delta, \delta_1, \delta_2)$ be a finite dimensional covariant Hom-bialgebra. Then the tuple $(A^*, \alpha^*, \Delta^*, \mu^*, \delta_1^*, \delta_2^*)$ is a dual covariant Hom-bialgebra, where*

$$\langle \Delta^*(\phi \otimes \psi), a \rangle = \langle \phi \otimes \psi, \Delta(a) \rangle, \quad \langle \mu^*(\phi), a \otimes b \rangle = \langle \phi, \mu(a \otimes b) \rangle, \quad \langle \delta_i^*(\phi \otimes \psi), a \rangle = \langle \phi \otimes \psi, \delta_i(a) \rangle, \quad \text{for } i = 1, 2.$$

Proof. Since (A, α, Δ) is a Hom-coassociative coalgebra, we have $(A^*, \alpha^*, \Delta^*)$ is a Hom-associative algebra. On the other hand, (A, α, μ) is a Hom-associative algebra and A is finite dimensional implies that (A^*, α^*, μ^*) is a Hom-coassociative coalgebra [23]. Next, for $i = 1, 2$, we observe that

$$\begin{aligned} \mu^* \circ \delta_i^* &= (\delta_i \circ \mu)^* = ((\mu \otimes \alpha) \circ (\alpha \otimes \delta_i))^* + ((\alpha \otimes \mu) \circ (\delta_i \otimes \alpha))^* \\ &= (\alpha^* \otimes \delta_i^*) \circ (\mu^* \otimes \alpha^*) + (\delta_i^* \otimes \alpha^*) \circ (\alpha^* \otimes \mu^*). \end{aligned}$$

This shows that δ_1^*, δ_2^* are coderivations on the Hom-coassociative coalgebra A^* with values in the bicomodule $A^* \otimes A^*$. Finally, it remains to prove that $\Delta^* : A^* \otimes A^* \rightarrow A^*$ is a covariant coderivation with respect to (δ_1^*, δ_2^*) . Note that

$$\begin{aligned} \mu^* \circ \Delta^* &= (\Delta \circ \mu)^* = ((\mu \otimes \alpha) \circ (\alpha \otimes \delta_1))^* + ((\alpha \otimes \mu) \circ (\Delta \otimes \alpha))^* \\ &= (\alpha^* \otimes \delta_1^*) \circ (\mu^* \otimes \alpha^*) + (\Delta^* \otimes \alpha^*) \circ (\alpha^* \otimes \mu^*). \end{aligned}$$

Similarly, we can show that $\mu^* \circ \Delta^* = (\alpha^* \otimes \Delta^*) \circ (\mu^* \otimes \alpha^*) + (\delta_2^* \otimes \alpha^*) \circ (\alpha^* \otimes \mu^*)$. Hence the proof. \square

Given a Hom-associative algebra (A, α, μ) and a Hom-Yang-Baxter pair (r, s) , we will construct a covariant Hom-bialgebra. We need some more notations. Let (A, α, μ) be a Hom-associative algebra. Then $M = A^{\otimes n}$ with the linear map $\beta = \alpha^{\otimes n} : A^{\otimes n} \rightarrow A^{\otimes n}$ can be given an A -bimodule structure with actions

$$\begin{aligned} a \bullet (b_1 \otimes \cdots \otimes b_n) &= \alpha(a) \cdot b_1 \otimes \alpha(b_2) \otimes \cdots \otimes \alpha(b_n), \\ (b_1 \otimes \cdots \otimes b_n) \bullet a &= \alpha(b_1) \otimes \cdots \otimes \alpha(b_{n-1}) \otimes b_n \cdot \alpha(a). \end{aligned}$$

This generalizes the A -bimodule structure on $A^{\otimes 2}$ given in (10). Moreover, for any $a \in A$ and $r = r_{(1)} \otimes r_{(2)} \in A^{\otimes 2}$, we define $ar, ra \in A^{\otimes 2}$ by

$$ar = a \cdot r_{(1)} \otimes \alpha(r_{(2)}) \quad \text{and} \quad ra = \alpha(r_{(1)}) \otimes r_{(2)} \cdot a.$$

5.7. Theorem. *Let (A, α, μ) be a Hom-associative algebra and $r, s \in A^{\otimes 2}$ be two α -invariant elements of $A^{\otimes 2}$. Define maps $\Delta', \delta_r, \delta_s : A \rightarrow A \otimes A$ by*

$$(11) \quad \Delta'(a) = ar - sa, \quad \delta_r(a) = ar - ra, \quad \delta_s(a) = as - sa.$$

Then $(A, \alpha, \mu, \Delta', \delta_r, \delta_s)$ is a covariant Hom-bialgebra if and only if

$$(12) \quad a \bullet (r_{13}r_{12} - r_{12}r_{23} + s_{23}r_{13}) = (s_{13}r_{12} - s_{12}s_{23} + s_{23}s_{13}) \bullet a.$$

Proof. First we have

$$\begin{aligned} \Delta'(\alpha(a)) &= \alpha(a) \cdot r_{(1)} \otimes \alpha(r_{(2)}) - \alpha(s_{(1)}) \otimes s_{(2)} \cdot \alpha(a) \\ &= \alpha(a) \cdot \alpha(r_{(1)}) \otimes \alpha^2(r_{(2)}) - \alpha^2(s_{(1)}) \otimes \alpha(s_{(2)}) \cdot \alpha(a) \\ &= (\alpha \otimes \alpha)(a \cdot r_{(1)} \otimes \alpha(r_{(2)}) - \alpha(s_{(1)}) \otimes s_{(2)} \cdot a) = (\alpha \otimes \alpha)\Delta'(a). \end{aligned}$$

This shows that Δ' commutes with α . Similarly, δ_r and δ_s also commutes with α . Next, we claim that δ_r is a derivation. We observe that

$$\begin{aligned} \delta_r \circ \mu(a \otimes b) &= \delta_r(a \cdot b) = (a \cdot b) \cdot r_{(1)} \otimes \alpha(r_{(2)}) - \alpha(r_{(1)}) \otimes r_{(2)} \cdot (a \cdot b) \\ &= (a \cdot b) \cdot \alpha(r_{(1)}) \otimes \alpha^2(r_{(2)}) - \alpha^2(r_{(1)}) \otimes \alpha(r_{(2)}) \cdot (a \cdot b) \\ (13) \quad &= \alpha(a) \cdot (b \cdot r_{(1)}) \otimes \alpha^2(r_{(2)}) - \alpha^2(r_{(1)}) \otimes (r_{(2)} \cdot a) \cdot \alpha(b). \end{aligned}$$

On the other hand,

$$\begin{aligned}
 & ((\mu \otimes \alpha) \circ (\alpha \otimes \delta_r) + (\alpha \otimes \mu) \circ (\delta_r \otimes \alpha))(a \otimes b) \\
 &= (\mu \otimes \alpha)(\alpha(a) \otimes b \cdot r_{(1)} \otimes \alpha(r_{(2)}) - \alpha(a) \otimes \alpha(r_{(1)}) \otimes r_{(2)} \cdot b) \\
 &\quad + (\alpha \otimes \mu)(a \cdot r_{(1)} \otimes \alpha(r_{(2)}) \otimes \alpha(b) - \alpha(r_{(1)}) \otimes r_{(2)} \cdot a \otimes \alpha(b)) \\
 (14) \quad &= \alpha(a) \cdot (b \cdot r_{(1)}) \otimes \alpha^2(r_{(2)}) - \alpha(a) \cdot \alpha(r_{(1)}) \otimes \alpha(r_{(2)} \cdot b) \\
 &\quad + \alpha(a \cdot r_{(1)}) \otimes \alpha(r_{(2)}) \cdot \alpha(b) - \alpha^2(r_{(1)}) \otimes (r_{(2)} \cdot a) \cdot \alpha(b).
 \end{aligned}$$

It follows from (13) and (14) that δ_r is a derivation. Replacing r by s , we get that δ_s is a derivation. Thus, to show that $(A, \alpha, \mu, \Delta', \delta_r, \delta_s)$ is a covariant Hom-bialgebra, we need to verify that Δ' is Hom-coassociative. In the next, we will see that the Hom-coassociativity of Δ' is equivalent to the condition (12) that proves the theorem. For $r = r_{(1)} \otimes r_{(2)}$ and $s = s_{(1)} \otimes s_{(2)}$, we have

$$\begin{aligned}
 (\Delta' \otimes \alpha)(\Delta' a) &= (\Delta' \otimes \alpha)(a \cdot r_{(1)} \otimes \alpha(r_{(2)}) - \alpha(s_{(1)}) \otimes s_{(2)} \cdot a) \\
 &= \Delta'(a \cdot r_{(1)}) \otimes \alpha^2(r_{(2)}) - \alpha^{\otimes 2}(\Delta'(s_{(1)})) \otimes \alpha(s_{(2)}) \cdot \alpha(a) \\
 &= \alpha(a) \cdot (r_{(1)} \cdot \tilde{r}_{(1)}) \otimes \alpha^2(\tilde{r}_{(2)}) \otimes \alpha^2(r_{(2)}) - \alpha^2(s_{(1)}) \otimes (s_{(2)} \cdot a) \cdot \alpha(r_{(1)}) \otimes \alpha^2(r_{(2)}) \\
 &\quad - \alpha(s_{(1)} \cdot \tilde{r}_{(1)}) \otimes \alpha^2(\tilde{r}_{(2)}) \otimes \alpha(s_{(2)}) \cdot \alpha(a) + \alpha^2(\tilde{s}_{(1)}) \otimes \alpha(\tilde{s}_{(2)} \cdot s_{(1)}) \otimes \alpha(s_{(2)}) \cdot \alpha(a) \\
 (15) \quad &= a \bullet (r_{13}r_{12}) - \alpha^2(s_{(1)}) \otimes (s_{(2)} \cdot a) \cdot \alpha(r_{(1)}) \otimes \alpha^2(r_{(2)}) - (s_{13}r_{12}) \bullet a + (s_{12}s_{23}) \bullet a.
 \end{aligned}$$

Similarly,

$$\begin{aligned}
 (\alpha \otimes \Delta')(\Delta' a) &= (\alpha \otimes \Delta')(a \cdot r_{(1)} \otimes \alpha(r_{(2)}) - \alpha(s_{(1)}) \otimes s_{(2)} \cdot a) \\
 &= \alpha(a) \cdot \alpha(r_{(1)}) \otimes \alpha^{\otimes 2}(\Delta'(r_{(2)})) - \alpha^2(s_{(1)}) \otimes \Delta'(s_{(2)} \cdot a) \\
 &= \alpha(a) \cdot \alpha(r_{(1)}) \otimes \alpha(r_{(2)} \cdot \tilde{r}_{(1)}) \otimes \alpha^2(\tilde{r}_{(2)}) - \alpha(a) \cdot \alpha(r_{(1)}) \otimes \alpha^2(s_{(1)}) \otimes \alpha(s_{(2)} \cdot r_{(2)}) \\
 &\quad - \alpha^2(s_{(1)}) \otimes \alpha(s_{(2)}) \cdot (a \cdot r_{(1)}) \otimes \alpha^2(r_{(2)}) + \alpha^2(s_{(1)}) \otimes \alpha^2(\tilde{s}_{(1)}) \otimes (\tilde{s}_{(2)} \cdot s_{(2)}) \cdot \alpha(a) \\
 (16) \quad &= a \bullet (r_{12}r_{23}) - a \bullet (s_{23}r_{13}) - \alpha^2(s_{(1)}) \otimes (s_{(2)} \cdot a) \cdot \alpha(r_{(1)}) \otimes \alpha^2(r_{(2)}) + (s_{23}s_{13}) \bullet a.
 \end{aligned}$$

It follows from (15) and (16) that

$$(17) \quad (\Delta' \otimes \alpha)(\Delta' a) - (\alpha \otimes \Delta')(\Delta' a) = a \bullet (r_{13}r_{12} - r_{12}r_{23} + s_{23}r_{13}) - (s_{13}r_{12} - s_{12}s_{23} + s_{23}s_{13}) \bullet a.$$

This shows that Δ' is Hom-coassociative if and only if (12) holds. This completes the proof. \square

It follows from the above theorem that a Hom-Yang-Baxter pair (r, s) on a Hom-associative algebra (A, α, μ) induces a covariant Hom-bialgebra $(A, \alpha, \mu, \Delta', \delta_r, \delta_s)$, called quasitriangular covariant bialgebra.

When $\alpha = \text{id}$, one recover the notion of quasitriangular covariant bialgebra introduced in [5]. On the other hand, if $r = s$ (i.e. r is a solution of Hom-Yang-Baxter equation), one get quasitriangular infinitesimal Hom-bialgebra introduced in [25].

The following theorem gives some characterizations of quasitriangular covariant Hom-bialgebras.

5.8. Theorem. *Let (A, α, μ) be a Hom-associative algebra and r, s be two α -invariant elements of $A^{\otimes 2}$. Let $\Delta', \delta_r, \delta_s$ be defined by (11). Then the followings are equivalent:*

- (i) $(A, \alpha, \mu, \Delta', \delta_r, \delta_s)$ is a quasitriangular covariant Hom-bialgebra;
- (ii) $(\alpha \otimes \Delta')(r) = r_{13}r_{12}$ and $(\Delta' \otimes \alpha)(s) = -s_{23}s_{13}$;
- (iii) The diagrams

$$(18) \quad \begin{array}{ccc} A^* \otimes A^* & \xrightarrow{\rho_1^r \otimes \rho_1^r} & A \otimes A \\ \Delta'^* \downarrow & & \downarrow \mu^{\text{op}} \\ A^* & \xrightarrow{\rho_2^r} & A \end{array} \qquad \begin{array}{ccc} A^* \otimes A^* & \xrightarrow{\lambda_1^s \otimes \lambda_1^s} & A \otimes A \\ \Delta'^* \downarrow & & \downarrow -\mu^{\text{op}} \\ A^* & \xrightarrow{\lambda_2^s} & A \end{array}$$

are commutative. Here $\rho_1^r, \rho_2^r, \lambda_1^s, \lambda_2^s : A^* \rightarrow A$ are maps defined respectively by

$$\begin{aligned}\rho_1^r(\phi) &= r_{(1)}\langle\phi, \alpha(r_{(2)})\rangle, & \rho_2^r(\phi) &= \alpha(r_{(1)})\langle\phi, r_{(2)}\rangle, \\ \lambda_1^s(\phi) &= \langle\phi, \alpha(s_{(1)})\rangle, & \lambda_2^s(\phi) &= \langle\phi, s_{(1)}\rangle\alpha(s_{(2)}).\end{aligned}$$

Proof. (i) \iff (ii). By definition, the tuple $(A, \alpha, \mu, \Delta', \delta_r, \delta_s)$ is a quasitriangular covariant Hom-bialgebra if and only if (r, s) is a Hom-Yang-Baxter pair. This holds if and only if

$$\begin{aligned}(\alpha \otimes \Delta')(r) &= \alpha(r_{(1)}) \otimes \Delta'(r_{(2)}) \\ &= \alpha(r_{(1)}) \otimes r_{(2)} \cdot \tilde{r}_{(1)} \otimes \alpha(\tilde{r}_{(2)}) - \alpha(r_{(1)}) \otimes \alpha(s_{(1)}) \otimes s_{(2)} \cdot r_{(2)} \\ &= r_{12}r_{23} - s_{23}r_{13} = r_{13}r_{12},\end{aligned}$$

and

$$\begin{aligned}(\Delta' \otimes \alpha)(s) &= \Delta'(s_{(1)}) \otimes \alpha(s_{(2)}) \\ &= s_{(1)} \cdot r_{(1)} \otimes \alpha(r_{(2)}) \otimes \alpha(s_{(2)}) - \alpha(\tilde{s}_{(1)}) \otimes \tilde{s}_{(2)} \cdot s_{(1)} \otimes \alpha(s_{(2)}) \\ &= s_{13}r_{12} - s_{12}s_{23} = -s_{23}s_{13}.\end{aligned}$$

(i) \iff (iii). Note that $r_{13}r_{12} - r_{12}r_{23} + s_{23}r_{13} = 0$ and $s_{13}r_{12} - s_{12}s_{23} + s_{23}s_{13} = 0$ if and only if

$$\langle \text{id}_A \otimes \phi \otimes \psi, r_{13}r_{12} - r_{12}r_{23} + s_{23}r_{13} \rangle = 0 \quad \text{and} \quad \langle \phi \otimes \psi \otimes \text{id}_A, s_{13}r_{12} - s_{12}s_{23} + s_{23}s_{13} \rangle = 0,$$

for all $\phi, \psi \in A^*$. First we have

$$\begin{aligned}r_{13}r_{12} - r_{12}r_{23} + s_{23}r_{13} &= r_{13}r_{12} - (\alpha \otimes \Delta')(r) \\ &= r_{(1)} \cdot \tilde{r}_{(1)} \otimes \alpha(\tilde{r}_{(2)}) \otimes \alpha(r_{(2)}) - \alpha(r_{(1)}) \otimes \Delta'(r_{(2)}).\end{aligned}$$

Hence

$$\begin{aligned}\langle \text{id}_A \otimes \phi \otimes \psi, r_{13}r_{12} - r_{12}r_{23} + s_{23}r_{13} \rangle &= r_{(1)} \cdot \tilde{r}_{(1)}\langle\phi, \alpha(\tilde{r}_{(2)})\rangle\langle\psi, \alpha(r_{(2)})\rangle - \alpha(r_{(1)})\langle\Delta'^*(\phi \otimes \psi), r_{(2)}\rangle \\ &= \mu^{\text{op}}((\rho_1^r)^{\otimes 2}(\phi \otimes \psi)) - \rho_2^r(\Delta'^*(\phi \otimes \psi)).\end{aligned}$$

Similarly,

$$s_{13}r_{12} - s_{12}s_{23} + s_{23}s_{13} = (\Delta' \otimes \alpha)(s) + s_{23}s_{13} = \Delta'(s_{(1)}) \otimes \alpha(s_{(2)}) + \alpha(s_{(1)}) \otimes \alpha(\tilde{s}_{(1)}) \otimes \tilde{s}_{(2)} \cdot s_{(2)}.$$

Hence

$$\begin{aligned}\langle \phi \otimes \psi \otimes \text{id}_A, s_{13}r_{12} - s_{12}s_{23} + s_{23}s_{13} \rangle &= \langle \Delta'^*(\phi \otimes \psi), s_{(1)}\rangle\alpha(s_{(2)}) + \langle \phi, \alpha(s_{(1)})\rangle\langle\psi, \alpha(\tilde{s}_{(1)})\rangle\tilde{s}_{(2)} \cdot s_{(2)} \\ &= \lambda_2^s(\Delta'^*(\phi \otimes \psi)) + \mu^{\text{op}}((\lambda_1^s)^{\otimes 2}(\phi \otimes \mu)).\end{aligned}$$

Hence the result follows. \square

Let $(A, \alpha, \mu, \Delta, \delta_1, \delta_2)$ be a covariant Hom-bialgebra. Next we consider perturbations of the coproduct Δ and derivations δ_1, δ_2 by a suitable pair (r, s) of α -invariant elements of $A^{\otimes 2}$. These perturbations generalize the perturbation theory of quasi-Hopf algebras studied by Drinfel'd [12,13]. The following result also generalizes the result of Yau for infinitesimal Hom-bialgebras [25].

5.9. Theorem. *Let $(A, \alpha, \mu, \Delta, \delta_1, \delta_2)$ be a covariant Hom-bialgebra and (r, s) be a pair of α -invariant elements of $A^{\otimes 2}$. Then $(A, \alpha, \mu, \Delta + \Delta', \delta_1 + \delta_r, \delta_2 + \delta_s)$ is also a covariant Hom-bialgebra if and only if*

$$(19) \quad a \bullet ((\alpha \otimes \Delta)^-(r) - (r_{13}r_{12} - r_{12}r_{23} + s_{23}r_{13})) - ((\alpha \otimes \Delta)^-(s) - (s_{13}r_{12} - s_{12}s_{23} + s_{23}s_{13})) \bullet a \\ = s_{23}\Delta(a)_{13} + \Delta(a)_{13}r_{12}.$$

Proof. First note that each $\Delta', \delta_r, \delta_s$ commutes with α . Therefore, the sums $\Delta + \Delta', \delta_1 + \delta_r, \delta_2 + \delta_s$ also commute with α . Likewise, δ_r and δ_s are derivations, so the sums $\delta_1 + \delta_r$ and $\delta_2 + \delta_s$. We claim that $\Delta + \Delta'$

is a covariant derivation with respect to $(\delta_1 + \delta_r, \delta_2 + \delta_s)$. We have

$$\begin{aligned} (\Delta + \Delta') \circ \mu &= \Delta \circ \mu + \Delta' \circ \mu \\ &= (\mu \otimes \alpha) \circ (\alpha \otimes \Delta) + (\alpha \otimes \mu) \circ (\delta_2 \otimes \alpha) + (\mu \otimes \alpha) \circ (\alpha \otimes \Delta') + (\alpha \otimes \mu) \circ (\delta_s \otimes \alpha) \\ &= (\mu \otimes \alpha) \circ (\alpha \otimes (\Delta + \Delta')) + (\alpha \otimes \mu) \circ ((\delta_2 + \delta_s) \otimes \alpha). \end{aligned}$$

Similarly, we can show that $(\Delta + \Delta') \circ \mu = (\mu \otimes \alpha) \circ (\alpha \otimes (\delta_1 + \delta_r)) + (\alpha \otimes \mu) \circ ((\Delta + \Delta') \otimes \alpha)$. Hence we proved our claim. Thus, we are left with the Hom-coassociativity of $\Delta + \Delta'$. We show that $\Delta + \Delta'$ is Hom-coassociative if and only if (19) holds. First observe that

$$(20) \quad (\alpha \otimes (\Delta + \Delta')) \circ (\Delta + \Delta') = (\alpha \otimes \Delta) \circ \Delta + (\alpha \otimes \Delta) \circ \Delta' + (\alpha \otimes \Delta') \circ \Delta + (\alpha \otimes \Delta') \circ \Delta'.$$

Similarly,

$$(21) \quad (\Delta + \Delta') \otimes \alpha \circ (\Delta + \Delta') = (\Delta \otimes \alpha) \circ \Delta + (\Delta \otimes \alpha) \circ \Delta' + (\Delta' \otimes \alpha) \circ \Delta + (\Delta' \otimes \alpha) \circ \Delta'.$$

Since Δ is Hom-coassociative, we have $(\alpha \otimes \Delta) \circ \Delta = (\Delta \otimes \alpha) \circ \Delta$. Therefore, it follows from (20) and (21) that $(\Delta + \Delta')$ is Hom-coassociative if and only if

$$(\alpha \otimes \Delta - \Delta \otimes \alpha) \circ \Delta' + (\alpha \otimes \Delta' - \Delta' \otimes \alpha) \circ \Delta = (\Delta' \otimes \alpha - \alpha \otimes \Delta') \circ \Delta',$$

or equivalently,

$$\begin{aligned} &(\alpha \otimes \Delta)^- \circ \Delta'(a) + (\alpha \otimes \Delta')^- \circ \Delta(a) \\ &= a \bullet (r_{13}r_{12} - r_{12}r_{23} + s_{23}r_{13}) - (s_{13}r_{12} - s_{12}s_{23} + s_{23}s_{13}) \bullet a \quad (\text{by (17)}). \end{aligned}$$

For $r = r_{(1)} \otimes r_{(2)}$ and $s = s_{(1)} \otimes s_{(2)}$, we have

$$\begin{aligned} &(\alpha \otimes \Delta)^- \circ \Delta'(a) \\ &= (\alpha \otimes \Delta - \Delta \otimes \alpha)(a \cdot r_{(1)} \otimes \alpha(r_{(2)}) - \alpha(s_{(1)}) \otimes s_{(2)} \cdot a) \\ &= \alpha(a) \cdot \alpha(r_{(1)}) \otimes \Delta(\alpha(r_{(2)})) - \alpha^2(s_{(1)}) \otimes \Delta(s_{(2)} \cdot a) - \Delta(a \cdot r_{(1)}) \otimes \alpha^2(r_{(2)}) + \Delta(\alpha(s_{(1)})) \otimes \alpha(s_{(2)}) \cdot \alpha(a) \\ &= \alpha(a) \cdot \alpha(r_{(1)}) \otimes \alpha^{\otimes 2}(\Delta(r_{(2)})) - \alpha^2(s_{(1)}) \otimes s_2 \bullet \Delta(a) - \alpha^2(s_{(1)}) \otimes \Delta(s_{(2)}) \bullet a \\ &\quad - a \bullet \Delta(r_{(1)}) \otimes \alpha^2(r_{(2)}) - \Delta(a) \bullet r_{(1)} \otimes \alpha^2(r_{(2)}) + \alpha^{\otimes 2}(\Delta(s_{(1)})) \otimes \alpha(s_{(2)}) \cdot \alpha(a) \\ &= a \bullet ((\alpha \otimes \Delta)(r)) - s_{12}\Delta(a)_{23} - ((\alpha \otimes \Delta)(s)) \bullet a - a \bullet ((\Delta \otimes \alpha)(r)) - \Delta(a)_{12}r_{23} + ((\Delta \otimes \alpha)(s)) \bullet a \\ &= a \bullet (\alpha \otimes \Delta)^-(r) - (\alpha \otimes \Delta)^-(s) \bullet a - s_{12}\Delta(a)_{23} - \Delta(a)_{12}r_{23}. \end{aligned}$$

On the other hand, by writing $\Delta(a) = a_{(1)} \otimes a_{(2)}$, we get

$$\begin{aligned} &(\alpha \otimes \Delta')^- \circ \Delta(a) \\ &= (\alpha \otimes \Delta' - \Delta' \otimes \alpha)(a_{(1)} \otimes a_{(2)}) \\ &= \alpha(a_{(1)}) \otimes a_{(2)} \cdot r_{(1)} \otimes \alpha(r_{(2)}) - \alpha(a_{(1)}) \otimes \alpha(s_{(1)}) \otimes s_{(2)} \cdot a_{(2)} \\ &\quad - a_{(1)} \cdot r_{(1)} \otimes \alpha(r_{(2)}) \otimes \alpha(a_{(2)}) + \alpha(s_{(1)}) \otimes s_{(2)} \cdot a_{(1)} \otimes \alpha(a_{(2)}) \\ &= \Delta(a)_{12}r_{23} - s_{23}\Delta(a)_{13} - \Delta(a)_{13}r_{12} + s_{12}\Delta(a)_{23}. \end{aligned}$$

By adding the above two identities, we get

$$\begin{aligned} &(\alpha \otimes \Delta)^- \Delta'(a) + (\alpha \otimes \Delta')^- \circ \Delta(a) \\ &= a \bullet ((\alpha \otimes \Delta)^-(r)) - ((\alpha \otimes \Delta)^-(s)) \bullet a - s_{23}\Delta(a)_{13} - \Delta(a)_{13}r_{12}. \end{aligned}$$

This is same as $a \bullet (r_{13}r_{12} - r_{12}r_{23} + s_{23}r_{13}) - (s_{13}r_{12} - s_{12}s_{23} + s_{23}s_{13}) \bullet a$ (equivalently, $(\Delta + \Delta')$ is Hom-coassociative) if and only if (19) holds. Hence the proof. \square

The above theorem has some nice consequences.

(i) (Taking $\alpha = \text{id}$) Given a covariant bialgebra $(A, \mu, \Delta, \delta_1, \delta_2)$ and a pair (r, s) of elements of $A^{\otimes 2}$, the theorem gives a necessary and sufficient condition for $(A, \mu, \Delta + \Delta', \delta_1 + \delta_r, \delta_2 + \delta_s)$ to be a covariant bialgebra.

(ii) (Taking $\Delta = \delta_1 = \delta_2 = 0$) Given a Hom-associative algebra (A, α, μ) and a pair (r, s) of α -invariant elements of $A^{\otimes 2}$, the tuple $(A, \alpha, \mu, \Delta', \delta_r, \delta_s)$ is a covariant Hom-bialgebra if and only if

$$a \bullet (r_{13}r_{12} - r_{12}r_{23} + s_{23}r_{13}) = (s_{13}r_{12} - s_{12}s_{23} + s_{23}s_{13}) \bullet a.$$

Therefore, in this case, we recover Theorem 5.7.

Acknowledgements. The research is supported by the fellowship of Indian Institute of Technology (IIT) Kanpur. The author thanks the Institute for support.

REFERENCES

- [1] F. Ammar, Z. Ejbehi and A. Makhlouf, Cohomology and Deformations of Hom-algebras, *J. Lie Theory* 21 (2011), no. 4, 813-836.
- [2] M. Aguiar, Infinitesimal Hopf algebras, Contemporary Mathematics, New trends in Hopf algebra theory (La Falda, 1999), 1-29, *Contemp. Math.*, 267, Amer. Math. Soc., Providence, RI, (2000).
- [3] M. Aguiar, Pre-Poisson algebras, *Lett. Math. Phys.* 54 (2000), 263-277.
- [4] G. Baxter, An analytic problem whose solution follows from a simple algebraic identity, *Pacific J. Math.* 10 (1960), 731-742.
- [5] T. Brzeziński, Rota-Baxter systems, dendriform algebras and covariant bialgebras, *J. Algebra* 460 (2016) 1-25.
- [6] P. Cartier, On the structure of free Baxter algebras, *Advances in Math.* 9 (1972), 253-265.
- [7] A. Connes and D. Kreimer, Renormalization in quantum field theory and the Riemann-Hilbert problem. I. The Hopf algebra structure of graphs and the main theorem, *Comm. Math. Phys.* 210 (2000), no. 1, 249-273.
- [8] A. Das, Homotopy G -algebra structure on the cochain complex of hom-type algebras, *C. R. Math. Acad. Sci. Paris* 356 (2018), no. 11, 1090-1099.
- [9] A. Das, Hom-associative algebras up to homotopy, *J. Algebra* 556 (2020) 836-878.
- [10] A. Das, Deformations of associative Rota-Baxter operators, *J. Algebra* 560 (2020) 144-180.
- [11] A. Das, Cohomology and deformations of hom-dendriform algebras and coalgebras, *Colloq. Math.* DOI: 10.4064/cm8049-10-2019.
- [12] V. G. Drinfel'd, On quasitriangular quasi-Hopf algebras and a certain group closely connected with $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$, *Leningrad Math. J.* 2 (1991) 829-860.
- [13] V. G. Drinfel'd, On the structure of quasitriangular quasi-Hopf algebras, *Func. Anal. Appl.* 26 (1992) 63-65.
- [14] K. Ebrahimi-Fard and L. Guo, Free Rota-Baxter algebras and rooted trees, *J. Alg. Appl.* Vol. 07, No. 02, pp. 167-194 (2008).
- [15] L. Guo, An introduction to Rota-Baxter algebra, Surveys of Modern Mathematics, 4. *International Press, Somerville, MA; Higher Education Press, Beijing*, 2012.
- [16] L. Guo and W. Keigher, Baxter algebras and shuffle products, *Adv. Math.* 150 (2000), 117-149.
- [17] J. T. Hartwig, D. Larsson and S. D. Silvestrov, Deformations of Lie algebras using σ -derivations, *J. Algebra* 295 (2006), 314-361.
- [18] S. A. Joni and G.-C. Rota, Coalgebras and bialgebras in combinatorics, *Stud. Appl. Math.* 61 (1979), no. 2, 93-139.
- [19] L. Liu, A. Makhlouf, C. Menini and F. Panaite, $\{\sigma, \tau\}$ -Rota-Baxter operators, infinitesimal Hom-bialgebras and the associative (Bi)Hom-Yang-Baxter equation, *Canad. Math. Bull.* Vol. 62 (2), 2019 pp. 355-372.
- [20] L. Liu, A. Makhlouf, C. Menini and F. Panaite, Rota-Baxter operators on BiHom-associative algebras and related structures, *Colloq. Math.* 161 (2020), 263-294.
- [21] J.-L. Loday, Dialgebras and related operads, 7-66, Lecture Notes in Math., 1763, *Springer, Berlin*, 2001.
- [22] A. Makhlouf, Hom-dendriform algebras and Rota-Baxter Hom-algebras, *Operads and universal algebra*, 147-171, Nankai Ser. Pure Appl. Math. Theoret. Phys., 9, *World Sci. Publ., Hackensack, NJ*, 2012.
- [23] A. Makhlouf and S. Silvestrov, Hom-algebras and Hom-coalgebras, *J. Alg. Appl.* Vol. 09, No. 04, pp. 553-589 (2010).
- [24] G.-C. Rota, Baxter algebras and combinatorial identities, I, II, *Bull. Amer. Math. Soc.* 75 (1969), 325-329; *ibid.* 75 1969 330-334.
- [25] D. Yau, Infinitesimal Hom-bialgebras and Hom-Lie bialgebras, arXiv:1001.5000 [math.RA]

DEPARTMENT OF MATHEMATICS AND STATISTICS, INDIAN INSTITUTE OF TECHNOLOGY, KANPUR 208016, UTTAR PRADESH, INDIA.

E-mail address: apurbadas348@gmail.com