

A complete characterisation of vertex-multiplications of trees with diameter 5

W.H.W. Wong*, E.G. Tay

National Institute of Education
Nanyang Technological University, Singapore

Abstract

Koh and Tay [4] introduced a new family of graphs, G vertex-multiplications, as an extension of complete n -partite graphs. They proved a fundamental classification of G vertex-multiplications into three classes $\mathcal{C}_0, \mathcal{C}_1$ and \mathcal{C}_2 . It was shown in [5] that any vertex-multiplication of a tree with diameter at least 3 does not belong to the class \mathcal{C}_2 . Furthermore, for vertex-multiplications of trees with diameter 5, some necessary and sufficient conditions for \mathcal{C}_0 were established. In this paper, we give a complete characterisation of vertex-multiplications of trees with diameter 5 in \mathcal{C}_0 and \mathcal{C}_1 .

1. Introduction

Let G be a graph with vertex set $V(G)$ and edge set $E(G)$. In this paper, we consider graphs G with no loops nor parallel edges, unless otherwise stated. For any vertices $v, x \in V(G)$, the *distance* from v to x , $d_G(v, x)$, is defined as the length of a shortest path from v to x . For $v \in V(G)$, its *eccentricity* $e_G(v)$ is defined as $e_G(v) := \max\{d_G(v, x) \mid x \in V(G)\}$. The *diameter* of G , denoted by $d(G)$, is defined as $d(G) := \max\{e_G(v) \mid v \in V(G)\}$ while the *radius* of G , denoted by $r(G)$, is defined as $r(G) := \min\{e_G(v) \mid v \in V(G)\}$. The above notions are defined similarly for a digraph D . A vertex x is said to be *reachable* from another vertex v if $d_D(v, x) < \infty$. For a digraph D , the *outset* and *inset* of a vertex $v \in V(D)$ are defined to be $O_D(v) := \{x \in V(D) \mid v \rightarrow x\}$ and $I_D(v) := \{y \in V(D) \mid y \rightarrow v\}$ respectively. If there is no ambiguity, we shall omit the subscript for the above notations.

An *orientation* D of a graph G is a digraph obtained from G by assigning a direction to every edge $e \in E(G)$. An orientation D of G is said to be *strong* if every two vertices in $V(D)$ are mutually reachable. An edge $e \in E(G)$ is a *bridge* if $G - e$ is disconnected. Robbins' well-known One-way Street Theorem [9] states the following.

Theorem 1.1 (*Robbins [9]*)

Let G be a connected graph. Then, G has a strong orientation if and only if G is bridgeless.

Given a connected and bridgeless graph G , let $\mathcal{D}(G)$ be the family of strong orientations of G . The *orientation number* of G is defined as

$$\bar{d}(G) := \min\{d(D) \mid D \in \mathcal{D}(G)\}.$$

The general problem of finding the orientation number of a connected and bridgeless graph is very difficult. Moreover, Chvátal and Thomassen [2] proved that it is NP-hard to determine if a graph admits an orientation of diameter 2. Hence, it is natural to focus

*Corresponding author. Email: williewong088@gmail.com.

on special classes of graphs. The orientation number was evaluated for various classes of graphs, such as the complete graphs [1, 6, 8] and complete bipartite graphs [3, 10].

In 2000, Koh and Tay [4] introduced a new family of graphs, G vertex-multiplications, and extended the results on complete n -partite graphs. Let G be a given connected graph with vertex set $V(G) = \{v_1, v_2, \dots, v_n\}$. For any sequence of n positive integers (s_i) , a G vertex-multiplication, denoted by $G(s_1, s_2, \dots, s_n)$, is the graph with vertex set $V^* = \bigcup_{i=1}^n V_i$ and edge set E^* , where V_i 's are pairwise disjoint sets with $|V_i| = s_i$, for $i = 1, 2, \dots, n$, and for any $u, v \in V^*$, $uv \in E^*$ if and only if $u \in V_i$ and $v \in V_j$ for some $i, j \in \{1, 2, \dots, n\}$ with $i \neq j$ such that $v_i v_j \in E(G)$. For instance, if $G \cong K_n$, then the graph $G(s_1, s_2, \dots, s_n)$ is a complete n -partite graph with partite sizes s_1, s_2, \dots, s_n . Also, we say G is a parent graph of $G(s_1, s_2, \dots, s_n)$.

For $i = 1, 2, \dots, n$, we denote the x -th vertex in V_i by (x, v_i) , i.e. $V_i = \{(x, v_i) \mid x = 1, 2, \dots, s_i\}$. Hence, two vertices (x, v_i) and (y, v_j) in V^* are adjacent in $G(s_1, s_2, \dots, s_n)$ if and only if $i \neq j$ and $v_i v_j \in E(G)$. For convenience, we write $G^{(s)}$ in place of $G(s, s, \dots, s)$ for any positive integer s , and it is understood that the number of s 's is equal to the order of G , n . Thus, $G^{(1)}$ is simply the graph G itself.

The following theorem by Koh and Tay [4] provides a fundamental classification on G vertex-multiplications.

Theorem 1.2 (Koh and Tay [4])

Let G be a connected graph of order $n \geq 3$. If $s_i \geq 2$ for $i = 1, 2, \dots, n$, then $d(G) \leq \bar{d}(G(s_1, s_2, \dots, s_n)) \leq d(G) + 2$.

In view of Theorem 1.2, all graphs of the form $G(s_1, s_2, \dots, s_n)$, with $s_i \geq 2$ for all $i = 1, 2, \dots, n$, can be classified into three classes \mathcal{C}_j , where

$$\mathcal{C}_j = \{G(s_1, s_2, \dots, s_n) \mid \bar{d}(G(s_1, s_2, \dots, s_n)) = d(G) + j\},$$

for $j = 0, 1, 2$. Henceforth, we assume $s_i \geq 2$ for $i = 1, 2, \dots, n$, unless otherwise stated. The following lemma was found useful in proving Theorem 1.2.

Lemma 1.3 (Koh and Tay [4])

Let μ_i, λ_i be integers such that $\mu_i \leq \lambda_i$ for $i = 1, 2, \dots, n$. If the graph $G(\mu_1, \mu_2, \dots, \mu_n)$ admits an orientation F in which every vertex v lies on a cycle of length not exceeding m , then $\bar{d}(G(\lambda_1, \lambda_2, \dots, \lambda_n)) \leq \max\{m, d(F)\}$.

Koh and Tay [5] further investigated vertex-multiplications of trees. Since trees with diameter at most 2 are parent graphs of complete bipartite graphs, which are completely solved, they considered trees of diameter at least 3. On the other hand, Ng and Koh [7] examined vertex-multiplications of cycles. To continue the discussion, we need some notations.

From here onwards, we consider a tree T of diameter 5 with vertex set $V(T) = \{v_1, v_2, \dots, v_n\}$. We let v_1 and v_2 be the two central vertices of T , i.e. $e_T(v_k) = \text{rad}(T) = 3$ for $k = 1, 2$. For $k = 1, 2$, denote the neighbours of v_k , excluding v_{3-k} , by $[i]_k$. i.e. $N_T(v_k) - \{v_{3-k}\} = \{[i]_k \mid i = 1, 2, \dots, \text{deg}_T(v_k) - 1\}$. For each $i = 1, 2, \dots, \text{deg}_T(v_k) - 1$, we denote the neighbours of $[i]_k$, excluding v_k itself, by $[\alpha, i]_k$. i.e. $N_T([i]_k) - \{v_k\} = \{[\alpha, i]_k \mid \alpha = 1, 2, \dots, \text{deg}_T([i]_k) - 1\}$. Also, we denote the set of **Non-Leaf** neighbours of v_k to be $NL_k := \{[i]_k \in V(T) \mid 1 \leq i \leq \text{deg}_T(v_k) - 1 \text{ and } [i]_k \text{ is not an end-vertex}\}$ for $k = 1, 2$. Of course, $|NL_k| \geq 1$, where $k = 1, 2$, for any tree T with $d(T) = 5$.

Example 1.4 Let T be the tree of diameter 5 shown in Figure 1. We label the vertices as described above. Note also $NL_1 = \{[1]_1, [2]_1\}$ and $NL_2 = \{[2]_2\}$.




Figure 1: Labelling vertices in T

In the following theorem, Koh and Tay [5] proved some necessary and sufficient conditions for $T(s_1, s_2, \dots, s_n) \in \mathcal{C}_0$, where T is a tree of diameter 5.

Theorem 1.5 (Koh and Tay [5])

Let T be a tree with diameter 5 and its central vertices be v_1 and v_2 . Suppose $A = \{x \in V(T) \mid d_T(x, u) = 5 = d_T(x, v) \text{ for some } u, v \in V(T), u \neq v\}$. Then,

- (a) $T(s_1, s_2, \dots, s_n) \in \mathcal{C}_0 \cup \mathcal{C}_1$.
- (b) If $|A| \leq 1$, then $T(s_1, s_2, \dots, s_n) \in \mathcal{C}_0$.
- (c) If $\deg_T(v) \leq 2$ for all $v \notin \{v_1, v_2\}$ and $|A| \geq 2$, then $T^{(2)} \in \mathcal{C}_1$.

In this paper, we give a complete characterisation of vertex-multiplications of trees with diameter 5 in \mathcal{C}_0 and \mathcal{C}_1 (see Theorem 2.6).

2. Main results

For convenience, we shall introduce some notations. Let D be an orientation of $T(s_1, s_2, \dots, s_n)$ with $s_i \geq 2$ for $1 \leq i \leq n$. If v_p and v_q , $1 \leq p, q \leq n$ and $p \neq q$, are adjacent vertices in T , then for each i , $1 \leq i \leq s_p$, we denote by $O_D^{v_q}((i, v_p)) := \{(j, v_q) \mid (i, v_p) \rightarrow (j, v_q), 1 \leq j \leq s_q\}$ and $I_D^{v_q}((i, v_p)) := \{(j, v_q) \mid (j, v_q) \rightarrow (i, v_p), 1 \leq j \leq s_q\}$. If there is no ambiguity, we shall omit the subscript D for the above notations.

In the vertex-multiplication graph $G := T(s_1, s_2, \dots, s_n)$ of T , the integer s_i corresponds to the vertex v_i , $i = 1, 2, \dots, n$. We will loosely use the two denotations of a vertex, for example, if $v_i = [j]_k$, then $s_i = s_{[j]_k}$. For any $v \in V(G)$, we set $(\mathbb{N}_s, v) := \{(1, v), (2, v), \dots, (s, v)\}$.

Proposition 2.1 Let T be a tree of diameter 5. If $s_1 \geq 3$ or $s_2 \geq 3$, then $G := T(s_1, s_2, \dots, s_n) \in \mathcal{C}_0$.

Proof: WLOG, assume $s_1 \geq 3$. Let H be a subgraph of G , where $s_1 = 3$ and $s_i = 2$ for all $i \neq 1$ in H . Define an orientation D of H as follows (see Figure 2).

$$\begin{aligned} (2, [i]_1) &\rightarrow \{(1, [\alpha, i]_1), (2, [\alpha, i]_1)\} \rightarrow (1, [i]_1), \\ (2, v_1) &\rightarrow (1, [i]_1) \rightarrow \{(1, v_1), (3, v_1)\}, \text{ and} \\ \{(1, v_1), (2, v_1)\} &\rightarrow (2, [i]_1) \rightarrow (3, v_1), \end{aligned}$$

for all $i = 1, 2, \dots, \deg_T(v_1) - 1$ and all $\alpha = 1, 2, \dots, \deg_T([i]_1) - 1$.

$$\begin{aligned} (2, [j]_2) &\rightarrow \{(1, [\beta, j]_2), (2, [\beta, j]_2)\} \rightarrow (1, [j]_2), \text{ and} \\ (2, [j]_2) &\rightarrow (1, v_2) \rightarrow (1, [j]_2) \rightarrow (2, v_2) \rightarrow (2, [j]_2), \end{aligned}$$

for all $j = 1, 2, \dots, \deg_T(v_2) - 1$ and all $\beta = 1, 2, \dots, \deg_T([j]_2) - 1$. Also,

$$\begin{aligned} \{(1, v_1), (3, v_1)\} &\rightarrow (1, v_2) \rightarrow (2, v_1), \text{ and} \\ (3, v_1) &\rightarrow (2, v_2) \rightarrow \{(1, v_1), (2, v_1)\}. \end{aligned}$$

We emphasize that the above orientation applies to the vertices $[i]_1$ and $[j]_2$ even if they are end-vertices. It suffices to verify $d(D) = 5$ and every vertex in $V(D)$ lies in a directed C_4 , where D is as shown in Figure 2. Then, by Lemma 1.3, $\bar{d}(G) \leq \max\{d(D), 4\} = 5$.

□




Figure 2: Orientation D for H .

The next two lemmas will be found useful in proving our propositions to come.

Lemma 2.2 *Let D be an orientation of $T(s_1, s_2, \dots, s_n)$ with $s_1 = s_2 = 2$. If $d(D) = 5$, then*

- (i) $|O^{v_1}((p, [i]_1))| \geq 1$ and $|I^{v_1}((p, [i]_1))| \geq 1$ for all $i = 1, 2, \dots, \deg_T(v_1) - 1$ and $p = 1, 2, \dots, s_{[i]_1}$, and
(ii) $|O^{v_2}((q, [j]_2))| \geq 1$ and $|I^{v_2}((q, [j]_2))| \geq 1$ for all $j = 1, 2, \dots, \deg_T(v_2) - 1$ and $q = 1, 2, \dots, s_{[j]_2}$.

Proof: (i) follows from the fact that $d_D((p, [i]_1), (1, [1, j]_2)) = d_D((1, [1, j]_2), (p, [1]_1)) = 4$, where $[j]_2$ is not an end-vertex in T . By symmetry, (ii) follows from (i). □

Lemma 2.3 (Duality)

Let D be an orientation of a graph G . Let \tilde{D} be the orientation of G such that $uv \in E(\tilde{D})$ if and only if $vu \in E(D)$. Then, $d(\tilde{D}) = d(D)$.

Proof: Suppose not. Then, there exist some vertices $u, v \in V(\tilde{D})$ such that $d_{\tilde{D}}(u, v) > d(D)$. By definition of \tilde{D} , $d_D(v, u) = d_{\tilde{D}}(u, v)$. It follows that $d_D(v, u) > d(D)$, a contradiction. □

Proposition 2.4 *Let T be a tree of diameter 5 and $m_k = \min\{s_{[i]_k} \mid [i]_k \in NL_k\}$ for $k = 1, 2$. Suppose $G := T(s_1, s_2, \dots, s_n)$ satisfy $2 \leq m_1 \leq 3$ or $2 \leq m_2 \leq 3$, and $s_1 = s_2 = 2$. Then,*

$$G \in \mathcal{C}_0 \iff |NL_k| = 1 \text{ for some } k = 1, 2.$$

Proof: (\Rightarrow) Suppose $|NL_k| \geq 2$ for all $k = 1, 2$. WLOG, assume $2 \leq m_1 \leq 3$, and $m_k = s_{[1]_k}$ where $[1]_k \in NL_k$ for $k = 1, 2$.

Case 1. $|I^{v_2}((p, v_1))| = 2$ for some $p = 1, 2$.

WLOG, assume $\{(1, v_2), (2, v_2)\} \rightarrow (2, v_1)$. Since for all $i = 1, 2, \dots, \deg_T(v_1) - 1$, and $q = 1, 2, \dots, s_{[i]_1}$, $d_D((q, [i]_1), (1, [1, 1]_2)) \leq 5$ and by Lemma 2.2, we have $(2, v_1) \rightarrow (q, [i]_1) \rightarrow (1, v_1)$. Then, $d_D((1, [1, 1]_1), (1, [1, i]_1)) > 5$ for any $[i]_1 \in NL_1$, $i \neq 1$, a contradiction.

Case 2. $|O^{v_2}((p, v_1))| = 2$ for some $p = 1, 2$.

By Duality Lemma and Case 1, this case follows.

Case 3. $|I^{v_1}((p, v_2))| = |I^{v_2}((p, v_1))| = 1$ for all $p = 1, 2$.

WLOG, assume $(1, v_1) \rightarrow (1, v_2) \rightarrow (2, v_1) \rightarrow (2, v_2) \rightarrow (1, v_1)$.

Subcase 3.1. $|O((1, [1, 1]_1))| = 1$.

Assume WLOG that $O((1, [1, 1]_1)) = \{(1, [1]_1)\}$. By Lemma 2.2, $|O^{v_1}((1, [1]_1))| = 1$. By symmetry, we may assume $O^{v_1}((1, [1]_1)) = \{(1, v_1)\}$. Since for all $i = 1, 2, \dots, \deg_T(v_2) - 1$, and $q = 1, 2, \dots, s_{[i]_2}$, $d_D((1, [1, 1]_1), (q, [i]_2)) \leq 5$ and by Lemma 2.2, we have $(1, v_2) \rightarrow (q, [i]_2) \rightarrow (2, v_2)$. However, $d_D((1, [1, 1]_2), (1, [1, j]_2)) > 5$ for any $[j]_2 \in NL_2$, $j \neq 1$, a contradiction.

Subcase 3.2. $|I((1, [1, 1]_1))| = 1$.

By Duality Lemma and Subcase 3.1, this subcase also results in a contradiction.

(\Leftarrow) Assume WLOG that $|NL_1| = 1$. Let H be a subgraph of G , where $s_i = 2$ for all i in H . Define an orientation D of H as follows (see Figure 3).

$$(1, [\alpha, 1]_1) \rightarrow (1, [1]_1) \rightarrow (2, [\alpha, 1]_1) \rightarrow (2, [1]_1) \rightarrow (1, [\alpha, 1]_1), \text{ and}$$

$$\{(1, [i]_1), (2, [i]_1)\} \rightarrow (1, v_1) \rightarrow \{(1, v_2), (2, v_2)\} \rightarrow (2, v_1) \rightarrow \{(1, [i]_1), (2, [i]_1)\},$$

for all $i = 1, 2, \dots, \deg_T(v_1) - 1$ and $\alpha = 1, 2, \dots, \deg_T([1]_1) - 1$.

$$(2, [j]_2) \rightarrow \{(1, [\beta, j]_2), (2, [\beta, j]_2)\} \rightarrow (1, [j]_2), \text{ and}$$

$$(1, [j]_2) \rightarrow (1, v_2) \rightarrow (2, [j]_2) \rightarrow (2, v_2) \rightarrow (1, [j]_2),$$

for all $j = 1, 2, \dots, \deg_T(v_2) - 1$ and $\beta = 1, 2, \dots, \deg_T([j]_2) - 1$.

We emphasize that the above orientation applies to the vertices $[i]_1$ and $[j]_2$ even if they are end-vertices. It suffices to verify $d(D) = 5$ and every vertex in $V(D)$ lies in a directed C_4 , where D is as shown in Figure 3. Then, by Lemma 1.3, $\bar{d}(G) \leq \max\{d(D), 4\} = 5$.

□




Figure 3: Orientation D for H .

Proposition 2.5 *Let T be a tree of diameter 5 and $m_k = \min\{s_{[i]_k} \mid [i]_k \in NL_k\}$ for $k = 1, 2$. If $m_1 \geq 4$ and $m_2 \geq 4$, then $G := T(s_1, s_2, \dots, s_n) \in \mathcal{C}_0$.*

Proof: Let H be a subgraph of G , where $s_{[i]_k} = 4$ for all $[i]_k \in NL_k$, $k = 1, 2$, and $s_j = 2$ otherwise. Define an orientation D of H as follows (see Figure 4). For $k = 1, 2$,

$$\begin{aligned} & \{(3, [i]_k), (4, [i]_k)\} \rightarrow \{(1, [\alpha, i]_k), (2, [\alpha, i]_k)\} \rightarrow \{(1, [i]_k), (2, [i]_k)\}, \\ & (2, v_k) \rightarrow \{(1, [i]_k), (3, [i]_k)\} \rightarrow (1, v_k) \rightarrow \{(2, [i]_k), (4, [i]_k)\} \rightarrow (2, v_k), \text{ and} \\ & (1, v_1) \rightarrow (1, v_2) \rightarrow (2, v_1) \rightarrow (2, v_2) \rightarrow (1, v_1), \end{aligned}$$

for all $i = 1, 2, \dots, \deg_T(v_k) - 1$ and $\alpha = 1, 2, \dots, \deg_T([i]_k) - 1$.

We emphasize that the above orientation applies to the vertices $[i]_1$ and $[j]_2$ even if they are end-vertices. Note that for any $p, q, r, k = 1, 2$, $1 \leq i, j \leq \deg_T(v_k)$, $i \neq j$, $\alpha = 1, 2, \dots, \deg_T([i]_k) - 1$, and $\beta = 1, 2, \dots, \deg_T([j]_k) - 1$, we have $d_D((p, [\alpha, i]_k), (q, v_k)) = 2$ and $d_D((q, v_k), (r, [\beta, j]_k)) = 2$ so that $d_D((p, [\alpha, i]_k), (r, [\beta, j]_k)) \leq d_D((p, [\alpha, i]_k), (q, v_k)) + d_D((q, v_k), (r, [\beta, j]_k)) = 4$. Furthermore, in view of the orientation's similarity, it suffices to verify $d(D) = 5$ and every vertex in $V(D)$ lies in a directed C_4 , where D is as shown in Figure 4. Hence, by Lemma 1.3, $\bar{d}(G) \leq \max\{d(D), 4\} = 5$. □




Figure 4: Orientation D for H .

The findings in this section provide a complete characterisation of vertex-multiplications of trees of diameter 5 in \mathcal{C}_0 and \mathcal{C}_1 . We summarise them in Theorem 2.6.

Theorem 2.6 Let $G := T(s_1, s_2, \dots, s_n)$ with $d(T) = 5$, and $m_k = \min\{s_{[i]_k} \mid i \in NL_k\}$ for $k = 1, 2$.

Underlying conditions	$G \in \mathcal{C}_0$ or \mathcal{C}_1 ?	Reference
$s_i \geq 3$ for some $i = 1, 2$	$G \in \mathcal{C}_0$.	Proposition 2.1
$m_1 \geq 4$ and $m_2 \geq 4$	$G \in \mathcal{C}_0$.	Proposition 2.5
$s_1 = s_2 = 2$, and $2 \leq m_i \leq 3$ for some $i = 1, 2$	$G \in \mathcal{C}_0 \iff NL_k = 1$ for some $k = 1, 2$.	Proposition 2.4

Table 1: Summary for $T(s_1, s_2, \dots, s_n)$ with $d(T) = 5$.

Koh and Tay [5] also showed that vertex-multiplications of trees with diameter at least 6 belong to \mathcal{C}_0 . Hence, it remains open to characterise vertex-multiplications of trees, with diameter 3 and 4, belonging to \mathcal{C}_0 . We conclude the paper by proposing the following problem.

Problem 2.7 For trees T with $d(T) = 3$ (and $d(T) = 4$ resp.), characterise the tree vertex-multiplications $T(s_1, s_2, \dots, s_n)$ that belong to \mathcal{C}_0 .

Acknowledgement

The first author would like to thank the National Institute of Education, Nanyang Technological University of Singapore, for the generous support of the Nanyang Technological University Research Scholarship.

References

- [1] Boesch F., Tindell R., *Robbins's Theorem for Mixed Multigraphs*, The American Mathematical Monthly, 87, (1980), 716-719.
- [2] Chvátal V., Thomassen C., *Distances in orientations of graphs*, Journal of Combinatorial Theory B, 24, (1978), 61-75.
- [3] Gutin G., *m-sources in complete multipartite digraphs*, Vestsi Acad. Navuk BSSR Ser. Fiz-Mat, 5, (1989), 101-106 (in Russian).
- [4] Koh K.M., Tay E.G., *On optimal orientations of G vertex-multiplications*, Discrete Mathematics, 219, (2000), 153-171.
- [5] Koh K.M., Tay E.G., *On optimal orientations of tree vertex-multiplications*, Australasian Journal of Combinatorics, 34, (2006), 69-87.
- [6] Maurer S.B., *The king chicken theorems*, Mathematics Magazine, 53, (1980), 67-80.
- [7] Ng K.L., Koh K.M. *On optimal orientation of cycle vertex multiplications*, Discrete Mathematics, 297, (2005), 104-118.

- [8] Plesnik J., *Diametrically critical tournaments*, Časopis pro pěstování matematiky, 100, (1975), 361-370.
- [9] Robbins H.E., *A Theorem on Graphs, with an Application to a Problem of Traffic Control*, The American Mathematical Monthly, 46, (1939), 281-283.
- [10] Šoltés L., *Orientations of graphs minimizing the radius or the diameter*, Math. Slovaca, 36, (1986), 289-296.