THE NIRENBERG PROBLEM FOR CONICAL SINGULARITIES

LISANDRA HERNANDEZ-VAZQUEZ

ABSTRACT. We propose a new approach to the question of prescribing Gaussian curvature on the 2-sphere with at least three conical singularities and angles less than 2π , the main result being sufficient conditions for a positive function of class at least C^2 to be the Gaussian curvature of such a conformal conical metric on the round sphere. Our methods particularly differ from the variational approach in that they don't rely on the Moser-Trudinger inequality. Along the way, we also prove a general precompactness theorem for compact Riemann surfaces with at least three conical singularities and angles less than 2π .

1. Introduction

Conical surfaces have been extensively studied by many in different contexts (see [PSSW20, Ram18, MRS15, Che98] for some examples). This class of objects consists of Riemann surfaces Σ , without boundary, equipped with a Riemannian metric g, points $p_1, \dots p_n$ and neighborhoods U_i of each p_i such that

- (1) g is of class C^k on $\Sigma \{p_1, \ldots, p_n\}$
- (2) there exist local holomorphic coordinates $z_i:U_i\to\mathbb{C}$ on each U_i and bounded functions $\phi_i(z_i)$ of class C^k such that

(1.1)
$$g = e^{2\phi_i(z_i)}|z_i - z_i(p_i)|^{2\beta_i}|dz_i|^2$$

where $\beta_i > -1$ for all *i*. The points $p_1, \dots p_n$ are called *cone points* with cone angles $\theta_i := 2\pi(\beta_i + 1)$. If a metric g on Σ satisfies the conditions above, we say g is a conical metric representing the divisor $\beta = \sum_{i=1}^{n} \beta_i p_i$, and write (Σ, g, β) for brevity. Many of the usual topological invariants defined for smooth surfaces extend to conical ones. For instance, one defines the *generalized Euler characteristic* for the conical surface (Σ, g, β) by

$$\chi(\Sigma,\beta) := \chi(\Sigma) + \sum_{i=1}^{n} \beta_i$$

and can further prove a corresponding Gauss-Bonnet theorem ([Tro91]), which states that if K_g is the Gaussian curvature of a conformal conical metric g (see Section 2 for the definition) and dA_g is its area element, then

(1.2)
$$\int_{\Sigma} K_g dA_g = 2\pi \chi(\Sigma) + 2\pi \sum_i \beta_i = 2\pi \chi(\Sigma, \beta)$$

A classical problem for conical surfaces is characterizing those smooth functions which arise as Gaussian curvatures of a pointwise conformal metric. Equivalently, one asks for necessary and sufficient conditions for existence of solutions to the Gaussian curvature equation of a pointwise conformal metric $\tilde{g} = e^{2u}g$, with g a conical metric representing a given divisor β . Such equation is given by

$$(1.3) K = e^{-2u}(K_g - \Delta_g u)$$

where K_g is the curvature of the metric g. In [Tro91], Troyanov proves several existence and uniqueness results analogous to the ones shown by Kazdan-Warner in [KW75] for smooth Riemann surfaces. Since the pioneering work of Troyanov in [Tro91], where he uses a variational method to find solutions to (1.3), several other methods have been employed. These include complex analytic ideas [Ere04], Ricci flow [MRS15], minmax theory [CM12] and recently, synthetic geometric methods when the surface is a sphere [MP19].

Date: June 1, 2018.

In the case of constant curvature, there is a complete existence theory developed over the years [Ere04, McO88, LT92, Tro91] for conical surfaces with at least three conical singularities and angles less than 2π . In particular, it has been observed by many that a necessary condition for the existence and uniqueness of such conformal conical metric of constant curvature 1 on S^2 is

(1.4)
$$\sum_{i \neq j} \beta_i < \beta_j, \text{ for all } j$$

Condition (1.4) is known as the *Troyanov condition* and we refer the reader to [MW15] for a geometric interpretation. A particularly delicate case is when the conical surface is a sphere, referred to in the literature as the *singular or conical* Nirenberg problem. The classical Nirenberg problem asks to characterize those smooth functions on the sphere which arise as the Gaussian curvature of a metric that is pointwise conformal to the round metric of curvature 1. Even though much progress has been done over the years (see for instance, [CY87, Han90, Ji04, DL95, CGY93, CL93]), the problem still remains open in full generality. In passing to the singular case, one now allows metrics in a given conformal class to have conical singularities. While there are some results on this topic in the literature [dMLS15, Tro91], many questions remain unanswered. In this work we propose a new approach to the singular Nirenberg problem when there are at least three cone points and the angles are less than 2π . Using our methods we find sufficient conditions for a function K to arise as the Gaussian curvature of a conformal conical metric in this setting. Specifically, we show

Theorem 1.1. Suppose $n \ge 3$, and $\beta = \sum_{i=1}^n \beta_i p_i$ is a divisor on S^2 satisfying the Troyanov condition (1.4) and there exists i, j, k distinct for which $\beta_i, \beta_j, \beta_k$ are all distinct. Assume $\chi(S^2, \beta) > 0$ and let g_β be the unique conical metric on S^2 representing the divisor β of Gaussian curvature $K_\beta = 1$. Then a function K on S^2 is the Gaussian curvature of a metric g conformal to g_β if K is a positive function in $C^{k-2,\alpha}_{\gamma-2}, k \ge 2, \alpha \in (0,1)$, where $\gamma = (\gamma_1, \ldots, \gamma_n) \in \mathbb{R}^n, \gamma_i \neq \frac{m}{\beta_i}, \gamma_i > 0$ and m is an integer.

The space $C_{\gamma-2}^{k-2,\alpha}$ consists of Hölder continuous functions which are (k-2)-times differentiable and satisfy growth conditions near the cone points that are determined by the weights γ (for a precise definition see Section 2). Following the ideas in recent work of Anderson [And17], our approach is to choose appropriate Banach spaces such that the differential operator defined by (1.3) is a proper Fredholm map of index zero. For such operators, there is a well-defined notion of degree, and one can perform a degree count in order to study the surjectivity of this map. In establishing properness, a preliminary step is a compactness theorem for conical surfaces. Although compactness theorems for conical surfaces have been shown in [Deb20] for the uniform topology and in [Ram18] for the $C^{m,\alpha}$ topology but angles less than π , to our knowledge no result of the form required for our proof of Thm 1.1 exists in the literature. Thus in Section 2, we show

Theorem 1.2. : Let Σ be a compact Riemann surface without boundary and fix a divisor $\beta = \sum_{j=1}^{n} \beta_j p_j$ on Σ such that $-1 < \beta_j < 0$ for all j. If $M_i = (\Sigma, g_i, \beta)$ is a sequence of smooth conformal conical metrics on Σ representing the divisor β such that there exist constants $D_0, \Lambda, v_0 > 0$ for which

- (1) $diam(M_i) \leq D_0$
- (2) there exists $t_0 > 0$ such that $vol_{g_i}(B(r)) \ge v_0$ for every $r \le t_0$
- (3) $||K_{g_i}(x)||_0 \le \Lambda$ away from the cone points

Then for any $\gamma \in \mathbb{R}^n$, there exists a subsequence of (g_i) , $C^{2,\alpha}$ diffeomorphisms $F_i : \Sigma \to \Sigma$ and a $C_{\gamma}^{1,\alpha}$ conformal conical metric g representing a divisor β' such that

(1.5)
$$||(F_i^*g_i)_{st} - g_{st}||_{1,\alpha;\gamma} \to 0$$
 as $i \to \infty$, where $\beta' = \sum_{j=1}^m \beta'_j q_j$ with $-1 < \beta'_j \le 0$, $m \le n$.

This result, however, only guarantees control of the metrics $g = e^{2u_i}g_{\beta}$ in a weighted $C^{m,\alpha}$ topology modulo diffeomorphisms. In order to obtain properness of the differential operator defined by (1.3), one

needs to ensure the conformal factors u_i themselves converge on a subsequence. As in [And17], one can further show that also in the conical case we actually have control of the metrics modulo the conformal group of g_{β} , where Astala's theorem ([Ast94]) plays a central role in this step. To obtain that the conformal factors u_i themselves converge, we rely on the fact that a conformal conical metric on a sphere with at least three cone points has compact conformal group (see Thm 3.1).

Acknowledgements. I would first like to thank my advisor Michael Anderson for all his help and advice in the completion of this project. I would also like to thank Dror Varolin and Frederik Benirschke for their time in discussing aspects of this work with me.

2. Compactness Theorems For Conical Surfaces

As mentioned in the introduction, a C^k ($k \ge 2$) conformal conical metric has an associated curvature function defined on the complement of the cone points. In fact, using (1.1), we can explicitly compute the Gaussian curvature in a neighborhood of any given point as stated in the following lemma.

Lemma 2.1. Let Σ be a compact Riemann surface and g a conformal conical metric representing the divisor β . If z is a holomorphic coordinate in a neighborhood of $p \in \Sigma$ such that z(p) = 0 and the conformal conical metric is given locally by $g = e^{2u}|z|^{2\beta}|dz|^2$, then the Gaussian curvature K of g satisfies

$$(2.1) K = -\frac{e^{-2u}\Delta u}{|z|^{2\beta}}$$

for $z \neq 0$, where Δ is the Euclidean laplacian.

A natural question is whether there is an associated compactness or precompactness theorem for the class of conical Riemann surfaces with curvature, diameter and volume bounds. We address this question following the ideas in the classical work of Anderson ([And90a], see also [AC92]). To begin, a chart *H* will be referred to as an *isothermal chart* if there exists a holomorphic coordinate *z* such that

$$(2.2) (H^{-1})^* g = e^{2\phi(z)} \prod_{i=1}^m |z - z^j|^{2\alpha_j} |dz|^2$$

Observe that the existence of such charts is guaranteed by the uniformization theorem.

In order to directly apply our results in Section 3, we will work with weighted Hölder spaces, which we define here as follows. To set things up, we first introduce a slightly different but equivalent definition of a conical surface as the one given in the introduction. Let Σ be a compact Riemann surface without boundary and fix a C^k , $k \ge 2$, metric g_0 on Σ , points $p_1, \ldots p_n$. Define $\Sigma^* = \Sigma - \{p_1, \ldots p_n\}$. For R > 0 small enough, let $B_R(p)$ denote a geodesic ball in (Σ, g_0) of radius R centered at p. Set $\Sigma_R := \Sigma - \bigcup_{i=1}^n \bar{B}_R(p_i)$, where $\bar{B}_R(p_i)$ denotes the closure of $\bar{B}_R(p_i)$ in (Σ, g_0) .

Definition 2.2. A smooth function $\rho: \Sigma^* \to (0,1]$ is a *radius function* on (Σ^*, g_0) , if for R > 0 small enough, $\rho \equiv 1$ on Σ_R and for each $i = 1, \ldots n$, there exist holomorphic coordinates z_i on $B_R(p_i)$ such that $\rho(z_i) = O(|z_i|)$ on $B_R(p_i)$.

We further define ρ^{γ} for $\gamma = (\gamma_1, \dots, \gamma_n)$ in \mathbb{R}^n as follows

$$\rho^{\gamma} = \rho^{\gamma_i} \text{ on } B_R(p_i)$$

$$\rho^{\gamma} \equiv 1 \text{ otherwise}$$

Moreover, for $a \in \mathbb{R}$, set $\gamma + a := (\gamma_1 + a, \dots, \gamma_n + a)$. We say $\gamma \le \delta$, $\gamma, \delta \in \mathbb{R}^n$ provided that $\gamma_i \le \delta_i$ for each $i = 1, \dots, n$.

Definition 2.3. We say g_{β} is a *conical metric* on (Σ, g_0) of class C^k $(k \ge 2)$ representing the divisor $\beta = \sum_{i=1}^n \beta_i p_i$ if g_0 is of class C^k and there exists a smooth radius function $\rho : \Sigma^* \to (0, 1]$ such that

$$(2.3) g_{\beta} = \rho^{2\beta} g_0$$

Moreover, g is called a *conformal conical metric* on (Σ, g_0) of class C^k representing the divisor β if there exists a smooth positive function $u : \Sigma \to \mathbb{R}$ such that

$$(2.4) g = e^{2u}g_{\beta}$$

where g_{β} is a conical metric on Σ of class C^k representing the divisor β .

For g_{β} a conical metric on (Σ, g_0) representing a divisor β and ρ a radius function on (Σ^*, g_0) such that $g_{\beta} = \rho^{2\beta}g_0$, if $u \in C^k_{loc}(\Sigma^*)$ and $\gamma \in \mathbb{R}^n$, set

(2.5)
$$||u||_{C^k_{\gamma}} := \sum_{j=0}^k \sup_{x \in \Sigma^*} |\rho(x)^{-\gamma+j} \nabla^j u(x)|$$

Define the space of $C^{k,\alpha}_{\gamma}(\Sigma,\beta)$ functions on (Σ,β) to be

(2.6)
$$C_{\gamma}^{k,\alpha}(\Sigma,\beta) = \left\{ u \in C_{loc}^{k}(\Sigma^{*}) : ||u||_{k,\alpha;\gamma} < \infty \right\}$$

where the norm $\|\cdot\|_{k,\alpha;\gamma}$ is given by

(2.7)
$$||u||_{k,\alpha;\gamma} := ||u||_{C^{k,\alpha}_{\gamma}} = ||u||_{C^k_{\gamma}} + [\nabla^k u]_{\alpha,\gamma-k}$$

and

$$(2.8) \qquad [\nabla^k u]_{\alpha;\gamma} = \sup_{\substack{x \neq y, d(x,y) < in \ i(x)}} \min(\rho(x)^{-\gamma}, \rho(y)^{-\gamma}) \frac{|\nabla^k u(x) - \nabla^k u(y)|}{d(x,y)^{\alpha}}$$

It is well known that the normed spaces $(C_{\gamma}^{k,\alpha}, \|\cdot\|_{k,\alpha;\gamma})$ are Banach for any γ (see for instance [Pac06]). One can further define the weighted Sobolev spaces $W_{\gamma}^{k,p}$, as in [Beh11]. Next, we introduce the notion of the *isothermal radius*, which plays the same role as that of the harmonic radius in [And90a, AC92].

Definition 2.4. The Isothermal Radius: Let (Σ, g, β) be a complete Riemann surface without boundary with g a conformal conical metric on Σ representing the divisor $\beta = \sum_{i=1}^{n} \beta_i p_i$. Let $x \in \Sigma$. Given a constant C > 1, $\alpha \in (0, 1), \gamma \in \mathbb{R}^n$, we define the isothermal radius $r_I = r_I(g, x, C, \alpha, \gamma)$ as the largest number such that on the geodesic ball $B(x, r_I(x))$ there exists an isothermal coordinate chart $H : B(x, r_I(x)) \to B_0(R) \subset \mathbb{C}$ with

(2.9)
$$(H^{-1})^* g = e^{2\phi(z)} \prod_{i=1}^m |z - z^j|^{2\alpha_j} |dz|^2$$

where z is a holomorphic coordinate on $B_0(R)$, $z^j = H(p_j)$ correspond to cone points, $1 \le m \le n$, $-1 < \alpha_j$ and $\phi(z) : B_0(R) \to \mathbb{R}$ is smooth and satisfies

A1.
$$\frac{1}{C} \le \phi(z) \le C$$
B1.
$$\sum_{0 \le |\mu| \le 1} r_I^{-\gamma + |\mu|} \sup_{x} |\partial^{\mu} \phi(x)| + \sum_{|\mu| = 1} r_I^{-\gamma + \alpha + 1} \sup_{x \ne y} \frac{|\partial^{\mu} \phi(x) - \partial^{\mu} \phi(y)|}{d(x, y)^{\alpha}} \le C - 1$$

We also define the isothermal radius of (Σ, β, g) by

(2.10)
$$r_I(\Sigma) = \inf_{\substack{x \in \Sigma \\ A}} r_I(x)$$

Observe that α_j either coincide with the β_j or are zero in neighborhoods with no conical points. In the following lemma we prove properties of the isothermal radius that will be needed for the upcoming blow-up argument in the proof of Theorem 2.9. All of these facts are true for the harmonic radius and the proofs presented here are essentially the same as in the works of [And90a, AC92, HH97], only slightly modified to fit our setting.

Lemma 2.5. Let (Σ, β, g) be a Riemann surface with a conformal conical metric g representing the divisor β and let $r_I : \Sigma \to \mathbb{R}$ be the isothermal radius. Then the following hold

- (1) $r_I(x)$ is positive and pointwise continuous on Σ
- (2) If $F:(\Sigma,g)\to(\Sigma',g')$ is an isometry, then

$$F^*r_I(\Sigma') = r_I(\Sigma)$$

(3) r_I scales as the distance: $r_I(\lambda^2 g, x) = \lambda r_I(g, x)$

Proof. Proof of (1). For the positivity, observe that for any $x \in \Sigma$ we can find a $0 \ge \mu > -1$, a neighborhood U of x and a holomorphic coordinate $z: U \to \mathbb{C}$ such that $g = e^{2\phi}|z|^{2\mu}|dz|^2$ for some smooth $\phi(z): U \to \mathbb{R}$. Clearly, if x is any of the cone points, such coordinates exist by definition once we choose $\mu = \frac{\theta}{2\pi}$ where θ is the cone angle. If on the other hand x is a smooth point, then such coordinates are guaranteed by the Uniformization Theorem: we can find a neighborhood U of x and a holomorphic coordinate $z: U \to \mathbb{C}$ such that $g = e^{2\phi}|dz|^2$. Finally observe that conditions A1, B1 always hold on a fixed conical surface in a given isothermal chart.

For the continuity, given any x close enough to y, we can find a ball of radius a centered at x which contains all the cone points in $B(y, r_I(y))$ and is contained in $B(y, r_I(y))$. Therefore if z is a holomorphic coordinate in $B(y, r_I(y))$ such that

$$g = e^{2\phi(z)} \prod_{i=1}^{m} |z - z_i|^{2\alpha_i} |dz|^2$$

with $\phi(z)$ satisfying conditions A1, B1 of Definition 2.4, by restricting the coordinates z to B(x, a), we get a holomorphic coordinate on this ball such that in this coordinate the same function $\phi(z)$ still satisfies conditions A1, B1 for the same C, α, γ . Therefore $a \le r_I(x)$. Using this one can directly show

$$|r_I(y) - r_I(x)| = |a + d(x, y) - r_I(x)| \le \epsilon$$

as wanted.

Proof of (2). We will actually prove something stronger than the statement of (2): computing the isothermal radius with respect to g' at a point F(x) gives the same result as computing the isothermal radius with respect to g at the point x. To begin, fix $x \in \Sigma$ and let $H': B(F(x), r_I F(x)) \to B_0(R)$ be an isothermal coordinate chart where

$$(2.11) (H'^{-1})^* g' = e^{2\phi(z)} \prod_{i=1}^m |z - z_i|^{2\alpha_i} |dz|^2$$

and $\phi(z)$ satisfies conditions A1, B1 of Definition 2.4. The set $U = F^{-1}(B(F(x), r_I F(x)))$ is open in Σ and contains x. If B(x, R) is the largest geodesic ball centered at x which is still contained in U, then $H := H' \circ F : B(x, R) \to B_0(R)$ is an isothermal coordinate chart for the metric g on B(x, R). Moreover, the metric g on the ball B(x, R) has the form (2.9) above, where the conformal factor can be computed to be $2\phi(F(z)) + \|DF\|^2$ and satisfies the bounds A1, B1 in these coordinates since $\|DF\|_{(F^{-1})^*g} = 1$. We have thus found a ball centered at x in which all the conditions of the definition of the isothermal radius are satisfied, so we conclude that its radius $R \le r_I(x)$. In particular, $r_I(F(x), g') \le r_I(x, g)$. To obtain the opposite inequality, we just follow the same argument with F^{-1} in place of F.

Proof of (3). We want to show that $r_I(\lambda^2 g, x) = \lambda r_I(g, x)$ for any nonzero λ . Start with a conical metric g and after having chosen a holomorphic coordinate for which we can write the metric g as in (2.9) of Definition

2.4, with $\phi(z)$ satisfying the bounds A1, B1, it is straightforward to check that the metric $\lambda^2 g$ satisfies the same bounds in the coordinates $w = \lambda z$.

We now turn to the question of convergence. For $k \geq 2$, $\alpha \in (0,1)$, $\gamma \in \mathbb{R}^n$, we say a conformal conical metric g on (Σ, g_0) representing the divisor β is of class $C_{\gamma}^{k,\alpha}$ if in an isothermal chart the coefficients g_{ij} of g are bounded in $C_{\gamma}^{k,\alpha}$. Moreover, a sequence of conformal conical metrics (Σ, g_i, β) of class $C_{\gamma}^{k,\alpha}$ converges in $C_{\gamma}^{k,\alpha}$ to a surface (Σ', g) provided that there exists a sequence of $C_{\gamma}^{k+1,\alpha}$ diffeomorphisms $F_i: \Sigma' \to \Sigma$ such that for all i large enough

in any chart of a C^{∞} subatlas of the complete C^{∞} atlas of Σ . The following lemma addresses the continuity of the isothermal radius in the $C^{1,\alpha}_{\nu}$ topology.

Lemma 2.6. For $\alpha \in (0,1)$, $\gamma \in \mathbb{R}^n$, the isothermal radius is continuous under $C_{\gamma}^{1,\alpha}$ convergence of a sequence of conical metrics (Σ, g_i, β) representing the divisor β .

Proof. Let (Σ, g_i, β) be a sequence of conformal conical metrics on Σ representing the divisor β and $x \in \Sigma$. As before, $\Sigma^* = \Sigma - \{p_1, \dots p_n\}$, where $p_1, \dots p_n$ are the cone points. Fix $\alpha \in (0, 1)$, $\gamma \in \mathbb{R}^n$. Assume that the sequence (g_i) converges in $C_{\gamma}^{1,\alpha}$ to a $C_{\gamma}^{1,\alpha}$ metric g on Σ . By the continuity of the isothermal radius we mean explicitly that the following two inequalities hold: given C > 1

$$(2.13) r_I(g, x, C) \ge \limsup_{i \to \infty} r_I(g_i, x, C)$$

and for any $0 < \epsilon < C - 1$

$$(2.14) r_I(g, x, C - \epsilon) \le \liminf_{i \to \infty} r_I(g_i, x, C)$$

For simplicity, set $r_i := r_I(g_i, x, C)$ and let $H_i : B(x, r_i) \to \mathbb{C}$, $H_i(y) = (H_i^1(y), H_i^2(y)) = z_i \in \mathbb{C}$ be isothermal coordinate charts in which the metrics g_i satisfy A1, B1 of Definition 2.4.

We begin with some preliminary claims, the first being that for any $r \le \limsup r_i$, a subsequence of the isothermal charts H_i converges in $C^{2,\alpha}$ to an isothermal chart $H: B(x,r) \to \mathbb{C}$, where B(x,r) is a geodesic ball for the metric g. To this end, suppose $(x_1(y), x_2(y)) \in \mathbb{C}$ is any given local coordinate chart on B(x,r). Observe that it is implicit in the fact that the charts H_i are isothermal that the coordinate functions H_i^k , for k = 1, 2 are harmonic. In other words,

(2.15)
$$(g_i)^{st} \frac{\partial^2 H_i^k}{\partial x_s \partial x_t} = (g_i)^{st} (\Gamma_i)_{st}^l \frac{\partial H_i^k}{\partial x_t}$$

where $(g_i)^{st}$ are the components of g_i in the coordinates (x_1, x_2) and $(\Gamma_i)_{st}^l$ are the Christoffel symbols for g_i in these coordinates. Now, condition A1 of Definition (2.4) implies that in the charts H_i , the components of the metrics g_i satisfy

(2.16)
$$\frac{1}{C} \prod_{i=1}^{m_i} |z_i - z_i^j|^{\alpha_j} \delta_{kl} \le (g_i)_{kl} \le C \prod_{i=1}^{m_i} |z_i - z_i^j|^{\alpha_j} \delta_{kl}$$

where we write $z_i = H_i^1 + iH_i^2$ and the inequality holds as bilinear forms. It then follows from (2.16) and the fact that the metrics converge in $C_{\gamma}^{1,\alpha}$ that the charts H_i are bounded in C^1 on Σ^* . Using standard elliptic estimates for (2.15) (see [GT83], for instance), we obtain that for each k = 1, 2, the sequence (H_i^k) is bounded in $C^{2,\alpha}(\Sigma^*)$. Therefore, by the Arzela-Ascoli theorem, we have that for each k = 1, 2, the sequences (H_i^k) converge weakly on a subsequence in $C^{2,\alpha'}$ on Σ^* for $\alpha' \leq \alpha$. In fact, repeating this argument for $H_q^k - H_n^k$ in place of H_i^k one can see that for each k = 1, 2, the sequences (H_i^k) are in fact Cauchy. Therefore, they converge strongly on a subsequence in $C_{loc}^{2,\alpha}$ to a limiting map $H: B(x,r) - \{p_1, \dots p_m\} \to \mathbb{C}$, where

 $H_i(p_j) = z^j$ for all *i*. Since the property of being an isothermal chart is preserved under $C^{2,\alpha}$ convergence, H is an isothermal chart for the metric g. Moreover, since there are finitely many cone points, by passing to a subsequence, we may assume m_i in (2.16) is independent of i and $z_i^j = z^j$ for all i. Therefore, g can be written in the chart given by H as

(2.17)
$$(H^{-1})^* g = e^{2\phi(z)} \prod_{j=1}^m |z - z^j|^{\alpha_j} |dz|^2$$

where $\phi(z)$ is a smooth function that satisfies conditions A1, B1 of Definition 2.4. Hence for any $r \le \limsup r_i$, a subsequence of the isothermal charts H_i converges in $C^{2,\alpha}$ to an isothermal chart $H: B(x,r) \to \mathbb{C}$ for the metric g. Observe that this argument also shows that if a sequence of conformal conical metrics representing a fixed divisor β converges in $C_{\gamma}^{1,\alpha}$ to a metric g, then g has at most as many cone points as the sequence g_i and no other types of singularities.

Our second preliminary claim is that if (Σ, g, β) is any complete Riemann surface without boundary with a conformal conical metric g representing the divisor β , $x \in \Sigma$, $y \in \mathbb{R}^n$, then for any $1 \le C' \le C < \infty$,

(2.18)
$$r_I(C')(x) \le r_I(C)(x)$$

and for any C > 1

(2.19)
$$\lim_{\epsilon \to 0^+} r_I(C + \epsilon)(x) = r_I(C)(x)$$

The first inequality follows from the definition, hence to prove the claim, it's enough to show that for any C > 1

(2.20)
$$\limsup_{\epsilon \to 0^+} r_I(C + \epsilon, x) \le r_I(C, x)$$

Fix $r < \limsup r_I(C + \epsilon, x)$. For a decreasing sequence of $\epsilon > 0$ converging to 0, there are isothermal coordinate charts H_{ϵ} on $B_x(r)$ satisfying conditions A1, B1 of Definition 2.4 with $C + \epsilon$ in place of C and C in place of C in place of C and C in place of C in

We're now ready to prove the first inequality (2.13). As before, let $r_i = r_I(g_i)$. We may suppose $\limsup r_i > 0$. The arguments above show that convergence of the metrics in $C_{\gamma}^{1,\alpha}$ implies convergence of the isothermal charts H_i in $C_{loc}^{2,\alpha}$. Once again, the bounds A1, B1 are preserved, so that $r_I(g, C) \ge r$ for any $r \le \limsup r_i$. Therefore we get the first inequality $r_I(g, C) \ge \limsup r_I(g_i, C)$.

Now fix $r < r_I(g,C)$. To obtain the second inequality (2.14), let $H: B(x,r) \to B_0$ be an isothermal coordinate chart for g, so that $(H^{-1})^*g = e^{2\phi(z)} \prod |z-z^j|^{2\alpha_j}|dz|^2$, with z a holomorphic coordinate on $B_0 \subset \mathbb{C}$. Let Δ_i be the Laplace operator for the metric g_i . In the coordinate z, the Laplacian for the metrics g_i has the form

(2.21)
$$\Delta_i = e^{-2(\phi_i(z) + \sum_{j=1}^m \alpha_j \log|z - z^j|)} \Delta$$

where Δ is the Euclidean laplacian. As observed before, we may assume the cone points z^j are the same for each i after passing to a subsequence. Now, if w_i are solutions to

$$\Delta_i w_i = 0 \text{ on } B$$

$$(2.23) w_i = z \text{ on } \partial B$$

then the functions $u_i := z - w_i$ are harmonic and vanish on the boundary of B. Since, for each i, the function z also solves the boundary value problem (2.22-2.23) it follows from uniqueness that in fact $u_i = 0$. Therefore, $\lim_{i \to \infty} ||u_i||_{2,\alpha} = 0$ and we have that for any compact subset $B' \subset B$ and for any i, i is an isothermal coordinate chart for g_i . Now, since the metrics g_i converge to g in $C_{\gamma}^{1,\alpha}$, i is an isothermal

coordinate chart for g_i in which the bounds of Definition 2.4 are satisfied with constants $C_i \to C$ as $i \to \infty$. Using (2.18), (2.19), we have that for any $\epsilon > 0$

$$r \le \liminf r_I(g_i, C_i) \le \liminf r_I(g_i, C + \epsilon)$$

Since $r \le r_I(g, C)$ was arbitrary, this ends the proof of the second inequality.

An important property of the harmonic radius in the smooth case is that Euclidean space has infinite harmonic radius. The isothermal radius satisfies an analogous condition, but in this case the model space is a flat Riemann surface M with finitely many cone points and angles less than 2π , which is noncompact, complete and of *quadratic area growth*, in the sense that for any $x \in M$ and any r > 0

$$(2.24) \frac{1}{V}r^2 \le vol(B(r,x)) \le Vr^2$$

Theorem 2.7. Any noncompact conical surface with a finite number of conical singularities and angles less than 2π which is flat, complete and of quadratic area growth has infinite isothermal radius.

Proof. Suppose M is a complete flat conical surface of quadratic area growth, so that there exist p_1, \ldots, p_n in M such that near each p_i we can find a coordinate z and a harmonic function u with $g = e^{2u}|z - z_i|^{2\beta_i}|dz|^2$ where $z_i = z(p_i)$ and (2.24) is satisfied. If we smooth out the singularities p_i , the resulting surface M' is still noncompact and complete. Moreover, since the cone angles are less than 2π , the curvature can only increase upon smoothing the singularities, thus M' has Gaussian curvature $K \ge 0$. It further follows from the volume comparison theorem of Bishop-Gromov [Pet16] that if the original (singular) surface has quadratic area growth, then any smoothing M' has at most quadratic area growth.

Now fix a smoothing M' of M and let \tilde{M} be its universal cover. Since M' is complete and noncompact, the pullback of the metric on M' to \tilde{M} by the covering map makes \tilde{M} into a simply connected, complete, noncompact surface with Gaussian curvature $K \geq 0$. A complete surface is hyperbolic if it admits a positive Green's function. On the other hand, a theorem of Yau [Yau75] (see also [BF42]) asserts that a complete surface of nonnegative Gaussian curvature admits no non-constant positive superharmonic functions. Thus \tilde{M} cannot be hyperbolic and by the uniformization theorem, we have \tilde{M} is parabolic, i.e. it is conformally equivalent to the complex plane.

We actually claim that M' is simply connected, so that M' is parabolic. By Bishop-Gromov again, the universal cover \tilde{M} of M' has at most quadratic area growth. Moreover, since M' has at least quadratic area growth, by Proposition 1.2 in [And90b] we then have that M' is the quotient of \tilde{M} by a finite group of isometries Γ . Since M' is smooth, Γ must be trivial.

Since smoothing out the singularities doesn't change the topology or the conformal structure, M is also simply connected and parabolic. Therefore, there exists a global coordinate z such that the metric on M has the form $e^{2v}|dz|^2$. Since M has conical singularities, $e^{2v} = e^{2u} \prod |z - z_i|^{2\alpha_i}$. Hence, there exist global coordinates on M for which the metric has the form

$$(2.25) g = e^{2u} \prod |z - z_i|^{2\alpha_i} |dz|^2$$

where u is harmonic. At this point, we have shown that a noncompact, complete, flat conical surface with cone angles less than 2π and quadratic area growth is conformally equivalent to the complex plane. Our final claim is that the function u in (2.25) has to be constant. To see this, define

$$(2.26) h = \prod |z - z_i|^{-\alpha_i} g$$

where g is the metric in (2.25). In other words, $h = e^{2u}|dz|^2$. Since g is flat by assumption, the function u is harmonic. Therefore, the Gaussian curvature K_h of h satisfies $K_h = e^{-2u}\Delta u = 0$. On the other hand, since the cone angles are less than 2π , $\alpha_i < 0$ for all i. Therefore, if γ is any C^1 curve, then far away from the cone points,

$$(2.27) \qquad \int h(\dot{\gamma}(t),\dot{\gamma}(t))^{\frac{1}{2}}dt = \int \prod |z(\gamma(t)) - z_i(\gamma(t))|^{\frac{-\alpha_i}{2}} g(\dot{\gamma}(t),\dot{\gamma}(t))^{\frac{1}{2}}dt \ge \int g(\dot{\gamma}(t),\dot{\gamma}(t))^{\frac{1}{2}}dt$$

where the inequality follows since $\prod |z(\gamma(t)) - z_i(\gamma(t))|^{\frac{-\alpha_i}{2}} >> 1$ whenever $|z(\gamma(t)) - z_i(\gamma(t))| >> 1$. The assumption that g is complete together with (2.27) now imply that h is complete.

Now, let $F: T_0\mathbb{C} \to \mathbb{C}$ be the exponential map of the origin for the metric h. By the above arguments, the smooth metric h is flat and complete, so by Cartan-Hadarmard's theorem (see [Pet16], Thm 22) the exponential map F is in fact a diffeomorphism of \mathbb{C} that satisfies

$$(2.28) F^*h = e^{2u(0)}|dz|^2$$

In other words,

$$F^*h = e^{u \circ F} |\partial F + \bar{\partial} F|^2 = e^{u \circ F} (|\partial F|^2 + |\bar{\partial} F|^2 + 2Re\partial F\bar{\partial} F) = e^{2u(0)} |dz|^2$$

The last equality implies that either $\partial F = 0$ or $\bar{\partial} F = 0$, and since the orientation of the tangent space is the same as the base manifold, we must have that $\bar{\partial} F = 0$, so F is holomorphic. A standard result in complex analysis is that holomorphic diffeomorphisms of the plane are affine linear maps. Given that F preserves the origin (it is the exponential map at the origin), we conclude that it must be of the form

$$F(\zeta) = c\zeta$$

for some $c \neq 0$. But then $e^{u \circ F} = e^{u \circ (c\zeta)} = e^{2u(0)}$, so u has to be constant.

The sequence of arguments above now show that M admits global coordinates for which the metric has the form

$$g = C \prod |z - z_i|^{2\alpha_i} |dz|^2$$

from which it follows that the isothermal radius must be infinite.

The following is a generalization of $C^{1,\alpha}$ convergence to weighted $C^{1,\alpha}_{\gamma}$ convergence of conical metrics on Riemann surfaces.

Theorem 2.8. Let $M_i = (\Sigma, g_i, \beta)$ be a sequence of complete, conformal conical surfaces with metrics g_i representing the divisor β . Let $\{x_i\} \in M_i$ be a sequence of points. Given $\Lambda > 0, C > 1, \alpha \in (0, 1)$, suppose that

- (1) for any i, $||K(g_i)||_0 \le \Lambda$ away from the cone points
- (2) there exists r > 0 such that for any sequence of points (y_i) in M_i there is an isothermal chart $H_i: \Omega_n \to B_0(r)$ where Ω_i is some open set in M_i and $B_0(r) \subset \mathbb{C}$ such that for any i, there exists ϕ_i smooth, with $\frac{1}{C} \leq \phi_i(z) \leq C$ such that

(2.29)
$$(H_i^{-1})^* g_i = e^{2\phi_i(z)} \prod_{j=1}^m |z - z_j|^{2\alpha_j} |dz|^2$$

where z are holomorphic coordinates on $B_0(r)$ and $z_j = H_n(p_j)$ correspond to the cone points p_j

(3) for $\gamma \in \mathbb{R}^n$, a subsequence of $(H_i^{-1})^* g_i$ converges in $C_{\gamma}^{1,\alpha}$ on $B_0(r)$

Then there exists a complete Riemannian manifold M of class $C^{2,\alpha}$, there exists a conformal conical metric g of class $C^{1,\alpha}_{\gamma}$ and a point $x \in M$ such that the following holds: for any compact domain $D \subset M$ with $x \in D$ there exist, up to passing to a subsequence, compact domains $D_i \subset M_i$ with points $x_i \in D_i$ and $C^{2,\alpha}$ diffeomorphisms $\Phi_n : D \to D_i$ satisfying

(2.30)
$$\lim_{i \to \infty} (\Phi_i^{-1})(x_i) = x$$

(2.31) $\Phi_i^* g_i$ converges in $C_{\gamma}^{1,\alpha}$ in any chart of the induced $C^{2,\alpha}$ complete atlas of D.

Proof. The proof is exactly as in the smooth case (see [HH97, Pet87] and [And90a] for a summarized version) since conical surfaces are considered to have singularities only in the metric sense, as can be seen from the local coordinate expression in (2.29) above.

Theorem 2.9. : Let Σ be a compact Riemann surface without boundary and fix a divisor $\beta = \sum_{j=1}^{n} \beta_j p_j$ on Σ such that $-1 < \beta_j < 0$ for all j. If $M_i = (\Sigma, g_i, \beta)$ is a sequence of smooth conformal conical metrics on Σ representing the divisor β such that there exist constants $D_0, \Lambda, v_0 > 0$ for which

- (1) $diam(M_i) \leq D_0$
- (2) there exists $t_0 > 0$ such that $vol_{g_i}(B(r)) \ge v_0$ for every $r \le t_0$
- (3) $||K_{g_i}(x)||_0 \le \Lambda$ away from the cone points

Then for any $\gamma \in \mathbb{R}^n$, there exists a subsequence of (g_i) , $C^{2,\alpha}$ diffeomorphisms $F_i : \Sigma \to \Sigma$ and a $C^{1,\alpha}_{\gamma}$ conformal conical metric g representing a divisor β' such that

$$||(F_i^*g_i)_{st} - g_{st}||_{1,\alpha;\gamma} \to 0$$

as
$$i \to \infty$$
, where $\beta' = \sum_{i=1}^{m} \beta'_{i} q_{j}$ with $-1 < \beta'_{i} \le 0$, $m \le n$.

Proof. The first part of the proof is a blow-up argument to show that, under the hypotheses of the theorem, there is a uniform lower bound on the isothermal radius. So to begin, let D_0 , Λ , v_0 be positive constants as in the statement of the theorem and $\alpha \in (0, 1)$. Given a conical Riemann surface (Σ, g, β) and $C, t_0 > 0$ satisfying

$$(2.33) ||K||_0 \le \Lambda$$

$$(2.34) vol_{g}(B(r)) \ge v_{0} > 0 \ \forall r \le t_{0}$$

we show that there exists $r_0 = r_0(\Lambda, v_0) > 0$ such that for every $x \in B(r)$ and any $\gamma \in \mathbb{R}^n$

$$(2.35) \frac{r_I(x, C, \alpha)}{d_g(x, \partial B(r))} \ge r_0 > 0$$

where $r_I(x, C, p)$ is as usual the isothermal radius. Indeed, if (2.35) does not hold, there exists a sequence of conical metrics $M_i = (\Sigma_i, g_i, \beta)$ representing the same divisor β with Gaussian curvature $|K_i|_0 \le \Lambda$, there exists a sequence of balls $B_i = B_i(r) \subset M_i$ of radius $r \le t_0$, there exists $\gamma \in \mathbb{R}^n$ and there exists a sequence of points $x_i \in B_i$ such that

$$\frac{r_i(x_i)}{d_i(x_i, \partial B_i)} \to 0$$

where $d_i = d_{g_i}$ is the induced distance on M_i from g_i and $r_i(x) = r_I(g_i, x)$ is the isothermal radius of the metric g_i at x.

Set $R_i(x) := \frac{r_i(x)}{d_i(x,\partial B_i)}$. By the same arguments as in [HH97, And90a], we may as well assume the points x_i minimize $R_i(x)$. Rescaling the metrics g_i as

$$(2.37) h_i = \frac{1}{r_i(x_i)^2} g_i$$

we get

(2.38)
$$\lim_{i \to \infty} ||K_{(B_i, h_i)}||_0 = r_i(x_i)^2 ||K_i||_0 \le r_i(x_i)^2 \Lambda \to 0$$

$$\lim_{i \to \infty} vol(B_i, h_i) = \infty$$

(2.40)
$$\lim_{i \to \infty} d(x_i, \partial B_i) = \infty$$

where (2.37) holds away from the cone points. By Lemma 2.5 (3), the isothermal radius scales as the distance under rescalings of the metric, thus the isothermal radius of the new metrics h_i satisfies

$$(2.41) r'_i(x_i) := r_I(h_i, x_i) = 1$$

Moreover, for every $y \in B_i$ and for every i

$$r_i'(y) = \frac{r_i(y)}{r_i(x_i)} \ge \frac{d_i(y, \partial B_i)}{d_i(x_i, \partial B_i)} = \frac{d_i'(y, \partial B_i)}{d_i'(x_i, \partial B_i)}$$

(where the first inequality follows since $R_i(y) \ge R_i(x_i)$ and d'_i is the induced distance from h_i). Set

(2.42)
$$\delta_i := \frac{1}{d_i'(x_i, \partial B_i)}$$

Then $\lim_{i\to\infty} \delta_i = 0$ and for all $y \in B(x_i, \frac{1}{2\delta_i})$ (the geodesic ball for the metric h_i with center x_i) we have

$$(2.43) r_i'(y) \ge \frac{1}{2}$$

Hence, the isothermal radius of the rescaled metrics is bounded from below and is at most 1. Now we claim (B_i, x_i, h_i) converges in $C_{\gamma}^{1,\alpha}$ uniformly on compact sets to a complete (noncompact) manifold (M, y, h). First, the argument above implies that given $R < \infty$, $r_i'(y) \ge \frac{1}{2}$ on $B(x_i, R)$ for i large enough. Thus given $R < \infty$ and a sequence (q_i) in $B(x_i, R)$ we can find isothermal charts $H_i : \Omega_i \to B_0(\frac{1}{2\sqrt{C}})$ centered at q_i such that

$$(2.44) (H_i^{-1})^* h_i = e^{2\phi_i(z_i)} \prod_{j=1}^{m_i} |z_i - z_i^j|^{2\alpha_j} |dz|^2$$

where $\phi_i(z_i): B_0(\frac{1}{2\sqrt{C}}) \to \mathbb{R}$ are smooth and bounded, the integers $m_i = m_i(q_i)$ and the real numbers α_j satisfy $1 \le m_i \le n, -1 < \alpha_j \le 0$ for all $i = 1, \ldots$ and $1 \le j \le m_i$. As before, z_i is a holomorphic coordinate on $B_0 = B_0(\frac{1}{2\sqrt{C}})$ and $z_i^j = H_i(p_j)$ (p_j are cone points). Moreover,

$$(2.45) \frac{1}{C} \le \phi_i(z_i) \le C$$

(2.46)
$$\|\phi_i(z_i)\|_{1,\alpha;\gamma} \le F(C)$$

with F depending only on C. By (2.46) we have that $(\phi_i(z_i))$ are bounded in $C_{\gamma}^{1,\alpha}$ on the ball B_0 , so after passing to a subsequence we can assume they converge weakly in $C_{\gamma}^{1,\alpha}$ to some ϕ by the Arzela-Ascoli Theorem. In particular, the metrics $(H_i^{-1})^*h_i =: h_i'$ converge weakly in $C_{\gamma}^{1,\alpha}$ on B_0 . We claim that in fact the metrics h_i' converge strongly in $C_{\gamma}^{1,\alpha}$. Indeed, from Lemma 2.1, we have that the Gaussian curvature K_i of the metrics h_i in the coordinates z_i is given by

(2.47)
$$K_{i}(z_{i}) = \frac{e^{-2\phi_{i}(z_{i})}\Delta\phi_{i}(z_{i})}{\prod_{i=1}^{m_{i}}|z_{i}-z_{i}^{j}|^{2\alpha_{j}}}$$

As we observed in the proof of Lemma 2.6, we can assume $z_i^j = z^j$ and $m_i = m$ are independent of i by passing to a subsequence. By (2.38), $||K_{(B_i,h_i)}||_0 \to 0$ away from the cone points, so that $||(H_i^{-1})^*K_i||_0 = ||K_i(z_i)||_0 \to 0$ as $i \to \infty$ for all $z_i \neq z^j \in B_i$. Now, the limit $\phi(z)$ of the sequence $(\phi_i(z_i))$ solves

(2.48)
$$0 = \frac{e^{-2\phi(z)}\Delta\phi(z)}{\prod_{j=1}^{m}|z-z^{j}|^{2\alpha_{j}}}$$

weakly. Applying standard elliptic estimates to (2.48), we get $\phi(z)$ is actually smooth. In fact, using the same arguments as in the proof of Lemma 2.6 (second paragraph following (2.16)), we get that the convergence is actually in the *strong* $C_{\gamma}^{1,\alpha}$ topology, hence the metrics h'_{i} converge strongly in $C_{\gamma}^{1,\alpha}$.

It now follows from the arguments used to prove Theorem 2.8 (see for instance Proposition 12 in [HH97]) that there exists a $C^{2,\alpha}$ manifold $M, y \in M$ and a $C^{1,\alpha}_{\gamma}$ conformal conical metric h on M such that for any compact domain $D \subset M$ with $y \in D$ and after passing to a subsequence, there exist compact domains $D_i \subset B_i$ and $Y_i \in D_i$ and Y_i

$$\lim_{i \to \infty} (\Phi_i^{-1})(y_i) = y$$

(2.50)
$$||(\Phi_i^{-1})^* h_i - h||_{1,\alpha;\gamma} \to 0 \text{ in } D$$

where (2.50) is in the sense that in any chart of the complete induced atlas on D the components of $(\Phi_i^{-1})^*h_i$ converge in $C_{\gamma}^{1,\alpha}$ to the components of h.

We now claim the pointed limit (M,h) is flat, conical and complete. First, the completeness follows from (2.40). To see (M,h) is flat, given a compact domain D in M, set $\hat{h}_i := \Phi_i^* h_i$ and for a given $x \in D$, let $U_i : B_x(r) \to \mathbb{C}$, r > 0, be an isothermal coordinate chart for \hat{h}_i satisfying A1, B1 of Definition 2.4. The convergence of the \hat{h}_i implies convergence in $C^{2,\alpha}$ of the charts U_i to a limiting chart $U : B_x(r) \to \mathbb{C}$, by the same arguments as in Lemma 2.6. If we write $(U^{-1})^*h = e^{2\psi(z)} \prod_{j=1}^m |z - z_j|^{2\alpha_j} |dz|^2$, then going back to (2.48) we have

(2.51)
$$0 = \frac{e^{-2\psi(z)}\Delta\psi(z)}{\prod_{j=1}^{m}|z-z_{j}|^{2\alpha_{j}}}$$

It follows from standard elliptic estimates then that the limit metric h is smooth on Σ^* . Since (2.51) is also the equation for the Gaussian curvature of h, we also have that it is flat. Moreover, assumption (2) in the Theorem along with the Bishop-Gromov volume comparison Theorem imply that

$$(2.52) \frac{vol(B(s))}{V^{\Lambda}(s)} \ge \frac{v}{V^{\Lambda}(t_0)}$$

for all $s \le t_0$, where V^{Λ} is the volume of a geodesic ball in constant curvature Λ and vol(B(s)) is taken in the g_i metric. Using scaling properties of volume and the convergence in $C_{\gamma}^{1,\alpha}$, we have

$$\frac{vol_M(B(s))}{s^2} \ge v' > 0$$

for all s > 0. On the other hand, since (M, h) is flat with cone angles less than 2π , the volume of a ball of radius s measured in the metric h has to be less than the volume of a ball of radius s measured with respect to the standard metric on \mathbb{R}^2 , hence

$$\frac{vol_M(B(s))}{s^2} \le v'$$

and we conclude that the volume growth must be exactly quadratic.

To obtain a contradiction, observe that since the h_i converge to h in $C_{\gamma}^{1,\alpha}$, by Lemma 2.6, we have that

$$(2.55) r_I(C', p, y) \le \lim_{i \to \infty} r_I(C, p, y_i)$$

for some C' < C, but by construction $r_I(C, p, y_i) = 1$, while a flat, noncompact, complete surface with finitely many conical singularities and of quadratic area growth has infinite isothermal radius for any C', as follows from Theorem 2.7.

Therefore, there is a uniform lower bound on the isothermal radius, which in turn allows us to apply Theorem 2.8 to conclude the existence of a limit with the desired properties. In the notation of Theorem 2.8, with $D = B_x(R)$, $R > D_0$, we get that for i large enough, $D_i = M_i$, and up to passing to a subsequence there exist diffeomorphisms $\Phi_i : M \to M_i$ such that $\Phi_i^* g_i$ satisfies conditions (2.30-2.31) of Theorem 2.8. In particular, M has a smooth structure coming for instance from one of the diffeomorphisms Φ_i with M_i . \square

3. Conformal Conical Metrics on the 2-sphere

In this section we are concerned with conformal conical metrics on the 2-sphere with angles less than 2π and at least three conical singularities. In the language of Definition 2.3, let $n \ge 3$, $\beta = \sum_{i=1}^{n} \beta_i p_i$ be a divisor on S^2 such that $-1 < \beta_i < 0$ for each i = 1, ..., n and g_β be a conical metric on S^2 representing the divisor β . We begin with some results concerning the conformal geometry of (S^2, β) .

Define $Iso(S^2, g_\beta) = \{ \phi \in Diff(\Sigma) : \phi^* g_\beta = g_\beta \}$, which forms a group under composition. A diffeomorphism $\psi : S^2 \to S^2$ is called a conformal transformation of (S^2, g_β) if $\psi^*(g_\beta) \in [g_\beta]$, i.e. if there exists a

function $u: S^2 \to \mathbb{R}$ which is smooth and positive and such that

$$\psi^* g_{\beta} = e^{2u} g_{\beta}$$

It is clear that the set of all conformal transformations forms a group under composition. We denote it by $Conf(S^2, g_\beta)$. Observe that if $\psi \in Diff(S^2)$, then ψ^*g_β is always a conical metric with the same number of conical singularities as g_β . Suppose $\phi \in Conf(S^2, g_\beta)$. Then by definition,

$$\phi^* g_\beta = \eta_\phi^2 g_\beta$$

where $\eta_{\phi} = |D\phi| > 0$. Now suppose $g = e^{2u}g_{\beta}$. We have

$$\phi^* g = e^{2(\phi^* u)} \phi^* g_\beta = e^{2(u \circ \phi + \log \eta_\phi)} g_\beta$$

Hence the conformal group $Conf(S^2, g_\beta)$ acts on the conformal factors $u \in C^{k,\alpha}_{\gamma}$ by

$$(3.1) (\phi, u) \to u \circ \phi + \log \eta_{\phi}$$

On the other hand, $Conf(S^2, g_\beta)$ acts on the functions $K \in C^{k-2,\alpha}_{\gamma-2}$ by precomposition, i.e.

$$(3.2) (\phi, K) \to K \circ \phi$$

The group of diffeomorphisms of the sphere S^2 which are $C^{k-2,p}$ is denoted by $Diff^{k-2,p}$ and acts on the curvature functions again by precomposition, as in (3.2).

Theorem 3.1. Suppose $\beta = \sum_{i=1}^{n} \beta_i p_i$ is a divisor on S^2 and g_{+1} is the round metric of curvature 1 on S^2 . If $n \ge 3$ and g_{β} is a conical metric on (S^2, g_{+1}) representing the divisor β , then $Conf(S^2, g_{\beta})$ is finite. Moreover, if there exists i, j, k distinct, such that $\beta_i, \beta_j, \beta_k$ are all distinct, then the conformal group $Conf(S^2, g_{\beta}) = 0$.

Proof. To begin, observe that every conformal transformation $\psi \in Conf(S^2 - \{p_1, p_2, \dots p_n\}, g_{+1})$ has an extension $\tilde{\psi} \in Conf(S^2, g_{+1})$. This follows from the fact that any conformal map $\psi \in Conf(S^2 - \{p_1, p_2, \dots p_n\}, g_{+1})$ can be viewed as a biholomorphism $\mathbb{C} - \{q_1, \dots, q_{n-1}\} \to \mathbb{C} - \{q_1, \dots, q_{n-1}\}$ after conjugation with the stereographic projection from, say, p_n .

To be precise, let σ_{p_n} be the sterographic projection $S^2 - \{p_n\} \to \mathbb{C}$ and suppose $q_i = \sigma_{p_n}(p_i)$ for $i = 1, \ldots, n-1$. The restriction of σ_{p_n} to the punctured sphere $S^2 - \{p_1, \ldots, p_n\}$ gives a diffeomorphism with $\mathbb{C} - \{q_1, \ldots, q_{n-1}\}$. In particular, if $\psi \in Conf(S^2 - \{p_1, \ldots, p_n\}, g_{+1})$, then

$$\bar{\psi} := \sigma_{p_n}^{-1} \circ \psi \circ \sigma_{p_n} : \mathbb{C} - \{q_1, \dots q_{n-1}\} \to \mathbb{C} - \{q_1, \dots q_{n-1}\}$$

is a biholomorphism. We claim now that the points q_i are removable singularities for $\bar{\psi}$. Indeed, if $\bar{\psi}$ had an essentially singularity at any of the points q_i , then by Picard's Theorem, $\bar{\psi}$ would take on all possible values with at most one exception on any neighborhood of q_i , infinitely often, but this would contradict injectivity. The second observation is that ψ has at worst one pole of order one. Again, any higher order pole is excluded because of injectivity. If there were two simple poles at say q_i, q_j , then $\bar{\psi}(q_i) = \infty$, $\bar{\psi}(q_j) = \infty$, but since ψ is an open map, it would map a punctured neighborhood of q_i to a neighborhood of ∞ and a punctured neighborhood of q_j to a neighborhood of ∞ . The fact that these two neighborhoods must intersect contradicts injectivity once again.

It then follows that $\bar{\psi}$ extends to a biholomorphism of the Riemann sphere, hence it corresponds to a conformal map $\tilde{\psi}$ of $S^2(1)$.

Now, observe that if $p_1, p_2 \dots, p_n$ are cone points for the conical metric g_{β} , then

$$Conf(S^2 - \{p_1, \dots, p_n\}, g_\beta) = Conf(S^2 - \{p_1, \dots, p_n\}, g_{+1})$$

This follows from the fact that g_{β} is conformal to g_{+1} by definition.

Finally, we conclude that the group $Conf(S^2 - \{p_1, \ldots, p_n\}, g_{+1})$ is finite if $n \ge 3$. Indeed, if $\psi \in Conf(S^2 - \{p_1, \ldots, p_n\}, g_{+1})$, then its extension $\tilde{\psi}$ to a conformal map of the round sphere fixes the set $\{p_1, \ldots, p_n\}$. Suppose $\tilde{\psi}(p_1) = p_i, \tilde{\psi}(p_2) = p_j, \tilde{\psi}(p_3) = p_k$, where $i, j, k \in (1, \ldots, n)$ are all distinct. Since $\tilde{\psi}$ is a Mobius transformation, its values are uniquely determined after specifying the image of the points

 p_1, p_2, p_3 . Since there are only finitely many choices for p_i, p_j, p_k , the collection of extensions of $\psi \in$ $Conf(S^2 - \{p_1, \dots, p_n\}, g_{+1})$ is finite. In particular, the group $Conf(S^2 - \{p_1, \dots, p_n\}, g_{+1})$ is finite.

Now, let $supp(\beta) = \{p_1, \dots p_n\}$. Define $F : Conf(S^2, \beta, g) \to Conf(S^2 - supp(\beta), g)$ by $F(\phi) =$ $\phi|_{S^2-supp(\beta)}$. Since any $\phi \in Conf(S^2,\beta,g)$ fixes the set $supp(\beta)$, we have that F is a surjective group homomorphism. Moreover,

$$ker(F) = \{ \phi \in Conf(S^2, \beta, g) : F(\phi) = \phi|_{S^2 - supp(\beta)} = Id|_{S^2 - supp(\beta)} \}$$

The only freedom is in where the points $p_1, \dots p_n$ are sent, and we know ϕ fixes them on (S^2, β, g) . Hence we have ker(F) is isomorphic to a subgroup of S_n , the symmetric group on n elements. Finally, since $Conf(S^2 - supp(\beta), g)$ is finite for $n \ge 3$, and ker(F) is also finite, we must have $Conf(S^2, \beta, g)$ is finite. If there are three distinct angles, any conformal map has to fix them, but every conformal map of the unit disk fixing three points is the identity. So the conformal group must be trivial in this case. This concludes the proof of the theorem.

Remark. The condition that $n \ge 3$ is only sufficient. In the examples below we show that there is a metric on $S^2(1)$ with two conical singularities and noncompact conformal group.

3.1. **Examples.** Example 1: The football. Let $\Sigma = S^2(1)/\mathbb{Z}_k$ where \mathbb{Z}_k acts by rotations. The quotient is known as the American football and it is a topological sphere with two conical singularities, each of angle $\frac{2\pi}{k}$. If \bar{g} is the induced metric on the quotient, i.e. $\pi^*\bar{g}=g_{+1}$, where $\pi:S^2\to S^2/\mathbb{Z}_k$ is the quotient map, then $Conf(S^2/\mathbb{Z}_k,\bar{g})$ is noncompact. Indeed, let $\phi_{\lambda}:S^2\to S^2$ be $\phi_{\lambda}(x)=\sigma_p^{-1}\circ\delta_{\lambda}\circ\sigma_p$ where $\delta_{\lambda}(x)=\lambda x$ $(\lambda > 0)$ is a dilation of \mathbb{R}^3 and σ_p is the stereographic projection from p (the composition is extended to the whole sphere by sending $p \to p$). The action of \mathbb{Z}_k on S^2 can be viewed as an action of \mathbb{Z}_k on \mathbb{C} after identifying $S^2 - p$ with the complex plane via the stereographic projection. From this point of view, for each element $[m] \in \mathbb{Z}_k$ we get a map $\psi_m(z) = \zeta^m \cdot z$, where ζ is a kth root of unity. Then one can check that

$$\phi_{\lambda} \circ \psi_m = \psi_m \circ \phi_{\lambda}$$

for every $[m] \in \mathbb{Z}_k$. In particular, the map ϕ_{λ} descends to the quotient and it will be a conformal map of $(S^2/\mathbb{Z}_k,\bar{g}).$

Example 2. Variation on the Football. Another way to obtain the American football of Example 1 is by cutting out two neighborhoods of say, the north and south pole and gluing back two different cones, e.g we can replace a neighborhood U_1 of the north pole by a quotient of the disk D/\mathbb{Z}_n and a neighborhood U_2 of the south pole by the quotient D/\mathbb{Z}_m . When n=m, upon choosing appropriate gluing maps and metrics on the cone pieces, this space can be realized as the quotient S^2/\mathbb{Z}_n of Example 1. In the case when $n \neq m$, there is no metric of constant curvature with conical singularities at the poles ([Tro91]).

Example 3. Double of a Spherical Triangle. Let T be a spherical triangle in $S^2(1)$ with angles $\alpha = \frac{2\pi}{n}, \beta = \frac{2\pi}{m}, \gamma = \frac{2\pi}{p}$. Construct the double of T by identifying T with itself via the identity. The resulting space M is a topological sphere with 3 conical singularities of angles 2α , 2β , 2γ . The conformal group is the dihedral group D_6 .

3.2. **Prescribing Gaussian Curvature on the 2-sphere.** For $\gamma = (\gamma_1, \dots \gamma_n)$, let $C_{\gamma}^{k,\alpha}$ be the Banach space of $C_{\gamma}^{k,\alpha}$ functions $u: S^2 \to \mathbb{R}$ considered as conformal factors of $g = e^{2u}g_{\beta}$ and let $C_{\gamma-2}^{k-2,\alpha}$ be the Banach space of $C_{\gamma-2}^{k-2,\alpha}$ functions K. Our main goal is to study the image of the *curvature map* π , defined to be the map $C_{\gamma}^{k,\alpha} \to C_{\gamma-2}^{k-2,\alpha}$ sending

$$u \mapsto K_{c}$$

As before, if g is a conformal conical metric on (S^2, g_{+1}) , then

$$g = e^{2u} g_{\beta} = e^{2u} \rho^{2\beta} g_{+1}$$

where ρ is a radius function as in Definition 2.2. The Gaussian curvature of g is then

$$K_g = K(e^{2u}g_\beta) = e^{-2u}(K_\beta - \Delta_\beta u)$$

where Δ_{β} is the Laplacian with respect to the conical metric g_{β} and K_{β} the Gaussian curvature of g_{β} . One can compute

$$\Delta_{\beta} = \rho^{-2\beta} \Delta_{+1}$$

$$(3.4) K_{\beta} = \rho^{-2\beta} (1 - \beta \Delta_{+1} \log \rho)$$

Observe that the function $\beta \Delta_{+1} \log \rho$ is defined to be $\beta_i \Delta_{+1} \rho$ in a neighborhood of the cone point p_i and vanishes identically away from the cone points (since $\rho \equiv 1$).

Recall that a C^1 map $F: \mathcal{B}_1 \to \mathcal{B}_2$ is a Fredholm map between Banach manifolds \mathcal{B}_i if the differential

$$D_{u}F(h) := \frac{d}{dt}F(u+th)\bigg|_{t=0} = \lim_{t\to 0} \frac{F(u+th) - F(u)}{t}$$

is a Fredholm operator at each $u \in \mathcal{B}_i$. As it is well-known, Fredholm maps between Banach spaces are bounded linear operators characterized by having finite-dimensional kernel and cokernel. The index of a Fredholm operator is defined as

$$ind(F) = dim(KerF) - dim(coKerF)$$

Moreover, the index of the Fredholm map F is defined to be the index of its differential, which is independent of the choice of u. For more on the theory of Fredholm maps on Banach manifolds, see [ET70, Nir01].

Theorem 3.2. Let (Σ, g, β) be a conical surface with g representing the divisor $\beta = \sum_{j=1}^{n} \beta_{j} p_{j}$. If $\gamma = (\gamma_{1}, \ldots, \gamma_{n}) \in \mathbb{R}^{n}$ satisfies $\gamma_{i} > 0$ and $\gamma_{i} \neq \frac{m}{\beta_{j}}$ for any $i, j \in (1, \ldots, n)$, where m is an integer, then the curvature map π is Fredholm of index 0.

Proof. Fix $u \in C_{\gamma}^{k,\alpha}$. If we set $g = e^{2u}g_{\beta}$, then as observed above

(3.5)
$$\pi(u) = e^{-2u} (K_{\beta} - \Delta_{\beta} u)$$

Let $h \in C_{\gamma}^{k,\alpha}$, thought of as the tangent space to $C_{\gamma}^{k,\alpha}$ at u. Then

$$D_{u}\pi(h) = \frac{d}{dt}\Big|_{t=0}\pi(u+th)$$

$$= \frac{d}{dt}\left(e^{-2(u+th)}(K_{\beta} - \Delta_{\beta}(u+th))\right)\Big|_{t=0}$$

$$= -2hK_{\sigma} - e^{-2u}\Delta_{\beta}h$$

with $K_g = e^{-2u}(K_\beta - \Delta_\beta u)$, which is the Gaussian curvature of g. At u = 0, we get $K_g = K_\beta$, hence

$$(3.6) D_0\pi(h) = -2hK_\beta - \Delta_\beta h$$

$$= -\rho^{-2\beta} (2h(1 - \beta \Delta_{+1} \log \rho)) + \Delta_{+1}h)$$

If we let $a := 2(1 - \beta \Delta_{+1} \log \rho)$, then

(3.8)
$$-L_{\beta}(h) := D_0 \pi(h) = \rho^{-2\beta} (ah + \Delta_{+1}h)$$

It is known that if $\gamma = (\gamma_1, \dots, \gamma_n) \in \mathbb{R}^n$ satisfies $\gamma_i \neq \frac{m}{\beta_j}$ for any $i, j \in (1, \dots, n)$, where m is an integer, then the linear operator $L_\beta : C_\gamma^{k,p} \to C_{\gamma-2}^{k-2,p}$ is Fredholm [MW15, Beh11]. We further claim L_β is formally self-adjoint. Observe there is a natural inner product on (S^2, β) given by

(3.9)
$$\langle u, v \rangle = \int_{S^2} u \cdot v \rho^{2\beta} dV_{+1}$$

Thus for all $u, v \in C_{\gamma}^{k,\alpha}$ we have

(3.10)
$$\langle v, L_{\beta} u \rangle = \int_{S^2} v(\rho^{-2\beta}(\Delta_{+1}u + au))\rho^{2\beta} dV_{+1}$$

$$= \int_{S^2} v(\Delta_{+1}u + au)dV_{+1}$$

(3.12)
$$= \int_{S^2} v(\Delta_{+1}u + au)dV_{+1}$$

$$= \int_{S^2} u(\Delta_{+1}v + av)dV_{+1}$$

(3.14)
$$= \int_{S^2} u \rho^{-2\beta} (\Delta_{+1} v + a v) \rho^{2\beta} dV_{+1}$$

$$(3.15) = \langle L_{\beta} v, u \rangle$$

The integration by parts in (3.12)-(3.13) needs some justification, so fix R > 0 small enough and let $B_R(p_i)$ be a geodesic ball of radius R around the cone point p_i . Let $S_R(p_i)$ denote the circle of radius R with center p_i , $\partial_\nu u$ denotes the normal derivative of u, and dS the volume element of $S_R(p_i)$. We then have

$$\int_{S^{2}-\{p_{1}...p_{n}\}} v\Delta_{+1}u \quad dV_{+1} = \lim_{R \to 0} \int_{S^{2}-\bigcup_{i=1}^{n} B_{R}(p_{i})} v\Delta_{+1}u \quad dV_{+1}
= \lim_{R \to 0} \left(\int_{S^{2}-\bigcup_{i=1}^{n} B_{R}(p_{i})} u\Delta_{+1}v \quad dV_{+1} + \int_{\bigcup_{i=1}^{n} S_{R}(p_{i})} u\partial_{\nu}v - v\partial_{\nu}u \quad dS \right)
= \int_{S^{2}-\{p_{1},...p_{n}\}} u\Delta_{+1}v \quad dV_{+1} + \lim_{R \to 0} \sum_{i=1}^{n} \int_{S_{R}(p_{i})} u\partial_{\nu}v - v\partial_{\nu}u \quad dS$$

Now, since $u \in C^{k,p}_{\gamma}(S^2 - \bigcup_{i=1}^n B_R(p_i))$ for all R > 0 small enough, it follows that $\partial_{\gamma} u \in C^{k-1,p}_{\gamma-1}(S^2 - \bigcup_{i=1}^n B_R(p_i))$, i.e. there is a C > 0 such that $\|u\|_{C^{l,\alpha}_{\gamma}} \leq C$. In particular, $\|\partial_{\gamma} u\|_{C^{l-1,\alpha}_{\gamma-1}} \leq C$. It follows from the definition of these norms (see 2.14) that

$$\sup_{x \in S^2 - \{p_1, \dots p_n\}} \rho^{-\gamma} |u| \le C$$

and

$$\sup_{x \in S^2 - \{p_1, \dots p_n\}} \rho^{-(\gamma - 1)} |\partial_{\nu} u| \le C_1$$

and similarly for v. Therefore

$$(3.16) \qquad \left| \int_{S_{R}(p_{i})} u \partial_{\nu} v - v \partial_{\nu} u \right| dS \leq \int_{S_{R}(p_{i})} |u| |\partial_{\nu} v| dS + \int_{S_{R}(p_{i})} |v| |\partial_{\nu} u| dS$$

$$= \int_{S_{R}(p_{i})} \rho^{-\gamma} |u| \rho^{-\gamma+1} |\partial_{\nu} v| \rho^{2\gamma-1} dS + \int_{S_{R}(p_{i})} \rho^{-\gamma} |v| \rho^{-\gamma+1} |\partial_{\nu} u| \rho^{2\gamma-1} dS$$

$$\leq 2C' \int_{S_{R}(p_{i})} \rho^{2\gamma-1} dS$$

$$(3.18) \leq 2C' \int_{S_R(p_i)} \rho^{2\gamma - 1} dS$$

$$(3.19) \leq C(\delta)R^{2\gamma}$$

Thus, provided $\gamma > 0$, taking the limit as $R \to 0$, we see that the boundary terms disappear, as wanted. This concludes the proof that the map L_{β} is formally self-adjoint and Fredholm, from which it follows that the map π is a Fredholm map of index 0.

3.2.1. Properness of the map π .

Theorem 3.3. Let C_+ be the subspace of $C_{\gamma-2}^{k-2,\alpha}$ consisting of positive curvature functions K. Define $\mathcal{U}=\pi^{-1}(C_+)$. If $\gamma>0$, then the map $\pi_0:\mathcal{U}\to C_+$ defined as the restriction of π to \mathcal{U} is proper.

Proof. If $K_i \to K \in C_+$, then the sequence K_i is bounded in $C_{\gamma-2}^{k-2,\alpha}$. In particular, K_i is bounded in $C_{\gamma-2}^0$. Hence there exist a constant K such that $||K_i||_0 \le K$ away from the cone points. Moreover, since the Euler characteristic is positive, we have using Gauss-Bonnet (1.2)

$$(3.20) 2\pi\chi(S^2,\beta) = \int_{S^2} K_i dvol_{g_i} \le K_0 \cdot area(S^2,g_i)$$

so we get

$$area(S^2, g_i) \ge \frac{2\pi\chi(S^2, \beta)}{K_0} > 0$$

On the other hand, Myers' theorem ([Pet16]) implies that the diameter of each conical surface is finite (since curvature is assumed positive). Furthermore, since the sequence K_i converges in C_+ to a positive function K, we can find constants D_0 , v_0 such that

- (1) $vol(g_i) \ge v_0$
- (2) $diam(g_i) \leq D_0$

for all i. Under these bounds we can now directly apply Theorem 2.9 to conclude that there exists a sequence of diffeomorphisms $F_i: S^2 \to S^2$ such that

$$(3.21) (F_i^* g_i)_{rs} \to (g_\infty)_{rs}$$

in $C_{\gamma}^{1,\alpha}$, where g_{∞} is a conical metric on S^2 with m conical singularities of angles $0 < \theta \le 2\pi$. By passing to a subsequence if necessary, we may assume that the F_i are orientation preserving. On the other hand, since g_{∞} is the limit of a sequence of conical metrics in the same conformal class, we claim there exists a diffeomorphism $\psi: S^2 \to S^2$ and a smooth, positive function $u: S^2 \to \mathbb{R}$ such that $\psi^*g = e^{2u}g_{\beta}$. To see this, suppose \mathfrak{C} denotes the space of conformal classes on the punctured sphere $S^2 - \{p_1, \ldots, p_n\}$, i.e., two smooth (incomplete) metrics h_1, h_2 on $S^2 - \{p_1, \ldots, p_n\}$ represent the same point in \mathfrak{C} if there exists a positive smooth function u such that $h_1 = e^{2u}h_2$. The group $Diff_+$ of orientation preserving diffeomorphisms of S^2 acts on \mathfrak{C} by

$$(\psi, [h]) \rightarrow \psi^*[h] = [\psi^*h]$$

Since it is not true in general that $[\psi^*h] = [h]$, we consider the moduli space $\mathcal{M} = \mathfrak{C}/Diff_+$. This space corresponds to Teichmuller space [Pet19, MW15], since the mapping class group of the sphere is trivial (so every orientation preserving diffeomorphism is isotopic to the identity). Observe then that every element of the sequence g_i , when considered as smooth metrics on $S^2 - \{p_1, \dots p_n\}$, corresponds to the same point in \mathfrak{C} , namely the conformal class $[g_\beta]$. If ψ_i is a sequence of orientation preserving diffeomorphisms, then $\psi_i^*[g_i] = [g_i]$ as elements of \mathcal{M} , hence the classes $\psi_i^*g_i$ and g_β correspond to the same point in \mathcal{M} . Since the topology of Teichmuller space is Hausdorff [Pet19], the sequence $\psi_i^*g_i$ is the constant sequence $[g_\beta] \in \mathcal{M}$. Thus the limit of the $\psi_i^*g_i$ must be in the conformal class of g_β modulo diffeomorphisms, i.e. there exists a diffeomorphism of S^2 such that the limit g satisfies

$$\psi^*g=e^{2u}g_\beta$$

for some positive smooth function u on S^2 , as claimed. In particular, there is a diffeomorphism Ψ of S^2 such that $\Psi^*g_{\infty} = e^{2u}g_{\beta}$. After precomposing F_i with Ψ^{-1} , we may as well assume that we have a sequence F_i of diffeomorphisms of S^2 such that

(3.22)
$$F_i^*(g_i) = F_i^*(e^{2u_i}g_{\beta}) \to e^{2u}g_{\beta}$$

in $C_{\gamma}^{1,\alpha}$, where u is some positive smooth function on S^2 . As in the arguments preceding Proposition 2.5 in [And17], one now has that the F_i converge on a subsequence to the identity modulo the action of the

conformal group, i.e. there exist conformal maps $\phi_i \in Conf(S^2, g_\beta)$ such that $\phi_i^{-1} \circ F_i$ converge to the identity. It follows from Proposition 2.5 in [And17] that the functions $\phi_i^* u_i$ are uniformly bounded in $C_\gamma^{1,\alpha} \cap W_\delta^{2,p}$, for $\gamma > 0$ and some $\delta \leq \gamma$ (see [Beh11] for a definition of weighted Sobolev spaces). Observe that nothing really changes in the presence of conical singularities since the diffeomorphisms F_i are still quasiconformal when restricted to the punctured sphere $S^2 - \{p_1, \ldots, p_n\}$ (see [LV73]). By the Arzela-Ascoli theorem, the uniform bound on the sequence $\phi_i^* u_i$ implies convergence on a subsequence to a limit in $C_\gamma^{1,\alpha}$. Moreover, since the conformal group $Conf(S^2, g_\beta)$ is finite, we actually have that $\{u_i\}$ themselves (sub)converge to a limit $u \in C_\gamma^{1,\alpha}$ which satisfies

$$\Delta_{g_{\beta}}u = K_{\beta} - Ke^{2u}$$

weakly. Given that the Gaussian curvatures K_i of the metrics g_i are assumed to be in $C_{\gamma-2}^{k-2,\alpha}$, a bootstrapping argument using Proposition 2.7 in [Beh11] implies $u \in C_{\gamma}^{k,\alpha}$. This then completes the proof that the map π_0 is proper.

3.2.2. Degree Computations. We conclude this section with a result providing sufficient conditions for a function K to arise as the Gaussian curvature of a conformal conical metric on S^2 having at least three conical singularities and angles less than 2π . As mentioned in the introduction, a necessary condition for the existence of a *constant* curvature conical metric on S^2 having at least three conical singularities and angles less than 2π is

(3.24)
$$\sum_{i \neq i} \beta_i < \beta_j, \text{ for all } j$$

In fact, Luo-Tian have shown in [LT92] that if the generalized Euler characteristic is positive, then this condition is sufficient and necessary for uniqueness and existence. Under these assumptions, we can now compute the degree of the curvature map given our previous results. Recall that if F is a proper Fredholm map of index 0 between open subsets of Banach spaces, one can define its *degree* by the formula

$$deg(F) = \sum_{x \in F^{-1}(y)} sign(D_x F)$$

where y is any regular value of F and the sign is \pm according to whether $D_x F$ preserves or reverses orientation. By definition, y is a regular value if $D_x F$ is an isomorphism for all $x \in F^{-1}(y)$. In particular, points with empty preimage are always regular values. We refer the reader to [Nir01] for more on the degree theory of Fredholm maps on Banach manifolds.

of Fredholm maps on Banach manifolds. Let $C = C_+ \cap C_{\gamma-2}^{k-2,\alpha}$, where $\gamma_i > 0$, $\gamma_i \neq \frac{m}{\beta_j}$ for any $(i,j) \in (1,\dots n)$. Recall that the restrictions on γ guarantee that the curvature map is proper and Fredholm of index 0. We have,

Theorem 3.4. Suppose $n \ge 3$, and $\beta = \sum_{i=1}^{n} \beta_i p_i$ is a divisor on S^2 satisfying the Troyanov condition (3.24) and there exists i, j, k distinct for which $\beta_i, \beta_j, \beta_k$ are all distinct. Assume $\chi(S^2, \beta) > 0$ and let g_β be the unique conical metric on S^2 representing the divisor β of Gaussian curvature $K_\beta = 1$. Then a function K on S^2 is the Gaussian curvature of a metric g conformal to g_β if and only if $K \in C$.

Proof. Suppose $K \in C$. We want to show there exists a function u such that $e^{2u}g_{\beta}$ has Gaussian curvature K, where g_{β} is the unique conformal conical metric with Gaussian curvature 1. The existence of such a metric is equivalent to the existence of a solution u to the equation

(3.25)
$$K = e^{-2u}(1 - \Delta_{\beta}u)$$

In the language of this section, it is enough to show that the restriction π_0 of the curvature map to $\pi^{-1}(C)$ has deg = 1. The assumption that $K \in C$ guarantees that the map π_0 is a proper Fredholm map of index 0 (see Theorems 3.3, 3.2). Observe that for given γ , α , k satisfying the conditions of the theorem, the subset of

 $C_{\gamma-2}^{k-2,\alpha}$ consisting of positive functions is convex, thus C is path-connected and there is a well-defined notion of degree. Clearly the function $K=1 \in C$. On the other hand, the preimage of K=1 under π_0 is given by all solutions to the equation

$$(3.26) 1 = e^{-2u}(1 - \Delta_{\beta}u)$$

By Theorem 2 in [LT92] there exists a unique conical metric g on S^2 representing the divisor β of constant curvature 1. Since u = 0 is a solution, it follows that the preimage $\pi_0^{-1}(1) = \{0\}$.

Now, the kernel of the differential of the curvature map π_0 under the assumption that g_{β} has Gaussian curvature 1 is given by solutions of

(3.27)
$$D_0 \pi(h) = -2h - \Delta_{\beta} h = 0$$

We now argue that the first eigenvalue λ of the problem

$$\Delta_{\beta}h = -\lambda h$$

satisfies $\lambda \ge 2$. Moreover, if the lowest possible eigenvalue is achieved, namely $\lambda = 2$, then there exists a non-constant solution to the equation Hess(f) = -fg. To see this, we follow the same ideas as in the works of Lichnerowicz and Obata [Lic58, Oba62] which have now become standard. Using Bochner's formula away from the cone points, we can write

(3.29)
$$\frac{1}{2}\Delta_{\beta}|\nabla h|^2 = |Hess(h)|^2 + g_{\beta}(\nabla \Delta h, \nabla h) + K_{\beta}|\nabla h|^2$$

Using Scharwz inequality and the fact that h is an eigenfunction we get

$$(3.30) |Hess(h)|^2 \ge \frac{1}{2}(\Delta h)^2 = -\frac{\lambda}{2}h\Delta_{\beta}h$$

Combining this with Bochner's formula we get the inequality

(3.31)
$$\Delta_{\beta} |\nabla h|^2 \ge -\frac{\lambda}{2} h \Delta_{\beta} h - \lambda |\nabla h|^2 + |\nabla h|^2$$

We now claim

(3.32)
$$\int_{S^2 - \{p_1, \dots p_n\}} \Delta_{\beta} f dvol_{\beta} = 0$$

holds for any function $f \in C^{k-1,\alpha}_{\gamma-1}$. For R > 0 small enough, let $B_R(p_k)$ be a geodesic ball of radius R centered at the cone point p_k . Then

$$\begin{split} \int_{S^{2}-\{p_{1},\dots p_{n}\}} \Delta_{\beta} f dvol_{\beta} &= \int_{S^{2}-\{p_{1},\dots p_{n}\}} \rho^{-2\beta} \Delta_{+1} f \rho^{2\beta} dvol_{+1} = \lim_{R \to 0} \int_{S^{2}-\cup_{k=1}^{n} B_{R}(p_{k})} \Delta_{+1} f dvol_{+1} \\ &= \lim_{R \to 0} \int_{S^{2}-\cup_{k=1}^{n} B_{R}(p_{k})} div(\nabla f) dvol_{+1} \\ &= \lim_{R \to 0} \sum_{k=1}^{n} \int_{S_{R}(p_{k})} (\nabla f \cdot v) dS \end{split}$$

where in the last equality we have used the divergence theorem, with S_R denoting the boundary of $B_R(p_k)$, ν the normal to each boundary circle and dS the area element. As before, the assumption that $f \in C^{k-1,\alpha}_{\gamma-1}$ implies that $\sup_{S^2-\{p_1,\dots,p_n\}} \rho^{-(\gamma-1)+1} |\nabla f| \leq C$, so that

$$\int_{S_R} |\rho^{-\gamma+2} \nabla f \cdot \nu| \rho^{\gamma-2} dS \le C' R^{\gamma}$$

Since $\gamma > 0$, taking the limit as $R \to 0$ we get the desired result in (3.32).

Now, using (3.32) in combination with (3.31), we obtain the inequality

$$0 = \int_{S^2 - \{p_1, \dots p_n\}} \Delta_{\beta} |\nabla h|^2 dvol_{\beta} \ge \int_{S^2 - \{p_1, \dots p_n\}} -\frac{\lambda}{2} h \Delta_{\beta} h - \lambda |\nabla h|^2 + |\nabla h|^2 dvol_{\beta}$$
$$= \left(\frac{\lambda}{2} - \lambda + 1\right) \left(\int_{S^2 - \{p_1, \dots p_n\}} |\nabla h|^2 dvol_{\beta}\right)$$

where the integration by parts is justified as in the proof of Theorem 3.2. The previous inequality then shows that

$$(3.33) \qquad \qquad -\frac{\lambda}{2} + 1 \le 0$$

so that $\lambda \ge 2$. Moreover, if $\lambda = 2$, then the inequalities become equality, which forces the trace free part of the Hessian of h to vanish, implying that h solves the equation

$$(3.34) Hess(h) = \phi g$$

One can further show that $\phi = -h$. Observe that (3.34) implies the existence of a nonconstant solution to the equation

$$\mathcal{L}_{\nabla h}g = -hg$$

Now suppose $h \in Ker(D_0\pi_0)$, that is, h solves $-2h = \Delta_\beta h = \rho^{-2\beta}\Delta_{+1}h$, thus h is an eigenfuction corresponding to the lowest possible eigenvalue. The discussion above shows that h satisfies (3.35), in other words, ∇h is a conformal Killing field on (S^2, g, β) . Equivalently, this means the locally defined flow of ∇h preserves the conformal structure. Therefore, there exists a nontrivial one-parameter group of conformal transformations. Since the conformal group $Conf(S^2, \beta)$ is trivial, we must have h = 0. Thus K = 1 is a regular value of the curvature map π_0 . It now follows that $deg\pi_0 = 1$, as wanted.

References

- [AC92] Michael T. Anderson and Jeff Cheeger. " C^{α} -compactness for manifolds with Ricci curvature and injectivity radius bounded below." J. Differential Geom., 35(2):265–281, 1992.
- [And90a] Michael T. Anderson. "Convergence and rigidity of manifolds under Ricci curvature bounds." *Invent. Math.*, 102(2):429–445, 1990.
- [And90b] Michael T. Anderson. "On the topology of complete manifolds of nonnegative Ricci curvature." *Topology*, 29(1):41–55, 1990.
- [And17] Michael T. Anderson. "The Nirenberg problem of prescribed gauss curvature on S²." Preprint arXiv: 1707.02938, 2017
- [Ast94] Kari Astala. "Area distortion of quasiconformal mappings." Acta Math., 173(1):37–60, 1994.
- [Beh11] Tapio Behrndt. "On the cauchy problem for the heat equation on riemannian manifolds with conical singularities." *The Ouarterly Journal of Mathematics*, 64, 05 2011.
- [BF42] Ch. Blanc and F. Fiala. "Le type d'une surface et sa courbure totale." Comment. Math. Helv., 14:230–233, 1942.
- [CGY93] Sun-Yung A. Chang, Matthew J. Gursky, and Paul C. Yang. "The scalar curvature equation on 2- and 3-spheres." *Calc. Var. Partial Differential Equations*, 1(2):205–229, 1993.
- [Che98] Xiuxiong Chen. "Weak limits of Riemannian metrics in surfaces with integral curvature bound." *Calc. Var. Partial Differential Equations*, 6(3):189–226, 1998.
- [CL93] Kung Ching Chang and Jia Quan Liu. "On Nirenberg's problem." *Internat. J. Math.*, 4(1):35–58, 1993.
- [CM12] Alessandro Carlotto and Andrea Malchiodi. "Weighted barycentric sets and singular Liouville equations on compact surfaces." J. Funct. Anal., 262(2):409–450, 2012.
- [CY87] Sun-Yung Alice Chang and Paul C. Yang. "Prescribing Gaussian curvature on S²." Acta Math., 159(3-4):215–259, 1987.
- [Deb20] Clément Debin. "A compactness theorem for surfaces with bounded curvature." *J. Inst. Math. Jussieu*, 19(2):597–645, 2020.
- [DL95] Wei Yue Ding and Jia Quan Liu. "A note on the problem of prescribing Gaussian curvature on surfaces." *Trans. Amer. Math. Soc.*, 347(3):1059–1066, 1995.
- [dMLS15] Francesca de Marchis and Rafael López-Soriano. "Existence and non existence results for the singular nirenberg problem." Preprint: arXiv 1507.08090, 2015

- [Ere04] A. Eremenko. "Metrics of positive curvature with conic singularities on the sphere." *Proc. Amer. Math. Soc.*, 132(11):3349–3355, 2004.
- [ET70] K. D. Elworthy and A. J. Tromba. "Degree theory on Banach manifolds." In Nonlinear Functional Analysis (Proc. Sympos. Pure Math., Vol. XVIII, Part 1, Chicago, Ill., 1968), pages 86–94. Amer. Math. Soc., Providence, R.I., 1970.
- [GT83] D. Gilbarg and N. Trudinger. Elliptic Partial Differential Equations of Second Order, 2nd Edition. Grundlehren Series, vol. 224, Springer Verlag, Berlin, 1983.
- [Han90] Zheng-Chao Han. "Prescribing Gaussian curvature on S²." Duke Math. J., 61(3):679–703, 1990.
- [HH97] E. Hebey and M. Herzlich. "Harmonic coordinates, harmonic radius and convergence of Riemannian manifolds." *Rend. Mat. Appl.* (7), 17(4):569–605 (1998), 1997.
- [Ji04] Min Ji. "On positive scalar curvature on S²." Calc. Var. Partial Differential Equations, 19(2):165–182, 2004.
- [KW75] Jerry L. Kazdan and F. W. Warner. "Existence and conformal deformation of metrics with prescribed Gaussian and scalar curvatures." *Ann. of Math.* (2), 101:317–331, 1975.
- [Lic58] André Lichnerowicz. Géométrie des groupes de transformations. Travaux et Recherches Mathématiques, III. Dunod, Paris. 1958.
- [LT92] Feng Luo and Gang Tian. "Liouville equation and spherical convex polytopes." *Proc. Amer. Math. Soc.*, 116(4):1119–1129, 1992.
- [LV73] O. Lehto and K. I. Virtanen. Quasiconformal mappings in the plane. Springer-Verlag, New York-Heidelberg, second edition, 1973. Translated from the German by K. W. Lucas, Die Grundlehren der mathematischen Wissenschaften, Band 126.
- [McO88] Robert C. McOwen. "Point singularities and conformal metrics on Riemann surfaces." *Proc. Amer. Math. Soc.*, 103(1):222–224, 1988.
- [MP19] Gabriele Mondello and Dmitri Panov. "Spherical surfaces with conical points: systole inequality and moduli spaces with many connected components." *Geom. Funct. Anal.*, 29(4):1110–1193, 2019.
- [MRS15] Rafe Mazzeo, Yanir A. Rubinstein, and Natasa Sesum. "Ricci flow on surfaces with conic singularities." *Anal. PDE*, 8(4):839–882, 2015.
- [MW15] Rafe Mazzeo and Hartmut Weiss. "Teichmller theory for conic surfaces" Preprint arXiv: 1509.07608, 2015
- [Nir01] Louis Nirenberg. Topics in nonlinear functional analysis, volume 6 of Courant Lecture Notes in Mathematics. New York University, Courant Institute of Mathematical Sciences, New York, 2001. Chapter 6 by E. Zehnder, Notes by R. A. Artino, Revised reprint of the 1974 original.
- [Oba62] Morio Obata. "Certain conditions for a Riemannian manifold to be isometric with a sphere." *J. Math. Soc. Japan*, 14:333–340, 1962.
- [Pac06] Frank Pacard. "Analysis in weighted spaces: preliminary version." *Teheran (Iran)*, pp.75, 3rd cycle,, 2006. cel-00392164.
- [Pet87] Stefan Peters. "Convergence of Riemannian manifolds." Compositio Math., 62(1):3–16, 1987.
- [Pet16] Peter Petersen. *Riemannian geometry*, volume 171 of *Graduate Texts in Mathematics*. Springer, Cham, third edition, 2016.
- [Pet19] Bram Petri. Teichmüller theory, Lecture notes, 2019. http://www.math.uni-bonn.de/people/bpetri/t_tt/ Teich190704.pdf
- [PSSW20] D. H. Phong, Jian Song, Jacob Sturm, and Xiaowei Wang. "The Ricci flow on the sphere with marked points." J. Differential Geom., 114(1):117–170, 2020.
- [Ram18] Daniel Ramos. "Ricci flow on cone surfaces." Port. Math., 75(1):11-65, 2018.
- [Tro91] Marc Troyanov. "Prescribing curvature on compact surfaces with conical singularities." *Trans. Amer. Math. Soc.*, 324(2):793–821, 1991.
- [Yau75] Shing Tung Yau. "Harmonic functions on complete Riemannian manifolds." *Comm. Pure Appl. Math.*, 28:201–228, 1975.

STONY BROOK UNIVERSITY

E-mail address: lisandra@math.stonybrook.edu