

# HOWE DUALITY OF THE SYMMETRIC GROUP AND A MULTISET PARTITION ALGEBRA

ROSA ORELLANA AND MIKE ZABROCKI

ABSTRACT. We introduce the *multiset partition algebra*,  $\text{MP}_{r,k}(x)$ , that has bases elements indexed by multiset partitions, where  $x$  is an indeterminate and  $r$  and  $k$  are non-negative integers. This algebra can be realized as a diagram algebra that generalizes the partition algebra. When  $x$  is an integer greater or equal to  $2r$ , we show that  $\text{MP}_{r,k}(x)$  is isomorphic to a centralizer algebra of the symmetric group,  $S_n$ , acting on the polynomial ring on the variables  $x_{ij}$ ,  $1 \leq i \leq n$  and  $1 \leq j \leq k$ . We describe the representations of  $\text{MP}_{r,k}(x)$ , branching rule and restriction of its representations in the case that  $x$  is an integer greater or equal to  $2r$ .

## CONTENTS

1. Introduction	1
2. Preliminaries and Definitions	4
2.1. Multisets and multiset partitions	4
2.2. Diagrams	5
2.3. Colored multiset partitions	5
2.4. Symmetric Functions	6
3. The generic multiset partition algebra	8
4. $\text{MP}_{r,k}(n)$ is a centralizer algebra of $\mathbb{C}S_n$	11
4.1. An $S_n$ action on polynomials	11
4.2. The centralizer of the action	12
4.3. $\mathcal{A}_{r,k}(n)$ is isomorphic to $\text{MP}_{r,k}(n)$ for $n \geq 2r$	14
5. Irreducible representations of $\mathcal{A}_{r,k}(n)$	16
5.1. The branching rule	20
5.2. Restriction of $W_{\mathcal{A}_{r,k}(n)}^\lambda$	22
References	23

## 1. INTRODUCTION

Let  $V_n$  be an  $n$  dimensional vector space, then Schur-Weyl duality is a fundamental property in representation theory that relates the representations of the general linear

---

Work supported by NSF grants DMS-1300512 and DMS-1700058, and by NSERC.

group  $GL_n(\mathbb{C})$  and the symmetric group algebra  $\mathbb{C}S_k$  as they both act on the tensor space

$$V_n^{\otimes k} = \underbrace{V_n \otimes V_n \otimes \cdots \otimes V_n}_{k \text{ times}} .$$

The duality of these actions implies that

$$(1.1) \quad V_n^{\otimes k} \cong \bigoplus_{\lambda} W_{GL_n(\mathbb{C})}^{\lambda} \otimes W_{\mathbb{C}S_k}^{\lambda}$$

where for a group or an algebra  $A$ , we use the notation  $W_A^{\lambda}$  to represent an irreducible representation of  $A$  and the direct sum is over all partitions  $\lambda$  of  $k$ .

This duality applies to algebras acting on spaces other than  $V_n^{\otimes k}$  which centralize each other. For instance, if we let  $V_{n,k} = (\mathbb{C}^n)^k$  denote the  $nk$  dimensional vector space of sequences  $(v_1, v_2, \dots, v_k)$  where  $v_i \in \mathbb{C}^n$ . There is an action of  $GL_n(\mathbb{C})$  and of  $GL_k(\mathbb{C})$  on  $V_{n,k}$  which mutually commute. Let  $\mathcal{P}(V)$  be the algebra of polynomial functions on  $V$  and

$$\mathcal{P}^r(V) = \{f \in \mathcal{P}(V) : f(zv) = z^r f(v) \text{ for } z \in \mathbb{C}^{\times}\}$$

be the homogeneous polynomials of degree  $r$ . The  $GL_n(\mathbb{C}) \times GL_k(\mathbb{C})$  action extends to  $\mathcal{P}^r(V_{n,k})$  [GW2, Section 5.6.2]. Then the analogous decomposition of  $\mathcal{P}^r(V_{n,k})$  into  $GL_n(\mathbb{C}) \times GL_k(\mathbb{C})$ -modules is

$$(1.2) \quad \mathcal{P}^r(V_{n,k}) \cong \bigoplus_{\lambda} W_{GL_n(\mathbb{C})}^{\lambda} \otimes W_{GL_k(\mathbb{C})}^{\lambda}$$

where the sum is over all partitions  $\lambda$  of  $r$  of length at most  $\min(n, k)$ . It is this duality that is referred to as ‘Howe duality’ [Ho].

The decompositions in the isomorphisms in Equations (1.1) and (1.2) create a correspondence between representations of one algebra (or group) and the representations of the dual algebra (or group). In addition, the multiplicities of irreducible representations of one algebra correspond to dimensions of irreducible representations of the dual algebra.

In the 1990’s, Martin [Mar1] introduced the partition algebra,  $P_k(n)$ , as a generalization of the Temperley-Lieb algebra that could be used to study the transfer matrix of certain statistical mechanics models. In the case that  $n \geq 2k$  the partition algebra is the algebra whose action centralizes the action of  $S_n \subseteq GL_n(\mathbb{C})$  as the subgroup of permutation matrices acting on  $V_n^{\otimes k}$  [Mar4, MR, J].

In this case we also have a decomposition of  $V_n^{\otimes k}$  into  $\mathbb{C}S_n \times P_k(n)$  irreducible representations,

$$(1.3) \quad V_n^{\otimes k} \cong \bigoplus_{\lambda} W_{\mathbb{C}S_n}^{\lambda} \otimes W_{P_k(n)}^{\lambda} .$$

One application of this decomposition is that it helps to establish the relationship between multiplicities occurring in the decomposition of the tensor product of  $S_n$  irreducible representations (i.e. Kronecker coefficients) to the multiplicities occurring in the restriction of  $P_k(n)$  irreducibles to Young subalgebras [BDO]. The multiplicities for tensor products of polynomial representations of  $GL_n$  are known as the Littlewood-Richardson coefficients and they are better understood than the Kronecker coefficients. One goal of the research in this

area has been to develop an understanding of algebras, combinatorics, and the representation theory associated to  $P_k(n)$  to help understand the representation theory of the symmetric group in general [COSSZ, E, Mar2, Mar4, ME, Ha, HJ, HL, HR, BH1, BH2, BHH].

In this paper we introduce a multiset partition algebra  $\text{MP}_{r,k}(x)$  such that, for  $n \geq 2r$ ,  $\text{MP}_{r,k}(n)$  is the centralizer algebra of the symmetric group  $S_n \subseteq GL_n(\mathbb{C})$  when it acts on  $\mathcal{P}^r(V_{n,k})$ . That is, if we denote the irreducible representations of this algebra by  $W_{\text{MP}_{r,k}(n)}^\lambda$ , then as a  $\mathbb{C}S_n \times \text{MP}_{r,k}(n)$  module,

$$(1.4) \quad \mathcal{P}^r(V_{n,k}) \cong \bigoplus_{\lambda} W_{\mathbb{C}S_n}^\lambda \otimes W_{\text{MP}_{r,k}(n)}^\lambda .$$

We begin by introducing the generic multiset partition algebra  $\text{MP}_{r,k}(x)$  that depends on a parameter  $x$  and uses multiset partitions to index basis elements. We show that when  $x = n$ , an integer greater than or equal to  $2r$ ,  $\text{MP}_{r,k}(n)$  is isomorphic to  $\mathcal{A}_{r,k}(n) := \text{End}_{S_n}(\mathcal{P}^r(V_{n,k}))$ , the centralizer algebra of the action of the symmetric group  $S_n$  as the subgroup of permutation matrices inside of  $GL_n(\mathbb{C})$  when it acts on  $\mathcal{P}^r(V_{n,k})$ .

Our hope is that this algebra can provide some insight into some aspects of the representation theory of the symmetric group that are still not well understood. Indeed, we show that the multiplicities when we restrict an irreducible multiset partition algebra representation to sums of Young-type subalgebras are equal to the Kronecker coefficients.

A recent paper by Narayanan, Paul and Srivastava [NPS, Pa] describes the centralizer algebra  $\text{End}_{S_n}(\mathcal{P}^{\alpha_1}(V_n) \otimes \mathcal{P}^{\alpha_2}(V_n) \otimes \cdots \otimes \mathcal{P}^{\alpha_k}(V_n))$  for a fixed weak composition  $\alpha$  of length  $k$ . This should be isomorphic to a subalgebra of the multiset partition algebra introduced in this paper, but their product does not precisely agree with ours and so the relationship is left open.

We did not consider in the definition of this algebra an extension corresponding to the linear transformations in  $\text{Hom}(\mathcal{P}^r(V_{n,k}), \mathcal{P}^s(V_{n,k}))$  where  $r, s$  are non-negative integers which commute with the action of  $S_n$ . This centralizer algebra would depend on 4 parameters,  $r, s, k, n$ , and is spanned by multiset partitions with  $r$  elements from  $\{1, 2, \dots, k\}$  and  $s$  elements from  $\{\bar{1}, \bar{2}, \dots, \bar{k}\}$ . This adds a bit to the complexity of the presentation, but we did not have much more to say about it beyond a formula for the dimension (see the comment in Remark 5.5).

The main results in this paper are:

- (1) A definition of the generic multiset partition algebra  $\text{MP}_{r,k}(x)$  and a proof that the product is associative (Proposition 3.4).
- (2) A definition of the centralizer algebra  $\mathcal{A}_{r,k}(n)$  and a proof that  $\text{MP}_{r,k}(n) \cong \mathcal{A}_{r,k}(n)$  when  $n \geq 2r$  (Theorem 4.7).
- (3) A generating function formula for the dimension of  $\mathcal{A}_{r,k}(n)$  (Corollary 5.4).
- (4) A generating function formula for the dimension of an irreducible  $\mathcal{A}_{r,k}(n)$  indexed by a partition  $\lambda$  of  $n$  (Proposition 5.2).
- (5) A formula for the branching coefficient of an irreducible  $\mathcal{A}_{r,k}(n)$  to an  $\mathcal{A}_{r-d,k-1}(n)$ -module (Theorem 5.6).

- (6) A proof that the multiplicities of an irreducible  $\mathcal{A}_{r,k+\ell}(n)$ -module restricted to a direct sum of  $\mathcal{A}_{d,k}(n) \otimes \mathcal{A}_{r-d,\ell}(n)$ -modules are the Kronecker coefficients (Theorem 5.10).

In Section 2 of this paper, we begin by introducing some of the combinatorial notation of multisets, multiset partitions, diagrams and colored multiset partitions and diagrams that we will need to encode the basis elements and their product. Section 3 presents the definitions of the generic form of the multiset partition algebra. Then in Section 4 we show that the generic multiset partition algebra is isomorphic to the centralizer algebra  $\mathcal{A}_{r,k}(n)$  when  $x = n \geq 2r$ . In Section 5 we give some enumerative results and formulae for the dimensions of the irreducible representations of  $\mathcal{A}_{r,k}(n)$ . In addition, we describe a branching rule and show that Kronecker coefficients occur when restricting irreducible representations to direct sum of Young-type subalgebras.

**Acknowledgement:** The authors would like to thank Yohana Solomon whose helpful conversations advanced some of the arguments in this paper.

## 2. PRELIMINARIES AND DEFINITIONS

**2.1. Multisets and multiset partitions.** A *multiset* is a collection of unordered elements and the elements can be repeated. The collection of elements in a multiset will be enclosed in  $\{\!\{, \}\!\}$  to differentiate this structure from a set. We say that  $S$  is a multiset of  $[k] := \{1, 2, \dots, k\}$  of size  $r$  if  $S$  contains  $r$  elements (counting repetitions) from the set  $[k]$  (e.g.  $\{\!\{1, 1, 3, 3, 3\}\!\}$  is a multiset of  $[3]$  with 5 elements). The multiset  $S$  will sometimes be represented using exponential notation  $S = \{\!\{1^{a_1}, 2^{a_2}, \dots, k^{a_k}\}\!\}$  where the exponent  $a_i$  indicates that the element  $i$  occurs with multiplicity  $a_i$  in  $S$ . If the size of the multiset is  $r$  then  $a_1 + a_2 + \dots + a_k = r$ . The set of multisets is endowed with the operation of taking union and if  $S = \{\!\{1^{a_1}, 2^{a_2}, \dots, k^{a_k}\}\!\}$  and  $T = \{\!\{1^{b_1}, 2^{b_2}, \dots, k^{b_k}\}\!\}$  are multisets, then  $S \uplus T = \{\!\{1^{a_1+b_1}, 2^{a_2+b_2}, \dots, k^{a_k+b_k}\}\!\}$ . Multisets (like the monomials that they represent) are not uniquely totally ordered. To assume a convention, given two multisets  $S$  and  $T$  we say that  $S < T$  if  $S$  is empty and  $T$  is not or  $\max(S) < \max(T)$  or  $\max(S) = \max(T) = m$  and  $S \setminus \{\!\{m\}\!\} < T \setminus \{\!\{m\}\!\}$ . We call this the *last letter order*.

A *multiset partition* of a multiset  $S$  is a multiset of multisets,  $\{\!\{S^{(1)}, S^{(2)}, \dots, S^{(d)}\}\!\}$ , such that  $S^{(1)} \uplus S^{(2)} \uplus \dots \uplus S^{(d)} = S$  where each of the  $S^{(i)}$  are non-empty multisets. We will assume the convention that if  $\pi = \{\!\{S^{(1)}, S^{(2)}, \dots, S^{(d)}\}\!\}$  then the parts of  $\pi$  are listed in last letter order. The nonempty multisets  $S^{(i)}$  in a multiset partition are called *blocks* of the multiset partition. The *length* of a multiset partition is the number of blocks (counted with multiplicity) and will be denoted by  $\ell(\pi)$ . The *content* of a multiset partition is the multiset  $S$  that it partitions. For example,  $\{\!\{\!\{1\}\!\}, \{\!\{1, 2\}\!\}, \{\!\{1, 2\}\!\}, \{\!\{3, 3\}\!\}\!\}$  is a multiset partition with content  $\{\!\{1^3, 2^2, 3^2\}\!\}$  and length 4.

For a multiset  $S$  which is a block of a multiset partition  $\pi$ , let  $m_S(\pi)$  represent the number of times that  $S$  occurs in  $\pi$ . If  $S^{(1)}, S^{(2)}, \dots, S^{(r)}$  are the distinct multisets that appear in the multiset partition  $\pi$ , define


$$m(\pi)! = m_{S^{(1)}}(\pi)! m_{S^{(2)}}(\pi)! \cdots m_{S^{(r)}}(\pi)! .$$

In this article we are interested in multiset partitions of multisets of the set  $[k] \cup [\bar{k}] := \{1, 2, \dots, k\} \cup \{\bar{1}, \bar{2}, \dots, \bar{k}\}$  such that  $r$  elements come from  $[k]$  and  $r$  elements come from  $[\bar{k}]$ . Thus the content of the multiset partition is a multiset of size  $2r$ . A multiset partition of this type is said to have *size*  $r$  and *maximum value*  $k$ . We will denote the set of multiset partitions of size  $r$  and max value  $k$  by  $\Pi_{r,k}$ . The notation  $\Pi_{r,k,n}$  will represent the subset of  $\Pi_{r,k}$  of multiset partitions with maximum length  $n$ .

**2.2. Diagrams.** The notion of a diagram that we use here comes from diagram algebras (e.g. the Brauer algebra or partition algebra). This is a graph theoretic representation of multiset partitions in the same sense that diagrams have classically been used to represent basis elements of centralizer algebras.

A multiset partition in  $\Pi_{r,k,n}$  can be represented by a graph with two horizontal rows with  $r$  vertices each. The vertices are labeled with the elements in the parts of the multisets in such a way that the top row has only unbarred labels and the bottom row has only barred labels and the numbers on both rows are weakly increasing from left to right according to the order  $1 < 2 < \dots < k < \bar{1} < \bar{2} < \dots < \bar{k}$ . Two vertices in this graph are connected by a path if they are in the same block of the same multiset partition. Thus, it follows that each block of the multiset partition defines a connected component of the graph. We remark that there are many graphs that can represent the same multiset partition, in fact any two graphs that have the same labeled connected components will represent the same multiset partition. We define the *diagram* of a multiset partition as the equivalence class of graphs that satisfy the above conditions. In our examples the blocks of the multiset partition will be connected by paths or cycles. However, as it is usual in diagram algebras, we only care that there is a connected component containing the vertices for each block of the multiset partition and not how they are connected.

**Example 2.1.** Consider the multiset partition  $\pi = \{\{1\}, \{1, 1\}, \{2, \bar{1}, \bar{1}\}, \{4, \bar{2}, \bar{4}\}, \{\bar{4}\}\}$  of  $[4] \cup [\bar{4}]$  which is an element of  $\Pi_{5,4}$  of length 5, i.e., an element in  $\Pi_{5,4,n}$  with  $n \geq 5$ . This can also be represented by any of the following graphs:




These three graphs all represent the same multiset partition. The first two are isomorphic as labeled graphs, but the last one is not isomorphic as a graph. All three are equivalent as multiset partitions as we only care about using the connectivity to represent the blocks of the multiset partition. We note that there are more labeled graphs that can be used to represent this multiset partition.


**2.3. Colored multiset partitions.** In order to define the product in our algebra we will need colored multiset partitions. Let  $n$  be a non-negative integer. A *colored multiset partition* is an element in  $\pi$  in  $\Pi_{r,k,n}$  together with an assignment of a label from  $[n]$  to each block of  $\pi$ , such that each block has a different label (these labels are the ‘colors’).

More precisely, given a multiset partition  $\pi = \{\{S^{(1)}, \dots, S^{(d)}\} \in \Pi_{r,k,n}$  with  $S^{(i)} \leq S^{(i+1)}$  and  $\mathbf{c} = (c_1, \dots, c_d) \in [n]^d$  and a sequence such that  $c_i \neq c_j$  if  $i \neq j$  and  $c_i < c_{i+1}$  if  $S^{(i)} = S^{(i+1)}$ , then the notation  $\pi^{\mathbf{c}} = \{S_{c_1}^{(1)}, \dots, S_{c_d}^{(d)}\}$  will denote that the color  $c_i$  was assigned to block  $S^{(i)}$ . We remark that a colored multiset partition is a set since the colors are distinct and so the colored multisets are also distinct. Notice that  $n \geq d$ , otherwise it is not possible to assign distinct colors to all the parts. We will say that  $\pi^{\mathbf{c}}$  is a coloring of  $\pi$ .

**Example 2.2.** Let  $n = 7$ ,  $\pi = \{\{\{1\}\}, \{1, 1\}, \{2, \bar{1}, \bar{1}\}, \{\bar{4}\}, \{4, \bar{2}, \bar{4}\}\}$  and  $\mathbf{c} = (1, 3, 5, 2, 6)$ ,  $\pi^{(1,3,5,2,6)} = \{\{\{1\}\}_1, \{1, 1\}_3, \{2, \bar{1}, \bar{1}\}_5, \{\bar{4}\}_2, \{4, \bar{2}, \bar{4}\}_6\}$  is a colored multiset partition. This can also be represented using diagrams using bold integers to indicate the colors of the connected component.



**Remark 2.3.** If we order the blocks using the last letter order and since  $\mathbf{c}$  is an ordered sequence, we get a unique colored multiset partition for each multiset partition in  $\Pi_{r,k,n}$  and  $\mathbf{c}$ . However, when we draw the diagram of such a colored multiset partition we could list the equal blocks colored with different colors in any order.



Let  $\pi \in \Pi_{r,k,n}$  and  $\pi^{\mathbf{c}}$  a coloring of  $\pi$ , denote by  $\pi_{top}^{\mathbf{c}}$  to be the colored multiset partition restricted to the entries in  $[k]$  while  $\pi_{bot}^{\mathbf{c}}$  will be the colored multiset partition restricted to the entries in  $[\bar{k}]$  and then ignoring the bars. Similarly, we will use  $\pi_{top}$  and  $\pi_{bot}$  to represent the corresponding restrictions for uncolored multiset partitions. Consider  $\pi^{(1,3,5,6,2)}$  from Example 2.2, then  $\pi_{top}^{\mathbf{c}} = \{\{\{1\}\}_1, \{1, 1\}_3, \{2\}_5, \{4\}_6\}$  and  $\pi_{bot}^{\mathbf{c}} = \{\{1, 1\}_5, \{4\}_2, \{2, 4\}_6\}$ , notice that we have left out the bars in  $\pi_{bot}^{\mathbf{c}}$ .

We use the notation  $\pi^{\mathbf{c}} \rightarrow \pi$  if  $\pi$  is obtained from  $\pi^{\mathbf{c}}$  by ignoring the colors. That is, if  $\pi^{\mathbf{c}} = \{S_{c_1}^{(1)}, S_{c_2}^{(2)}, \dots, S_{c_d}^{(d)}\}$  for some multisets  $S^{(i)}$  and values  $c_i \in [n]$  then  $\pi = \{\{S^{(1)}, S^{(2)}, \dots, S^{(d)}\}\}$ . In this case we also say that  $\pi$  is the underlying multiset partition of  $\pi^{\mathbf{c}}$ .

**2.4. Symmetric Functions.** A *partition* is a sequence  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_{\ell(\lambda)}$  where  $\ell(\lambda)$  denotes the number of nonzero terms (called parts) of  $\lambda$ . To indicate that  $\lambda$  is a partition we use the notation  $\lambda \in \text{Par}$ . If in addition, we have that  $\lambda_1 + \lambda_2 + \dots + \lambda_{\ell(\lambda)} = n$ , we say that  $\lambda$  is a partition of  $n$  and we use  $\lambda \in \text{Par}_n$  to indicate that  $\lambda$  is a partition of  $n$ . In general, we use greek letters such as  $\lambda$  and  $\mu$  to denote partitions.

To prove the results about the representation theory of our algebra we will use results from the theory of symmetric functions. For this we need to introduce notation and well known results in the theory, these results can be found in [Mac], [Sta]. We define the ring of symmetric functions as the polynomial ring

$$\Lambda = \mathbb{Q}[p_1, p_2, p_3, \dots]$$

where for each  $i \geq 1$ , the  $p_i$  is the power sum generator of degree  $i$ . For any partition  $\lambda$ ,  $p_\lambda = p_{\lambda_1} \cdots p_{\lambda_{\ell(\lambda)}}$ . The ring  $\Lambda$  has distinguished bases  $\{p_\lambda\}_{\lambda \in \text{Par}}$ ,  $\{e_\lambda\}_{\lambda \in \text{Par}}$ ,  $\{h_\lambda\}_{\lambda \in \text{Par}}$ ,  $\{m_\lambda\}_{\lambda \in \text{Par}}$  and  $\{s_\lambda\}_{\lambda \in \text{Par}}$  as the power sum, elementary, complete homogeneous, monomial and Schur bases respectively. Let  $z_\lambda = \prod_{i \geq 1} i^{m_i(\lambda)} m_i(\lambda)!$  where  $m_i(\lambda)$  is the number of parts of size  $i$  in the partition  $\lambda$ . We use the Hall-scalar product on symmetric functions which is defined so that  $\langle s_\lambda, s_\mu \rangle = \langle p_\lambda, p_\mu / z_\mu \rangle = \langle h_\lambda, m_\mu \rangle = \delta_{\lambda\mu}$ , where  $\delta_{\lambda\mu}$  is 1 if  $\lambda = \mu$  and zero otherwise.

In addition to the ring structure of  $\Lambda$  as a polynomial ring, there is a second product called the *Kronecker product* defined on the power sum basis by  $\frac{p_\mu}{z_\mu} * \frac{p_\lambda}{z_\lambda} = \delta_{\lambda\mu} \frac{p_\lambda}{z_\lambda}$ . It is defined so that for all  $\lambda, \mu$  and  $\nu$  which are partitions of a positive integer  $n$ ,

$$\langle s_\lambda * s_\mu, s_\nu \rangle = \dim \text{Hom}(W_{\mathbb{C}S_n}^\nu, W_{\mathbb{C}S_n}^\lambda \otimes W_{\mathbb{C}S_n}^\mu) := g_{\lambda\mu\nu} .$$

where  $W_{\mathbb{C}S_n}^\lambda$  denotes the irreducible representation of the symmetric group algebra,  $\mathbb{C}S_n$ , indexed by the partition  $\lambda$ . The coefficients  $g_{\lambda\mu\nu}$  are known as *Kronecker coefficients*.

We will make use of ‘plethystic notation’ for some of the symmetric function expressions (see [LR] for an expository reference). Let  $E(x_1, x_2, \dots; q) = E(X; q)$  be an algebraic expression in indeterminates  $q, x_1, x_2, \dots$ , then for an element  $f \in \Lambda$ , the evaluation of a symmetric function at an expression  $E(X; q)$  is denoted  $f[E(X; q)]$  and is defined by expanding  $f$  in the power sum basis and replacing each  $p_r$  by  $E(x_1^r, x_2^r, \dots; q^r)$ . That is, if  $f = \sum_\lambda c_\lambda p_\lambda$ , then

$$f[E(X; q)] = \sum_\lambda c_\lambda \prod_{i=1}^{\ell(\lambda)} E(x_1^{\lambda_i}, x_2^{\lambda_i}, \dots; q^{\lambda_i}) .$$

We will use  $X$  (or  $Y$ ) to represent the sum  $x_1 + x_2 + x_3 + \dots$  (or  $y_1 + y_2 + y_3 + \dots$ ) and use freely that  $\mathbb{Q}[p_1, p_2, p_3, \dots] \cong \mathbb{Q}[p_1[X], p_2[X], p_3[X], \dots]$ .

The following symmetric function identities will be used to obtain a generating function expression in Section 5. In the Lemma, for any integer  $r$ , the  $s_r$  denotes the Schur function indexed by partition consisting of one part  $r$ .

**Lemma 2.4.** [Mac, Ch. 1, Sec. 4 and 5] *Let  $\Omega = 1 + s_1 + s_2 + s_3 + \cdots$ ,*

$$(a) \quad \Omega[XY] = \prod_{i \geq 1} \prod_{j \geq 1} \frac{1}{1 - x_i y_j} = \sum_{\lambda \in \text{Par}} s_\lambda[X] s_\lambda[Y],$$

$$(b) \quad \langle \Omega[XY], f[X] \rangle = f[Y],$$

$$(c) \quad \Omega[X + Y] = \Omega[X] \Omega[Y] \text{ or } s_r[X + Y] = \sum_{a=0}^r s_a[X] s_{r-a}[Y].$$

### 3. THE GENERIC MULTISSET PARTITION ALGEBRA

Let  $r$  and  $k$  be non-negative integers and  $x$  be a nonzero indeterminate. In Section 2.1 we defined  $\Pi_{r,k}$  as the set of multiset partitions of size  $r$  and maximum value  $k$ . Define  $\text{MP}_{r,k}(x)$  to be the  $\mathbb{C}(x)$ -span of elements indexed by multiset partitions with coefficients in  $\mathbb{C}(x)$ , i.e,  $\text{span}_{\mathbb{C}(x)}\{X_\pi \text{ for } \pi \in \Pi_{r,k}\}$ . We set  $\text{MP}_{0,0}(x) = \mathbb{C}(x)$  and we identify it with the linear span of the empty multiset partition.

In what follows we will define a product on basis elements,  $X_\pi$ , where we identify the diagram for  $\pi$  with the basis element. This product will make  $\text{MP}_{r,k}(x)$  an associative algebra over  $\mathbb{C}(x)$  with an identity element.

For  $\gamma \in \Pi_{r,k}$ , let  $\bar{\gamma}$  be the multiset partition of  $[\bar{k}] \cup [\bar{k}]$  formed by adding an extra bar to all the entries of  $\gamma$ . Let  $A$  be a set and  $S$  a multiset, then let  $S|_A$  be the restriction of the multiset  $S$  to the elements which are in  $A$  (e.g.  $S|_A = \{\{i \in S : i \in A\}\}$ ) and for a multiset partition  $\gamma$ ,  $\gamma|_A$  is the (possibly empty) multiset partition  $\{\{S|_A : S \in \gamma\}\}$ , that is  $\gamma|_A$  is the multiset partition  $\gamma$  restricted to the elements in  $A$ .

The expression for the product will involve multiset partitions with entries in  $[k] \cup [\bar{k}] \cup [\bar{k}]$ . Let  $\Gamma_{r,k}$  denote the set of multiset partitions with  $r$  entries from  $[k]$ ,  $r$  entries from  $[\bar{k}]$  and  $r$  entries from  $[\bar{k}]$  and let  $\nu$  be an element of  $\Gamma_{r,k}$ . For each distinct multiset  $S$  in  $\nu|_{[k] \cup [\bar{k}]}$ , let  $\nu_S$  be the multiset partition consisting of the parts  $T$  of  $\nu$  such that  $T|_{[k] \cup [\bar{k}]} = S$ , that is,

$$\nu_S = \{\{T \in \nu : T|_{[k] \cup [\bar{k}]} = S\}\}.$$

Also let  $\beta = \{\{S \in \nu : \forall i \in S, i \in [\bar{k}]\}\}$  be the (possibly empty) multiset partition with just the parts of  $\nu$  which have elements only in  $[\bar{k}]$ . Then define the coefficient

$$(3.1) \quad a_\nu = \prod_S \ell(\nu_S)! / m(\nu_S)!$$

where the product is over the distinct multisets  $S$  which appear in  $\nu|_{[k] \cup [\bar{k}]}$ . Also define

$$(3.2) \quad b_\nu(x) = (x - \ell(\nu|_{[k] \cup [\bar{k}]}))_{\ell(\beta)} / m(\beta)!$$

where  $(a)_r = a(a-1) \cdots (a-r+1)$ .

**Example 3.1.** Let  $r = 9$  and  $k = 2$  and consider, for example, the multiset partition  $\nu = \{\{1\}, \{1, \bar{1}\}, \{1, \bar{2}\}, \{1, \bar{2}\}, \{1, \bar{2}\}, \{1, 2, \bar{1}\}, \{1, 2, \bar{1}, \bar{1}\}, \{\bar{1}\}, \{\bar{1}\}, \{\bar{1}, \bar{2}\}, \{\bar{1}, \bar{2}, \bar{1}\},$



$\{\{\bar{2}, \bar{2}\}, \{\{\bar{2}, \bar{2}\}\}\}$ . Now  $\nu|_{[k] \cup [\bar{k}]} = \{\{\{1\}, \{1\}, \{1\}, \{1\}, \{1, \bar{2}\}, \{1, 2, \bar{1}\}, \{1, 2, \bar{1}, \bar{1}\}, \{\bar{1}\}, \{\bar{2}, \bar{2}\}, \{\{\bar{2}, \bar{2}\}\}\}$  has 6 distinct multisets and has length 10. Below we list the distinct multisets  $S$  of  $\nu|_{[k] \cup [\bar{k}]}$  and the corresponding  $\nu_S$ .

$S$	$\nu_S$
$\{\{1\}\}$	$\{\{\{1\}, \{1, \bar{1}\}, \{1, \bar{2}\}, \{1, \bar{2}\}\}\}$
$\{\{1, \bar{2}\}\}$	$\{\{\{1, \bar{2}\}\}\}$
$\{\{1, 2, \bar{1}\}\}$	$\{\{\{1, 2, \bar{1}\}\}\}$
$\{\{1, 2, \bar{1}, \bar{1}\}\}$	$\{\{\{1, 2, \bar{1}, \bar{1}\}\}\}$
$\{\{\bar{1}\}\}$	$\{\{\{\bar{1}, \bar{2}, \bar{1}\}\}\}$
$\{\{\bar{2}, \bar{2}\}\}$	$\{\{\{\bar{2}, \bar{2}\}, \{\bar{2}, \bar{2}\}\}\}$

From this table we conclude that  $a_\nu = 12$  (for the multiset  $S = \{\{1\}\}$  is  $\ell(\nu_S)!/m(\nu_S)! = 12$ , while from each of the other multisets  $S$ ,  $\ell(\nu_S)!/m(\nu_S)! = 1$ ).

The multiset partition  $\beta$  consisting of the parts with only entries in  $[\bar{2}]$  is  $\beta = \{\{\{\bar{1}\}, \{\bar{1}\}, \{\bar{1}, \bar{2}\}\}\}$ , and hence  $b_\nu(x) = (x - 10)(x - 11)(x - 12)/2$

Let  $\pi$  and  $\gamma$  be elements of  $\Pi_{r,k}$ , define the product in  $\text{MP}_{k,r}(x)$  by

$$(3.3) \quad X_\pi \cdot X_\gamma = \sum_{\nu} a_\nu b_\nu(x) X_{\nu|_{[k] \cup [\bar{k}]}}$$

where the sum is over all multiset partitions  $\nu \in \Gamma_{r,k}$  such that  $\nu|_{[k] \cup [\bar{k}]} = \pi$  and  $\nu|_{[\bar{k}] \cup [\bar{k}]} = \bar{\gamma}$  and where, by convention, we consider  $\nu|_{[k] \cup [\bar{k}]}$  to be a multiset partition with entries in  $[k] \cup [\bar{k}]$  by removing the second bar on elements in  $[\bar{k}]$ . By definition, this product is zero if  $\pi|_{[\bar{k}]} \neq \bar{\gamma}|_{[\bar{k}]}$  (that is, if the lower row of  $\pi$  does not have the same multisets as the upper row of  $\bar{\gamma}$ ).

The following example is relatively small, but shows the dependence on the multiset partitions in  $\Gamma_{r,k}$ .

**Example 3.2.** Let  $\pi = \{\{\{1\}, \{1, \bar{1}\}, \{\bar{1}\}\}\} \in \Pi_{2,1}$ , then to calculate  $X_\pi^2 \in \text{MP}_{2,1}(x)$ , there are four multiset partitions  $\nu$  which contribute to the sum with  $\nu|_{[1] \cup [\bar{1}]} = \pi$  and  $\nu|_{[\bar{1}] \cup [\bar{1}]} = \bar{\pi}$ , they are:

$$\begin{aligned} & \{\{\{1\}, \{\bar{1}\}, \{\bar{1}\}, \{1, \bar{1}, \bar{1}\}\}\} \\ & \{\{\{1, \bar{1}\}, \{\bar{1}\}, \{1, \bar{1}, \bar{1}\}\}\} \\ & \{\{\{1\}, \{1, \bar{1}\}, \{\bar{1}, \bar{1}\}, \{\bar{1}\}\}\} \\ & \{\{\{1, \bar{1}\}, \{1, \bar{1}\}, \{\bar{1}, \bar{1}\}\}\} \end{aligned}$$

We construct the following table where each of the three columns contains the diagrams representing the three multiset partitions  $\nu$  and data representing their contribution to the sum in Equation (3.3).

three row msp $\nu$	
$\nu _{[k] \cup [\bar{k}]} \text{ and } [\bar{k}] \rightarrow [\bar{k}]$	
coefficient $a_\nu b_\nu(x)$	$x - 3 \quad 2(x - 2) \quad 4 \quad 1$

Therefore

$$X_\pi^2 = (x - 2)X_{\{\{1\}, \{\bar{1}\}, \{1, \bar{1}\}\}} + 2(x - 2)X_{\{\{1, \bar{1}\}, \{1, \bar{1}\}\}} + 4X_{\{\{1\}, \{1\}, \{\bar{1}\}, \{\bar{1}\}\}}.$$

**Example 3.3.** Let  $r = 4$ ,  $k = 2$  and consider the multiset partitions  $\pi = \{\{1, 1\}, \{\bar{1}, \bar{2}\}, \{1, 2, \bar{1}, \bar{2}\}\}$ , and  $\gamma = \{\{1, 2\}, \{\bar{1}, \bar{1}\}, \{1, 2, \bar{1}, \bar{1}\}\}$ . There are four multiset partitions  $\nu$  with entries in  $\{1, 2, \bar{1}, \bar{2}, \bar{1}, \bar{2}\}$  which contribute to the terms in the product  $X_\pi X_\gamma$ .

three row msp $\nu$	
$\nu _{[k] \cup [\bar{k}]} \text{ and } [\bar{k}] \rightarrow [\bar{k}]$	
coefficient $a_\nu b_\nu(x)$	$(x - 3) \quad (x - 2) \quad 1 \quad 2$

Therefore

$$X_\pi X_\gamma = (x - 3)X_{\{\{1, 1\}, \{\bar{1}, \bar{1}\}, \{1, 2, \bar{1}, \bar{1}\}\}} + (x - 2)X_{\{\{1, 1, \bar{1}, \bar{1}\}, \{1, 2, \bar{1}, \bar{1}\}\}} + X_{\{\{1, 1, \bar{1}, \bar{1}\}, \{1, 2\}, \{\bar{1}, \bar{1}\}\}} + 2X_{\{\{1, 1\}, \{\bar{1}, \bar{1}\}, \{1, 2\}, \{\bar{1}, \bar{1}\}\}}.$$

Observe that by definition this product is a well-defined algebra product. In addition, for any  $\pi, \gamma, \tau \in \Pi_{r, k}$  the coefficient of  $X_\tau$  in a product  $X_\pi X_\gamma$  is always a polynomial in the variable  $x$ .

**Proposition 3.4.** *The product defined in Equation (3.3) is associative.*

*Proof.* Proving associativity of this product directly is difficult since we would need to give an interpretation to a product of three basis elements. That is, we would have to find a formula for the coefficients occurring in these triple products. Thus, we show associativity indirectly using the fact that the algebra  $\text{MP}_{r, k}(x)$  is isomorphic to an endomorphism algebra  $\mathcal{A}_{r, k}(n)$  (see Section 4 for details), for specialized values of  $x = n$  with  $n$  and integer

such that  $n \geq 2r$ , see Theorem 4.7. The algebra  $\mathcal{A}_{r,k}(n)$  has a basis  $\{O_\pi \mid \pi \in \Pi_{r,k}\}$  and in Theorem 4.7 we show that  $X_\pi$  corresponds to  $O_\pi$  under the isomorphism.

For any  $\pi, \gamma, \zeta, \tau \in \Pi_{r,k}$ , the coefficient of  $X_\tau$  in  $(X_\pi X_\gamma)X_\zeta - X_\pi(X_\gamma X_\zeta)$  is a polynomial in  $x$ . For all  $x = n$  where  $n$  is an integer which is greater than or equal to  $2r$ , by Theorem 4.7 this polynomial evaluates to 0 because

$$\begin{aligned} & \text{the coefficient of } X_\tau \text{ in } (X_\pi X_\gamma)X_\zeta - X_\pi(X_\gamma X_\zeta) \\ &= \text{the coefficient of } O_\tau \text{ in } (O_\pi O_\gamma)O_\zeta - O_\pi(O_\gamma O_\zeta) \end{aligned}$$

and because the product of the  $O_\pi$  is associative in  $\mathcal{A}_{r,k}(n)$ . Since this holds for an infinite number of values if  $x = n$ , as a polynomial in  $x$  the coefficient must be the 0 polynomial.  $\square$

A multiset of  $[k] \cup [\bar{k}]$  is self-symmetric if replacing  $i$  with  $\bar{i}$  and  $\bar{i}$  with  $i$  is the same multiset. A multiset partition is self-symmetric if each of the parts are individually self-symmetric. For example  $\{\{1, 1, \bar{1}, \bar{1}\}, \{1, 2, 2, \bar{1}, \bar{2}, \bar{2}\}, \{1, \bar{1}\}, \{2, 3, \bar{2}, \bar{3}\}\}$  is self-symmetric, but for instance,  $\{\{1, 2, \bar{2}\}, \{2, \bar{1}, \bar{2}\}\}$  is not.

**Proposition 3.5.** *For  $r, k > 0$ , the element  $I_{r,k} = \sum_{\pi} X_\pi$  where the sum is over all self-symmetric multiset partitions  $\pi \in \Pi_{r,k}$  is the identity element of  $\mathbb{MP}_{r,k}(x)$ .*

*Proof.* Fix a  $\gamma \in \Pi_{r,k}$  and to compute  $I_{r,k}X_\gamma$  we know that for any self symmetric  $\pi$ ,  $X_\pi X_\gamma = 0$  unless  $\pi|_{[\bar{k}]} = \bar{\gamma}|_{[\bar{k}]}$ . Therefore  $I_{r,k}X_\gamma = X_\pi X_\gamma$  where  $\pi$  is the self-symmetric element of  $\Pi_{r,k}$  such that  $\pi|_{[\bar{k}]} = \bar{\gamma}|_{[\bar{k}]}$ .

The product  $X_\pi X_\gamma$  is the sum over all  $\nu$  such that  $\nu|_{[k] \cup [\bar{k}]} = \pi$  and  $\nu|_{[\bar{k}] \cup [\bar{k}]} = \bar{\gamma}$ . Notice that if we unbar the multisets in  $\nu|_{[\bar{k}]}$ , the result is the multiset partition  $\nu|_{[k]}$  it follows that  $\nu|_{[k] \cup [\bar{k}]} = \gamma$  (if we remove one bar from the double barred elements). Thus, it follows that  $\nu = \{S \uplus \{i : \bar{i} \in S\} : S \in \bar{\gamma}\}$  is the single  $\nu \in \Gamma_{r,k}$  that satisfies this condition.

The multiset  $\{S \in \nu : \forall i \in S, i \in [\bar{k}]\}$  is empty since  $\pi$  is self-symmetric so each multiset with an  $\bar{i} \in [\bar{k}]$  also contains the corresponding  $i \in [k]$ , therefore  $b_\nu(x) = 1$  by Equation (3.2). For each multiset  $S$  in  $\nu|_{[k] \cup [\bar{k}]}$ , the multisets in  $\nu_S$  will consist of only the multiset  $S \uplus \{\bar{i} : i \in S|_{[k]}\}$  (possibly with a multiplicity) and hence  $\ell(\nu_S)! = m(\nu_S)!$  and by Equation (3.1),  $a_\nu = 1$ . It follows that  $I_{r,k}X_\gamma = X_\pi X_\gamma = X_\gamma$ .

Since the transformation which sends  $X_\pi \mapsto X_{\bar{\pi}}$  is an algebra antihomomorphism, a right identity exists and the right and left identity of an algebra must be the same.  $\square$

#### 4. $\mathbb{MP}_{r,k}(n)$ IS A CENTRALIZER ALGEBRA OF $\mathbb{CS}_n$

In this section we introduce a centralizer algebra of the symmetric group. Our objective is to show that this algebra is isomorphic to  $\mathbb{MP}_{r,k}(n)$ .

**4.1. An  $S_n$  action on polynomials.** Let  $V_{n,k} = (\mathbb{C}^n)^k$  denote the  $nk$  dimensional vector space of sequences  $(v_1, v_2, \dots, v_k)$  where  $v_j \in \mathbb{C}^n$ . Notice that  $V_{n,k}$  can be identified with the space of  $n \times k$  matrices  $M_{n,k}$ , where each  $v_j$  is a column vector. With this identification, we can define  $e_{ij}$  as the  $n \times k$  matrix with 1 in the  $ij$ -th entry and zeros everywhere else, then  $\{e_{ij} \mid 1 \leq i \leq n, 1 \leq j \leq k\}$  is a basis for  $V_{n,k}$ .

The entries  $x_{ij}$  of a matrix  $X$  in  $M_{n,k}$  can be viewed as functions on  $V_{n,k}$ . Then  $\mathcal{P}(V_{n,k})$  is the polynomial ring in the commuting variables

$$\{x_{ij} \mid 1 \leq i \leq n, 1 \leq j \leq k\}.$$

We will denote by  $\mathcal{P}^r(V_{n,k})$  the space of polynomials of degree  $r$  (the symmetric tensor). This space has as basis the monomials in the  $x_{ij}$  of degree  $r$ , we order the monomials in weakly decreasing order in such a way that  $j_1 \leq j_2 \leq \dots \leq j_r$  and  $i_s \leq i_t$  if  $j_s = j_t$ . Assuming this order, then

$$\{x_{i_1 j_1} x_{i_2 j_2} \cdots x_{i_r j_r} \mid 1 \leq i_s \leq n \text{ and } 1 \leq j_t \leq k\}$$

is an ordered basis for  $\mathcal{P}^r(V_{n,k})$ . Thus, the dimension of  $\mathcal{P}^r(V_{n,k})$  is  $\binom{n+k+r-1}{r}$ .

Using the identification of  $V_{n,k}$  with the  $n \times k$  matrices, then  $GL_n$  acts on  $V_{n,k}$  via left multiplication. This action induces an action of  $GL_n$  on  $\mathcal{P}(V_{n,k})$ , for the details of this action see [GW2], Section 5.6.2. Now the symmetric group,  $S_n$ , embeds in  $GL_n$  as permutation matrices and thus acts on  $V_{n,k}$ ; for  $\sigma \in S_n$  and  $e_{ij}$  a basis element of  $V_{n,k}$ , we have

$$\sigma \cdot e_{i,j} = e_{\sigma(i),j}.$$

With the identification of  $e_{ij}$  as a matrix unit, this action corresponds to permutation of rows of the matrix or equivalently left multiplication by the permutation matrix  $\sigma$ . Then by restriction,  $S_n$  acts on  $\mathcal{P}(V_{n,k})$ , this action of  $S_n$  has a simple description on monomials. For  $x_{i_1 j_1} x_{i_2 j_2} \cdots x_{i_r j_r} \in \mathcal{P}^r(V_{n,k})$  and  $\sigma \in S_n$ , we have

$$(4.1) \quad \sigma \cdot x_{i_1 j_1} x_{i_2 j_2} \cdots x_{i_r j_r} = x_{\sigma^{-1}(i_1) j_1} x_{\sigma^{-1}(i_2) j_2} \cdots x_{\sigma^{-1}(i_r) j_r}.$$

**Remark 4.1.** This action decomposes  $\mathcal{P}^r(V_{n,k})$  into invariant subspaces indexed by multisets of size  $r$  with elements from the set  $\{1, \dots, k\}$ . These multisets can be identified with a weakly increasing sequence  $(j_1, j_2, \dots, j_r)$  which correspond to the second index of the variables  $x_{ij}$ . In particular, the subspace of monomials with  $j_s = s$  can be identified with the tensor product  $(\mathbb{C}^n)^{\otimes r}$  and the action of  $S_n$  is the diagonal action on tensor space. Recall that the centralizer of the diagonal action of  $S_n$  on tensor space is isomorphic to the partition algebra when  $n \geq 2r$  [J].

In Section 5, when we describe the branching rule and the restriction to simplify the presentation we will use the following notation. For a subset  $S \subseteq \{1, 2, \dots, k\}$ , we will denote  $V_{n,S} \subseteq V_{n,k}$  where  $V_{n,S}$  is the span of elements  $(v_1, v_2, \dots, v_k)$  where  $v_j = 0$  if  $j \notin S$ . That is,  $V_{n,\{1,2,\dots,k\}} = V_{n,k}$  and  $V_{n,S}$  is the span of the basis elements  $\{e_{ij} : 1 \leq i \leq n, j \in S\}$ .

**4.2. The centralizer of the action.** In this section we are interested in the algebra,  $\mathcal{A}_{r,k}(n) := \text{End}_{\mathbb{C}S_n}(\mathcal{P}^r(V_{n,k}))$ , of endomorphisms that commute with the action of  $S_n$  on  $\mathcal{P}^r(V_{n,k})$ , see Equation (4.1). That is,

$$\mathcal{A}_{r,k}(n) = \{f : \mathcal{P}^r(V_{n,k}) \rightarrow \mathcal{P}^r(V_{n,k}) \mid f\sigma = \sigma f \text{ for all } \sigma \in S_n\}.$$

First we want to describe a basis for this algebra. In order to do this, we use the fact that an operator  $f \in \text{End}(\mathcal{P}^r(V_{n,k}))$  is determined by a  $\binom{n+k+r-1}{r} \times \binom{n+k+r-1}{r}$  matrix  $A = (A_{(\mathbf{i}, \mathbf{j})}^{(\mathbf{i}', \mathbf{j}')} )$ , where  $(\mathbf{i}, \mathbf{j}) = ((i_1, j_1), \dots, (i_r, j_r))$  satisfying  $i_s \in [n]$  and  $j_s \in [k]$  ordered such

that  $j_1 \leq j_2 \leq \dots \leq j_r$  and  $i_s \leq i_t$  if  $j_s = j_t$ . Similarly  $(\mathbf{i}', \mathbf{j}') = ((i'_1, \bar{j}_1), \dots, (i'_r, \bar{j}_r))$ , where  $i'_a \in [n]$  and  $\bar{j}_b \in [\bar{k}]$ . The reason we place an overline is to make sure that we emphasize that the primed correspond to the column indices. The order for the column indices is the same as for the unbarred. Abusing notation, we will use  $\mathbf{j}$  as both a sequence ordered in weakly increasing order and as a multiset of the set  $[k]$  with  $r$  elements. Similarly,  $\mathbf{j}'$  denotes a multiset of  $[\bar{k}]$  and a weakly increasing sequence.

To simplify notation we set  $x_{(\mathbf{i}, \mathbf{j})} = x_{i_1 j_1} x_{i_2 j_2} \cdots x_{i_r j_r}$ , in this notation,  $\sigma x_{(\mathbf{i}, \mathbf{j})} = x_{(\sigma^{-1}(\mathbf{i}), \mathbf{j})}$  where for  $\sigma \in S_n$ ,  $\sigma^{-1}(\mathbf{i}) = (\sigma^{-1}(i_1), \sigma^{-1}(i_2), \dots, \sigma^{-1}(i_r))$ . Therefore, if  $A$  is a matrix corresponding to an endomorphism in  $\text{End}(\mathcal{P}^r(V_{n,k}))$ , then

$$Ax_{(\mathbf{i}, \mathbf{j})} = \sum_{(\mathbf{i}', \mathbf{j}')} A_{(\mathbf{i}, \mathbf{j})}^{(\mathbf{i}', \mathbf{j}')} x_{(\mathbf{i}', \mathbf{j}')} .$$

**Lemma 4.2.** *For  $r, k, n > 0$ , the element  $A \in \mathcal{A}_{r,k}(n)$  if and only if  $A_{(\mathbf{i}, \mathbf{j})}^{(\mathbf{i}', \mathbf{j}')} = A_{(\sigma(\mathbf{i}), \mathbf{j})}^{(\sigma(\mathbf{i}'), \mathbf{j}')}$  for all  $\sigma \in S_n$  and  $(\mathbf{i}, \mathbf{j})$  and  $(\mathbf{i}', \mathbf{j}')$  are as specified above.*

*Proof.* A straightforward computation show that  $\sigma Ax_{(\mathbf{i}, \mathbf{j})} = \sum_{(\mathbf{i}', \mathbf{j}')} A_{(\mathbf{i}, \mathbf{j})}^{(\mathbf{i}', \mathbf{j}')} x_{(\sigma^{-1}(\mathbf{i}'), \mathbf{j}')}$  and  $A\sigma x_{(\mathbf{i}, \mathbf{j})} = \sum_{(\mathbf{i}'', \mathbf{j}'')} A_{(\sigma^{-1}(\mathbf{i}), \mathbf{j})}^{(\mathbf{i}'', \mathbf{j}'')} x_{(\mathbf{i}'', \mathbf{j}'')}$ . Since  $A \in \mathcal{A}_{r,k}(n)$ ,  $A\sigma = \sigma A$ . Hence, by equating coefficients, the claim follows.  $\square$

The main consequence of Lemma 4.2 is that  $A$  commutes with the action of  $S_n$  on  $\mathcal{P}^r(V_{n,k})$  if and only if the matrix entries of  $A$  are equal on  $S_n$ -orbits. We now describe these orbits and relate them to the combinatorics described in the previous sections.

Observe that each fixed pair  $((\mathbf{i}, \mathbf{j}), (\mathbf{i}', \mathbf{j}'))$  determines a matrix unit  $E_{(\mathbf{i}, \mathbf{j})}^{(\mathbf{i}', \mathbf{j}')} \in \text{End}(\mathcal{P}^r(V_{n,k}))$  with 1 in the  $((\mathbf{i}, \mathbf{j}), (\mathbf{i}', \mathbf{j}'))$  position and zeros everywhere else. Therefore, the elements in  $\mathcal{A}_{r,k}(n)$  can be written as linear combinations of these matrix units. Each fixed pair  $((\mathbf{i}, \mathbf{j}), (\mathbf{i}', \mathbf{j}'))$  determines a partition of the multiset  $\mathbf{j} \uplus \mathbf{j}'$  into at most  $n$  blocks obtained by placing  $j_a$  and  $j_b$  in the same block if and only if  $i_a = i_b$ . Two pairs  $((\mathbf{i}, \mathbf{j}), (\mathbf{i}', \mathbf{j}'))$  and  $((\mathbf{s}, \mathbf{t}), (\mathbf{s}', \mathbf{t}'))$  are in the same  $S_n$  orbit if they give rise to the same multiset partition.

Recall that in Section 2.3 we introduced the notion of a colored multiset partition. This is a data structure that encodes the objects which index the matrix units in  $\text{End}(\mathcal{P}^r(V_{n,k}))$ . Each pair  $((\mathbf{i}, \mathbf{j}), (\mathbf{i}', \mathbf{j}'))$  corresponds to a colored multiset partition, where the colors are determined by the  $\mathbf{i}$  and  $\mathbf{i}'$  and the underlying multiset partition is determined by the  $\mathbf{j}$  and  $\mathbf{j}'$ . This means that each colored multiset partition corresponds to a matrix unit. Hence we can define  $E_{\pi^c} := E_{(\mathbf{i}, \mathbf{j})}^{(\mathbf{i}', \mathbf{j}')}$  where the pair  $((\mathbf{i}, \mathbf{j}), (\mathbf{i}', \mathbf{j}'))$  is the sequence of pairs encoded by the colored multiset partition  $\pi^c$ .

**Example 4.3.** If  $r = 5$ ,  $k = 4$  and  $n = 7$ , let  $(\mathbf{i}, \mathbf{j}) = ((1, 1), (3, 1), (3, 1), (5, 2), (6, 4))$  and  $(\mathbf{i}', \mathbf{j}') = ((5, \bar{1}), (5, \bar{1}), (6, \bar{2}), (6, \bar{4}), (2, \bar{4}))$ . The pair  $((\mathbf{i}, \mathbf{j}), (\mathbf{i}', \mathbf{j}'))$  is an index for a matrix unit which determines a colored multiset partition  $\pi^{(1,3,5,2,6)} = \{ \{ \{ 1 \} \}_1, \{ \{ 1, 1 \} \}_3, \{ \{ 2, \bar{1}, \bar{1} \} \}_5, \{ \{ \bar{4} \} \}_2, \{ \{ 4, \bar{2}, \bar{4} \} \}_6 \}$  of  $[4] \cup [\bar{4}]$ .

The pair  $((2, 1), (4, 1), (4, 1), (1, 2), (3, 4)), ((1, \bar{1}), (1, \bar{1}), (3, \bar{2}), (3, \bar{4}), (5, \bar{4}))$  is another index corresponding to the colored multiset partition is  $\pi^{(2,4,1,5,3)}$  and has the same underlying multiset partition  $\pi$ .

Using the notation that we introduced at the end of Section 2.3, the product of two matrix units  $E_{\pi^c}$  and  $E_{\gamma^{c'}}$  is

$$E_{\pi^c} \cdot E_{\gamma^{c'}} = \delta_{\tau_{top}^{c''}, \pi_{top}^c} \delta_{\tau_{bot}^{c''}, \gamma_{bot}^c} \delta_{\pi_{bot}^{c''}, \gamma_{top}^{c'}} E_{\tau^{c''}}$$

where  $\delta_{A,B}$  is the Kronecker delta. For a permutation  $\sigma \in S_n \subseteq \text{End}(\mathcal{P}^r(V_{n,k}))$  which acts on  $\mathcal{P}^r(V_{n,k})$ , we have that  $\sigma E_{\pi^c} = E_{\pi^{\sigma(c)}} \sigma$ .

Let  $\pi$  be a multiset partition with  $r$  elements from  $[k]$  and  $r$  elements from  $[\bar{k}]$ , then define

$$(4.2) \quad O_\pi := \sum_{\pi^c \rightarrow \pi} E_{\pi^c}$$

where the sum runs over all colored multiset set partitions  $\pi^c$  that give rise to the same multiset partition  $\pi$ . Notice that since the colors that index the multisets are all in  $[n]$ , there can be at most  $n$  parts in the set partition  $\pi$ , otherwise the sum is empty and the element is 0.

**Proposition 4.4.** *(The orbit basis) Recall that  $\Pi_{r,k,n}$  is the set of all multiset partitions with at most  $n$  parts of a multiset with  $r$  elements from the set  $[k]$  and  $r$  elements from  $[\bar{k}]$ . For  $r, k, n > 0$ , the set of elements  $\{O_\pi \mid \pi \in \Pi_{r,k,n}\}$  is a basis of  $\mathcal{A}_{r,k}(n)$ .*

*Proof.* Since if  $\pi^c \rightarrow \pi$ , then  $\pi^{\sigma(c)} \rightarrow \pi$  for all  $\sigma \in S_n$ , it follows that for all  $\sigma \in S_n$  that  $\sigma O_\pi = O_\pi \sigma$  and hence  $O_\pi \in \mathcal{A}_{r,k}(n)$ . By Lemma 4.2, we have that for any  $A \in \mathcal{A}_{r,k}(n)$ ,  $A$  is a linear combination of the  $O_\pi$ . Since each  $O_\pi$  is the sum of a disjoint set of matrix units, these matrices are linearly independent and therefore the collection is a basis.  $\square$

It follows that the dimension of  $\mathcal{A}_{r,k}(n)$  is equal to  $|\Pi_{r,k,n}|$ . The number of multiset partitions has been previously studied and asymptotic formulas have been obtained when the multiset partitions are counted by type, see [Bend] for details. A generating function formula for this dimension is given in Corollary 5.4.

**4.3.  $\mathcal{A}_{r,k}(n)$  is isomorphic to  $\text{MP}_{r,k}(n)$  for  $n \geq 2r$ .** In the previous section we showed that  $\mathcal{A}_{r,k}(n)$  has a basis indexed by the elements in  $\Pi_{r,k,n}$ . If  $n \geq 2r$ ,  $\Pi_{r,k,n} = \Pi_{r,k}$  and this shows that  $\mathcal{A}_{r,k}(n)$  and  $\text{MP}_{r,k}(n)$  are isomorphic as vector spaces. In this section we will show that they are isomorphic as algebras.

**Lemma 4.5.** *For positive integers  $r, k, n$  and  $\pi, \gamma \in \Pi_{r,k,n}$ , the product on the orbit basis can be expressed as*

$$(4.3) \quad O_\pi \cdot O_\gamma = \sum_{c: \pi^c \rightarrow \pi} \sum_{c': \gamma^{c'} \rightarrow \gamma} \sum_{\tau \in \Pi_{r,k,n}} \delta_{\tau_{top}^{c'}, \pi_{top}^c} \delta_{\pi_{top}^c, \tau_{top}^{(1,2,\dots,\ell(\tau))}} \delta_{\gamma_{bot}^{c'}, \tau_{bot}^{(1,2,\dots,\ell(\tau))}} O_\tau \cdot$$

*Proof.* Using Equation (4.2), we expand  $O_\pi$  and  $O_\gamma$  using colored multiset partitions, we have

$$\begin{aligned} O_\pi \cdot O_\gamma &= \sum_{\mathbf{c}:\pi^{\mathbf{c}} \rightarrow \pi} \sum_{\mathbf{c}':\gamma^{\mathbf{c}'} \rightarrow \gamma} E_{\pi^{\mathbf{c}}} E_{\gamma^{\mathbf{c}'}} \\ &= \sum_{\mathbf{c}:\pi^{\mathbf{c}} \rightarrow \pi} \sum_{\mathbf{c}':\gamma^{\mathbf{c}'} \rightarrow \gamma} \sum_{\tau} \sum_{\mathbf{c}'':\tau^{\mathbf{c}''} \rightarrow \tau} \delta_{\gamma_{top}^{\mathbf{c}'}, \pi_{bot}^{\mathbf{c}}} \delta_{\pi_{top}^{\mathbf{c}}, \tau_{bot}^{\mathbf{c}''}} \delta_{\gamma_{bot}^{\mathbf{c}'}, \tau_{bot}^{\mathbf{c}''}} E_{\tau^{\mathbf{c}''}}. \end{aligned}$$

Now since the orbit basis is defined so that the support is disjoint when expressed in the basis elements  $\{E_{\pi^{\mathbf{c}}}\}$  of  $\text{End}(\mathcal{P}^r(M_{n,k}))$ , we only need take a coefficient of a representative element from the expression. That is, in any expression in  $\text{End}_{\mathbb{C}S_n}(\mathcal{P}^r(M_{n,k}))$ , the coefficient of  $O_\tau$  is equal to the coefficient of  $E_{\tau^{\mathbf{c}}}$  for any coloring  $\mathbf{c}$ . Hence we can take  $\mathbf{c} = (1, 2, \dots, \ell(\tau))$  as a representative element of that orbit, and Equation (4.3) follows.  $\square$

Now the following lemma shows that the product in Equation (4.3) can be transformed as a sum over elements of  $\Gamma_{r,k}$ , i.e., multisets with  $r$  elements from each in  $[k]$ ,  $[\bar{k}]$  and  $[\bar{k}]$ . Recall that a coloring  $\mathbf{c}$  of a multiset partition  $\pi$  is a sequence with all distinct entries in  $[n]$ . The entries have to be distinct because two blocks of the multiset partition  $\pi$  cannot have the same color.

**Lemma 4.6.** *Fix positive integers  $r, k, n$  and elements  $\pi, \gamma, \tau \in \Pi_{r,k,n}$ . There is a bijection between pairs of colorings  $(\mathbf{c}, \mathbf{c}') \in [n]^{\ell(\pi)} \times [n]^{\ell(\gamma)}$ , each with distinct entries, of  $\pi$  and  $\gamma$  such that  $\gamma_{top}^{\mathbf{c}'} = \pi_{bot}^{\mathbf{c}}$ ,  $\pi_{top}^{\mathbf{c}} = \tau^{(1,2,\dots,\ell(\tau))}$  and  $\gamma_{bot}^{\mathbf{c}'} = \tau^{(1,2,\dots,\ell(\tau))}$  and pairs  $(\nu, \mathbf{c}'')$  with  $\nu \in \Gamma_{r,k}$ ,  $\mathbf{c}'' \in [n]^{\ell(\nu)}$  with distinct entries such that  $\nu|_{[k] \cup [\bar{k}]} = \pi$ ,  $\nu|_{[\bar{k}] \cup [\bar{k}]} = \bar{\gamma}$ ,  $\nu^{\mathbf{c}''}|_{[k] \cup [\bar{k}]} = \tau^{(1,2,\dots,\ell(\tau))}$ .*

*Proof.* Consider a pair of colored multiset partitions  $(\pi^{\mathbf{c}}, \gamma^{\mathbf{c}'})$  which satisfies  $\gamma_{top}^{\mathbf{c}'} = \pi_{bot}^{\mathbf{c}}$ ,  $\pi_{top}^{\mathbf{c}} = \tau^{(1,2,\dots,\ell(\tau))}$  and  $\gamma_{bot}^{\mathbf{c}'} = \tau^{(1,2,\dots,\ell(\tau))}$ . This pair of colorings uniquely determines a multiset partition  $\nu \in \Gamma_{r,k}$ . That is,  $\nu$  is the unique multiset partition such that there is a coloring  $\mathbf{c}''$  with  $\nu^{\mathbf{c}''}|_{[k] \cup [\bar{k}]} = \pi^{\mathbf{c}}$  and  $\nu^{\mathbf{c}''}|_{[\bar{k}] \cup [\bar{k}]} = \bar{\gamma}^{\mathbf{c}'}$ . To find  $\nu$ , start with  $\pi$  and then add elements of  $[\bar{k}]$  to the multisets from  $\pi$  if the multiset from  $\pi^{\mathbf{c}}$  shares the same color as the multiset in  $\gamma^{\mathbf{c}'}$ . The multisets with only elements from  $[\bar{k}]$  in  $\gamma$  correspond to multisets with only elements in  $[\bar{k}]$  in  $\nu$  and their coloring is determined by  $\gamma^{\mathbf{c}'}$ . That is, the  $\mathbf{c}$  and  $\mathbf{c}'$  determine  $\mathbf{c}''$ , a coloring of  $\nu$  such that  $\tau^{(1,2,\dots,\ell(\tau))} = \nu^{\mathbf{c}''}|_{[k] \cup [\bar{k}]}$ . Moreover, a coloring of  $\nu$  also determines the colored multiset partitions  $\pi^{\mathbf{c}}$  and  $\gamma^{\mathbf{c}'}$  by reversing the process.  $\square$

**Theorem 4.7.** *For positive integers  $n, k, r$  such that  $n \geq 2r$ , the algebra  $\text{MP}_{r,k}(n)$  is isomorphic to  $\mathcal{A}_{r,k}(n)$ .*

*Proof.* We need to show that there is a bijective homomorphism. We know that  $\{X_\pi \mid \pi \in \Pi_{r,k}\}$  is a basis for  $\text{MP}_{r,k}(n)$  and  $\{O_\pi \mid \pi \in \Pi_{r,k,n}\}$  is a basis for  $\mathcal{A}_{r,k}(n)$  and since  $n \geq 2r$ ,  $\Pi_{r,k} = \Pi_{r,k,n}$ . Then we define a linear map  $\phi : \text{MP}_{r,k}(n) \rightarrow \mathcal{A}_{r,k}(n)$  on the basis  $\phi(X_\pi) = O_\pi$  and extend this map linearly.





of  $k$  and  $r$ , it may be the case that for only a subset of the partitions  $\lambda$  will  $W_{\mathbb{C}S_n}^\lambda$  occur with non-zero multiplicity in  $\mathcal{P}^r(V_{n,k})$ . Let

$$(5.1) \quad W_{\mathcal{A}_{r,k}(n)}^\lambda := \text{Hom}(W_{\mathbb{C}S_n}^\lambda, \mathcal{P}^r(V_{n,k}))$$

represent the irreducible  $\mathcal{A}_{r,k}(n)$ -module indexed by the partition  $\lambda$ .

For any partition  $\lambda$  of  $n$ , we define the Reynolds operator  $R^\lambda = \frac{1}{n!} \sum_{\sigma \in S_n} \chi^\lambda(\sigma) \sigma$ . The operator  $R^\lambda$  projects a  $\mathbb{C}S_n$ -module to the component of the module consisting of the direct sum of  $\mathbb{C}S_n$  irreducibles indexed by the partition  $\lambda$ . This means that for a  $\mathbb{C}S_n$ -module  $W$ , we have

$$(5.2) \quad W \cong \bigoplus_{\lambda \in \text{Par}_n} R^\lambda W .$$

In the case that  $W = \mathcal{P}^r(V_{n,k})$  the action of  $\mathcal{A}_{r,k}(n)$  commutes with the  $S_n$  action and  $R^\lambda W$  is both a  $\mathbb{C}S_n$  and a  $\mathcal{A}_{r,k}(n)$ -module and so we have that

$$(5.3) \quad R^\lambda \mathcal{P}^r(V_{n,k}) \cong W_{\mathcal{A}_{r,k}(n)}^\lambda \otimes W_{\mathbb{C}S_n}^\lambda .$$

The following proposition gives the dimension of an irreducible  $\mathcal{A}_{r,k}(n)$  representation in terms of a Schur function evaluated at a series. It also gives precise conditions for when this dimension is non-zero. The plethystic notation is defined in Section 2.4.

**Proposition 5.2.** *For positive integers  $n, k, r$  and  $\lambda \in \text{Par}_n$ ,*

$$(5.4) \quad \dim W_{\mathcal{A}_{r,k}(n)}^\lambda = \text{coefficient of } q^r \text{ in } s_\lambda \left[ \frac{1}{(1-q)^k} \right] .$$

Moreover, this coefficient is non-zero (that is, the module  $W_{\mathcal{A}_{r,k}(n)}^\lambda$  exists) if and only if  $r \geq \sum_{j \geq 1} (j-1) \cdot \sum_{i=t_{j-1}+1}^{t_j} \lambda_i$  where  $t_d = \binom{k+0-1}{0} + \binom{k+1-1}{1} + \dots + \binom{k+d-2}{d-1}$  and using the convention that  $\lambda_d = 0$  for  $d > \ell(\lambda)$ .

*Proof.* Considering  $\mathcal{P}^r(V_{n,k}) = \mathcal{P}^r(V_{n,\{1,2,\dots,k\}})$  as a  $GL_n$ -module,

$$\mathcal{P}^r(V_{n,\{1,2,\dots,k\}}) = \bigoplus_{r_1+r_2+\dots+r_k=r} \mathcal{P}^{r_1}(V_{n,\{1\}}) \otimes \mathcal{P}^{r_2}(V_{n,\{2\}}) \otimes \dots \otimes \mathcal{P}^{r_k}(V_{n,\{k\}})$$

where the sum is over all sequences of non-negative integers  $(r_1, r_2, \dots, r_k)$  whose sum is  $r$ . The  $GL_n$  character of a diagonal matrix acting on a basis  $\mathcal{P}^d(V_{n,\{j\}})$  is equal to  $s_d[X_n]$  where the  $X_n = x_1 + x_2 + \dots + x_n$  are a sum of  $n$  variables which can be specialized to the eigenvalues of the element of  $GL_n$ . Therefore this decomposition shows that the character of the action on  $\mathcal{P}^r(V_{n,\{1,2,\dots,k\}})$  is (by Lemma 2.4 part (c)),

$$\sum_{r_1+r_2+\dots+r_k=r} s_{r_1}[X_n] s_{r_2}[X_n] \dots s_{r_k}[X_n] = s_r[kX_n] .$$

Now Scharf and Thibon's result [ST, Theorem 4.1] (that, in turn, follows from a theorem due to Littlewood [Li, Theorem XI]) states that the multiplicity of an irreducible representation indexed by  $\lambda$  is equal to  $\langle s_r[kX], s_\lambda[\Omega[X]] \rangle$ .

When evaluating symmetric functions at expressions in variables  $q$ , we consider  $kq = \underbrace{q + q + \cdots + q}_{k \text{ times}}$  then by Lemma 2.4 part (c),  $\Omega[kq] = \Omega[q]^k = \frac{1}{(1-q)^k}$ . Since the coefficient

of  $q^r$  in  $\Omega[kqX] = \sum_{d \geq 0} q^d s_d[kX]$  is  $s_r[kX]$ , the multiplicity that we have just calculated is equal to the coefficient of  $q^r$  in the series:

$$(5.5) \quad \langle \Omega[kqX], s_\lambda[\Omega[X]] \rangle = s_\lambda[\Omega[kq]] = s_\lambda \left[ \frac{1}{(1-q)^k} \right]$$

where the equalities follow by applying Lemma 2.4 part (b).

Fix a positive integer  $k$  and a partition  $\lambda$  of  $n$ . To determine if the multiplicity of  $W_{\mathbb{C}S_n}^\lambda$  in  $\mathcal{P}^r(V_{n,k})$  is non-zero for a given  $r$  (or equivalently, to determine if there is a module  $W_{\mathcal{A}_{r,k}(n)}^\lambda$ ) we are asking if the coefficient of  $q^r$  is non-zero in  $s_\lambda[1/(1-q)^k]$ .

Let  $a_{r,k}^\lambda$  be equal to the number of multiset tableaux of shape  $\lambda$  with  $r$  entries from 1 to  $k$ . We know by Theorem 3.1 in [OZ4] that this is the multiplicity of  $W_{\mathbb{C}S_n}^\lambda$ . There is an injection of the tableaux enumerated by  $a_{r,k}^\lambda$  into the tableaux enumerated by  $a_{r+1,k}^\lambda$  by adding an extra entry  $k$  in the highest corner of the tableau. Hence we can say that if  $a_{r,k}^\lambda$  is non-zero, then  $a_{r+1,k}^\lambda$  is non-zero.

By Equation (5.4), it remains to identify the smallest value of  $r$  such that the coefficient of  $q^r$  in  $s_\lambda[1/(1-q)^k]$  is non-zero. First note that

$$1/(1-q)^k = 1 + \underbrace{q + q + \cdots + q}_{\binom{k+1-1}{1} \text{ times}} + \underbrace{q^2 + q^2 + \cdots + q^2}_{\binom{k+2-1}{2} \text{ times}} + \underbrace{q^3 + q^3 + \cdots + q^3}_{\binom{k+3-1}{3} \text{ times}} + \cdots .$$

The expression  $s_\lambda[1/(1-q)^k]$  is equal to the monomial expansion of  $s_\lambda[X]$  with  $x_1 = 1$ , the variables  $x_2$  through  $x_{\binom{k+1-1}{1}+1}$  equal to  $q$ , the variables  $x_{2+\binom{k+1-1}{1}}$  through  $x_{1+\binom{k+1-1}{1}+\binom{k+2-1}{2}}$  equal to  $q^2$ , and in general the variables  $x_{t_{j-1}+1}$  through  $x_{t_j}$  equal to  $q^{j-1}$ . The term of smallest degree in this expression will be the leading term of  $s_\lambda[X]$  which is  $x_1^{\lambda_1} x_2^{\lambda_2} \cdots x_{\ell(\lambda)}^{\lambda_{\ell(\lambda)}}$  and that degree will be equal to

$$\sum_{j \geq 1} (j-1) \cdot \sum_{i=t_{j-1}+1}^{t_j} \lambda_i .$$

If  $r$  is smaller than this value, then the coefficient of  $q^r$  in  $s_\lambda[1/(1-q)^k]$  is equal to 0, otherwise the coefficient is non-zero.  $\square$

**Remark 5.3.** The series on the right hand side of Equation (5.4) is perhaps more explicitly written as

$$s_\lambda \left[ \frac{1}{(1-q)^k} \right] = \sum_{\gamma \in \text{Par}_n} \frac{\chi_{S_n}^\lambda(\gamma)}{z_\gamma} \prod_{i=1}^{\ell(\gamma)} \frac{1}{(1-q^i)^k}$$

where  $\chi_{S_n}^\lambda(\gamma)$  are the values of the irreducible symmetric group characters indexed by  $\lambda$  at an element of cycle type  $\gamma$ . The special case when  $k = 1$  is due to Aiken [Aik] and in this

case the result is the multiplicity of a symmetric group irreducible in the polynomial ring in a single set of  $n$  variables (see [Sta] Example 7.73 for details).

By Theorem 3.1 of [OZ4], if  $n$  be a positive integer and  $\lambda \in \text{Par}_n$ , then the dimension of the irreducible  $\mathcal{A}_{r,k}(n)$ -module indexed by  $\lambda$  (assuming that it exists) is equal to the number of multiset valued tableaux of shape  $\lambda$  with  $r$  values from 1 through  $k$  (where a multiset tableaux is a column strict tableaux whose entries are multisets ordered by a total order).

**Corollary 5.4.** *For positive integers  $n, k, r$ , the dimension of  $\mathcal{A}_{r,k}(n)$  is equal to the coefficient of  $z^n q^r t^r$  in*

$$(5.6) \quad \prod_{i \geq 0} \prod_{j \geq 0} \frac{1}{(1 - zq^i t^j)^{\binom{k+i-1}{i} \binom{k+j-1}{j}}} .$$

**Remark 5.5.** To use this generating function for effective computation of the dimension, one should take the finite product of  $0 \leq i, j \leq r$  and expand the series in  $z$ . The coefficients of  $z^n$  will be a polynomial in  $q, t$  and the coefficient of  $q^r t^r$  will be the dimension of  $\mathcal{A}_{r,k}(n)$ . However, like the partition algebra, the dimensions of the algebras grow quickly in the parameters  $k$  and  $r$  and stabilize for  $n \geq 2r$  (e.g. as a reference point and example,  $\dim \mathcal{A}_{3,4}(3) = 22736$ ,  $\dim \mathcal{A}_{3,4}(4) = 33712$ ,  $\dim \mathcal{A}_{3,4}(5) = 36912$ , and  $\dim \text{MP}_{3,4}(x) = \dim \mathcal{A}_{3,4}(n) = 37312$  for  $n \geq 6$ ). The other coefficients of  $q^a t^b$  will be the dimension of the homomorphisms from the degree  $a$  polynomials to the degree  $b$  polynomials in the variables  $x_{ij}$  with  $1 \leq i \leq n$  and  $1 \leq j \leq k$  which commute with the action of the symmetric group.

*Proof.* (of Corollary 5.4) It is a consequence of Theorem 5.1, but also by the insertion described in [COSSZ] there is a bijection between  $\Pi_{r,k,n}$  and pairs of multiset tableaux that directly show that

$$(5.7) \quad \sum_{\lambda \in \text{Par}_n} \left( \dim W_{\mathcal{A}_{r,k}(n)}^\lambda \right)^2 = \dim \mathcal{A}_{r,k}(n) .$$

We will build on this equation to prove that the coefficient of  $z^n q^r t^r$  in Equation (5.6) is equal to  $\dim \mathcal{A}_{r,k}(n)$ .

We have by Lemma 2.4, that

$$\begin{aligned} \prod_{i \geq 0} \prod_{j \geq 0} \frac{1}{(1 - zq^i t^j)^{\binom{k+i-1}{i} \binom{k+j-1}{j}}} &= \Omega \left[ \sum_{i, j \geq 0} \binom{k+i-1}{i} \binom{k+j-1}{j} zq^i t^j \right] \\ &= \Omega \left[ \frac{z}{(1-q)^k (1-t)^k} \right] \\ &= \sum_{n \geq 0} \sum_{\lambda \in \text{Par}_n} z^n s_\lambda \left[ \frac{1}{(1-q)^k} \right] s_\lambda \left[ \frac{1}{(1-t)^k} \right] . \end{aligned}$$

Hence, by Proposition 5.2 the coefficient of  $z^n q^r t^r$  in the right hand side is equal to  $\sum_{\lambda \in \text{Par}_n} \left( \dim W_{\mathcal{A}_{r,k}(n)}^\lambda \right)^2 = \dim \mathcal{A}_{r,k}(n)$ .  $\square$

5.1. **The branching rule.** The coefficients in the product of Schur functions

$$s_{\tau(1)} s_{\tau(2)} \cdots s_{\tau(r)} = \sum_{\nu} c_{\tau(1)\tau(2)\dots\tau(r)}^{\nu} s_{\nu}$$

are known as the Littlewood-Richardson coefficients. We will use these coefficients to express what is known as a branching rule for the multiplicity of an irreducible for  $\mathcal{A}_{r-d,k-1}(n)$ -module in an irreducible of  $\mathcal{A}_{r,k}(n)$ -module.

Note that the polynomial ring decomposes as

$$(5.8) \quad \mathcal{P}^r(V_{n,\{1,2,\dots,k\}}) \cong \bigoplus_{d=0}^r \mathcal{P}^{r-d}(V_{n,\{1,2,\dots,k-1\}}) \otimes \mathcal{P}^d(V_{n,\{k\}})$$

where the action of  $\mathbb{C}S_n$  is diagonal on the tensors and we separate the variables with the second index  $k$  and the parameter  $d$  represents the total degree of the variables  $x_{ik}$  with  $1 \leq i \leq n$ . This decomposition of the polynomial ring induces an inclusion  $\phi : \mathcal{A}_{r-d,k-1}(n) \hookrightarrow \mathcal{A}_{r,k}(n)$  where for  $\pi \in \Pi_{r-d,k-1,n}$ ,  $O_{\pi} \in \mathcal{A}_{r-d,k-1}(n)$  and

$$\phi(O_{\pi}) = \sum_{\nu} O_{\nu}$$

where the sum is over  $\nu \in \Pi_{r,k,n}$  with the restriction of the entries of  $\nu$  to  $[k-1] \cup [\overline{k-1}]$  is  $\pi$  and when  $\nu$  is restricted to  $\{k, \overline{k}\}$  the result is a self symmetric set partition with  $d$  values  $k$  and  $d$  values  $\overline{k}$ .

**Theorem 5.6.** *For positive integers  $r, k, n$  and for all partitions  $\lambda, \mu \in \text{Par}_n$  and  $d \geq 0$ , the multiplicity of  $W_{\mathcal{A}_{r-d,k-1}(n)}^{\mu}$  in  $W_{\mathcal{A}_{r,k}(n)}^{\lambda}$  is equal to*

$$\dim \text{Hom}(W_{\mathbb{C}S_n}^{\lambda}, W_{\mathbb{C}S_n}^{\mu} \otimes \mathcal{P}^d(V_{n,\{k\}})) = \sum_{\alpha} \sum_{\tau^{(*)}} c_{\tau^{(*)}}^{\mu} c_{\tau^{(*)}}^{\lambda}$$

where the sum is over sequences of non-negative integers  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_{d+1})$  such that  $\sum_{i=1}^{d+1} (i-1)\alpha_i = d$  and  $\sum_{i=1}^{d+1} \alpha_i = n$  and  $\tau^{(*)} = (\tau^{(1)}, \tau^{(2)}, \dots, \tau^{(d+1)})$  where for each  $1 \leq i \leq d+1$ ,  $\tau^{(i)}$  is a partition of  $\alpha_i$ .

When  $d = 1$ , the conditions on the sum impose that  $\alpha = (n-1, 1)$  and  $\sum_{\tau^{(1)} \vdash n-1} c_{\tau^{(1)}(1)}^{\mu} c_{\tau^{(1)}(1)}^{\lambda}$

is equal to the number of ways of removing a corner cell from the partition  $\mu$  and then adding a corner cell to the result to obtain the partition  $\lambda$ . This is precisely the branching rule for the partition algebra [Ha, Mar4].

**Remark 5.7.** Note that the analogous branching rule for  $GL_k$ -modules would state how  $W_{GL_k}^{\lambda}$  for  $\lambda \in \text{Par}_r$  decomposes as a direct sum of modules  $W_{GL_{k-1}}^{\mu}$  where  $\mu$  is a partition of size smaller than or equal to  $r$ . In this case (see for instance [GW2, Section 12.2.3]), the branching rule is known as the Pieri rule and the multiplicity of  $W_{GL_{k-1}}^{\mu}$  in  $W_{GL_k}^{\lambda}$  is 1 if  $\lambda$  and  $\mu$  are ‘interlaced’ or ‘differ by a horizontal strip’ and is equal to 0 otherwise. Theorem 5.6 is a formula for the multiplicity of  $W_{\mathcal{A}_{r-d,k-1}(n)}^{\mu}$  in  $W_{\mathcal{A}_{r,k}(n)}^{\lambda}$  which could be greater than 1.

**Remark 5.8.** In [OZ2, Proposition 21] the authors gave a combinatorial interpretation for the multiplicity appearing in Theorem 5.6 in terms of tableaux that simultaneously satisfy two types of lattice conditions. The application of that expression is Theorem 13 of [OZ2] which is a multiset tableaux interpretation of

$$\dim \operatorname{Hom}(W_{\mathbb{C}S_n}^\gamma, W_{\mathbb{C}S_n}^\lambda \otimes \mathcal{P}^{\alpha_1}(V_{n,1}) \otimes \mathcal{P}^{\alpha_2}(V_{n,1}) \otimes \cdots \otimes \mathcal{P}^{\alpha_{\ell(\alpha)}}(V_{n,1})) .$$

*Proof.* (of Theorem 5.6) The subalgebra  $\mathcal{A}_{r-d,k-1}(n)$  acts exclusively on the part of the polynomial ring where the  $j$  index of the  $x_{ij}$  variables has  $j \neq k$ , so we have  $\mathcal{P}^{r-d}(V_{n,k-1}) = \bigoplus_{\mu} W_{\mathcal{A}_{r-d,k-1}(n)}^{\mu} \otimes W_{\mathbb{C}S_n}^{\mu}$ . Then by Equation (5.1) and (5.8),

$$\begin{aligned} W_{\mathcal{A}_{r,k}(n)}^{\lambda} &\cong \bigoplus_{d=0}^r \operatorname{Hom}(W_{\mathbb{C}S_n}^{\lambda}, \mathcal{P}^{r-d}(V_{n,\{1,2,\dots,k-1\}}) \otimes \mathcal{P}^d(V_{n,\{k\}})) \\ &\cong \bigoplus_{d=0}^r \bigoplus_{\mu \in \operatorname{Par}_n} W_{\mathcal{A}_{r-d,k-1}(n)}^{\mu} \otimes \operatorname{Hom}(W_{\mathbb{C}S_n}^{\lambda}, W_{\mathbb{C}S_n}^{\mu} \otimes \mathcal{P}^d(V_{n,\{k\}})) . \end{aligned}$$

Therefore restricting the action of  $\mathcal{A}_{r,k}(n)$  to the action of  $\bigoplus_{d=0}^r \mathcal{A}_{r-d,k-1}(n)$  acting on  $W_{\mathcal{A}_{r,k}(n)}^{\lambda}$ , the multiplicity of  $W_{\mathcal{A}_{r-d,k-1}(n)}^{\mu}$  will be equal to the multiplicity of  $W_{\mathbb{C}S_n}^{\lambda}$  in  $W_{\mathbb{C}S_n}^{\mu} \otimes \mathcal{P}^d(V_{n,\{k\}})$ .

By Proposition 5.2, the multiplicity of  $W_{\mathbb{C}S_n}^{\nu}$  in  $\mathcal{P}^d(V_{n,\{k\}})$  is the coefficient of  $q^d$  in  $s_{\nu} \left[ \frac{1}{1-q} \right]$ , so then  $\dim \operatorname{Hom}(W_{\mathbb{C}S_n}^{\lambda}, W_{\mathbb{C}S_n}^{\mu} \otimes \mathcal{P}^d(V_{n,\{k\}}))$  is equal to the coefficient of  $q^d$  in  $\sum_{\nu \in \operatorname{Par}_n} s_{\nu} \left[ \frac{1}{1-q} \right] \langle s_{\lambda}, s_{\mu} * s_{\nu} \rangle$ , where  $*$  indicates the Kronecker product. Then by a change of basis calculation,

$$\sum_{\nu \in \operatorname{Par}_n} s_{\nu} \left[ \frac{1}{1-q} \right] \langle s_{\lambda}, s_{\mu} * s_{\nu} \rangle = \sum_{\gamma \in \operatorname{Par}_n} m_{\gamma} \left[ \frac{1}{1-q} \right] \langle s_{\lambda}, s_{\mu} * h_{\gamma} \rangle .$$

Since  $m_{\gamma} \left[ \frac{1}{1-q} \right]$  is equal to  $m_{\gamma}[X]$  with  $x_i$  replaced with  $q^{i-1}$ , then

$$m_{\gamma} \left[ \frac{1}{1-q} \right] = \sum_{\beta} q^{\sum_i (i-1)\beta_i}$$

where the sum is over sequences  $\beta$  of non-negative integers that are permutations of the partition  $\gamma$  (with trailing zeros). Therefore the coefficient of  $q^d$  in this expression follows by applying [Mac, Example 23(d), p. 130] to show that

$$\sum_{\alpha} \langle s_{\lambda}, s_{\mu} * h_{\alpha} \rangle = \sum_{\alpha} \sum_{\tau^{(*)}} c_{\tau^{(1)}\tau^{(2)}\dots\tau^{(d+1)}}^{\mu} c_{\tau^{(1)}\tau^{(2)}\dots\tau^{(d+1)}}^{\lambda}$$

where the sum is over all sequences of non-negative integers  $\alpha = (\alpha_1, \alpha_2, \dots, \alpha_{d+1})$  that sum to  $n$  such that  $\sum_{i=1}^{d+1} (i-1)\alpha_i = d$  and where the inner sum on the right hand side is over sequence of partitions  $\tau^{(*)} = (\tau^{(1)}, \tau^{(2)}, \dots, \tau^{(d+1)})$  where  $\tau^{(i)} \in \operatorname{Par}_{\alpha_i}$  for  $1 \leq i \leq d+1$ .  $\square$

**5.2. Restriction of  $W_{\mathcal{A}_{r,k}(n)}^\lambda$ .** In this section we view  $\mathcal{A}_{d,k}(n) \otimes \mathcal{A}_{r-d,\ell}(n)$  as a subalgebra of  $\mathcal{A}_{r,k+\ell}(n)$ . In order to define the embedding we need some notation suppose that  $\gamma$  is a set partition of a multiset containing only elements in  $\{k+1, k+2, \dots, k+\ell\} \cup \{\overline{k+1}, \overline{k+2}, \dots, \overline{k+\ell}\}$ . Then the *k-standardization* of  $\gamma$  is the multiset partition in  $\Pi_{r,\ell,n}$  obtained by subtracting  $k$  (resp.  $\overline{k}$ ) from all elements in  $\gamma$ .

The embedding of  $\mathcal{A}_{d,k}(n) \otimes \mathcal{A}_{r-d,\ell}(n)$  in  $\mathcal{A}_{r,k+\ell}(n)$  is defined by mapping the element  $O_\pi \otimes O_\gamma \in \mathcal{A}_{d,k}(n) \otimes \mathcal{A}_{r-d,\ell}(n)$ , where  $\pi \in \Pi_{d,k,n}$  and  $\gamma \in \Pi_{r-d,\ell,n}$ , to the sum of elements  $\sum_{\tau \in \Pi_{r,k+\ell,n}} O_\tau$  where the sum ranges over all  $\tau$  such that the restriction of  $\tau$  to  $[k] \cup \overline{[k]}$  is  $\pi$  and the  $k$ -standardization of  $\tau$  restricted to the set  $\{k+1, k+2, \dots, k+\ell\} \cup \{\overline{k+1}, \overline{k+2}, \dots, \overline{k+\ell}\}$  is  $\gamma$ .

**Example 5.9.** Consider the multiset partitions  $\pi = \{\{\{1\}\}, \{\{\overline{1}\}\}\}$  and  $\tau = \{\{\{1, \overline{1}\}\}\}$ , then  $O_\pi \otimes O_\tau \in \mathcal{A}_{1,1}(n) \otimes \mathcal{A}_{1,1}(n)$  as an element of  $\mathcal{A}_{2,1+1}(n)$  is equal to  $O_{\{\{\{1\}\}, \{\{\overline{1}\}\}, \{\{2, \overline{2}\}\}\}} + O_{\{\{\{1,2, \overline{2}\}\}, \{\{\overline{1}\}\}\}} + O_{\{\{\{1\}\}, \{\{2, \overline{1}, \overline{2}\}\}\}}.$

The next theorem shows that the multiplicity of a tensor of irreducible multiset partition algebra modules in the restriction from  $\mathcal{A}_{r,k+\ell}(n)$  to  $\mathcal{A}_{d,k}(n) \otimes \mathcal{A}_{r-d,\ell}(n)$  is given by the Kronecker coefficients. Recall that these coefficients are the multiplicities which appear in the tensor of irreducible symmetric group algebra modules. That is, if we denote  $g_{\lambda\nu\gamma} = \dim \text{Hom}(W_{\mathbb{C}S_n}^\lambda, W_{\mathbb{C}S_n}^\nu \otimes W_{\mathbb{C}S_n}^\gamma)$ , then we have that the decomposition of the restriction is given by the following theorem.

**Theorem 5.10.** *Let  $r, k, \ell$  and  $n$  be positive integers and  $\lambda$  be a partition of  $n$ , then the restriction of the irreducible  $W_{\mathcal{A}_{r,k+\ell}(n)}^\lambda$  to  $\bigoplus_{d=0}^r \mathcal{A}_{d,k}(n) \otimes \mathcal{A}_{r-d,\ell}(n) \subseteq \mathcal{A}_{r,k+\ell}(n)$  has the decomposition*

$$\text{Res}_{\bigoplus_{d=0}^r \mathcal{A}_{d,k}(n) \otimes \mathcal{A}_{r-d,\ell}(n)}^{\mathcal{A}_{r,k+\ell}(n)} W_{\mathcal{A}_{r,k+\ell}(n)}^\lambda \cong \bigoplus_{d=0}^r \bigoplus_{\nu \in \text{Par}_n} \bigoplus_{\gamma \in \text{Par}_n} (W_{\mathcal{A}_{d,k}(n)}^\nu \otimes W_{\mathcal{A}_{r-d,\ell}(n)}^\gamma)^{\oplus g_{\lambda\nu\gamma}}.$$

*Proof.* Recall from Equation (5.1) and for any  $d \geq 0$  that  $\mathcal{A}_{d,k}(n)$  is the centralizer of  $\mathbb{C}S_n$  acting on  $\mathcal{P}^d(V_{n,\{1,2,\dots,k\}})$ , so

$$(5.9) \quad \mathcal{P}^d(V_{n,\{1,2,\dots,k\}}) \cong \bigoplus_{\nu \in \text{Par}_n} W_{\mathbb{C}S_n}^\nu \otimes W_{\mathcal{A}_{d,k}(n)}^\nu.$$

Therefore as an  $\mathbb{C}S_n \otimes \mathcal{A}_{d,k}(n) \otimes \mathbb{C}S_n \otimes \mathcal{A}_{r-d,\ell}(n)$ -module that

$$\begin{aligned} & \mathcal{P}^d(V_{n,\{1,2,\dots,k\}}) \otimes \mathcal{P}^{r-d}(V_{n,\{k+1,k+2,\dots,k+\ell\}}) \\ & \cong \bigoplus_{\nu \in \text{Par}_n} \bigoplus_{\gamma \in \text{Par}_n} \left( W_{\mathbb{C}S_n}^\nu \otimes W_{\mathcal{A}_{d,k}(n)}^\nu \right) \otimes \left( W_{\mathbb{C}S_n}^\gamma \otimes W_{\mathcal{A}_{r-d,\ell}(n)}^\gamma \right). \end{aligned}$$

Now since

$$\mathcal{P}^r(V_{n,\{1,2,\dots,k+\ell\}}) \cong \bigoplus_{d=0}^r \mathcal{P}^d(V_{n,\{1,2,\dots,k\}}) \otimes \mathcal{P}^{r-d}(V_{n,\{k+1,k+2,\dots,k+\ell\}}),$$

then as a  $\bigoplus_{d=0}^r \mathcal{A}_{d,k}(n) \otimes \mathcal{A}_{r-d,\ell}(n)$ -module,

$$\begin{aligned} & \text{Res}_{\bigoplus_{d=0}^r \mathcal{A}_{d,k}(n) \otimes \mathcal{A}_{r-d,\ell}(n)}^{\mathcal{A}_{r,k+\ell}(n)} W_{\mathcal{A}_{r,k+\ell}(n)}^\lambda \\ & \cong \bigoplus_{d=0}^r \bigoplus_{\nu \in \text{Par}_n} \bigoplus_{\gamma \in \text{Par}_n} \text{Hom}(W_{\mathbb{C}S_n}^\lambda, W_{\mathbb{C}S_n}^\nu \otimes W_{\mathbb{C}S_n}^\gamma) \otimes W_{\mathcal{A}_{d,k}(n)}^\nu \otimes W_{\mathcal{A}_{r-d,\ell}(n)}^\gamma. \end{aligned}$$

The theorem follows since  $g_{\lambda\nu\gamma}$  is the dimension of  $\text{Hom}(W_{\mathbb{C}S_n}^\lambda, W_{\mathbb{C}S_n}^\nu \otimes W_{\mathbb{C}S_n}^\gamma)$ . □

REFERENCES

[Aik] A. C. Aitken, *On induced permutation matrices and the symmetric group*, Proc. Edinburgh Math. Soc. vol 5, issue 1, (1937), 1–13.

[Bend] E. Bender, *Partitions of multisets*, Discrete Math. 9, (1974) 301–311.

[BH1] G. Benkart, T. Halverson, *Partition algebras and the invariant theory of the symmetric group*, in *Recent trends in algebraic combinatorics*, Assoc. Women Math. Ser., Springer, 2019, Vol 16, pp. 1–41.

[BH2] *Partition algebras  $P_k(n)$  with  $2k > n$  and the fundamental theorems of invariant theory for the symmetric group  $S_n$* , J. Lond. Math. Soc. (2), 99, (2019), pp. 194–224.

[BHH] G. Benkart, T. Halverson, N. Harmon, *Dimensions of irreducible modules for partition algebras and tensor power multiplicities for symmetric and alternating groups*, J. Algebr. Comb. (2017) 46, pp. 77–108. <https://doi.org/10.1007/s10801-017-0748-4>.

[BDE] C. Bowman, M. De Visscher, and J. Enyang, *The co-Pieri rule for Kronecker coefficients*, arXiv:1710.04523.

[BDO] C. Bowman, M. De Visscher, R. Orellana, *The partition algebra and the Kronecker coefficients*, Trans. Amer. Math. Soc. 367 (2015), no. 5, 3647–3667.

[COSSZ] L. Colmenarejo, R. Orellana, F. Saliola, A. Schilling, M. Zabrocki, *An insertion algorithm on multiset partitions with applications to diagram algebras*, Journal of Algebra, Volume 557, 1 September 2020, Pages 97–128. <https://doi.org/10.1016/j.jalgebra.2020.04.010>

[E] J. Enyang, *A seminormal form for partition algebras*, J. Combin. Theory Ser. A **120** (2013), 1737–1785.

[F] F. G. Frobenius, *Über die Charaktere der symmetrischen Gruppe*, Sitzungsberichte der Königlich Preussischen Akademie Wissenschaften zu Berlin (1900) 516–34 (Ges. Abhandlungen, **3**, 148–66).

[Ge] I. Gessel, *Enumerative Applications of Symmetric Functions*, Publ. I.R.M.A. Strasbourg, 1987, 229/S-08 Actes 17e Séminaire Lotharingien, p. 5–21.

[GW1] R. Goodman, N. R. Wallach, *Representations and invariants of the classical groups*, Encyclopedia of Mathematics and its Applications, vol. 68, Cambridge University Press, Cambridge, 1998.

[GW2] R. Goodman, N. R. Wallach, *Symmetry, Representations, and Invariants*, Graduate Texts in Mathematics, Springer, 2009.

[Ha] T. Halverson, *Characters of the Partition Algebras*, J. Algebra, Volume 238, Issue 2, 15 April 2001, pp. 502–533.

[HJ] T. Halverson and T.N. Jacobson, *Set-partition tableaux and representations of diagram algebras*, Alg. Comb., Volume 3 (2020) no. 2, pp. 509–538. <https://doi.org/10.5802/alco.102>

[HL] T. Halverson, T. Lewandowski, *RSK Insertion for Set Partitions and Diagram Algebras*, Elect. J. Comb., Volume 11, Issue 2 (2004–6) (The Stanley Festschrift volume).

[HR] T. Halverson, A. Ram, *Partition Algebras*, Europ. J. of Combinatorics 26 (2005) pp. 869–921.

[Ho] R. Howe, *Perspectives on invariant theory: Schur duality, multiplicity-free actions and beyond*, The Schur lectures (1992) (Tel Aviv), Israel Math. Conf. Proc., vol. 8, Bar-Ilan Univ., Ramat Gan, 1995, pp. 1–182.

[J] V.F.R. Jones, *The Potts model and the symmetric group*, Subfactors: Proceedings of the Taniguchi Symposium on Operator Algebras, Kyuzeso, 1993. World Scientific, River Edge (1994) pp. 259–267.

- [Li] D. E. Littlewood, *Products and Plethysms of Characters with Orthogonal, Symplectic and Symmetric Groups*, *Canad. J. Math.*, **10**, 1958, 17–32.
- [LR] N. A. Loehr, J. B. Remmel, *A computational and combinatorial exposé of plethystic calculus*, *J. Alg. Comb.*, March 2011, Volume 33, Issue 2, pp. 163–198.
- [Mac] I. G. Macdonald, *Symmetric Functions and Hall Polynomials*, Second Edition, Oxford University Press, second edition, 1995.
- [Mar1] P. Martin, *Representations of graph Temperley-Lieb algebras*, *Publ. Res. Inst. Math. Sci.*, 26 (1990), pp. 485–503.
- [Mar2] P. Martin, *Potts Models and Related Problems in Statistical Mechanics*, World Scientific, Singapore (1991).
- [Mar3] P. Martin, *Temperley-Lieb algebras for nonplanar statistical mechanics—The partition algebra construction*, *J. Knot Theory Ramifications*, 3 (1994), pp. 51–82.
- [Mar4] P. Martin, *The structure of the partition algebras*, *J. Algebra*, 183 (1996), pp. 319–358.
- [ME] P. Martin, A. Elgamal, *The structure of the partition algebras*, *J. Algebra* **183** (1996), 319–358.
- [MR] P. Martin and G. Rollet, *The Potts model representation and a Robinson-Schensted correspondence for the partition algebra*, *Compositio Math.* 112 (1998), 237–254.
- [NPS] S. Narayanan, D. Paul, S. Srivastava, *The Multiset Partition Algebra*, [arXiv:1903.10809](https://arxiv.org/abs/1903.10809).
- [OZ] R. Orellana, M. Zabrocki, *Symmetric group characters as symmetric functions*, [arXiv:1605.06672](https://arxiv.org/abs/1605.06672).
- [OZ2] R. Orellana, M. Zabrocki, *Products of characters of the symmetric group*, *Journal of Combinatorial Theory, Series A*, Volume 165, July 2019, Pages 299–324.
- [OZ3] R. Orellana, M. Zabrocki, *The Hopf structure of symmetric group characters as symmetric functions*, [arXiv:1901.00378](https://arxiv.org/abs/1901.00378).
- [OZ4] R. Orellana, M. Zabrocki, *A combinatorial model for the decomposition of multivariate polynomial rings as  $S_n$ -modules*, *Elect. J. Combin.*, to appear, [arXiv:1906.01125](https://arxiv.org/abs/1906.01125).
- [OEIS] N. J. A. Sloane, *The On-Line Encyclopedia of Integer Sequences*. Published electronically at <http://oeis.org>, 2013.
- [Pa] D. Paul, *The Multiset Partition Algebra*, Ph.D. Thesis, The Institute of Mathematical Sciences, Chennai, 2020.
- [Pr] C. Procesi, *Lie Groups: An approach through invariants and representations*, Springer, 2007.
- [sage] W. A. Stein et al. *Sage Mathematics Software (Version 6.10)*, The Sage Development Team, 2016, <http://www.sagemath.org>.
- [sage-combinat] The Sage-Combinat community. *Sage-Combinat: enhancing Sage as a toolbox for computer exploration in algebraic combinatorics*, <http://combinat.sagemath.org>, 2008.
- [ST] T. Scharf, J. Y. Thibon, *A Hopf-algebra approach to inner plethysm*. *Adv. in Math.* 104 (1994), pp. 30–58.
- [Sta] R. Stanley, *Enumerative Combinatorics, Vol. 2*, Cambridge University Press, 1999.

DARTMOUTH COLLEGE, MATHEMATICS DEPARTMENT, HANOVER, NH 03755, USA  
*E-mail address:* [rosa.c.orellana@dartmouth.edu](mailto:rosa.c.orellana@dartmouth.edu)

DEPARTMENT OF MATHEMATICS AND STATISTICS, YORK UNIVERSITY, TORONTO, ONTARIO M3J 1P3, CANADA  
*E-mail address:* [zabrocki@mathstat.yorku.ca](mailto:zabrocki@mathstat.yorku.ca)