CUBIC-QUINTIC NLS: SCATTERING BEYOND THE VIRIAL THRESHOLD

ROWAN KILLIP, JASON MURPHY, AND MONICA VISAN

ABSTRACT. We consider the nonlinear Schrödinger equation in three space dimensions with combined focusing cubic and defocusing quintic nonlinearity. This problem was considered previously by Killip, Oh, Pocovnicu, and Visan, who proved scattering for the whole region of the mass/energy plane where the virial quantity is guaranteed to be positive. In this paper we prove scattering in a larger region where the virial quantity is no longer guaranteed to be sign definite.

1. INTRODUCTION

We consider the cubic-quintic NLS in three space dimensions:

$$\begin{cases} i\partial_t u + \Delta u = -|u|^2 u + |u|^4 u, \\ u|_{t=0} = u_0 \in H^1(\mathbb{R}^3), \end{cases}$$
(1.1)

which describes the evolution of a complex field u under the Hamiltonian

$$E(u) = \int_{\mathbb{R}^3} \frac{1}{2} |\nabla u(x)|^2 + \frac{1}{6} |u(x)|^6 - \frac{1}{4} |u(x)|^4 \, dx.$$

This evolution also conserves the mass, defined by

$$M(u) = \int_{\mathbb{R}^3} |u(x)|^2 \, dx.$$

We work here in $H^1(\mathbb{R}^3)$ because this is precisely the class of initial data for which both the mass and energy are finite. The initial value problem (1.1) was shown to be globally well-posed for such data by Zhang in [18], building on the paper [3] that treated the purely quintic nonlinearity.

We are concerned here with the long-time behavior of solutions. For small initial data, standard arguments demonstrate that solutions scatter both forward and backward in time. This means that there are functions $u_{\pm} \in H^1$ so that

$$\lim_{t \to +\infty} \|u(t) - e^{it\Delta}u_{\pm}\|_{H^1} = 0.$$
(1.2)

On the other hand, our equation admits solitary wave solutions which evidently do not scatter.

The natural question then arises of determining the sharp threshold at which scattering breaks down. There is already a large body of work on determining such thresholds for a wide variety of models, building on the paradigm introduced in [11]; see [1, 2, 4–8, 10, 13, 15], for example. This approach has two main steps: First one shows that the threshold for scattering is witnessed by a minimal counterexample that (by virtue of its minimality) is almost-periodic (modulo symmetries). One

then uses the virial identity (or close analogue) to prove that such almost periodic solutions cannot exist for initial data below the soliton threshold.

The virial identity for (1.1) takes the following form:

$$\frac{d}{dt} \left\langle u(t), \frac{1}{4i} (x \cdot \nabla + \nabla \cdot x) u(t) \right\rangle = V(u(t)),$$

$$V(u) = \int |\nabla u(x)|^2 + |u(x)|^6 - \frac{3}{4} |u(x)|^4 \, dx.$$
(1.3)

where

Many of the results of [12] are most readily framed with reference to Figure 1. This mass-energy diagram is purely schematic; while precise numerics are presented in [12], the salient features live at very different length scales and so cannot be represented intelligibly on a single graph.



FIGURE 1. Schematic depiction of the mass-energy plane showing soliton solutions (heavy curve), non-soliton virial obstruction (light curve), and the scattering region \mathcal{R} from [12].

The heavy line in Figure 1 traces the mass-energy curve of ground-state soliton solutions. By ground-state soliton, we mean an H^1 -solution P_{ω} to

$$-\Delta P_{\omega} + |P_{\omega}|^4 P_{\omega} - |P|^2 P_{\omega} + \omega P_{\omega} = 0$$

that is radially symmetric decreasing. Such solutions exist if and only if $0 < \omega < \frac{3}{16}$ and are unique for each such ω ; see [12].

This mass-energy curve $\omega \mapsto (M(P_{\omega}), E(P_{\omega}))$ continues beyond the edges of Figure 1. The convex branch is asymptotic to the M axis, while the concave branch continues toward infinite mass and infinitely negative energy. Note that we have only plotted the positive energy part of the mass-energy plane, since it is readily shown that no scattering solution can have negative energy. We write m_2 for the point were the curve of solitons crosses the mass axis. It is shown in [12] that for masses less than m_2 , all solutions (other than zero) have positive energy. While individual solutions with mass larger than m_2 may scatter, scattering cannot be guaranteed for any mass-energy pair in this regime.

The mass m_0 is defined as the least mass at which it is possible to have V(u) = 0 without $u \equiv 0$. The first big surprise is that m_0 is strictly smaller than the smallest mass of any soliton (which coincides with the cusp in the heavy curve). This was proved rigorously in [12]. It is also shown there that $27m_0^2 = 16m_2^2$.

This leads us to the natural question: What are these minimal objects that achieve zero virial? The answer, discovered in [12], is that they are exotic rescalings of soliton solutions in the sense that the rescaling preserves neither mass nor energy. (Due to the combined nonlinearity, equation (1.1) has no scaling symmetry.) These rescaled solitons are defined by the equation

$$R_{\omega}(x) := \sqrt{\frac{1+\beta(\omega)}{4\beta(\omega)}} P_{\omega}\left(\frac{3(1+\beta(\omega))}{4\sqrt{3\beta(\omega)}}x\right), \quad \text{where} \quad \beta(\omega) := \frac{\|P_{\omega}\|_{L^{6}}^{6}}{\|\nabla P_{\omega}\|_{L^{2}}^{2}}.$$
(1.4)

The mass-energy curve of these rescaled solitons is shown as the light weight curve in Figure 1.

Consider now the open region \mathcal{R} shown in Figure 1. The boundary of \mathcal{R} , which is not included in \mathcal{R} , is comprised partly of the mass-energy curve of solitons (heavy curve), partly by the curve of rescaled solitons (light curve), and partly by the mass m_0 (dotted line).

In [12], it is shown that V(u) > 0 for every state u whose mass and energy lies in the region \mathcal{R} . In fact, the region \mathcal{R} is the maximal region with this property in the following sense: Defining the function

$$E_{\min}^{V}(m) = \inf\{E(u) : u \in H^{1}(\mathbb{R}^{3}), \ M(u) = m, \text{ and } V(u) = 0\},$$
(1.5)

we have $\mathcal{R} = \{(m, e) : e < E_V(m)\}$. Note that $E_{\min}^V(m) = \infty$ for $m < m_0$.

In this paper, we prove that scattering holds in a still larger region \mathcal{B} of the mass-energy plane:

Theorem 1.1. There is an open region $\mathcal{B} \subseteq \mathbb{R}^2$ so that every solution u to (1.1) with $(M(u), E(u)) \in \mathcal{B}$ belongs to $L^{10}(\mathbb{R} \times \mathbb{R}^3)$ and so scatters (in both time directions) in the sense of (1.2). The region \mathcal{B} is larger than \mathcal{R} : (a) There is an $m > m_0$ so that $\mathcal{B} \supseteq (0, m) \times (0, \infty)$; and

(b) If $(m, e) \in \partial \mathcal{R}$ is not achieved by a soliton, then $(m, e) \in \mathcal{B}$.

With reference to Figure 1, we see from (a) that the whole dotted line on the boundary may be moved to the right (remaining vertical!). Part (b) then completes our promise to expand the region of scattering in every place where this is not rigorously forbidden by the existence of solitons (which evidently do not scatter). That is, we extend the scattering region across the entire portion of $\partial \mathcal{R}$ comprised by the light-weight curve.

In Figure 1 we see that for $m \geq m_0$, the left portion of $\partial \mathcal{R}$ is delimited by rescaled solitons and the right portion by solitons. The mass at which this transition takes place is marked m_1 . The validity of this description was shown in [12] via numerics. The purely analytical arguments given there show only that the left-most portion is made up of rescaled solitons and that the right-most part is made up solely of solitons; the possibility that roles are exchanged multiple times is not excluded. The scattering claim in Theorem 1.1 follows from a more quantitative assertion that we will prove, namely, that the spacetime bound

$$\|u\|_{L^{10}_{t,\sigma}(\mathbb{R}\times\mathbb{R}^3)} \le C(M(u), E(u)) \tag{1.6}$$

holds for every solution u with $(M(u), E(u)) \in \mathcal{B}$. Here C is some unspecified function. The same type of bound was proven in [12] for $(m, e) \in \mathcal{R}$ and indeed for other equations in many papers preceding it. One novelty of [12] is the appearance of two parameters (both mass and energy) in RHS(1.6). It is worth emphasizing that due to the concavity of the soliton portion of $\partial \mathcal{R}$, the arguments in [12] cannot be based on a free energy of the form $E + \lambda M$ as had proven successful for a number of similar problems. The need to induct on two parameters leads us to consider the following partial order on mass-energy pairs:

Definition 1.2. We write $(m, e) \preccurlyeq (m_0, e_0)$ to indicate that $m \le m_0$ and $e \le e_0$. That is, (m, e) is southwest of (m_0, e_0) in the mass-energy plane.

With these preliminaries set, we can establish the following proposition. As we will explain, this is essentially a recapitulation of the concentration compactness analysis in [12].

Proposition 1.3. Fix $0 < m < m_2$ and e > 0. Then exactly one of the two following possibilities hold:

(i) There exists $\varepsilon > 0$ so that

 $\|u\|_{L^{10}_{t,r}(\mathbb{R}\times\mathbb{R}^3)} \le \varepsilon^{-1} \quad whenever \quad (M(u), E(u)) \preccurlyeq (m+\varepsilon, e+\varepsilon).$ (1.7)

(ii) There is a global solution $u \in C_t(\mathbb{R}; H^1)$ to (1.1) with $(M(u), E(u)) \preccurlyeq (m, e)$. Moreover, for this solution there is a function $c : \mathbb{R} \to \mathbb{R}^3$ so that

 $\{u(t, x-c(t)): t \in \mathbb{R}\}\$ is precompact in $H^1(\mathbb{R}^3)$ and c(t) = o(t) as $t \to \pm \infty$. (1.8)

Proof. If (i) holds at the original pair (m, e), then the matter is settled: Scattering holds and so (1.8) cannot.

Henceforth, we assume that (i) does not hold for any $\varepsilon > 0$ and must prove (ii). For this purpose, we may safely replace (m, e) with any of its \preccurlyeq -predecessors. We choose (m, e) to be a \preccurlyeq -minimal pair for which (i) fails.

As (i) fails at (m, e), there must be a sequence of solutions $u_n(t)$ to (1.1) with

$$M(u_n) \to m$$
, $E(u_n) \to e$, and $\|u_n\|_{L^{10}_{t,r}(\mathbb{R} \times \mathbb{R}^3)} \to \infty$.

Applying [12, Proposition 9.1] exactly as in the proof of [12, Theorem 9.6] we obtain the existence of a solution u satisfying the precompactness claim in (1.8). This compactness relies on the minimality of (m, e). That the spatial center function c(t) must be o(t) also follows from this minimality; see [12, Proposition 10.2]. \Box

We are now ready to discuss how we are to prove Theorem 1.1. The argument has two main phases. In phase one, we prove that a point in $\partial \mathcal{R}$ can only support an almost periodic solution (that is, one satisfying (1.8)) if it supports a solitary wave solution. It then follows from Proposition 1.3 that scattering extends to a small neighborhood of any point $(m, e) \in \partial \mathcal{R}$ that is not achieved by a soliton. The key observation driving the phase-one argument is that while individual functions with mass-energy belonging to $\partial \mathcal{R}$ may have vanishing virial, the only trajectories that maintain zero virial are the soliton solutions. Actually, this is not quite enough: In order to preclude almost periodic solutions we need a quantitative lower bound.

CUBIC-QUINTIC NLS

Evidently, this cannot hold pointwise in time; nevertheless, we will be able to prove that such a bound does hold for the integral over fixed-size time intervals.

While phase one provides a resulting scattering region that fulfills part (b) of Theorem 1.1, there is no reason to believe it satisfies (a). However, it does show that given E_* , there is an $m = m(E_*) > m_0$ so that scattering holds in the rectangle $(0, m) \times (0, E_*)$. To complete phase two of the proof, we show that there is a suitable choice of E_* that allows us to employ a different argument to prove that scattering extends to the entire strip $(0, m) \times (0, \infty)$.

Acknowledgments

R. K. was supported by NSF grant DMS-1856755. J. M. was supported by a Simons Collaboration Grant. M. V. was supported by NSF grant DMS-1763074.

2. Preliminaries

In order to prove scattering, we must preclude solutions to (1.1) with the property (1.8). Like many of our predecessors, we will do this using a localized version of the virial identity. The novel aspect of this paper is *how* we falsify (2.1).

Proposition 2.1 (The localized virial argument). If u is a solution to (1.1) satisfying (1.8), then

$$\lim_{T \to \infty} \frac{1}{T} \int_0^T V(u(t)) \, dt = 0, \tag{2.1}$$

where V(u) is as in (1.3).

Proof. This is the essence of the standard localized virial argument; details can be found in [12, p. 542–543]. The only difference is that there the argument is by contradiction since it is already known that for the solutions of interest, V(u(t)) is bounded away from zero (uniformly for $t \in \mathbb{R}$).

In order to upgrade certain qualitative statements from [12] to quantitative ones, we employ the following from [9]:

Proposition 2.2 (Profile decomposition for Gagliardo–Nirenberg). Let $\{u_n\}$ be a bounded sequence in $H^1(\mathbb{R}^3)$. Passing to a subsequence if necessary, there exist $J^* \in \{0, 1, 2, ...\} \cup \{\infty\}$, profiles $\phi^j \in H^1 \setminus \{0\}$, and positions $\{x_n^j\} \subset \mathbb{R}^3$, so that

$$u_n(x) = \sum_{j=1}^{J} \phi^j(x - x_n^j) + w_n^J(x)$$
(2.2)

for each finite $0 \leq J \leq J^*$. Moreover, the following hold:

$$\begin{split} &\lim_{J \to J^*} \limsup_{n \to \infty} \|w_n^J\|_{L^4} = 0, \\ &\sup_J \limsup_{n \to \infty} \lim_{n \to \infty} \left| \|u_n\|_{\dot{H}^1}^2 - \sum_{j=1}^J \|\phi^j\|_{\dot{H}^1}^2 - \|w_n^J\|_{\dot{H}^1}^2 \right| = 0, \\ &\sup_J \limsup_{n \to \infty} \lim_{n \to \infty} \left| \|u_n\|_{L^q}^q - \sum_{j=1}^J \|\phi^j\|_{L^q}^q - \|w_n^J\|_{L^q}^q \right| = 0 \quad for \quad q \in \{2, 4, 6\}. \end{split}$$

Recall that m_0 denotes the least (non-zero) mass at which V(u) = 0 is possible. This quantity is determined through a variational problem analyzed in [12], which will be important for analyzing the vertical portion of $\partial \mathcal{R}$.

Proposition 2.3 (Interpolation inequality). Every $u \in H^1(\mathbb{R}^3)$ satisfies

$$\frac{3}{4} \int |u|^4 \, dx \le \sqrt{\frac{M(u)}{m_0}} \int |\nabla u|^2 + |u|^6 \, dx. \tag{2.3}$$

Moreover, there is a non-empty finite set $\Omega \subset (0, \frac{3}{16})$ which characterizes optimizers thus: $M(u) = m_0$ and equality holds in (2.3) if and only if $u(x) = e^{i\theta}R_{\omega}(x-c)$ for some $c \in \mathbb{R}^3$, $\theta \in \mathbb{R}$, and $\omega \in \Omega$.

Remark 2.4. The numerics in [12] show that Ω is actually comprised of a single point; this would also be a consequence of Conjecture 2.6 of that paper. This is what is depicted in Figure 1. While the numerics are very stable and compelling, we do not have a rigorous proof that there is only one value of ω so that $M(R_{\omega}) = m_0$. Thus, we will not make this assumption in what follows.

Proof. Choosing $\alpha = 1$ in [12, Proposition 3.1] shows that the optimal constant in the inequality

$$\|u\|_{L^4}^4 \le C \|u\|_{L^2} \|\nabla u\|_{L^2}^{\frac{3}{2}} \|u\|_{L^6}^{\frac{3}{2}}$$
(2.4)

is realized. Moreover, every optimizer takes the form $u(x) = e^{i\theta} \lambda R_{\omega}(\rho(x-c))$ with $\theta \in \mathbb{R}, \lambda > 0, \rho > 0, c \in \mathbb{R}^3$, and $\beta(\omega) = 1$. In [12], this is written with P_{ω} , rather than R_{ω} , but these assertions are equivalent in view of (1.4).

The value of m_0 was determined in [12] from the optimal constant C in (2.4) via

$$m_0 = \sqrt{3} \left[\frac{16}{9C}\right]^2 \tag{2.5}$$

for details, see [12, Lemma 3.3 and Theorem 5.2].

From [12, Theorem 2.2] we see that β is an analytic function of $\omega \in (0, \frac{3}{16})$ and that $\beta(\omega) \to 0$ as $\omega \to 0$ and $\beta(\omega) \to \infty$ as $\omega \to \frac{3}{16}$. Thus choosing Ω to denote the set of all ω that arise as optimizers, we are guaranteed that Ω is a finite set. Note that, at this time, we cannot claim that every solution to $\beta(\omega) = 1$ corresponds to an optimizer.

The inequality (2.3) follows from and (2.4) via Young's inequality in the form

 $3^{1/4}ab \le \frac{3}{4}[a^{4/3} + b^4]$ with equality iff $a^{4/3} = 3b^4$.

This requirement for equality places one constraint on the scaling parameters ρ and λ . Combining this with the requirement that $M(u) = m_0$ then guarantees that $\lambda = 1$ and $\rho = 1$ as stated in the proposition. Details of these computations can also be found in [12, Lemma 5.5].

3. Proof of Theorem 1.1

Let us write $\partial \mathcal{R}_s$ to represent the portion of $\partial \mathcal{R}$ represented by solitary waves:

$$\partial \mathcal{R}_s := \left\{ (m, e) \in \partial \mathcal{R} : (m, e) = \left(M(P_\omega), E(P_\omega) \right) \text{ for some } \omega \in (0, \frac{3}{16}) \right\}.$$

The remainder of the boundary will be denoted $\partial \mathcal{R}_r$. Our first goal in this section is to prove Proposition 3.4, which shows that scattering holds in a neighborhood of any mass-energy pair $(m, e) \in \partial \mathcal{R}_r$.

Before we can begin on the novel portion of the analysis, we need one more lemma recapitulating material developed in [12]:

CUBIC-QUINTIC NLS

Lemma 3.1. Fix $(m, e) \in \partial \mathcal{R}_r$ and let $u \in H^1$ satisfy $(M(u), E(u)) \preccurlyeq (m, e)$. Then $V(u) \ge 0$. Moreover, if V(u) = 0, then either $u \equiv 0$ or $u(x) = e^{i\theta} \mathcal{R}_{\omega}(x-c)$ for some $c \in \mathbb{R}^3$ and $\theta \in [0, 2\pi)$. The value of ω is uniquely determined by E(u).

Proof. When $m = m_0$, both $V(u) \ge 0$ and the identification of cases of equality follow from Proposition 2.3. When $m > m_0$, we rely instead on [12, Theorem 5.6], which identifies optimizers for (1.5) at fixed mass, and [12, Theorem 5.2], which shows strict monotonicity of $m \mapsto E_{\min}^V(m)$.

Finally, the monotonicity [12, Equation (5.19)] guarantees that $E(R_{\omega})$ uniquely determines ω .

For our purposes, it is not enough merely to understand which functions achieve zero virial. Rather, we need to understand minimizing sequences:

Lemma 3.2. Fix $(m, e) \in \partial \mathcal{R}_r$ and let u_n be a sequence in $H^1(\mathbb{R}^3)$ with $M(u_n) = m$ and $E(u_n) = e$. If $V(u_n) \to 0$ then there are $x_n \in \mathbb{R}^3$ and $\theta_n \in [0, 2\pi)$ so that

$$e^{i\theta_n}u_n(x-x_n) \longrightarrow R_{\omega}(x) \quad in \ H^1(\mathbb{R}^3),$$
(3.1)

where ω is uniquely determined by $E(R_{\omega}) = e$.

Proof. We apply the profile decomposition of Proposition 2.2 to the sequence u_n . The remaining analysis will be confined to the resulting subsequence where (2.2) holds. This suffices, for if the claim were false, there would be a sequence u_n without any subsequence satisfying (3.1).

From the conclusions of Proposition 2.2, we know that the profiles satisfy

$$\limsup_{J \to J^*} \limsup_{n \to \infty} \left[M(w_n^J) + \sum_{j=1}^J M(\phi^j) \right] = m, \qquad \sum_{j=1}^{J^*} E(\phi^j) \le e, \tag{3.2}$$

and moreover,

$$0 = \limsup_{J \to J^*} \limsup_{n \to \infty} \left[\|\nabla w_n^J\|_{L^2}^2 + \|w_n^J\|_{L^6}^6 + \sum_{j=1}^J V(\phi_j) \right].$$
(3.3)

From these relations and Lemma 3.1, we see that any ϕ^j (which are always non-zero) must agree with some (a priori *j*-dependent) rescaled soliton R_{ω} up to a translation and phase rotation. Modifying the symmetry parameters in (2.2), if necessary, we may assume that the profile is exactly R_{ω} . Lemma 3.1 and (3.3) also show that $\nabla w_n^J \to 0$ in L^2 .

In view of the mass and energy constraints in (3.2) and the monotonicity of $m \mapsto E_{\min}^V(m)$ for $m \ge m_0$, we are left with two possibilities: Either there is exactly one profile $\phi^1 = R_{\omega}$ and $w_n \to 0$ in L^2 -sense, or $u_n = w_n$ for all n. The former case yields (3.1). The latter case is incompatible with $V(u_n) \to 0$, because $w_n \to 0$ in L^4 and so $\liminf_{n\to\infty} V(w_n) \ge E(w_n) \equiv e$.

Recalling that our ambition is to contradict (2.1) (and thereby preclude solutions obeying (1.8)), we can now see how Lemma 3.2 helps: It shows that the virial will remain bounded away from zero unless the trajectory passes near R_{ω} . To handle the remaining case, we introduce the notation

$$\rho_{\omega}(u) := \inf \left\{ \left\| u(x) - e^{i\theta} R_{\omega}(x-c) \right\|_{H^1} : c \in \mathbb{R}^3, \ \theta \in \mathbb{R} \right\}$$
(3.4)

and prove the following:

Lemma 3.3. Fix $\omega \in (0, \frac{3}{16})$. Then there exists $\delta > 0$ so that every solution u to (1.1) with $\rho_{\omega}(u(0)) \leq \delta$ satisfies

$$\int_{-1}^{1} V(u(t)) dt \ge \delta. \tag{3.5}$$

Proof. The crux of the proof is the following: If v is the solution to (1.1) with initial data $v(0) = R_{\omega}$, then

$$\int_{-1}^{1} V(v(t)) \, dt > 0. \tag{3.6}$$

We will prove this shortly. As V is continuous on H^1 , it follows from this assertion and local well-posedness that there is a $\delta > 0$ so that (3.5) holds for every solution with $||u(0) - R_{\omega}||_{H^1} \leq \delta$. This extends to solutions with $\rho_{\omega}(u) \leq \delta$ due to the gauge and translation symmetries of (1.1).

We now turn our attention to (3.6), arguing by contradiction. From Lemma 3.2, we know $V(v(t)) \ge 0$ and thus the failure of (3.6) ensures that $V(v(t)) \equiv 0$ for $t \in [-1, 1]$. But this in turn guarantees

$$v(t,x) = e^{i\theta(t)} R_{\omega}(x - x(t))$$

for some (continuous) functions $\theta(t)$ and x(t). In view of the uniqueness of H^1 solutions to (1.1), the rotational symmetry of (1.1) and the initial data v(0) guarantee that $x(t) \equiv 0$. Analogously, combining gauge symmetry and time-translation invariance guarantee that $\theta(t+s) = \theta(t) + \theta(s)$ whenever $s, t, s+t \in [-1,1]$. This in turn shows that $\theta(t) = \xi t$ for some $\xi \in \mathbb{R}$ and consequently that $v(t, x) = e^{i\xi t}R_{\omega}(x)$ for $t \in [-1,1]$. By gauge invariance this shows v to be a solitary wave solution for all $t \in \mathbb{R}$, which is inconsistent with the assumption $(M(R_{\omega}), E(R_{\omega})) \in \partial \mathcal{R}_r$. \Box

With Lemmas 3.2 and 3.3 in place, we are now ready to demonstrate the existence of the new, larger scattering region that satisfies property (b) listed in Theorem 1.1, completing phase one of our argument. The region will be further enlarged in Proposition 3.5 so that property (a) also holds.

Proposition 3.4. Given any pair $(m, e) \in \partial \mathcal{R}_r$, there exist $\varepsilon > 0$ and $C < \infty$ so that for any solution u to (1.1),

 $(M(u), E(u)) \preccurlyeq (m + \varepsilon, e + \varepsilon) \quad implies \quad ||u||_{L^{10}(\mathbb{R} \times \mathbb{R}^3)} \le C.$

Proof. In view of Proposition 1.3, it suffices to show that there are no solutions u satisfying (1.8) with $(M(u), E(u)) \preccurlyeq (m, e)$. This in turn will be effected by falsifying (2.1).

Suppose that u is a solution satisfying (1.8) and $(M(u), E(u)) \preccurlyeq (m, e)$. By the main theorem from [12], we must have $(M(u), E(u)) \in \partial \mathcal{R}_r$. We first consider the case that this mass-energy pair does *not* coincide with that of any R_{ω} . This assumption combined with Lemma 3.2 shows that $V(u(t_n)) \rightarrow 0$ is impossible for any sequence of times t_n . Thus it follows that

$$V(u(t)) \gtrsim 1$$
 uniformly for $t \in \mathbb{R}$, (3.7)

which clearly contradicts (2.1).

Suppose now that the mass and energy of u do coincide with those of a (necessarily unique) rescaled solitary wave R_{ω} . Let $\delta > 0$ be that given by Lemma 3.3 for this value of ω . By Lemma 3.2, there exists $\varepsilon > 0$ so that

$$V(u(t)) \ge \varepsilon$$
 whenever $\rho_{\omega}(u(t)) \ge \delta.$ (3.8)

Now given T > 0 large, let $K_T = \{t \in [1, T - 1] : \rho_{\omega}(u(t)) \ge \delta\}$. The (trivial) onedimensional Besicovich covering lemma shows that one can find a finite collection of times $t_i \in K_T$ so that

$$K_T \subseteq K_T^* := \bigcup \left[t_i - 1, t_i + 1 \right]$$

and no $t \in \mathbb{R}$ belongs to more than two intervals $[t_i - 1, t_i + 1]$. Applying (3.5) to each such interval, (3.8) on $[1, T - 1] \setminus K_T^*$, and neglecting the positive contribution of any points not included, we find

$$\int_0^T V(u(t)) dt \ge \frac{1}{2} \delta |K_T^*| + \varepsilon \big| [1, T-1] \setminus K_T^* \big|.$$

As neither δ nor ε depend on T, this conclusion is inconsistent with (2.1) and so completes the proof of the proposition in the case that the pair $(m, e) \in \partial \mathcal{R}_r$ coincides with that of some rescaled soliton.

Proposition 3.5. There exists $m > m_0$ and a function $C : [0, \infty) \to \mathbb{R}$ so that any solution u to (1.1) with $M(u) \le m$ satisfies

$$\|u\|_{L^{10}_{t,\sigma}(\mathbb{R}\times\mathbb{R}^3)} \le C(E(u)). \tag{3.9}$$

Proof. Proposition 3.4 together with a simple covering argument shows that for any choice of E_* , there is an $m > m_0$ so that (3.9) holds for all solutions with $M(u) \le m$ and $E(u) \le E_*$.

We choose $E_* = m_2$, which is guaranteed to be larger than the corresponding m due to the existence of solitary waves. It follows that if $M(u) \leq m$ and $E(u) \geq E_*$, then

$$V(u) = 2E(u) - \frac{3}{64}M(u) + \frac{1}{3}\int |u|^2 \left[|u|^2 - \frac{3}{8}\right]^2 dx \ge m_2.$$

It is evident from Proposition 2.1 that no solution to (1.1) can satisfy both these mass-energy constraints and (1.8). On the other hand, m was chosen so that no solutions can satisfy (1.8) with $(M(u), E(u)) \preccurlyeq (m, E_*)$. This proves the proposition, since Proposition 1.3 shows that the failure of (3.9) would produce exactly the type of solution that we have just precluded.

References

- T. Akahori, S. Ibrahim, H. Kikuchi, and H. Nawa, Existence of a ground state and scattering for a nonlinear Schrödinger equation with critical growth. Selecta Math. (N.S.) 19 (2013), no. 2, 545–609.
- [2] T. Akahori and H. Nawa, Blowup and scattering problems for the nonlinear Schrödinger equations. Kyoto J. Math. 53 (2013), no. 3, 629–672.
- [3] J. Colliander, M. Keel, G. Staffilani, H. Takaoka, and T. Tao, Global well-posedness and scattering for the energy-critical nonlinear Schrödinger equation in ℝ³. Ann. of Math. (2) 167 (2008), no. 3, 767–865.
- [4] B. Dodson, Global well-posedness and scattering for the focusing, cubic Schrödinger equation in dimension d = 4. Ann. Sci. Ec. Norm. Supér. (4) 52 (2019), no. 1, 139–180.
- [5] B. Dodson, Global well-posedness and scattering for the mass critical nonlinear Schrödinger equation with mass below the mass of the ground state. Adv. Math. 285 (2015), 1589–1618.
- [6] T. Duyckaerts, J. Holmer, and S. Roudenko, Scattering for the non-radial 3D cubic nonlinear Schrödinger equation. Math. Res. Lett. 15 (2008), no. 6, 1233–1250.
- [7] D. Fang, J. Xie, and T. Cazenave, Scattering for the focusing energy-subcritical nonlinear Schrödinger equation. Sci. China. Math. 54 (2011), no. 10, 2037–2062.
- [8] C. Guevara, Global behavior of finite energy solutions to the d-dimensional focusing nonlinear Schrödinger equation. Appl. Math. Res. Express. AMRX 2014, no. 2, 177–243.

- [9] T. Hmidi and S. Keraani, Blowup theory for the critical nonlinear Schrödinger equations revisited. Int. Math. Res. Not. 2005, 2815–2828.
- [10] J. Holmer and S. Roudenko, A sharp condition for scattering of the radial 3d cubic nonlinear Schrödinger equation. Comm. Math. Phys. 282 (2008), 435–467.
- [11] C. E. Kenig and F. Merle, Global well-posedness, scattering and blowup for the energy-critical focusing non-linear wave equation. Acta. Math. 201 (2008), no. 2, 147–212.
- [12] R. Killip, T. Oh, O. Pocovnicu, and M. Visan, Solitons and scattering for the cubic-quintic nonlinear Schrödinger equation on ℝ³. Arch. Ration. Mech. Anal. 225 (2017), no. 1, 469–548.
- [13] R. Killip and. M. Visan, The focusing energy-critical nonlinear Schrödinger equation in dimensions five and higher. Amer. J. Math. 132 (2010), no. 2, 361–424.
- [14] R. Killip and M. Visan, Nonlinear Schrödinger equations at critical regularity. Evolution equations, 325–437, Clay Math. Proc., 17, Amer. Math. Soc., Providence, RI, 2013.
- [15] C. Miao, G. Xu, and L. Zhao, The dynamics of the 3D radial NLS with the combined terms. Comm. Math. Phys. 318 (2013), no. 3, 767–808.
- [16] K. Nakanishi and W. Schlag, Global dynamics above the ground state energy for the cubic NLS equation in 3D. Calc. Var. Partial Differential Equations 44 (2012), no. 1-2, 1-45.
- [17] T. Tao, M. Visan, and X. Zhang, The nonlinear Schrödinger equation with combined powertype nonlinearities. Comm. Partial Differential Equations 32 (2007), no. 7-9, 1281–1343.
- [18] X. Zhang, On the Cauchy problem of 3 D energy-critical Schrödinger equations with subcritical perturbations. J. Differ. Equ. 230 (2006), no. 2, 422–445.

DEPARTMENT OF MATHEMATICS, UCLA *E-mail address*: killip@math.ucla.edu

DEPARTMENT OF MATHEMATICS & STATISTICS, MISSOURI S&T *E-mail address*: jason.murphy@mst.edu

DEPARTMENT OF MATHEMATICS, UCLA *E-mail address*: visan@math.ucla.edu