

SHEAF THEORETIC COMPACTIFICATIONS OF THE SPACE OF RATIONAL QUARTIC PLANE CURVES

KIRYONG CHUNG

ABSTRACT. Let R_4 be the space of rational plane curves of degree 4. In this paper, we obtain a sheaf theoretic compactification of R_4 via the space of α -semistable pairs on \mathbb{P}^2 and its birational relations through wall-crossings of semistable pairs. We obtain the Poincaré polynomial of the compactified space.

1. INTRODUCTION

1.1. Compactifications and results. Let R_d be the space of *irreducible* rational plane curves of degree d . The space R_d is an irreducible variety of dimension $3d - 1$ ([KP01]). One of very famous compactification of R_d is so called the moduli space of stable maps, which is a key role in the curve counting theory. For the detailed description, see [FP97]. This compactification is motivated by viewing the rational plane curve with its unique normalization map (up to isomorphism). In this paper, our aim is to obtain a sheaf-theoretic compactification of R_d and its birational relations. In this case, the curve is regarded as the pair of the direct image sheaf of the normalization map and its canonical section. In detail,

- (1) let $M(dm + 1)$ be the moduli space of stable sheaves F with Hilbert polynomial $\chi(F(m)) = dm + 1$ and
- (2) $M^\alpha(dm + 1)$ be the moduli space of α -semistable pairs (s, F) with Hilbert polynomial $\chi(F(m)) = dm + 1$.

For explicit definition and properties, see Section 2. Note that all of these moduli spaces are projective scheme ([Sim94, He98, LP93]). For the irreducible plane curve C of degree d , let $n : C^v \rightarrow C$ be the normalization map such that $C^v \cong \mathbb{P}^1$. Then the pair (resp. sheaf $n_*\mathcal{O}_{C^v}$) $(s, n_*\mathcal{O}_{C^v})$ with the unique section $s \in H^0(n_*\mathcal{O}_{C^v})$ is α -stable (resp. stable) for any positive parameter $\alpha > 0$ ([CC11, Proposition 3.18]). That is, we have natural injective maps

$$R_d \subset M(dm + 1); [C] \mapsto n_*\mathcal{O}_{C^v} \text{ and } R_d \subset M^\alpha(dm + 1); [C] \mapsto (s, n_*\mathcal{O}_{C^v}).$$

Let us denote \mathbf{M}_d (resp. \mathbf{M}_d^α) by the closure of the rational curves space R_d in the moduli space $M(dm + 1)$ (resp. $M^\alpha(dm + 1)$).

Problem 1.1. *Study the geometry \mathbf{M}_d via its variations spaces \mathbf{M}_d^α .*

Remark 1.2. For the degree is $d \leq 3$, one can easily see that all of α -stable sheaves spaces \mathbf{M}_d^α are isomorphic to \mathbf{M}_d . For $d = 1$ and 2, the space \mathbf{M}_d is isomorphic to the complete linear system

2010 *Mathematics Subject Classification.* 14E15, 14E05, 14M15.

Key words and phrases. Plane rational curves, Compactification, Birational geometry.

$|\mathcal{O}_{\mathbb{P}^2}(d)|$. The space \mathbf{M}_3 is isomorphic to a \mathbb{P}^6 -bundle over \mathbb{P}^2 . Here the fiber \mathbb{P}^6 parameterizes the rational cubic curves passing through a singular point \mathbb{P}^2 .

For the higher degree cases since there does not have any systematic method to study the geometry of \mathbf{M}_d , we do the work for case $d = 4$ firstly. We will prove that there exists a divisorial contraction among \mathbf{M}_d^α . Let us state main theorem of this paper. For sufficiently large parameter $\alpha \gg 1$, let denote by $\mathbf{M}_d^{\alpha \gg 1} := \mathbf{M}_d^+$. It is known that the space \mathbf{M}_d^+ is isomorphic to the relative Hilbert scheme of $\frac{(d-1)(d-2)}{2}$ -points over the complete linear system $|\mathcal{O}_{\mathbb{P}^2}(d)|$ ([He98, Section 4.4]). We call that a birational contraction is a *smooth* blow-down whenever the two birational varieties are locally smooth around the exceptional divisor and its image in the target space.

Theorem 1.3 (Proposition 3.6). *Under the above notation, there exist a smooth blow-down*

$$p_R : \mathbf{M}_4^+ \longrightarrow \mathbf{M}_4$$

where the exceptional divisor parameterizes the pair of the (degenerated) rational cubic curve and the colinear three points.

The key ingredient of the the proof of Theorem 1.3 is the analysis of the restrictions of pair wall-crossing of $\mathbf{M}^\alpha(4m+1)$. Using this fact, one can compute the virtual Poincaré polynomial of the space \mathbf{M}_4 (Corollary 4.4).

Corollary 1.4. *The virtual Poincaré polynomial of \mathbf{M}_4 is given by*

$$1 + 2t^2 + 5t^4 + 8t^6 + 11t^8 + 12t^{10} + 13t^{12} + 13t^{14} + 11t^{16} + 7t^{18} + 3t^{20} + t^{22}.$$

Question 1.5. Find out a smooth resolution of the space \mathbf{M}_4 and study its geometry in the view point of curve counting theory.

In the introduction part of the paper [DH88], S. Diaz and J. Harris has already mentioned the necessity of a *good* compactification of the space R_d for the intersection theory.

1.2. Stream of the paper. In section 2, we explicitly propose relevant spaces and review its properties. In section 3, we prove our main theorem by analyzing the wall-crossing of α -stable pairs on \mathbb{P}^2 . In the last section, as presenting the special fibres by using Macaulay2 ([GS]), we calculate the Poincaré polynomial of \mathbf{M}_4 .

Acknowledgement. The author would like to thank Dawei Chen for suggesting this topic and Jinhyung Park for helpful discussions and comments.

2. RELEVANT MODULI SPACES

In this section, we collect known facts for the proof of Theorem 1.3

2.1. Kronecker quiver and Hilbert scheme of cubes. Let $\mathbf{N} := \mathbf{N}(3; 2, 3)$ be the moduli space of the space of sheaf homomorphisms (so called, the Kronecker quivers)

$$(1) \quad \mathcal{O}_{\mathbb{P}^2}(-2)^{\oplus 2} \longrightarrow \mathcal{O}_{\mathbb{P}^2}(-1)^{\oplus 3}$$

up to the action of the automorphism group $\mathrm{GL}_2 \times \mathrm{GL}_3/\mathbb{C}^*$. It is known that \mathbf{N} is a smooth projective variety of dimension 6 ([Dre87]). On the other hand, let Z be a finite subscheme of \mathbb{P}^2 of length 3. Since a resolution of a general ideal sheaf $I_Z(1)$ twisted by $\mathcal{O}_{\mathbb{P}^2}(1)$ is of the form (1), the moduli space \mathbf{N} is birational to the Hilbert scheme $\mathbf{H}[3]$ of three points in \mathbb{P}^2 . The classification of locus in $\mathbf{H}[3]$ are very well-known. We list some loci which we will use later. Let x, y, z be the homogeneous coordinates of \mathbb{P}^2 .

- a) Let D_1 be the locus of *colinear* three points on \mathbb{P}^2 which is isomorphic to a \mathbb{P}^3 -bundle over \mathbb{P}^2 .
- b) Let D_2 be the locus of *curvelinear* three points which is isomorphic to $\mathbb{C}^* \times \mathbb{C}$ -bundle over \mathbb{P}^2 . The general element is of the form $\langle x^2, xy, y^2 - zx \rangle$.
- c) Let $D_3 := \overline{D}_2 \setminus D_2$ be the locus of the *non-curvelinear* three points which is isomorphic to \mathbb{P}^2 . The general element is of the form $\langle x^2, xy, y^2 \rangle$.

An explicit birational relation between $\mathbf{H}[3]$ and \mathbf{N} has been studied in [LQZ03].

Proposition 2.1 ([LQZ03, Section 6]). *Under the above notation, there exists a smooth blow-down morphism*

$$t : \mathbf{H}[3] \longrightarrow \mathbf{N}$$

which contracts the fiber \mathbb{P}^3 of the exceptional divisor D_1 .

2.2. The moduli of stable maps space. Let C be a projective connected reduced curve. A map $f : C \rightarrow \mathbb{P}^r$ is called by *stable* if C has at worst nodal singularities and $|\mathrm{Aut}(f)| < \infty$. Let \mathcal{K}_d be the moduli space of isomorphism classes of stable maps $f : C \rightarrow X$ with genus $g(C) = 0$ and $\deg(f) = d$. The space \mathcal{K}_d has been widely studied in [FP97]. In special, the space is compact and naturally contains the space R_d as a dense open subset because the normalization map of an irreducible rational curve has the trivial automorphism.

2.3. Summary of the result in [CC17]. Let X be a projective variety with a fixed ample line bundle $\mathcal{O}_X(1)$. Let

$$\chi(F(m)) = \sum_{k=0}^{\dim X} \dim H^k(F(m))$$

be the Hilbert polynomial of the coherent sheaf $F \in \mathrm{Coh}(X)$.

Definition 2.2. A pair (s, F) , $s \in H(F)$ is called α -*semistable* if F is pure and for any subsheaf $F' \subset F$, the inequality

$$(2) \quad \frac{\chi(F'(m)) + \delta \cdot \alpha}{r(F')} \leq \frac{\chi(F(m)) + \alpha}{r(F)}$$

holds for $m \gg 0$. Here $\delta = 1$ if the section s factors through F' and $\delta = 0$ otherwise. When the strict inequality holds, (s, F) is called an α -*stable* pair.

It is well-known that there exists a projective scheme $M^\alpha(P(m))$ parameterizing S -equivalence classes of α -semistable pairs with Hilbert polynomial $P(m)$ ([LP93, Theorem 4.12] and [He98, Theorem 2.6]). By decreasing the stability parameter α , one can obtain a sequence of flips among the moduli spaces of α -stable pairs.

As we let $\alpha = 0$ in (2), one can obtain the usual concept of semistable sheaves and thus the moduli space $M(P(m))$ which parameterizes S -equivalence classes of semistable sheaves on X with Hilbert polynomial $P(m)$ (For construction and its properties, see [Sim94]). In this paper, we deal with the case $X = \mathbb{P}^2$ and $P(m) = 4m + 1$. In [CC17, CM15], the authors studied the geometric properties of $M^\alpha(4m + 1)$ and $M(4m + 1)$ using the wall-crossing of pairs. It turns out that there exists a unique wall at $\alpha = 3$ ([CC17, Lemma 3.1]). Let us summarize the result for later use.

- (i) ([He98, Section 4.4] and [PT10]) Let $M^+(4m + 1) := M^\alpha(4m + 1)$ for $\alpha > 3$. The moduli space M_4^+ of α -stable pairs is isomorphic to the relative Hilbert scheme $\mathcal{H}[3]$ of three points on the complete linear system $|\mathcal{O}_{\mathbb{P}^2}(4)|$. For (s, F) be a stable pair, the corresponding point in the relative Hilbert scheme $\mathcal{H}[3]$ is given by

$$M^+ \rightarrow \mathcal{H}[3], (s, F) \mapsto (Z, C), Z \subset C = \text{Supp}(F), l(Z) = 3.$$

- (ii) ([CC17, Proposition 4.4]) Let $M^-(4m + 1) := M^\alpha(4m + 1)$ for $\alpha < 3$. The forgetful map $r : M^- \rightarrow M : (s, F) \mapsto F$ is a smooth blow-up along the Brill-Noether locus $\{[F] \in M_4 \mid \dim H^0(F) = 2\}$.
- (iii) ([CC17, Section 3.1]) The single flip over $M^3(4m + 1)$ is a composition of a smooth blow-up and a smooth blow-down. Let us denote Y^+ by the blow-up center in M^+ . Then Y^+ is isomorphic to a \mathbb{P}^3 -bundle over $\mathbb{P}^2 \times \mathbb{P}^9$ where the base parameterizes the pair of line and cubic curves and the fiber parameterizes three points on that line.

Remark 2.3. The image of the blow-up center Y^- in M^- by the forgetful map r in (ii) parameterizes the stable sheaf F which fits into a non-split extension:

$$0 \rightarrow \mathcal{O}_C \rightarrow F \rightarrow \mathcal{O}_L \rightarrow 0$$

for a line L and a cubic curve C . By a simple computation, we see that the image $r(Y^-)$ is isomorphic to a $\mathbb{P}(\text{Ext}_{\mathbb{P}^2}^1(\mathcal{O}_L, \mathcal{O}_C)) \cong \mathbb{P}^2$ -bundle over $\mathbb{P}^2 \times \mathbb{P}^9$. Also,

$$h^0(F) = \begin{cases} 1 & \text{if } [F] \in \mathbb{P}^2 \setminus \mathbb{P}^1; \\ 2 & \text{if } [F] \in \mathbb{P}^1. \end{cases}$$

and thus $Y^- \cong r(Y^-)$ by (ii).

In summary, there exists a commutative diagram

$$(3) \quad \begin{array}{ccccc} & & \widetilde{M(4m+1)} & & \\ & \swarrow p^+ & & \searrow p^- & \\ M^+(4m+1) & & & & M^-(4m+1) \\ & \swarrow & \text{a single flip} & \searrow & \\ & & \text{-----} & & \\ & \swarrow & & \searrow & \\ & & M^3(4m+1) & & M(4m+1) \\ & & & & \downarrow r \end{array}$$

Let \mathbf{M}_4^α be the closure of the space R_4 of $\mathbf{M}_4^\alpha(4m+1)$ for each $\alpha \in \mathbb{Q}^+$. Then the diagram (3) has been restricted to

(4)

$$\begin{array}{ccccc}
 & & \widetilde{\mathbf{M}}_4 & & \\
 & p^+ & \swarrow & \searrow & p^- \\
 & & \mathbb{R} & & \\
 \mathbf{M}_4^+ & \xleftarrow{\quad} & & \xrightarrow{\quad} & \mathbf{M}_4^- \\
 & \searrow & & \swarrow & \\
 & & \mathbf{M}_4^0 & & \\
 & \swarrow & & \searrow & \\
 \mathbf{H}[3] & \xrightarrow{\quad} & & \xrightarrow{\quad} & \mathbf{N}
 \end{array}$$

$\begin{array}{c} \parallel \\ \pi^+ \\ \parallel \end{array}$
 $\begin{array}{c} \cong \\ r \\ \cong \end{array}$
 $\begin{array}{c} \parallel \\ \pi^- \\ \parallel \end{array}$

where $\pi^+ : \mathbf{M}_4^+ \rightarrow \mathbf{H}[3]$ is the restriction map of the structure morphism of the relative Hilbert scheme $\mathcal{H}[3] \rightarrow \mathbf{H}[3]$. The remaining one of this paper is to prove that

- the restrictions of maps p^+ and r are an isomorphism,
- the composition of the restriction maps

$$p_R := r \circ p^- \circ (p^+)^{-1} : \mathbf{M}_4^+ \rightarrow \mathbf{M}_4$$

is a smooth blow-up map and

- there exists a generically bundle morphism $\pi^- : \mathbf{M}_4 \rightarrow \mathbf{N}$ such that $\pi^- \circ p_R = t \circ \pi^+$.

3. PROOF OF THEOREM 1.3

In this section, we analyze the restricted diagram in (3) during the wall-crossing. One of main method is to use the elementary modification of sheaves coming from the family of stable maps.

3.1. Some lemmas via stable maps.

Lemma 3.1 ([CC11, Example 6.1]). *For the irreducible plane curve C , let $f : C^v \rightarrow C$ be the normalization map and $s \in \text{Hom}(\mathcal{O}_C, f_*\mathcal{O}_{C^v}) \cong \mathbf{H}^0(\mathcal{O}_{C^v}) = \mathbb{C}$ be the canonical section. Then the pair $(s, f_*\mathcal{O}_{C^v})$ is α -stable for all $\alpha \in \mathbb{Q}^+$. In special, $f_*\mathcal{O}_{C^v}$ is stable.*

The correspondence between R_4 and the relative Hilbert scheme $\mathcal{H}[3]$ is given by the following way. For the curve C , the direct image sheaf $f_*\mathcal{O}_{C^v}$ fits into the short exact sequence

$$0 \rightarrow \mathcal{O}_C \rightarrow f_*\mathcal{O}_{C^v} \rightarrow \mathcal{Q} \rightarrow 0$$

such that $\dim \mathcal{Q} = 0$. By taking the dual $\text{Hom}^\bullet(-, \mathcal{O}_C)$ in the exact sequence,

$$(5) \quad 0 \rightarrow \text{Hom}_C(f_*\mathcal{O}_{C^v}, \mathcal{O}_C) \rightarrow \text{Hom}_C(\mathcal{O}_C, \mathcal{O}_C) = \mathcal{O}_C \rightarrow \text{Ext}_C^1(\mathcal{Q}, \mathcal{O}_C) \rightarrow 0$$

we have a pair (Z, C) , $Z \subset C$ such that the subscheme Z is defined by the (conductor) ideal sheaf $I_{Z,C} = \text{Hom}_C(f_*\mathcal{O}_{C^v}, \mathcal{O}_C)$ (cf. [PT10, Proposition B.5]).

Example 3.2. Let $f : \mathbb{P}^1 \rightarrow \mathbb{P}^2; [s : t] \mapsto [g_3(s, t)s : g_3(s, t)t : g_4(s, t)]$ be a map such that $g_i(s, t)$ are homogenous polynomial with respect to s, t of degree i . Then it is straightforward to check that the conductor ideal sheaf is defined by

$$I_Z = \langle x^2, xy, y^2 \rangle$$

where x, y, z are homogenous coordinates of \mathbb{P}^2 .

Recall that \mathbf{M}_4 (resp. \mathbf{M}_4^α) is the closure of R_4 in $M(4m + 1)$ (resp. $M^\alpha(4m + 1)$) respectively.

Proposition 3.3. For each closed point $[F] \in \mathbf{M}_4$,

$$\dim H^0(F) = 1.$$

Hence the forgetful map r is an isomorphism $\mathbf{M}_4^- \xrightarrow{r} \mathbf{M}_4$.

Proof. Let \mathcal{K}_4 be the moduli space of stable maps of degree 4 into \mathbb{P}^2 . Let

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{\mathbf{f}} & \mathbb{P}^2 \\ \downarrow \pi & & \\ \mathcal{K}_4 & & \end{array}$$

be a flat family of stable maps parameterized by \mathcal{K}_4 . Let $\Phi := (\pi, \mathbf{f}) : \mathcal{C} \rightarrow \mathcal{K}_4 \times \mathbb{P}^2$ be the push-out map. Then the direct image sheaf $\mathcal{F} := \Phi_* \mathcal{O}_{\mathcal{C}}$ is the flat family of pure sheaves parameterized by \mathcal{K}_4 . The general sheaves parameterized by \mathcal{K}_4 is stable by Lemma 3.1. Also, $H^0(\mathcal{F}|_{[f] \times \mathbb{P}^2}) \cong H^0(f_* \mathcal{O}_{\mathcal{C}}) = \mathbb{C}$. The claim is proved by the below Lemma 3.4. \square

Lemma 3.4. Let S be a small disk in \mathbb{C} containing 0. Let \mathcal{F} be a flat family of sheaves on

$$\mathbb{P}^r \times S \rightarrow S$$

with Hilbert polynomial $\chi(\mathcal{F}(m)) = dm + 1$. Let \mathcal{F} be stable sheaves over S except the origin $0 \in S$. Furthermore, $h^0(\mathcal{F}|_s) = 1$ for all $s \in S$. Then the modified stable sheaf at the origin has a unique non-zero global section.

Proof. Following the proof of [HL10, Theorem 2.B.1], it is enough to show that, after the single elementary modification of sheaf \mathcal{F} along $0 \in S$, the resulting center fiber has a one-dimensional global section space. Let

$$G \subset F := \mathcal{F}|_0$$

be the non-zero maximal destabilizing subsheaf of F . Then we have a short exact sequence

$$0 \rightarrow G \rightarrow F \rightarrow H := F/G \rightarrow 0.$$

Let

$$\chi(H(m)) = d'm + \chi' \text{ and } \chi(G(m)) = (d - d')m + 1 - \chi'.$$

Then by the maximality of G , we have

$$\frac{1 - \chi'}{d - d'} \geq \frac{1}{d} \Leftrightarrow \frac{d'}{d} \geq \chi'.$$

Since $h^1(F) = 0$ and G is supported on a curve in \mathbb{P}^r , $h^1(H) = 0$. That is, $h^0(H) = \chi' \geq 0$. Also, by above inequality, $\chi' \leq 1$. If $\chi' = 1$, then it is contradict to the assumption of G . Thus $h^0(H) = 0$.

This implies that $h^1(G) = 0$ because $h^1(F) = 0$. Now, for the modified sheaf \mathcal{F}' on $\mathbb{P}^r \times S$, the central fiber fits into the short exact sequence

$$0 \rightarrow H \rightarrow \mathcal{F}'|_{\mathbb{P}^r \times \{0\}} \rightarrow G \rightarrow 0.$$

Therefore $h^1(\mathcal{F}'|_{\mathbb{P}^r \times \{0\}}) = 0$. That is, $h^0(\mathcal{F}'|_{\mathbb{P}^r \times \{0\}}) = 1$. \square

3.2. Proof of Theorem 1.3. One of simple observation is that the canonical form of rational quartic plane curve with the prescribed singular points ($[1 : 0 : 0]$, $[0 : 1 : 0]$, $[0 : 0 : 1]$) can be written as

$$F(x, y, z) = a_0x^2y^2 + a_1y^2z^2 + a_2z^2x^2 + a_3x^2yz + a_4xy^2z + a_5xyz^2$$

where x, y, z are the homogenous coordinate of \mathbb{P}^2 and $a_0, a_1, \dots, a_5 \in \mathbb{C}$. Hence the projection map $\pi^+ : \mathbf{M}_4^+ \rightarrow \mathbf{H}[3]$ in the diagram (4) is generically \mathbb{P}^5 -fibration over $\mathbf{H}[3]$.

Lemma 3.5. *The map*

$$\pi^+ : \mathbf{M}_4^+ \rightarrow \mathbf{H}[3]$$

has a \mathbb{P}^5 -fibration over $\mathbf{H}[3] \setminus D_1 \sqcup D_3$.

Proof. Let \mathcal{Z} be the universal subscheme of $\mathbf{H}[3]$ and $q : \mathbf{H}[3] \times \mathbb{P}^2 \rightarrow \mathbf{H}[3]$ be the projection onto the first component. Then, the push-forward sheaf $\mathcal{E} := q_*(I_{\mathcal{Z}}^2 \boxtimes \mathcal{O}_{\mathbb{P}^2}(4))$ is locally free of rank 6 away from the union $D_1 \sqcup D_3$. Indeed, it is straightforward to check that

$$h^0(I_{\mathcal{Z}}^2(4)) = \begin{cases} 6 & \text{if } [Z] \in \mathbf{H}[3] \setminus D_1 \sqcup D_3; \\ 5 & \text{if } [Z] \in D_1; \\ 7 & \text{if } [Z] \in D_3. \end{cases}$$

and thus the claim is proved. Now let $\mathbb{P}(\mathcal{E}) := \text{Proj}(\text{Sym}^\bullet(\mathcal{E})) \rightarrow \mathbf{H}[3]$ be the projective scheme over $\mathbf{H}[3]$. Then $\mathbb{P}(\mathcal{E})$ is a subscheme of \mathbf{M}_4^+ (cf. [He98]). Since R_4 is irreducible and the canonical form of rational quartic curve, R_4 is a subscheme of $\mathbb{P}(\mathcal{E})$. Thus $\mathbf{M}_4^+ = \overline{R_4} \subset \mathbb{P}(\mathcal{E})$. Therefore \mathbf{M}_4^+ is also a \mathbb{P}^5 -fibration away from $D_1 \sqcup D_3$. \square

Proposition 3.6. (1) *The map*

$$\pi^- : \mathbf{M}_4 \rightarrow \mathbf{N}$$

is a \mathbb{P}^5 -fibration over $\mathbf{N} \setminus t(D_3)$.

(2) *Furthermore, the birational morphism $p_R : \mathbf{M}_4^+ \rightarrow \mathbf{M}$ is a smooth blow-down of $(\pi^+)^{-1}(D_1)$ such that there exists a commutative diagram*

$$\begin{array}{ccc} \mathbf{M}_4^+ & \xrightarrow{p_R} & \mathbf{M}_4 \\ \pi^+ \downarrow & & \downarrow \pi^- \\ \mathbf{H}[3] & \xrightarrow{t} & \mathbf{N}. \end{array}$$

Proof. Recall that the image of the blow-up center Y^- by the forgetful map $r : \mathbf{M}^-(4m+1) \rightarrow \mathbf{M}(4m+1)$ is isomorphic to a \mathbb{P}^2 -bundle over $\mathbb{P}^2 \times \mathbb{P}^9$ where the base space parameterizes the pair (L, C) and the fiber is $\mathbb{P}(\text{Ext}^1(\mathcal{O}_L, \mathcal{O}_C))$ for the line L and cubic curve C . Considering the image of the stable maps parameterized by \mathcal{K}_4 , the support of the sheaf $F \in \mathbf{M} \cap r(Y^-)$ must be $\text{Supp}(F) = L^2 \cup Q$ for some line L and conic Q .

Step 1. Through the elementary modification of stable maps, we prove that

$$\mathbf{M} \cap r(Y^-) \cong \mathbb{P}^2 \times \mathbb{P}^5 := \Gamma$$

where \mathbb{P}^2 (resp. \mathbb{P}^5) parameterizes lines (resp. conics) in \mathbb{P}^2 . Main stream of the proof is the same as that of [CCM14, Proposition 5.10]. For the convenience of reader we address the detail. Let Δ be the locus of the stable maps $f : C \rightarrow \mathbb{P}^2$ of degree 4 such that the image is $f(C) = L \cup Q$ where L (resp. Q) is a line (resp. conic). One can regard f as the composition map

$$f = g \circ h : C \xrightarrow{h} C' \xrightarrow{g} \mathbb{P}^2,$$

where C' is a pair of lines. Here h is a $2 : 1$ -covering from L_1 and a bijection from L_2 . Also, g is a partial normalization map such that the image is $g(C') = L \cup Q$. By applying the octahedron axiom for the complex $f^*\Omega_{\mathbb{P}^2} \rightarrow h^*\Omega_{C'} \rightarrow \Omega_C$, we see that the normal space of Δ in $\mathbf{M}(\mathbb{P}^2, 4)$ at $[f]$ is identified with

$$N_{\Delta/\mathbf{M}(\mathbb{P}^2, 4), [f]} \cong \text{Ext}_{C'}^1([g^*\Omega_{\mathbb{P}^2} \rightarrow \Omega_{C'}], \mathcal{O}_L(-1)).$$

On the other hand, let $i : L \subset C'$ is the inclusion map. By the octahedron axiom for the complex $(i \circ g)^*\Omega_{\mathbb{P}^2} \rightarrow i^*\Omega_{C'} \rightarrow \Omega_L$ again, one can easily see that there exists a quasi-isomorphism between

$$i^*[g^*\Omega_{\mathbb{P}^2} \rightarrow \Omega_{C'}] \cong N_{L/\mathbb{P}^2}^*(-1)[1]$$

and thus

$$N_{\Delta/\mathbf{M}(\mathbb{P}^2, 4), [f]} \cong \text{Hom}(N_{L/\mathbb{P}^2}^*(-1), \mathcal{O}_L(-1)) \cong \text{Hom}(I_L(-1), \mathcal{O}_L(-1)).$$

By analyzing the Kodaira-Spencer map

$$(6) \quad \phi : N_{\Delta/\mathbf{M}(\mathbb{P}^2, 4), [f]} \rightarrow \text{Ext}^1(g_*\mathcal{O}_C, \mathcal{O}_L(-1)),$$

one can prove that ϕ is injective and thus each modified sheaf is non-split (cf. [CK11, Lemma 4.10]). Hence after the modification, we have a flat family of stable sheaves parameterized by $\mathbb{P}(N_{\Delta/\mathbf{M}(\mathbb{P}^2, 4)}) = \mathbb{P}^1$ -bundle over Δ . For $w \in \mathbb{P}^1$, let F_w be the stable sheaf which fits into a exact sequence

$$0 \rightarrow \mathcal{O}_L(-1) \rightarrow F_w \rightarrow g_*\mathcal{O}_C \rightarrow 0.$$

By Proposition 3.3, $h^0(F_w) = 1$ for each point $[w] \in \mathbb{P}^1$. Hence $F_w \cong F_{w'}$ for any point $[w], [w'] \in \mathbb{P}^1$ by Bézout's Theorem (Remark 2.3). By some diagram chasing, we know that F_w fits into the short exact sequence

$$0 \rightarrow \mathcal{O}_{L \cup Q} \rightarrow F_w \rightarrow \mathcal{O}_L \rightarrow 0$$

and thus the claim is proved. Hence the composition map $t \circ \pi^+ \circ p^+ : \widetilde{\mathbf{M}(4m+1)} \rightarrow \mathbf{N}$ factors through $\mathbf{M}_4^- = \mathbf{M}_4$ by the rigidity lemma. That is, we have a morphism

$$\pi^- : \mathbf{M}_4 \rightarrow \mathbf{N}.$$

Note that the fiber of $[L] \in t(D_1) \subset \mathbf{N}$ is constantly a \mathbb{P}^5 which parameterizes conics. By Lemma 3.5, we know that the map π^- is a \mathbb{P}^5 -fibration except the locus D_3 .

Step 2. By (1), the normal space of Γ in \mathbf{M}_4 at $[F_w] \in \mathbf{M}_4$ is naturally isomorphic to

$$N_{\Gamma/\mathbf{M}_4, F_w} \cong N_{t(D_1)/\mathbf{N}, [\pi^-(F_w)]}.$$

This implies that the blow-up of $\mathbf{M}_4 = \mathbf{M}_4^+$ along Γ is nothing but the strict transformation of the blow-up morphism $p^- : \widetilde{\mathbf{M}(4m+1)} \rightarrow \mathbf{M}^-(4m+1)$ because the projectivation of the normal

space $N_{t(D_1)/\mathbf{N}, [\pi^-(F_w)]}$ determines three points on the line L . In special, the exceptional divisor E in $\widetilde{\mathbf{M}}^+$ is a \mathbb{P}^3 -bundle over Γ . Then the image $p^+(E) \cong E \subset Y^+$ is still divisor in \mathbf{M}^+ and thus $\widetilde{\mathbf{M}}^+ \stackrel{p^+}{\cong} \mathbf{M}^+$. Note that E is the inverse image $(\pi^+)^{-1}(D_1)$ by its construction. This finishes the proof of (2). \square

4. POINCARÉ POLYNOMIAL OF \mathbf{M}_4

In this section we compute the Poincaré polynomial of \mathbf{M}_4 by using Proposition 3.6.

4.1. Fibre over the non-curvilinear points. In this section, we describe the fiber of the morphism p over Y_1 by using the computer algebra system, Macaulay2 ([GS]) and hence we compute the virtual Poincaré polynomial of \mathbf{M}_4 .

Proposition 4.1. *Each fiber $\pi^- : \mathbf{M}_4 \rightarrow \mathbf{N}$ over the closed point $[Z] \in D_1$ is isomorphic to a projective space \mathbb{P}^8 .*

Proof. We assume that $I_Z = \langle x^2, xy, y^2 \rangle$. We prove the claim by presenting a local neighborhood of \mathbf{M}_4^+ containing the fiber $p^{-1}([Z])$ with the help of the computer system. Let $a_1, a_2, b_1, b_2, c_1, c_2$ be a local affine chart of $\mathbf{H}[3]$ at $[I_Z] = [\langle x^2, xy, y^2 \rangle]$ ([Hai02, Section 3.1]). The ideal of subschemes parameterizing the points of $\mathbf{H}[3]$ around $[Z]$ is generated by three quadric polynomials:

- i) $q_1 := x^2 - a_1x - a_2y - a_2(b_1 - c_2) - b_2(b_2 - a_1)$;
- ii) $q_2 := xy - b_1x - b_2y - a_2c_1 + b_1b_2$;
- iii) $q_3 := y^2 - c_1x - c_2y - c_1(b_2 - a_1) - b_1(b_1 - c_2)$.

Note that $\langle q_1, q_2, q_3 \rangle|_{\{a_i=b_i=c_i=0\}} = I_Z$ for $i = 1, 2$. Let

$$f(x, y, z) = d_0x^4 + d_1x^3y + d_2x^2y^2 + d_3xy^3 + \dots + d_8y^3z + d_9x^2z^2 + d_{10}y^2z^2 + d_{11}xyz^2 + d_{12}xz^3 + d_{13}yz^3 + d_{14}z^4$$

be the homogeneous polynomial defining the plane quartic curve. Let f_x be the partial derivatives of f with respect to x (similarly, f_y and f_z). The rationality condition of the plane curve provides us the nine resultant conditions

$$J = \langle \text{Res}(f_x, q_i), \text{Res}(f_y, q_i), \text{Res}(f_z, q_i) \rangle \quad \text{for } i = 1, 2, 3.$$

Note that $f_x(x, y, 1) \equiv s(x, y) \pmod{(q_1, q_2, q_3)}$ for some quadric polynomial $s(x, y)$ (similarly, $f_y(x, y, 1)$ and $f_z(x, y, 1)$). Hence the resultant can be easily computed by using the computer algebra system, Macaulay2 ([GS]) and obtain that the ideal J is the defining equation of \mathbf{M}^+ around $(\pi^+)^{-1}([Z])$. The special fiber $(\pi^+)^{-1}([Z])$ is nothing but

$$J + \langle a_1, a_2, b_1, b_2, c_1, c_2 \rangle = \langle d_9, d_{10}, d_{11}, d_{12}, d_{13}, d_{14} \rangle$$

which defines the space \mathbb{P}^8 . Thus we finish the proof of the claim. \square

Remark 4.2. The fiber $[Z]$ by π^- in the statement of Proposition 4.1 can be described by the stable map. As we have seen in Example 3.2, the conductor ideal of the map

$$f : \mathbb{P}^1 \rightarrow \mathbb{P}^2, \quad f([s : t]) = [g_3(s, t)s : g_3(s, t)t : g_4(s, t)]$$

is given by $\langle x^2, xy, y^2 \rangle$ for the general choice of $g_3(s, t)$ and $g_4(s, t)$. Also the supporting curve $C = f(\mathbb{P}^1)$ is defined by the ideal $I_C = \langle g_3(x, y)z - g_4(x, y) \rangle$. Hence the fiber $p^{-1}([Z])$ contains an open part of $\mathbb{P}(\mathbb{H}^0(\mathcal{O}_{\mathbb{P}^1}(3) \oplus \mathcal{O}_{\mathbb{P}^1}(4)))$.

4.2. The virtual Poincaré polynomial of \mathbf{M}_4 . In this subsection, we calculate the virtual Poincaré polynomial of \mathbf{M} by using Theorem 1.3. Let X be a quasi-projective variety. For the Hodge-Deligne polynomial $E_c(X)(u, v)$ for compactly supported cohomology of X , let

$$P(X) := E_c(X)(-t, -t)$$

be the *virtual* Poincaré polynomial of X . For the motivic properties of the virtual Poincaré polynomial, see [Muñ08, Theorem 2.2] and [BJ12, Lemma 3.1].

Proposition 4.3. (1) $P(\mathbb{P}^n) = \frac{t^{2n+2}-1}{t^2-1}$.

(2) $P(X) = P(Z) + P(X \setminus Z)$ for any closed subset $Z \subset X$.

(3) $P(X) = P(F) \cdot P(B)$ for the étal locally trivial fibration $X \rightarrow B$ with the constant fiber isomorphic to the Grassmannian variety $\text{Gr}(k, n)$. Here $\text{Gr}(k, n)$ parameterizes k -dimensional sub-vector spaces in the vector space \mathbb{C}^n .

Corollary 4.4. The virtual Poincaré polynomials of \mathbf{M}_4 is given by

$$1 + 2t^2 + 5t^4 + 8t^6 + 11t^8 + 12t^{10} + 13t^{12} + 13t^{14} + 11t^{16} + 7t^{18} + 3t^{20} + t^{22}.$$

Proof. By $P(\mathbf{H}[3]) = 1 + 2t^2 + 5t^4 + 6t^6 + 5t^8 + 2t^{10} + t^{12}$ ([ELB88]) and Proposition 2.1,

$$P(\mathbf{N}) = P(\mathbf{H}[3]) + (1 - P(\mathbb{P}^3)) \cdot P(\mathbb{P}^2) = 1 + t^2 + 3t^4 + 3t^6 + 3t^8 + t^{10} + t^{12}.$$

Also, from Proposition 3.6, Proposition 4.1 and Proposition 4.3, we have

$$P(\mathbf{M}) = P(\mathbb{P}^5) \cdot (P(\mathbf{N}) - P(\mathbb{P}^2)) + P(\mathbb{P}^8) \cdot P(\mathbb{P}^2).$$

Cooking up these ones, we obtain the results. □

REFERENCES

- [BJ12] B. Bakker and A. Jorza. Higher rank stable pairs on k3 surfaces. *Commun. Number Theory Phys.*, 6(4):805–847, 2012. [10](#)
- [CC11] Dawei Chen and Izzet Coskun. Towards Mori’s program for the moduli space of stable maps. *Amer. J. Math.*, 133(5):1389–1419, 2011. [1](#), [5](#)
- [CC17] Jinwon Choi and Kiryong Chung. Moduli spaces of -stable pairs and wall-crossing on \mathbb{P}^2 . *J. Math. Soc. Jap.*, 68(2):685–789, 2017. [3](#), [4](#)
- [CCM14] Jinwon Choi, Kiryong Chung, and Mario Maican. Moduli of sheaves supported on quartic space curves. arXiv:1409.1449, 2014. [8](#)
- [CK11] Kiryong Chung and Young-Hoon Kiem. Hilbert scheme of rational cubic curves via stable maps. *Amer. J. Math.*, 133(3):797–834, 2011. [8](#)
- [CM15] Kiryong Chung and Han-Bom Moon. Chow ring of the moduli space of stable sheaves supported on quartic curves. arXiv:1506.00298, 2015. [4](#)
- [DH88] Steven Diaz and Joe Harris. Geometry of the severi variety. *Trans. Amer. Math. Soc.*, 309(1):1–34, 1988. [2](#)
- [Dre87] J.-M. Drezet. Fibrés exceptionnels et variétés de modules de faisceaux semi-stables sur $\mathbf{P}_2(\mathbb{C})$. *J. Reine Angew. Math.*, 380:14–58, 1987. [3](#)
- [ELB88] G. Elencwajg and P. Le Barz. Explicit computations in $\text{Hilb}^3\mathbb{P}^2$. In *Algebraic geometry (Sundance, UT, 1986)*, volume 1311 of *Lecture Notes in Math.*, pages 76–100. Springer, Berlin, 1988. [10](#)

- [FP97] W. Fulton and R. Pandharipande. Notes on stable maps and quantum cohomology. In *Algebraic geometry—Santa Cruz 1995*, volume 62 of *Proc. Sympos. Pure Math.*, pages 45–96. Amer. Math. Soc., Providence, RI, 1997. [1](#), [3](#)
- [GS] Daniel R. Grayson and Michael E. Stillman. Macaulay2, a software system for research in algebraic geometry. Available at <http://www.math.uiuc.edu/Macaulay2/>. [2](#), [9](#)
- [Hai02] Mark Haiman. Notes on macdonald polynomials and the geometry of hilbert schemes. In Sergey Fomin, editor, *Symmetric Functions 2001: Surveys of Developments and Perspectives*, pages 1–64, Dordrecht, 2002. Springer Netherlands. [9](#)
- [He98] Min He. Espaces de modules de systèmes cohérents. *Internat. J. Math.*, 9(5):545–598, 1998. [1](#), [2](#), [3](#), [4](#), [7](#)
- [HL10] Daniel Huybrechts and Manfred Lehn. *The geometry of moduli spaces of sheaves*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, second edition, 2010. [6](#)
- [KP01] B. Kim and R. Pandharipande. The connectedness of the moduli space of maps to homogeneous spaces. In *Symplectic geometry and mirror symmetry (Seoul, 2000)*, pages 187–201. World Sci. Publ., River Edge, NJ, 2001. [1](#)
- [LP93] Joseph Le Potier. Systèmes cohérents et structures de niveau. *Astérisque*, (214):143, 1993. [1](#), [3](#)
- [LQZ03] Wei-Ping Li, Zhenbo Qin, and Qi Zhang. Curves in the Hilbert schemes of points on surfaces. In *Vector bundles and representation theory (Columbia, MO, 2002)*, volume 322 of *Contemp. Math.*, pages 89–96. Amer. Math. Soc., Providence, RI, 2003. [3](#)
- [Muñ08] Vicente Muñoz. Hodge polynomials of the moduli spaces of rank 3 pairs. *Geom. Dedicata*, 136:17–46, 2008. [10](#)
- [PT10] R. Pandharipande and R. P. Thomas. Stable pairs and BPS invariants. *J. Amer. Math. Soc.*, 23(1):267–297, 2010. [4](#), [5](#)
- [Sim94] Carlos T. Simpson. Moduli of representations of the fundamental group of a smooth projective variety. I. *Inst. Hautes Études Sci. Publ. Math.*, (79):47–129, 1994. [1](#), [4](#)

DEPARTMENT OF MATHEMATICS EDUCATION, KYUNGPOOK NATIONAL UNIVERSITY, 80 DAEHAKRO, BUKGU, DAEGU 41566, KOREA

E-mail address: krchung@knu.ac.kr