CANONICAL DECOMPOSITIONS OF ABELIAN GROUPS

PHILL SCHULTZ

ABSTRACT. Every torsion–free abelian group of finite rank has two essentially unique complete direct decompositions whose summands come from specific classes of groups.

1. INTRODUCTION

An intriguing feature of abelian group theory is that torsion-free groups, even those of finite rank, may fail to have unique complete decompositions; see for example [Fuchs, 2015, Chapter 12, §5] or [Mader and Schultz, 2018]). Therefore it is interesting to show that every such group has essentially unique complete decompositions with summands from certain identifiable classes. For example, it was shown in [Mader and Schultz, 2018] that every finite rank torsion-free group Ghas a Main Decomposition, $G = G_{cd} \oplus G_{cl}$ where G_{cd} is completely decomposable and G_{cl} has no rank 1 direct summand; G_{cd} is unique up to isomorphism and G_{cl} unique up to near isomorphism.

Let \mathcal{G} be the class of torsion-free abelian groups of finite rank and let $G \in \mathcal{G}$. The aim of this paper is to show that:

- $G = G_{sd} \oplus G_{ni}$ where G_{sd} is a direct sum of strongly indecomposable groups and G_{ni} has no strongly indecomposable direct summand. G_{sd} is unique up to isomorphism and G_{ni} up to near isomorphism.
- $G = G^{(fq)} \oplus G^{(rq)} \oplus G^{(dq)}$, where $G^{(fq)}$ is an extension of a completely decomposable group by a finite group, $G^{(rq)}$ is an extension of a completely decomposable group by an infinite reduced torsion group, and $G^{(dq)}$ is an extension of a completely decomposable group by an infinite torsion group which has a divisible summand. Each of the three summands is unique up to isomorphism.

The first case, dealt with in Section 3, I call a *Principal Decomposition*. Since rank 1 groups are strongly indecomposable, a Principal Decomposition is a refinement of a Main Decomposition; the second case, the subject of Section 5, I call a *Torsion Quotient Decomposition*. Of course the Principal Decomposition can be applied to each summand of the Torsion Quotient Decomposition and vice versa, leading to decompositions of the form $\bigoplus G_j^{(i)}$, and it does not matter in which order the canonical decompositions are applied.

Although this paper is intended to address particular problems in abelian group theory, in Section 6, I show that the results can be couched in categorical form, so

²⁰¹⁰ Mathematics Subject Classification. 20K15, 20K25, 16D70.

 $Key\ words\ and\ phrases.$ finite rank torsion–free abelian group; strongly indecomposable; direct decomposition.

they have immediate applications to larger classes of modules, for example torsion–free finite rank modules over an integral domain.

2. NOTATION

Unless otherwise signalled, the word 'group' denotes torsion-free abelian group of finite rank. Every group is isomorphic to a subgroup of the infinite dimensional rational vector space $V = \mathbb{Q}^{\mathbb{N}}$; define \mathcal{G} to be the set of finite rank subgroups of V. If $S \subseteq V$, $\langle S \rangle$ denotes the additive subgroup generated by S, [S] the subspace generated by S and S^G_* the group $[S] \cap G$. Thus [G] is the subspace of V generated by G and if $S \subseteq G$ then S^G_* is the pure subgroup of G generated by S. When there is no ambiguity, we just write S_* , and we adopt the usual convention on singletons, namely $b_* = \langle b \rangle_*$ for $b \in G$.

Apart from side-stepping foundational problems, these conventions have the advantage that homomorphisms $G \to H$ can be identified with their unique extensions $[G] \to [H]$. In particular, groups G and H are isomorphic if and only if there is an automorphism of V mapping G onto H, so the automorphism group $\operatorname{Aut}(G)$ coincides with the the stabilizer $\operatorname{Aut}_G([G])$ of the non-singular linear transformations of [G]. If $r \in \mathbb{Q}^*$, the non-zero rationals, then $rG \in \mathcal{G}$. Multiplication by r = a/bis an automorphism of [G] which is an automorphism of G when G is divisible by b. By r = a/b, we intend that $0 \neq a \in \mathbb{Z}$ and $0 \neq b \in \mathbb{N}$ with $\operatorname{gcd}\{a, b\} = 1$. Another advantage of our conventions is that integrally independent subsets of Gare rationally independent subsets of [G] contained in G, so we may omit the first adjective. The *rank* of G is the cardinality of some, and hence every, maximal independent subset.

Since every group is a direct sum of a unique divisible and a reduced group, we limit consideration to reduced groups. A decomposition of G into non-zero direct summands is called *complete* if these summands are indecomposable. Since rank(G) is finite, G always has complete decompositions, in general not unique even up to isomorphism.

A rank 1 group is a group isomorphic to a non-zero subgroup of the rationals \mathbb{Q} , a completely decomposable (cd) group is a direct sum of rank 1 groups, and an almost completely decomposable (acd) group is one containing a completely decomposable group as a subgroup of finite index.

Groups G and H in \mathcal{G} are quasi-equal, denoted $G \doteq H$, if there exists $r \in \mathbb{Q}^*$ such that rG = H, and quasi-isomorphic, denoted $G \approx H$, if $G' \cong H'$ for some $G' \doteq G$ and $H' \doteq H$. Thus if H and K are pure subgroups of G then $H \doteq K$ if and only if H = K. Clearly quasi-isomorphic groups have the same rank.

G is *quasi-decomposable* if $G \doteq H \oplus K$ for non-zero *H* and $K \in \mathcal{G}$, and *H* and *K* are called quasi-summands. *G* is *strongly indecomposable* (si) if it has no proper quasi-decomposition. Equivalently, *G* is si if and only if for all $k \in \mathbb{N}$, kG is indecomposable. A direct sum $\bigoplus_{i \in [t]} H_i$ with each H_i si, is called a *strong decomposition*. It follows from [Fuchs, 2015, Theorem 9.9] that strong decompositions are unique up to isomorphism, i.e., if $\bigoplus_i H_i = \bigoplus_j K_j$ are strong decompositions, then their summands can be paired so that $H_i \cong K_j$.

It was shown in [Schultz, 2020, Section 6] that every non-zero $G \in \mathcal{G}$ contains as a subgroup of finite index a strong decomposition $\bigoplus_{i \in [t]} H_i$ in which each H_i is pure in G. We call such a subgroup a Jónsson subgroup of G, denoted J(G). By the finite index property, $G = (J(G))_*$. For example, by [Mader, 2000, Chapter 4], if G is an acd group, then regulating subgroups are Jónsson subgroups.

Clearly rank 1 groups are si and it was shown in [Schultz, 2020, Theorem 7.2] that $J \in \mathcal{G}$ of rank > 1 is si if and only if J is an extension of a cd group H by an infinite torsion group T such that J/H has no decomposition lifting to J.

Extending the well known notion of type ([Mader, 2000, §2.2]) from rank 1 groups to si groups in general, the isomorphism class of an si group is called a *Type*, so types are special cases of Types. Denote by $\mathcal{T}(D)$ the set of Types of a strong decomposition D. By Lady's Theorem [Schultz, 2020, Corollary 6.5], $\mathcal{T}(D)$ is finite and a strong decomposition has the form $D = \bigoplus_{J \in \mathcal{T}} J^{n_J}$ where $\mathcal{T} = \mathcal{T}(D)$ and each $n_J \in \mathbb{N}$. Such a decomposition, which is unique up to isomorphism, is called a *homogeneous decomposition* of D, and the summands J^{n_J} are called the *homogeneous parts* of D.

Near isomorphism is an equivalence on \mathcal{G} weaker than isomorphism but stronger than quasi-isomorphism. There are several equivalent characterisations, see for example [Mader, 2000, §9.1], of which the most useful in our case is that G is nearly isomorphic to H if there exists $K \in \mathcal{G}$ such that $K \oplus G \cong K \oplus H$.

We define an ostensibly stronger equivalence by defining groups G and H to be strongly stably isomorphic, denoted $G \cong_{ss} H$, if there exists a strong decomposition K such that $K \oplus G \cong K \oplus H$. It is known, [Mader, 2000, Corollary 9.1.9], that for acd groups, near and strongly stable isomorphism are equivalent; however, it is still undetermined whether there exist nearly isomorphic groups G and H which fail to be strongly stably isomorphic.

A subgroup H of G is full if rank $(H) = \operatorname{rank}(G)$; equivalently, H is full in G if and only if G/H is torsion, and hence if and only if $H_* = G$. In particular, J(G) is full in G.

Unexplained notation comes from the standard references [Fuchs, 2015] or [Mader, 2000].

3. The Principal Decomposition

Let H be a strong decomposition with Type set \mathcal{J} , so that H has a homogeneous decomposition $\bigoplus_{J \in \mathcal{J}} J^{n_J}$. By the uniqueness property of strong decompositions, each decomposition $H = L \oplus K$ has the form $L = \bigoplus_{J \in \mathcal{J}} J^{m_J}$ and $K = \bigoplus_{J \in \mathcal{J}} J^{\ell_J}$ where m_J and ℓ_J are non-negative integers such that $m_J + \ell_J = n_J$.

Recall that a Principal Decomposition of $G \in \mathcal{G}$ is a decomposition $G = G_{sd} \oplus G_{ni}$ in which G_{sd} is a strong decomposition and G_{ni} has no si summand.

Proposition 3.1. Let $G = D \oplus B = A \oplus C$ where D and A are strong decompositions and B and C have no si summand. Then $D \cong A$ and $B \cong_{ss} C$.

Proof. Let π_A and π_C be the projections corresponding to the second decomposition. Now $D = D(\pi_A + \pi_C) = D\pi_A \oplus D\pi_C$ so $D = J(D) \cong J(D\pi_A) \oplus J(D\pi_C)$. Since D is a strong decomposition while C contains no si summand, $D \cong J(D\pi_A)$ is isomorphic to a summand of A. Similarly, A is isomorphic to a summand of D. Since G has finite rank, $A \cong D$ and hence by definition, $B \cong_{ss} C$.

Theorem 3.2. Every $G \in \mathcal{G}$ has a Principal Decomposition $G = G_{sd} \oplus G_{ni}$, in which G_{sd} is unique up to isomorphism, and G_{ni} up to strongly stable isomorphism.

Proof. The theorem is vacuously true if G is a strong decomposition or has no strongly indecomposable summands, so assume neither of these extreme cases occur.

PHILL SCHULTZ

Let \mathcal{J} be the set of all strong decompositions which are summands of G. Let G_{sd} be an element of \mathcal{J} of maximum rank, and let G_{ni} be any direct complement of G_{sd} . Then G_{ni} has no strongly decomposable summand, so this is a Principal Decomposition. By Proposition 3.1, G_{sd} is unique up to isomorphism, and G_{ni} up to strongly stable isomorphism.

Corollary 3.3. All summands of G which are maximum rank strong decompositions are isomorphic.

Remark 3.4. Principal Decompositions do not respect direct decompositions, in the following sense: Let $G = H \oplus K \in \mathcal{G}$. Then in general, $G_{sd} \neq H_{sd} \oplus K_{sd}$. For example there is a well known example [Fuchs, 2015, Theorem 5.2] of a rank 4 group G having a Principal Decomposition with rank $(G_{sd}) = 1$ such that $G = H \oplus K$ where both H and K are indecomposable of rank 2.

4. Ranges and Torsion Quotients

We call a maximal independent subset of G a basis. Bases of groups were studied in [Schultz, 2020, Proposition 2.2], where it was shown that if B is a basis of G then every $0 \neq a \in G$ has a unique representation $a = k^{-1} \sum_{b \in B} n_b b$ where $n_b \in \mathbb{Z}, k \in \mathbb{N}$ and $\gcd\{k, n_b : b \in B\} = 1$.

A range of G is a full cd subgroup whose the rank 1 summands are pure in G. If B is a basis of G, the group $(B)_* = \bigoplus_{b \in B} b_*$ is a range, and all ranges are of the form $(B)_*$ for some basis B.

Since $(B)_*$ is full in $G, G/(B)_*$ is a torsion group, called the *torsion quotient of* G with respect to $(B)_*$.

Lemma 4.1. With the notation above, $(B)_* = J(G)$ if and only if $(B)_*$ has finite index in G.

Proof. If $(B)_* = J(G)$, then by definition, $(B)_*$ has finite index in G. Conversely, since $(B)_*$ is a strong decomposition each of whose summands is pure in G, if $(B)_*$ has finite index in G then $(B)_* = J(G)$.

Proposition 4.2. Let $G \in \mathcal{G}$ have basis B such that $G/(B)_*$ is infinite and reduced. Then for every basis C, $G/(C)_*$ is infinite and reduced.

Proof. Let $\{x_i + (B)_* : i \in \mathbb{N}\}$ be an infinite independent set in $G/(B)_*$. Suppose by way of contradiction that $\{x_i + (C)_* : i \in \mathbb{N}\}$ is dependent in $G/(C)_*$, say $\sum_{j \in [n]} r_j x_j = c \in (C)_*$. Let c have B-representation $k^{-1} \sum_{i \in [t]} n_i b_i$. Then $\sum_{i \in [n]} r_j x_j \in (B)_*$, contradicting independence of $\{x_i + (B)_*\}$.

Now suppose that $G/(C)_*$ has a divisible summand $\cong \mathbb{Z}(p^{\infty})$ for some prime p. Then there exist $\{y_i : i \in \mathbb{N}\}$ in G such that $py_0 \in (C)_*$ and $py_{i+1} - y_i \in (C)_*$. Each y_i has B-representation $z_i = k_i^{-1} \sum_{j \in [t_i]} n_{ij} b_{ij}$, so that $\{z_0, z_i : i \in \mathbb{N}\}$ generates a summand of $G/(B)_* \cong \mathbb{Z}(p^{\infty})$. This contradiction implies that $G/(C)_*$ is reduced.

Corollary 4.3. Let G have torsion quotient $T = G/(B)_*$. Then the property that T is finite or has a divisible summand is independent of the choice of basis B. \Box

A useful property relating decompositions of G to decompositions of J(G), proved in [Schultz, 2020, Proposition 6.4], is the following:

Proposition 4.4. Let $G \in \mathcal{G}$.

- (1) If $G = H \oplus K$, then $J(G) = J(H) \oplus J(K)$.
- (2) If $J(G) = L \oplus M$, then $G = L_* \oplus M_*$ if and only if $L_* \oplus M_*$ is pure in G.

Proposition 4.5. Let J be a strongly indecomposable group. Then either J has rank 1, or for every basis B of J, $J/(B)_*$ is infinite and has no decomposition lifting to J.

Proof. Rank 1 groups are evidently si, so assume $\operatorname{rank}(J) > 1$. Let B be a basis of J, so $(B)_*$ is a full cd subgroup of J and hence $T = J/(B)_*$ is torsion. If T is finite, then $J \doteq (B)_*$ so $\operatorname{rank}(J) = 1$.

Suppose then that T is infinite. If T has no decomposition lifting to J, then by [Schultz, 2020, Theorem 7.2], J is si, while if T has such a decomposition, then by definition, J is decomposable, a contradiction.

5. The Torsion Quotient Decomposition

We shall use the following notation:

- \mathcal{J} : the set of strongly indecomposable groups in \mathcal{G} , and $\oplus \mathcal{J}$: finite direct sums from \mathcal{J} , i.e., strong decompositions;
- $\mathcal{J}^{(fq)}$: the subset of strong decompositions in \mathcal{J} having finite torsion quotient, and $\oplus \mathcal{J}^{(fq)}$: finite direct sums from \mathcal{J} ;
- $\mathcal{J}^{(rq)}$: the subset of strong decompositions in \mathcal{J} having reduced torsion quotient, and $\oplus \mathcal{J}^{(rq)}$: finite direct sums from \mathcal{J} .

Proposition 5.1. The three sets in the hierarchy of subclasses

$$\oplus \mathcal{J}^{(fq)} \subset \oplus \mathcal{J}^{(rq)} \subset \oplus \mathcal{J},$$

are closed under isomorphism, direct sums and direct summands.

Proof. Let $G \in \oplus \mathcal{J}$ and $\theta \colon G \to H$ an isomorphism. Then by [Schultz, 2020, Section 6] $H \in \oplus \mathcal{J}$. If $G \in \oplus \mathcal{J}^{(fq)}$ or $\oplus \mathcal{J}^{(rq)}$ then by Lemma 4.2, $H \in \oplus \mathcal{J}^{(fq)}$ or $\oplus \mathcal{J}^{(rq)}$ respectively.

The closure of each of the sets under direct sums follows directly from the definitions.

Let $H = K \oplus L \in \oplus \mathcal{J}$. Then H has a basis $B = C \cup D$ in which C is a basis of K and D a basis of L. Hence $H/(B)_* \cong K/(C)_* \oplus L/(D)_*$ and $K \in \oplus \mathcal{J}$. Furthermore, $K \in \oplus \mathcal{J}^{(fq)}$ or $\oplus \mathcal{J}^{(rq)}$ if H is. \Box

To show that we have a proper hierarchy, we introduce the following notation:

Let $\mathbb{P} = \{P_0, \dots, P_n\}$ be a partition of the set of primes into n+1 infinite sets. For $i \in [n]$, let τ_i be the type which is infinite for all $p \in P_i$ and zero elsewhere. Let $\{v_i : i \in [n]\}$ be independent elements in V, and $U = \bigoplus_i [v_i] \cong \mathbb{Q}^n$.

For all $i \in [n]$, let $b_i = \tau_i v_i$, $(B)_* = \bigoplus_i b_{i*}$. and $T = U/(B)_*$.

Proposition 5.2. For every $n \in \mathbb{N}$ there exist H in $\mathcal{J}^{(rq)} \setminus \mathcal{J}^{(fq)}$ and $K \in \mathcal{J} \setminus \mathcal{J}^{(rq)}$, each of rank n.

Proof. With the notation above, T contains a summand isomorphic to $\bigoplus_{p \in P_0} \mathbb{Z}(p^{\infty})$ and hence a subgroup $A \cong \bigoplus_{p \in P_0} \mathbb{Z}(p)$ and for each $q \in P_0$ a summand $A_q \cong \mathbb{Z}(q^{\infty})$. Note that no element of $(B)_*$ is divisible by all $p \in P_0$ and no element has infinite q-height.

Let $H = A\eta^{-1}$ and $K_q = A_q \eta^{-1}$. Then by Proposition 4.4 (2), H and K_q are strongly indecomposable with range $(B)_*$ and $H/(B)_* \cong A \in \mathcal{J}^{rq} \setminus \mathcal{J}^{fq}$ and $K_q/(B)_* \in \mathcal{J} \setminus \mathcal{J}^{rq}$ as required.

We use the hierarchy to define the Torsion Quotient Decomposition. Let

- $\mathcal{G}^{(fq)} = \{ G \in \mathcal{G} \colon J(G) \in \oplus \mathcal{J}^{(fq)} \}$
- $\mathcal{G}^{(rq)} = \{ G \in \mathcal{G} : J(G) \in \oplus \mathcal{J}^{(rq)} \setminus \oplus \mathcal{J}^{(fq)} \}$ $\mathcal{G}^{(dq)} = \{ G \in \mathcal{G} : J(G) \in \oplus \mathcal{J} \setminus \oplus \mathcal{J}^{(rq)} \}$

Note that $\mathcal{G}^{(fq)}$ is the set of acd groups and all torsion quotients of each $G \in \mathcal{G}^{(dq)}$ contain a divisible summand.

Proposition 5.3. The sets $\mathcal{G}^{(fq)}$, $\mathcal{G}^{(rq)}$ and $\mathcal{G}^{(dq)}$ form a partition of \mathcal{G} and are closed under isomorphism, direct sums and direct summands;

Proof. That they form a partition of \mathcal{G} is clear by definition, and their closure properties follow from Proposition 4.4 (1) and Proposition 5.1. \square

A Torsion Quotient Decomposition of G is a decomposition

$$G = G^{(fq)} \oplus G^{(rq)} \oplus G^{(dq)}$$

where $G^{(fq)} \in \mathcal{G}^{(fq)}$, $G^{(rq)} \in \mathcal{G}^{(rq)}$ and $G^{(dq)} \in \mathcal{G}^{(dq)}$.

Theorem 5.4. Let $G \in \mathcal{G}$. Then G has a Torsion Quotient Decomposition.

If $G = G^{(fq)} \oplus G^{(rq)} \oplus G^{(dq)} = H^{(fq)} \oplus H^{(rq)} \oplus H^{dq}$ are Torsion Quotient Decompositions of G, then $G^{(fq)} \cong H^{(fq)}$, $G^{rq} \cong H^{(rq)}$ and $G^{(dq)} \cong H^{(dq)}$.

Proof. Let $G \in \mathcal{G}$. Let $G^{(fq)}$ be a $\mathcal{G}^{(fq)}$ summand of G of maximal rank, and let D be any complementary summand, so that $D \in \mathcal{G}^{(rq)} \cup \mathcal{G}^{(dq)}$: let $G^{(rq)}$ be a $\mathcal{G}^{(rq)}$ summand of D of maximal rank, and let $G^{(dq)}$ be any complementary summand of $G^{(fq)} \oplus \mathcal{G}^{(rq)}$, so that $G^{(dq)} \in \mathcal{G}^{(dq)}$ and $G = G^{(fq)} \oplus G^{(rq)} \oplus G^{(dq)}$.

Since Jónsson subgroups are unique up to isomorphism, $J(G) \cong J(G^{(fq)}) \oplus$ $J(G^{(rq)}) \oplus J(G^{(dq)}) \cong J(H^{(fq)}) \oplus J(H^{(rq)}) \oplus J(H^{(dq)})$ with $J(G^{(fq)})$ and $J(H^{(fq)}) \in J(H^{(fq)})$ $\oplus \mathcal{G}^{(fq)}, J(G^{(rq)}) \text{ and } J(H^{(rq)}) \in \mathcal{G}^{(rq)} \text{ and } J(G^{(dq)}) \text{ and } J(H^{(dq)}) \in \mathcal{G}^{(dq)}.$

Thus $J(G^{(fq)}) \cong J(H^{(fq)})$ and hence $G^{(fq)} \approx H^{(fq)}$. Since summands of G are pure, $G^{(fq)} \cong H^{(fq)}$, and similarly for the other summands, $G^{(rq)} \cong H^{(rq)}$ and $G^{(dq)} \cong H^{(dq)}.$

Remark 5.5. It follows from the definitions that the Torsion Quotient Decomposition of G respects direct sums, in the sense that if $G = H \oplus K$ then $G^{(fq)} =$ $H^{(fq)} \oplus K^{(fq)}, \ G^{(rq)} = H^{(rq)} \oplus K^{(rq)} \text{ and } G^{(dq)} = H^{(dq)} \oplus K^{(dq)}$

5.1. The Principal and Torsion Quotient Decompositions are Compatible. Since the decompositions are independent of each other, both can be applied, to achieve canonical decompositions with 6 terms. We show that these terms are independent of the order in which the decompositions are applied.

Proposition 5.6. Let $G \in \mathcal{G}$ and let $* \in \{fq, rq, dq\}$. Then $(G_{sd})^{(*)} = (G^{(*)})_{sd}$ and $(G_{ni})^{(*)} = (G^{(*)})_{ni}$

Proof. The subgroup $G_{sd} \cap G^{(*)}$ of G can be considered as both a maximal summand of G_{sd} contained in $\mathcal{G}^{(*)}$, i.e., $(G_{sd})^{(*)}$ and a maximal summand of $G^{(*)}$ contained in \mathcal{G}_{sd} , i.e. $(G^{(*)})_{sd}$.

Consequently, $G_{ni} \cap G^{(*)}$ can be considered as both a complement of a maximal summand of G_{sd} contained in $\mathcal{G}^{(*)}$, i.e., $(G_{ni})^{(*)}$ and a complement of a maximal summand of $G^{(*)}$ contained in \mathcal{G}_{sd} , i.e. $(G^*)_{ni}$.

6. RANKED ATOMIC CATEGORIES

In the preceding development, we used no properties of \mathcal{G} other than the fact that it forms a *ranked atomic category* in the following sense:

Let \mathcal{C} be an additive category having a set of indecomposable objects \mathcal{D} .

(1) C is *atomic* if every object is a coproduct of finitely many objects in D;

(2) \mathcal{C} is ranked if there is a function rank: objects of $\mathcal{C} \to \mathbb{N}$ satisfying

- $\operatorname{rank}(H) = 0$ if and only if $H = \{0\}$,
- $H \cong K$ implies $\operatorname{rank}(H) = \operatorname{rank}(K)$, and
- $\operatorname{rank}(H \oplus K) = \operatorname{rank}(H) + \operatorname{rank}(K).$

The category of finite rank modules over an integral domain, and more generally categories of modules with a composition series over an arbitrary ring, are examples of ranked atomic categories.

Let \mathcal{C} be a ranked atomic category, and \mathcal{D} the set of indecomposable objects of \mathcal{C} . For any subset \mathcal{A} of \mathcal{D} let $Add(\mathcal{A})$ be the closure of \mathcal{A} under isomorphism, direct sums and summands.

A partition $\{\mathcal{A}, \mathcal{B}\}$ of \mathcal{D} is called *summand disjoint* if no non-zero element of $Add(\mathcal{A})$ is isomorphic to an element of $Add(\mathcal{B})$ and *vice versa*.

Proposition 6.1. Let C be a ranked atomic category and $\{A, B\}$ a summand disjoint partition of the indecomposable objects of C. Then

- (1) every object G of C is a direct sum $G = A \oplus B$ with $A \in Add(A)$ and $B \in Add(B)$;
- (2) if also $G = A' \oplus B'$ with $A' \in Add(\mathcal{A})$ and $B' \in Add(\mathcal{B})$, then $A \cong A'$ and $B \cong B'$.

Proof. (1) This is clear if $G \in Add(\mathcal{A})$ or $Add(\mathcal{B})$, so we may assume G has summands from both $Add(\mathcal{A})$ and $Add(\mathcal{B})$. Let \mathcal{C} be the set of all summands of G which are elements of $Add(\mathcal{A})$ and let A be an element of \mathcal{C} of maximal rank, so $G = A \oplus B$ where $0 \neq B$ has no summand isomorphic to an element of $Add(\mathcal{A})$. Then every indecomposable summand of B is in \mathcal{B} , so $B \in Add(\mathcal{B})$.

(2) Each of A, A', B and B' is a direct sum of indecomposables. Let C be an indecomposable summand of A, so $C \in \mathcal{A}$. Now $C \cong D \oplus E$ with $D \leq A'$ and $E \leq B'$. Since C is indecomposable and B' has no summand from $\mathcal{A}, C \cong D$. Thus every indecomposable summand of A, and hence A itself is isomorphic to a summand of A'.

By a symmetrical argument, A' is isomorphic to a summand of A; and similarly, B is isomorphic to a summand of B' and B' to a summand of B. Since rank is preserved by isomorphism, $A \cong A'$ and $B \cong B'$.

It is straightforward to extend the definition of summand disjoint partition to partitions with more than two parts and to prove the corresponding analogues of Proposition 6.1 of such partitions. Thus we conclude:

Theorem 6.2. Let C be a ranked atomic category and $\{A_i : i \in [n]\}$ a summand disjoint partition of its indecomposable objects. Then every object G of C has a decomposition $G = \bigoplus_{i \in [n]} G_i$ with $G_i \in \text{Add}(A_i)$ which is unique up to isomorphism.

PHILL SCHULTZ

Proof. Both existence and uniqueness follow by a routine induction from Proposition 6.1.

References

 [Fuchs, 2015] L. Fuchs, Abelian Groups, Springer Monographs in Mathemaics, Springer 2015.
[Mader, 2000] A. Mader, Almost completely decomposable abelian groups, Gordon and Breach, Algebra, Logic and Applications Vol. 13, Amsterdam, 1999.

[Mader and Schultz, 2018] Completely decomposable direct summands of torsion-free abelian groups of finite rank, PAMS 146, (2018), 93–96.

[Schultz, 2020] Phill Schultz, *Torsion-free abelian groups revisited*, Rend. Sem. Univ. Padova, (to appear 2020).

(phill.schultz@uwa.edu.au) THE UNIVERSITY OF WESTERN AUSTRALIA