

# AS-REGULAR ALGEBRAS FROM ACYCLIC SPHERICAL HELICES

SHINNOSUKE OKAWA AND KAZUSHI UEDA

ABSTRACT. We discuss a construction of AS-regular algebras from acyclic spherical helices, generalizing works of Bondal and Polishchuk [BP93] and Van den Bergh [VdB11].

Let  $\mathcal{C}$  be an idempotent-complete pretriangulated dg category over a field  $\mathbf{k}$ . See e.g. [Kel06] and references therein for basic definitions on dg categories. For  $X, Y \in \mathfrak{Ob}(\mathcal{C})$ , we write the dg vector space of morphisms in  $\mathcal{C}$  and its  $i$ -th cohomology as  $\mathrm{hom}(X, Y)$  and  $\mathrm{Hom}^i(X, Y)$  respectively.

A  $\mathcal{C}$ -module is a dg functor  $\mathcal{C}^{\mathrm{op}} \rightarrow \mathrm{Mod} \mathbf{k}$  to the dg category of dg vector spaces. A dg functor  $F: \mathcal{C} \rightarrow \mathcal{C}'$  will be identified with its *graph bimodule*  $\Gamma_F \in \mathrm{Mod}(\mathcal{C}^{\mathrm{op}} \otimes \mathcal{C}')$ .

Assume that  $\mathcal{C}$  is *proper* in the sense that  $\mathrm{hom}(X, Y) \in \mathrm{per} \mathbf{k}$  for any  $X, Y \in \mathfrak{Ob}(\mathcal{C})$ . Assume further that  $\mathcal{C}$  has a Serre functor  $\mathbb{S}$  in the sense of [BK89]. The graph of the Serre functor is given by

$$(1) \quad \mathrm{hom}(Y, \mathbb{S}(X)) = \mathrm{hom}(X, Y)^*,$$

where  $(-)^* := \mathrm{hom}_{\mathbf{k}}(-, \mathbf{k}): (\mathrm{Mod} \mathbf{k})^{\mathrm{op}} \rightarrow \mathrm{Mod} \mathbf{k}$ .

A prototypical example of an idempotent-complete proper pretriangulated dg category admitting a Serre functor is the category of perfect complexes on a proper Gorenstein scheme  $X$ . The Serre functor is the tensor product with the canonical bundle followed by the shift by the dimension of  $X$ ;  $\mathbb{S}(-) = (-) \otimes \omega_X[\dim X]$ .

Let  $d$  be a natural number. An object  $S$  of  $\mathcal{C}$  is said to be *spherical of dimension  $d$*  if  $\mathbb{S}(S) \simeq S[d]$  and

$$(2) \quad \mathrm{Hom}^i(S, S) \cong \begin{cases} \mathbf{k} & i = 0, d, \\ 0 & \text{otherwise.} \end{cases}$$

The *spherical twist* along  $S$  is the functorial cone  $T_S$  over the evaluation morphism  $\mathrm{ev}: \mathrm{hom}(S, -) \otimes S \rightarrow \mathrm{id}_{\mathcal{C}}$ . It is an autoequivalence of the dg category  $\mathrm{per} \mathcal{C}$  of perfect  $\mathcal{C}$ -modules, whose inverse is given by the *dual twist*  $T_S^{\vee}$  defined as the cone over the coevaluation morphism  $\mathrm{coev}: \mathrm{id}_{\mathcal{C}} \rightarrow \mathrm{hom}(-, S)^{\vee} \otimes S$  shifted by  $-1$  [ST01].

Given a sequence  $\mathbf{S} = (S_i)_{i=1}^{\ell}$  of spherical objects, let  $\mathcal{B} = \mathcal{B}_{\mathbf{S}}$  be the full subcategory of  $\mathcal{C}$  consisting of  $\{S_i\}_{i=1}^{\ell}$ . We identify  $\mathrm{per} \mathcal{B}$  with the full idempotent-complete pretriangulated subcategory of  $\mathcal{C}$  generated by  $\mathcal{B}$ .

**Definition 1.** A sequence  $\mathbf{S} = (S_i)_{i=1}^{\ell}$  of spherical objects is a *spherical collection* if the Serre functor of  $\mathcal{C}$  preserves  $\mathrm{per} \mathcal{B}$  and acts as the shift functor there;  $\mathbb{S}_{\mathcal{C}}|_{\mathrm{per} \mathcal{B}} \simeq [d]$ .

The number  $\ell$  of objects in the collection is called the *length* of the collection.

The *directed subcategory*  $\mathcal{A} = \mathcal{A}_{\mathbf{S}}$  of  $\mathcal{B}$  is the dg category whose set of objects is  $\{E_i\}_{i=1}^{\ell}$  and whose dg vector spaces of morphisms are given by

$$(3) \quad \mathrm{hom}_{\mathcal{A}}(E_i, E_j) := \begin{cases} \mathrm{hom}_{\mathcal{B}}(S_i, S_j) & i < j, \\ \mathbf{k} \cdot \mathrm{id}_{E_i} & i = j, \\ 0 & i > j. \end{cases}$$

The differentials and the compositions of morphisms in  $\mathcal{A}$  are inherited from that of  $\mathcal{B}$  in the obvious way.

Since the collection  $\mathbf{S}$  is spherical, the  $\mathcal{A}$ -bimodule  $\mathcal{B}/\mathcal{A}$ , defined as the cone of the natural morphism  $\mathcal{A} \rightarrow \mathcal{B}$  of  $\mathcal{A}$ -bimodules, is isomorphic to the shift  $\mathcal{A}^*[-d]$  of the graph  $\mathcal{A}^* := \mathrm{hom}_{\mathbf{k}}(\mathcal{A}, \mathbf{k})$  of the Serre functor. In other words, the category  $\mathcal{B}$  is a *noncommutative anticanonical divisor* of  $\mathcal{A}$  in the sense of [Sei17].

The sequence  $(E_i)_{i=1}^\ell$  is a full exceptional collection of per  $\mathcal{A}$ . The existence of a full exceptional collection implies that per  $\mathcal{A}$  is proper and *smooth* (in the sense that the diagonal bimodule is perfect), so that per  $\mathcal{A}$  has a Serre functor.

Let  $\iota_*: \text{per } \mathcal{B} \rightarrow \text{per } \mathcal{A}$  be the dg functor sending a  $\mathcal{B}$ -module to the same module considered as an  $\mathcal{A}$ -module by the embedding  $\mathcal{A} \hookrightarrow \mathcal{B}$ . The graph of the left adjoint functor  $\iota^*: \text{per } \mathcal{A} \rightarrow \text{per } \mathcal{B}$  is  $\mathcal{B}$  considered as an  $\mathcal{A}^{\text{op}} \otimes \mathcal{B}$ -module. The right adjoint functor is given by  $\iota^! = \mathbb{S}_{\mathcal{B}} \circ \iota^* \circ \mathbb{S}_{\mathcal{A}}^{-1} \simeq \iota^* \circ \mathbb{S}_{\mathcal{A}}^{-1}[d]$ .

The *dual twist*  $T_{\iota_*}^\vee$  along  $\iota_*$  is defined as the shift by  $-1$  of the cone over the unit  $\text{id}_{\mathcal{A}} \rightarrow \iota_* \circ \iota^*$  of the adjunction. The graph of the functor  $\iota_* \circ \iota^*$  is  $\mathcal{B}$  considered as an  $\mathcal{A}$ -bimodule, so that the graph of the dual twist is the  $\mathcal{A}$ -bimodule  $\mathcal{B}/\mathcal{A}[-1]$ , which is the graph of  $\mathbb{S}[-d-1]$ . The *twist*  $T_{\iota_*}$  along  $\iota_*$ , defined as the cone over the counit  $\iota_* \circ \iota^! \rightarrow \text{id}_{\mathcal{A}}$  of the adjunction, is right adjoint to  $T_{\iota_*}^\vee$  by [AL17, Proposition 5.3], and hence is quasi-isomorphic to  $\mathbb{S}^{-1}[d+1]$ .

The *dual cotwist*  $C_{\iota_*}^\vee$  along  $\iota_*$ , defined as the cone over the counit  $\iota^* \circ \iota_* \rightarrow \text{id}_{\mathcal{B}}$ , is described explicitly as the (iterated) cone of

$$(4) \quad \text{hom}(S_\ell, -) \otimes \text{hom}(S_{\ell-1}, S_\ell) \otimes \cdots \otimes \text{hom}(S_1, S_2) \otimes S_1 \rightarrow \cdots \\ \rightarrow \bigoplus_{1 \leq i_1 < i_2 \leq \ell} \text{hom}(S_{i_2}, -) \otimes \text{hom}(S_{i_1}, S_{i_2}) \otimes S_{i_1} \rightarrow \bigoplus_{i=1}^{\ell} \text{hom}(S_i, -) \otimes S_i \rightarrow \text{id}_{\mathcal{B}},$$

which is quasi-isomorphic to the composition of spherical twists;

$$(5) \quad C_{\iota_*}^\vee \simeq T_{S_1} \circ T_{S_2} \circ \cdots \circ T_{S_\ell}.$$

Since both  $T_{\iota_*}$  and  $C_{\iota_*}^\vee$  are equivalences, the functor  $\iota_*$  is spherical in the sense of [AL17, Theorem 1.1]. We expect that  $\iota_*$  has a *relative Calabi–Yau structure* in the sense of [BD19], which is a slight strengthening of an identification of the spherical twist with a shift of the inverse Serre functor.

For  $1 \leq i \leq \ell - 1$ , the *right mutation* of a spherical collection  $\mathbf{S} = (S_j)_{j=1}^\ell$  at position  $i$  is defined by

$$(6) \quad R_i(\mathbf{S}) := (S_1, \dots, S_{i-1}, S_{i+1}, T_{S_{i+1}}^\vee S_i, S_{i+2}, \dots, S_\ell).$$

The inverse operation is the *left mutation* defined by

$$(7) \quad L_i(\mathbf{S}) := (S_1, \dots, S_{i-1}, T_{S_i} S_{i+1}, S_i, S_{i+2}, \dots, S_\ell).$$

The *left mutation*  $L_i(\mathcal{A})$  of  $\mathcal{A}$  at position  $i$  is the full subcategory of per  $\mathcal{A}$  consisting of  $E_j$  for  $j \neq i+1$  and

$$(8) \quad L_{E_i} E_{i+1} := \text{Cone} \left( \text{hom}(E_i, E_{i+1}) \otimes E_i \xrightarrow{\text{ev}} E_{i+1} \right).$$

The passage to the directed subcategory commutes with mutations;  $\mathcal{A}_{L_i(\mathbf{S})} \simeq L_i(\mathcal{A}_{\mathbf{S}})$ .

The *helix* generated by an exceptional collection  $(E_i)_{i=1}^\ell$  is the sequence  $(E_i)_{i \in \mathbb{Z}}$  of exceptional objects of per  $\mathcal{A}$  satisfying

$$(9) \quad E_{i-\ell} = L_{E_{i-\ell+1}} \circ L_{E_{i-\ell+2}} \circ \cdots \circ L_{E_{i-1}}(E_i)[-d-1]$$

for any  $i \in \mathbb{Z}$ . The shift is chosen in such a way that  $E_{i-\ell} \cong \mathbb{S}(E_i)[-d-1]$ , so that if per  $\mathcal{A} \simeq D^b \text{coh } X$  for a smooth projective variety  $X$  of dimension  $d+1$ , then one has  $E_{i-\ell} \cong E_i \otimes \omega_X$ . The length  $\ell$  of the exceptional collection generating the helix is called the *period* of the helix. For any  $i \in \mathbb{Z}$ , the exceptional collection  $(E_j)_{j=i}^{i+\ell-1}$  consisting of  $\ell$  consecutive objects in the helix is called a *foundation* of the helix. We say that a helix  $(E_i)_{i \in \mathbb{Z}}$  is *acyclic* if  $\text{Hom}^{\neq 0}(E_i, E_j) = 0$  for any  $-\infty < i < j < \infty$ .

Given an invertible  $\mathcal{A}$ -bimodule  $\theta$ , the *tensor algebra* over  $\mathcal{A}$  is defined by

$$(10) \quad T_{\mathcal{A}}(\theta) := \bigoplus_{i=0}^{\infty} \theta^{\otimes_{\mathcal{A}} i}$$

$$(11) \quad = \mathcal{A} \oplus \theta \oplus (\theta \otimes_{\mathcal{A}} \theta) \oplus \cdots$$

$$(12) \quad \simeq \bigoplus_{i=0}^{\infty} \bigoplus_{j,k=1}^{\ell} \text{hom}(E_j, \theta^i(E_k)).$$

In addition to the cohomological grading coming from that of  $\mathcal{A}$  and  $\theta$ , the tensor algebra  $T_{\mathcal{A}}(\theta)$  has an additional grading such that  $\theta^{\otimes_{\mathcal{A}} i}$  has degree  $i$ . The positively graded part  $T_{\mathcal{A}}(\theta)^+ := \bigoplus_{i=1}^{\infty} \theta^{\otimes_{\mathcal{A}} i} \simeq \theta \otimes_{\mathcal{A}} T_{\mathcal{A}}(\theta)$  is the kernel (the cone shifted by  $-1$ ) of the natural morphism  $\sigma^{\sharp}: T_{\mathcal{A}}(\theta) \rightarrow \mathcal{A}$ .

The category  $\text{per } T_{\mathcal{A}}(\theta)$  is a generalization of the derived category of coherent sheaves on the total space of a line bundle, and the pull-back  $\sigma_*: \text{per } \mathcal{A} \rightarrow \text{per } T_{\mathcal{A}}(\theta)$  along  $\sigma^{\sharp}$  is a generalization of the push-forward along the zero section. Its left adjoint  $\sigma^*: \text{per } T_{\mathcal{A}}(\theta) \rightarrow \text{per } \mathcal{A}$  is given by tensoring  $\mathcal{A}$  over  $T_{\mathcal{A}}(\theta)$ . Since the graph of  $\sigma^* \circ \sigma_*$  is

$$(13) \quad \text{Cone}(\theta \otimes_{\mathcal{A}} T_{\mathcal{A}}(\theta) \rightarrow T_{\mathcal{A}}(\theta)) \otimes_{T_{\mathcal{A}}(\theta)} \mathcal{A} \simeq \text{Cone}(\theta \rightarrow \mathcal{A}),$$

the dual cotwist along  $\sigma_*$  is the autoequivalence  $\theta[1]$  of  $\text{per } \mathcal{A}$ . See also [Seg18] for a related construction.

The push-forward  $\pi^* := (-) \otimes_{\mathcal{A}} T_{\mathcal{A}}(\theta): \text{per } \mathcal{A} \rightarrow \text{per } T_{\mathcal{A}}(\theta)$  along the structure morphism  $\pi^{\sharp}: \mathcal{A} \rightarrow T_{\mathcal{A}}(\theta)$  of the tensor algebra is a generalization of the pull-back along the projection to the base space. For any pair  $(E, F)$  of objects of  $\text{per } \mathcal{A}$ , one has

$$(14) \quad \text{hom}(\sigma_* E, \pi^* F) \simeq \text{hom}(\pi^* E \otimes_{T_{\mathcal{A}}(\theta)} \text{Cone}(\theta \otimes_{\mathcal{A}} T_{\mathcal{A}}(\theta) \rightarrow T_{\mathcal{A}}(\theta)), \pi^* F)$$

$$(15) \quad \simeq \text{hom}(\pi^* E, \pi^* F \otimes_{T_{\mathcal{A}}(\theta)} \text{Cone}(T_{\mathcal{A}}(\theta) \rightarrow \theta^{-1} \otimes_{\mathcal{A}} T_{\mathcal{A}}(\theta))[-1])$$

$$(16) \quad \simeq \text{hom}(\pi^* E, \sigma_*(\theta^{-1}(F)))[-1]$$

$$(17) \quad \simeq \text{hom}(E, \theta^{-1}(F))[-1].$$

The dg category  $\text{gr } T_{\mathcal{A}}(\theta)$  of graded perfect  $T_{\mathcal{A}}(\theta)$ -modules is a generalization of the derived category of equivariant coherent sheaves on the total space of a line bundle with respect to the  $\mathbb{G}_m$ -action by fiberwise dilation. The dg  $\mathbb{Z}$ -algebra  $A = \bigoplus_{i,j=-\infty}^{\infty} A_{ij}$  defined by

$$(18) \quad A_{i+k\ell, j+m\ell} = \text{hom}(\theta^m(E_j), \theta^k(E_i))$$

for  $1 \leq i, j \leq \ell$  and  $-\infty < m \leq k < \infty$  is  $\ell$ -periodic, and the 1-periodic dg  $\mathbb{Z}$ -algebra  $A' = \bigoplus_{i,j=-\infty}^{\infty} A'_{ij}$  defined by

$$(19) \quad A'_{km} = \bigoplus_{i=(k-1)\ell+1}^{k\ell} \bigoplus_{j=(m-1)\ell+1}^{m\ell} A_{ij}$$

is the dg  $\mathbb{Z}$ -algebra associated to the  $\mathbb{Z}$ -graded algebra  $T_{\mathcal{A}}(\theta)$ . One has quasi-equivalences

$$(20) \quad \text{gr } T_{\mathcal{A}}(\theta) \simeq \text{mod } A' \simeq \text{mod } A$$

of dg categories.

Let  $(F_i)_{i=\ell}^1$  be the exceptional collection in  $\text{per } \mathcal{A}$  right dual to  $(E_i)_{i=1}^{\ell}$  so that  $\text{hom}(E_i, F_j) \simeq \mathbf{k}$  if  $i = j$  and 0 otherwise [Bon89, Section 7]. Then  $E_i$  (resp.  $F_i$ ) represents the  $i$ -th projective (resp. simple)  $\mathcal{A}$ -module, and  $\pi^* E_i$  (resp.  $\sigma_* F_i$ ) represents the  $i$ -th projective (resp. simple)  $T_{\mathcal{A}}(\theta)$ -module.

The dg algebra  $T_{\mathcal{A}}(\theta)$  in the case  $\theta = \mathbb{S}^{-1}[d+1]$  is known as the *derived  $(d+1)$ -preprojective algebra*, which is a model of the *trivial Calabi–Yau completion* [Kel11];

$$(21) \quad \Pi_{d+1}(\mathcal{A}) := T_{\mathcal{A}}(\mathbb{S}^{-1}[d+1]).$$

It is given explicitly as

$$(22) \quad \bigoplus_{i=0}^{\infty} \bigoplus_{j,k=1}^{\ell} \text{hom}(E_j, \mathbb{S}^{-i}(E_k)[i(d+1)]) \simeq \bigoplus_{j=1}^{\ell} \bigoplus_{k=1}^{\infty} \text{hom}(E_j, E_k),$$

and also known as the *rolled-up helix dg algebra*. The dg  $\mathbb{Z}$ -algebra  $A$  in this case will be called the *helix dg algebra*.

For  $1 \leq i, j \leq \ell$ , one has

$$(23) \quad \text{hom}(\sigma_* F_i, \pi^* E_j) \simeq \text{hom}(F_i, \mathbb{S}(E_j)[-d-1])[-1]$$

$$(24) \quad \simeq \text{hom}(E_j, F_i)[-d-2]$$

$$(25) \quad \simeq \begin{cases} \mathbf{k}[-d-2] & i = j, \\ 0 & \text{otherwise.} \end{cases}$$

An object of  $\text{gr } T_{\mathcal{A}}(\theta)$  is said to be a *torsion* if the grading is bounded. Let  $\text{tor } T_{\mathcal{A}}(\theta)$  be the full subcategory of  $\text{gr } T_{\mathcal{A}}(\theta)$  consisting of torsion modules, and  $\text{qgr } T_{\mathcal{A}}(\theta) := \text{gr } T_{\mathcal{A}}(\theta) / \text{tor } T_{\mathcal{A}}(\theta)$  be the dg quotient.

**Proposition 2.** *One has an equivalence  $\text{qgr } T_{\mathcal{A}}(\theta) \simeq \text{per } \mathcal{A}$ .*

*Proof.* One has a semiorthogonal decomposition

$$(26) \quad \text{tor } T_{\mathcal{A}}(\theta) = \langle \mathcal{S}_i \rangle_{i \in \mathbb{Z}},$$

where  $\mathcal{S}_i \simeq \text{per } \mathcal{A}$  is the admissible subcategory consisting of graded modules concentrated in degree  $i$ . One has

$$(27) \quad \text{gr } T_{\mathcal{A}}(\theta) = \langle \dots, \mathcal{S}_{-2}, \mathcal{S}_{-1}, (\text{gr } T_{\mathcal{A}}(\theta))_{\geq 0} \rangle$$

just as in [Orl09, Lemma 14], where  $(\text{gr } T_{\mathcal{A}}(\theta))_{\geq 0}$  is the full subcategory of  $\text{gr } T_{\mathcal{A}}(\theta)$  consisting of non-negatively graded modules. It follows from the quasi-isomorphism

$$(28) \quad \mathcal{A} \simeq \text{Cone}(T_{\mathcal{A}}(\theta)^+ \rightarrow T_{\mathcal{A}}(\theta))$$

of graded  $T_{\mathcal{A}}(\theta)$ -modules that the free module  $T_{\mathcal{A}}(\theta)$  is right orthogonal to  $\langle \mathcal{S}_i \rangle_{i \geq 0}$ , and that its shift  $T_{\mathcal{A}}(\theta)(i)$  for any  $i < 0$  is in the full pretriangulated subcategory generated by  $T_{\mathcal{A}}(\theta)$  and  $\langle \mathcal{S}_i \rangle_{i \geq 0}$ . Hence  $T_{\mathcal{A}}(\theta)$  generates the right orthogonal to  $\langle \mathcal{S}_i \rangle_{i \geq 0}$  in  $(\text{gr } T_{\mathcal{A}}(\theta))_{\geq 0}$ , so that  $\text{qgr } T_{\mathcal{A}}(\theta)$  is equivalent to the full subcategory  $\langle T_{\mathcal{A}}(\theta) \rangle$  of  $\text{gr } T_{\mathcal{A}}(\theta)$ , which in turn is equivalent to  $\text{per } \mathcal{A}$ .  $\square$

**Definition 3.** A *spherical helix* generated by a spherical collection  $(S_i)_{i=1}^{\ell}$  is a sequence  $(S_i)_{i \in \mathbb{Z}}$  of spherical objects of  $\mathcal{C}$  satisfying

$$(29) \quad S_{i-\ell} = T_{S_{i-\ell+1}} \circ T_{S_{i-\ell+2}} \circ \dots \circ T_{S_{i-1}}(S_i)[-d-1]$$

for any  $i \in \mathbb{Z}$ .

The length  $\ell$  of the spherical collection generating the spherical helix is called the *period* of the spherical helix. For any  $i \in \mathbb{Z}$ , the spherical collection  $(S_j)_{j=i}^{i+\ell-1}$  consisting of  $\ell$  consecutive objects in the spherical helix is called a *foundation* of the spherical helix  $\mathcal{H}$ .

**Definition 4.** A spherical helix  $(S_i)_{i \in \mathbb{Z}}$  is *acyclic* if  $\text{Hom}^{\neq 0}(S_i, S_j) = 0$  for any  $-\infty < i < j < \infty$ .

One has

$$(30) \quad \text{hom}(S_i, S_j) \simeq \text{hom}(\iota^* E_i, \iota^* E_j)$$

$$(31) \quad \simeq \text{hom}(E_i, \iota_* \iota^* E_j)$$

$$(32) \quad \simeq \text{hom}(E_i, \text{Cone}(E_{j-\ell} \rightarrow E_j)),$$

which immediately yields the following:

**Theorem 5.** *If the spherical helix  $(S_i)_{i \in \mathbb{Z}}$  is acyclic, then the helix  $(E_i)_{i \in \mathbb{Z}}$  is acyclic.*

Recall that a connected  $\mathbb{Z}$ -algebra  $A = \bigoplus_{i,j=-\infty}^{\infty} A_{ij}$  over  $\mathbf{k}$  is said to be *AS-Gorenstein* if

$$(33) \quad \sum_{j,k=-\infty}^{\infty} \dim \text{Hom}^k(S_{i,A}, P_{j,A}) = 1$$

for any  $i \in \mathbb{Z}$ , where  $S_{i,A}$  and  $P_{i,A}$  are the  $i$ -th simple and projective  $A$ -modules respectively. It follows from (25) that the acyclicity of the helix  $(E_i)_{i \in \mathbb{Z}}$  implies the AS-Gorenstein property of the helix  $\mathbb{Z}$ -algebra  $A$ . Since  $\mathcal{A}$  is smooth and  $\theta$  is perfect, the tensor algebra  $T_{\mathcal{A}}(\theta)$  is smooth, so

that  $A$  has finite global dimension (see e.g. [Kel11, Theorem 4.8]). The Hilbert polynomial of  $A$  is determined by the projective resolutions of the simples.

*Acknowledgement:* S. O. was partially supported by JSPS Grant-in-Aid for Scientific Research No.16H05994, 16H02141, 16H06337, 18H01120, 20H01797. K. U. was partially supported by JSPS Grant-in-Aid for Scientific Research No.16H03930.

## REFERENCES

- [AL17] Rina Anno and Timothy Logvinenko, *Spherical DG-functors*, J. Eur. Math. Soc. (JEMS) **19** (2017), no. 9, 2577–2656. MR 3692883
- [BD19] Christopher Brav and Tobias Dyckerhoff, *Relative Calabi-Yau structures*, Compos. Math. **155** (2019), no. 2, 372–412. MR 3911626
- [BK89] A. I. Bondal and M. M. Kapranov, *Representable functors, Serre functors, and reconstructions*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 6, 1183–1205, 1337. MR MR1039961 (91b:14013)
- [Bon89] A. I. Bondal, *Representations of associative algebras and coherent sheaves*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 1, 25–44. MR MR992977 (90i:14017)
- [BP93] A. I. Bondal and A. E. Polishchuk, *Homological properties of associative algebras: the method of helices*, Izv. Ross. Akad. Nauk Ser. Mat. **57** (1993), no. 2, 3–50. MR 1230966 (94m:16011)
- [Kel06] Bernhard Keller, *On differential graded categories*, International Congress of Mathematicians. Vol. II, Eur. Math. Soc., Zürich, 2006, pp. 151–190. MR MR2275593 (2008g:18015)
- [Kel11] ———, *Deformed Calabi-Yau completions*, J. Reine Angew. Math. **654** (2011), 125–180, With an appendix by Michel Van den Bergh. MR 2795754
- [Orl09] Dmitri Orlov, *Derived categories of coherent sheaves and triangulated categories of singularities*, Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II, Progr. Math., vol. 270, Birkhäuser Boston Inc., Boston, MA, 2009, pp. 503–531. MR 2641200 (2011c:14050)
- [Seg18] Ed Segal, *All autoequivalences are spherical twists*, Int. Math. Res. Not. IMRN (2018), no. 10, 3137–3154. MR 3805198
- [Sei17] Paul Seidel, *Fukaya  $A_\infty$ -structures associated to Lefschetz fibrations. II*, Algebra, geometry, and physics in the 21st century, Progr. Math., vol. 324, Birkhäuser/Springer, Cham, 2017, pp. 295–364. MR 3727564
- [ST01] Paul Seidel and Richard Thomas, *Braid group actions on derived categories of coherent sheaves*, Duke Math. J. **108** (2001), no. 1, 37–108. MR MR1831820 (2002e:14030)
- [VdB11] Michel Van den Bergh, *Noncommutative quadrics*, International Mathematics Research Notices. IMRN (2011), no. 17, 3983–4026. MR MR2836401 (2012m:14004)

DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, MACHIKANNEYAMA 1-1, TOYONAKA, OSAKA, 560-0043, JAPAN.

*E-mail address:* okawa@math.sci.osaka-u.ac.jp

GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-1 KOMABA, MEGURO-KU, TOKYO, 153-8914, JAPAN.

*E-mail address:* kazushi@ms.u-tokyo.ac.jp