

# New partition function recurrences

Robson da Silva  
 Universidade Federal de São Paulo  
 São José dos Campos - SP, 12247-014, Brazil  
 silva.robson@unifesp.br

Pedro Diniz Sakai  
 Universidade Federal de São Paulo  
 São José dos Campos - SP, 12247-014, Brazil  
 pedro.sakai@unifesp.br

## Abstract

We present Euler-type recurrence relations for some partition functions. Some of our results present new recurrences for the number of unrestricted partitions of  $n$ , denote by  $p(n)$ . Others establish recurrences for partition functions not yet considered.

**keywords:** Partition, overpartition, recurrence relation, generating function  
**MSC:** 05A17, 11P81

## 1 Introduction

A partition of an integer  $n$  is a finite set of positive integers  $\{\lambda_1, \dots, \lambda_s\}$  such that  $n = \lambda_1 + \dots + \lambda_s$ . The  $\lambda_i$ s are called the parts of the partition. The number of partitions of  $n$  is usually denote by  $p(n)$ , with  $p(0) = 1$  by convention. For example, we have  $p(4) = 5$  since there are five partitions of 4, namely:

$$4, 3 + 1, 2 + 2, 2 + 1 + 1, 1 + 1 + 1 + 1.$$

The generating function of  $p(n)$ , due to Euler, is given by (see [1, eq. (1.1.6)])

$$\sum_{n=0}^{\infty} p(n)q^n = \prod_{k=1}^{\infty} \frac{1}{1 - q^k}. \quad (1)$$

So, one can obtain the values of  $p(n)$  by expanding the right-hand side of (1) and extracting the coefficient of  $q^n$ . Another way to obtain  $p(n)$  was found by Euler after he proved the following identity (known as Euler's pentagonal number theorem):

$$\sum_{n=-\infty}^{\infty} (-1)^n q^{n(3n-1)/2} = \prod_{k=1}^{\infty} (1 - q^k). \quad (2)$$

Indeed, multiplying (1) and (2) we obtain

$$\sum_{n=0}^{\infty} p(n)q^n \sum_{n=-\infty}^{\infty} (-1)^n q^{n(3n-1)/2} = 1,$$

from which the following recurrence for  $p(n)$  is derived after extracting the coefficient of  $q^n$  from its both sides:

$$\begin{aligned} p(n) - p(n-1) - p(n-2) + p(n-5) + p(n-7) - p(n-12) - p(n-15) \\ + \cdots + (-1)^j p(n-j(3j-1)/2) + (-1)^j p(n-j(3j+1)/2) + \cdots \\ = \begin{cases} 1, & \text{if } n = 0, \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

The numbers  $j(3j \pm 1)/2$  are the pentagonal numbers.

Some subsequent works bring new recurrence relations for  $p(n)$  and other partition functions. Ewell [4, Theorem 2], for instance, presents the following recurrence for  $p(n)$  involving the triangular numbers

$$\begin{aligned} p(n) - p(n-1) - p(n-3) + p(n-6) + p(n-10) - p(n-15) - p(n-21) \\ + \cdots + (-1)^j p(n-j(2j-1)) + (-1)^j p(n-j(2j+1)) + \cdots \\ = \begin{cases} 0, & \text{if } n \text{ is odd,} \\ q(n/2), & \text{if } n \text{ is even,} \end{cases} \end{aligned}$$

where  $q(n)$  denotes the number of partitions of  $n$  into distinct parts. Merca [6] derived two new recurrence relations for  $p(n)$ , which allowed him to obtain a more efficient method to compute the parity of  $p(n)$ . Ono, Robbins, and Wilson [8] presented recurrence relations for some partition functions, including  $q(n)$ ,  $q_O(n)$  the number of partitions into distinct and odd parts,  $p_E(n)$  the number of partitions into an even number of parts, and  $p_O(n)$  the number of partitions into an odd number of parts. Recently, Choliy, Kolitsch, and Sills [2] found a number of new recurrences for  $p(n)$ , including

$$\begin{aligned} p(n) - p(n-1) - p(n-2) + p(n-4) + p(n-8) - p(n-9) - p(n-18) \\ + \cdots + (-1)^j p(n-j^2) + (-1)^j p(n-2j^2) + \cdots \\ = \begin{cases} 0, & \text{if } n \text{ is odd,} \\ qq(n), & \text{if } n \text{ is even,} \end{cases} \end{aligned}$$

and

$$\begin{aligned} p(n) - 2p(n-1) + 2p(n-4) - 2p(n-9) + 2p(n-16) + \cdots \\ + (-1)^j 2p(n-j^2) + \cdots = (-1)^n qq(n). \end{aligned}$$

where  $qq(n)$  denotes the number of partitions of  $n$  into distinct odd parts. Additional recurrence relations for partition functions can be found in [2, 4, 5, 6, 7, 8].

In this paper, using some classical identities and generating function manipulations, we provide a number of new recurrence relations for  $p(n)$ ,  $qq(n)$ ,  $\bar{p}(n)$  the number of overpartitions of  $n$ ,  $p_o(n)$  the number of partitions of  $n$  into odd parts, and  $p_m^c(n)$  the number of partitions of  $n$  into parts congruent to  $\pm c$  modulo  $m$ . For some of these functions, it is the first time that recurrence relations are presented.

## 2 Preliminaries

We recall Ramanujan's theta functions

$$f(a, b) := \sum_{n=-\infty}^{\infty} a^{\frac{n(n+1)}{2}} b^{\frac{n(n-1)}{2}}, \text{ for } |ab| < 1, \quad (3)$$

$$\psi(q) := f(q, q^3) = \sum_{n=0}^{\infty} q^{n(n+1)/2}. \quad (4)$$

In the proofs of some of our results, we will need Jacobi triple product identity given by (see [1, Theorem 1.3.3])

$$\sum_{n=-\infty}^{\infty} z^n q^{n^2} = (-zq; q^2)_{\infty} (-q/z; q^2)_{\infty} (q^2; q^2)_{\infty}.$$

where we use the following standard  $q$ -series notation:

$$\begin{aligned} (a; q)_0 &= 1, \\ (a; q)_n &= (1-a)(1-aq) \cdots (1-aq^{n-1}), \forall n \geq 1, \\ (a; q)_{\infty} &= \lim_{n \rightarrow \infty} (a; q)_n, |q| < 1. \end{aligned}$$

Using (3), we can rewrite Jacobi triple product identity in the form

$$f(a, b) = (-a; ab)_{\infty} (-b; ab)_{\infty} (ab; ab)_{\infty}. \quad (5)$$

An important consequence of (5) is following identity (see [1, Eq. (1.3.14)])

$$\psi(q) = \frac{(q^2; q^2)_{\infty}}{(q; q^2)_{\infty}}. \quad (6)$$

We also recall the well-known Euler's pentagonal number theorem (see [1, Corollary 1.3.5]):

$$(q; q)_{\infty} = \sum_{n=-\infty}^{\infty} (-1)^n q^{n(3n-1)/2}. \quad (7)$$

### 3 Main results

In what follows, we let  $t_j^e$  (resp.  $t_j^o$ ) denote the  $j$ -th even (resp. odd) triangular number. So,  $t_1^e = 0$ ,  $t_1^o = 1$ ,  $t_2^e = 3$ ,  $t_2^o = 6$ ,  $t_3^e = 10$ ,  $t_3^o = 15$ , etc.

**Theorem 1.** *For all even integer  $n \geq 1$ , we have*

$$p(n/2) + p((n-6)/2) + p((n-10)/2) + p((n-28)/2) + p((n-36)/2) + p((n-66)/2) + p((n-78)/2) + \cdots + p((n-t_j^e)/2) + \cdots = p_d(n), \quad (8)$$

where  $p_d(n)$  denotes the number of partitions of  $n$  into distinct parts. For all odd integer  $n \geq 1$ , we have

$$p((n-1)/2) + p((n-3)/2) + p((n-15)/2) + p((n-21)/2) + p((n-45)/2) + p((n-55)/2) + \cdots + p((n-t_j^o)/2) + \cdots = p_o(n), \quad (9)$$

where  $p_o(n)$  denotes the number of partitions of  $n$  into odd parts

*Proof.* Initially, we note that

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{1+(-1)^n}{2} p_d(n) q^n &= \frac{1}{2} \left( \sum_{n=0}^{\infty} p_d(n) q^n + \sum_{n=0}^{\infty} p_d(n) (-q)^n \right) \\ &= \frac{1}{2} \left( \prod_{k=1}^{\infty} (1+q^k) + \prod_{k=1}^{\infty} (1+(-1)^k q^k) \right) \\ &= \frac{1}{2} \left( \prod_{k=1}^{\infty} (1+q^k) + \prod_{k=1}^{\infty} (1+q^{2k})(1-q^{2k-1}) \right) \end{aligned}$$

and

$$\begin{aligned} \prod_{k=1}^{\infty} (1+q^k) &= \prod_{k=1}^{\infty} (1+q^k) \frac{(1-q^k)}{(1-q^k)} \frac{(1-q^{2k-1})}{(1-q^{2k-1})} \\ &= \prod_{k=1}^{\infty} \frac{(1-q^{2k-1})(1-q^{2k})}{(1-q^k)(1-q^{2k-1})} = \prod_{k=1}^{\infty} \frac{1}{1-q^{2k}} \psi(q). \end{aligned}$$

We also have

$$\begin{aligned} \prod_{k=1}^{\infty} (1+q^{2k})(1-q^{2k-1}) &= \prod_{k=1}^{\infty} (1+q^{2k})(1-q^{2k-1}) \frac{(1-q^{2k})}{(1-q^k)(1+q^k)} \\ &= \prod_{k=1}^{\infty} \frac{(1-q^{2k-1})(1-q^{2k})}{(1-q^k)(1+q^{2k-1})} = \prod_{k=1}^{\infty} \frac{1}{1-q^{2k}} \psi(-q). \end{aligned}$$

It follows that

$$\begin{aligned}
\sum_{n=0}^{\infty} \frac{1+(-1)^n}{2} p_d(n) q^n &= \frac{1}{2} \left( \prod_{k \geq 1} \frac{1}{1-q^{2k}} \psi(q) + \prod_{k=1}^{\infty} \frac{1}{1-q^{2k}} \psi(-q) \right) \\
&= \left( \prod_{k=1}^{\infty} \frac{1}{1-q^{2k}} \right) \frac{1}{2} \left( \psi(q) + \psi(-q) \right) \\
&= \left( \prod_{k=1}^{\infty} \frac{1}{1-q^{2k}} \right) \frac{1}{2} \left( \sum_{j=0}^{\infty} q^{\frac{j(j+1)}{2}} + \sum_{j=0}^{\infty} (-q)^{\frac{j(j+1)}{2}} \right).
\end{aligned}$$

The parity of the exponent  $\frac{j(j+1)}{2}$  is given by

$$\frac{4i(4i+1)}{2} = 8i^2 + 2i, \quad (10)$$

$$\frac{(4i-1)(4i-1+1)}{2} = 8i^2 - 2i, \quad (11)$$

$$\frac{(4i-2)(4i-2+1)}{2} = 8i^2 - 6i + 1, \quad (12)$$

$$\frac{(4i-3)(4i-3+1)}{2} = 8i^2 - 10i + 3. \quad (13)$$

The even triangular numbers are given by (10) and (11), while (12) and (13) represent the odd triangular numbers. Thus

$$\sum_{j=0}^{\infty} q^{\frac{j(j+1)}{2}} = \sum_{i=0}^{\infty} q^{8i^2+2i} + q^{8i^2-2i} + q^{8i^2-6i+1} + q^{8i^2-10i+3} \quad (14)$$

and

$$\sum_{j=0}^{\infty} (-q)^{\frac{j(j+1)}{2}} = \sum_{i=0}^{\infty} q^{8i^2+2i} + q^{8i^2-2i} - q^{8i^2-6i+1} - q^{8i^2-10i+3}, \quad (15)$$

which yields

$$\begin{aligned}
\sum_{n=0}^{\infty} \frac{1+(-1)^n}{2} p_d(n) q^n &= \left( \prod_{k=1}^{\infty} \frac{1}{1-q^{2k}} \right) \frac{1}{2} \left( \sum_{i=0}^{\infty} 2q^{8i^2+2i} + 2q^{8i^2-2i} \right) \\
&= \left( \sum_{k=0}^{\infty} p(k) q^{2k} \right) \left( \sum_{j=0}^{\infty} q^{t_j^e} \right) \\
&= \sum_{k=0}^{\infty} \left( \sum_{j=0}^{\infty} p(k) q^{2k+t_j^e} \right).
\end{aligned}$$

Now we extract the coefficient of  $q^n$  on both sides of the above equation to obtain

$$\sum_{n=0}^{\infty} p_d(n) q^n = \sum_{n=0}^{\infty} \left( \sum_{j=0}^{\infty} p((n-t_j^e)/2) \right) q^n,$$

which completes the proof of (8).

In order to prove (9), we begin with

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{1 - (-1)^n}{2} p_o(n) q^n &= \frac{1}{2} \left( \sum_{n=0}^{\infty} p_o(n) q^n - \sum_{n=0}^{\infty} p_o(n) (-q)^n \right) \\ &= \frac{1}{2} \left( \prod_{k=1}^{\infty} \frac{1}{(1 - q^{2k-1})} - \prod_{k=1}^{\infty} \frac{1}{(1 - (-q)^{2k-1})} \right) \\ &= \frac{1}{2} \left( \prod_{k=1}^{\infty} \frac{1}{(1 - q^{2k-1})} - \prod_{k=1}^{\infty} \frac{1}{(1 + q^{2k-1})} \right). \end{aligned}$$

We note that

$$\prod_{k=1}^{\infty} \frac{1}{(1 - q^{2k-1})} = \prod_{k=1}^{\infty} \frac{1}{(1 - q^{2k-1})} \frac{(1 - q^{2k})}{(1 - q^{2k})} = \prod_{k=1}^{\infty} \frac{1}{1 - q^{2k}} \psi(q)$$

and

$$\prod_{k=1}^{\infty} \frac{1}{(1 + q^{2k-1})} = \prod_{k=1}^{\infty} \frac{1}{(1 + q^{2k-1})} \frac{(1 - q^{2k})}{(1 - q^{2k})} = \prod_{k=1}^{\infty} \frac{1}{1 - q^{2k}} \psi(-q).$$

It follows that

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{1 - (-1)^n}{2} p_o(n) q^n &= \frac{1}{2} \left( \prod_{k=1}^{\infty} \frac{1}{1 - q^{2k}} \psi(q) - \prod_{k=1}^{\infty} \frac{1}{1 - q^{2k}} \psi(-q) \right) \\ &= \left( \prod_{k=1}^{\infty} \frac{1}{1 - q^{2k}} \right) \frac{1}{2} \left( \psi(q) - \psi(-q) \right) \\ &= \left( \prod_{k=1}^{\infty} \frac{1}{1 - q^{2k}} \right) \frac{1}{2} \left( \sum_{j=0}^{\infty} q^{\frac{j(j+1)}{2}} - \sum_{j=0}^{\infty} (-q)^{\frac{j(j+1)}{2}} \right). \end{aligned}$$

By (14) and (15), we have

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{1 - (-1)^n}{2} p_o(n) q^n &= \left( \prod_{k=1}^{\infty} \frac{1}{1 - q^{2k}} \right) \frac{1}{2} \left( \sum_{i=0}^{\infty} 2q^{8i^2 - 6i + 1} + 2q^{8i^2 - 10i + 3} \right) \\ &= \sum_{k=0}^{\infty} p(k) q^{2k} \sum_{j=0}^{\infty} q^{t_j^o} = \sum_{k=0}^{\infty} \sum_{j=0}^{\infty} p(k) q^{2k + t_j^o}. \end{aligned}$$

Extracting the coefficient of  $q^n$  in the identity above, we obtain

$$\sum_{n=0}^{\infty} p_o(n) q^n = \sum_{n=0}^{\infty} \left( \sum_{j=0}^{\infty} p((n - t_j^o)/2) \right) q^n,$$

from which (9) follows.  $\square$

We recall that an overpartition of  $n$ , introduced in [3], is a partition in which the first occurrence of a number may be overlined. For instance, there are eight overpartitions of 3:

$$3, \overline{3}, 2 + 1, \overline{2} + 1, 2 + \overline{1}, \overline{2} + \overline{1}, 1 + 1 + 1, \overline{1} + 1 + 1.$$

In the next three results, we present recurrence relations for the number of overpartitions of  $n$ , denoted by  $\overline{p}(n)$ .

**Theorem 2.** *For all  $n \geq 0$ , we have*

$$\begin{aligned} & \overline{p}(n) - 2\overline{p}(n-1) + 2\overline{p}(n-4) - 2\overline{p}(n-9) + 2\overline{p}(n-16) - 2\overline{p}(n-25) + \\ & 2\overline{p}(n-36) - \cdots + 2(-1)^j \overline{p}(n-j^2) + \cdots = \begin{cases} 1, & \text{if } n = 0, \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

*Proof.* We recall from [3] that the generating function for overpartitions is given by

$$\sum_{n=0}^{\infty} \overline{p}(n) q^n = \prod_{k=1}^{\infty} \frac{(1+q^k)}{(1-q^k)}.$$

We note that

$$\begin{aligned} \prod_{k=1}^{\infty} (1-q^k) &= \prod_{k=1}^{\infty} (1-q^k) \frac{(1-q^{2k-1})}{(1-q^{2k-1})} \frac{(1-q^{2k})}{(1-q^{2k})} \\ &= \prod_{k=1}^{\infty} \frac{(1-q^k)(1-q^{2k-1})(1-q^{2k})}{(1-q^{2k})(1-q^{2k-1})} \\ &= \prod_{k=1}^{\infty} \frac{(1-q^{2k-1})^2(1-q^{2k})}{(1-q^{2k-1})} \\ &= \varphi(-q) \prod_{k=1}^{\infty} \frac{1}{(1-q^{2k-1})}. \end{aligned}$$

By Euler's identity, we have

$$\prod_{k=1}^{\infty} (1-q^k) = \varphi(-q) \prod_{k=1}^{\infty} (1+q^k),$$

and, then,

$$\prod_{k=1}^{\infty} \frac{1}{(1-q^k)} = \frac{1}{\varphi(-q)} \prod_{k=1}^{\infty} \frac{1}{(1+q^k)}.$$

Hence we obtain the following equivalent equalities

$$\begin{aligned}
\sum_{n=0}^{\infty} \bar{p}(n)q^n &= \frac{\prod_{k=1}^{\infty} (1+q^k)}{\varphi(-q) \prod_{k=1}^{\infty} (1+q^k)} \\
\varphi(-q) \sum_{n=0}^{\infty} \bar{p}(n)q^n &= 1 \\
\left(1 + 2 \sum_{k=1}^{\infty} (-1)^k q^{k^2}\right) \sum_{n=0}^{\infty} \bar{p}(n)q^n &= 1 \\
\sum_{n=0}^{\infty} \left(\bar{p}(n)q^n + 2 \sum_{k=1}^{\infty} (-1)^k \bar{p}(n)q^{k^2+n}\right) &= 1 \\
\sum_{n=0}^{\infty} \left(\bar{p}(n) + 2 \sum_{k=1}^{\infty} (-1)^k \bar{p}(n-k^2)\right) q^n &= 1.
\end{aligned}$$

The result now follows from the last equality.  $\square$

Our second recurrence for  $\bar{p}(n)$  involves  $p_d(n)$ .

**Theorem 3.** *For all  $n \geq 1$ , we have*

$$\begin{aligned}
&\bar{p}(n) - \bar{p}(n-1) - \bar{p}(n-2) + \bar{p}(n-5) + \bar{p}(n-7) - \dots \\
&\dots + (-1)^j (\bar{p}(n-j(3j-1)/2) + \bar{p}(n-j(3j+1)/2)) + \dots = p_d(n).
\end{aligned}$$

*Proof.* We have

$$\begin{aligned}
\prod_{k=1}^{\infty} (1+q^k) &= \prod_{k=1}^{\infty} \frac{(1+q^k)}{(1-q^k)} \prod_{k=1}^{\infty} (1-q^k) \\
&= \sum_{k=1}^{\infty} \bar{p}(k)q^k \sum_{k=-\infty}^{\infty} (-1)^j q^{\frac{j(3j-1)}{2}} \\
&= \sum_{k=0}^{\infty} \bar{p}(k)q^k \left(1 + \sum_{j=1}^{\infty} (-1)^j q^{\frac{j(3j-1)}{2}} + \sum_{j=1}^{\infty} (-1)^j q^{\frac{j(3j+1)}{2}}\right) \\
&= \sum_{k=0}^{\infty} \left(\bar{p}(k)q^k + \sum_{j=1}^{\infty} (-1)^j \bar{p}(k)q^{k+\frac{j(3j-1)}{2}} + \sum_{j=1}^{\infty} (-1)^j \bar{p}(k)q^{k+\frac{j(3j+1)}{2}}\right).
\end{aligned}$$

Therefore,

$$\sum_{n=0}^{\infty} p_d(n)q^n = \sum_{n=0}^{\infty} \left(\bar{p}(n) + \sum_{j=1}^{\infty} (-1)^j (\bar{p}(n-j(3j-1)/2) + \bar{p}(n-j(3j+1)/2))\right) q^n,$$

from which the proof follows by comparing coefficients of  $q^n$  on both sides of the last equation.  $\square$



If  $\overline{p}_d(n)$  denotes the number of overpartitions of  $n$  into distinct parts, then we have the following recurrence for  $\overline{p}(n)$ .

**Theorem 4.** *For all  $n \geq 0$ , we have*

$$\begin{aligned} & \overline{p}(n) - \overline{p}(n-2) - \overline{p}(n-4) + \overline{p}(n-10) + \overline{p}(n-14) - \dots \\ & \dots + (-1)^j (\overline{p}(n-j(3j-1)) + \overline{p}(n-j(3j+1))) + \dots = \overline{p}_d(n). \end{aligned}$$

*Proof.* By (7) we have

$$\begin{aligned} \sum_{k=0}^{\infty} \overline{p}_d(n) q^n &= \prod_{k=1}^{\infty} (1+q^k)^2 = \prod_{k=1}^{\infty} \frac{(1-q^{2k})(1+q^k)}{1-q^k} \\ &= (q^2; q^2)_{\infty} \sum_{k=0}^{\infty} \overline{p}(k) q^k = \sum_{k=0}^{\infty} \overline{p}(k) q^k \sum_{j=-\infty}^{\infty} (-1)^j q^{j(3j-1)} \\ &= \sum_{k=0}^{\infty} \overline{p}(k) q^k \left( 1 + \sum_{j=1}^{\infty} (-1)^j q^{j(3j-1)} + \sum_{j=1}^{\infty} (-1)^j q^{j(3j+1)} \right) \\ &= \sum_{n=0}^{\infty} \left( \overline{p}(n) + \sum_{j=1}^{\infty} (-1)^j (\overline{p}(n-j(3j-1)) + \overline{p}(n-j(3j+1))) \right) q^n. \end{aligned}$$

Thus, the result follows from extracting the coefficient of  $q^n$  on both sides of the last equality.  $\square$

Now we prove a recurrence relations satisfied by  $qq(n)$ , the number of partitions of  $n$  into distinct odd parts.

**Theorem 5.** *For all  $n \geq 1$ , we have*

$$\begin{aligned} & qq(n) - qq(n-4) - qq(n-8) + qq(n-20) + qq(n-28) - qq(n-48) - \\ & qq(n-60) + \dots + (-1)^j (qq(n-2j(3j-1)) + qq(n-2j(3j+1))) \\ & \dots = \begin{cases} 1, & \text{if } n \text{ is a triangular number,} \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

*Proof.* It is easy to see that

$$\prod_{k=1}^{\infty} (1+q^{2k-1}) = \prod_{k=1}^{\infty} \frac{(1+q^k)}{(1+q^{2k})} = \prod_{k=1}^{\infty} \frac{(1+q^k)(1-q^{2k})}{(1-q^{4k})}.$$

Thus,

$$\sum_{j=0}^{\infty} qq(j) q^j = \prod_{k=1}^{\infty} \frac{(1+q^k)(1-q^{2k})}{(1-q^{4k})},$$

which can be rewritten as

$$\prod_{k=1}^{\infty} (1 - q^{4k}) \sum_{j=0}^{\infty} qq(j)q^j = \sum_{i=0}^{\infty} p_d(i)q^i \prod_{k \geq 1} (1 - q^{2k}).$$

That is to say

$$(q^4; q^4)_{\infty} \sum_{j=0}^{\infty} qq(j)q^j = (q^2; q^2)_{\infty} \sum_{i=0}^{\infty} p_d(i)q^i.$$

Then, by (7), we have

$$\begin{aligned} (q^4; q^4)_{\infty} \sum_{j=0}^{\infty} qq(j)q^j &= \left( 1 + \sum_{k=1}^{\infty} (-1)^k q^{2j(3j-1)} + \sum_{k=1}^{\infty} (-1)^k q^{2j(3j+1)} \right) \sum_{j=0}^{\infty} qq(j)q^j \\ &= \sum_{n=0}^{\infty} \left( qq(n) + \sum_{k=0}^{\infty} (-1)^k (qq(n - 2j(3j - 1)) + qq(n - 2j(3j + 1))) \right) q^n. \end{aligned}$$

On the other hand, we have

$$\begin{aligned} (q^2; q^2)_{\infty} \sum_{i=0}^{\infty} p_d(i)q^i &= \sum_{i=0}^{\infty} p_d(i)q^i \left( 1 + \sum_{k=1}^{\infty} (-1)^k q^{k(3k-1)} + \sum_{k=1}^{\infty} (-1)^k q^{k(3k+1)} \right) \\ &= \sum_{n=0}^{\infty} \left( p_d(n) + \sum_{k=1}^{\infty} (-1)^k (p_d(n - k(3k - 1)) + p_d(n - k(3k + 1))) \right) q^n. \end{aligned}$$

Hence,

$$\begin{aligned} &\sum_{n=0}^{\infty} \left( qq(n) + \sum_{k=1}^{\infty} (-1)^k (qq(n - 2j(3j - 1)) + qq(n - 2j(3j + 1))) \right) q^n \\ &= \sum_{n=0}^{\infty} \left( p_d(n) + \sum_{k=1}^{\infty} (-1)^k (p_d(n - k(3k - 1)) + p_d(n - k(3k + 1))) \right) q^n. \end{aligned}$$

The result follows from extracting the coefficient of  $q^n$  on both sides of the last equation and using Theorem 1 of [8].  $\square$

The next theorem presents a recurrence for the number of partitions into odd parts, denoted by  $p_o(n)$ .

**Theorem 6.** For all  $n \geq 0$ , we have

$$\begin{aligned} & p_o(n) - p_o(n-1) - p_o(n-5) + p_o(n-8) + p_o(n-16) - \cdots \\ & \cdots + (-1)^j [p_o(n-j(3j-2)) + p_o(n-j(3j+2))] + \cdots \\ & \cdots = \begin{cases} 1, & \text{if } n \text{ is 3 times a triangular number,} \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

*Proof.* Setting  $a = -q$  and  $b = -q^5$  in (3) and (5) we obtain

$$(q; q^6)_\infty (q^5; q^6)_\infty (q^6; q^6)_\infty = f(-q, -q^5) = \sum_{j=-\infty}^{\infty} (-q)^{\frac{j(j+1)}{2}} (-q^5)^{\frac{j(j-1)}{2}},$$

from which it follows that

$$\begin{aligned} \sum_{j=-\infty}^{\infty} (-1)^j q^{j(3j-2)} &= \prod_{k=1}^{\infty} (1 - q^{6k-5})(1 - q^{6k-1})(1 - q^{6k}) \\ &= \prod_{k=1}^{\infty} (1 - q^{6k-5})(1 - q^{6k-1})(1 - q^{6k}) \frac{(1 - q^{6k-3})}{(1 - q^{6k-3})} \\ &= \prod_{k=1}^{\infty} \frac{(1 - q^{2k-1})(1 - q^{6k})}{(1 - q^{6k-3})}. \end{aligned} \quad (16)$$

Then, by (4), we have

$$\begin{aligned} \psi(q^3) &= \prod_{k=1}^{\infty} \frac{1}{(1 - q^{2k-1})} \sum_{j=-\infty}^{\infty} (-1)^j q^{j(3j-2)} \\ &= \sum_{i=0}^{\infty} p_o(i) q^i \left( 1 + \sum_{j=1}^{\infty} (-1)^j q^{j(3j-2)} + \sum_{j=1}^{\infty} (-1)^j q^{j(3j+2)} \right) \\ &= \sum_{n=0}^{\infty} \left( p_o(n) + \sum_{j=1}^{\infty} (-1)^j (p_o(n-j(3j-2)) + p_o(n-j(3j+2))) \right) q^n. \end{aligned}$$

The result follows by comparing the coefficients of  $q^n$  on both sides of the last expression.  $\square$

Let  $\ell$  be a positive integer. A partition of  $n$  having no part divisible by  $\ell$  is called an  $\ell$ -regular partition of  $n$ . Let  $b_\ell(n)$  denote the number of  $\ell$ -regular partition of  $n$ . The generating function of  $b_\ell(n)$  is

$$\sum_{n=0}^{\infty} b_\ell(n) q^n = \frac{(q^\ell; q^\ell)_\infty}{(q; q)_\infty}.$$

Our next result is a recurrence relation for  $p(n)$  involving  $b_\ell(n)$ .

**Theorem 7.** Let  $\ell \geq 1$ . For all  $n \geq 0$ , we have

$$\begin{aligned} & p(n) - p(n - \ell) - p(n - 2\ell) + p(n - 5\ell) + p(n - 7\ell) - \cdots \\ & \cdots + (-1)^j [p(n - \ell j(3j - 1)/2) + p(n - \ell j(3j + 1)/2)] + \cdots = b_\ell(n). \end{aligned}$$

*Proof.* We have

$$\begin{aligned} & \sum_{n=0}^{\infty} b_\ell(n) q^n \\ &= \prod_{k=1}^{\infty} \frac{(1 - q^{\ell k})}{(1 - q^k)} \\ &= \sum_{n=0}^{\infty} p(n) q^n \left( \sum_{j=-\infty}^{\infty} (-1)^j q^{\ell \frac{j(3j-1)}{2}} \right) \\ &= \sum_{n=0}^{\infty} p(n) q^n \left( 1 + \sum_{j=1}^{\infty} (-1)^j q^{\ell \frac{j(3j-1)}{2}} + \sum_{j=1}^{\infty} (-1)^j q^{\ell \frac{j(3j+1)}{2}} \right) \\ &= \sum_{n=0}^{\infty} \left( p(n) + \sum_{j=1}^{\infty} (-1)^j [p(n)(n - \ell j(3j - 1)/2) + p(n - \ell j(3j + 1)/2)] \right) q^n, \end{aligned}$$

from which the result follows.  $\square$

We close this section with a recurrence relation for the number of partitions of  $n$  having parts congruent to  $\pm c \pmod{m}$ .

**Theorem 8.** Given integers  $a$  and  $m \geq 1$ , we let  $p_m^c(n)$  denote the number of partitions of  $n$  having parts congruent to  $\pm c$  modulo  $m$ . Then, for all  $n \geq 0$ ,

$$\begin{aligned} & p_m^c(n) - p_m^c(n - (m - c)) - p_m^c(n - c) + p_m^c(n - (3m - 2c)) \\ & + p_m^c(n - (m + 2c)) + \cdots + (-1)^j p_m^c(n - (mj^2 + (m - 2c)j)/2) \\ & + (-1)^j p_m^c(n - (mj^2 - (m - 2c)j)/2) + \cdots = \begin{cases} 1, & \text{if } n = mk_j^e, \\ -1, & \text{if } n = mk_j^o, \\ 0, & \text{otherwise,} \end{cases} \end{aligned}$$

where  $k_j^e$  (resp.  $k_j^o$ ) is the  $j$ -th even (resp. odd) pentagonal number.

*Proof.* Setting  $c = -q^{m-c}$  and  $b = -q^c$  in (3) and (5), we obtain

$$\begin{aligned} & (q^{m-c}; q^m)_\infty (q^c; q^m)_\infty (q^m; q^m)_\infty = f(-q^{m-c}, -q^c) \\ & = \sum_{j=-\infty}^{\infty} (-q^{m-c})^{\frac{j(j+1)}{2}} (-q^c)^{\frac{j(j-1)}{2}}, \end{aligned}$$

which yields

$$\sum_{j=-\infty}^{\infty} (-1)^j q^{\frac{mj^2+(m-2c)j}{2}} = \prod_{k=1}^{\infty} (1 - q^{mk-c})(1 - q^{mk-(m-c)})(1 - q^{mk}). \quad (17)$$

The generating function for  $p_m^c$  is given by

$$\sum_{i=0}^{\infty} p_m^c(i) q^i = \prod_{k=1}^{\infty} \frac{1}{(1 - q^{mk-c})(1 - q^{mk-(m-c)})}.$$

Hence, we can rewrite (17) as

$$\prod_{k=1}^{\infty} \frac{1}{(1 - q^{mk-c})(1 - q^{mk-(m-c)})} \sum_{j=-\infty}^{\infty} (-1)^j q^{\frac{mj^2+(m-2c)j}{2}} = \prod_{k=1}^{\infty} (1 - q^{mk}),$$

or, equivalently,

$$\sum_{i=0}^{\infty} p_m^c(i) q^i \sum_{j=-\infty}^{\infty} (-1)^j q^{\frac{mj^2+(m-2c)j}{2}} = \sum_{j=-\infty}^{\infty} (-1)^j q^{\frac{mj(3j-1)}{2}}.$$

This last identity yields

$$\begin{aligned} & \sum_{i=0}^{\infty} p_m^c(i) q^i \left( 1 + \sum_{j=1}^{\infty} (-1)^j q^{\frac{mj^2+(m-2c)j}{2}} + \sum_{j=1}^{\infty} (-1)^j q^{\frac{mj^2-(m-2c)j}{2}} \right) \\ &= \sum_{n=0}^{\infty} \left( p_m^c(n) + \sum_{j=1}^{\infty} (-1)^j p_m^c(n - (mj^2 + (m-2c)j)/2) \right. \\ & \quad \left. + (-1)^j p_m^c(n - (mj^2 - (m-2c)j)/2) \right) q^n. \end{aligned}$$

Therefore,

$$\begin{aligned} \sum_{j=-\infty}^{\infty} (-1)^j q^{\frac{mj(3j-1)}{2}} &= \sum_{n=0}^{\infty} \left( p_m^c(n) + \sum_{j \geq 1} (-1)^j p_m^c(n - (mj^2 + (m-2c)j)/2) \right. \\ & \quad \left. + (-1)^j p_m^c(n - (mj^2 - (m-2c)j)/2) \right) q^n, \end{aligned}$$

which completes the proof.  $\square$

As special cases of the above theorem, we have the following corollaries which provide recurrence relations for the number of partitions that appear in the well-known Rogers-Ramanujan's identities.

**Corollary 9.** *Let  $p_{R1}(n)$  denote the number of partitions of  $n$  whose parts are congruent to  $\pm 1$  modulo 5 and let  $p_{R2}(n)$  denote the number of partitions of  $n$*

whose parts are congruent to  $\pm 2$  modulo 5. Then, for all  $n \geq 0$ ,

$$\begin{aligned} & p_{R1}(n) - p_{R1}(n-1) - p_{R1}(n-4) + p_{R1}(n-7) + p_{R1}(n-13) - \dots \\ & \dots + (-1)^j [p_{R1}(n - j(5j-3)/2) + p_{R1}(n - j(5j+3)/2)] + \dots \\ & \dots = \begin{cases} 1, & \text{if } n = 5k_j^e \\ -1, & \text{if } n = 5k_j^o \\ 0, & \text{otherwise,} \end{cases} \end{aligned}$$

and

$$\begin{aligned} & p_{R2}(n) - p_{R2}(n-1) - p_{R2}(n-2) + p_{R2}(n-5) + p_{R2}(n-7) - \dots \\ & \dots + (-1)^j [p_{R2}(n - j(3j-1)/2) + p_{R2}(n - j(3j+1)/2)] + \dots \\ & \dots = \begin{cases} 1, & \text{if } n = h_j^e \\ -1, & \text{if } n = h_j^o \\ 0, & \text{otherwise,} \end{cases} \end{aligned}$$

where  $h_j^e$  (resp.  $h_j^o$ ) is the  $j$ -th heptagonal number with  $j$  even (resp. odd).

**Corollary 10.** Let  $s_1(n)$  denote the number of partitions of  $n$  having congruent to  $\pm 1$  modulo 6 and let  $s_2(n)$  denote the number of partitions of  $n$  whose parts are congruent to  $\pm 2$  modulo 6. Then, for all  $n \geq 0$ ,

$$\begin{aligned} & s_1(n) - s_1(n-1) - s_1(n-5) + s_1(n-8) + s_1(n-16) - \dots \\ & \dots + (-1)^j [s_1(n - j(3j-2)) + s_1(n - j(3j+2))] + \dots \\ & = s_2(n) - s_2(n-2) - s_2(n-4) + s_2(n-10) + s_2(n-14) - \dots \\ & \dots + (-1)^j [s_2(n - j(3j-1)) + s_2(n - j(3j+1))] + \dots \\ & \dots = \begin{cases} 1, & \text{if } n = 6k_j^e \\ -1, & \text{if } n = 6k_j^o \\ 0, & \text{otherwise.} \end{cases} \end{aligned}$$

## 4 Acknowledgments

This research was supported by São Paulo Research Foundation (FAPESP) (grant no. 2019/10742-0).

## References

- [1] B. C. Berndt, *Number Theory in the Spirit of Ramanujan*, American Mathematical Society, 2006.
- [2] Y. Choliy, L. W. Kolitsch, and A. V. Sills, *Partition recurrences*, *Integers* **18** (2018), Article A1.

- [3] S. Corteel and J. Lovejoy, *Overpartitions*, Trans. Amer. Math. Soc. **356** (2004), 1623–1635.
- [4] J. A. Ewell, *Partition recurrences*, J. Combin. Theory Ser. A **14** (1973), 125–127.
- [5] J. A. Ewell, *Another recurrence for the partition function*, JP J. Algebra, Number Theory, App. **4** (2004), 147–152.
- [6] M. Merca, *New recurrences for Euler’s partition function*, Turk. J. Math. **41** (2017), 1184–1190.
- [7] D. Nyirenda, *On parity and recurrences for certain partition functions*, Contrib. Discrete Math. **15** (2020), 72–79.
- [8] K. Ono, N. Robbins, and B. Wilson, *Some recurrences for arithmetical functions*, J. Indian Math. Soc. **62** (1996), pages 29–50.