CONSTRUCTING SPAN CATEGORIES FROM CATEGORIES WITHOUT PULLBACKS

DAVID WEISBART¹ AND ADAM M. YASSINE²

¹Department of Mathematics University of California, Riverside ²Department of Mathematics Bowdoin College

In memory of Professor V.S.Varadarajan.

ABSTRACT. Span categories provide an abstract framework for formalizing mathematical models of certain systems. The mathematical descriptions of some systems, such as classical mechanical systems, require categories that do not have pullbacks, and this limits the utility of span categories as a formal framework. Given categories \mathscr{C} and \mathscr{C}' and a functor \mathcal{F} from \mathscr{C} to \mathscr{C}' , we introduce the notion of an \mathcal{F} -pullback of a cospan in \mathscr{C} , as well as the notion of span tightness of \mathcal{F} . If \mathcal{F} is span tight, then we can form a generalized span category $\text{Span}(\mathscr{C}, \mathcal{F})$ and circumvent the technical difficulty of \mathscr{C} failing to have pullbacks. Composition in $\text{Span}(\mathscr{C}, \mathcal{F})$ uses \mathcal{F} -pullbacks rather than pullbacks and in this way differs from the category $\text{Span}(\mathscr{C})$, but reduces to it when both \mathscr{C} has pullbacks and \mathcal{F} is the identity functor.

Contents

2. Span Categories3.3. Some Categories that do not have Pullbacks6.4. Composition by F-Pullbacks14.5. Generalized Span Category18.6. Examples20.References20.	1.	Introduction	1
3. Some Categories that do not have Pullbacks64. Composition by F-Pullbacks145. Generalized Span Category156. Examples20References20	2.	Span Categories	3
4. Composition by F-Pullbacks145. Generalized Span Category156. Examples20References20	3.	Some Categories that do not have Pullbacks	6
5. Generalized Span Category186. Examples20References20	4.	Composition by \mathcal{F} -Pullbacks	14
6. Examples20References20	5.	Generalized Span Category	15
References 20	6.	Examples	20
	References		20

1. INTRODUCTION

The last few decades have seen significant application of category theory to modeling systems. The basic idea is to model a system by a morphism in an appropriate category, where composition of morphisms describes the assembly of systems into larger or more complicated systems. Earlier works have used span and cospan categories to study the composition of

E-mail address: ¹weisbart@math.ucr.edu, ²a.yassine@bowdoin.edu.

physical systems. For example, Baez and Pollard [3] used cospans to study reaction networks. Haugseng [8] used spans to study classical topological field theories. A span in a category \mathscr{C} is a pair of morphisms in \mathscr{C} with the same source. Bénabou [5, p. 22] credits Yoneda [13] with introducing the notion of a span in the category of categories. Bénabou [5] proved that if \mathscr{C} is a category with pullbacks, then there is a bicategory, Span(\mathscr{C}), whose objects, morphisms, and 2-morphisms are the respective objects, spans, and maps of spans in \mathscr{C} . View this bicategory as a category, a *span category*, by ignoring the bicategory structure and taking isomorphism classes of spans in \mathscr{C} , that Section 2 defines, as the morphisms. Leinster [10, p. 26] discussed bicategories and Bénabou's work.

Denote by Set the category whose objects are sets, whose morphisms are functions between sets, and where composition of functions defines the composition of morphisms. Suppose that X, Y, and Z are sets and that f and g are functions that respectively map X and Y to the set Z. Take ρ_X and ρ_Y to be the canonical projections

$$\rho_X \colon X \times Y \to X$$
 and $\rho_Y \colon X \times Y \to Y$,

and take π_X and π_Y to be the respective restrictions of ρ_X and ρ_Y to the fibered product $X \times_Z Y$, the subset of $X \times Y$ consisting of all elements on which f is equal to g. Section 4 discusses the universal properties of the fibered product in **Set**. These properties give a prescription for composing certain spans in **Set** in a way that seems strikingly similar to the way in which classical mechanical systems appear to compose.

Baez initiated the current line of research by proposing that the study of classical mechanics might have a foundation in category theory. In particular, he suggested that classical mechanical systems could be morphisms in an appropriate span category, where composition of morphisms using fibered products would describe the composition of the systems [2]. However, modeling classical mechanical systems necessitates working with spans in categories other than **Set**, where the fibered product lacks the universal properties that it has in **Set**. The current paper provides a way to circumvent this technical obstruction by generalizing the construction of a span category. Generalized span categories differ from span categories in the way that they define composition. The composite of spans may not be defined in a given category \mathscr{C} because composition requires the existence of pullbacks and \mathscr{C} may not have pullbacks. The rough idea is to map \mathscr{C} by a functor \mathcal{F} into another category \mathscr{C}' . The composite of two spans in \mathscr{C} will be a span in \mathscr{C} with the property that its image under \mathcal{F} is a composite in \mathscr{C}' of the the images under \mathcal{F} of the spans being composed in \mathscr{C} . If the functor \mathcal{F} satisfies a certain property, namely that it is span tight, then composition will be well defined for isomorphism classes of spans in \mathscr{C} .

Section 2 defines a span, a cospan, and an isomorphism class of spans. The language it employs is arguably nonstandard from a category theorist's perspective but we have found it helpful both for presenting the results to nonspecialists in category theory and for our specific applications. Section 3 proves that some categories that are important in the study of classical mechanics do not have pullbacks, which leads to the present goal of generalizing the definition of a span category. Denote by Diff the category whose objects are smooth manifolds and whose morphisms are smooth functions. Since two submanifolds of a given manifold may not intersect transversally, the fibered product of two manifolds is not necessarily an embedded submanifold of their product. For reasons related to this, Diff does not have pullbacks. This technical difficulty that Spivak [12] encountered parallels a central technical difficulty of the current paper.

Suppose that \mathscr{C} and \mathscr{C}' are categories and \mathcal{F} is a functor from \mathscr{C} to \mathscr{C}' . Section 4 defines an \mathcal{F} -pullback of a cospan in \mathscr{C} and the span tightness of the functor \mathcal{F} , as well as the composite of two spans along an \mathcal{F} -pullback. While the notion of an \mathcal{F} -pullback generalizes the notion of a pullback in a way that is sufficient for the current setting, without an additional condition on \mathcal{F} it is not enough to provide a method for composing isomorphism classes of spans. Section 5 proves that if the functor \mathcal{F} is span tight, then there exists a category $\text{Span}(\mathscr{C}, \mathcal{F})$ whose objects are the objects in \mathscr{C} and whose morphisms are isomorphism classes of spans in \mathscr{C} . Composition in this generalized span category is defined using \mathcal{F} -pullbacks and appears to depend on the functor \mathcal{F} . In [4], we apply the tools that we develop to the study of classical mechanics. Section 6 briefly discusses some examples of generalized span categories.

Acknowledgements. We thank Professor John C. Baez for initially suggesting this project to us, for his valuable guidance, and for his enlightening instruction. We thank him as well for helping us to understand his motivation for this project and the orientation of this current project in his larger research program. We thank Professors Daniel Cicala and Christina Vasilakopoulou for many insightful discussions on category theory.

We dedicate this work to Professor V.S. Varadarajan, a dear mentor and friend, whose memory we will always treasure.

2. Span Categories

2.1. Spans and their Isomorphism Classes. A span in a category \mathscr{C} is a pair of morphisms in \mathscr{C} with the same source and a *cospan* in \mathscr{C} is a pair of morphisms in \mathscr{C} with the same target. For any span S in \mathscr{C} , write

$$S = (s_L, s_R),$$

where S_L , S_R , and S_A are objects in \mathscr{C} ,

$$s_L \colon S_A \to S_L$$
, and $s_R \colon S_A \to S_R$.

Utilize the same notation if S is a cospan, but where s_L and s_R map S_L and S_R to S_A , respectively. The span S and the cospan C are these respective diagrams:



For any span or cospan S in \mathscr{C} , refer to the objects S_A , S_L , and S_R in \mathscr{C} as the *apex*, *left foot*, and *right foot* of S, respectively.

Definition 2.1. A span S in \mathscr{C} is *paired with* a cospan C in \mathscr{C} if

$$C_L = S_L, \quad C_R = S_R, \quad \text{and} \quad c_L \circ s_L = c_R \circ s_R.$$

Definition 2.2. Take S and Q to be spans in \mathscr{C} with S_L equal to Q_L and S_R equal to Q_R . A span morphism in \mathscr{C} from S to Q is a morphism Φ in \mathscr{C} from S_A to Q_A with

$$s_L = q_L \circ \Phi$$
 and $s_R = q_R \circ \Phi$.

A span isomorphism in \mathscr{C} from S to Q is a span morphism that is additionally an isomorphism.

View a span S paired with a cospan C and view a span morphism Φ from a span S to a span Q, respectively, as these commutative diagrams:



Proposition 2.3. For any span isomorphism Φ , the inverse Φ^{-1} is also a span isomorphism. Furthermore, a composite of span morphisms is a span morphism.

2.2. Pullbacks in a Category \mathscr{C} . This subsection introduces the notion of a pullback of a cospan and defines what it means for a category to have pullbacks.

Definition 2.4. A span S in \mathscr{C} is a *pullback of a cospan* C *in* \mathscr{C} if it is paired with C and if for any other span Q in \mathscr{C} that is also paired with C, there exists a unique span morphism Φ in \mathscr{C} from Q to S.

The morphism Φ in Definition 2.4 is the unique span morphism so that this diagram commutes:



Definition 2.5. A category \mathscr{C} has pullbacks if for any cospan C in \mathscr{C} , there is a span S in \mathscr{C} that is a pullback of C and S is unique up to a span isomorphism in \mathscr{C} .

The diagram consisting of a pullback S of a cospan C together with C is a *pullback square*. We have found it useful to separately define the parts of a pullback square.

2.3. Examples of Categories that have Pullbacks. Denote by Top the category whose objects are topological spaces and whose morphisms are continuous functions. The categories Set and Top are examples of categories that have pullbacks, as MacLane [11] discusses and Awodey [1] discusses more specifically for Set. We provide a proof here for the convenience of the reader.

Let C be a cospan in Set, and let ρ_L and ρ_R be the canonical projections

 $\rho_L \colon C_L \times C_R \to C_L$ and $\rho_R \colon C_L \times C_R \to C_R$.

Denote by S_A the fibered product

$$C_L \times_{C_A} C_R := \{ (x, y) \in C_L \times C_R : (c_L \circ \rho_L)(x, y) = (c_R \circ \rho_R)(x, y) \}.$$

Take S_L and S_R to be equal to C_L and C_R , respectively, and let s_L and s_R be the respective restrictions of ρ_L and ρ_R to the set S_A . For any span P that is paired with C, take Φ to be the function

$$\Phi: P_A \to C_L \times C_R$$
 by $a \mapsto (p_L(a), p_R(a))$ $(\forall a \in P_A).$

The function Φ is the unique function from P_A to $C_L \times C_R$ such that

(1)
$$p_L = \rho_L \circ \Phi \quad \text{and} \quad p_R = \rho_R \circ \Phi.$$

The image of Φ is contained in S_A and so Φ is a span morphism from P to S. Since any other span morphism from P to S defines a function from P to $C_L \times C_R$ with the property given by (1), the function Φ is the unique span morphism from P to S, and so the span S is a pullback of the cospan C.

Suppose that C is a cospan in Top, and let ρ_L and ρ_R again be the canonical projections on $C_L \times C_R$. The product $C_L \times C_R$ with the product topology is a topological space. The fibered product S_A given above is a subset of $C_L \times C_R$ and is a topological space with the subspace topology. The projections s_L and s_R are continuous maps, and so (s_L, s_R) is a pullback of C. The proof of this fact is nearly the same as the proof in the setting of Set, with only an additional straightforward check that the mappings involved are continuous.

2.4. The Category Span(\mathscr{C}). Suppose that \mathscr{C} is a category that has pullbacks. Suppose that [S] and [Q] are isomorphism classes of spans with respective representatives S and Q, and S_R is equal to Q_L . Since \mathscr{C} has pullbacks, there is a span P that is a pullback of the cospan (s_R, q_L) . Define by $[(s_L \circ p_L, q_R \circ p_R)]$ the composite $[S] \circ [Q]$. Take the objects in \mathscr{C} to be the objects in Span(\mathscr{C}), the isomorphism classes of spans in \mathscr{C} to be the morphisms in Span(\mathscr{C}), and S_R and S_L to respectively be the source and target of the span [S]. Given an object X in \mathscr{C} and the identity morphism Id taking X to X, define by $[(\mathrm{Id}, \mathrm{Id})]$ the identity morphism in Span(\mathscr{C}) with X as both source and target. It is well known that Span(\mathscr{C}) is a category [5]. Our treatment in Section 4 of generalized span categories specializes in the case when \mathscr{C} has pullbacks to give a proof that Span(\mathscr{C}) is a category. If \mathscr{C} does not have pullbacks, then the existence of P is not guaranteed. The next section will demonstrate that some categories that are important in classical mechanics, and more generally in differential geometry, do not have pullbacks.

3. Some Categories that do not have Pullbacks

3.1. **Background.** Refer to [9] as a standard reference for smooth manifold theory. The current section reviews some well-known definitions in order to explicitly establish language and notational conventions.

An *m*-dimensional manifold is a triple $(M, \mathcal{T}_M, \mathcal{A}_M)$ such that:

- (1) M is a set;
- (2) \mathcal{T}_M is a topology for M that is Hausdorff and second countable;
- (3) \mathcal{A}_M is an atlas, a collection of homeomorphisms such that the domain of each element of \mathcal{A}_M is an open subset of M, the collection of domains of the elements of \mathcal{A}_M form an open cover for M, and the range of each element of \mathcal{A}_M is an open subset of \mathbb{R}^m .

If \mathcal{A}_M is maximal with respect to the property that for any ϕ and ψ in \mathcal{A}_M that have intersecting domains, the transition function $\phi \circ \psi^{-1}$ and its inverse are of class C^r (r-times continuously differentiable), then M is a C^r -manifold. Only the smooth case, when r is infinity, is relevant to this paper. Refer to the elements of \mathcal{A}_M as coordinates and refer to their domains as charts. It is customary to denote by M a manifold $(M, \mathcal{T}_M, \mathcal{A}_M)$, and we generally follow this convention, except when it is important to explicitly distinguish between the manifold, the topological space associated to the manifold, and the underlying set. Reference to the manifold M, the topological space M, and the underlying set M, will respectively be a reference to the triple $(M, \mathcal{T}_M, \mathcal{A}_M)$, the pair (M, \mathcal{T}_M) , and the set M. Denote by Diff the category whose objects are smooth manifolds and whose morphisms are smooth maps between smooth manifolds.

3.2. Some Functors that preserve Pullbacks. Suppose that \mathscr{C} is a locally small category and that X is an object in \mathscr{C} . Denote by $\operatorname{Hom}(X, -)$ the hom functor from \mathscr{C} to Set. A functor \mathcal{F} that takes \mathscr{C} to Set is *representable* if there is an object B in \mathscr{C} so that \mathcal{F} is naturally isomorphic to $\operatorname{Hom}(B, -)$.

The categories Diff, Top, and Set are locally small and there are forgetful functors, each to be ambiguously denoted by \mathcal{F} , from Diff to Top and from Top to Set given by

$$\mathcal{F}(M, \mathcal{T}_M, \mathcal{A}_M) = (M, \mathcal{T}_M) \text{ and } \mathcal{F}(M, \mathcal{T}_M) = M$$

The morphisms in Diff and Top are entirely determined by their action on the underlying sets and so the forgetful functor in each case maps a given source category to a subcategory of the target category. The functor obtained by composing the above forgetful functors is the forgetful functor, denoted again by \mathcal{F} , from Diff to Set.

A functor \mathcal{F} from a category \mathscr{C} to a category \mathscr{C}' preserves pullbacks if for any cospan Cin \mathscr{C} , if S is a pullback of C, then $\mathcal{F}(S)$ is a pullback of $\mathcal{F}(C)$. The following lemma is a special case of a more general result that guarantees that representable functors preserve pullbacks [6, p. 64]. The proof of Lemma 3.1 is included for the convenience of the reader, because the language in our definition of a pullback is slightly different from Borceux's [6]. **Lemma 3.1.** For any locally small category \mathscr{C} and any object B in \mathscr{C} , the functor $\operatorname{Hom}(B, -)$ preserves pullbacks, where

$$\operatorname{Hom}(B,-)\colon \mathscr{C} \to \mathsf{Set}.$$

Proof. Take X and Y to be objects in \mathscr{C} . For any morphism f in \mathscr{C} from X to Y, denote by \tilde{f} the morphism $\operatorname{Hom}(B, f)$ that is defined to act on any β in $\operatorname{Hom}(B, X)$ by

$$\tilde{f}(\beta) = f \circ \beta.$$

Suppose that C is a cospan in \mathscr{C} and that S is a pullback of C. Since \mathscr{C} is locally small, the functor $\operatorname{Hom}(B, -)$ maps the cospan C to a cospan $\operatorname{Hom}(B, C)$ in Set, taking the pair (c_L, c_R) to the pair $(\tilde{c}_L, \tilde{c}_R)$. It similarly maps the span S to the span $\operatorname{Hom}(B, S)$. Since S is a pullback of C, it is paired with C and so

$$c_L \circ s_L = c_R \circ s_R.$$

Therefore, for any ψ in Hom (B, S_A) ,

$$(\tilde{c}_L \circ \tilde{s}_L)(\psi) = c_L \circ s_L \circ \psi = c_R \circ s_R \circ \psi = (\tilde{c}_R \circ \tilde{s}_R)(\psi),$$

and so the span $\operatorname{Hom}(B, S)$ is paired with the cospan $\operatorname{Hom}(B, C)$.

Denote respectively by ρ_L and ρ_R the canonical projections from $\text{Hom}(B, C_L) \times \text{Hom}(B, C_R)$ to $\text{Hom}(B, C_L)$ and $\text{Hom}(B, C_R)$, and by Q_A the set

$$\operatorname{Hom}(B, C_L) \times_{\operatorname{Hom}(B, C_A)} \operatorname{Hom}(B, C_R) \\= \left\{ \alpha \in \operatorname{Hom}(B, C_L) \times \operatorname{Hom}(B, C_R) \colon (\tilde{c}_L \circ \rho_L)(\alpha) = (\tilde{c}_R \circ \rho_R)(\alpha) \right\}.$$

Take q_L and q_R to be the respective restrictions of ρ_L and ρ_R to Q_A . Denote by Q the span (q_L, q_R) in Set, a pullback of the cospan Hom(B, C).

For any α in Q_A , there are morphisms α_L and α_R in \mathscr{C} that respectively map B to C_L and C_R , and α is equal to (α_L, α_R) . Furthermore,

$$c_L \circ \alpha_L = \tilde{c}_L(\alpha_L) = (\tilde{c}_L \circ q_L)(\alpha) = (\tilde{c}_R \circ q_R)(\alpha) = \tilde{c}_R(\alpha_R) = c_R \circ \alpha_R.$$

The pair (α_L, α_R) is, therefore, a span in \mathscr{C} that is paired with C and, since S is a pullback of C, there is a unique span morphism ϕ_{α} in \mathscr{C} from (α_L, α_R) to S that maps B to S_A . Take Φ to be the function from Q to $\operatorname{Hom}(B, S)$ that is defined for each α in Q_A by

$$\Phi(\alpha) = \phi_{\alpha}$$

The morphism ϕ_{α} is a span morphism, implying that

$$s_L \circ \phi_{\alpha} = \alpha_L$$
 and $s_R \circ \phi_{\alpha} = \alpha_R$.

These equalities further imply that

$$(\tilde{s}_L \circ \Phi)(\alpha) = s_L \circ \phi_\alpha, \quad \alpha_L = q_L(\alpha), \quad (\tilde{s}_R \circ \Phi)(\alpha) = s_R \circ \phi_\alpha, \quad \text{and} \quad \alpha_R = q_R(\alpha),$$

and so

$$(\tilde{s}_L \circ \Phi)(\alpha) = q_L(\alpha)$$
 and $(\tilde{s}_R \circ \Phi)(\alpha) = q_R(\alpha)$.

The morphism Φ in Set is, therefore, a span morphism and is unique since ϕ_{α} is uniquely determined. Since Q is a pullback of Hom(B, C), the span Hom(B, S) is as well, and so Hom(B, -) maps pullbacks in \mathscr{C} to pullbacks in Set.

Take $(\mathbf{1}, \mathcal{T}_{\mathbf{1}}, \mathcal{A}_{\mathbf{1}})$ to be the singleton manifold with its unique discrete topology. Denote by $\mathsf{Diff}((\mathbf{1}, \mathcal{T}_{\mathbf{1}}, \mathcal{A}_{\mathbf{1}}), -)$ the functor that maps any given manifold $(X, \mathcal{T}_X, \mathcal{A}_X)$ to the set of functions $\mathsf{Diff}((\mathbf{1}, \mathcal{T}_{\mathbf{1}}, \mathcal{A}_{\mathbf{1}}), (X, \mathcal{T}_X, \mathcal{A}_X))$. Compress notation by writing $\mathsf{Diff}(\mathbf{1}, X)$ instead of $\mathsf{Diff}((\mathbf{1}, \mathcal{T}_{\mathbf{1}}, \mathcal{A}_{\mathbf{1}}), (X, \mathcal{T}_X, \mathcal{A}_X))$ and $\mathsf{Diff}(\mathbf{1}, -)$ instead of $\mathsf{Diff}((\mathbf{1}, \mathcal{T}_{\mathbf{1}}, \mathcal{A}_{\mathbf{1}}), -)$. Lemma 3.2 is well known, but we include it for the reader's convenience.

Lemma 3.2. The forgetful functor \mathcal{F} from Diff to Set is naturally isomorphic to the functor Diff(1, -).

Proof. For any object $(X, \mathcal{T}_X, \mathcal{A}_X)$ in Diff, take η_X to be the function from Diff $(\mathbf{1}, X)$ to X that is given for any h in Diff $(\mathbf{1}, X)$ by

$$\eta_X(h) = h(\mathbf{1}).$$

The function η_X is invertible, with inverse η_X^{-1} that is given for any x in X by

$$\eta_X^{-1}(x) = h_x \colon \mathbf{1} \mapsto x,$$

hence η_X is an isomorphism.

Suppose that g is in Diff(1, X) and f is a smooth function from $(X, \mathcal{T}_X, \mathcal{A}_X)$ to $(Y, \mathcal{T}_Y, \mathcal{A}_Y)$. The point $\eta_X(g)$ in X is equal to g(1) and $\mathcal{F}(f)(g(1))$ is equal to f(g(1)). Furthermore, Diff(1, f) takes g to $f \circ g$ and η_Y takes $(f \circ g)$ to $(f \circ g)(1)$, which is equal to f(g(1)), and so this diagram is commutative:

$$\begin{array}{c|c} \mathsf{Diff}(\mathbf{1}, X) & \xrightarrow{\eta_X} & \mathcal{F}(X, \mathcal{T}_X, \mathcal{A}_X) \\ \mathsf{Diff}(\mathbf{1}, f) & & & & \downarrow \mathcal{F}(f) \\ & & & & \downarrow \mathcal{F}(f) \\ \mathsf{Diff}(\mathbf{1}, Y) & \xrightarrow{\eta_Y} & \mathcal{F}(Y, \mathcal{T}_Y, \mathcal{A}_Y) \end{array}$$

The functor $\mathsf{Diff}(1, -)$ is, therefore, naturally isomorphic to \mathcal{F} .

Lemma 3.1 and Lemma 3.2 together imply Proposition 3.3.

Proposition 3.3. The forgetful functor \mathcal{F} from Diff to Set preserves pullbacks.

3.3. SurjSub does not have Pullbacks. Denote by SurjSub the category whose objects are smooth manifolds and whose morphisms are smooth surjective submersions. This category is important in the study of classical mechanical systems because a map that takes the configuration space of a classical mechanical system to the configuration space of a subsystem should be a surjective submersion. The category SurjSub is an example of a category that does not have pullbacks. To see this, denote the one and two point manifolds respectively

by 1 and 2. Take f to be the unique map from 2 to 1. Denote by Id the identity map from 2 to 2. The span (Id, Id) is paired with the cospan (f, f).

Suppose that π_L and π_R are the canonical projections from $\mathbf{2} \times_1 \mathbf{2}$ to $\mathbf{2}$. Suppose that S is a pullback of (f, f) in SurjSub. Proposition 3.3 implies that the image of S under the forgetful functor from Diff to Set is the span (π_L, π_R) and that there is a unique Φ in SurjSub so that this diagram commutes:



However, $\mathbf{2} \times_{\mathbf{1}} \mathbf{2}$ is isomorphic to $\mathbf{2} \times \mathbf{2}$, a set with four elements, and so Φ is a surjection from a set with only two elements to a set with four elements, which is not possible. The cospan (f, f) does not have a pullback in SurjSub, and so SurjSub does not have pullbacks.

The technical obstructions that arise in working with categories of smooth manifolds with surjective morphisms are elementary, as the above example indicates. To avoid any misunderstanding, notice that examples of cospans that do not have pullbacks may be constructed in a similar way that involve connected manifolds with nontrivial smooth structures. For example, take f to be a projection from \mathbb{R}^4 to \mathbb{R}^2 . Take π_L and π_R to be the restrictions to the subset $\mathbb{R}^4 \times_{\mathbb{R}^2} \mathbb{R}^4$ of the respective projections ρ_L and ρ_R that are defined for each xand y in \mathbb{R}^4 by

$$\rho_L(x,y) = x$$
 and $\rho_R(x,y) = y$.

Take Id to be the identity map from \mathbb{R}^4 to \mathbb{R}^4 . The span (Id, Id) is paired with (f, f), but there is no smooth surjection from \mathbb{R}^4 to the fibered product since the fibered product is six-dimensional. This counterexample is relevant to demonstrating that the category of Riemannian manifolds with surjective Riemannian submersions and the category of symplectic manifolds with surjective Poisson maps also do not have pullbacks. These categories play a fundamental role in [4].

3.4. Diff does not have Pullbacks. Suppose throughout this subsection that f and g are morphisms in Diff that have mutual target $(Z, \mathcal{T}_Z, \mathcal{A}_Z)$ and respective sources $(X, \mathcal{T}_X, \mathcal{A}_X)$ and $(Y, \mathcal{T}_Y, \mathcal{A}_Y)$. Recall that π_X and π_Y are the respective projections from the set $X \times_Z Y$ to X and Y. Take \mathcal{T}_S to be the subspace topology on $X \times_Z Y$ that $X \times_Z Y$ inherits from the product topology on $X \times Y$ and with respect to which π_X and π_Y are both continuous. View the functions f and g as functions in Top that have the topological space (Z, \mathcal{T}_Z) as their mutual target and the topological spaces (X, \mathcal{T}_X) and (Y, \mathcal{T}_Y) as their respective sources. Suppose that $(W, \mathcal{T}_W, \mathcal{A}_W)$ is an embedded submanifold of $(Z, \mathcal{T}_Z, \mathcal{A}_Z)$. Refer to [9, p. 143-144] for further discussion of transversality and, in particular, for the proof of Proposition 3.5.

Definition 3.4. The smooth function f is transverse to W if, for every x in $f^{-1}(W)$, the spaces $T_{f(x)}W$ and $df(T_xX)$ together span $T_{f(x)}Z$. The smooth functions f and g are transverse if, for every point x in X and y in Y with f(x) and g(y) both equal to z,

$$\mathrm{d}f(T_xX) + \mathrm{d}g(T_yY) = T_zZ.$$

Proposition 3.5. Suppose that X and Z are smooth manifolds and W is an embedded submanifold of Z. If f is a smooth map from X to Z that is transverse to W, then $f^{-1}(W)$ is an embedded submanifold of X whose codimension is equal to the codimension of W in Z.

Proposition 3.6. If f and g are transverse, then the fibered product $X \times_Z Y$ is a smooth embedded submanifold of codimension equal to the dimension of Z. Furthermore, the span (π_X, π_Y) in Diff is a pullback of (f, g).

Proof. Denote by Δ_Z the diagonal $\{(z, z) : z \in Z\}$ of $Z \times Z$, an embedded submanifold of $Z \times Z$. The function $f \times g$, with

$$f \times g \colon X \times Y \to Z \times Z$$
 by $(x, y) \mapsto (f(x), g(y)),$

is smooth and $(f \times g)^{-1}(\Delta_Z)$ is equal to $X \times_Z Y$. Since f and g are transverse, the function $f \times g$ is transverse to Δ_Z . Proposition 3.5 implies that $X \times_Z Y$ is a smooth manifold with codimension in $X \times Y$ equal to the dimension of Δ_Z . The dimension of Δ_Z is equal to that of Z, and so the codimension of $X \times_Z Y$ in $X \times Y$ is equal to the dimension of Z.

To show that (π_X, π_Y) is a pullback of (f, g), suppose that S is a span in Diff that is paired with (f, g). Define for each s in S_A the span morphism Φ from S to (π_X, π_Y) by

$$\Phi(s) = (s_L(s), s_R(s)).$$

Take Φ' to be another span morphism from S to (π_X, π_Y) . For any s in S_A ,

 $\pi_X(\Phi'(s)) = s_L(s)$ and $\pi_Y(\Phi'(s)) = s_R(s),$

and so $\Phi'(s)$ is equal to $\Phi(s)$. Therefore, the morphism Φ' is equal to Φ . Since Φ is unique, (π_X, π_Y) is a pullback.

If f and g are in SurjSub with mutual target Z, then they are transverse and so Proposition 3.6 implies the following:

Proposition 3.7. If (f,g) is a cospan in SurjSub, then the fibered product $X \times_Z Y$ is a smooth embedded submanifold of $X \times Y$ of dimension dim $(X \times_Z Y)$, where

$$\dim(X \times_Z Y) = \dim(X) + \dim(Y) - \dim(Z).$$

For the following proposition, take (f, g) to be a cospan in Diff but where the maps f and g are not assumed to be transverse:

Proposition 3.8. If S is a span in Diff that is a pullback of (f, g), and if (π_X, π_Y) and (s_L, s_R) are span isomorphic as spans in Top, then $X \times_Z Y$ has a manifold structure.

Proof. Take Φ to be the unique span isomorphism in **Top** from S to (π_X, π_Y) . The homeomorphism Φ transports the manifold structure of S_A to $X \times_Z Y$, giving it a manifold structure as well.

If S is a span in Diff that is paired with (f, g), then the map Φ , that is defined for each s in S_A by

$$\Phi(s) = (s_L(s), s_R(s)),$$

is a smooth map from S_A to $X \times Y$. If $X \times_Z Y$ is an embedded submanifold of $X \times Y$, then Φ is a smooth map from S_A to $X \times_Z Y$ and is the unique such map, implying the following proposition:

Proposition 3.9. If $X \times_Z Y$ is an embedded submanifold of $X \times Y$, then (π_X, π_Y) is a span in Diff and a pullback of (f, g).

Propositions 3.6 and 3.9 together imply the following proposition:

Proposition 3.10. If (f,g) is a cospan in Diff and f and g are transverse, then (π_X, π_Y) is a pullback of (f,g) in Diff.

The following example demonstrates that $X \times_Z Y$ may be a manifold and (π_X, π_Y) may a pullback of a cospan (f, g) in Diff, but $X \times_Z Y$ is not an embedded submanifold of $X \times Y$. In light of Proposition 3.6, such an example requires the functions f and g to be non-transverse. It also shows that the forgetful functor from Diff to Top does not preserve pullbacks.

Example 3.11. Take X and Z to be equal to \mathbb{R} and take Y to be the one point manifold $\{y_0\}$. Suppose that f is a smooth function from X to Z, (a_n) is a sequence in \mathbb{R} that converges to a point a_0 that is not equal to a_n for any natural number n, and the zero set of f is the set $\{a_0\} \cup \{a_n : n \in \mathbb{N}\}$. Take g to be the zero function from Y to Z. The set $X \times_Z Y$ is the subset $\{(a_0, y_0)\} \cup \{(a_n, y_0) : n \in \mathbb{N}\}$ of $X \times Y$.

The span (π_X, π_Y) is paired with the cospan (f, g) in Top, where π_X and π_Y are the restrictions to $X \times_Z Y$ of the continuous projections ρ_X and ρ_Y that respectively take $X \times Y$ to X and Y, where $X \times Y$ is endowed with the product topology. The span (π_X, π_Y) is a pullback of (f, g), where $X \times_Z Y$ is endowed with the subspace topology \mathcal{T}_S . In this topology, each set $\{(a_n, y_0)\}$ is an open set, where n varies over \mathbb{N} . Any open set containing (a_0, y_0) contains infinitely many points.

Now view (f,g) as a cospan in Diff, so that $X \times_Z Y$ is a manifold. Each point in $X \times_Z Y$ must be contained in a neighborhood that is homeomorphic to a point, and so $X \times_Z Y$ must be endowed with the discrete topology \mathcal{T}_D . In this case, the manifold $X \times_Z Y$ is not an embedded submanifold of $X \times Y$ since its topology is not the subspace topology. Nevertheless, the span (π_X, π_Y) is a pullback of (f, g) in Diff.

The topology on the image of the manifold $X \times_Z Y$ under the forgetful functor from Diff to Top is \mathcal{T}_D , which is a finer topology than \mathcal{T}_S . The forgetful functor \mathcal{F} from Diff to Top maps the pullback (π_X, π_Y) , where maps π_X and π_Y have the manifold $(X \times_Z Y, \mathcal{T}_D, \mathcal{A}_{X \times_Z Y})$ as their common source, to the span (π_X, π_Y) , where the maps have the topological space $(X \times_Z Y, \mathcal{T}_D)$ as their common source. The identity map taking $(X \times_Z Y, \mathcal{T}_D)$ to $(X \times_Z Y, \mathcal{T}_S)$ is a continuous span morphism from (π_X, π_Y) to (π_X, π_Y) , but the inverse is not continuous. This demonstrates that the forgetful functor from Diff to Top does not preserve pullbacks.

The maps f and g in Example 3.11 are not transverse because f is smooth and constant on a sequence that is convergent to a_0 , and so $df(T_{a_0}\mathbb{R})$ is zero-dimensional. This example demonstrates that f and g may be non-transverse morphisms in Diff, but (f, g) nevertheless has a pullback in Diff. The forgetful functor \mathcal{F} from Diff to Set preserves pullbacks, and so if Q is a span in Diff and a pullback of (f, g), then $\mathcal{F}(Q)$ is a span in Set that is a pullback of $(\mathcal{F}(f), \mathcal{F}(g))$. Since Set has pullbacks, there is a span isomorphism in Set from $\mathcal{F}(Q)$ to $(\mathcal{F}(\pi_X), \mathcal{F}(\pi_Y))$. This span isomorphism is only a bijection, and there should be no expectation that it preserves topological structure.

The category Top also has pullbacks, and so if (f, g) is a cospan in Top, then a pullback of (f, g) will exist and, in fact, the span (π_X, π_Y) in Top is a pullback of (f, g) where the maps π_X and π_Y have $(X \times_Z Y, \mathcal{T}_S)$ as their common source. Since the forgetful functor from Diff to Top does not preserve pullbacks, there is no guarantee that Q being a pullback of (f, g) implies that it is a pullback when mapped by a forgetful functor to Top, as is the case in Example 3.11.

The former discussion, and the example that precedes it, demonstrate that there is some subtlety involved in determining that Diff does not have pullbacks. The proof of Proposition 3.13 presents an elementary counterexample. The functions given in the statement of Lemma 3.12 form this commutative diagram:



Lemma 3.12. For any cospan (f,g) in Diff and for any pullback Q of (f,g) in Diff, there is a bijective span morphism Φ in Top from $\mathcal{F}(Q)$ to (π_X, π_Y) , where \mathcal{F} is the forgetful functor from Diff to Top, and $X \times_Z Y$ is endowed with the subspace topology.

To clarify, Lemma 3.12 guarantees the existence of a bijective span morphism Φ in Top, but Φ is not guaranteed to be an isomorphism in Top since its inverse need not be continuous.

Proof. Take Q to be a span in Diff that is a pullback of (f, g). Define for each a in Q_A the function Φ by

(2)
$$\Phi(a) = (q_L(a), q_R(a)).$$

The map Φ from Q_A to $X \times Y$ is smooth because the functions q_L and q_R are smooth. The span Q is paired with (f, g), implying that the range of Φ is $X \times_Z Y$, and so Φ is a continuous function from Q_A to $X \times_Z Y$, where $X \times_Z Y$ is endowed with the subspace topology. Proposition 3.3 implies that the forgetful functor \mathcal{G} from Diff to Set preserves pullbacks, therefore $\mathcal{G}(\Phi)$ is a span morphism in Set from $\mathcal{G}(Q)$ to (π_X, π_Y) , where the pair of projections is viewed only as a pair of maps in Set. The span Q is a pullback in Diff, hence $\mathcal{G}(Q)$ is a span in Set that is a pullback of (f, g), and so the map $\mathcal{G}(\Phi)$ is a bijection. Maps between manifolds and topological spaces are determined by their behavior on the underlying sets, hence $\mathcal{F}(\Phi)$ is a continuous bijection.

Although the fact that Diff does not have pullbacks is commonly cited in the literature, we found it difficult to locate a detailed proof of this fact and so present one here.

Proposition 3.13. The category Diff does not have pullbacks.

Proof. Take X, Y, and Z to be equal to \mathbb{R} . Take f and g to be the respective functions from X to Z and Y to Z given for each x in X and y in Y by

$$f(x) = x^2 \quad \text{and} \quad g(y) = y^2.$$

Denote by Id the identity function from \mathbb{R} to \mathbb{R} . For any pullback Q in Diff of (f, g), since the span (Id, Id) is paired with (f, g), there is a span morphism Φ_1 in Diff from (Id, Id) to Q. The composite $q_L \circ \Phi_1$ is equal to Id, and so it is injective. Therefore, Φ_1 is injective. Since Φ_1 is a continuous injection from \mathbb{R} to Q_A , the dimension of Q_A is at least 1.

The fibered product $X \times_Z Y$ is the set

$$X \times_Z Y = \{(x, y) \colon |x| = |y|\}.$$

Take \mathcal{F} to be the forgetful functor from Diff to Set. The span (π_X, π_Y) is a pullback in Set of $(\mathcal{F}(f), \mathcal{F}(g))$. There are intervals I and J, each equal to either $(-\infty, 0)$ or $(0, \infty)$, so that the set U is nonempty, where U is given by

$$U = q_L^{-1}(I) \cap q_R^{-1}(J).$$

Otherwise, the span $(0_X, 0_Y)$ would be a pullback in Diff of (f, g), where 0_X and 0_Y are the respective zero functions from the one point manifold **1** to X and Y. Since \mathcal{F} preserves pullbacks, there would then be a bijection between the underlying set **1** and $X \times_Z Y$.

The set U is an open subset of Q_A as a non-empty intersection of open sets and is, therefore, a manifold. The restricted functions $f|_I$ and $g|_J$ are surjective submersions onto $(0,\infty)$. Proposition 3.7 implies that the fibered product $I \times_{(0,\infty)} J$ is a one-dimensional manifold. The span (π_I, π_J) is a pullback of $(f|_I, g|_J)$. Since $(q_L|_U, q_R|_U)$ is paired with $(f|_I, g|_J)$, there is a span morphism Φ_2 in Diff from $(q_L|_U, q_R|_U)$ to (π_I, π_J) . The function Φ_2 is just the restriction of Φ , given by (2), to U. Since Φ_2 is an injection, U has dimension at most 1, and so the dimension of U is equal to 1.

The dimension of the manifold Q_A is also equal to 1 since Q_A contains U as an open subset, and so Q_A is homeomorphic to either a line, an open interval, a half-open interval, or a circle, [7]. The subspace $X \times_Z Y \setminus \{(0,0)\}$ of $X \times Y$ has four connected components, but the preimage $\Phi^{-1}(X \times_Z Y \setminus \{(0,0)\})$ is the set Q_A with one point removed, and so has either one or two connected components. However, Lemma 3.12 implies that Φ is continuous. Since $X \times_Z Y \setminus \{(0,0)\}$ cannot have more connected components than its preimage, Q cannot be a pullback of (f,g), and so (f,g) does not have a pullback in Diff.

4. Composition by \mathcal{F} -Pullbacks

Assume, henceforth, that \mathscr{C} and \mathscr{C}' are categories and that \mathcal{F} is a functor from \mathscr{C} to \mathscr{C}' . For any span S in \mathscr{C} , denote by $\mathcal{F}(S)$ the span $(\mathcal{F}(s_L), \mathcal{F}(s_R))$ in \mathscr{C}' . For any cospan C in \mathscr{C} , denote by $\mathcal{F}(C)$ the cospan $(\mathcal{F}(c_L), \mathcal{F}(c_R))$ in \mathscr{C}' .

4.1. *F*-Pullbacks and Span Tightness.

Definition 4.1. The category \mathscr{C} has \mathcal{F} -pullbacks in \mathscr{C}' if for any cospan C in \mathscr{C} , there is a span S in \mathscr{C} that is paired with C and the span $\mathcal{F}(S)$ is a pullback of the cospan $\mathcal{F}(C)$ in \mathscr{C}' . In this case, the span S is an \mathcal{F} -pullback of C.

Note that if \mathscr{C}' is equal to \mathscr{C} and \mathcal{F} is the identity functor, then an \mathcal{F} -pullback is simply a pullback.

Definition 4.2. Suppose that S and Q are spans in \mathscr{C} such that:

- (1) $S_R = Q_L;$
- (2) there is a span P in \mathscr{C} that is a pullback of the cospan (s_R, q_L) .

The composite of S and Q along P is the span in \mathscr{C} given by

 $S \circ_P Q = (s_L \circ p_L, q_R \circ p_R).$

If P is an \mathcal{F} -pullback, then the span $S \circ_P Q$ is an \mathcal{F} -pullback composite of S and Q along P.

These morphisms determine the construction of the composite span $S \circ_P Q$ along the \mathcal{F} -pullback P:



The composite span $S \circ_P Q$ is this diagram:



Definition 4.3. Suppose that \mathscr{C} has \mathcal{F} -pullbacks in \mathscr{C}' . The functor \mathcal{F} is span tight if for any \mathcal{F} -pullbacks S and Q of the same cospan, the unique span isomorphism Φ from $\mathcal{F}(S)$ to $\mathcal{F}(Q)$ is $\mathcal{F}(\Psi)$ for some span isomorphism Ψ from S to Q.

4.2. \mathcal{F} -Pullbacks of SurjSub. Suppose that X, Y, and Z are smooth manifolds. Suppose further that f is a surjective submersion from X to Z and that g is a surjective submersion from Y to Z. Proposition 3.7 implies that the fibered product $X \times_Z Y$ is an embedded submanifold of $X \times Y$. Again, denote by ρ_X and ρ_Y the respective projections from $X \times Y$ to X and Y, and let π_X and π_Y be their respective restrictions to $X \times_Z Y$.

Proposition 4.4. For any cospan (f,g) in SurjSub, the span (π_X, π_Y) is a pullback in Diff of (f,g).

Proof. For any span Q in Diff that is paired with the cospan (f, g), take Φ to be the map from Q_A to $X \times Y$ that is given for any a in Q_A by

$$\Phi(a) = (q_L(a), q_R(a)).$$

This map is smooth as a product of smooth maps and unique since Diff has products. Furthermore, for any a in Q_A ,

$$(f \circ \rho_X \circ \Phi)(a) = f(q_L(a))$$
 and $(g \circ \rho_Y \circ \Phi)(a) = g(q_R(a))$.

Since Q is paired with (f,g), $f(q_L(a))$ is equal to $g(q_R(a))$, and so $\Phi(a)$ is in $X \times_Z Y$. Therefore, the span (π_X, π_Y) is a pullback in Diff.

Note that while SurjSub is a subcategory of Diff, the category SurjSub does not have pullbacks. Take \mathcal{F} to be the inclusion functor from SurjSub to Diff. For any cospan (f,g) in SurjSub, where f and g have respective sources X and Y and both maps have target Z, Proposition 4.4 implies that the span (π_X, π_Y) is an \mathcal{F} -pullback of the cospan (f, g). Take Q to be another \mathcal{F} -pullback of (f, g). There is a span isomorphism Φ from $\mathcal{F}(Q)$ to $(\mathcal{F}(\pi_X), \mathcal{F}(\pi_Y))$, since both spans are pullbacks of $(\mathcal{F}(f), \mathcal{F}(g))$ in Diff. Every diffeomorphism is a surjective submersion, and this implies Theorem 4.5.

Theorem 4.5. The inclusion functor from SurjSub to Diff is span tight.

5. GENERALIZED SPAN CATEGORY

Define the objects in $\text{Span}(\mathscr{C}, \mathcal{F})$ to be the objects in \mathscr{C} and the isomorphism classes of spans in \mathscr{C} to be the morphisms in $\text{Span}(\mathscr{C}, \mathcal{F})$. For any isomorphism class of spans [S] in $\text{Span}(\mathscr{C}, \mathcal{F})$, identify S_R and S_L respectively to be the source and target of [S]. Define composition of isomorphism classes of spans by

$$\left[S^{1}\right]\circ\left[S^{2}\right]=\left[S^{1}\circ_{P}S^{2}\right],$$

where $S^1 \circ_P S^2$ is an \mathcal{F} -pullback composite of S^1 and S^2 . Theorem 5.1 is the main result of the section and the lemmata that follow simplify the proof of the theorem.

Theorem 5.1. If \mathcal{F} is a span tight functor from \mathscr{C} to \mathscr{C}' , then $\operatorname{Span}(\mathscr{C}, \mathcal{F})$ is a category.

If the functor \mathcal{F} from \mathscr{C} to \mathscr{C}' is span tight and S and Q are spans in \mathscr{C} with S_R equal to Q_L , then there is an \mathcal{F} -pullback P of the cospan (s_R, q_L) , and so there is an \mathcal{F} -pullback composite of S and Q along P. The \mathcal{F} -pullback P is, however, only defined up to a span isomorphism Φ . The following lemma shows that changing P, up to an isomorphism, changes the resulting composite span only up to a span isomorphism in \mathscr{C} .

Lemma 5.2. Suppose that \mathcal{F} is span tight, that S and Q are spans in \mathscr{C} , and that $S \circ_{P^i} Q$ is an \mathcal{F} -pullback composite, with i equal to 1 or 2. There is a span isomorphism Φ in \mathscr{C} from $S \circ_{P^1} Q$ to $S \circ_{P^2} Q$.

Proof. Since P^1 and P^2 are both \mathcal{F} -pullbacks of the cospan (s_R, q_L) , there is a span isomorphism Φ in \mathscr{C}' from $\mathcal{F}(P^1)$ to $\mathcal{F}(P^2)$. Since \mathcal{F} is span tight, there is a span isomorphism Ψ in \mathscr{C} from P^1 to P^2 with $\mathcal{F}(\Psi)$ equal to Φ , and so

(3)
$$p_L^1 = p_L^2 \circ \Psi$$
 and $p_R^1 = p_R^2 \circ \Psi$.

Left compose s_L with both sides of the left equality of (3) and q_R with both sides of the right equality of (3) to obtain the respective equalities

$$s_L \circ p_L^1 = s_L \circ p_L^2 \circ \Psi$$
 and $q_R \circ p_R^1 = q_R \circ p_R^2 \circ \Psi$

that demonstrate that Ψ is a span isomorphism from $S \circ_{P^1} Q$ to $S \circ_{P^2} Q$.

Lemma 5.3. Suppose that \mathcal{F} is span tight, that S^i and Q^i are spans in \mathscr{C} , and that $S^i \circ_{P^i} Q^i$ is an \mathcal{F} -pullback composite, with i equal to 1 or 2. Suppose that S^1 and Q^1 are respectively span isomorphic to S^2 and Q^2 . There is a span isomorphism in \mathscr{C} between spans $S^1 \circ_{P^1} Q^1$ and $S^2 \circ_{P^2} Q^2$.

This diagram includes many of the mappings involved in the proof of Lemma 5.3:



Proof. Take α and β to be span isomorphisms from S^1 to S^2 and from Q^1 to Q^2 , respectively. The span P^1 is an \mathcal{F} -pullback of (s_R^1, q_L^1) . Since α and β are span morphisms, the span

 $(\alpha \circ p_L^1, \beta \circ p_R^1)$ is paired with (s_R^2, q_L^2) . Since $\mathcal{F}(P^2)$ is a pullback of $(\mathcal{F}(s_R^2), \mathcal{F}(q_L^2))$, there is a span morphism Φ_1 in \mathscr{C}' from $(\mathcal{F}(\alpha \circ p_L^1), \mathcal{F}(\beta \circ p_R^1))$ to $\mathcal{F}(P^2)$.

If T is a span in \mathscr{C}' that is paired with $(\mathcal{F}(s_R^1), \mathcal{F}(q_L^1))$, then there is a span morphism Φ_2 in \mathscr{C}' from T to $\mathcal{F}(P^1)$. The composite $\Phi_1 \circ \Phi_2$ is a span morphism from T to $(\mathcal{F}(\alpha^{-1} \circ p_L^2), \mathcal{F}(\beta^{-1} \circ p_R^2))$, which is also paired with $(\mathcal{F}(s_R^1), \mathcal{F}(q_L^1))$. Uniqueness of the pullback of $(\mathcal{F}(s_R^1), \mathcal{F}(q_L^1))$ up to a span isomorphism implies that there is a span isomorphism Φ_3 in \mathscr{C}' from $(\mathcal{F}(\alpha^{-1} \circ p_L^2), \mathcal{F}(\beta^{-1} \circ p_R^2))$ to $\mathcal{F}(P^1)$. Since \mathcal{F} is span tight, there is a span isomorphism Ψ in \mathscr{C} such that $\mathcal{F}(\Psi)$ is Φ_3 . Use the fact that Ψ is a span isomorphism to obtain the equalities

(4)
$$\alpha^{-1} \circ p_L^2 = p_L^1 \circ \Psi \quad \text{and} \quad \beta^{-1} \circ p_R^2 = p_R^1 \circ \Psi.$$

The equalities

$$s_L^2 = s_L^1 \circ \alpha^{-1}$$
 and $q_R^2 = q_R^1 \circ \beta^{-1}$

together with (4) imply that

$$s_L^2 \circ p_L^2 = s_L^1 \circ \alpha^{-1} \circ p_L^2 = s_L^1 \circ p_L^1 \circ \Psi$$

and similarly that

$$q_R^2 \circ p_R^2 = q_R^1 \circ p_R^1 \circ \Psi,$$

and so Ψ is a span isomorphism from $S^2 \circ_{P^2} Q^2$ to $S^1 \circ_{P^1} Q^1$.

Lemma 5.3 generalizes Lemma 5.2 and reduces to Lemma 5.2 when S^1 is equal to S^2 , when Q^1 is equal to Q^2 , and when P^1 and P^2 are pullbacks that are not necessarily equal to each other.

Lemma 5.4. Suppose that \mathcal{F} is span tight and that S, Q, and T are spans in \mathscr{C} with S_R equal to Q_L and Q_R equal to T_L . Suppose that $S \circ_{P^1} Q$ and $Q \circ_{P^4} T$ are \mathcal{F} -pullback composites and that $(S \circ_{P^1} Q) \circ_{P^2} T$ and $S \circ_{P^3} (Q \circ_{P^4} T)$ are also \mathcal{F} -pullback composites. There is a span isomorphism from $(S \circ_{P^1} Q) \circ_{P^2} T$ to $S \circ_{P^4} (Q \circ_{P^3} T)$.

The mappings involved in the proof of Lemma 5.4 appear in this commutative diagram:



and in this commutative diagram:



Proof. Suppose that P^1 is an \mathcal{F} -pullback of the cospan (s_R, q_L) , that P^3 is an \mathcal{F} -pullback of the cospan (q_R, t_L) , and that P is an \mathcal{F} -pullback of the cospan (p_R^1, p_L^3) , where

$$P_L = P_A^1$$
 and $P_R = P_A^3$.

Suppose further that P^2 is an \mathcal{F} -pullback of the cospan $(q_R \circ p_R^1, t_L)$ and that P^4 is an \mathcal{F} -pullback of the cospan $(s_R, q_L \circ p_L^3)$. Since P^2 is an \mathcal{F} -pullback of the cospan $(q_R \circ p_R^1, t_L)$, the span $(p_R^1 \circ p_L^2, p_R^2)$ is paired with the cospan (q_R, t_L) , and so $(\mathcal{F}(p_R^1 \circ p_L^2), \mathcal{F}(p_R^2))$ is paired with the cospan $(\mathcal{F}(q_R), \mathcal{F}(t_L))$. The span P^3 is an \mathcal{F} -pullback, which implies the existence of a span morphism Φ_1 in \mathscr{C}' from $(\mathcal{F}(p_R^1 \circ p_L^2), \mathcal{F}(p_R^2))$ to $\mathcal{F}(P^3)$. The span $(\mathcal{F}(p_L^2), \Phi_1)$ is paired with $(\mathcal{F}(p_R^1), \mathcal{F}(p_L^3))$, and so there is a span morphism Φ_2 in \mathscr{C}' from $(\mathcal{F}(p_L^2), \Phi_1)$ to $\mathcal{F}(P)$.

If U is a span paired with $(q_R \circ p_R^1, t_L)$, then there is a span morphism Φ_3 in \mathscr{C}' from $\mathcal{F}(U)$ to $\mathcal{F}(P^2)$. The composite $\Phi_2 \circ \Phi_3$ is a span morphism in \mathscr{C}' from $\mathcal{F}(U)$ to $(\mathcal{F}(p_L), \mathcal{F}(p_R^3 \circ p_R))$, and so $(\mathcal{F}(p_L), \mathcal{F}(p_R^3 \circ p_R))$ is a pullback in \mathscr{C}' of the cospan $(\mathcal{F}(q_R \circ p_R^1), \mathcal{F}(t_L))$. There is, therefore, a span isomorphism in \mathscr{C}' from $(\mathcal{F}(p_L), \mathcal{F}(p_R^3 \circ p_R))$ to $\mathcal{F}(P^2)$. Span tightness of \mathcal{F} implies that there is a span isomorphism Ψ_1 in \mathscr{C} from $(p_L, p_R^3 \circ p_R)$ to P^2 , with

$$p_R^2 \circ \Psi_1 = p_R^3 \circ p_R$$
 and so $t_R \circ p_R^2 \circ \Psi_1 = t_R \circ p_R^3 \circ p_R$

The equality

 $p_L^2 \circ \Psi_1 = p_L$ implies that $s_L \circ p_L^1 \circ p_L^2 \circ \Psi_1 = s_L \circ p_L^1 \circ p_L.$

The isomorphism Ψ_1 in \mathscr{C} is, therefore, a span isomorphism, with

$$\left(s_L \circ p_L^1 \circ p_L, t_R \circ p_R^3 \circ p_R\right) = \left(s_L \circ p_L^1 \circ p_L^2 \circ \Psi_1, t_R \circ p_R^2 \circ \Psi_1\right),$$

and so $(s_L \circ p_L^1 \circ p_L, t_R \circ p_R^3 \circ p_R)$ and $(S \circ_{P^1} Q) \circ_{P^2} T$ are span isomorphic.

A similar argument shows that $(s_L \circ p_L^1 \circ p_L, t_R \circ p_R^3 \circ p_R)$ and $S \circ_{P^4} (Q \circ_{P^3} T)$ are span isomorphic, where P^4 is an \mathcal{F} -pullback of the cospan $(s_R, q_L \circ p_L^3)$. Since $(S \circ_{P^1} Q) \circ_{P^2} T$ and $S \circ_{P^4} (Q \circ_{P^3} T)$ are span isomorphic to the same span, Proposition 2.3 implies Lemma 5.4. \Box Proof of Theorem 5.1. To prove the theorem, it suffices to show that the composition of morphisms in $\text{Span}(\mathscr{C}, \mathcal{F})$ is well defined, satisfies the left and right unit laws, and is associative.

For any two isomorphism classes of spans $[S^1]$ and $[S^2]$, where the source of $[S^1]$ is the target of $[S^2]$, if S^1 and S^2 are, respectively, representatives of $[S^1]$ and $[S^2]$, then there is an \mathcal{F} -pullback P of (s_R^1, s_L^2) . Hence, there exists a composite $S^1 \circ_P S^2$. Lemma 5.2 implies that the equivalence class $[S^1 \circ_P S^2]$ is independent of P. Lemma 5.3 additionally implies that $[S^1 \circ_P S^2]$ is independent of the choice of representatives S^1 and S^2 . Furthermore, S_R^2 is the source of $[S^1] \circ [S^2]$ and S_L^1 is the target of $[S^1] \circ [S^2]$, and so the composition \circ is well defined.

Suppose that [S] is an isomorphism class of spans in \mathscr{C} and that $[I_{S_R}]$ is the isomorphism class of spans containing (Id_{S_R}, Id_{S_R}) , where

$$\mathrm{Id}_{S_R}\colon S_R\to S_R$$

is the identity map from S_R to S_R . Take P to be the span (Id_{S_A}, s_R) . For any span Q in \mathscr{C}' that is paired with $(\mathcal{F}(s_R), \mathcal{F}(\mathrm{Id}_{S_R}))$,

$$\mathcal{F}(s_R) \circ q_L = \mathcal{F}(\mathrm{Id}_{S_R}) \circ q_R = q_R$$

and so q_L is a span morphism in \mathscr{C}' from Q to $\mathcal{F}(P)$. Given any other span morphism Φ in \mathscr{C}' from Q to $\mathcal{F}(P)$,

$$q_L = \mathcal{F}(\mathrm{Id}_{S_A}) \circ \Phi = \mathrm{Id}_{\mathcal{F}(S_A)} \circ \Phi = \Phi.$$

The span morphism in \mathscr{C}' from Q to $\mathcal{F}(P)$ is unique, which implies that $\mathcal{F}(P)$ is a pullback in \mathscr{C}' of $(\mathcal{F}(s_R), \mathcal{F}(\mathrm{Id}_{S_R}))$, and so P is an \mathcal{F} -pullback of (s_R, Id_{S_R}) .

The composite $S \circ_P I_{S_R}$ is determined by these mappings:



and is this diagram:



Since composition is well defined and $S \circ_P I_{S_R}$ is span isomorphic in \mathscr{C} to S, the composite $[S] \circ [I_{S_R}]$ is equal to [S]. Similar arguments will show that $[I_{S_L}] \circ [S]$ is equal to [S], and so $\operatorname{Span}(\mathscr{C}, \mathcal{F})$ has both a right and left unit law.

Lemma 5.4 implies that \circ is associative.

6. Examples

6.1. Categories that have Pullbacks. Suppose that \mathscr{C} is a category that has pullbacks, and let \mathcal{F} be the identity functor from \mathscr{C} to \mathscr{C} . The functor \mathcal{F} is span tight and so $\operatorname{Span}(\mathscr{C}, F)$ is a category. Since every \mathcal{F} -pullback of a cospan is a pullback of a cospan, the category $\operatorname{Span}(\mathscr{C}, \mathcal{F})$ is the category $\operatorname{Span}(\mathscr{C})$. In this way, the concept of a generalized span category $\operatorname{Span}(\mathscr{C}, \mathcal{F})$ generalizes the notion of a span category and reduces to it when \mathscr{C} has pullbacks and \mathcal{F} is the identity functor.

6.2. Smooth Manifolds and Surjective Submersions. Suppose that \mathcal{F} is the inclusion functor from SurjSub to Diff. Theorems 4.5 and 5.1 together imply that Span(SurjSub, \mathcal{F}) is a category.

6.3. Classical Mechanics. In [4], we worked in the categories RiemSurj, whose objects are Riemannian manifolds and whose morphisms are surjective Riemannian submersions, and SympSurj, whose objects are symplectic manifolds and whose morphisms are surjective Poisson maps. Unlike SurjSub, these categories are not subcategories of Diff. However, the forgetful functors from these categories into Diff are still span tight, and so it is possible to construct generalized span categories in these settings which are critical to the study of classical mechanics. In [4], we introduced the notion of an *augmented generalized span category*. Such categories are critical to the categorification of classical mechanics and the study of the functoriality of the Legendre transformation.

References

- [1] Awodey, S.: Category Theory. Oxford University Press New York (2010). (Referred to on page 5.)
- [2] Baez, J. C.: Personal communication. (Referred to on page 2.)
- Baez, J. C., Pollard, B.: A compositional framework for reaction networks. Rev. Math. Phys. 29 (2017). (Referred to on page 2.)
- Baez, J.C., Weisbart, D. and Yassine, A.M.: Open Systems in Classical Mechanics. J. Math Phys. 62, 042902 (2021). (Referred to on page 3, 9, 20.)
- [5] Bénabou, J.: Introduction to bicategories. Reports of the Midwest Category Seminar, Singapore (Lecture Notes in Mathematics) 47, Springer Berlin Heidelberg, 1967. (Referred to on page 2, 5.)
- [6] Borceux, F.: Handbook Of Categorical Algebra I: Basic Category Theory. Cambridge University Press (1994). (Referred to on page 6.)
- [7] Gale, D.: The classification of 1-manifolds: a take-home exam. The American Mathematical Monthly. Vol. 94, No. 2 (Feb., 1987), pp. 170-175. (Referred to on page 13.)
- [8] Haugseng, R.: Iterated spans and classical topological field theories. Math. Z. 289 (2018), 1427–1488. (Referred to on page 2.)
- [9] Lee, J.: Introduction To Smooth Manifolds. Springer Berlin (2015). (Referred to on page 6, 10.)
- [10] Leinster, T.: Higher Operads, Higher Categories. London Mathematics Society Lecture Note Series 298, Cambridge University Press (2004). (Referred to on page 2.)
- [11] MacLane, S.: Categories For The Working Mathematician. Springer Berlin (1998). (Referred to on page 5.)

- [12] Spivak, D.: Derived smooth manifolds. Duke Math. J. Volume 153, Number 1 (2010), 55–128. (Referred to on page 3.)
- [13] Yoneda, N.: On Ext and exact sequences. J. Fac. Sci. Univ. Tokyo, Sec. I 7 (1954), 193–227. (Referred to on page 2.)