

New curvature tensors along Riemannian submersions

Mehmet Akif Akyol and Gülhan Ayar

Abstract. In 1966, B. O'Neill [The fundamental equations of a submersion, Michigan Math. J., Volume 13, Issue 4 (1966), 459-469.] obtained some fundamental equations and curvature relations between the total space, the base space and the fibres of a submersion. In the present paper, we define new curvature tensors along Riemannian submersions such as Weyl projective curvature tensor, concircular curvature tensor, conharmonic curvature tensor, conformal curvature tensor and M -projective curvature tensor, respectively. Finally, we obtain some results in case of the total space of Riemannian submersions has umbilical fibres for any curvature tensors mentioned by the above.

Mathematics Subject Classification (2010). Primary 53C15, 53B20.

Keywords. Riemannian submersion, Weyl projective curvature tensor, M -projective curvature tensor, concircular curvature tensor, conformal curvature tensor, conharmonic curvature tensor.

1. Introduction and Preliminaries

In differential geometry, an important tool to define the curvature of n -dimensional spaces (such as Riemannian manifolds) is the Riemannian curvature tensor. The tensor has played an important role both general relativity and gravity. In this manner, Mishra in [10] defined some new curvature tensors on Riemannian manifolds such as concircular curvature tensor, conharmonic curvature tensor, conformal curvature tensor, respectively. Taking into account the paper of Mishra, Pokhariyal and Mishra defined the Weyl projective curvature tensor on Riemannian manifolds [13]. Afterwards, Ojha defined M -projective curvature tensor [11].

Curvature tensors also play an important role in physics. Relevance in physics of the tensors considered in our work and some of the important studies that focus on the geometric properties of curvature tensors are; in 1988 conditions of conharmonic curvature tensor of Kaehler hypersurfaces in

complex space forms has been analysed by M. Doric et al. [19]. By the way, the relativistic significance of concircular curvature tensor has been studied by Ahsan [18] and this tensor has been explored in that study. Finally in 2018 based on Einsteins geodesic postulate, projectively related connections on a space-time manifold and the closely related Weyl projective tensor has been examined detailed by G. Hall [17].

Riemannian submersion appears to have been studied and its differential geometry has first defined by O'Neill 1966 and Gray 1967 [12]. We note that Riemannian submersions have been studied widely not only in mathematics, but also in theoretical physics because of their applications in the Yang-Mills theory, Kaluza Klein theory, super gravity, relativity and superstring theories (see [2], [3], [7], [8], [9], [16]). Most of the studies related to Riemannian submersion can be found in the books ([5], [15]). In 1966, B. O'Neill has defined a paper related to some fundamental equations of a submersion. In that paper, he has given some curvature relations on Riemannian submersions.

In this study, in addition to the curvature relations previously defined on Riemannian submersion, we investigate new curvature tensors on a Riemannian submersion and the curvature properties of these tensors. In the present paper, in the first part of our study, the basic definitions and theorems that we will use throughout the paper are given. In sections 2-6 include the Weyl projective curvature tensor, concircular curvature tensor, conharmonic curvature tensor, conformal curvature tensor and M -projective curvature tensor relations for a Riemannian submersion respectively. Also various results are obtained by examining the conditions for having total umbilical fibers.

Now, we will give the basic definitions and theorems without proofs that we will use throughout the paper.

Definition 1.1. Let (M, g) and (N, g_N) be Riemannian manifolds, where $\dim(M) > \dim(N)$. A surjective mapping $\pi : (M, g) \rightarrow (N, g_N)$ is called a *Riemannian submersion* [12] if:

(S1) The rank of π equals $\dim(N)$.

In this case, for each $q \in N$, $\pi^{-1}(q) = \pi_q^{-1}$ is a k -dimensional submanifold of M and called a *fiber*, where $k = \dim(M) - \dim(N)$. A vector field on M is called *vertical* (resp. *horizontal*) if it is always tangent (resp. orthogonal) to fibers. A vector field X on M is called *basic* if X is horizontal and π -related to a vector field X_* on N , i.e. , $\pi_*(X_p) = X_{*\pi(p)}$ for all $p \in M$, where π_* is derivative or differential map of π . We will denote by \mathcal{V} and \mathcal{H} the projections on the vertical distribution $\ker\pi_*$, and the horizontal distribution $\ker\pi_*^\perp$, respectively. As usual, the manifold (M, g) is called *total manifold* and the manifold (N, g_N) is called *base manifold* of the submersion $\pi : (M, g) \rightarrow (N, g_N)$.

(S2) π_* preserves the lengths of the horizontal vectors.

This condition is equivalent to say that the derivative map π_* of π , restricted to $\ker\pi_*^\perp$, is a linear isometry.

If X and Y are the basic vector fields, π -related to X_N, Y_N , we have the following facts:

1. $g(X, Y) = g_N(X_N, Y_N) \circ \pi$,
2. $h[X, Y]$ is the basic vector field π -related to $[X_N, Y_N]$,
3. $h(\nabla_X Y)$ is the basic vector field π -related to $\nabla^N_{X_N} Y_N$,

for any vertical vector field V , $[X, V]$ is the vertical.

The geometry of Riemannian submersions is characterized by O'Neill's tensors \mathcal{T} and \mathcal{A} , defined as follows:

$$\mathcal{T}_E F = v\nabla_{vE} hF + h\nabla_{vE} vF, \quad (1.1)$$

$$\mathcal{A}_E F = v\nabla_{hE} hF + h\nabla_{hE} vF \quad (1.2)$$

for any vector fields E and F on M , where ∇ is the Levi-Civita connection, v and h are orthogonal projections on vertical and horizontal spaces, respectively.

We now recall the following curvature relations for a Riemannian submersion from [5] and [12].

Theorem 1.2. (M, g) and (G, g') Riemannian manifolds,

$$\pi : (M, g) \rightarrow (G, g')$$

a Riemannian submersion and R^M , R^G and \hat{R} be Riemannian curvature tensors of M, G and $(\pi^{-1}(x), \hat{g}_x)$ fibre respectively. In this case, there are the following equations for any $U, V, W, F \in \chi^v(M)$ and $X, Y, Z, H \in \chi^h(M)$

$$\begin{aligned} g(R^M(X, Y)Z, H) &= g(R^G(X, Y)Z, H) + 2g(A_X Y, A_Z H) \\ &\quad - g(A_Y Z, A_X H) + g(A_X Z, A_Y H), \end{aligned} \quad (1.3)$$

$$\begin{aligned} g(R^M(X, Y)Z, V) &= -g((\nabla_Z A)_X Y, V) - g(A_X Y, T_V Z) \\ &\quad + g(A_Y Z, T_V X) - g(A_X Z, T_V Y), \end{aligned} \quad (1.4)$$

$$\begin{aligned} g(R^M(X, Y)V, W) &= g((\nabla_V A)_X Y, W) - g((\nabla_W A)_X Y, V) \\ &\quad + g(A_X V, A_Y W) + g(A_X W, A_Y V) \\ &\quad - g(T_V X, T_W Y) + g(T_W X, T_V Y), \end{aligned} \quad (1.5)$$

$$\begin{aligned} g(R^M(X, V)Y, W) &= g((\nabla_X T)_V W, Y) - g((\nabla_V A)_X Y, W) \\ &\quad - g(T_V X, T_W Y) + g(A_X V, A_Y W), \end{aligned} \quad (1.6)$$

$$g(R^M(U, V)W, X) = g((\nabla_U T)_V W, X) - g((\nabla_V T)_U W, X) \quad (1.7)$$

and

$$g(R^M(U, V)W, F) = g(\hat{R}(U, V)W, F) + g(T_U W, T_V F) - g(T_V W, T_U F). \quad (1.8)$$

Definition 1.3. [5] Let (M, g) be a Riemannian manifold and a local orthonormal frame of the vertical distribution ν is $\{U_j\}_{1 \leq j \leq r}$. Then N , the horizontal vector field on (M, g) is locally defined by

$$N = \sum_{j=1}^r T_{U_j} U_j.$$

Proposition 1.4. Let (M, g) and (G, g') Riemannian manifolds,

$$\pi : (M, g) \rightarrow (G, g')$$

a Riemannian submersion and $\{X_i, U_j\}$ be a π -compatible frame.

In this case, for any $U, V \in \chi^v(M)$ and $X, Y \in \chi^h(M)$, the Ricci tensor S^M satisfies the following equations [5]:

$$(i) \quad S^M(U, V) = \hat{S}(U, V) - g(N, T_U V) + \sum_i \{g((\nabla_{X_i} T)_U V, X_i) + g(A_{X_i} U, A_{X_i} V)\}, \quad (1.9)$$

$$(ii) \quad S^M(X, Y) = S^G(X', Y') \circ \pi + \frac{1}{2} \{g(\nabla_X N, Y) + g(\nabla_Y N, X)\} - 2 \sum_i g(A_X X_i, A_Y X_i) - \sum_j g(T_{U_j} X, T_{U_j} Y), \quad (1.10)$$

$$(iii) \quad S^M(U, X) = g(\nabla_U N, X) - \sum_j g(\nabla_{U_j} T)_{U_j} U, X) + \sum_i \{g((\nabla_{X_i} A)_{X_i} X, U) - 2g(A_X X_i, T_U X_i)\}. \quad (1.11)$$

Proposition 1.5. [5] Let's take the scalar curvatures of $(M, g), (G, g')$ Riemannian manifolds and $x \in G, \pi^{-1}(x)$ fibre with r^M, r^G and \hat{r} , respectively. In a

$$\pi : (M, g) \rightarrow (G, g')$$

Riemannian submersion, (M, g) depends on the scalar curvature of the Riemannian manifold r^G and the scalar curve of any lift \hat{r} . In this case

$$r^M = \hat{r} + r^G \circ \pi - \|N\|^2 - \|A\|^2 - \|T\|^2 + 2 \sum_i g(\nabla_{X_i} N, X_i). \quad (1.12)$$

2. Weyl projective curvature tensor along a Riemannian submersion

In this section, we examine the Weyl projective curvature tensor relations between the total space, the base space and fibres on a Riemannian submersion. We also give a corollary in case of the Riemannian submersion has totally umbilical fibres.

Definition 2.1. [10] Let take an n -dimensional differentiable manifold M with differentiability class C^∞ . In the n -dimensional space V_n , the tensor

$$P^*(X, Y)Z = R^M(X, Y)Z - \frac{1}{n-1}\{S^M(Y, Z)X - S^M(X, Z)Y\}.$$

is called Weyl projective curvature tensor, where Ricci tensor of total space denoted by S^M .

Now, we have the following main theorem.

Theorem 2.2. *Let (M, g) and (G, g') Riemannian manifolds,*

$$\pi : (M, g) \rightarrow (G, g')$$

a Riemannian submersion and R^M, R^G and \hat{R} be Riemannian curvature tensors, S^M, S^G and \hat{S} be Ricci tensors of M, G and the fibre respectively. Then for any $U, V, W, F \in \chi^v(M)$ and $X, Y, Z, H \in \chi^h(M)$, we have the following relations for Weyl projective curvature tensor:

$$\begin{aligned} g(P^*(X, Y)Z, H) &= g(R^G(X, Y)Z, H) + 2g(A_X Y, A_Z H) - g(A_Y Z, A_X H) + g(A_X Z, A_Y H) \\ &- \frac{1}{n-1} \left\{ g(X, H) \left[S^G(Y', Z') \circ \pi + \frac{1}{2} (g(\nabla_Y N, Z) + g(\nabla_Z N, Y)) \right. \right. \\ &- 2 \sum_i g(A_Y X_i, A_Z X_i) - \sum_j g(T_{U_j} Y, T_{U_j} Z) \left. \right] \\ &- g(Y, H) \left[S^G(X', Z') \circ \pi + \frac{1}{2} (g(\nabla_X N, Z) + g(\nabla_Z N, X)) \right. \\ &- 2 \sum_i g(A_X X_i, A_Z X_i) - \sum_j g(T_{U_j} X, T_{U_j} Z) \left. \right] \left. \right\}, \end{aligned}$$

$$\begin{aligned} g(P^*(X, Y)Z, V) &= -g((\nabla_Z A)_X Y, V) - g(A_X Y, T_V Z) + g(A_Y Z, T_V X) \\ &- g(A_X Z, T_V Y), \end{aligned}$$

$$\begin{aligned} g(P^*(X, Y)V, W) &= g((\nabla_V A)_X Y, W) - g((\nabla_W A)_X Y, V) + g(A_X V, A_Y W) \\ &- g(A_X W, A_Y V) - g(T_V X, T_W Y) + g(T_W X, T_V Y), \end{aligned}$$

$$\begin{aligned} g(P^*(X, V)Y, W) &= g((\nabla_X T)_V W, Y) + g((\nabla_V A)_X Y, W) - g(T_V X, T_W Y) \\ &+ g(A_X V, A_Y W), \end{aligned}$$

$$g(P^*(U, V)W, X) = g((\nabla_U T)_V W, X) - g((\nabla_V T)_U W, X)$$

and

$$\begin{aligned}
g(P^*(U, V)W, F) &= g(\hat{R}(U, V)W, F) + g(T_U W, T_V F) - g(T_V W, T_U F) \\
&\quad - \frac{1}{n-1} \left\{ g(F, U) \left[\hat{S}(V, W) - g(N, T_V W) \right. \right. \\
&\quad \left. \left. + \sum_i g((\nabla_{X_i} T)_V W, X_i) + g(A_{X_i} V, A_{X_i} W) \right] \right. \\
&\quad \left. - g(F, V) \left[\hat{S}(U, W) - g(N, T_U W) \right. \right. \\
&\quad \left. \left. + \sum_i g((\nabla_{X_i} T)_U W, X_i) + g(A_{X_i} U, A_{X_i} W) \right] \right\}.
\end{aligned}$$

Proof. We only give the proof of the 1st equation of this theorem. The following equations are obtained inner production with H to P^* and using (1.3) and (1.10) equations.

$$\begin{aligned}
g(R^M(X, Y)Z, H) &= g(R^G(X, Y)Z, H) + 2g(A_X Y, A_Z H) - g(A_Y Z, A_X H) \\
&\quad + g(A_X Z, A_Y H),
\end{aligned}$$

$$\begin{aligned}
S^M(Y, Z) &= S^G(Y', Z') \circ \pi + \frac{1}{2} \{g(\nabla_Y N, Z) + g(\nabla_Z N, Y)\} \\
&\quad - 2 \sum_i g(A_Y Y_i, A_Z Y_i) - \sum_j (T_{U_j} Y, T_{U_j} Z)
\end{aligned}$$

and

$$\begin{aligned}
S^M(X, Z) &= S^G(Y', Z') \circ \pi + \frac{1}{2} \{g(\nabla_X N, Z) + g(\nabla_Z N, X)\} \\
&\quad - 2 \sum_i g(A_X X_i, A_Z X_i) - \sum_j (T_{U_j} X, T_{U_j} Z).
\end{aligned}$$

When these equations are substituted in P^* , the given result is obtained. Other equations are similarly proved by using Theorem 1.2 and Proposition 1.4. \square

Corollary 2.3. *Let $\pi : (M, g) \rightarrow (G, g')$ be a Riemannian submersion, where (M, g) and (G, g') Riemannian manifolds. If the Riemannian submersion has total umbilical fibres, that is $N = 0$, then the Weyl projective curvature tensor*

is given by

$$\begin{aligned}
& g(P^*(X, Y)Z, H) \\
&= g(R^G(X, Y)Z, H) + 2g(A_X Y, A_Z H) - g(A_Y Z, A_X H) + g(A_X Z, A_Y H) \\
&- \frac{1}{n-1} \left\{ g(X, H) \left[S^G(Y', Z') \circ \pi - 2 \sum_i g(A_Y X_i, A_Z X_i) - \sum_j g(T_{U_j} Y, T_{U_j} Z) \right] \right. \\
&- \left. g(Y, H) \left[S^G(X', Z') \circ \pi - 2 \sum_i g(A_X X_i, A_Z X_i) - \sum_j g(T_{U_j} X, T_{U_j} Z) \right] \right\},
\end{aligned}$$

and

$$\begin{aligned}
& g(P^*(U, V)W, F) \\
&= g(\hat{R}(U, V)W, F) + g(T_U W, T_V F) - g(T_V W, T_U F) \\
&- \frac{1}{n-1} \left\{ g(F, U) \left[\hat{S}(V, W) + \sum_i (g((\nabla_{X_i} T)_V W, X_i) + g(A_{X_i} V, A_{X_i} W)) \right] \right. \\
&- \left. g(F, V) \left[\hat{S}(U, W) + \sum_i (g((\nabla_{X_i} T)_U W, X_i) + g(A_{X_i} U, A_{X_i} W)) \right] \right\}.
\end{aligned}$$

for any $U, V, W, F \in \chi^v(M)$ and $X, Y, Z, H \in \chi^h(M)$,

3. Concircular curvature tensor along a Riemannian submersion

In this section, curvature relations of concircular curvature tensor in a Riemannian submersion are examined. In particular, we show that the Riemannian submersion with concircular curvature tensor has no the totally umbilical fibres.

Definition 3.1. In the n -dimensional space V_n , the tensor

$$C^*(X, Y, Z, H) = R^M(X, Y, Z, H) - \frac{r^M}{n(n-1)} [g(X, H)g(Y, Z) - g(Y, H)g(X, Z)],$$

is called concircular curvature tensor, where scalar tensor denoted by r^M [10].

Now, we have the following main theorem.

Theorem 3.2. Let (M, g) and (G, g') Riemannian manifolds,

$$\pi : (M, g) \rightarrow (G, g')$$

a Riemannian submersion and R^M , R^G and \hat{R} be Riemannian curvature tensors, r^M , r^G and \hat{r} be scalar curvature tensors of M , G and the fibre

respectively. Then for any $U, V, W, F \in \chi^v(M)$ and $X, Y, Z, H \in \chi^h(M)$, we have the following relations

$$\begin{aligned} g(C^*(X, Y)Z, H) &= g(R^G(X, Y)Z, H) + 2g(A_X Y, A_Z H) \\ &\quad - g(A_Y Z, A_X H) + g(A_X Z, A_Y H) \\ &\quad - \frac{r^M}{n(n-1)} \left\{ g(Y, Z)g(X, H) - g(X, Z)g(Y, H) \right\}, \end{aligned}$$

$$\begin{aligned} g(C^*(X, Y)Z, V) &= -g((\nabla_Z A)_X Y, V) - g(A_X T, T_V Z) \\ &\quad + g(A_Y Z, T_V X) - g(A_X Z, T_V Y), \end{aligned}$$

$$\begin{aligned} g(C^*(X, Y)V, W) &= g((\nabla_V A)_X Y, W) - g((\nabla_W A)_X Y, V) + g(A_X V, A_Y W) \\ &\quad - g(A_X W, A_Y V) - g(T_V X, T_W Y) + g(T_W X, T_V Y), \end{aligned}$$

$$\begin{aligned} g(C^*(X, V)Y, W) &= g((\nabla_X T)_V W, Y) + g((\nabla_V A)_X Y, W) - g(T_V X, T_W Y) \\ &\quad + g(A_X Y, A_Y W) - \frac{r^M}{n(n-1)} \{-g(X, Y)g(V, W)\}, \end{aligned}$$

$$g(C^*(U, V)W, X) = g((\nabla_U T)_V W, X) - g((\nabla_V T)_U W, X)$$

and

$$\begin{aligned} g(C^*(U, V)W, F) &= g(\hat{R}(U, V)W, F) + g(T_U W, T_V F) - g(T_V W, T_U F) \\ &\quad - \frac{r^M}{n(n-1)} \left\{ g(V, W)g(U, F) - g(U, W)g(V, F) \right\} \end{aligned}$$

where

$$r^M = \hat{r} + r^G \circ \pi - \|A\|^2 - \|T\|^2.$$

Proof. Let's prove the 2^{nd} equation of this theorem. Taking inner product C^* with V then we have

$$g(C^*(X, Y)Z, V) = g(R(X, Y)Z, V) - \frac{r^M}{n(n-1)} \{g(Y, Z)g(X, V) - g(X, Z)g(Y, V)\}.$$

Then using equation (1.4), we get

$$\begin{aligned} g(C^*(X, Y)Z, V) &= -g((\nabla_Z A)_X Y, V) - g(A_X T, T_V Z) \\ &\quad + g(A_Y Z, T_V X) - g(A_X Z, T_V Y). \end{aligned}$$

which completes the proof of the second equation. Other equations are similarly proved by using Theorem 1.2, Proposition 1.4 and Proposition 1.5. \square

Corollary 3.3. *Let $\pi : (M, g) \rightarrow (G, g')$ be a Riemannian submersion, where (M, g) and (G, g') Riemannian manifolds. Then the concircular curvature tensor of Riemannian submersion has no total umbilical fibres.*

4. Conharmonic curvature tensor along a Riemannian submersion

In this section, curvature relations of conharmonic curvature tensor in a Riemannian submersion are examined.

Definition 4.1. In the n -dimensional space V_n , the tensor

$$L^*(X, Y, Z, H) = R^M(X, Y, Z, H) - \frac{1}{n-2}[g(Y, Z)Ric(X, H) - g(X, Z)Ric(Y, H)],$$

is called conharmonic curvature tensor, where Ricci tensor denoted by Ric [10].

In a similar way, we have the following main theorem.

Theorem 4.2. Let (M, g) and (G, g') Riemannian manifolds,

$$\pi : (M, g) \rightarrow (G, g')$$

a Riemannian submersion and R^M , R^G and \hat{R} be Riemannian curvature tensors, S^M , S^G and \hat{S} be Ricci tensors of M , G and the fibre respectively. Then for any $U, V, W, F \in \chi^v(M)$ and $X, Y, Z, H \in \chi^h(M)$, we have the following relations

$$\begin{aligned} &g(L^*(X, Y)Z, H) \\ &= g(R^G(X, Y)Z, H) + 2g(A_X Y, A_Z H) - g(A_Y Z, A_X H) + g(A_X Z, A_Y H) \\ &- \frac{1}{(n-2)} \left\{ g(Y, Z) \left[S^G(X', H') \circ \pi + \frac{1}{2} (g(\nabla_X N, H) + g(\nabla_H N, X)) \right. \right. \\ &- 2 \sum_i g(A_X X_i, A_H X_i) - \sum_j g(T_{U_j} X, T_{U_j} H) \left. \right] \\ &- g(X, Z) \left[S^G(Y', H') \circ \pi + \frac{1}{2} (g(\nabla_Y N, H) + g(\nabla_H N, Y)) \right. \\ &- 2 \sum_i g(A_Y X_i, A_H X_i) - \sum_j g(T_{U_j} Y, T_{U_j} H) \left. \right] \\ &+ g(X, H) \left[S^G(Y', Z') \circ \pi + \frac{1}{2} (g(\nabla_Y N, Z) + g(\nabla_Z N, Y)) \right. \\ &- 2 \sum_i g(A_Y X_i, A_Z X_i) - \sum_j g(T_{U_j} Y, T_{U_j} Z) \left. \right] \\ &- g(Y, H) \left[S^G(X', Z') \circ \pi + \frac{1}{2} (g(\nabla_X N, Z) + g(\nabla_Z N, X)) \right. \\ &- 2 \sum_i g(A_X X_i, A_Z X_i) - \sum_j g(T_{U_j} X, T_{U_j} Z) \left. \right] \left. \right\}, \end{aligned}$$

$$\begin{aligned}
& g(L^*(X, Y)Z, V) \\
&= -g((\nabla_Z A)_X Y, V) - g(A_X T, T_V Z) + g(A_Y Z, T_V X) - g(A_X Z, T_V Y) \\
&- \frac{1}{(n-2)} \left\{ g(Y, Z) \left[g(\nabla_X N, V) - \sum_j g((\nabla_{U_j} T)_{U_j} X, V) \right. \right. \\
&+ \left. \left. \sum_i (g((\nabla_{X_i} A)_{X_i} X, V) - 2g(A_V X_i, T_X X_i)) \right] \right. \\
&- g(X, Z) \left[g(\nabla_Y N, V) - \sum_j g((\nabla_{U_j} T)_{U_j} Y, V) \right. \\
&+ \left. \left. \sum_i (g((\nabla_{X_i} A)_{X_i} Y, V) - 2g(A_V X_i, T_Y X_i)) \right] \right\},
\end{aligned}$$

$$\begin{aligned}
g(L^*(X, Y)V, W) &= g((\nabla_V A)_X Y, W) - g((\nabla_W A)_X Y, V) + g(A_X V, A_Y W) \\
&- g(A_X W, A_Y V) - g(T_V X, T_W Y) + g(T_W X, T_V Y),
\end{aligned}$$

$$\begin{aligned}
& g(L^*(X, V)Y, W) \\
&= g((\nabla_X T)_V W, Y) + g((\nabla_V A)_X Y, W) - g(T_V X, T_W Y) + g(A_X Y, A_Y W) \\
&- \frac{1}{(n-2)} \left\{ -g(V, W) \left[S^G(X', Y') \circ \pi + \frac{1}{2} (g(\nabla_X N, Y) + g(\nabla_Y N, X)) \right. \right. \\
&- 2 \sum_i g(A_X X_i, A_Y X_i) - \sum_j g(T_{U_j} X, T_{U_j} Y) \left. \right] \\
&- g(X, Y) \left[\hat{S}(V, W) - g(N, T_V W) + \sum_i (g((\nabla_{X_i} T)_V W, X_i) \right. \\
&+ \left. \left. g(A_{X_i} V, A_{X_i} W)) \right] \right\},
\end{aligned}$$

$$\begin{aligned}
g(L^*(U, V)W, X) &= g((\nabla_U T)_V W, X) - g((\nabla_V T)_U W, X) \\
&- \frac{1}{(n-2)} \left\{ g(V, W) \left[g(\nabla_U N, X) - \sum_j g((\nabla_{U_j} T)_{U_j} U, X) \right. \right. \\
&+ \sum_i \{g((\nabla_{X_i} A)_{X_i} X, U) - 2g(A_X X_i, T_U X_i)\} \left. \right] \\
&- g(U, W) \left[g(\nabla_V N, X) - \sum_j g((\nabla_{U_j} T)_{U_j} V, X) \right. \\
&+ \left. \left. \sum_i \{g((\nabla_{X_i} A)_{X_i} X, V) - 2g(A_X X_i, T_V X_i)\} \right] \right\}
\end{aligned}$$

and

$$\begin{aligned}
g(L^*(U, V)W, F) &= g(\hat{R}(U, V)W, F) + g(T_U W, T_V F) - g(T_V W, T_U F) \\
&\quad - \frac{1}{(n-2)} \left\{ g(V, W) \left[\hat{S}(U, F) - g(N, T_U F) \right. \right. \\
&\quad \left. \left. + \sum_i (g((\nabla_{X_i} T)_U F, X_i) + g(A_{X_i} U, A_{X_i} F)) \right] \right. \\
&\quad \left. - g(U, W) \left[\hat{S}(V, F) - g(N, T_V F) \right. \right. \\
&\quad \left. \left. + \sum_i (g((\nabla_{X_i} T)_V F, X_i) + g(A_{X_i} V, A_{X_i} F)) \right] \right. \\
&\quad \left. + g(F, U) \left[\hat{S}(U, V) - g(N, T_U V) \right. \right. \\
&\quad \left. \left. + \sum_i (g((\nabla_{X_i} T)_U V, X_i) + g(A_{X_i} U, A_{X_i} V)) \right] \right. \\
&\quad \left. - g(F, V) \left[\hat{S}(U, W) - g(N, T_U W) \right. \right. \\
&\quad \left. \left. + \sum_i (g((\nabla_{X_i} T)_U W, X_i) + g(A_{X_i} U, A_{X_i} W)) \right] \right\}.
\end{aligned}$$

Proof. Let's prove the 3th equation of this theorem. The following equations are obtained inner production with W to L^* and by using equation (1.5)

$$\begin{aligned}
g(L^*(X, Y)V, W) &= g(R^M(X, Y)V, W) - \frac{1}{n-2} \{ g(X, W)S(Y, V) \\
&\quad - g(Y, W)S(X, V) + g(Y, V)S(X, W) - g(X, V)S(Y, W) \}.
\end{aligned}$$

One can easily obtain the other equations by using Theorem 1.2 and Proposition 1.4. \square

Corollary 4.3. *Let $\pi : (M, g) \rightarrow (G, g')$ be a Riemannian submersion, where (M, g) and (G, g') Riemannian manifolds. If the Riemannian submersion has total umbilical fibres, that is $N = 0$, then the conharmonic curvature tensor*

is given by

$$\begin{aligned}
& g(L^*(X, Y)Z, H) \\
&= g(R^G(X, Y)Z, H) + 2g(A_X Y, A_Z H) - g(A_Y Z, A_X H) + g(A_X Z, A_Y H) \\
&- \frac{1}{(n-2)} \left\{ g(Y, Z) \left[S^G(X', H') \circ \pi - 2 \sum_i g(A_X X_i, A_H X_i) - \sum_j g(T_{U_j} X, T_{U_j} H) \right] \right. \\
&- g(X, Z) \left[S^G(Y', H') \circ \pi - 2 \sum_i g(A_Y X_i, A_H X_i) - \sum_j g(T_{U_j} Y, T_{U_j} H) \right] \\
&+ g(X, H) \left[S^G(Y', Z') \circ \pi - 2 \sum_i g(A_Y X_i, A_Z X_i) - \sum_j g(T_{U_j} Y, T_{U_j} Z) \right] \\
&\left. - g(Y, H) \left[S^G(X', Z') \circ \pi - 2 \sum_i g(A_X X_i, A_Z X_i) - \sum_j g(T_{U_j} X, T_{U_j} Z) \right] \right\},
\end{aligned}$$

$$\begin{aligned}
g(L^*(X, Y)Z, V) &= -g((\nabla_Z A)_X Y, V) - g(A_X T, T_V Z) + g(A_Y Z, T_V X) - g(A_X Z, T_V Y) \\
&- \frac{1}{(n-2)} \left\{ g(Y, Z) \left[- \sum_j g((\nabla_{U_j} T)_{U_j} X, V) \right. \right. \\
&+ \left. \sum_i (g((\nabla_{X_i} A)_{X_i} X, V) - 2g(A_V X_i, T_X X_i)) \right] \\
&- g(X, Z) \left[- \sum_j g((\nabla_{U_j} T)_{U_j} Y, V) \right. \\
&\left. \left. + \sum_i (g((\nabla_{X_i} A)_{X_i} Y, V) - 2g(A_V X_i, T_Y X_i)) \right] \right\},
\end{aligned}$$

$$\begin{aligned}
& g(L^*(X, V)Y, W) \\
&= g((\nabla_X T)_V W, Y) + g((\nabla_V A)_X Y, W) - g(T_V X, T_W Y) + g(A_X Y, A_Y W) \\
&- \frac{1}{(n-2)} \left\{ -g(V, W) \left[S^G(X', Y') \circ \pi - 2 \sum_i g(A_X X_i, A_Y X_i) - \sum_j g(T_{U_j} X, T_{U_j} Y) \right] \right. \\
&\left. - g(X, Y) \left[\hat{S}(V, W) + \sum_i (g((\nabla_{X_i} T)_V W, X_i) + g(A_{X_i} V, A_{X_i} W)) \right] \right\},
\end{aligned}$$

$$\begin{aligned}
g(L^*(U, V)W, X) &= g((\nabla_U T)_V W, X) - g((\nabla_V T)_U W, X) \\
&\quad - \frac{1}{(n-2)} \left\{ g(V, W) \left[\sum_j g(\nabla_{U_j} T)_{U_j} U, X \right] \right. \\
&\quad \left. + \sum_i \{ g((\nabla_{X_i} A)_{X_i} X, U) - 2g(A_X X_i, T_U X_i) \} \right] \\
&\quad - g(U, W) \left[\sum_j g(\nabla_{U_j} T)_{U_j} V, X \right] \\
&\quad \left. + \sum_i \{ g((\nabla_{X_i} A)_{X_i} X, V) - 2g(A_X X_i, T_V X_i) \} \right\}
\end{aligned}$$

and

$$\begin{aligned}
&g(L^*(U, V)W, F) \\
&= g(\hat{R}(U, V)W, F) + g(T_U W, T_V F) - g(T_V W, T_U F) \\
&\quad - \frac{1}{(n-2)} \left\{ g(V, W) \left[\hat{S}(U, F) \right. \right. \\
&\quad \left. \left. + \sum_i (g((\nabla_{X_i} T)_U F, X_i) + g(A_{X_i} U, A_{X_i} F)) \right] \right. \\
&\quad \left. - g(U, W) \left[\hat{S}(V, F) + \sum_i (g((\nabla_{X_i} T)_V F, X_i) + g(A_{X_i} V, A_{X_i} F)) \right] \right. \\
&\quad \left. + g(F, U) \left[\hat{S}(U, V) + \sum_i (g((\nabla_{X_i} T)_U V, X_i) + g(A_{X_i} U, A_{X_i} V)) \right] \right. \\
&\quad \left. - g(F, V) \left[\hat{S}(U, W) + \sum_i (g((\nabla_{X_i} T)_U W, X_i) + g(A_{X_i} U, A_{X_i} W)) \right] \right\}.
\end{aligned}$$

5. Conformal curvature tensor along a Riemannian submersion

In this section, we find some curvature relations of conformal curvature tensor in a Riemannian submersion and give a corollary in case of the Riemannian submersion has totally umbilical fibres.

Definition 5.1. In the n -dimensional space V_n , the tensor

$$\begin{aligned}
V^*(X, Y, Z, H) &= R^M(X, Y, Z, H) - \frac{1}{n-2} [g(X, H)Ric(Y, Z) - g(Y, H)Ric(X, Z) \\
&\quad + g(Y, Z)Ric(X, H) - g(X, Z)Ric(Y, H)] \\
&\quad + \frac{r^M}{(n-1)(n-2)} [g(X, H)g(Y, Z) - g(Y, H)g(X, Z)],
\end{aligned}$$

is called conformal curvature tensor, where Ricci tensor and scalar tensor denoted by Ric and r^M respectively [10].

Theorem 5.2. Let (M, g) and (G, g') Riemannian manifolds,

$$\pi : (M, g) \rightarrow (G, g')$$

a Riemannian submersion and R^M , R^G and \hat{R} be Riemannian curvature tensors, S^M , S^G and \hat{S} be Ricci tensors and r^M , r^G and \hat{r} be scalar curvature tensors of M , G and the fibre respectively. Then for any $U, V, W, F \in \chi^v(M)$ and $X, Y, Z, H \in \chi^h(M)$, we have the following relations

$$\begin{aligned}
& g(V^*(X, Y)Z, H) \\
&= g(R^G(X, Y)Z, H) + 2g(A_X Y, A_Z H) - g(A_Y Z, A_X H) + g(A_X Z, A_Y H) \\
&- \frac{1}{(n-2)} \left\{ g(X, H) \left[S^G(Y', Z') \circ \pi + \frac{1}{2} (g(\nabla_Y N, Z) + g(\nabla_Z N, Y)) \right. \right. \\
&- 2 \sum_i g(A_Y X_i, A_Z X_i) - \left. \sum_j g(T_{U_j} Y, T_{U_j} Z) \right] \\
&- g(Y, H) \left[S^G(X', Z') \circ \pi + \frac{1}{2} (g(\nabla_X N, Z) + g(\nabla_Z N, X)) \right. \\
&- 2 \sum_i g(A_X X_i, A_Z X_i) - \left. \sum_j g(T_{U_j} X, T_{U_j} Z) \right] \\
&+ g(Y, Z) \left[S^G(X', H') \circ \pi + \frac{1}{2} (g(\nabla_X N, H) + g(\nabla_H N, X)) \right. \\
&- 2 \sum_i g(A_X X_i, A_H X_i) - \left. \sum_j g(T_{U_j} X, T_{U_j} H) \right] \\
&- g(X, Z) \left[S^G(Y', H') \circ \pi + \frac{1}{2} (g(\nabla_Y N, H) + g(\nabla_H N, Y)) \right. \\
&- 2 \sum_i g(A_Y X_i, A_H X_i) - \left. \sum_j g(T_{U_j} Y, T_{U_j} H) \right] \left. \right\} \\
&+ \frac{r^M}{(n-1)(n-2)} \{g(Y, Z)g(X, H) - g(X, Z)g(Y, H)\},
\end{aligned}$$

$$\begin{aligned}
g(V^*(X, Y)Z, V) &= -g((\nabla_Z A)_X Y, V) - g(A_X T, T_V Z) + g(A_Y Z, T_V X) - g(A_X Z, T_V Y) \\
&- \frac{1}{(n-2)} \left\{ g(Y, Z) \left[g(\nabla_X N, V) - \sum_j g((\nabla_{U_j} T)_{U_j} X, V) \right. \right. \\
&+ \left. \sum_i (g((\nabla_{X_i} A)_{X_i} X, V) - 2g(A_V X_i, T_X X_i)) \right] \\
&- g(X, Z) \left[g(\nabla_Y N, V) - \sum_j g((\nabla_{U_j} T)_{U_j} Y, V) \right. \\
&+ \left. \left. \sum_i (g((\nabla_{X_i} A)_{X_i} Y, V) - 2g(A_V X_i, T_Y X_i)) \right] \right\},
\end{aligned}$$

$$g(V^*(X, Y)V, W) = g((\nabla_V A)_X Y, W) - g((\nabla_W A)_X Y, V) + g(A_X V, A_Y W) \\ - g(A_X W, A_Y V) - g(T_V X, T_W Y) + g(T_W X, T_V Y),$$

$$g(V^*(X, V)Y, W) \\ = g((\nabla_X T)_V W, Y) + g((\nabla_V A)_X Y, W) - g(T_V X, T_W Y) + g(A_X Y, A_Y W) \\ - \frac{1}{(n-2)} \left\{ -g(V, W) \left[S^G(X', Y') \circ \pi + \frac{1}{2} (g(\nabla_X N, Y) + g(\nabla_Y N, X)) \right] \right. \\ \left. - 2 \sum_i g(A_X X_i, A_Y X_i) - \sum_j g(T_{U_j} X, T_{U_j} Y) \right] \\ - g(X, Y) \left[\hat{S}(V, W) - g(N, T_V W) + \sum_i (g((\nabla_{X_i} T)_V W, X_i) \right. \\ \left. + g(A_{X_i} V, A_{X_i} W)) \right] \left. \right\},$$

$$g(V^*(U, V)W, X) = g((\nabla_U T)_V W, X) - g((\nabla_V T)_U W, X) \\ - \frac{1}{(n-2)} \left\{ g(V, W) \left[g(\nabla_U N, X) - \sum_j g(\nabla_{U_j} T)_{U_j} U, X \right] \right. \\ \left. + \sum_i \{g((\nabla_{X_i} A)_{X_i} X, U) - 2g(A_X X_i, T_U X_i)\} \right] \\ - g(U, W) \left[g(\nabla_V N, X) - \sum_j g(\nabla_{U_j} T)_{U_j} V, X \right] \\ \left. + \sum_i \{g((\nabla_{X_i} A)_{X_i} X, V) - 2g(A_X X_i, T_V X_i)\} \right] \left. \right\}$$

and

$$g(V^*(U, V)W, F) = g(\hat{R}(U, V)W, F) + g(T_U W, T_V F) - g(T_V W, T_U F) \\ - \frac{1}{(n-2)} \left\{ g(F, U) \left[\hat{S}(V, W) - g(N, T_V W) \right. \right. \\ \left. \left. + \sum_i (g(\nabla_{X_i} T)_V W, X_i + g(A_{X_i} V, A_{X_i} W)) \right] \right. \\ - g(F, V) \left[\hat{S}(U, W) - g(N, T_U W) + \sum_i (g((\nabla_{X_i} T)_U W, X_i) + g(A_{X_i} U, A_{X_i} W)) \right] \\ + g(V, W) \left[\hat{S}(U, F) - g(N, T_U F) + \sum_i (g((\nabla_{X_i} T)_U F, X_i) + g(A_{X_i} U, A_{X_i} F)) \right] \\ - g(U, W) \left[\hat{S}(V, F) - g(N, T_V F) + \sum_i (g((\nabla_{X_i} T)_V F, X_i) + g(A_{X_i} V, A_{X_i} F)) \right] \left. \right\} \\ + \frac{r^M}{(n-1)(n-2)} \{g(V, W)g(U, F) - g(U, W)g(V, F)\}$$

where

$$r^M = \hat{r} + r^G \circ \pi - \|N\|^2 - \|A\|^2 - \|T\|^2 + 2 \sum_i g(\nabla_{X_i} N, X_i).$$

Proof. Let's prove the 4th equation of this theorem. The following equations are obtained inner production with W to V^*

$$\begin{aligned} g(V^*(X, V)Y, W) &= g(R^M(X, V)Y, W) - \frac{1}{n-2} \{g(X, W)S^M(V, Y) \\ &\quad - g(V, W)S^M(X, Y) + g(V, Y)S(X, W) - g(X, Y)S^M(V, W)\} \\ &\quad + \frac{r^M}{(n-1)(n-2)} \{g(X, W)g(Y, V) - g(X, V)g(Y, W)\}. \end{aligned}$$

Then using equations (1.6)-(1.10), we have the desired result. From the Theorem 1.2, Proposition 1.4 and Proposition 1.5 the above equations are obtained. \square

Corollary 5.3. *Let $\pi : (M, g) \rightarrow (G, g')$ be a Riemannian submersion, where (M, g) and (G, g') Riemannian manifolds. If the Riemannian submersion has total umbilical fibres, that is $N = 0$, then the conformal curvature tensor is given by*

$$\begin{aligned} &g(V^*(X, Y)Z, H) \\ &= g(R^G(X, Y)Z, H) + 2g(A_X Y, A_Z H) - g(A_Y Z, A_X H) + g(A_X Z, A_Y H) \\ &\quad - \frac{1}{(n-2)} \left\{ g(X, H) \left[S^G(Y', Z') \circ \pi - 2 \sum_i g(A_Y X_i, A_Z X_i) - \sum_j g(T_{U_j} Y, T_{U_j} Z) \right] \right. \\ &\quad - g(Y, H) \left[S^G(X', Z') \circ \pi - 2 \sum_i g(A_X X_i, A_Z X_i) - \sum_j g(T_{U_j} X, T_{U_j} Z) \right] \\ &\quad + g(Y, Z) \left[S^G(X', H') \circ \pi - 2 \sum_i g(A_X X_i, A_H X_i) - \sum_j g(T_{U_j} X, T_{U_j} H) \right] \\ &\quad \left. - g(X, Z) \left[S^G(Y', H') \circ \pi - 2 \sum_i g(A_Y X_i, A_H X_i) - \sum_j g(T_{U_j} Y, T_{U_j} H) \right] \right\} \\ &\quad + \frac{r^M}{(n-1)(n-2)} \{g(Y, Z)g(X, H) - g(X, Z)g(Y, H)\}, \end{aligned}$$

$$\begin{aligned} &g(V^*(X, Y)Z, V) \\ &= -g((\nabla_Z A)_X Y, V) - g(A_X T, T_V Z) + g(A_Y Z, T_V X) - g(A_X Z, T_V Y) \\ &\quad - \frac{1}{(n-2)} \left\{ g(Y, Z) \left[- \sum_j g((\nabla_{U_j} T)_{U_j} X, V) + \sum_i (g((\nabla_{X_i} A)_{X_i} X, V) - 2g(A_V X_i, T_X X_i)) \right] \right. \\ &\quad \left. - g(X, Z) \left[- \sum_j g((\nabla_{U_j} T)_{U_j} Y, V) + \sum_i (g((\nabla_{X_i} A)_{X_i} Y, V) - 2g(A_V X_i, T_Y X_i)) \right] \right\}, \end{aligned}$$

$$\begin{aligned}
& g(V^*(X, V)Y, W) \\
&= g((\nabla_X T)_V W, Y) + g((\nabla_V A)_X Y, W) - g(T_V X, T_W Y) + g(A_X Y, A_Y W) \\
&- \frac{1}{(n-2)} \left\{ -g(V, W) \left[S^G(X', Y') \circ \pi - 2 \sum_i g(A_X X_i, A_Y X_i) - \sum_j g(T_{U_j} X, T_{U_j} Y) \right] \right. \\
&- \left. g(X, Y) \left[\hat{S}(V, W) + \sum_i (g((\nabla_{X_i} T)_V W, X_i) + g(A_{X_i} V, A_{X_i} W)) \right] \right\},
\end{aligned}$$

$$\begin{aligned}
g(V^*(U, V)W, X) &= g((\nabla_U T)_V W, X) - g((\nabla_V T)_U W, X) \\
&- \frac{1}{(n-2)} \left\{ g(V, W) \left[\sum_j g(\nabla_{U_j} T)_{U_j} U, X \right] \right. \\
&+ \left. \sum_i \{g((\nabla_{X_i} A)_{X_i} X, U) - 2g(A_X X_i, T_U X_i)\} \right] \\
&- g(U, W) \left[\sum_j g(\nabla_{U_j} T)_{U_j} V, X \right] \\
&+ \left. \sum_i \{g((\nabla_{X_i} A)_{X_i} X, V) - 2g(A_X X_i, T_V X_i)\} \right\}
\end{aligned}$$

and

$$\begin{aligned}
g(V^*(U, V)W, F) &= g(\hat{R}(U, V)W, F) + g(T_U W, T_V F) - g(T_V W, T_U F) \\
&- \frac{1}{(n-2)} \left\{ g(F, U) \left[\hat{S}(V, W) \right. \right. \\
&+ \left. \left. \sum_i (g(\nabla_{X_i} T)_V W, X_i + g(A_{X_i} V, A_{X_i} W)) \right] \right. \\
&- g(F, V) \left[\hat{S}(U, W) + \sum_i (g((\nabla_{X_i} T)_U W, X_i) + g(A_{X_i} U, A_{X_i} W)) \right] \\
&+ g(V, W) \left[\hat{S}(U, F) + \sum_i (g((\nabla_{X_i} T)_U F, X_i) + g(A_{X_i} U, A_{X_i} F)) \right] \\
&- \left. \left. g(U, W) \left[\hat{S}(V, F) + \sum_i (g((\nabla_{X_i} T)_V F, X_i) + g(A_{X_i} V, A_{X_i} F)) \right] \right\} \right. \\
&+ \frac{r^M}{(n-1)(n-2)} \{g(V, W)g(U, F) - g(U, W)g(V, F)\}
\end{aligned}$$

where

$$r^M = \hat{r} + r^G \circ \pi - \|A\|^2 - \|T\|^2.$$

Finally, we investigate the M -projective curvature tensor on a Riemannian submersion and give a corollary in case of the totally umbilical fibres.

6. M -projective curvature tensor along a Riemannian submersion

In this section, curvature relations of M -projective curvature tensor in a Riemannian submersion are examined and obtain a corollary using the curvature tensor.

Definition 6.1. Let take an n -dimensional differentiable manifold M^n with differentiability class C^∞ . In 1971 on a n -dimensional Riemannian manifold, ones [13] defined a tensor field W^* as

$$W^*(X, Y)Z = R^M(X, Y)Z - \frac{1}{2(n-1)}[S^M(Y, Z)X - S^M(X, Z)Y + g(Y, Z)QX - g(X, Z)QY]$$

tensor W^* as M -projective curvature tensor.

In addition, on an n -dimensional Riemannian manifold M^n the Ricci operator Q is defined by

$$S^M(X, Y) = g(QX, Y).$$

Theorem 6.2. Let (M, g) and (G, g') Riemannian manifolds,

$$\pi : (M, g) \rightarrow (G, g')$$

a Riemannian submersion and R^M , R^G and \hat{R} be Riemannian curvature tensors, S^M , S^G and \hat{S} be Ricci tensors of M , G and the fibre respectively. Then for any $U, V, W, F \in \chi^v(M)$ and $X, Y, Z, H \in \chi^h(M)$, we have the

following relations for M -projective curvature tensor:

$$\begin{aligned}
& g(W^*(X, Y)Z, H) \\
&= g(R^G(X, Y)Z, H) + 2g(A_X Y, A_Z H) - g(A_Y Z, A_X H) + g(A_X Z, A_Y H) \\
&- \frac{1}{2(n-1)} \left\{ g(X, H) \left[S^G(Y', Z') \circ \pi + \frac{1}{2} (g(\nabla_Y N, Z) + g(\nabla_Z N, Y)) \right. \right. \\
&- 2 \sum_i g(A_Y X_i, A_Z X_i) - \sum_j g(T_{U_j} Y, T_{U_j} Z) \left. \right] \\
&- g(Y, H) \left[S^G(X', Z') \circ \pi + \frac{1}{2} (g(\nabla_X N, Z) + g(\nabla_Z N, X)) \right. \\
&- 2 \sum_i g(A_X X_i, A_Z X_i) - \sum_j g(T_{U_j} X, T_{U_j} Z) \left. \right] \\
&+ g(Y, Z) \left[S^G(X', H') \circ \pi + \frac{1}{2} (g(\nabla_X N, H) + g(\nabla_H N, X)) \right. \\
&- 2 \sum_i g(A_X X_i, A_H X_i) - \sum_j g(T_{U_j} X, T_{U_j} H) \left. \right] \\
&- g(X, Z) \left[S^G(Y', H') \circ \pi + \frac{1}{2} (g(\nabla_Y N, H) + g(\nabla_H N, Y)) \right. \\
&- 2 \sum_i g(A_Y X_i, A_H X_i) - \sum_j g(T_{U_j} Y, T_{U_j} H) \left. \right] \left. \right\},
\end{aligned}$$

$$\begin{aligned}
g(W^*(X, Y)Z, V) &= -g((\nabla_Z A)_X Y, V) - g(A_X Y, T_V Z) + g(A_Y Z, T_V X) - g(A_X Z, T_V Y) \\
&- \frac{1}{2(n-1)} \left\{ g(Y, Z) \left[g(\nabla_X N, V) - \sum_j g((\nabla_{U_j} T)_{U_j} X, V) \right. \right. \\
&+ \sum_i (g((\nabla_{X_i} A)_{X_i} X, V) - 2g(A_V X_i, T_X X_i)) \left. \right] \\
&- g(X, Z) \left[g(\nabla_Y N, V) - \sum_j g((\nabla_{U_j} T)_{U_j} Y, V) \right. \\
&+ \sum_i (g((\nabla_{X_i} A)_{X_i} Y, V) - 2g(A_V X_i, T_Y X_i)) \left. \right] \left. \right\},
\end{aligned}$$

$$\begin{aligned}
g(W^*(X, Y)V, W) &= g((\nabla_V A)_X Y, W) - g((\nabla_W A)_X Y, V) + g(A_X V, A_Y W) \\
&- g(A_X W, A_Y V) - g(T_V X, T_W Y) + g(T_W X, T_V Y),
\end{aligned}$$

$$\begin{aligned}
g(W^*(X, V)Y, W) &= g((\nabla_X T)_V W, Y) + g((\nabla_V A)_X Y, W) - g(T_V X, T_W Y) + g(A_X Y, A_Y W) \\
&\quad - \frac{1}{2(n-1)} \left\{ -g(V, W) \left[S^G(X', Y') \circ \pi + \frac{1}{2} (g(\nabla_X N, Y) + g(\nabla_Y N, X)) \right] \right. \\
&\quad - 2 \sum_i g(A_X X_i, A_Y X_i) - \sum_j g(T_{U_j} X, T_{U_j} Y) \left. \right] - g(X, Y) \left[\hat{S}(V, W) - g(N, T_V W) \right. \\
&\quad \left. + \sum_i (g((\nabla_{X_i} T)_V W, X_i) + g(A_{X_i} V, A_{X_i} W)) \right] \left. \right\},
\end{aligned}$$

$$\begin{aligned}
g(W^*(U, V)W, X) &= g((\nabla_U T)_V W, X) - g((\nabla_V T)_U W, X) \\
&\quad - \frac{1}{2(n-1)} \left\{ g(V, W) \left[g(\nabla_U N, X) - \sum_j g(\nabla_{U_j} T)_{U_j} U, X \right] \right. \\
&\quad \left. + \sum_i \{g((\nabla_{X_i} A)_{X_i} X, U) - 2g(A_X X_i, T_U X_i)\} \right] \\
&\quad - g(U, W) \left[g(\nabla_V N, X) - \sum_j g(\nabla_{U_j} T)_{U_j} V, X \right] \\
&\quad \left. + \sum_i \{g((\nabla_{X_i} A)_{X_i} X, V) - 2g(A_X X_i, T_V X_i)\} \right] \left. \right\}
\end{aligned}$$

and

$$\begin{aligned}
g(W^*(U, V)W, F) &= g(\hat{R}(U, V)W, F) + g(T_U W, T_V F) - g(T_V W, T_U F) \\
&\quad - \frac{1}{2(n-1)} \left\{ g(F, U) \left[\hat{S}(V, W) - g(N, T_V W) \right. \right. \\
&\quad \left. \left. + \sum_i (g(\nabla_{X_i} T)_V W, X_i + g(A_{X_i} V, A_{X_i} W)) \right] \right. \\
&\quad - g(F, V) \left[\hat{S}(U, W) - g(N, T_U W) + \sum_i (g((\nabla_{X_i} T)_U W, X_i) + g(A_{X_i} U, A_{X_i} W)) \right] \\
&\quad + g(V, W) \left[\hat{S}(U, F) - g(N, T_U F) + \sum_i (g((\nabla_{X_i} T)_U F, X_i) + g(A_{X_i} U, A_{X_i} F)) \right] \\
&\quad \left. \left. - g(U, W) \left[\hat{S}(V, F) - g(N, T_V F) + \sum_i (g((\nabla_{X_i} T)_V F, X_i) + g(A_{X_i} V, A_{X_i} F)) \right] \right] \right\}.
\end{aligned}$$

Proof. Let's prove the 6th equation of this theorem. The following equations are obtained inner production with F to W^* and using (1.8) and (1.9) equations.

$$\begin{aligned}
g(W^*(U, V)W, F) &= g(R^M(U, V)W, F) - \frac{1}{2(n-1)} \left\{ g(F, U) S^M(U, V) \right. \\
&\quad \left. - g(F, V) S^M(U, W) + g(V, W) S^M(U, F) - g(U, W) S^M(V, F) \right\},
\end{aligned}$$

$$g(R^M(U, V)W, F) = g(\hat{R}(U, V)W, F) + g(T_U W, T_V F) - g(T_V W, T_U F)$$

and

$$S^M(U, V) = \hat{S}(U, V) - g(N, T_U V) + \sum_i \{g((\nabla_{X_i} T)_U V, X_i) + g(A_{X_i} U, A_{X_i} V)\}.$$

When these equations are substituted in W^* , the given result is obtained. Other equations are similarly proved by using Theorem 1.2 and Proposition 1.4. \square

Corollary 6.3. *Let $\pi : (M, g) \rightarrow (G, g')$ be a Riemannian submersion, where (M, g) and (G, g') Riemannian manifolds. If the Riemannian submersion has total umbilical fibres, that is $N = 0$, then the M -projective curvature tensor is given by*

$$\begin{aligned} &g(W^*(X, Y)Z, H) \\ &= g(R^G(X, Y)Z, H) + 2g(A_X Y, A_Z H) - g(A_Y Z, A_X H) + g(A_X Z, A_Y H) \\ &- \frac{1}{2(n-1)} \left\{ g(X, H) \left[S^G(Y', Z') \circ \pi - 2 \sum_i g(A_Y X_i, A_Z X_i) - \sum_j g(T_{U_j} Y, T_{U_j} Z) \right] \right. \\ &- g(Y, H) \left[S^G(X', Z') \circ \pi - 2 \sum_i g(A_X X_i, A_Z X_i) - \sum_j g(T_{U_j} X, T_{U_j} Z) \right] \\ &+ g(Y, Z) \left[S^G(X', H') \circ \pi - 2 \sum_i g(A_X X_i, A_H X_i) - \sum_j g(T_{U_j} X, T_{U_j} H) \right] \\ &\left. - g(X, Z) \left[S^G(Y', H') \circ \pi - 2 \sum_i g(A_Y X_i, A_H X_i) - \sum_j g(T_{U_j} Y, T_{U_j} H) \right] \right\}, \end{aligned}$$

$$\begin{aligned} g(W^*(X, Y)Z, V) &= -g((\nabla_Z A)_X Y, V) - g(A_X Y, T_V Z) + g(A_Y Z, T_V X) - g(A_X Z, T_V Y) \\ &- \frac{1}{2(n-1)} \left\{ g(Y, Z) \left[- \sum_j g((\nabla_{U_j} T)_{U_j} X, V) \right. \right. \\ &\left. \left. + \sum_i (g((\nabla_{X_i} A)_{X_i} X, V) - 2g(A_V X_i, T_X X_i)) \right] \right. \\ &- g(X, Z) \left[- \sum_j g((\nabla_{U_j} T)_{U_j} Y, V) \right. \\ &\left. \left. + \sum_i (g((\nabla_{X_i} A)_{X_i} Y, V) - 2g(A_V X_i, T_Y X_i)) \right] \right\}, \end{aligned}$$

$$\begin{aligned}
g(W^*(X, V)Y, W) &= g((\nabla_X T)_V W, Y) + g((\nabla_V A)_X Y, W) - g(T_V X, T_W Y) + g(A_X Y, A_Y W) \\
&\quad - \frac{1}{2(n-1)} \left\{ -g(V, W) \left[S^G(X', Y') \circ \pi - 2 \sum_i g(A_X X_i, A_Y X_i) - \sum_j g(T_{U_j} X, T_{U_j} Y) \right] \right. \\
&\quad \left. - g(X, Y) \left[\hat{S}(V, W) + \sum_i (g((\nabla_{X_i} T)_V W, X_i) + g(A_{X_i} V, A_{X_i} W)) \right] \right\},
\end{aligned}$$

$$\begin{aligned}
g(W^*(U, V)W, X) &= g((\nabla_U T)_V W, X) - g((\nabla_V T)_U W, X) \\
&\quad - \frac{1}{2(n-1)} \left\{ g(V, W) \left[\sum_j g(\nabla_{U_j} T)_{U_j} U, X \right] \right. \\
&\quad \left. + \sum_i \{g((\nabla_{X_i} A)_{X_i} X, U) - 2g(A_X X_i, T_U X_i)\} \right] \\
&\quad - g(U, W) \left[\sum_j g(\nabla_{U_j} T)_{U_j} V, X \right] \\
&\quad \left. + \sum_i \{g((\nabla_{X_i} A)_{X_i} X, V) - 2g(A_X X_i, T_V X_i)\} \right\}
\end{aligned}$$

and

$$\begin{aligned}
&g(W^*(U, V)W, F) \\
&= g(\hat{R}(U, V)W, F) + g(T_U W, T_V F) - g(T_V W, T_U F) \\
&\quad - \frac{1}{2(n-1)} \left\{ g(F, U) \left[\hat{S}(V, W) + \sum_i (g(\nabla_{X_i} T)_V W, X_i + g(A_{X_i} V, A_{X_i} W)) \right] \right. \\
&\quad - g(F, V) \left[\hat{S}(U, W) + \sum_i (g(\nabla_{X_i} T)_U W, X_i) + g(A_{X_i} U, A_{X_i} W) \right] \\
&\quad + g(V, W) \left[\hat{S}(U, F) + \sum_i (g((\nabla_{X_i} T)_U F, X_i) + g(A_{X_i} U, A_{X_i} F)) \right] \\
&\quad \left. - g(U, W) \left[\hat{S}(V, F) + \sum_i (g((\nabla_{X_i} T)_V F, X_i) + g(A_{X_i} V, A_{X_i} F)) \right] \right\}.
\end{aligned}$$

Remark 6.4. The authors investigate new curvature tensors along Riemannian submersions and obtain some results by using totally umbilical fibres. Therefore, it will be worth examining new curvature tensors along Riemannian submersions.

Acknowledgement. Both authors would like to thank Prof. Dr. Gabriel Eduard Vilcu for some useful comments and questions that help to clarify some statements of the original manuscript.

References

- [1] D. E. Blair, Riemannian geometry of contact and symplectic manifolds, Progress in Mathematics, 203. Birkhuser Boston, Inc., Boston, MA, (2002).

- [2] Bourguignon, J. P. and Lawson, H. B., *Stability and isolation phenomena for Yang-Mills fields*, Commun. Math. Phys. 79, (1981), 189-230.
- [3] Bourguignon, J. P. and Lawson, H. B. *A mathematician's visit to Kaluza-Klein theory*, Rend. Sem. Mat. Univ. Politec. Torino, Special Issue (1989), 143-163.
- [4] B. Y. Chen, *Geometry of Submanifolds*, Marcel Dekker, Inc. New York (1973).
- [5] M. Falcitelli, S. Ianuş, A.M. Pastore, *Riemannian submersions and related topics*, World Scientific Publishing Co., Inc., River Edge, NJ, 2004.
- [6] Ghosh, T. Koufogiorgos and R. Sharma, Conformally flat contact metric manifolds, J. Geom., 70(2001), 66–76.
- [7] Ianus, S. and Visinescu, M., *Kaluza-Klein theory with scalar fields and generalized Hopf manifolds*, Class. Quantum Gravity 4, (1987), 1317-1325.
- [8] Ianus, S. and Visinescu, M., *Space-time compactification and Riemannian submersions*, In: Rassias, G.(ed.) The Mathematical Heritage of C. F. Gauss, (1991), 358-371, World Scientific, River Edge.
- [9] Mustafa, M. T., *Applications of harmonic morphisms to gravity*, J. Math. Phys., 41, (2000), 6918-6929.
- [10] R.S. Mishra, *H-Projective curvature tensor in Kahler manifold*, Indian J. Pure Appl. Math. (1970) Vol. 1, 336-340.
- [11] R. H. Ojha, A note on the M -projective curvature tensor, Indian J. Pure Appl. Math. 8, 12 (1975), 1531–1534.
- [12] B. O'Neill, *The Fundamental Equations of a Submersions*, Michigan Math. J., 13 (1966), 459–469.
- [13] G. P. Pokhariyal and R. S. Mishra, *Curvature tensor and their relativistic significance II*, Yokohama Mathematical Journal 19 (1971), 97–103.
- [14] S. Sasaki and Y. Hatakeyama, On differentiable manifolds with certain structures which are closely related to almost contact structure, Tohoku Math. J., 13 (1961), 281–294.
- [15] B. Şahin., *Riemannian submersions, Riemannian maps in Hermitian Geometry and their applications*, Elsevier, Academic Press, (2017).
- [16] Watson, B., *G, G'-Riemannian submersions and nonlinear gauge field equations of general relativity*, In: Rassias, T. (ed.) Global Analysis - Analysis on manifolds, dedicated M. Morse. Teubner-Texte Math., 57, (1983), 324-349, Teubner, Leipzig.
- [17] G. Hall, Einstein's geodesic postulate, projective relatedness and Weyl's projective tensor. Mathematical physics, 27–35, World Sci. Publ., Hackensack, NJ, 2018
- [18] Z. Ahsan, S.A. Siddiqui, Concircular curvature tensor and fluid spacetimes. Internat. J. Theoret. Phys. 48 (2009), no. 11, 3202-3212
- [19] Doric, M.; Petrovic-Torgasev, M.; Verstraelen, L. Conditions on the conharmonic curvature tensor of Kaehler hypersurfaces in complex space forms. Publ. Inst. Math. (Beograd) (N.S.) 44(58) (1988), 97–108.

Mehmet Akif Akyol
Bingol University
Faculty of Arts and Sciences,
Department of Mathematics
12000, Bingöl, Turkey
e-mail: mehmetakifakyol@bingol.edu.tr

Gülhan Ayar
Karaman Mehmet Bey University
Department of Mathematics
70000, Karaman, Turkey
e-mail: gulhanayar@kmu.edu.tr