

The mod k chromatic index of graphs is $O(k)^*$

Fábio Botler,[†] Lucas Colucci and Yoshiharu Kohayakawa[‡]

Abstract

Let $\chi'_k(G)$ denote the minimum number of colors needed to color the edges of a graph G in a way that the subgraph spanned by the edges of each color has all degrees congruent to 1 (mod k). Scott [*Discrete Math.* 175, 1-3 (1997), 289–291] proved that $\chi'_k(G) \leq 5k^2 \log k$, and thus settled a question of Pyber [*Sets, graphs and numbers* (1992), pp. 583–610], who had asked whether $\chi'_k(G)$ can be bounded solely as a function of k . We prove that $\chi'_k(G) = O(k)$, answering affirmatively a question of Scott.

Throughout this paper, unless stated otherwise, $k \geq 2$ denotes an integer. All graphs considered here are simple, and $e(G)$ denotes the number of edges in the graph G . A χ'_k -coloring of G is a coloring of the edges of G in which the subgraph spanned by the edges of each color has all degrees congruent to 1 (mod k), and we denote by $\chi'_k(G)$ the minimum number of colors in a χ'_k -coloring of G . Pyber [3] proved that $\chi'_2(G) \leq 4$ for every graph G and asked whether $\chi'_k(G)$ is bounded by some function of k only. Scott [4] proved that $\chi'_k(G) \leq 5k^2 \log k$ for every graph G , and in turn asked if $\chi'_k(G)$ is in fact bounded by a linear function of k . This would be best possible apart from the multiplicative constant, as $\chi'_k(K_{1,k}) = k$. In this paper, we answer Scott's question affirmatively.

We shall make use of the following two results.

Lemma 1 (Mader [2]). *If $k \geq 1$, G is a graph on n vertices, and $e(G) \geq 2kn$, then G contains a k -connected subgraph.*

Lemma 2 (Thomassen [5]). *If $k \geq 1$ and G is a $(12k - 7)$ -edge-connected graph with an even number of vertices, then G has a spanning subgraph in which each vertex has degree congruent to $k \pmod{2k}$.*

We say that a graph G is k -divisible if k divides the degree of every vertex of G . Lemma 2 thus guarantees the existence of a k -divisible spanning subgraph in G when G is $(12k - 7)$ -edge-connected.

Lemma 3. *If G is a graph on n vertices and does not contain a non-empty k -divisible subgraph, then $e(G) < 2(12k - 6)n$.*

Proof. Let G be a graph and suppose that $e(G) \geq 2(12k - 6)n$. Lemma 1 tells us that G contains a $(12k - 6)$ -connected subgraph H . If H has an odd number of vertices, let $H' = H - v$ for an arbitrary

*This research has been partially supported by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil - CAPES - Finance Code 001. F. Botler is supported by CNPq (423395/2018-1) and by FAPERJ (211.305/2019). Y. Kohayakawa is partially supported by CNPq (311412/2018-1, 423833/2018-9) and FAPESP (2018/04876-1). The research that led to this paper started at WoPOCA 2019, which was financed by FAPESP (2015/11937-9) and CNPq (425340/2016-3, 423833/2018-9). FAPERJ is the Rio de Janeiro Research Foundation. FAPESP is the São Paulo Research Foundation. CNPq is the National Council for Scientific and Technological Development of Brazil.

[†]Programa de Engenharia de Sistemas e Computação, COPPE, Universidade Federal do Rio de Janeiro, Brazil

[‡]Instituto de Matemática e Estatística, Universidade de São Paulo, Brazil

vertex $v \in V(H)$. Otherwise, let $H' = H$. Then H' has an even number of vertices and is $(12k - 7)$ -connected, and hence is $(12k - 7)$ -edge-connected also. Lemma 2 tells us that H' contains a non-empty k -divisible subgraph and therefore so does G . \square

Given an integer d , we say that a graph G is d -degenerate if there is an ordering v_1, \dots, v_n of its vertices for which the number of neighbors of v_i in $V_i = \{v_1, \dots, v_{i-1}\}$ is at most d . In this case, the neighbors of v_i in V_i are its *left neighbors* and the neighbors of v_i in $\{v_{i+1}, \dots, v_n\}$ are its *right neighbors*. An edge uv is a *left edge* of v if u is a left neighbor of v . A left edge uv of v is a *right edge* of u .

Lemma 4. *Let G be a d -degenerate graph. Then $\chi'_k(G) \leq 4d + 2k - 2$.*

Proof. Let $V(G) = \{v_1, \dots, v_n\}$ be an ordering of $V(G)$ as above. In what follows, we color G by coloring the right edges of v_i for each $i \in \{1, \dots, n-1\}$ in turn, so that at each step we have a χ'_k -coloring of the graph spanned by the right edges of v_1, \dots, v_i with the following properties: for each $1 \leq j \leq n$, each of the colored, left edges of v_j is colored with a distinct color, and the colors used on the right edges of v_j are distinct from the colors used on its left edges. We proceed by induction on i . Let $i \in \{1, \dots, n-1\}$, and suppose that, for every $j < i$, the right edges of v_j is colored as above. This implies that all the left edges of v_i are colored and no right edge of v_i is colored. In what follows, we color the set S of right edges of v_i while keeping the properties of the partial coloring.

Since G is d -degenerate, v_i has at most d left edges, and hence we have at least $3d + 2k - 2$ colors to use on its right edges. We partition these colors arbitrarily into sets A and B so that $|A| = d + k$ and $|B| \geq 2d + k - 2$. Let $j > i$ and suppose v_j is a (right) neighbor of v_i . Note that at most $d - 1$ left edges of v_j are colored. We say that a color c is *forbidden* at v_j if a left edge of v_j is colored with c , and we call the colors in A that are not forbidden at v_j *available* at v_j .

Let S^* be a maximal subset of S that can be colored with colors in A in a way that (a) each right edge $v_i v_j \in S^*$ is colored with a color available at v_j , and (b) the number of edges in S^* colored with any given color is congruent to 1 (mod k). Let $\bar{S} = S \setminus S^*$ be the set of the remaining edges in S . We claim that $|\bar{S}| < |A|$. Assume for a contradiction that $|\bar{S}| \geq |A|$. For each edge $e = v_i v_j \in \bar{S}$, let A_e be the set of colors available at v_j , and for each color $x \in A$, let \bar{S}_x be the set of edges e in \bar{S} for which $x \in A_e$. Note that $\sum_{e \in \bar{S}} |A_e| = \sum_{x \in A} |\bar{S}_x|$ and that $|A_e| \geq |A| - (d - 1) = d + k - d + 1 = k + 1$ for every $e \in \bar{S}$. Therefore

$$(k + 1)|A| \leq (k + 1)|\bar{S}| \leq \sum_{e \in \bar{S}} |A_e| = \sum_{x \in A} |\bar{S}_x| \leq |A| \max\{|\bar{S}_x| : x \in A\},$$

whence $\max\{|\bar{S}_x| : x \in A\} \geq k + 1$. Let $z \in A$ be such that $|\bar{S}_z| = \max\{|\bar{S}_x| : x \in A\}$. If some edge in S^* is colored with z , then we color k edges in \bar{S} with color z . If no edge in S^* is colored with z , then we color $k + 1$ edges in \bar{S} with color z . In both cases we obtain a contradiction to the maximality of S^* . This shows that, indeed, $|\bar{S}| < |A| = d + k$.

Finally, we color the edges of \bar{S} consecutively and with distinct colors in B . This is possible, since for each $v_i w \in \bar{S}$ there are at most $d - 1 + |\bar{S}| - 1 \leq 2d + k - 3 < |B|$ colors of B that are forbidden (the colors forbidden at w plus the colors of B used on previous edges of \bar{S}). \square

Our main result is a consequence of Lemmas 3 and 4.

Theorem 5. *For every graph G we have $\chi'_k(G) \leq 198k - 101$.*

Proof. Let H be a maximal subgraph of G for which $\deg_H(v) \equiv 1 \pmod{k}$ for every $v \in V(H)$, and let $G' = G \setminus E(H)$. The maximality of H implies that $V(G) \setminus V(H)$ is independent, and that every

vertex in $V(H)$ has at most $k - 1$ neighbors in $V(G) \setminus V(H)$. Moreover, $G'[V(H)]$ has no non-empty k -divisible subgraph. By Lemma 3, every $J \subseteq G'[V(H)]$ has less than $2(12k - 6)|V(J)|$ edges, and hence its minimum degree is less than $48k - 24$. This implies that every subgraph of G' has a vertex of degree at most $49k - 25$, and hence G' is $(49k - 25)$ -degenerate. Lemma 4 tells us that G' has a χ'_k -coloring with at most $198k - 102$ colors. We then color $E(H)$ with a new color, and the result follows. \square

Alon, Friedland and Kalai [1] proved that when k is a prime power, the bound $2(12k - 6)n$ in Lemma 3 can be replaced by $(k - 1)n + 1$, and conjectured that this holds for every positive integer k . This result, together with Lemma 4, implies that $\chi'_k(G) \leq 14k - 9$ for any G and any prime power k .

Our arguments can be tweaked to give slightly better multiplicative constants, but we do not think it is worth pursuing this because we think we would be far from the truth still. Although we are not able to offer any strong evidence, we conjecture the following.

Conjecture 6. *There is a constant C such that $\chi'_k(G) \leq k + C$ for every graph G .*

We know that C in the conjecture above has to be at least 2, because one can prove that the graph G obtained from a $K_{k,k}$ by adding a universal vertex satisfies $\chi'_k(G) = k + 2$.

References

- [1] ALON, N., FRIEDLAND, S., AND KALAI, G. Regular subgraphs of almost regular graphs. *J. Combin. Theory Ser. B* 37, 1 (1984), 79–91.
- [2] MADER, W. Existenz n -fach zusammenhängender Teilgraphen in Graphen genügend grosser Kantendichte. *Abh. Math. Sem. Univ. Hamburg* 37 (1972), 86–97.
- [3] PYBER, L. Covering the edges of a graph by In *Sets, graphs and numbers (Budapest, 1991)* (1992), vol. 60 of *Colloq. Math. Soc. János Bolyai*, North-Holland, Amsterdam, pp. 583–610.
- [4] SCOTT, A. D. On graph decompositions modulo k . *Discrete Math.* 175, 1-3 (1997), 289–291.
- [5] THOMASSEN, C. Graph factors modulo k . *J. Combin. Theory Ser. B* 106 (2014), 174–177.