

SIGMA FUNCTIONS AND LIE ALGEBRAS OF SCHRÖDINGER OPERATORS

V. M. BUCHSTABER, E. YU. BUNKOVA

ABSTRACT. In the work by V. M. Buchstaber and D. V. Leikin [1] for any $g > 0$ is defined a system of $2g$ multidimensional Schrödinger equations in magnetic fields with quadratic potentials. This systems are equivalent to systems of heat equations in nonholonomic frame. It is proved that such a system determines the sigma function of the universal hyperelliptic curve of genus g . A polynomial Lie algebra with $2g$ Schrödinger operators $Q_0, Q_2, \dots, Q_{4g-2}$ as generators is introduced.

In this work for any $g > 0$ we obtain explicit expressions for Q_0, Q_2, Q_4 , and recurrent formulas for Q_{2k} with $k > 2$ expressing this operators as elements of the polynomial Lie algebra using Lie brackets of the operators Q_0, Q_2 , and Q_4 .

As an application we obtain explicit expressions for the operators $Q_0, Q_2, \dots, Q_{4g-2}$ for $g = 1, 2, 3, 4$.

*Dedicated to the memory of remarkable mathematician
Viktor Zelikovich Enolskii (1945 - 2019),
who made a great contribution to the development and applications
of the theory of multidimensional sigma functions.*

Viktor Enolskii's contribution to mathematics, mathematical and theoretical physics, including his results on the theory and applications of multidimensional sigma functions, is described in a memorial article to appear in the Notices of AMS.

1. INTRODUCTION

The heat equation, equivalent to Schrödinger equation, plays a fundamental role in the theory of Abelian functions. For example, part I of the monograph [2] begins with characterizing the elliptic theta function as a periodic fundamental solution of the one-dimensional heat equation.

Directly from the construction of the function $\theta(z, \Omega)$, where $z \in \mathbb{C}^g$, and Ω is a symmetric $(g \times g)$ matrix, follows that this function satisfies the classical system of multidimensional heat equations. In this system the times are the $g(g+1)/2$ independent parameters of the manifold of symmetric $(g \times g)$ matrices Ω . For this system, the vector fields along the times form a holonomic frame.

For $g > 3$ the dimension of the manifold of symmetric $(g \times g)$ matrices that are period matrices of a non-singular Riemann surface of genus g is less than $g(g+1)/2$. Here arises the well-known Riemann-Schottky problem, which has been solved by T. Shiota (see [3]) based on a hypothesis by S. P. Novikov, in terms of the Kadomtsev–Petviashvili equation.

Key words and phrases. Schrödinger operators, polynomial Lie algebras, differentiation of abelian functions over parameters.

This work is supported by the Russian Science Foundation under grant 20-11-19998.

Within the framework of F. Klein's problem, for each non-singular Riemann surface V of genus g , an entire function on \mathbb{C}^g is constructed, namely, the multidimensional sigma function. Its logarithmic derivatives of order 2 and higher generate the whole field of meromorphic functions on the Jacobian of the curve V (see [4]). In its first construction, this multidimensional sigma function was introduced as a modification of the multidimensional theta function. The efficiency of its implementation is achieved through the specially selected choice of the curve model. This results in an entire function whose series expansion in the vector argument z in \mathbb{C}^g gives a series over the ring of polynomials in the parameters of the equation of the curve. By choosing a family of curves (hyperelliptic or a wider family of (n, s) -curves), we can represent theta functions of these curves as modified sigma functions of these curves. An effective description of such modifications gives a solution to the Riemann–Schottky problem that is special for the family of curves under consideration. But, in this case, we cannot use the classical system of heat equations to characterize the obtained theta functions, since the vector fields along the times corresponding to the variations of the periods can intersect the discriminants of the curves. This is again connected to the Riemann–Schottky problem.

In [1] was constructed a system of heat equations that characterizes the multidimensional sigma functions. In this system the vector fields along the times corresponding to the variations of the parameters of the equations of the curves, form a non-holonomic frame. The Schrödinger equations in magnetic fields arise.

Naturally arises the question of an effective description of this system of equations. This work is devoted to the solution of this question in the case of a family of hyperelliptic curves.

2. THE PROBLEM IN THE CASE OF A FAMILY OF HYPERELLIPTIC CURVES

We consider hyperelliptic curves of genus $g \in \mathbb{N}$ in the model

$$\mathcal{V}_\lambda = \{(x, y) \in \mathbb{C}^2 : y^2 = x^{2g+1} + \lambda_4 x^{2g-1} + \lambda_6 x^{2g-2} + \dots + \lambda_{4g} x + \lambda_{4g+2}\}. \quad (1)$$

The curve depends on the parameters $\lambda = (\lambda_4, \lambda_6, \dots, \lambda_{4g}, \lambda_{4g+2}) \in \mathbb{C}^{2g}$. Let $\mathcal{B} \subset \mathbb{C}^{2g}$ be the subspace of parameters such that the curve \mathcal{V}_λ is nonsingular for $\lambda \in \mathcal{B}$. Then we have $\mathcal{B} = \mathbb{C}^{2g} \setminus \Sigma$, where Σ is the discriminant hypersurface of the universal curve.

For a meromorphic function f in \mathbb{C}^g the vector $\omega \in \mathbb{C}^g$ is a period if $f(z + \omega) = f(z)$ for all $z \in \mathbb{C}^g$. If a meromorphic function f has $2g$ independent periods in \mathbb{C}^g , then f is called an *abelian function*. Therefore an abelian function is a meromorphic function on the complex torus $T^g = \mathbb{C}^g/\Gamma$, where Γ is the lattice formed by the periods.

For each $\lambda \in \mathcal{B}$ the set of periods of holomorphic differentials on the curve \mathcal{V}_λ generates a lattice Γ_λ of rank $2g$ in \mathbb{C}^g . A *hyperelliptic function of genus g* (see [5, 6, 7]) is a meromorphic function in $\mathbb{C}^g \times \mathcal{B}$, such that for each $\lambda \in \mathcal{B}$ it's restriction on $\mathbb{C}^g \times \lambda$ is an abelian function, where the torus T^g is the Jacobian $\mathcal{J}_\lambda = \mathbb{C}^g/\Gamma_\lambda$ of the curve \mathcal{V}_λ . We denote by \mathcal{F} the field of hyperelliptic functions of genus g . For the properties of this field, see [6, 7].

We denote the coordinates in \mathbb{C}^g by $z = (z_1, z_3, \dots, z_{2g-1})$. The indices of the coordinates $z = (z_1, z_3, \dots, z_{2g-1}) \in \mathbb{C}^g$ and of the parameters $\lambda = (\lambda_4, \lambda_6, \dots, \lambda_{4g}, \lambda_{4g+2}) \in \mathbb{C}^{2g}$ determine their weights: $\text{wt } z_k = -k$, $\text{wt } \lambda_k = k$. We denote by P the ring of polynomials in $\lambda \in \mathcal{B} \subset \mathbb{C}^{2g}$.

We consider polynomial Lie algebras [8] of vector fields tangent to the discriminant Σ in \mathbb{C}^{2g} . Their generators $\{L_0, L_2, L_4, \dots, L_{4g-2}\}$ are the vector fields

$$L_{2k} = \sum_{s=2}^{2g+1} v_{2k+2, 2s-2}(\lambda) \frac{\partial}{\partial \lambda_{2k}}, \quad \text{where } v_{2k+2, 2s-2}(\lambda) \in P.$$

At a point $\lambda \in \mathcal{B}$ this vector fields determine a $2g$ -dimensional nonholonomic frame. The structure of a Lie algebra in the P -module with the generators $1, L_0, L_2, L_4, \dots, L_{4g-2}$ is determined by the polynomial matrices $V(\lambda) = (v_{2i, 2j}(\lambda))$, where $i, j = 1, \dots, 2g$, and $C(\lambda) = \{c_{2i, 2j}^{2k}(\lambda)\}$, where $i, j, k = 0, \dots, 2g-1$, such that

$$[L_{2i}, L_{2j}] = \sum_{k=0}^{2g-1} c_{2i, 2j}^{2k}(\lambda) L_{2k}, \quad [L_{2i}, \lambda_{2q}] = v_{2i+2, 2q-2}(\lambda), \quad [\lambda_{2q}, \lambda_{2r}] = 0. \quad (2)$$

Here λ_q is the operator of multiplication by the function λ_q in P .

We can find the explicit expressions for the matrix $V(\lambda)$ in [9, §4.1] (see also [8] and [10, Lemma 3.1]). For convenience, we assume that $\lambda_s = 0$ for all $s \notin \{0, 4, 6, \dots, 4g, 4g+2\}$ and $\lambda_0 = 1$. For $k, m \in \{1, 2, \dots, 2g\}$, $k \leq m$, we set

$$v_{2k, 2m}(\lambda) = \sum_{s=0}^{k-1} 2(k+m-2s) \lambda_{2s} \lambda_{2(k+m-s)} - \frac{2k(2g-m+1)}{2g+1} \lambda_{2k} \lambda_{2m},$$

and for $k > m$ we set $v_{2k, 2m}(\lambda) = v_{2m, 2k}(\lambda)$.

The vector field L_0 is the Euler vector field, namely, since $\text{wt } \lambda_{2k} = 2k$, we have

$$[L_0, \lambda_{2k}] = 2k \lambda_{2k}, \quad [L_0, L_{2k}] = 2k L_{2k}. \quad (3)$$

This determines the weights of the vector fields L_k , namely $\text{wt } L_{2k} = 2k$. The structure of a Lie algebra described above gives in this case a graded polynomial Lie algebra [8] that we denote by \mathcal{L}_L . The structure polynomials $c_{2i, 2j}^{2s}(\lambda)$ are described in [1, Theorem 2.5].

There is the well-known Lie–Witt algebra W_{\geq} over the field of complex numbers \mathbb{C} generated by the operators l_{2i} , where $i = 0, 1, 2, \dots$, with the commutation relations

$$[l_{2i}, l_{2j}] = 2(j - i)l_{2(i+j)}.$$

With respect to the bracket $[\cdot, \cdot]$ the Lie–Witt algebra W_{\geq} is generated by the three operators l_0, l_2, l_4 . The graded polynomial Lie algebra \mathcal{L}_L over P is a deformation of the Lie–Witt algebra W_{\geq} . It is as well generated by only three operators L_0, L_2 , and L_4 . The relation holds (see Lemma 4.3):

$$[L_2, L_{2k}] = 2(k - 1)L_{2k+2} + \frac{4(2g - k)}{(2g + 1)}(\lambda_{2k+2}L_0 - \lambda_4L_{2k-2}).$$

Now we introduce the Schrödinger operators. We consider the space \mathbb{C}^{3g} with coordinates (z, λ) . We denote by $\mathcal{C}(z, \lambda)$ the ring of differentiable functions in z and λ . We set

$$Q_{2k} = L_{2k} - H_{2k}, \quad k = 0, 1, 2, \dots, 2g - 1, \quad (4)$$

where

$$H_{2k} = \frac{1}{2} \sum \left(\alpha_{a,b}^{(k)}(\lambda) \partial_a \partial_b + 2\beta_{a,b}^{(k)}(\lambda) z_a \partial_b + \gamma_{a,b}^{(k)}(\lambda) z_a z_b \right) + \delta^{(k)}(\lambda), \quad (5)$$

and the summation is over odd a, b from 1 to $2g - 1$. In [1] a solution to the following problem is given:

Problem 2.1. *Find the sufficient conditions on the data $\{\alpha^{(i)}(\lambda), \beta^{(i)}(\lambda), \gamma^{(i)}(\lambda), \delta^{(i)}(\lambda)\}$ for the operators (4) to give a representation of Lie algebra (2) in the ring of operators on $\mathcal{C}(z, \lambda)$.*

Definition 2.1. The system of equations for $\varphi = \varphi(z, \lambda)$

$$Q_{2k}\varphi = 0 \quad (6)$$

is called the *system of heat equations*. The operators Q_{2k} are called *Schrödinger operators*.

We use the theory of hyperelliptic Kleinian functions (see [6, 11, 12, 4], and [13] for elliptic functions). Take the coordinates (z, λ) in $\mathbb{C}^g \times \mathcal{B} \subset \mathbb{C}^{3g}$. Let $\sigma(z, \lambda)$ be the hyperelliptic sigma function (or elliptic sigma function in genus $g = 1$ case). We denote $\partial_k = \frac{\partial}{\partial z_k}$. Following [5, 7, 14], we use the notation

$$\zeta_k = \partial_k \ln \sigma(z, \lambda), \quad \wp_{k_1, \dots, k_n} = -\partial_{k_1} \cdots \partial_{k_n} \ln \sigma(z, \lambda), \quad (7)$$

where $n \geq 2$, $k_s \in \{1, 3, \dots, 2g - 1\}$. The functions \wp_{k_1, \dots, k_n} provide us with examples of hyperelliptic functions. The field \mathcal{F} is the field of fractions of the ring of polynomials \mathcal{P} generated by the functions \wp_{k_1, \dots, k_n} , where $n \geq 2$, $k_s \in \{1, 3, \dots, 2g - 1\}$.

As shown in [1], a system of heat equations (6) for operators Q_{2k} that give a solution to Problem 1.1 determines the hyperelliptic sigma function $\sigma(z, \lambda)$, which allows to construct the hyperelliptic Kleinian functions theory starting from such operators.

The construction of the operators Q_{2i} in [1] uses the condition [1, equation (1.3)] stating that the commutator of operators $[Q_{2i}, Q_{2j}]$ is determined by a formula over P with the same coefficients as the formula for $[L_{2i}, L_{2j}]$, namely, the polynomial algebra generated by the operators Q_{2i} with $i = 0, 1, \dots$ is yet another realization of Lie–Witt algebra W_{\geq} deformation. Therefore, for an effective description of polynomial Lie algebra \mathcal{L}_Q one needs to obtain explicit formulas for Q_0, Q_2 , and Q_4 . These formulas are the main result of this work. As an application, we give the explicit form for differential operators in the case of universal hyperelliptic curve of genus 4.

3. GENERATING FUNCTIONS FOR SCHRÖDINGER OPERATORS

In this section, in the case of the model (1) we present the explicit solution to Problem 2.1 from [1, Theorem 2.6]. We will shift the original notation to be consistent with the formulas introduced in this work.

We denote by $q(a(x), b(x))$ the quotient in the Euclidean division of the polynomial $a(x)$ by the polynomial $b(x)$ and by $r(a(x), b(x))$ the residue in the Euclidean division of the polynomial $a(x)$ by the polynomial $b(x)$. We set (see (1))

$$f(x) = x^{2g+1} + \sum_{k=0}^{2g-1} \lambda_{2(2g+1-k)} x^k$$

and define the generating functions for the operators L_{2k} and H_{2k} :

$$L(x) = x^{2g-1} \sum_{k=0}^{2g-1} x^{-k} L_{2k}, \quad H(x) = x^{2g-1} \sum_{k=0}^{2g-1} x^{-k} H_{2k}. \quad (8)$$

For $R_i(x) = x^{g-i+1} \partial_x q(f(x), x^{2g-2i+2})$ we set

$$\begin{aligned} h(x) &= \sum_{i=1}^g x^{g-i} \partial_{2i-1} + R_i(x) z_{2i-1}, \\ t(x) &= \frac{1}{2} \sum_{i=1}^g (g-i+1) z_{2i-1}^2 q(R_i(x), x^{g-i+2}) + \\ &\quad + \sum_{i=1}^{g-1} \sum_{j=i+1}^g (g-j+1) z_{2j-1} q(x^{g-i} \partial_{2i-1} + R_i(x) z_{2i-1}, x^{g-j+2}). \end{aligned}$$

Then we have the relation (see [1, Theorem 2.6]):

$$H(x) = r \left(-\frac{1}{4} f''(x) + 2f(x)t(x) + \frac{1}{2} h(x) \circ h(x), f'(x) \right), \quad (9)$$

where \circ denotes the composition of operators.

The function $Q(x) = L(x) - H(x)$ is the generating function for Schrödinger operators.

Lemma 3.1. *For Schrödinger operators, in (5) we have*

$$\begin{aligned} \alpha_{a,b}^{(k)}(\lambda) &= 1, \quad \text{if } a+b=2k, \quad \text{and } a, b \in 2\mathbb{N}+1, \\ \alpha_{a,b}^{(k)}(\lambda) &= 0, \quad \text{if } a+b \neq 2k, \quad \text{and } a, b \in 2\mathbb{N}+1, \\ \delta^{(k)}(\lambda) &= \left(-\frac{1}{4}(2g-k+1)(2g-k) + \frac{1}{2} \left(g + \left\lfloor \frac{k+1}{2} \right\rfloor - k \right) \left(g - \left\lfloor \frac{k+1}{2} \right\rfloor \right) \right) \lambda_{2k}. \end{aligned}$$

Proof. The coefficient at $\partial_a \partial_b$ in $H(x)$ comes from the summand $\frac{1}{2} h(x) \circ h(x)$, by expanding we get $\sum_{i=1}^g \sum_{j=1}^g x^{2g-i-j} \partial_{2i-1} \partial_{2j-1}$. Thus from (8) we get the expressions for $\alpha_{a,b}^{(k)}(\lambda)$.

The expression for $\delta^{(k)}(\lambda)$ is obtained from the summands $-\frac{1}{4} f''(x)$ and $\frac{1}{2} h(x) \circ h(x)$. The first gives $-\frac{1}{4}(2g-k+1)(2g-k) \lambda_{2k}$ and the second gives $\frac{1}{2} (g + \lfloor \frac{k+1}{2} \rfloor - k) (g - \lfloor \frac{k+1}{2} \rfloor) \lambda_{2k}$, which results in the answer. \square

Remark 3.2. *We will later show that for Schrödinger operators, in (5) we have*

$$\begin{aligned} \beta_{a,b}^{(k)}(\lambda) &\text{ is a linear function in } \lambda, \\ \gamma_{a,b}^{(k)}(\lambda) &\text{ is a quadratic function in } \lambda. \end{aligned}$$

See Lemma 4.2.

4. EXPLICIT FORM FOR SCHRÖDINGER OPERATORS

Recall that in our notation $\lambda_s = 0$ for all $s \notin \{0, 4, 6, \dots, 4g, 4g + 2\}$ and $\lambda_0 = 1$.

Theorem 4.1. *We have the explicit expressions:*

$$\begin{aligned}
 H_0 &= \sum_{s=1}^g (2s-1)z_{2s-1}\partial_{2s-1} - \frac{g(g+1)}{2}; \\
 H_2 &= \frac{1}{2}\partial_1^2 + \sum_{s=1}^{g-1} (2s-1)z_{2s-1}\partial_{2s+1} - \frac{4}{2g+1}\lambda_4 \sum_{s=1}^{g-1} (g-s)z_{2s+1}\partial_{2s-1} + \\
 &\quad + \sum_{s=1}^g \left(\frac{2s-1}{2}\lambda_{4s} - \frac{2(g-s+1)}{2g+1}\lambda_4\lambda_{4s-4} \right) z_{2s-1}^2; \\
 H_4 &= \partial_1\partial_3 + \sum_{s=1}^{g-2} (2s-1)z_{2s-1}\partial_{2s+3} + \lambda_4 \sum_{s=1}^{g-1} (2s-1)z_{2s+1}\partial_{2s+1} - \\
 &\quad - \frac{6}{2g+1}\lambda_6 \sum_{s=1}^{g-1} (g-s)z_{2s+1}\partial_{2s-1} + \\
 &\quad + \sum_{s=1}^g \left((2s-1)\lambda_{4s+2} - \frac{3(g-s+1)}{2g+1}\lambda_6\lambda_{4s-4} \right) z_{2s-1}^2 + \\
 &\quad + \sum_{s=1}^{g-1} (2s-1)\lambda_{4s+4}z_{2s-1}z_{2s+1} - \frac{g(g-1)}{2}\lambda_4.
 \end{aligned}$$

Proof. The explicit expressions for $\alpha_{a,b}^{(k)}(\lambda)$ and $\delta^{(k)}(\lambda)$ are obtained in Lemma 3.1. We see that for $k = 0, 1, 2$ they coincide with the ones given in the Theorem. Recall $\lambda_0 = 1$, $\lambda_2 = 0$.

In (9) the coefficient at $z_{2j-1}\partial_{2i-1}$ is equal to

$$r(2(g-j+1)f(x)q(x^{g-i}, x^{g-j+2}) + x^{g-i}R_j(x), f'(x)).$$

For H_0 we are interested in the coefficient at x^{2g-1} of this polynomial, for H_2 in the coefficient at x^{2g-2} and for H_4 in the coefficient at x^{2g-3} .

For $j < i + 1$ the polynomial is equal to

$$r(x^{g-i}R_j(x), f'(x)) = x^{g-i}R_j(x)$$

because the degree of the polynomial $R_j(x)$ in x is $g + j - 1$ and the degree of $f'(x)$ is $2g$. Therefore we get the coefficient $(2j-1)$ in $j = i$ for H_0 , in $j = i - 1$ for H_2 , and in $j = i - 2$ for H_4 , as well as the coefficient $(2j-3)\lambda_4$ in $j = i \geq 2$ for H_4 , all the other coefficients being zero.

For $j = i + 1$ the polynomial is equal to

$$\begin{aligned}
 r(x^{g-i}R_j(x), f'(x)) &= r(x^{2g-2i}\partial_x q(f(x), x^{2g-2i}), f'(x)) = \\
 &= r((2i+1)x^{2g}, f'(x)) + \sum_{k=2g-2i+1}^{2g-1} (k-2g+2i)\lambda_{2(2g+1-k)}x^{k-1} = \\
 &= -\frac{2i+1}{2g+1} \sum_{k=0}^{2g-1} k\lambda_{2(2g+1-k)}x^{k-1} + \sum_{k=2g-2i+1}^{2g-1} (k-2g+2i)\lambda_{2(2g+1-k)}x^{k-1}.
 \end{aligned}$$

Therefore the coefficient for H_0 is zero, the coefficient for H_2 is $-\frac{4}{2g+1}(g-i)\lambda_4$, the coefficient for H_4 is $-\frac{6}{2g+1}(g-i)\lambda_6$.

For $j \geq i+2$ the polynomial is equal to

$$\begin{aligned} r(2(g-j+1)x^{j-i-2}f(x) + x^{2g-i-j+1}\partial_x q(f(x), x^{2g-2j+2}), f'(x)) = \\ = r(f'(x)x^{j-i+1} + \sum_{k=0}^{2g-2j+2} (2g-2j+2-k)\lambda_{2(2g+1-k)}x^{k+j-i-2}, f'(x)) = \\ = \sum_{k=0}^{2g-2j+2} (2g-2j+2-k)\lambda_{2(2g+1-k)}x^{k+j-i-2}. \end{aligned}$$

As $j \geq i+2 \geq 3$, the coefficients for H_0 , H_2 and H_4 are zero.

In (9) the coefficient at z_{2i-1}^2 is equal to

$$r\left((g-i+1)f(x)q(R_i(x), x^{g-i+2}) + \frac{1}{2}R_i(x)^2, f'(x)\right). \quad (10)$$

For $i=1$ we note that $R_1(x) = x^g$ and the expression (10) takes the form

$$\frac{1}{2}r(x^{2g}, f'(x)) = -\frac{1}{2(2g+1)} \sum_{k=1}^{2g-1} k\lambda_{2(2g+1-k)}x^{k-1}.$$

In this expression the coefficient at x^{2g-1} is zero, the coefficients at x^{2g-2} and x^{2g-3} provide the corresponding coefficients for H_2 and H_4 at z_1^2 .

For $i > 1$ we have

$$\begin{aligned} r\left((g-i+1)f(x)q(R_i(x), x^{g-i+2}) + \frac{1}{2}R_i(x)^2, f'(x)\right) = \\ = r\left(\frac{1}{2}R_i(x)x^{-g+i-1}(2(g-i+1)x^{-1}f(x) + x^{g-i+1}R_i(x)) - \right. \\ \left. - (g-i+1)f(x)x^{-1}r(x^{-g+i-1}R_i(x), x), f'(x)\right) = \\ = r\left(\frac{1}{2}R_i(x)x^{-g+i-1}\left(f'(x) + x^{-1}\sum_{k=0}^{2g-2i+1} (2g-2i+2-k)\lambda_{2(2g+1-k)}x^k\right) - \right. \\ \left. - (g-i+1)f(x)x^{-1}r(x^{-g+i-1}R_i(x), x), f'(x)\right) = \\ = \frac{1}{2}R_i(x)x^{-g+i-1}\left(x^{-1}\sum_{k=0}^{2g-2i+1} (2g-2i+2-k)\lambda_{2(2g+1-k)}x^k\right) - \\ - (g-i+1)r(x^{-g+i-1}R_i(x), x)x^{-1}(f(x) - x^{2g+1}) - \\ - (g-i+1)r(x^{-g+i-1}R_i(x), x)r(x^{2g}, f'(x)) = \\ = \frac{1}{2}R_i(x)x^{-g+i-1}\left(x^{-1}\sum_{k=0}^{2g-2i+1} (2g-2i+2-k)\lambda_{2(2g+1-k)}x^k\right) - \\ - (g-i+1)\lambda_{4(i-1)}x^{-1}\left(\sum_{k=0}^{2g-1} \lambda_{2(2g+1-k)}x^k\right) + \frac{(g-i+1)}{(2g+1)}\lambda_{4(i-1)}\left(\sum_{k=1}^{2g-1} k\lambda_{2(2g+1-k)}x^{k-1}\right). \end{aligned}$$

This is a polynomial of degree $2g-2$. Therefore the coefficient for H_0 is zero. The coefficients for H_2 and H_4 coincide with the corresponding coefficient given in the Theorem.

In (9) the coefficient at $z_{2i-1}z_{2j-1}$, $j > i$, is equal to

$$\begin{aligned} r(R_i(x)x^{-g+j-2}(2(g-j+1)f(x) + x^{g-j+2}R_j(x)), f'(x)) = \\ r\left(R_i(x)x^{-g+j-2}\left(xf'(x) + \sum_{k=0}^{2g-2j+1} (2g-2j+2-k)\lambda_{2(2g+1-k)}x^k\right), f'(x)\right) = \\ = R_i(x)x^{-g+j-2}\left(\sum_{k=0}^{2g-2j+1} (2g-2j+2-k)\lambda_{2(2g+1-k)}x^k\right). \end{aligned}$$

The degree of this polynomial is equal to $2g+i-j-2 \leq 2g-3$, and the equality holds for $j = i+1$. Therefore the coefficients for H_0 and H_2 are zero, and the coefficient for H_4 for $j = i+1$ is $(2i-1)\lambda_{4i+4}$. \square

Lemma 4.2. *For Schrödinger operators, in (5) we have*

$$\begin{aligned} \beta_{a,b}^{(k)}(\lambda) \text{ is a linear function in } \lambda, \\ \gamma_{a,b}^{(k)}(\lambda) \text{ is a quadratic function in } \lambda. \end{aligned}$$

The proof follows from the explicit expressions in the proof of Theorem 4.1.

Lemma 4.3. *The relation holds:*

$$[L_2, L_{2k}] = 2(k-1)L_{2k+2} + \frac{4(2g-k)}{(2g+1)}(\lambda_{2k+2}L_0 - \lambda_4L_{2k-2}). \quad (11)$$

Proof. We have

$$\begin{aligned} L_{2k} &= 2 \sum_{m=2}^{k+1} \left(\sum_{s=0}^{m-2} (k+m-2s)\lambda_{2s}\lambda_{2(k+m-s)} - \frac{(m-1)(2g-k)}{2g+1}\lambda_{2(k+1)}\lambda_{2(m-1)} \right) \frac{\partial}{\partial \lambda_{2m}} + \\ &+ 2 \sum_{m=k+2}^{2g+1} \left(\sum_{s=0}^k (k+m-2s)\lambda_{2s}\lambda_{2(k+m-s)} - \frac{(k+1)(2g-m+2)}{2g+1}\lambda_{2(k+1)}\lambda_{2(m-1)} \right) \frac{\partial}{\partial \lambda_{2m}}; \\ L_2 &= \sum_{m=2}^{2g+1} \left(2(m+1)\lambda_{2(m+1)} - \frac{4(2g-m+2)}{2g+1}\lambda_4\lambda_{2(m-1)} \right) \frac{\partial}{\partial \lambda_{2m}}. \end{aligned}$$

Now we obtain (11) by a direct calculation of the coefficients. \square

Corollary 4.4. *For $k = 3, 4, 5, \dots, 2g-1$ the formula holds*

$$Q_{2k} = \frac{1}{2(k-2)}[Q_2, Q_{2k-2}] - \frac{2(2g-k+1)}{(k-2)(2g+1)}(\lambda_{2k}Q_0 - \lambda_4Q_{2k-4}). \quad (12)$$

This determines the operators Q_{2k} for $k = 3, 4, 5, \dots, 2g-1$ recurrently and explicitly.

Proof. The operators Q_{2k} by construction (see Section 3) give a solution to Problem 2.1. The expression (12) is equivalent to (11) rewritten in Q_{2k} . \square

5. EXPLICIT FORMULAS FOR SCHRÖDINGER OPERATORS IN NONHOLONOMIC FRAME

The following operators can be found in [1, 5, 15, 16] for genus $g = 1$, in [5, 15, 16] for genus $g = 2$, in [10, 17] for genus $g = 3$. The genus $g = 4$ case is new. It follows from the formulas given in Section 4.

5.1. Schrödinger operators for genus $g = 1$.

In this case, the explicit formulas for $\{H_{2k}\}$ in (4) are

$$H_0 = z_1 \partial_1 - 1; \quad H_2 = \frac{1}{2} \partial_1^2 - \frac{1}{6} \lambda_4 z_1^2.$$

5.2. Schrödinger operators for genus $g = 2$.

In this case, the explicit formulas for $\{H_{2k}\}$ in (4) are

$$\begin{aligned} H_0 &= z_1 \partial_1 + 3z_3 \partial_3 - 3; \\ H_2 &= \frac{1}{2} \partial_1^2 - \frac{4}{5} \lambda_4 z_3 \partial_1 + z_1 \partial_3 - \frac{3}{10} \lambda_4 z_1^2 + \left(\frac{3}{2} \lambda_8 - \frac{2}{5} \lambda_4^2 \right) z_3^2; \\ H_4 &= \partial_1 \partial_3 - \frac{6}{5} \lambda_6 z_3 \partial_1 + \lambda_4 z_3 \partial_3 - \frac{1}{5} \lambda_6 z_1^2 + \lambda_8 z_1 z_3 + \left(3\lambda_{10} - \frac{3}{5} \lambda_4 \lambda_6 \right) z_3^2 - \lambda_4; \\ H_6 &= \frac{1}{2} \partial_3^2 - \frac{3}{5} \lambda_8 z_3 \partial_1 - \frac{1}{10} \lambda_8 z_1^2 + 2\lambda_{10} z_1 z_3 - \frac{3}{10} \lambda_4 \lambda_8 z_3^2 - \frac{1}{2} \lambda_6. \end{aligned}$$

5.3. Schrödinger operators for genus $g = 3$.

In this case, the explicit formulas for $\{H_{2k}\}$ in (4) are

$$\begin{aligned} H_0 &= z_1 \partial_1 + 3z_3 \partial_3 + 5z_5 \partial_5 - 6; \\ H_2 &= \frac{1}{2} \partial_1^2 - \frac{8}{7} \lambda_4 z_3 \partial_1 + \left(z_1 - \frac{4}{7} \lambda_4 z_5 \right) \partial_3 + 3z_3 \partial_5 + \\ &\quad - \frac{5}{14} \lambda_4 z_1^2 + \left(\frac{3}{2} \lambda_8 - \frac{4}{7} \lambda_4^2 \right) z_3^2 + \left(\frac{5}{2} \lambda_{12} - \frac{2}{7} \lambda_4 \lambda_8 \right) z_5^2; \\ H_4 &= \partial_1 \partial_3 - \frac{12}{7} \lambda_6 z_3 \partial_1 + \left(\lambda_4 z_3 - \frac{6}{7} \lambda_6 z_5 \right) \partial_3 + (z_1 + 3\lambda_4 z_5) \partial_5 - \frac{2}{7} \lambda_6 z_1^2 + \\ &\quad + \lambda_8 z_1 z_3 + \left(3\lambda_{10} - \frac{6}{7} \lambda_4 \lambda_6 \right) z_3^2 + 3\lambda_{12} z_3 z_5 + \left(5\lambda_{14} - \frac{3}{7} \lambda_6 \lambda_8 \right) z_5^2 - 3\lambda_4; \\ H_6 &= \frac{1}{2} \partial_3^2 + \partial_1 \partial_5 - \frac{9}{7} \lambda_8 z_3 \partial_1 - \frac{8}{7} \lambda_8 z_5 \partial_3 + (\lambda_4 z_3 + 2\lambda_6 z_5) \partial_5 - \frac{3}{14} \lambda_8 z_1^2 + 2\lambda_{10} z_1 z_3 + \\ &\quad + \left(\frac{9}{2} \lambda_{12} - \frac{9}{14} \lambda_4 \lambda_8 \right) z_3^2 + \lambda_{12} z_1 z_5 + 6\lambda_{14} z_3 z_5 + \left(\frac{3}{2} \lambda_4 \lambda_{12} - \frac{4}{7} \lambda_8^2 \right) z_5^2 - 2\lambda_6; \\ H_8 &= \partial_3 \partial_5 - \left(\frac{6}{7} \lambda_{10} z_3 - \lambda_{12} z_5 \right) \partial_1 - \frac{10}{7} \lambda_{10} z_5 \partial_3 + \lambda_8 z_5 \partial_5 - \frac{1}{7} \lambda_{10} z_1^2 + 3\lambda_{12} z_1 z_3 + \\ &\quad + \left(6\lambda_{14} - \frac{3}{7} \lambda_4 \lambda_{10} \right) z_3^2 + 2\lambda_{14} z_1 z_5 + \lambda_4 \lambda_{12} z_3 z_5 + \left(3\lambda_4 \lambda_{14} + \lambda_6 \lambda_{12} - \frac{5}{7} \lambda_8 \lambda_{10} \right) z_5^2 - \lambda_8; \\ H_{10} &= \frac{1}{2} \partial_5^2 - \left(\frac{3}{7} \lambda_{12} z_3 - 2\lambda_{14} z_5 \right) \partial_1 - \frac{5}{7} \lambda_{12} z_5 \partial_3 - \\ &\quad - \frac{1}{14} \lambda_{12} z_1^2 + 4\lambda_{14} z_1 z_3 - \frac{3}{14} \lambda_4 \lambda_{12} z_3^2 + 2\lambda_4 \lambda_{14} z_3 z_5 + \left(2\lambda_6 \lambda_{14} - \frac{5}{14} \lambda_8 \lambda_{12} \right) z_5^2 - \frac{1}{2} \lambda_{10}. \end{aligned}$$

5.4. Schrödinger operators for genus $g = 4$.

In this case, the explicit formulas for $\{H_{2k}\}$ in (4) are

$$\begin{aligned}
H_0 &= z_1 \partial_1 + 3z_3 \partial_3 + 5z_5 \partial_5 + 7z_7 \partial_7 - 10; \\
H_2 &= \frac{1}{2} \partial_1^2 + z_1 \partial_3 + 3z_3 \partial_5 + 5z_5 \partial_7 - \frac{4}{9} \lambda_4 (3z_3 \partial_1 + 2z_5 \partial_3 + z_7 \partial_5) - \\
&\quad - \frac{7}{18} \lambda_4 z_1^2 + \left(\frac{3}{2} \lambda_8 - \frac{2}{3} \lambda_4^2 \right) z_3^2 + \left(\frac{5}{2} \lambda_{12} - \frac{4}{9} \lambda_4 \lambda_8 \right) z_5^2 + \left(\frac{7}{2} \lambda_{16} - \frac{2}{9} \lambda_4 \lambda_{12} \right) z_7^2; \\
H_4 &= \partial_1 \partial_3 + z_1 \partial_5 + 3z_3 \partial_7 + \lambda_4 (z_3 \partial_3 + 3z_5 \partial_5 + 5z_7 \partial_7) - \frac{2}{3} \lambda_6 (3z_3 \partial_1 + 2z_5 \partial_3 + z_7 \partial_5) - \\
&\quad - \frac{1}{3} \lambda_6 z_1^2 + \lambda_8 z_1 z_3 + (3\lambda_{10} - \lambda_4 \lambda_6) z_3^2 + 3\lambda_{12} z_3 z_5 + \left(5\lambda_{14} - \frac{2}{3} \lambda_6 \lambda_8 \right) z_5^2 + \\
&\quad + 5\lambda_{16} z_5 z_7 + \left(7\lambda_{18} - \frac{1}{3} \lambda_6 \lambda_{12} \right) z_7^2 - 6\lambda_4; \\
H_6 &= \frac{1}{2} \partial_3^2 + \partial_1 \partial_5 - \frac{5}{3} \lambda_8 z_3 \partial_1 - \frac{16}{9} \lambda_8 z_5 \partial_3 + \\
&\quad + \left(\lambda_4 z_3 + 2\lambda_6 z_5 - \frac{8}{9} \lambda_8 z_7 \right) \partial_5 + (z_1 + 3\lambda_4 z_5 + 4\lambda_6 z_7) \partial_7 - \\
&\quad - \frac{5}{18} \lambda_8 z_1^2 + 2\lambda_{10} z_1 z_3 + \lambda_{12} z_1 z_5 - \left(\frac{5}{6} \lambda_4 \lambda_8 - \frac{9}{2} \lambda_{12} \right) z_3^2 + 6\lambda_{14} z_3 z_5 + 3\lambda_{16} z_3 z_7 + \\
&\quad + \left(\frac{3}{2} \lambda_4 \lambda_{12} - \frac{8}{9} \lambda_8^2 + \frac{15}{2} \lambda_{16} \right) z_5^2 + 10\lambda_{18} z_5 z_7 - \left(\frac{4}{9} \lambda_8 \lambda_{12} - \frac{5}{2} \lambda_4 \lambda_{16} \right) z_7^2 - \frac{9}{2} \lambda_6; \\
H_8 &= \partial_3 \partial_5 + \partial_1 \partial_7 - \left(\frac{4}{3} \lambda_{10} z_3 - \lambda_{12} z_5 \right) \partial_1 - \frac{20}{9} \lambda_{10} z_5 \partial_3 + \\
&\quad + \left(\lambda_8 z_5 - \frac{10}{9} \lambda_{10} z_7 \right) \partial_5 + (\lambda_4 z_3 + 2\lambda_6 z_5 + 3\lambda_8 z_7) \partial_7 - \\
&\quad - \frac{2}{9} \lambda_{10} z_1^2 + 3\lambda_{12} z_1 z_3 + 2\lambda_{14} z_1 z_5 + \lambda_{16} z_1 z_7 - \left(\frac{2}{3} \lambda_4 \lambda_{10} - 6\lambda_{14} \right) z_3^2 + \\
&\quad + (\lambda_4 \lambda_{12} + 9\lambda_{16}) z_3 z_5 + 6\lambda_{18} z_3 z_7 - \left(\frac{10}{9} \lambda_8 \lambda_{10} - \lambda_6 \lambda_{12} - 3\lambda_4 \lambda_{14} - 10\lambda_{18} \right) z_5^2 + \\
&\quad + 3\lambda_4 \lambda_{16} z_5 z_7 - \left(\frac{5}{9} \lambda_{10} \lambda_{12} - 2\lambda_6 \lambda_{16} - 5\lambda_4 \lambda_{18} \right) z_7^2 - 3\lambda_8; \\
H_{10} &= \frac{1}{2} \partial_5^2 + \partial_3 \partial_7 - (\lambda_{12} z_3 - 2\lambda_{14} z_5 - \lambda_{16} z_7) \partial_1 - \frac{5}{3} \lambda_{12} z_5 \partial_3 - \\
&\quad - \frac{4}{3} \lambda_{12} z_7 \partial_5 + (\lambda_8 z_5 + 2\lambda_{10} z_7) \partial_7 + \\
&\quad - \frac{1}{6} \lambda_{12} z_1^2 + 4\lambda_{14} z_1 z_3 + 3\lambda_{16} z_1 z_5 + 2\lambda_{18} z_1 z_7 - \left(\frac{1}{2} \lambda_4 \lambda_{12} - \frac{15}{2} \lambda_{16} \right) z_3^2 + \\
&\quad + (2\lambda_4 \lambda_{14} + 12\lambda_{18}) z_3 z_5 + \lambda_4 \lambda_{16} z_3 z_7 - \left(\frac{5}{6} \lambda_8 \lambda_{12} - 2\lambda_6 \lambda_{14} - \frac{9}{2} \lambda_4 \lambda_{16} \right) z_5^2 + \\
&\quad + (2\lambda_6 \lambda_{16} + 6\lambda_4 \lambda_{18}) z_5 z_7 - \left(\frac{2}{3} \lambda_{12}^2 - \frac{3}{2} \lambda_8 \lambda_{16} - 4\lambda_6 \lambda_{18} \right) z_7^2 - 2\lambda_{10};
\end{aligned}$$

$$\begin{aligned}
H_{12} &= \partial_5 \partial_7 - \left(\frac{2}{3} \lambda_{14} z_3 - 3 \lambda_{16} z_5 - 2 \lambda_{18} z_7 \right) \partial_1 - \left(\frac{10}{9} \lambda_{14} z_5 - \lambda_{16} z_7 \right) \partial_3 - \\
&\quad - \frac{14}{9} \lambda_{14} z_7 \partial_5 + \lambda_{12} z_7 \partial_7 - \\
&\quad - \frac{1}{9} \lambda_{14} z_1^2 + 5 \lambda_{16} z_1 z_3 + 4 \lambda_{18} z_1 z_5 - \left(\frac{1}{3} \lambda_4 \lambda_{14} - 9 \lambda_{18} \right) z_3^2 + \\
&\quad + 3 \lambda_4 \lambda_{16} z_3 z_5 + 2 \lambda_4 \lambda_{18} z_3 z_7 - \left(\frac{5}{9} \lambda_8 \lambda_{14} - 3 \lambda_6 \lambda_{16} - 6 \lambda_4 \lambda_{18} \right) z_5^2 + \\
&\quad + (\lambda_8 \lambda_{16} + 4 \lambda_6 \lambda_{18}) z_5 z_7 - \left(\frac{7}{9} \lambda_{12} \lambda_{14} - \lambda_{10} \lambda_{16} - 3 \lambda_8 \lambda_{18} \right) z_7^2 - \lambda_{12}; \\
H_{14} &= \frac{1}{2} \partial_7^2 - \left(\frac{1}{3} \lambda_{16} z_3 - 4 \lambda_{18} z_5 \right) \partial_1 - \left(\frac{5}{9} \lambda_{16} z_5 - 2 \lambda_{18} z_7 \right) \partial_3 - \frac{7}{9} \lambda_{16} z_7 \partial_5 - \\
&\quad - \frac{1}{18} \lambda_{16} z_1^2 + 6 \lambda_{18} z_1 z_3 - \frac{1}{6} \lambda_4 \lambda_{16} z_3^2 + 4 \lambda_4 \lambda_{18} z_3 z_5 - \left(\frac{5}{18} \lambda_8 \lambda_{16} - 4 \lambda_6 \lambda_{18} \right) z_5^2 + \\
&\quad + 2 \lambda_8 \lambda_{18} z_5 z_7 - \left(\frac{7}{18} \lambda_{12} \lambda_{16} - 2 \lambda_{10} \lambda_{18} \right) z_7^2 - \frac{1}{2} \lambda_{14}.
\end{aligned}$$

6. APPLICATION: DIFFERENTIATION OPERATORS FOR GENUS $g = 4$

Recall that \mathcal{F} is the field of hyperelliptic functions of genus g . In this section we consider the problem of constructing the Lie algebra of derivations of \mathcal{F} , i.e. to find $3g$ independent differential operators \mathcal{L} such that $\mathcal{L}\mathcal{F} \subset \mathcal{F}$. The exposition to the problem, as well as a general approach to the solution was developed in [15, 16]. An overview is given in [7]. In [18, 5, 14] an explicit solution to this problem has been obtained for $g = 1, 2, 3$. Here we give an explicit answer to this problem in the genus $g = 4$ case.

We introduce a ring of functions \mathcal{R}_φ . The generators of this graded ring over $\mathbb{Q}[\lambda]$ are the functions $\psi_{k_1 \dots k_n} = -\partial_{k_1} \dots \partial_{k_n} \ln \varphi$, where $n \geq 2$, $k_s \in \{1, 3, \dots, 2g - 1\}$, and $\text{wt } \psi_{k_1 \dots k_n} = k_1 + \dots + k_n$, $\text{wt } \lambda_k = k$. We introduce the operators:

$$\mathcal{L}_0 = L_0 - z_1 \partial_1 - 3z_3 \partial_3 - 5z_5 \partial_5 - 7z_7 \partial_7;$$

$$\mathcal{L}_2 = L_2 - \psi_1 \partial_1 + \frac{4}{3} \lambda_4 z_3 \partial_1 - \left(z_1 - \frac{8}{9} \lambda_4 z_5 \right) \partial_3 - \left(3z_3 - \frac{4}{9} \lambda_4 z_7 \right) \partial_5 - 5z_5 \partial_7;$$

$$\begin{aligned} \mathcal{L}_4 = L_4 - \psi_3 \partial_1 - \psi_1 \partial_3 - \\ + 2\lambda_6 z_3 \partial_1 - \left(\lambda_4 z_3 - \frac{4}{3} \lambda_6 z_5 \right) \partial_3 - \left(z_1 + 3\lambda_4 z_5 - \frac{2}{3} \lambda_6 z_7 \right) \partial_5 - (3z_3 + 5\lambda_4 z_7) \partial_7; \end{aligned}$$

$$\begin{aligned} \mathcal{L}_6 = L_6 - \psi_5 \partial_1 - \psi_3 \partial_3 - \psi_1 \partial_5 + \\ + \frac{5}{3} \lambda_8 z_3 \partial_1 + \frac{16}{9} \lambda_8 z_5 \partial_3 - \left(\lambda_4 z_3 + 2\lambda_6 z_5 - \frac{8}{9} \lambda_8 z_7 \right) \partial_5 - (z_1 + 3\lambda_4 z_5 + 4\lambda_6 z_7) \partial_7; \end{aligned}$$

$$\begin{aligned} \mathcal{L}_8 = L_8 - \psi_7 \partial_1 - \psi_5 \partial_3 - \psi_3 \partial_5 - \psi_1 \partial_7 + \left(\frac{4}{3} \lambda_{10} z_3 - \lambda_{12} z_5 \right) \partial_1 + \\ + \frac{20}{9} \lambda_{10} z_5 \partial_3 - \left(\lambda_8 z_5 - \frac{10}{9} \lambda_{10} z_7 \right) \partial_5 - (\lambda_4 z_3 + 2\lambda_6 z_5 + 3\lambda_8 z_7) \partial_7; \end{aligned}$$

$$\begin{aligned} \mathcal{L}_{10} = L_{10} - \psi_7 \partial_3 - \psi_5 \partial_5 - \psi_3 \partial_7 + \\ + (\lambda_{12} z_3 - 2\lambda_{14} z_5 - \lambda_{16} z_7) \partial_1 + \frac{5}{3} \lambda_{12} z_5 \partial_3 + \frac{4}{3} \lambda_{12} z_7 \partial_5 - (\lambda_8 z_5 + 2\lambda_{10} z_7) \partial_7; \end{aligned}$$

$$\begin{aligned} \mathcal{L}_{12} = L_{12} - \psi_7 \partial_5 - \psi_5 \partial_7 + \\ + \left(\frac{2}{3} \lambda_{14} z_3 - 3\lambda_{16} z_5 - 2\lambda_{18} z_7 \right) \partial_1 + \left(\frac{10}{9} \lambda_{14} z_5 - \lambda_{16} z_7 \right) \partial_3 + \frac{14}{9} \lambda_{14} z_7 \partial_5 - \lambda_{12} z_7 \partial_7; \end{aligned}$$

$$\mathcal{L}_{14} = L_{14} - \psi_7 \partial_7 + \left(\frac{1}{3} \lambda_{16} z_3 - 4\lambda_{18} z_5 \right) \partial_1 + \left(\frac{5}{9} \lambda_{16} z_5 - 2\lambda_{18} z_7 \right) \partial_3 + \frac{7}{9} \lambda_{16} z_7 \partial_5.$$

Denote the Lie algebra with this generators by $\mathcal{L}_\mathcal{L}$.

Theorem 6.1. *If the function φ satisfies the system of heat equations in a nonholonomic frame for genus 4, then the algebra $\mathcal{L}_\mathcal{L}$ is an algebra of derivations of the ring \mathcal{R}_φ .*

The proof follows the proof of Theorem 5.9 in [17].

Corollary 6.2. *For $g = 4$ the algebra $\mathcal{L}_\mathcal{L}$ for $\varphi = \sigma$ is an algebra of derivations of \mathcal{F} .*

We set

$$w_{0,1} = \psi_1, \quad w_{0,3} = 3\psi_3, \quad w_{2k,5} = 5\psi_5, \quad w_{0,7} = 7\psi_7,$$

$$w_{2,1} = \frac{1}{2}\psi_{111} + \psi_3 - \frac{7}{9}\lambda_4 z_1,$$

$$w_{2,3} = \frac{1}{2}\psi_{113} - \frac{4}{3}\lambda_4\psi_1 + 3\psi_5 + \left(3\lambda_8 - \frac{4}{3}\lambda_4^2\right) z_3,$$

$$w_{2,5} = \frac{1}{2}\psi_{115} - \frac{8}{9}\lambda_4\psi_3 + 5\psi_7 + \left(5\lambda_{12} - \frac{8}{9}\lambda_4\lambda_8\right) z_5,$$

$$w_{2,7} = \frac{1}{2}\psi_{117} - \frac{4}{9}\lambda_4\psi_5 + \left(7\lambda_{16} - \frac{4}{9}\lambda_4\lambda_{12}\right) z_7.$$

$$w_{4,1} = \psi_{113} + \psi_5 - \frac{2}{3}\lambda_6 z_1 + \lambda_8 z_3,$$

$$w_{4,3} = \psi_{133} - 2\lambda_6\psi_1 + \lambda_4\psi_3 + 3\psi_7 + \lambda_8 z_1 + (6\lambda_{10} - 2\lambda_4\lambda_6) z_3 + 3\lambda_{12} z_5,$$

$$w_{4,5} = \psi_{135} - \frac{4}{3}\lambda_6\psi_3 + 3\lambda_4\psi_5 + 3\lambda_{12} z_3 + \left(10\lambda_{14} - \frac{4}{3}\lambda_6\lambda_8\right) z_5 + 5\lambda_{16} z_7,$$

$$w_{4,7} = \psi_{137} - \frac{2}{3}\lambda_6\psi_5 + 5\lambda_4\psi_7 + 5\lambda_{16} z_5 + \left(14\lambda_{18} - \frac{2}{3}\lambda_6\lambda_{12}\right) z_7.$$

$$w_{6,1} = \frac{1}{2}\psi_{133} + \psi_{115} + \psi_7 - \frac{5}{9}\lambda_8 z_1 + 2\lambda_{10} z_3 + \lambda_{12} z_5;$$

$$w_{6,3} = \frac{1}{2}\psi_{333} + \psi_{135} - \frac{5}{3}\lambda_8\psi_1 + \lambda_4\psi_5 + 2\lambda_{10} z_1 - \left(\frac{5}{3}\lambda_4\lambda_8 - 9\lambda_{12}\right) z_3 + 6\lambda_{14} z_5 + 3\lambda_{16} z_7;$$

$$w_{6,5} = \frac{1}{2}\psi_{335} + \psi_{155} - \frac{16}{9}\lambda_8\psi_3 + 2\lambda_6\psi_5 + 3\lambda_4\psi_7 + \\ + \lambda_{12} z_1 + 6\lambda_{14} z_3 + \left(3\lambda_4\lambda_{12} - \frac{16}{9}\lambda_8^2 + 15\lambda_{16}\right) z_5 + 10\lambda_{18} z_7;$$

$$w_{6,7} = \frac{1}{2}\psi_{337} + \psi_{157} - \frac{8}{9}\lambda_8\psi_5 + 4\lambda_6\psi_7 + 3\lambda_{16} z_3 + 10\lambda_{18} z_5 - \left(\frac{8}{9}\lambda_8\lambda_{12} - 5\lambda_4\lambda_{16}\right) z_7.$$

$$w_{8,1} = \psi_{135} + \psi_{117} - \frac{4}{9}\lambda_{10} z_1 + 3\lambda_{12} z_3 + 2\lambda_{14} z_5 + \lambda_{16} z_7;$$

$$w_{8,3} = \psi_{335} + \psi_{137} - \frac{4}{3}\lambda_{10}\psi_1 + \lambda_4\psi_7 + \\ + 3\lambda_{12} z_1 - \left(\frac{4}{3}\lambda_4\lambda_{10} - 12\lambda_{14}\right) z_3 + (\lambda_4\lambda_{12} + 9\lambda_{16}) z_5 + 6\lambda_{18} z_7;$$

$$w_{8,5} = \psi_{355} + \psi_{157} + \lambda_{12}\psi_1 - \frac{20}{9}\lambda_{10}\psi_3 + \lambda_8\psi_5 + 2\lambda_6\psi_7 + 2\lambda_{14} z_1 + \\ + (\lambda_4\lambda_{12} + 9\lambda_{16}) z_3 - \left(\frac{20}{9}\lambda_8\lambda_{10} - 2\lambda_6\lambda_{12} - 6\lambda_4\lambda_{14} - 20\lambda_{18}\right) z_5 + 6\lambda_4\lambda_{16} z_7;$$

$$w_{8,7} = \psi_{357} + \psi_{177} - \frac{10}{9}\lambda_{10}\psi_5 + 3\lambda_8\psi_7 + \\ + \lambda_{16} z_1 + 6\lambda_{18} z_3 + 3\lambda_4\lambda_{16} z_5 - \left(\frac{10}{9}\lambda_{10}\lambda_{12} - 4\lambda_6\lambda_{16} - 10\lambda_4\lambda_{18}\right) z_7;$$

$$\begin{aligned}
w_{10,1} &= \frac{1}{2}\psi_{155} + \psi_{137} - \frac{1}{3}\lambda_{12}z_1 + 4\lambda_{14}z_3 + 3\lambda_{16}z_5 + 2\lambda_{18}z_7; \\
w_{10,3} &= \frac{1}{2}\psi_{355} + \psi_{337} - \lambda_{12}\psi_1 + \\
&\quad + 4\lambda_{14}z_1 - (\lambda_4\lambda_{12} - 15\lambda_{16})z_3 + (2\lambda_4\lambda_{14} + 12\lambda_{18})z_5 + \lambda_4\lambda_{16}z_7; \\
w_{10,5} &= \frac{1}{2}\psi_{555} + \psi_{357} + 2\lambda_{14}\psi_1 - \frac{5}{3}\lambda_{12}\psi_3 + \lambda_8\psi_7 + 3\lambda_{16}z_1 + \\
&\quad + (2\lambda_4\lambda_{14} + 12\lambda_{18})z_3 - \left(\frac{5}{3}\lambda_8\lambda_{12} - 4\lambda_6\lambda_{14} - 9\lambda_4\lambda_{16}\right)z_5 + (2\lambda_6\lambda_{16} + 6\lambda_4\lambda_{18})z_7; \\
w_{10,7} &= \frac{1}{2}\psi_{557} + \psi_{377} + \lambda_{16}\psi_1 - \frac{4}{3}\lambda_{12}\psi_5 + 2\lambda_{10}\psi_7 + \\
&\quad + 2\lambda_{18}z_1 + \lambda_4\lambda_{16}z_3 + (2\lambda_6\lambda_{16} + 6\lambda_4\lambda_{18})z_5 - \left(\frac{4}{3}\lambda_{12}^2 - 3\lambda_8\lambda_{16} - 8\lambda_6\lambda_{18}\right)z_7; \\
w_{12,1} &= \psi_{157} - \frac{2}{9}\lambda_{14}z_1 + 5\lambda_{16}z_3 + 4\lambda_{18}z_5; \\
w_{12,3} &= \psi_{357} - \frac{2}{3}\lambda_{14}\psi_1 + 5\lambda_{16}z_1 - \left(\frac{2}{3}\lambda_4\lambda_{14} - 18\lambda_{18}\right)z_3 + 3\lambda_4\lambda_{16}z_5 + 2\lambda_4\lambda_{18}z_7; \\
w_{12,5} &= \psi_{557} + 3\lambda_{16}\psi_1 - \frac{10}{9}\lambda_{14}\psi_3 + 4\lambda_{18}z_1 + \\
&\quad + 3\lambda_4\lambda_{16}z_3 - \left(\frac{10}{9}\lambda_8\lambda_{14} - 6\lambda_6\lambda_{16} - 12\lambda_4\lambda_{18}\right)z_5 + (\lambda_8\lambda_{16} + 4\lambda_6\lambda_{18})z_7; \\
w_{12,7} &= \psi_{577} + 2\lambda_{18}\psi_1 + \lambda_{16}\psi_3 - \frac{14}{9}\lambda_{14}\psi_5 + \lambda_{12}\psi_7 + \\
&\quad + 2\lambda_4\lambda_{18}z_3 + (\lambda_8\lambda_{16} + 4\lambda_6\lambda_{18})z_7 - \left(\frac{14}{9}\lambda_{12}\lambda_{14} - 2\lambda_{10}\lambda_{16} - 6\lambda_8\lambda_{18}\right)z_7; \\
w_{14,1} &= \frac{1}{2}\psi_{177} - \frac{1}{9}\lambda_{16}z_1 + 6\lambda_{18}z_3; \\
w_{14,3} &= \frac{1}{2}\psi_{377} - \frac{1}{3}\lambda_{16}\psi_1 + 6\lambda_{18}z_1 - \frac{1}{3}\lambda_4\lambda_{16}z_3 + 4\lambda_4\lambda_{18}z_5; \\
w_{14,5} &= \frac{1}{2}\psi_{577} + 4\lambda_{18}\psi_1 - \frac{5}{9}\lambda_{16}\psi_3 + 4\lambda_4\lambda_{18}z_3 - \left(\frac{5}{9}\lambda_8\lambda_{16} - 8\lambda_6\lambda_{18}\right)z_5 + 2\lambda_8\lambda_{18}z_7; \\
w_{14,7} &= \frac{1}{2}\psi_{777} + 2\lambda_{18}\psi_3 - \frac{7}{9}\lambda_{16}\psi_5 + 2\lambda_8\lambda_{18}z_5 - \left(\frac{7}{9}\lambda_{12}\lambda_{16} - 4\lambda_{10}\lambda_{18}\right)z_7.
\end{aligned}$$

We obtain a genus $g = 4$ analog of Theorems 4.1, 4.2 and 4.3 from [17].

Theorem 6.3. *For $g = 4$ a solution φ of the system of heat equations (6) gives a solution $(\psi_1, \psi_3, \psi_5, \psi_7) = (\partial_1 \ln \varphi, \partial_3 \ln \varphi, \partial_5 \ln \varphi, \partial_7 \ln \varphi)$ of the system of nonlinear differential equations that we call an analog of the Burgers equation for $g = 4$:*

$$\mathcal{L}_{2k}(\psi_1, \psi_3, \psi_5, \psi_7) = (w_{2k,1}, w_{2k,3}, w_{2k,5}, w_{2k,7}), \quad k = 0, 1, 2, 3, 4, 5, 6, 7. \quad (13)$$

THE PROOF is the result of direct computation.

REFERENCES

- [1] V. M. BUCHSTABER, D. V. LEYKIN, *Heat Equations in a Nonholonomic Frame*, *Funct. Anal. Appl.*, 38:2 (2004), 88–101.
- [2] D. MUMFORD, *Tata Lectures on Theta I, II*, *Progress in Mathematics*, Vol. 28, 43.
- [3] T. SHIOTA, *Characterization of Jacobian varieties in terms of soliton equations*, *Invent. Math.*, 83 (1986), 333–382.
- [4] V. M. BUCHSTABER, V. Z. ENOLSKII, D. V. LEIKIN, *Hyperelliptic Kleinian functions and applications*, “Solitons, Geometry and Topology: On the Crossroad”, *Adv. Math. Sci.*, AMS Transl., 179:2, Providence, RI, 1997, 1–34.
- [5] V. M. BUCHSTABER, *Polynomial dynamical systems and Korteweg–de Vries equation*, *Proc. Steklov Inst. Math.*, 294 (2016), 176–200.
- [6] V. M. BUCHSTABER, V. Z. ENOLSKII, D. V. LEIKIN, *Multi-Dimensional Sigma-Functions*, arXiv:1208.0990, 2012, 267 pp.
- [7] V. M. BUCHSTABER, V. Z. ENOLSKI, D. V. LEYKIN, *Sigma-functions: old and new results*, *Algebraic Geometry*, v. 2, LMS Lecture Note Series, Cambridge Univ. Press, 2019, arXiv: 1810.11079.
- [8] V. M. BUCHSTABER, D. V. LEIKIN, *Polynomial Lie Algebras*, *Funct. Anal. Appl.*, 36:4 (2002), 267–280.
- [9] V. I. ARNOLD, *Singularities of Caustics and Wave Fronts*, *Mathematics and its Applications*, vol. 62, Kluwer Academic Publisher Group, Dordrecht, 1990.
- [10] J. C. EILBECK, J. GIBBONS, Y. ONISHI, S. YASUDA, *Theory of Heat Equations for Sigma Functions*, arXiv:1711.08395, (2018).
- [11] H. F. BAKER, *On the hyperelliptic sigma functions*, *Amer. Journ. Math.* 20, 1898, 301–384.
- [12] V. M. BUCHSTABER, V. Z. ENOLSKII, D. V. LEIKIN, *Kleinian functions, hyperelliptic Jacobians and applications*, *Reviews in Mathematics and Math. Physics*, 10:2, Gordon and Breach, London, 1997, 3–120.
- [13] E. T. WHITTAKER, G. N. WATSON, *A Course of Modern Analysis*, Reprint of 4th (1927) ed., Vol 2. *Transcendental functions*, Cambridge Univ. Press, Cambridge, 1996.
- [14] E. YU. BUNKOVA, *Differentiation of genus 3 hyperelliptic functions*, *European Journal of Mathematics*, 4:1 (2018), 93–112, arXiv:1703.03947
- [15] V. M. BUCHSTABER, D. V. LEIKIN, *Differentiation of Abelian functions with respect to parameters*, *Russian Math. Surveys*, 62:4 (2007), 787–789.
- [16] V. M. BUCHSTABER, D. V. LEIKIN, *Solution of the Problem of Differentiation of Abelian Functions over Parameters for Families of (n, s) -Curves*, *Funct. Anal. Appl.*, 42:4, 2008, 268–278.
- [17] V. M. BUCHSTABER, E. YU. BUNKOVA, *Lie Algebras of Heat Operators in Nonholonomic Frame*, 2019, arXiv:1911.08266.
- [18] F. G. FROBENIUS, L. STICKELBERGER, *Über die Differentiation der elliptischen Functionen nach den Perioden und Invarianten*, *J. Reine Angew. Math.*, 92 (1882), 311–337.

STEKLOV MATHEMATICAL INSTITUTE OF RUSSIAN ACADEMY OF SCIENCES, MOSCOW, RUSSIA
E-mail address: buchstab@mi-ras.ru, bunkova@mi-ras.ru