p-Permutation Equivalences between Blocks of Group Algebras^{*}

Robert Boltje Department of Mathematics University of California Santa Cruz, CA 95064 U.S.A. boltje@ucsc.edu Philipp Perepelitsky Department of Mathematics University of California Santa Cruz, CA 95064 U.S.A. pperepel@ucsc.edu

July 17, 2020

Contents

1. Introduction 2. Subgroups of direct product groups 3. p-permutation modules 4. Block theoretic preliminaries 5. Brauer pairs for p-permutation modules 6. Extended tensor products and homomorphisms 7. Tensor products of p-permutation bimodules 8. Character groups and perfect isometries 9. Grothendieck groups of p-permutation modules and p-permutation equivalences 10. Brauer pairs of p-permutation equivalences 11. Fusion systems, local equivalences and finiteness 12. A character theoretic criterion and moving from left to right 13. Külshammer-Puig classes 14. The maximal module of a p-permutation equivalence 15. Connection with isotypies and splendid Rickard equivalences

Abstract

We extend the notion of a p-permutation equivalence between two p-blocks A and B of finite groups G and H, from the definition in [BX08] to a virtual p-permutation bimodule whose components have twisted diagonal vertices. It is shown that various invariants of A and B are preserved, including defect groups, fusion systems, and Külshammer-Puig classes. Moreover it is shown that p-permutation equivalences have additional surprising properties. They have only one constituent with maximal vertex and the set of ppermutation equivalences between A and B is finite (possibly empty). The paper uses new methods: a consequent use of module structures on subgroups of $G \times H$ arising from Brauer constructions which in general are not direct product subgroups, the necessary adaptation of the notion of tensor products between bimodules, and a general formula (stated in these new terms) for the Brauer construction of a tensor product of p-permutation bimodules.

1 Introduction

Let G and H be finite groups, \mathcal{O} a complete discrete valuation ring of characteristic 0 which contains a root of unity whose order is equal to the exponent of $G \times H$. We denote by K the field of fractions of \mathcal{O} , and by F its residue field whose characteristic we assume to be a prime p. Furthermore we assume that A is a block algebra of $\mathcal{O}G$ and B is a block algebra of $\mathcal{O}H$. Various authors have defined notions of equivalence between A and B (e.g. [Br90], [Br95], [R96], [P99], [BX08], [L09]). They are divided into two parts: those that are equivalences of categories (as for instance Morita equivalences, derived equivalences, splendid Rickard equivalences), and those that are isomorphisms between associated representation rings (as for instance perfect isometries and isotypies), preserving additional features on the representation ring level. The first attempt to define a strongest possible equivalence on a representation ring level goes back to [BX08], where a preliminary notion of a p-permutation equivalence was defined. [BX08] made the restrictive assumptions that the blocks A and B have a common

^{*}MR Subject Classification: 20C20, 19A22Keywords: *p*-permutation modules, trivial source modules, blocks of group algebras, fusion systems, perfect isometries, isotypies, splendid Rickard complexes.

defect group D, that certain fusion categories are equivalent, and some results on this notion were only proved under the hypothesis that D is abelian. This notion was soon after extended in [L09] to source algebras, see also [L18, Section 9.5].

In this paper we cast the net much wider and define a *p*-permutation equivalence as an element $\gamma \in T^{\Delta}(A, B)$, the representation group of finitely generated *p*-permutation (A, B)-bimodules whose indecomposable direct summands, when regarded as left $\mathcal{O}[G \times H]$ -modules, have twisted diagonal vertices, i.e., vertices of the form $\Delta(P, \phi, Q) := \{(\phi(y), y) \mid y \in Q\}$, the graph of an isomorphism $\phi : Q \xrightarrow{\sim} P$ between a *p*-subgroup Q of H and a *p*-subgroup P of G, with the property that

$$\gamma_{H}^{\cdot} \gamma^{\circ} = [A] \in T^{\Delta}(A, A) \quad \text{and} \quad \gamma^{\circ}_{G}^{\cdot} \gamma = [B] \in T^{\Delta}(B, B) ,$$

$$(1)$$

where $\gamma^{\circ} \in T^{\Delta}(B, A)$ is the \mathcal{O} -dual of γ and \vdots_{H} is induced by the tensor product over $\mathcal{O}H$, or equivalently, over B. We show that a p-permutation equivalence forces the defect groups and fusion systems of A and B to be isomorphic. In fact, via the notion of γ -Brauer pairs, it selects an isomorphism between defect groups and fusion systems, unique up to $G \times H$ -conjugation in a precise sense, see Theorem 1.1. Moreover, we show that a splendid Rickard equivalence between A and B induces a p-permutation equivalence, and that a p-permutation equivalence between A and B induces an isotypy, see Theorem 1.6. The goal of this paper is twofold: On the one hand, we want to understand what p-permutation equivalences can look like by finding necessary conditions on such an element $\gamma \in T^{\Delta}(A, B)$. On the other hand, we want to study which invariants of A and B are preserved under a p-permutation equivalence.

It turns out that the language of Brauer pairs (cf. 4.1(b) for a definition) is crucial for the study of p-permutation equivalences. We denote by $-^*$ the antipode $x \mapsto x^{-1}$ of any group algebra of a group X. Note that any (A, B)-bimodule belongs to the block algebra $A \otimes_{\mathcal{O}} B^*$ of $\mathcal{O}G \otimes \mathcal{O}H$, when viewed as $\mathcal{O}[G \times H] \cong \mathcal{O}G \otimes_{\mathcal{O}} \mathcal{O}H$ -module. A γ -Brauer pair is an $(A \otimes_{\mathcal{O}} B^*)$ -Brauer pair $(X, e \otimes f^*)$, where X is a p-subgroup of $G \times H$, e is a block idempotent of $\mathcal{O}C_G(p_1(X))$ and f is a block idempotent of $\mathcal{O}C_H(p_2(X))$, where $p_1: G \times H \to G$ and $p_2: G \times H \to H$ denote the canonical projections, which satisfies $\gamma(X, e \otimes f^*) \neq 0 \in T(FN_{G \times H}(X, e \otimes f^*))$. Here, $N_{G \times H}(X, e \otimes f^*)$ denotes the $G \times H$ -stabilizer of the Brauer pair $(X, e \otimes f^*)$, and the expression $\gamma(X, e \otimes f^*)$, is defined by applying the Brauer construction with respect to X (cf. 3.2(b)) to γ and then cutting with the idempotent $e \otimes f^*$. Note that X is necessarily a twisted diagonal subgroup $\Delta(P, \phi, Q)$ and that $C_{G \times H}(X) = C_G(P) \times C_H(Q)$, so that e is a block idempotent of $\mathcal{O}C_G(P)$ and f is a block idempotent of $\mathcal{O}C_H(Q)$. The following theorem shows that even though A and B are no longer required to have a *common* defect group, the element γ selects through the choice of a maximal γ -Brauer pair an isomorphism $\phi: E \xrightarrow{\sim} D$ between defect groups D and E of A and B, respectively. Recall that $(A \otimes_{\mathcal{O}} B^*)$ -Brauer pairs form a $G \times H$ -poset.

1.1 Theorem Assume that $\gamma \in T^{\Delta}(A, B)$ is a *p*-permutation equivalence between A and B.

(a) The set of γ -Brauer pairs is closed under $G \times H$ -conjugation and under taking smaller Brauer pairs. Moreover, the maximal γ -Brauer pairs form a single $G \times H$ -conjugacy class.

(b) Let $(\Delta(D, \phi, E), e \otimes f^*)$ be a maximal γ -Brauer pair. Then D is a defect group of A, E is a defect group of B, (D, e) is a maximal A-Brauer pair, (E, f) is a maximal B-Brauer pair, and the isomorphism $\phi \colon E \xrightarrow{\sim} D$ is an isomorphism between the fusion systems of B and A associated to (E, f) and (D, e), respectively.

The above theorem follows from the more precise Theorems 10.11 and 11.2.

The following theorem states additional restrictive properties of *p*-permutation equivalences.

1.2 Theorem Suppose that $\gamma \in T^{\Delta}(A, B)$ is a *p*-permutation equivalence between A and B and let $(\Delta(D, \phi, E), e \otimes f^*)$ be a maximal γ -Brauer pair.

(a) Every indecomposable (A, B)-bimodule appearing in γ has a vertex contained in $\Delta(D, \phi, E)$.

(b) Up to isomorphism, there exists a unique indecomposable (A, B)-bimodule M appearing in γ with vertex $\Delta(D, \phi, E)$. Its coefficient in γ is 1 or -1.

The module M in Theorem 1.2 is called the *maximal module* of γ . Theorem 1.2 follows from the stronger statements in Theorem 14.1 and 14.3.

The following theorem shows that Brauer constructions of *p*-permutation equivalences lead again to *p*-permutation equivalences or even Morita equivalences. It follows from the more precise statements in Theorem 11.4, Theorem 14.5, and Proposition 14.4.

1.3 Theorem Let $\gamma \in T^{\Delta}(A, B)$ be a *p*-permutation equivalence and let $(\Delta(P, \phi, Q), e \otimes f^*)$ be a γ -Brauer pair. Set $I := N_G(P, e)$ and $J := N_H(Q, f)$, let $Y := N_{G \times H}(\Delta(P, \phi, Q), e \otimes f^*) = N_{I \times J}(\Delta(P, \phi, Q))$, and let $\gamma' \in T(\mathcal{O}Y)$ denote the unique lift of $\gamma(\Delta(P, \phi, Q), e \otimes f^*) \in T(FY)$.

(a) $\operatorname{res}_{C_G(P)\times C_H(Q)}^Y(\gamma') \in T^{\Delta}(\mathcal{O}C_G(P)e, \mathcal{O}C_H(Q)f)$ is a p-permutation equivalence between $\mathcal{O}C_G(P)e$ and $\mathcal{O}C_H(Q)f$.

(b) $\operatorname{ind}_{V}^{I \times J}(\gamma') \in T^{\Delta}(\mathcal{O}Ie, \mathcal{O}Jf)$ is a *p*-permutation equivalence between $\mathcal{O}Ie$ and $\mathcal{O}Jf$.

(c) If $(\Delta(P, \phi, Q), e \otimes f^*)$ is a maximal γ -Brauer pair then $\gamma' = \pm [M']$ for an indecomposable p-permutation $\mathcal{O}Y$ -module M' which arises from the maximal module M of γ by $M' = M(\Delta(P, \phi, Q), e \otimes f^*)$. Moreover, $\operatorname{Res}_{C_G(P) \times C_H(Q)}^{Y}(M')$ induces a Morita equivalence between $\mathcal{O}C_G(P)e$ and $\mathcal{O}C_H(Q)f$, and $\operatorname{Ind}_Y^{I \times J}(M')$ induces a Morita equivalence between $\mathcal{O}Ie$ and $\mathcal{O}Jf$.

(d) Suppose that $(\Delta(P,\phi,Q), e \otimes f^*)$ is a maximal γ -Brauer pair, set $\tilde{Y} := N_{G \times H}(\Delta(P,\phi,Q))$ and let $\tilde{M} \in \mathcal{O}_Y \mod$ be the Green correspondent of M, then the p-permutation $(\mathcal{O}N_G(P), \mathcal{O}N_H(Q))$ -bimodule $\operatorname{Ind}_Y^{N_G(P) \times N_H(Q)}(\tilde{M})$ induces a Morita equivalence between the Brauer correspondents of A and B.

The following interesting additional properties of p-permutation equivalences follow from the more precise Theorems 11.10 and 12.3.

1.4 Theorem (a) The number of p-permutation equivalences between A and B is finite (possibly zero).

(b) If $\gamma \in T^{\Delta}(A, B)$ satisfies one of the two equations in (1) then it also satisfies the other.

Another invariant of a block algebra is given by the collection of Külshammer-Puig classes (see 4.6), one for every centric subgroup of a defect group in the associated fusion system. Since the fusion systems of A and Bare isomorphic, centric subgroups correspond. This gives a way to compare Külshammer-Puig classes of A and B. The following theorem follows from the more precise Theorem 13.4.

1.5 Theorem Suppose that $\gamma \in T^{\Delta}(A, B)$ is a *p*-permutation equivalence between A and B and let $(\Delta(P, \phi, Q), e \otimes f^*)$ be a γ -Brauer pair such that Z(P) is a defect group of the block algebra $\mathcal{O}C_G(P)e$. Set $I := N_G(P, e), J := N_H(Q, f), \overline{I} := I/PC_G(P)$ and $\overline{J} := J/QC_H(Q)$, and let $\kappa \in H^2(\overline{I}, F^{\times})$ and $\lambda \in H^2(\overline{J}, F^{\times})$ be the corresponding Külshammer-Puig classes of (P, e) and (Q, f). Then the isomorphism between \overline{I} and \overline{J} induced by $N_{G \times H}(\Delta(P, \phi, Q))$ (see Proposition 11.1) makes κ correspond to λ .

The following theorem is proved in Section 15.

1.6 Theorem (a) Suppose that the chain complex C_{\bullet} is a splendid Rickard equivalence between A and B (see Definition 15.1). Then the element $\gamma := \sum_{n \in \mathbb{Z}} (-1)^n [C_n] \in T^{\Delta}(A, B)$ is a p-permutation equivalence between A and B.

(b) Suppose that $\gamma \in T^{\Delta}(A, B)$ is a *p*-permutation equivalence between A and B and let $(\Delta(D, \phi, E), e \otimes f^*)$ be a maximal γ -Brauer pair. Then the Brauer constructions with respect to subgroups of $\Delta(D, \phi, E)$, yield an isotypy between A and B.

In [P99], Puig proved that some of the invariants (defect groups, fusion systems) of blocks considered here are preserved by splendid Rickard equivalences. Therefore, in view of Theorem 1.6(a), our results provide a significant improvement. We also use different techniques. That Külshammer-Puig classes, cf. Theorem 1.5 are preserved was not even known under the stronger hypothesis of a splendid Rickard equivalence.

One main point of view and crucial tool in this paper is that the Brauer construction of an (A, B)-bimodule with respect to a twisted diagonal subgroup $\Delta(P, \phi, Q)$ of $G \times H$ yields a module for the normalizer Y of $\Delta(P, \phi, Q)$. Rather than working with the restriction of this Brauer construction to $C_G(P) \times C_H(Q)$, we consistently work with the resulting $\mathcal{O}Y$ -module. This requires to lift the construction of tensor products of bimodules to a generalized tensor product functor

$$- \bigotimes_{H}^{X,Y} -: {}_{\Bbbk X} \operatorname{\mathsf{mod}} \times {}_{\Bbbk Y} \operatorname{\mathsf{mod}} \to {}_{\Bbbk [X*Y]} \operatorname{\mathsf{mod}}$$
(2)

for an arbitrary commutative ring \Bbbk and subgroups $X \leq G \times H$ and $Y \leq H \times K$, where $X * Y \leq G \times K$ is the composition of X and Y, viewed as correspondences between G and H, and H and K, respectively. This type of generalized tensor product was first used by Bouc in [Bc10b]. In Section 6 we develop the necessary properties of this generalized tensor product. Another main ingredient of our approach is a formula for the Brauer construction of the tensor product $M \otimes_{\mathcal{O}H} N$ of two *p*-permutation bimodules $M \in {}_{\mathcal{O}G}\mathsf{mod}_{\mathcal{O}H}$ and $N \in {}_{\mathcal{O}H}\mathsf{mod}_{\mathcal{O}K}$ in terms of a direct sum of tensor products of Brauer constructions of X and of Y. This formula goes back to earlier work in [BD12] and is refined to a block-wise version in Section 7.

The paper is arranged as follows. Section 2 recalls facts about subgroups X of direct product groups $G \times H$ and the composition $X * Y \leq G \times K$ if $Y \leq H \times K$. In Section 3 we recall the necessary preliminaries on p-permutation modules and add some technical lemmas that are used later. Preliminaries on blocks and Brauer pairs, together with some additional results on *p*-permutation modules in a block are given in Section 4. In Section 5 we introduce Brauer pairs for p-permutation modules, generalizing the concept of Brauer pairs of group algebras and blocks. We show that these Brauer pairs have very similar properties as the ones for blocks, cf. Proposition 5.3. In Section 6 we introduce the generalized tensor product (2) and prove basic properties of this construction. The main result in Section 7 is a formula (see Theorem 7.5) for the Brauer construction of the tensor product of two p-permutation bimodules. This formula incorporates the generalized tensor product and blocks. In Section 8 we recall Broué's notion of *perfect isometry* and prove several related results, while Section 9 introduces various representation groups associated to p-permutation modules, their relations with other representation groups, the notion of a Brauer pair for an element in the representation group of p-permutation modules, and the notion of a p-permutation equivalence. In Section 10 we study Brauer pairs of *p*-permutation equivalences and prove several surprising results; surprising, because one would not expect them to hold for virtual modules, but only for actual modules (cf. Proposition 10.8 and Theorem 10.11). Section 11 establishes that a p-permutation equivalence γ induces an isomorphism between the fusion systems of the underlying blocks (Theorem 11.2) and p-permutation equivalences on local levels through the Brauer construction with respect to a γ -Brauer pair (see Theorem 11.4). This section also contains character theoretic results that are interesting in their own right (see Proposition 11.8) and imply that irreducible characters that correspond via local equivalences have the same extension properties with respect to the inertia groups of their corresponding Brauer pairs. These properties lead to the finiteness of p-permutation equivalences between two given blocks. In Section 12 we prove a character theoretic criterion for an element in $T^{\Delta}(A, B)$ to be a p-permutation equivalence which leads to the equivalence of the two conditions in (1). That a p-permutation equivalence preserves the Külshammer-Puig classes is proved in Section 13. Section 14 establishes that every p-permutation equivalence has a maximal module and that the Green correspondent of the maximal module induces Morita equivalences between associated blocks on the local levels associated with the defect groups. Finally, in Section 15 we show that p-permutation equivalences are logically nested between splendid Rickard equivalences and isotypies.

1.7 Notation Throughout this paper we will use the following notation:

For a group G and $g \in G$ we write $c_g \colon G \to G$ or just g- for the conjugation map $x \mapsto gxg^{-1}$. For subgroups K and H of G, we write $K \leq_G H$ to denote that K is G-conjugate to a subgroup of H.

For a ring R we denote by Z(R), R^{\times} , J(R) its center, its unit group, and its Jacobson radical, respectively. Unadorned tensor products are taken over the ground ring that should be apparent from the context. For R-modules M and N, we write $M \mid N$ to indicate that M is isomorphic to a direct summand of N.

For a left G-set X, $H \leq G$, and $x \in X$, we write X^H for the set of H-fixed points of X, $[H \setminus X]$ for a set of representatives of the H-orbits of X, and $\operatorname{stab}_H(x)$ for the stabilizer of x in H.

Throughout this paper, $(\mathbb{K}, \mathcal{O}, F)$ denotes a *p*-modular system and π denotes a prime element of \mathcal{O} . We say that $(\mathbb{K}, \mathcal{O}, F)$ is *large enough* for a finite group G if \mathcal{O} contains a root of unity of order |G|. We write $\overline{\cdot}: \mathcal{O} \to F$ and $\overline{\cdot}: \mathcal{O}G \to FG$ for the natural epimorphisms. If M is an FG-module, we will view it without further explanation also as $\mathcal{O}G$ -module via restriction along $\mathcal{O}G \to FG$. Thus, expressions as $am \in M$ and $ab \in FG$ are defined for $a \in \mathcal{O}G$, $m \in M$ and $b \in FG$.

2 Subgroups of direct product groups

This section recalls basic facts and constructions related to subgroups of direct product groups. For more details the reader is referred to [Bc10a]

2.1 Let G, H, and K be finite groups and let $X \leq G \times H$ and $Y \leq H \times K$ be subgroups.

(a) We denote by $p_1: G \times H \to G$ and $p_2: G \times H \to H$ the canonical projections. Setting

 $k_1(X) := \{g \in G \mid (g,1) \in X\}$ and $k_2(X) := \{h \in H \mid (1,h) \in X\}$

one obtains normal subgroups $k_i(X)$ of $p_i(X)$ and canonical isomorphisms $X/(k_1(X) \times k_2(X)) \to p_i(X)/k_i(X)$, for i = 1, 2, induced by the projection maps p_i . The resulting isomorphism $\eta_X : p_2(X)/k_2(X) \xrightarrow{\sim} p_1(X)/k_1(X)$ satisfies $\eta_X(hk_2(X)) = gk_1(X)$ if and only if $(g, h) \in X$. Here, $(g, h) \in p_1(X) \times p_2(X)$.

(b) If $\phi \colon Q \xrightarrow{\sim} P$ is an isomorphism between subgroups $Q \leqslant H$ and $P \leqslant G$ then

$$\Delta(P,\phi,Q) := \{(\phi(y),y) \mid y \in Q\}$$

is a subgroup of $G \times H$. Subgroups arising this way will be called *twisted diagonal* subgroups of $G \times H$. For $P \leq G$ we also set $\Delta(P) := \Delta(P, \mathrm{id}_P, P) \leq G \times G$. Note that a subgroup $X \leq G \times H$ is twisted diagonal if and only if $k_1(X) = \{1\}$ and $k_2(X) = \{1\}$. Note also that for $(g, h) \in G \times H$ one has ${}^{(g,h)}\Delta(P, \phi, Q) = \Delta({}^{g}P, c_g \phi c_h^{-1}, {}^{h}Q)$.

(c) The subgroup $X^{\circ} := \{(h,g) \in H \times G \mid (g,h) \in X\}$ of $H \times G$ is called the *opposite* subgroup of X. Clearly, one has $(X^{\circ})^{\circ} = X$.

(d) The *composition* of X and Y is defined as

$$X * Y := \{ (g, k) \in G \times K \mid \exists h \in H : (g, h) \in X, (h, k) \in Y \}.$$

It is a subgroup of $G \times K$. Composition is associative. If $\Delta(P, \phi, Q) \leq G \times H$ and $\Delta(Q, \psi, R) \leq H \times K$ are twisted diagonal subgroups then $\Delta(P, \phi, Q) * \Delta(Q, \psi, R) = \Delta(P, \phi\psi, R) \leq G \times K$. Note that $(X * Y)^{\circ} = Y^{\circ} * X^{\circ}$, for arbitrary $X \leq G \times H$ and $Y \leq H \times K$.

The following lemma follows immediately from the definitions.

2.2 Lemma Let G, H and K be finite groups and let $X \leq G \times H$ and $Y \leq H \times K$ be subgroups.

- (a) One has $X * X^{\circ} = \Delta(p_1(X)) \cdot (k_1(X) \times \{1\}) = \Delta(p_1(X)) \cdot (\{1\} \times k_2(X)) = \Delta(p_1(X)) \cdot (k_1(X) \times k_2(X)).$
- (b) One has $X * X^{\circ} * X = X$.
- (c) If $p_1(X) \leq P \leq N_G(k_1(X))$ then $(\Delta(P) \cdot (k_1(X) \times \{1\})) * X = X$.
- (d) For any $q \in G$, $h \in H$, and $k \in K$, one has ${}^{(g,h)}X * {}^{(h,k)}Y = {}^{(g,k)}(X * Y)$.

2.3 Notation Let G and H be finite groups, let $\Delta(P, \phi, Q)$ be a twisted diagonal subgroup of $G \times H$, and let $S \leq N_G(P)$ and $T \leq N_H(Q)$. We denote by $N_{(S,\phi,T)}$ the subgroup of S consisting of all elements $g \in S$ such that there exists an element $h \in T$ satisfying $c_g \phi c_h = \phi$ as functions from Q to P. Note that if S contains $C_G(P)$ then also $N_{(S,\phi,T)}$ contains $C_G(P)$. Moreover, if $P \leq S$ and $Q \leq T$ then $P \leq N_{(S,\phi,T)}$. We further set $N_{\phi} := N_{(N_G(P),\phi,N_H(Q))}$. Note that this definition of N_{ϕ} corresponds to $N_{\phi^{-1}}$ in the literature on fusion systems, see for instance [AKO11].

The following proposition follows again immediately from the definitions and the conjugation formula in 2.1(b).

2.4 Proposition Let G and H be finite groups, let $\Delta(P, \phi, Q)$ be a twisted diagonal subgroup of $G \times H$, and let $C_G(P) \leq S \leq N_G(P)$ and $C_H(Q) \leq T \leq N_H(Q)$ be intermediate subgroups.

(a) One has $N_{G \times G}(\Delta(P)) = \Delta(N_G(P)) \cdot (C_G(P) \times \{1\}) = \Delta(N_G(P)) \cdot (\{1\} \times C_G(P)).$

(b) For $X := N_{G \times H}(\Delta(P, \phi, Q))$ one has $k_1(X) = C_G(P), k_2(X) = C_H(Q), p_1(X) = N_{\phi}, p_2(X) = N_{\phi^{-1}}.$

(c) For $X := N_{S \times T}(\Delta(P, \phi, Q))$ one has $k_1(X) = C_G(P)$, $k_2(X) = C_H(Q)$, $p_1(X) = N_{(S, \phi, T)}$, $p_2(X) = N_{(T, \phi^{-1}, S)}$.

3 *p*-permutation modules

In this section we recall module theoretic preliminaries and prove some results on *p*-permutation modules that will be needed later. For standard concepts of modular representation theory the reader is referred to [NT89].

We first recall concepts for modules over group rings $\Bbbk G$, where \Bbbk is an arbitrary commutative ring.

3.1 Let \Bbbk be a commutative ring and let G and H be finite groups.

(a) We always assume that, for a $(\Bbbk G, \Bbbk H)$ -bimodule M, the induced left and right \Bbbk -module structures coincide: $\alpha m = m\alpha$ for $m \in M$ and $\alpha \in \Bbbk$. One obtains an isomorphism between categories ${}_{kG} \operatorname{mod}_{\Bbbk H} \cong {}_{\Bbbk[G \times H]} \operatorname{mod}$ via the formula $(g, h)m = gmh^{-1}$ for $m \in M$ and $(g, h) \in G \times H$. We try to be consistent to

translate a left $G \times H$ -action into a (G, H)-biaction, i.e., we take the second component of the direct product to the right hand side. Similarly, we may view left $\Bbbk G$ -modules as right & G-modules via the anti-involution $-^*: kG \to kG, g \mapsto g^{-1}$.

(b) For a $(\Bbbk G, \Bbbk H)$ -bimodule M, we usually view its \Bbbk -dual $M^{\circ} := \operatorname{Hom}_{\Bbbk}(M, \Bbbk)$ as a $(\Bbbk H, \Bbbk G)$ -bimodule via (hfg)(m) := f(gmh), for $f \in M^{\circ}$, $m \in M$, $g \in G$ and $h \in H$. Similarly, if M is a left (resp. right) $\Bbbk G$ -module, we consider M° as right (resp. left) $\Bbbk G$ -module. However, we sometimes switch sides using the identification in (a). More generally, if $X \leq G \times H$ and M is a left $\Bbbk X$ -module, we can view M° as left $\Bbbk X^{\circ}$ -module via $((h,g)f)(m) = f((g^{-1}, h^{-1})m)$ for $f \in M^{\circ}$, $m \in M$ and $(g,h) \in X$.

(c) The *trivial* &G-module has underlying &-module & and satisfies $g\alpha = \alpha$ for all $g \in G$ and $\alpha \in \&$. We denote it by $\&_G$.

(d) If $H \leq G$, $g \in G$, and M is a $\Bbbk H$ -module, then we denote by ${}^{g}M$ the left $\Bbbk [{}^{g}H]$ -module with underlying \Bbbk -module M and ${}^{g}H$ -action given by restricting the H-action along the isomorphism $c_{g}^{-1} \colon {}^{g}H \to H$.

(e) For subgroups $Q \leq P \leq G$ and a $\Bbbk G$ -module M we denote by $M^P := \{m \in M \mid xm = m \text{ for all } x \in P\}$ the set of P-fixed points and by $\operatorname{tr}_Q^P : M^Q \to M^P$ the relative trace map defined by $m \mapsto \sum_{x \in [P/Q]} xm$.

(f) Let H be a subgroup of G, let $e \in \Bbbk H \cap Z(\Bbbk G)$ be an idempotent, and let N be a $\Bbbk H$ -module. Then $\operatorname{Ind}_{H}^{G}(eN) \cong e\operatorname{Ind}_{H}^{G}(N)$ as $\Bbbk G$ -modules. In fact, this follows from the obvious $\Bbbk G$ -module isomorphism $\Bbbk G \otimes_{\Bbbk H} eN \cong e(\Bbbk G \otimes_{\Bbbk H} N)$.

(g) Recall that a *permutation* &G-module M is a module that is isomorphic to &X for some finite left G-set X. Equivalently, M has a &-basis that is permuted by G.

Recall that $(\mathbb{K}, \mathcal{O}, F)$ denotes a *p*-modular system.

3.2 Let G be a finite group. The following constructions and statements will be used extensively throughout the paper.

- (a) One has a functor $\overline{?}$: $\mathcal{O}_G \mathsf{mod} \to {}_{FG}\mathsf{mod}$, given on objects by $M \mapsto \overline{M} := FG \otimes_{\mathcal{O}G} M \cong F \otimes_{\mathcal{O}} M \cong M/\pi M$.
- (b) For any p-subgroup $P \leq G$, one has a functor $-(P): \mathcal{O}_G \mathsf{mod} \to _{F[N_G(P)/P]} \mathsf{mod}$ given on objects by

$$M \mapsto M(P) := M^P / \left(\pi M^P + \sum_{Q < P} \operatorname{tr}_Q^P(M^Q) \right)$$

The module M(P) is called the *Brauer construction* of M at P. We often view M(P) as $F[N_G(P)]$ -module via inflation without notational indication. Similarly, one defines the Brauer construction M(P) of an FG-module M by $M(P) = M^P / \sum_{Q < P} \operatorname{tr}_Q^P(M^Q)$. The canonical map $M^P \to M(P)$ is denoted by Br_P^M , or just Br_P , and is called the *Brauer map*. Note that, for any intermediate subgroup $P \leq H \leq N_G(P)$ and any $\mathcal{O}G$ -module or FG-module M, one has $\operatorname{Res}_H^{N_G(P)}(M(P)) = (\operatorname{Res}_H^G(M))(P)$ as FH-modules.

(c) For a finite G-set X, the Brauer construction of the permutation $\mathcal{O}G$ -module $\mathcal{O}X$ can be described as follows: The composition of the canonical maps $\mathcal{O}[X^P] \to (\mathcal{O}X)^P \to (\mathcal{O}X)(P)$ induces an isomorphism of $F[N_G(P)/P]$ -modules $F[X^P] \xrightarrow{\sim} (\mathcal{O}X)(P)$, see [Br85, (1.1)(3)]. In the special case where $M = \mathcal{O}G$ and Gacts by conjugation, the map $\operatorname{Br}_{\mathcal{O}}^{\mathcal{O}G}$ translates under the above canonical isomorphism to the projection map $\operatorname{br}_P = \operatorname{br}_P^G : (\mathcal{O}G)^P \to FC_G(P), \sum_{g \in G} \alpha_g g \mapsto \sum_{g \in C_G(P)} \overline{\alpha}_g g$. This map is an \mathcal{O} -algebra homomorphism, called the Brauer homomorphism. Note that $\operatorname{br}_P(Z(\mathcal{O}G)) \subseteq Z(FC_G(P))^{N_G(P)} = Z(FN_G(P)) \cap FC_G(P)$.

(d) Let H be a subgroup of G, let M be an indecomposable $\mathcal{O}G$ -module (resp. FG-module) with vertex P and let N be an indecomposable $\mathcal{O}H$ -module (resp. FH-module) with vertex Q. If $M \mid \operatorname{Ind}_{H}^{G}(N)$ then $P \leq_{G} Q$. If $N \mid \operatorname{Res}_{H}^{G}(M)$ then $Q \leq_{G} P$.

Recall that an $\mathcal{O}G$ -module (resp. FG-module) M is called a *p*-permutation module if $\operatorname{Res}_P^G(M)$ is a permutation module for each *p*-subgroup P of G. An $(\mathcal{O}G, \mathcal{O}H)$ -bimodule (resp. (FG, FH)-bimodule) M is called a *p*-permutation bimodule if it is a *p*-permutation module when considered as $\mathcal{O}[G \times H]$ -module (resp. $F[G \times H]$ -module). If M is indecomposable its vertices form a conjugacy class of *p*-subgroups of $G \times H$.

In the next part of this section we recall some basic properties of p-permutation modules that will be used later. For more details on p-permutation modules we refer the reader to [Br85] and to [L18, Sections 5.10, 5.11].

3.3 Proposition Let G be a finite group.

- (a) Let M be an $\mathcal{O}G$ -module or an FG-module. Then the following are equivalent:
 - (i) M is a p-permutation module.

- (ii) M is isomorphic to a direct summand of a permutation module.
- (iii) Each indecomposable direct summand of M has the trivial module as source.

(b) The functor $M \mapsto \overline{M}$ induces a vertex-preserving bijection between the set of isomorphism classes of indecomposable *p*-permutation $\mathcal{O}G$ -modules and the set of isomorphism classes of indecomposable *p*-permutation FG-modules.

(c) For each p-subgroup P of G, the Brauer construction $M \mapsto M(P)$ induces a bijection between the set of isomorphism classes of indecomposable p-permutation $\mathcal{O}G$ -modules (resp. FG-modules) with vertex P and the set of isomorphism classes of indecomposable projective $F[N_G(P)/P]$ -modules. Moreover, if M is an indecomposable p-permutation FG-module with vertex P then the $F[N_G(P)]$ -module M(P) is the Green correspondent of M.

Proof See [Br85, (0.4), (3.2), (3.4), (3.5)].

3.4 Remark In view of (iii) in Part (a) of the previous proposition, *p*-permutation modules are often called *trivial source modules* in the literature. The reformulations of Part (a) allow to see quickly that the class of trivial source modules is closed under the usual constructions of restriction, induction, inflation, Brauer construction, $\otimes_{\mathcal{O}}$, \oplus , taking direct summands, and taking duals. Note that projective modules are *p*-permutation modules. Note also that Part (b) and the Krull-Schmidt theorem imply that the functor $\overline{?}: \mathcal{O}_{G} \mod \to {}_{FG} \mod \operatorname{induces}$ a bijection between the set of isomorphism classes of *p*-permutation $\mathcal{O}G$ -modules and the set of isomorphism classes of *p*-permutation $\mathcal{O}G$ -modules by $\mathcal{O}_{G} \operatorname{triv}$. Similarly we define ${}_{FG}\operatorname{triv}, {}_{O}\operatorname{g}\operatorname{triv}_{\mathcal{O}H}$, etc.

3.5 Proposition Let G be a finite group, let $M, N \in OG$ triv, and let P be a p-subgroup of G.

(a) One has $M(P)^{\circ} \cong (M^{\circ})(P)$ as $F[N_G(P)]$ -modules.

(b) One has canonical isomorphisms $M(P) \cong \overline{M}(P)$ and $(M \otimes_{\mathcal{O}} N)(P) \cong M(P) \otimes_F N(P)$ of $F[N_G(P)]$ modules. Moreover, if Q is a p-subgroup of $N_G(P)$ one has a canonical isomorphism $(M(P))(Q) \cong M(PQ)$ of $F[N_G(P) \cap N_G(Q)]$ -modules.

(c) The canonical map $\operatorname{Hom}_{\mathcal{O}G}(M, N) \to \operatorname{Hom}_{FG}(\overline{M}, \overline{N}), f \mapsto \overline{f}$, induces an *F*-linear isomorphism $F \otimes_{\mathcal{O}} \operatorname{Hom}_{\mathcal{O}G}(M, N) \xrightarrow{\sim} \operatorname{Hom}_{FG}(\overline{M}, \overline{N})$. In particular one has

 $\dim_{\mathbb{K}} \operatorname{Hom}_{\mathbb{K}G}(\mathbb{K} \otimes_{\mathcal{O}} M, \mathbb{K} \otimes_{\mathcal{O}} N) = \operatorname{rk}_{\mathcal{O}} \operatorname{Hom}_{\mathcal{O}G}(M, N) = \dim_{F} \operatorname{Hom}_{FG}(\overline{M}, \overline{N}).$

Proof (a) See the proof of [Br85, (2.4)(2)].

(b) In all three cases one obtains natural homomorphisms which are functorial in M (resp. M and N). Moreover, if M (resp. M and N) is a permutation module then 3.2(c) implies that these maps are isomorphisms. Thus, they are also isomorphisms for direct summands of permutation modules.

(c) This follows again immediately by reduction to the case of permutation modules.

The statements of the following lemma are folklore. We provide quick proofs for the reader's convenience and note that all statements also hold if M is a p-permutation FG-module, mutatis mutandis.

3.6 Lemma Let G be a finite group, let P be a p-subgroup of G, and let $M \in \mathcal{O}_G$ triv.

(a) Assume that M is indecomposable and that Q is a vertex of M. Then

$$M(P) \neq \{0\} \iff \mathcal{O}_P \mid \operatorname{Res}_P^G(M) \iff P \leqslant_G Q$$

(b) Assume that M is indecomposable, that Q is a vertex of M, and that P is normal in G. Then one has $P \leq Q$ if and only if P acts trivially on M.

(c) Assume that M is indecomposable and that P is normal in G. Then one of the two following must hold:

(i) P acts trivially on M and $M(P) = \overline{M}$, or

(ii) $\mathcal{O}_P \nmid \operatorname{Res}_P^G(M)$ and $M(P) = \{0\}.$

(d) There exists a decomposition $\operatorname{Res}_{N_G(P)}^G(M) = L \oplus N$ into $\mathcal{O}[N_G(P)]$ -submodules with the following property: P acts trivially on L and $\mathcal{O}_P \nmid \operatorname{Res}_P^{N_G(P)}(N)$. Moreover, $M(P) \cong \overline{L}$.

Proof (a) Let X be a P-set such that $\operatorname{Res}_{P}^{G}(M) \cong \mathcal{O}X$ as $\mathcal{O}P$ -modules. The first and second statement are both equivalent to $X^{P} \neq \emptyset$. Since the trivial $\mathcal{O}P$ -module has vertex P, 3.2(d) shows that the second statement implies the third. Conversely, since M has trivial source, one has $\mathcal{O}_{Q} \mid \operatorname{Res}_{Q}^{G}(M)$, and the third statement implies the second.

(b) If P acts trivially on M then $\mathcal{O}_P | \operatorname{Res}_P^G(M)$ and Part (a) implies $P \leq Q$. Conversely, since $M | \operatorname{Ind}_Q^G(F_Q)$, the Mackey formula and $P \leq Q$ imply that P acts trivially on M.

(c) Let Q be a vertex of M. If $P \leq Q$ then (i) holds by Part (b). If Q does not contain P then (ii) holds by Part (a).

(d) This follows by applying Part (c) and the last statement in 3.2(b) to each indecomposable direct summand of $\operatorname{Res}_{N_G(P)}^G(M)$.

The following Lemma is well-known and an easy exercise.

3.7 Lemma Let G be a finite group, $M \in {}_{FG}$ triv, $P \leq G$ a p-subgroup, and let $i \in (FG)^H$ be an idempotent, fixed under the conjugation action of a subgroup H of G containing P. Then, for each $m \in M^P$, one has $\operatorname{Br}_P^M(im) = \operatorname{br}_P^G(i)\operatorname{Br}_P^M(m)$. In particular, one obtains a canonical isomorphism $(iM)(P) \xrightarrow{\sim} \operatorname{br}_P^G(i)M(P)$ of $F[N_H(P)]$ -modules.

The following lemma will be used repeatedly.

3.8 Lemma (a) Let G be a finite group, let $M \in {}_{\mathcal{O}G}$ triv be indecomposable, and let P be a p-subgroup of G. Then each vertex of each indecomposable direct summand of the $F[N_G(P)]$ -module M(P) is contained in a vertex of M. (Note that $M(P) = \{0\}$ is possible, in which case the statement is vacuously true.)

(b) Let G and H be finite groups and let $M \in {}_{\mathcal{O}G}\mathsf{triv}_{\mathcal{O}H}$ be indecomposable with twisted diagonal vertices. Let X be any p-subgroup of $G \times H$. Then each vertex of each indecomposable direct summand of the $F[N_{G \times H}(X)]$ -module M(X) is again twisted diagonal.

Proof Part (b) is an immediate consequence of Part (a). In order to prove Part (a), note that $M(P) = \overline{M}(P) = (\operatorname{Res}_{N_G(P)}^G(\overline{M}))(P)$ and, by Lemma 3.6(d), the latter $F[N_G(P)]$ -module is isomorphic to a direct summand (possibly {0}) of $\operatorname{Res}_{N_G(P)}^G(\overline{M})$. Thus, Proposition 3.3(b) and 3.2(d) imply the result.

The following lemma will be used in the proof of Proposition 11.5.

3.9 Lemma Let G be a finite group, let Q be a p-subgroup of G, let $M \in {}_{F[N_G(Q)]}$ triv be indecomposable, and let R be a vertex of M. If P is a p-subgroup of $N_G(Q)$ satisfying $N_G(P) \leq N_G(Q)$ and $P \leq {}^{g}R$, for each $g \in G \setminus N_G(Q)$, then $(\operatorname{Ind}_{N_G(Q)}^G(M))(P) \cong M(P)$ as $F[N_G(P)]$ -modules.

Proof By 3.2(b), it suffices to show that $(\operatorname{Res}_{N_G(P)}^G(\operatorname{Ind}_{N_G(Q)}^G(M)))(P) \cong (\operatorname{Res}_{N_G(P)}^{N_G(Q)}(M))(P)$ as $F[N_G(P)]$ -modules. The Mackey formula yields $\operatorname{Res}_{N_G(P)}^G(\operatorname{Ind}_{N_G(Q)}^G(M)) \cong \bigoplus_{q \in [N_G(P) \setminus G/N_G(Q)]} L_g$, where

$$L_g := \operatorname{Ind}_{N_G(P) \cap {}^gN_G(Q)}^{N_G(P)} \left(\operatorname{Res}_{N_G(P) \cap {}^gN_G(Q)}^{{}^gN_G(Q)} ({}^gM) \right),$$

for $g \in G$. Since $L_1 = \operatorname{Res}_{N_G(P)}^{N_G(Q)}(M)$, it suffices to show that for each $g \in G \setminus N_G(Q)$ one has $L_g(P) = \{0\}$. Assume that there exists an element $g \in G \setminus N_G(Q)$ such that $L_g(P) \neq \{0\}$. By Lemma 3.6(a), there exists an indecomposable direct summand of L_g which has a vertex S that contains P. Two applications of 3.2(d) imply that there exists $x \in N_G(Q)$ and $y \in N_G(P)$ such that $S \leq ygxR$. Thus $P \leq ygxR$ and, since $y \in N_G(P)$, we obtain $P \leq gxR$, with $gx \notin N_G(Q)$, contradicting the hypothesis of the Lemma. The result now follows.

4 Block theoretic preliminaries

Throughout this section let G and H be finite groups and assume that $(\mathbb{K}, \mathcal{O}, F)$ is large enough for G and H and that $F = \mathcal{O}/(\pi)$ is algebraically closed.

4.1 Blocks, Brauer pairs, defect groups. (a) Recall that $\mathcal{O}G$ has a unique decomposition $\mathcal{O}G = B_1 \oplus \cdots \oplus B_t$ into indecomposable two-sided ideals, the blocks of $\mathcal{O}G$. If one decomposes the identity element of $\mathcal{O}G$ according to the block decomposition, $1 = e_1 + \cdots + e_t$, then e_1, \ldots, e_t are central, pairwise orthogonal idempotents of $\mathcal{O}G$ and e_i is an identity element of B_i , called the block idempotent of B_i . Thus, $B_i = e_i\mathcal{O}G$ is an \mathcal{O} -algebra in its own right. The block idempotents of $\mathcal{O}G$ are precisely the primitive idempotents of $Z(\mathcal{O}G)$. The identity element of a block B will be denoted by e_B . One obtains a bijection $B \mapsto e_B$ between the blocks of $\mathcal{O}G$ and the primitive idempotents of $Z(\mathcal{O}G)$. Similar statements hold for FG and the reduction map $\overline{?}$: $\mathcal{O}G \to FG$, $\sum_{g \in G} \alpha_g g \mapsto \overline{\alpha}_g g$ induces bijections $B_i \mapsto \overline{B}_i$ (resp. $e \mapsto \overline{e}$) between the blocks of $\mathcal{O}G$ and the blocks of $\mathcal{O}G$, if $B = e\mathcal{O}G$ for some central idempotent $e \neq 0$ of $\mathcal{O}G$, i.e., $B = \bigoplus_{i \in I} B_i$ and $e = \sum_{i \in I} e_i$ for a unique subset Iof $\{1, \ldots, t\}$. In this case $e_B := e$ is an identity of B. This way, sums of blocks and non-zero idempotents of $Z(\mathcal{O}G)$ are in bijective correspondence. Similarly, one defines sums of blocks of FG. Every block idempotents of $\mathcal{O}G \otimes \mathcal{O}H$ is of the form $e \otimes f$ for uniquely determined block idempotents e of $\mathcal{O}G$ and f of $\mathcal{O}H$. Note that we identify the \mathcal{O} -algebras $\mathcal{O}[G \times H]$ and $\mathcal{O}G \otimes_{\mathcal{O}} \mathcal{O}H$ via $(g, h) \mapsto g \otimes h$.

(b) Recall that a *Brauer pair* of FG is a pair (P, e), where P is a p-subgroup of G and e is a block idempotent of $F[C_G(P)]$. Note that the block idempotents of $F[C_G(P)]$ coincide with those of $F[PC_G(P)]$. The group Gacts by conjugation on the set of Brauer pairs of FG: ${}^g(P, e) := ({}^gP, {}^ge)$, for $g \in G$. We denote the G-stabilizer of the Brauer pair (P, e) by $N_G(P, e)$. Note that $PC_G(P) \leq N_G(P, e) \leq N_G(P)$. For Brauer pairs (P, e) and (Q, f) of FG, one writes $(Q, f) \leq (P, e)$ if $Q \leq P \leq N_G(Q, f)$ and $\operatorname{br}_P(f) \cdot e = e$ (or equivalently $\operatorname{br}_P(f) \cdot e \neq 0$). The transitive closure of the relation \leq on the set of Brauer pairs of FG is denoted by \leq . It is a partial order and is respected by G-conjugation.

If e is a central idempotent of FG then $br_P(e)$ is an $N_G(P)$ -stable central idempotent of $FC_G(P)$ and also a central idempotent of $FN_G(P)$, see 3.2(c). If B is a sum of blocks of FG, one says that a Brauer pair (P, e)is a B-Brauer pair if $br_P(e_B)e = e$, or equivalently, $br_P(e_B)e \neq 0$. Every Brauer pair is a B-Brauer pair for a unique block B of FG, and in this case $(\{1\}, e_B) \leq (P, e)$; see also Proposition 4.2(a) below. Let B be again a sum of blocks of FG. The set of B-Brauer pairs is closed under G-conjugation and if $(Q, f) \leq (P, e)$ are Brauer pairs of FG then (P, e) is a B-Brauer pair if and only if (Q, f) is a B-Brauer pair. This follows also immediately from Proposition 4.2(a) below. The set of B-Brauer pairs is denoted by $\mathcal{BP}(B)$.

For a sum *B* of blocks of $\mathcal{O}G$ we simply define $\mathcal{BP}(B) := \mathcal{BP}(\overline{B})$ and call them Brauer pairs of *B*. Thus, Brauer pairs by default are viewed as pairs (P, e), where *e* is an idempotent of a group algebra over *F*. Sometimes it is convenient to lift the idempotent *e* to an idempotent over \mathcal{O} . We denote the set of the resulting pairs by $\mathcal{BP}_{\mathcal{O}}(B)$ or $\mathcal{BP}_{\mathcal{O}}(\overline{B})$.

(c) A defect group of a block B of FG is a subgroup D of G, minimal with respect to the property that $e_B \in \operatorname{tr}_D^G((FG)^D)$, see [NT89, Section 5.1]. The defect groups of B form a G-conjugacy class of p-subgroups of G. A subgroup D of G is a defect group of B if and only if $\Delta(D)$ is a vertex of B, viewed as indecomposable $F[G \times G]$ -module, see [NT89, Theorem 10.8]. If P is a normal p-subgroup of G then P is contained in each defect group of each block B, see [NT89, Theorem 5.2.8], and e_B is contained in the F-span of the p'-elements of $C_G(P)$, see [NT89, Theorem 3.6.22(ii)]. Similarly one defines defect groups of blocks of $\mathcal{O}G$. All the above statements hold again over \mathcal{O} and defect groups don't change under reduction of blocks modulo π .

The following Proposition recalls more standard facts about Brauer pairs, see Theorem 1.8 and 1.14 in [BrP80].

4.2 Proposition (a) For each Brauer pair (P, e) of FG and each subgroup $Q \leq P$, there exists a unique Brauer pair (Q, f) of FG such that $(Q, f) \leq (P, e)$. In particular, if $(R, g) \leq (P, e)$ are Brauer pairs of FG and $R \leq Q \leq P$ then there exists a unique Brauer pair (Q, f) of FG satisfying $(R, g) \leq (Q, f) \leq (P, e)$.

(b) Let (Q, f) and (P, e) be Brauer pairs of FG. If $(Q, f) \leq (P, e)$ and $Q \leq P$ then $(Q, f) \leq (P, e)$.

(c) Let B be a block of FG. Then the maximal elements in the poset of B-Brauer pairs form a single full conjugacy class. Moreover, a B-Brauer pair (P, e) is a maximal B-Brauer pair if and only if P is a defect group of B.

The following Proposition is well-known. We give a proof for the reader's convenience. Note that $G \times G$ acts on FG via its $F[G \times G]$ -module structure and that G acts on FG via conjugation. These actions are linked via the diagonal embedding $\Delta \colon G \mapsto G \times G, g \mapsto (g, g)$, so that $(FG)^{\Delta(H)} = (FG)^H$ for all subgroups $H \leq G$.

4.3 Proposition Let B be a block of FG and let e_B denote the identity element of B. Furthermore, let (Q, e) be a B-Brauer pair and set $I := N_G(Q, e)$.

- (a) One has $B(\Delta(Q)) \cong F[C_G(Q)] \operatorname{br}_Q(e_B)$ as $F[N_{G \times G}(\Delta(Q))]$ -modules.
- (b) One has $eB(\Delta(Q))e \cong F[C_G(Q)]e$ as $F[N_{I\times I}(\Delta(G))]$ -modules.

Proof (a) Recall that $\operatorname{br}_Q \colon (FG)^Q \to F[C_G(Q)]$ is a surjective *F*-algebra homomorphism and an $F[N_{G\times G}(\Delta(Q))]$ -module homomorphism. Since $B^Q = (FG)^Q e_B$ and br_Q is multiplicative, we obtain $\operatorname{br}_Q(B^Q) = \operatorname{br}_Q((FG)^Q)\operatorname{br}_Q(e_B) = F[C_G(Q)]\operatorname{br}_Q(e_B)$. Moreover, we have $\operatorname{ker}(\operatorname{br}_Q) = \sum_{R < Q} \operatorname{tr}_R^Q((FG)^R)$. Thus, $\operatorname{ker}(\operatorname{br}_Q) \cap B^Q = \sum_{R < Q} \operatorname{tr}_R^Q(B^R)$. Altogether, we obtain

$$B(\Delta(Q)) = B^Q / \sum_{R < Q} \operatorname{tr}_R^Q(B^R) \cong F[C_G(Q)] \operatorname{br}_Q(e_B)$$

as $F[N_{G \times G}(\Delta(Q))]$ -modules.

Part (b) follows immediately from Part (a).

4.4 The fusion system of a block. Next we define the fusion system associated to a block, a structure and block invariant introduced by Puig. See [AKO11] for the definition of fusion systems, saturated fusion systems and basic facts about them. Let B be a block of FG and let (P, e) be a maximal B-Brauer pair. For $Q \leq P$, denote by e_Q the unique block idempotent of $FC_G(Q)$ such that $(Q, e_Q) \leq (P, e)$, cf. 4.2(a).

The fusion system of B, associated with (P, e), is the category \mathcal{F} whose objects are the subgroups of P, and whose morphism set $\operatorname{Hom}_{\mathcal{F}}(Q, R)$, for subgroups Q and R of P, is defined as the set of group homomorphisms arising as conjugation maps $c_g \colon Q \to R$, where $g \in G$ satisfies ${}^{g}(Q, e_Q) \leq (R, e_R)$. The category \mathcal{F} is a saturated fusion system on P, see for instance [AKO11, IV.3] for a proof.

Recall that, for a general fusion system \mathcal{F} on a *p*-group P, a subgroup Q of P is called *fully* \mathcal{F} -centralized (resp. *fully* \mathcal{F} -normalized) if $|C_P(Q)| \ge |C_P(Q')|$ (resp. $|N_P(Q)| \ge |N_P(Q')|$) for all subgroups Q' of P that are \mathcal{F} -isomorphic to Q. Recall also that a subgroup Q of P is called \mathcal{F} -centric if $C_P(Q') = Z(Q')$ for all Q' that are \mathcal{F} -isomorphic to Q.

We will need the following result that goes back to Alperin and Broué, see [AB79]. We'll use the formulation given in [L06, Theorems 2.4 and 2.5] and in [K07, Theorem 3.11(i)].

4.5 Proposition Let B be a block of FG, let (P, e) be a maximal B-Brauer pair, and let \mathcal{F} be the fusion system associated to B and (P, e). For every subgroup Q of P, denote by e_Q the unique primitive idempotent in $Z(F[C_G(Q)])$ such that $(Q, e_Q) \leq (P, e)$.

(a) A subgroup Q of P is fully \mathcal{F} -centralized if and only if $C_P(Q)$ is a defect group of $F[C_G(Q)]e_Q$. In this case $(C_P(Q), e_{QC_P(Q)})$ is a maximal Brauer pair of the block algebra $F[C_G(Q)]e_Q$. In particular, Q is \mathcal{F} -centric if and only if Z(Q) is a defect group of $F[C_G(Q)]e_Q$.

(b) A subgroup Q of P is fully \mathcal{F} -normalized if and only if $N_P(Q)$ is a defect group of the block algebra $F[N_G(Q, e_Q)]e_Q$. In this case $(N_P(Q), e_{N_P(Q)})$ is a maximal $F[N_G(Q, e_Q)]e_Q$ -Brauer pair.

4.6 Külshammer-Puig classes. Let (P, e) be a self-centralizing Brauer pair of FG, i.e., such that Z(P) is the defect group of the block $F[C_G(P)]e$. Set $I := N_G(P, e)$ and $\overline{I} := I/PC_G(P)$. By [NT89, Theorems 5.8.10 and 5.8.11], P is the defect group of the block $F[PC_G(P)]e$ and hence, by [NT89, Lemma 5.8.12], the block algebra $F[PC_G(P)]e$ has a unique simple module V. Since V is I-stable, the canonical cohomology class $\kappa \in H^2(\overline{I}, F^{\times})$, assigned to the data $PC_G(P) \leq I$ and V by Schur (see [NT89, Theorem 3.5.7]), is called the Külshammer-Puig class of (P, e).

Recall that a surjective group homomorphism $f: G \to \overline{G}$ induces two functors over any commutative ring \Bbbk : The *inflation* functor $\operatorname{Inf}_{\overline{G}}^{\underline{G}}: {}_{\Bbbk\overline{G}}\mathsf{Mod} \to {}_{\Bbbk G}\mathsf{Mod}$ which is given by restriction along the homomorphism f; and the *deflation* functor $\operatorname{Def}_{\overline{G}}^{\underline{G}}: {}_{\Bbbk\overline{G}}\mathsf{Mod} \to {}_{\Bbbk\overline{G}}\mathsf{Mod}$ which assigns to a $\Bbbk G$ -module M the largest factor-module on which $N := \ker(f)$ acts trivially. More explicitly, $\operatorname{Def}_{\overline{G}}^{\underline{G}}(M) = {}_{\Bbbk}\overline{G} \otimes_{\Bbbk G} M \cong M/I_N M$, where ${}_{\Bbbk}\overline{G}$ is viewed as $({}_{\Bbbk}\overline{G}, {}_{\Bbbk}G)$ -bimodule using f for the right module structure, and where I_N is the ideal of ${}_{\Bbbk}G$ generated by the elements $x - 1, x \in N$. Note that $I_N = \ker({}_{\Bbbk}G \to {}_{\Bbbk}\overline{G})$.

4.7 Proposition Let B be a block of FG with normal defect group $P \leq G$.

(a) For any B-module M one has an FG-module isomorphism $M/J(M) \cong \operatorname{Inf}_{G/P}^G \operatorname{Def}_{G/P}^G(M)$.

Π

(b) Let Q be a normal subgroup of G with $Q \leq P$. Then $M \mapsto \operatorname{Inf}_{G/P}^G \operatorname{Def}_{G/P}^G(M)$ induces a bijection between the set of isomorphism classes of indecomposable *p*-permutation *B*-modules with vertex Q and the set of isomorphism classes of simple *B*-modules.

Proof Denote by $\overline{:}: FG \to F[G/P]$ the canonical *F*-algebra homomorphism. Then, by [NT89, Theorems 5.8.10 and 5.8.7(ii)], \overline{B} is a non-zero sum of blocks of F[G/P] of defect 0 and therefore a semisimple *F*-algebra.

(a) As above, let I_P denote the ideal of FG generated by the elements $x - 1, x \in P$. It suffices to show that $I_PM = J(M)$. Since P is normal in G, P acts trivially on every simple B-module and I_P annihilates every simple B-module. Thus, $I_P \subseteq J(B)$ and $I_PM \subseteq J(B)M = J(M)$. For the converse it suffices to show that M/I_PM is semisimple as FG-module. But the FG-module M/I_PM is the inflation of a semisimple \overline{B} -module, and therefore semisimple.

(b) By Proposition 3.3(c) the Brauer construction $M \mapsto M(Q)$ and $\operatorname{Inf}_{G/Q}^G$ define inverse bijections between the set of isomorphism classes of indecomposable *p*-permutation *B*-modules with vertex *Q* and the set of isomorphism classes of projective indecomposable F[G/Q]-modules which after inflation belong to *B*. Each such projective indecomposable F[G/Q]-module is the projective cover P_V in $_{F[G/Q]}$ -module of a simple *B*-module *V*, viewed as F[G/Q]-module. Thus, using Part (a), it suffices to show that for $M = \operatorname{Inf}_{G/Q}^G(P_V)$ one has $M/J(M) \cong V$ as *FG*-modules. But this is clear, since the *FG*-submodules *U* of *M* are the same as the F[G/Q]-submodules of P_V and the factor module M/U is semisimple as *FG*-module if and only if P_V/U is semisimple as F[G/Q]-module.

4.8 Proposition Let B be a block of FG with central defect group P, let $Q \leq P$, and denote by $\overline{\cdot}$: FG \rightarrow F[G/Q] the natural surjective F-algebra homomorphism.

(a) The image $\overline{B} \subseteq F[G/Q]$ of B is a block of F[G/Q] with defect group P/Q. Up to isomorphism, \overline{B} has a unique simple module V.

(b) Up to isomorphism, there exists a unique indecomposable p-permutation B-module M with vertex Q. It is isomorphic to the inflation of the unique indecomposable projective \overline{B} -module P_V . Moreover, one has an F[G/Q]-module isomorphism $V \cong \operatorname{Inf}_{G/P}^{G/Q}\operatorname{Def}_{G/P}^G(M)$ and $\operatorname{Inf}_{G/Q}^G(V)$ is the unique simple B-module.

Proof (a) This follows from Theorems 5.8.10 and 5.8.11 and from Lemma 5.8.12 in [NT89].

(b) This follows from Proposition 4.7(b), Part (a), and Proposition 3.3(c).

5 Brauer pairs for *p*-permutation modules

Throughout this section, G denotes a finite group. In analogy to Brauer pairs for blocks, we introduce Brauer pairs for p-permutation modules.

5.1 Definition Let $M \in {}_{\mathcal{O}G}$ triv or $M \in {}_{FG}$ triv. We call a Brauer pair (P, e) of FG an M-Brauer pair if $M(P, e) := e \cdot M(P) \neq \{0\}$. Note that M(P, e) is an FIe-module, where $I := N_G(P, e)$. For $M \in {}_{\mathcal{O}G}$ triv the set of M-Brauer pairs coincides with the set of \overline{M} -Brauer pairs. It is denoted by $\mathcal{BP}(M)$ or $\mathcal{BP}(\overline{M})$. The corresponding set of Brauer pairs over \mathcal{O} will be denoted by $\mathcal{BP}_{\mathcal{O}}(M)$ or $\mathcal{BP}(\overline{M})$.

This generalizes the notion of *B*-Brauer pairs for a block *B* of $\mathcal{O}G$ in the following sense: A Brauer pair (P, e) of *FG* is a *B*-Brauer pair as defined in 4.1(b) if and only if $(\Delta(P), e \otimes e^*)$ is a *B*-Brauer pair of the indecomposable $\mathcal{O}[G \times G]$ -module *B* as defined above.

5.2 We will use the following Morita equivalence between block algebras in the proof of the next two propositions. Recall from [NT89, Theorem 5.5.12] that if (Q, f) is a Brauer pair of $\mathcal{O}G$, if $I := N_G(Q, f)$, and if $e := \operatorname{tr}_I^{N_G(Q)}(f)$ is the block idempotent of $\mathcal{O}[N_G(Q)]$ covering f, i.e., the unique block idempotent e of $\mathcal{O}[N_G(Q)]$ such that $ef \neq 0$, then one has a Morita equivalence between $\mathcal{O}[N_G(Q)]_e \mod \mathcal{O}_I f \mod \mathcal{O}_I f \mod \mathcal{O}[N_G(Q)]$ from the left with the $(\mathcal{O}If, \mathcal{O}[N_G(Q)]e)$ -bimodule $f\mathcal{O}[N_G(Q)] = f\mathcal{O}[N_G(Q)]e$. This functor is naturally isomorphic to the functor $f \cdot \operatorname{Res}_I^{N_G(Q)}$. Its inverse is given by tensoring with the $(\mathcal{O}[N_G(Q)]e, \mathcal{O}If)$ -bimodule $\mathcal{O}[N_G(Q)]f = e\mathcal{O}[N_G(Q)]f$. It is naturally isomorphic to the functor $\operatorname{Ind}_I^{N_G(Q)}$. This Morita equivalence induces a similar equivalence over F.

The following proposition generalizes standard facts on Brauer pairs for blocks to Brauer pairs for p-permutation modules. Part (c) can also be derived from Theorem 2.5 in [S90], but we give an independent proof.

5.3 Proposition Let M be a p-permutation FG-module.

(a) Assume that M belongs to a sum of blocks B of FG, i.e., $e_B M = M$. Then $\mathcal{BP}(M) \leq \mathcal{BP}(B)$.

(b) $\mathcal{BP}(M)$ is a G-stable ideal in the poset $\mathcal{BP}(FG)$, i.e., it is stable under G-conjugation and if $(Q, f) \leq (P, e)$ are Brauer pairs of FG such that (P, e) is an M-Brauer pair then also (Q, f) is an M-Brauer pair.

(c) Assume that M is indecomposable. Then the maximal M-Brauer pairs are precisely the M-Brauer pairs (P, e), where P is a vertex of M. Moreover, any two maximal M-Brauer pairs are G-conjugate.

Proof (a) By Lemma 3.7 we have $\{0\} \neq eM(P) = e((e_BM)(P)) = e \operatorname{br}_P(e_B)M(P)$, and therefore $e \operatorname{br}_P(e_B) \neq 0$. Thus, (P, e) is a *B*-Brauer pair.

(b) For any Brauer pair (P, e) of FG and any $g \in G$ one has ${}^{g}e \cdot M({}^{g}P) \cong {}^{g}(e \cdot M(P))$ as $F[N_{G}({}^{g}P, {}^{g}e)]$ -modules. Thus, (P, e) is an M-Brauer pair if and only if ${}^{g}(P, e)$ is an M-Brauer pair.

Now let $(Q, f) \leq (P, e)$ be Brauer pairs of FG and assume that (P, e) is an M-Brauer pair. In order to show that (Q, f) is an M-Brauer pair, we may assume that $(Q, f) \leq (P, e)$. Then $P \leq N_G(Q, f) =: I$ and $\operatorname{br}_P(f)e = e$. For the FI-module $f \cdot M(Q)$, Lemma 3.7 and Proposition 3.5(b) imply the following isomorphisms of $FC_G(P)$ -modules: $(f \cdot M(Q))(P) \cong \operatorname{br}_P(f) \cdot (M(Q)(P)) \cong \operatorname{br}_P(f) \cdot M(P)$. With this we obtain $\{0\} \neq e \cdot M(P) = e \cdot \operatorname{br}_P(f) \cdot M(P) \cong e \cdot (f \cdot M(Q))(P)$, which implies $f \cdot M(Q) \neq \{0\}$.

(c) First we claim that any two *M*-Brauer pairs of the form (P, e), where *P* is a vertex of *M*, are *G*-conjugate. As the vertices of *M* are *G*-conjugate, it suffices to fix a vertex *P* of *M* and to show that any two *M*-Brauer pairs of the form (P, e) are $N_G(P)$ -conjugate. By Proposition 3.3(c), the $F[N_G(P)]$ -module M(P) is the Green correspondent of *M*. Thus, by Lemma 5.5.4 in [NT89], $eM(P) \neq \{0\}$ if and only if the block $eF[C_G(P)]$ is covered by the block of $FN_G(P)$ to which M(P) belongs. But all these blocks $eF[C_G(P)]$ are $N_G(P)$ -conjugate, see Lemma 5.5.3 in [NT89], and the claim is proved.

In order to prove the statements in (c) it suffices now to show the following claim: Each M-Brauer pair (Q, f) is contained in some M-Brauer pair (P, e), where P is a vertex of M. First note that, since (Q, f) is an *M*-Brauer pair, we have $M(Q) \neq \{0\}$. This implies that Q is contained in a vertex P of M (see Lemma 3.6(a)). We proceed by induction on the index [P : Q]. If Q = P, the claim is trivially true. Assume now that Q < P. Set $I := N_G(Q, f)$. Since (Q, f) is an M-Brauer pair, we have $fM(Q) \neq \{0\}$ and there exists an indecomposable direct summand N of the $F[N_G(Q)]$ -module M(Q) such that $fN \neq \{0\}$. By 5.2, the FIf-module fN is indecomposable. Assume first that fN has vertex Q. Since, by 5.2, $fN \mid \operatorname{Res}_{I}^{N_{G}(Q)}(N)$ and $N \cong \operatorname{Ind}_{I}^{N_{G}(Q)}(fN)$, also N has vertex Q (see 3.2(d)). Since $N \mid M(Q) \mid \operatorname{Res}_{N_{G}(Q)}^{G}(M)$ (see Lemma 3.6(d)), also M has vertex Q by the Burry-Carlson-Puig Theorem (see [NT89, Theorem 4.4.6]), a contradiction. Thus, the FIf-module fN has a vertex R with Q < R. By Lemma 3.6(a) we have $(fN)(R) \neq \{0\}$. Since $fN \mid fM(Q)$, we also have $(fM(Q))(R) \neq \{0\}$. Since $\{0\} \neq (fM(Q))(R) \cong \operatorname{br}_R(f)M(R)$ as $F[C_G(R)]$ -modules (see Lemma 3.7) and Proposition 3.5(b)) we obtain $br_R(f) \neq 0$, and since $br_R(f)$ is central in $F[C_G(R)]$, there exists a block idempotent e of $FC_G(R)$ such that $e \operatorname{br}_R(f) = e$ and $e \cdot M(R) \neq \{0\}$. As $R \leq I$ and $e \operatorname{br}_R(f) = e$, we have $(Q, f) \triangleleft (R, e)$. Since $e \cdot M(R) \neq \{0\}$, also (R, e) is an M-Brauer pair. Applying the induction hypothesis to (R, e) we have $(R, e) \leq (P', e')$ for some M-Brauer pair (P', e') such that P' is a vertex of M. This concludes the proof.

5.4 Proposition Let M and N be indecomposable p-permutation FG-modules, suppose that $(P, e) \in \mathcal{BP}(FG)$ is both a maximal M-Brauer pair and a maximal N-Brauer pair, and set $I := N_G(P, e)$. Then M(P, e) and N(P, e) are indecomposable p-permutation FIe-modules. Moreover, $M \cong N$ if and only if $M(P, e) \cong N(P, e)$ as FIe-modules.

Proof This is an immediate consequence of Proposition 5.3(c), the Green correspondence, and the Morita equivalence from 5.2 (with (Q, f) replaced by (P, e)).

6 Extended tensor products and homomorphisms

Throughout this section, G, H, and K denote finite groups and \Bbbk denotes a commutative ring. We consider extended versions of tensor products and homomorphism sets of bimodules for group algebras and prove several basic facts about this construction.

6.1 (a) Let $X \leq G \times H$ and $Y \leq H \times K$. Further, let $M \in {}_{\Bbbk X}$ mod and $N \in {}_{\Bbbk Y}$ mod. Since $k_1(X) \times k_2(X) \leq X$, the k-module M can be viewed as $(\Bbbk[k_1(X)], \Bbbk[k_2(X)])$ -bimodule. Similarly, N can be considered as $(\Bbbk[k_1(Y)], \Bbbk[k_2(Y)])$ -bimodule, and $M \otimes_{\Bbbk[k_2(X) \cap k_1(Y)]} N$ is a $(\Bbbk[k_1(X)], \Bbbk[k_2(Y)])$ -bimodule. Note that $k_1(X) \times k_2(Y) \leq X * Y$ and that this bimodule structure can be extended to a $\Bbbk[X * Y]$ -module structure such that, for $(g, k) \in X * Y$, $m \in M$, and $n \in N$, one has

$$(g,k) \cdot (m \otimes n) = (g,h)m \otimes (h,k)n, \qquad (3)$$

where $h \in H$ is chosen such that $(g, h) \in X$ and $(h, k) \in Y$. To the best of our knowledge, this construction was first used in [Bc10b]. We will denote this extended tensor product by $M \bigotimes_{\Bbbk H}^{X,Y} N \in_{\Bbbk[X*Y]} \mathsf{mod}$ and obtain a functor $-\bigotimes_{\Bbbk H}^{X,Y} -: {}_{\Bbbk X}\mathsf{mod} \times_{\Bbbk Y}\mathsf{mod} \to_{\Bbbk[X*Y]}\mathsf{mod}$. A quick calculation shows that this construction is associative: If also L is a finite group, $Z \leq K \times L$, and $P \in_{\Bbbk Z}\mathsf{mod}$ then $(M \bigotimes_{\Bbbk H}^{X,Y} N) \bigotimes_{\Bbbk K}^{X*Y,Z} P$ and $M \bigotimes_{\Bbbk H}^{X,Y*Z} (N \bigotimes_{\Bbbk H}^{X,Y} P)$ are canonically isomorphic under $(m \otimes n) \otimes p \mapsto m \otimes (n \otimes p)$, for $m \in M$, $n \in N$ and $p \in P$. Clearly, $\bigotimes_{\Bbbk H}^{X,Y}$ also behaves distributively with respect to direct sums. Moreover, for any $g \in G$, $h \in H$, and $k \in K$ one has an isomorphism

$${}^{(g,k)}_{(M \underset{\Bbbk H}{\otimes} N)} (M \underset{\Bbbk H}{\overset{(g,h)}{\otimes} N}) \cong {}^{(g,h)}_{M} M \underset{\Bbbk H}{\overset{(g,h)}{\otimes} Y} {}^{(h,k)}_{(h,k)} N$$

$$(4)$$

(+ 1)

of $\mathbb{k}[{}^{(g,k)}(X * Y)]$ -modules, cf. Lemma 2.2(d).

(b) Let $X \leq H \times G$, $Y \leq H \times K$, $M \in {}_{\Bbbk X}$ mod, and $N \in {}_{\Bbbk Y}$ mod. Then, $M \in {}_{\Bbbk[k_1(X)]}$ mod ${}_{\Bbbk[k_2(X)]}$, $N \in {}_{\Bbbk[k_1(Y)]}$ mod ${}_{\Bbbk[k_2(Y)]}$ and consequently, $\operatorname{Hom}_{\Bbbk[k_1(X)\cap k_1(Y)]}(M,N) \in {}_{\Bbbk[k_2(X)]}$ mod ${}_{\Bbbk[k_2(Y)]}$. This bimodule structure can be extended to a $\Bbbk[X^{\circ} * Y]$ -module structure satisfying

$$((g,k) \cdot f)(m) = (h,k)f((h,g)^{-1}m),$$

for $(g,k) \in X^{\circ} * Y$, $f \in \operatorname{Hom}_{\Bbbk[k_1(X) \cap k_1(Y)]}(M, N)$, and $m \in M$, where $h \in H$ is chosen such that $(h,g) \in X$ and $(h,k) \in Y$. We leave the details of this straightforward verification to the reader. We will denote the resulting $\Bbbk[X^{\circ} * Y]$ -module by $\operatorname{LHom}_{\Bbbk H}^{X,Y}(M, N)$. The symbol LHom is used, since often G, H and K will coincide and it might not be clear if one uses homomorphisms of left modules or right modules.

(c) Let $X \leq G \times H$, $Y \leq K \times H$, $M \in {}_{\Bbbk X} \mathsf{mod}$, and $N \in {}_{\Bbbk Y} \mathsf{mod}$. Similarly as in (b), considering homomorphisms with respect to right module structures, we obtain $\operatorname{Hom}_{\Bbbk[k_2(X) \cap k_2(Y)]}(M, N) \in {}_{\Bbbk[k_1(Y)]} \mathsf{mod}_{\Bbbk[k_1(X)]}$. This bimodule structure can be extended to a $\Bbbk[Y * X^\circ]$ -module structure satisfying

$$((k,g) \cdot f)(m) = (k,h)f((g,h)^{-1}m),$$

for $(k,g) \in Y * X^{\circ}$, $f \in \operatorname{Hom}_{\Bbbk[k_2(X) \cap k_2(Y)]}(M,N)$, and $m \in M$, were $h \in H$ is chosen such that $(k,h) \in Y$ and $(g,h) \in X$. We denote the resulting $\Bbbk[Y * X^{\circ}]$ -module by $\operatorname{RHom}_{\Bbbk H}^{X,Y}(M,N)$.

(d) Let X, Y, M, and N be as in (c). Note that, by restriction along the flip isomorphism $\tau \colon X \to X^{\circ}$, $(g, h) \mapsto (h, g)$, we obtain a $\Bbbk X^{\circ}$ -module M^{τ} and similarly, a $\Bbbk Y^{\circ}$ -module N^{τ} . With these operations, one has the equality $\operatorname{RHom}_{\Bbbk H}^{X,Y}(M,N)^{\tau} = \operatorname{LHom}_{\Bbbk H}^{X^{\circ},Y^{\circ}}(M^{\tau},N^{\tau})$ of $\Bbbk [X * Y^{\circ}]$ -modules.

Note that the tensor product construction in 6.1(a) generalizes both the tensor product of bimodules (when $X = G \times H$ and $Y = H \times K$) and the internal tensor product of $\Bbbk G$ -modules (when $X = Y = \Delta(G)$).

For later use, we will state the following theorem due to Serge Bouc, see [Bc10b].

6.2 Theorem Let $X \leq G \times H$, $Y \leq H \times K$, $M \in {}_{\Bbbk X} \mod$ and $N \in {}_{\Bbbk Y} \mod$. Then one has an isomorphism

$$\operatorname{Ind}_{X}^{G \times H}(M) \otimes_{\Bbbk H} \operatorname{Ind}_{Y}^{H \times K}(N) \cong \bigoplus_{t \in [p_{2}(X) \setminus H/p_{1}(Y)]} \operatorname{Ind}_{X*}^{G \times K}(M \overset{X, \overset{(\cdot, \cdot)}{\otimes}}{\underset{\Bbbk H}{\otimes}} (t, 1)N)$$

of $(\Bbbk G, \Bbbk H)$ -bimodules.

In the sequel we need to establish a list of functorial properties of the extended construction of tensor products and homomorphism functors. The following proposition generalizes the usual adjunction between the tensor product and homomorphism functors.

6.3 Proposition Let G, H, K and L be finite groups, let $X \leq G \times H$, $Y \leq H \times K$ and $Z \leq G \times L$, and let $M \in {}_{\Bbbk X} \mathsf{mod}$, $N \in {}_{\Bbbk Y} \mathsf{mod}$ and $P \in {}_{\Bbbk Z} \mathsf{mod}$. Then there exists an isomorphism

$$\operatorname{LHom}_{\Bbbk G}^{X*Y,Z}\left(M \underset{\Bbbk H}{\overset{X,Y}{\otimes}} N, P\right) \cong \operatorname{LHom}_{\Bbbk H}^{Y,X^{\circ}*Z}\left(N, \operatorname{LHom}_{\Bbbk G}^{X,Z}(M,P)\right)$$

of $\mathbb{k}[(X * Y)^{\circ} * Z]$ -modules which is functorial in M, N and P.

Proof It is a straightforward verification that the functions

$$\operatorname{Hom}_{\Bbbk[k_1(X*Y)\cap k_1(Z)]} \left(M \otimes_{\Bbbk[k_2(X)\cap k_1(Y)]} N, P \right) \leftrightarrow \operatorname{Hom}_{\Bbbk[k_1(Y)\cap k_1(X^\circ*Z)]} \left(N, \operatorname{Hom}_{\Bbbk[k_1(X)\cap k_1(Z)]} (M, P) \right) \\ f \mapsto \left(n \mapsto \left(m \mapsto f(m \otimes n) \right) \right) \\ \left(m \otimes n \mapsto (f'(n))(m) \right) \leftrightarrow f'$$

are well-defined, mutually inverse homomorphisms of $k[(X * Y)^{\circ} * Z]$ -modules and natural in M, N and P.

A similar adjunction isomorphism exists for RHom and can be deduced from the above proposition via the functor $-\tau$, see 6.1(d).

In the special case where $Y = X^{\circ}$ and $Z = X * X^{\circ}$, the following proposition gives a different type of adjunction. Recall from Lemma 2.2(b) that $X * X^{\circ} * X = X$ and $X^{\circ} * X * X^{\circ} = X^{\circ}$.

6.4 Proposition Let $X \in G \times H$, $M \in {}_{\Bbbk X} \mod$, $N \in {}_{\Bbbk X^{\circ}} \mod$, and $P \in {}_{\Bbbk [X * X^{\circ}]} \mod$.

(a) There exists an isomorphism

$$\operatorname{Hom}_{\Bbbk[X*X^{\circ}]}(M \underset{\Bbbk H}{\overset{X,X^{\circ}}{\otimes}} N, P) \cong \operatorname{Hom}_{\Bbbk X^{\circ}}(N, \operatorname{LHom}_{\Bbbk G}^{X,X*X^{\circ}}(M, P))$$

of k-modules which is natural in M, N and P.

(b) There exists an isomorphism

$$\operatorname{Hom}_{\Bbbk[X*X^{\circ}]}(M \underset{\Bbbk H}{\overset{X,X^{\circ}}{\otimes}} N, P) \cong \operatorname{Hom}_{\Bbbk X}(M, \operatorname{RHom}_{\Bbbk G}^{X^{\circ}, X*X^{\circ}}(N, P))$$

of k-modules which is natural in M, N and P.

Proof (a) Again, it is straightforward to verify that the maps

$$\operatorname{Hom}_{\Bbbk[X*X^{\circ}]}(M \otimes_{\Bbbk k_{2}(X)} N, P) \leftrightarrow \operatorname{Hom}_{\Bbbk X^{\circ}}(N, \operatorname{Hom}_{\Bbbk k_{1}(X)}(M, P))$$
$$f \mapsto (n \mapsto (m \mapsto f(m \otimes n)))$$
$$(m \otimes n \mapsto (f'(n))(m)) \leftarrow f'$$

are well-defined, mutually inverse k-module homomorphisms which are natural in M, N and P.

(b) The maps analogous to the ones in (a) give again the desired isomorphisms.

The following lemma generalizes a well-known compatibility of the tensor product and induction in the special case that G = H = K, $X = Y = \Delta(G)$.

6.5 Lemma Let $X' \leq X \leq G \times H$ and $Y' \leq Y \leq H \times K$.

(a) Let $M' \in {}_{\Bbbk X'}$ mod and $N \in {}_{\Bbbk Y}$ mod. There exists a ${}_{\Bbbk}[X * Y]$ -module homomorphism

$$\alpha_1 \colon \operatorname{Ind}_{X'*Y}^{X*Y}(M' \underset{\Bbbk H}{\overset{X',Y}{\otimes}} N) \to \operatorname{Ind}_{X'}^X(M') \underset{\Bbbk H}{\overset{X,Y}{\otimes}} N$$

П

which maps $(g,k) \otimes (m' \otimes n)$ to $((g,h) \otimes m') \otimes (h,k)n$, for $(g,k) \in X * Y$, $m' \in M'$ and $n \in N$, where $h \in H$ is chosen such that $(g,h) \in X$ and $(h,k) \in Y$. The homomorphism α_1 is functorial in M' and N, and if $p_2(X) \leq p_1(Y)$ then α_1 is an isomorphism.

(b) Let $M \in {}_{\Bbbk X}$ mod and $N' \in {}_{\Bbbk Y'}$ mod. There exists a ${}_{\Bbbk}[X * Y]$ -module homomorphism

$$\alpha_2 \colon \operatorname{Ind}_{X*Y'}^{X*Y}(M \underset{\Bbbk H}{\overset{X,Y'}{\otimes}} N') \to M \underset{\Bbbk H}{\overset{X,Y}{\otimes}} \operatorname{Ind}_{Y'}^Y(N')$$

which maps $(g,k) \otimes (m \otimes n')$ to $(g,h)m \otimes ((h,k) \otimes n')$, for $(g,k) \in X * Y$, $m \in M$, and $n' \in N'$, where $h \in H$ is chosen such that $(g,h) \in X$ and $(h,k) \in Y$. The homomorphism α_2 is functorial in M and N', and if $p_2(X) \ge p_1(Y)$ then it is an isomorphism.

Proof We only prove Part (a). Part (b) is proved similarly. It is straightforward to verify that the map α_1 is well-defined and a natural $\mathbb{K}[X * Y]$ -module homomorphism. Now assume that $p_2(X) \leq p_1(Y)$. It is again straightforward to verify that one obtains a well-defined $\mathbb{K}[X * Y]$ -module homomorphism $\beta_1 \colon \operatorname{Ind}_{X'}^X(M') \bigotimes_{\mathbb{K}H}^{X,Y} N \to \operatorname{Ind}_{X'*Y}^{X',Y}(M' \bigotimes_{\mathbb{K}H}^{X',Y} N)$ by mapping $((g,h) \otimes m') \otimes n$ to $(g,k) \otimes (m' \otimes (h,k)^{-1}n)$, for $(g,h) \in X$, $m' \in M'$, and $n \in N$, were $k \in K$ is chosen such that $(h,k) \in Y$ (using $p_2(X) \leq p_1(Y)$). It is obvious that α_1 and β_1 are inverses.

 $\begin{array}{ll} \textbf{6.6} \quad \text{Let } X \leqslant G \times H \text{ and } M \in {}_{\Bbbk X} \textbf{mod.} \text{ Recall from Lemma 2.2(a) that } X \ast X^{\circ} = (k_1(X) \times \{1\}) \Delta(p_1(X)) = \\ \{(g,g') \in p_1(X) \times p_1(X) \mid gk_1(X) = g'k_1(X)\}. \text{ Moreover, } \Bbbk[k_1(X)] \text{ can be considered as left } \Bbbk[X \ast X^{\circ}]\text{-module } \\ \text{via } (g,g')n := gng'^{-1}, \text{ for } n \in k_1(X) \text{ and } (g,g') \in X \ast X^{\circ}. \text{ Thus, we obtain a } \Bbbk[X]\text{-module } \Bbbk[k_1(X)] \overset{X \ast X^{\circ}, X}{\underset{\& G}{\otimes}} M \\ \text{and a } \Bbbk[X^{\circ}]\text{-module LHom}_{\& G}^{X,X \ast X^{\circ}}(M, \Bbbk[k_1(X)]), \text{ since } X \ast X^{\circ} \ast X = X \text{ and } X^{\circ} \ast X \ast X^{\circ} = X^{\circ} \text{ (see Lemma 2.2(b))}. \\ \text{Similarly, } \Bbbk[k_2(X)] \text{ is a left } \Bbbk[X^{\circ} \ast X]\text{-module via } (h, h')n = hnh'^{-1} \text{ for } n \in k_2(X) \text{ and } (h, h') \in X^{\circ} \ast X. \end{array}$

Thus, one obtains a $\Bbbk[X]$ -module $M \overset{X,X^\circ*X}{\underset{\Bbbk H}{\otimes}} \Bbbk[k_2(X)]$ and a $\Bbbk[X^\circ]$ -module $\operatorname{RHom}_{\Bbbk H}^{X,X^\circ*X}(M,\Bbbk[k_2(X)])$.

We leave the straightforward proof of the following proposition to the reader.

6.7 Proposition Let $X \leq G \times H$ and $M \in {}_{\Bbbk X} \mathsf{mod}$.

(a) The map

$$M^{\circ} \to \operatorname{LHom}_{\Bbbk G}^{X, X * X^{\circ}}(M, \Bbbk[k_1(X)]), \quad \lambda \mapsto \left(m \mapsto \sum_{g \in k_1(X)} \lambda(g^{-1}m)g\right)$$

is a well-defined isomorphism of $\Bbbk X^{\circ}$ -modules. Its inverse maps the homomorphism f to $(m \mapsto t(f(m)))$, where t denotes the \Bbbk -linear extension of $k_1(X) \to \Bbbk$, $g \mapsto \delta_{g,1}$.

(b) The map

$$M^{\circ} \to \operatorname{RHom}_{\Bbbk H}^{X,X^{\circ}*X}(M, \Bbbk[k_2(X)])\,, \quad \lambda \mapsto \left(m \mapsto \sum_{h \in k_2(X)} \lambda(mh^{-1})h\right),$$

is a well-defined isomorphism of $\Bbbk X^{\circ}$ -modules. Its inverse maps f to $(m \mapsto t(f(m)))$.

(c) The maps $a \otimes m \mapsto am$ and $m \otimes a \mapsto ma$ define $\Bbbk[X]$ -module isomorphisms

$$\Bbbk[k_1(X)] \overset{X \ast X^{\circ}, X}{\underset{\Bbbk G}{\otimes}} M \to M \quad and \quad M \overset{X, X^{\circ} \ast X}{\underset{\Bbbk H}{\otimes}} \Bbbk[k_2(X)] \to M$$

with inverses given by $m \mapsto 1 \otimes m$ and $m \mapsto m \otimes 1$.

(d) Assume that $K \leq k_1(X)$ with $K \leq p_1(X)$, that $e \in Z(\Bbbk[k_1(X)])^{p_1(X)}$ is an idempotent, and that $M \in {}_{\Bbbk X(e \otimes 1)} \mod$. Then one has an isomorphism of $\Bbbk X$ -modules

$$\operatorname{Inf}_{X/(K\times\{1\})}^{X}\operatorname{Def}_{X/(K\times\{1\})}^{X}(M) = \Bbbk[X/(K\times\{1\})] \otimes_{\Bbbk X} M \cong \Bbbk[k_{1}(X)/K] \overline{e} \overset{X*X^{\circ},X}{\underset{\Bbbk G}{\otimes}} M,$$

given by $(g,h)(K \times \{1\}) \otimes m \mapsto \overline{e} \otimes (g,h)m$ with inverse $(gK)\overline{e} \otimes m \mapsto 1 \otimes (g,1)m$, where \overline{e} is the image of e under the canonical k-algebra homomorphism $\mathbb{k}[k_1(X)] \to \mathbb{k}[k_1(X)/K]$ and where $\mathbb{k}[k_1(X)/K]\overline{e}$ is a $\mathbb{k}[X * X^\circ]$ -module via $(g_1, g_2) \cdot a = g_1 a g_2^{-1}$ for $(g_1, g_2) \in X * X^\circ$ and $a \in \mathbb{k}[k_1(X)/K]\overline{e}$.

6.8 Proposition (a) Let $X \leq G \times H$, $Y \leq G \times K$, $M \in {}_{\Bbbk X} \mathsf{mod}$, and $N \in {}_{\Bbbk Y} \mathsf{mod}$. The map

$$\alpha_1 \colon \operatorname{LHom}_{\Bbbk G}^{X,X*X^{\circ}}(M, \Bbbk[k_1(X)]) \underset{\Bbbk G}{\overset{X^{\circ},Y}{\otimes}} N \to \operatorname{LHom}_{\Bbbk G}^{X,Y}(M,N), \quad f \otimes n \mapsto \left(m \mapsto t(f(m))n\right).$$

is a well-defined $\Bbbk[X^{\circ} * Y]$ -module homomorphism and functorial in M and N. Here t denotes the canonical projection map $\Bbbk[k_1(X)] \to \Bbbk[k_1(X) \cap k_1(Y)]$. If M is projective as left $\Bbbk[k_1(X)]$ -module then α_1 is an isomorphism.

(b) Let $X \leq G \times H$, $Y \leq K \times H$, $M \in {}_{\Bbbk X} \mathsf{mod}$, and $N \in {}_{\Bbbk Y} \mathsf{mod}$. The map

$$\alpha_2 \colon N \bigotimes_{\Bbbk H}^{Y, X^{\circ}} \operatorname{RHom}_{\Bbbk H}^{X, X^{\circ} * X}(M, \Bbbk[k_2(X)]) \to \operatorname{RHom}_{\Bbbk H}^{X, Y}(M, N), \quad n \otimes f \mapsto \left(m \mapsto nt(f(m))\right),$$

is a well-defined $\Bbbk[Y * X^{\circ}]$ -module homomorphism and functorial in M and N. Here, t denotes the canonical projection map $\Bbbk[k_2(X)] \to \Bbbk[k_2(X) \cap k_2(Y)]$. If M is projective as right $\Bbbk[k_2(X)]$ -module then α_2 is an isomorphism.

Proof (a) Set $A := k_1(X) \cap k_1(Y)$ and $B := k_1(X)$. It is straightforward to verify that

 $\alpha_1 \colon \operatorname{Hom}_{\Bbbk B} (M, \Bbbk B) \otimes_{\Bbbk A} N \to \operatorname{Hom}_{\Bbbk A} (M, N) \,, \quad (f \otimes n) \mapsto (m \mapsto t(f(m))n) \,,$

is well-defined and functorial in M and N. A careful but straightforward computation also shows that α_1 is a $\Bbbk[X^\circ * Y]$ -module homomorphism. Next assume that M is projective as left $\Bbbk B$ -module. Note that, for fixed N, the map α_1 is also a natural transformation between two additive contravariant functors $\Bbbk_B \mod \to \Bbbk \mod A$. Thus, in order to show that α_1 is an isomorphism it suffices to show this when $M = \Bbbk B$. Using the natural isomorphism $\Bbbk B \to \operatorname{Hom}_{\Bbbk B}(\Bbbk B, \Bbbk B), b \mapsto \rho_b$, with $\rho_b(b') = b'b$ for $b, b' \in B$, it suffices to show that the \Bbbk -linear map

$$\tilde{\alpha}_1 \colon \Bbbk B \otimes_{\Bbbk A} N \to \operatorname{Hom}_{\Bbbk A}(\Bbbk B, N), \quad b \otimes n \mapsto (b' \mapsto t(b'b)n)$$

is bijective. But this is easily verified: If $b_1, \ldots, b_d \in B$ are representatives of B/A, then the map

$$\tilde{\beta}_1 \colon \operatorname{Hom}_{\Bbbk A}(\Bbbk B, N) \to \Bbbk B \otimes_{\Bbbk A} N, \quad f' \mapsto \sum_{i=1}^d b_i \otimes f'(b_i^{-1}),$$

is an inverse to $\tilde{\alpha}_1$.

(b) This is proved in a similar way as Part (a).

6.9 Corollary Let $X \leq G \times H$, $M \in {}_{\Bbbk X} \mod$, $N \in {}_{\Bbbk X^{\circ}} \mod$, and $P \in {}_{\Bbbk [X * X^{\circ}]} \mod$.

(a) If M is projective as left $k[k_1(X)]$ -module then one has a k-module isomorphism

$$\operatorname{Hom}_{\Bbbk[X*X^{\circ}]}\left(M \underset{\Bbbk H}{\overset{X,X^{\circ}}{\otimes}} N, P\right) \cong \operatorname{Hom}_{\Bbbk X^{\circ}}\left(N, M^{\circ} \underset{\Bbbk G}{\overset{X^{\circ},X*X^{\circ}}{\otimes}} P\right).$$

(b) If N is projective as right $k[k_1(X)]$ -module then one has a k-module isomorphism

$$\operatorname{Hom}_{\Bbbk[X*X^{\circ}]}\left(M \underset{\Bbbk H}{\overset{X,X^{\circ}}{\otimes}} N, P\right) \cong \operatorname{Hom}_{\Bbbk X}\left(M, P \underset{\Bbbk G}{\overset{X*X^{\circ},X}{\otimes}} N^{\circ}\right)$$

Proof This follows immediately from Propositions 6.4, 6.8, and 6.7.

The following lemma will be used in the proof of Lemma 10.4. Recall the definition of $N_{(S,\phi,T)}$ from 2.3.

6.10 Lemma Let $\phi: Q \xrightarrow{\sim} P$ be an isomorphism between subgroups $Q \leq H$ and $P \leq G$. Furthermore, let $C_G(P) \leq S \leq N_G(P)$ and $C_H(Q) \leq T \leq N_H(Q)$ be intermediate groups. Set $X := N_{S \times S}(\Delta(P))$, $X' := \Delta(N_{(S,\phi,T)})(C_G(P) \times \{1\}) \leq G \times G$, and $Y := N_{S \times T}(\Delta(P,\phi,Q)) \leq G \times H$.

(a) One has $Y * Y^{\circ} = X' \leq X = X * X$, X * X' = X' and X * Y = X' * Y = Y.

(b) Let $M \in {}_{\Bbbk Y} \mod$, $N \in {}_{\Bbbk Y} \mod$, and $V, W \in {}_{\Bbbk X} \mod$. If N is projective as right $\Bbbk[C_G(P)]$ -module then one has a \Bbbk -linear isomorphism

$$\operatorname{Hom}_{\Bbbk Y} \left(V \underset{\Bbbk G}{\overset{X,Y}{\otimes}} M, W \underset{\Bbbk G}{\overset{X,Y}{\otimes}} N^{\circ} \right) \cong \operatorname{Hom}_{\Bbbk X} \left(V \underset{\Bbbk G}{\overset{X,X}{\otimes}} \operatorname{Ind}_{X'}^{X} (M \underset{\Bbbk H}{\overset{Y,Y^{\circ}}{\otimes}} N), W \right).$$

Proof (a) All assertions follow immediately from Lemma 2.2 and Proposition 2.4.

(b) By Part (a), Lemma 6.5(b), and the associativity property in 6.1(a) we obtain isomorphisms

$$V \underset{\Bbbk G}{\overset{X,X}{\otimes}} \mathrm{Ind}_{X'}^X (M \underset{\Bbbk H}{\overset{Y,Y^\circ}{\otimes}} N) \cong \mathrm{Ind}_{X*X'}^{X*X} \big(V \underset{\Bbbk G}{\overset{X,X'}{\otimes}} (M \underset{\Bbbk H}{\overset{Y,Y^\circ}{\otimes}} N) \big) \cong \mathrm{Ind}_{X'}^X \big((V \underset{\Bbbk G}{\overset{X,Y}{\otimes}} M) \underset{\Bbbk H}{\overset{X*Y,Y^\circ}{\otimes}} N \big)$$

of &X-modules. Thus, using again Part (a), the usual adjunction of Ind and Res, and Corollary 6.9(b), we obtain isomorphisms

$$\operatorname{Hom}_{\Bbbk X}\left(V \underset{\Bbbk G}{\overset{X,X}{\otimes}} \operatorname{Ind}_{X'}^{X} \left(M \underset{\Bbbk H}{\overset{Y,Y^{\circ}}{\otimes}} N\right), W\right) \cong \operatorname{Hom}_{\Bbbk X}\left(\operatorname{Ind}_{X'}^{X}\left(\left(V \underset{\Bbbk G}{\overset{X,Y}{\otimes}} M\right) \underset{\Bbbk H}{\overset{X*Y,Y^{\circ}}{\otimes}} N\right), W\right) \cong$$
$$\cong \operatorname{Hom}_{\Bbbk X'}\left(\left(V \underset{\Bbbk G}{\overset{X,Y}{\otimes}} M\right) \underset{\Bbbk H}{\overset{X*Y,Y^{\circ}}{\otimes}} N, \operatorname{Res}_{X'}^{X}(W)\right) \cong \operatorname{Hom}_{\Bbbk Y}\left(V \underset{\Bbbk G}{\overset{X,Y}{\otimes}} M, \operatorname{Res}_{X'}^{X}(W) \underset{\Bbbk G}{\overset{X,Y}{\otimes}} N\right)$$

of k-modules, since N is projective as right module for $\Bbbk[C_G(P)] = \Bbbk[k_1(Y^\circ)]$, see Proposition 2.4(c). Finally, $\operatorname{Res}_{X'}^X(W) \overset{X',Y}{\underset{\Bbbk G}{\otimes}} N^\circ = W \overset{X,Y}{\underset{\Bbbk G}{\otimes}} N^\circ$ as $\Bbbk[Y]$ -modules, since X' * Y = X * Y = Y by Part (a) and since $k_2(X') = C_G(P) = k_2(X)$ by Proposition 2.4(c). This completes the proof.

7 Tensor products of *p*-permutation bimodules

Throughout this section, G, H and K denote finite groups.

7.1 Let k be a commutative ring. The following observation will allow us to create situations where α_1 and α_2 in Lemma 6.5 are isomorphisms. Let $X \leq G \times H$, $Y \leq H \times K$, $M \in {}_{\Bbbk X} \mod$, and $N \in {}_{\Bbbk Y} \mod$. We define

$$\tilde{X} := \{ (g, h) \in X \mid \exists k \in K \colon (h, k) \in Y \}.$$

Then $\tilde{X} \leq X$, $p_2(\tilde{X}) \leq p_1(Y)$, $k_2(\tilde{X}) \cap k_1(Y) = k_2(X) \cap k_1(Y)$, and $\tilde{X} * Y = X * Y$. Moreover, one has $\operatorname{Res}_{\tilde{X}}^X(M) \overset{\tilde{X},Y}{\underset{\Bbbk H}{\otimes}} N = M \overset{X,Y}{\underset{\Bbbk H}{\otimes}} N$ as $\Bbbk[X * Y]$ -modules. Similarly, one can define $\tilde{Y} \leq Y$ with the analogous properties. The use of this method is illustrated in the proof of the following lemma.

For the remainder of this section we assume that \mathcal{O} is a complete discrete valuation ring of characteristic 0 with residue field F of characteristic p > 0.

7.2 Lemma Let $X \leq G \times H$ and $Y \leq H \times K$.

(a) If $M \in \mathcal{O}_X$ triv and $N \in \mathcal{O}_Y$ triv then $M \bigotimes_{\mathcal{O}_H}^{X,Y} N \in \mathcal{O}_{[X*Y]}$ triv.

(b) If $M \in \mathcal{O}_X \mod$ and $N \in \mathcal{O}_Y \mod$ are indecomposable with twisted diagonal vertices then each indecomposable direct summand of the $\mathcal{O}[X * Y]$ -module $M \bigotimes_{\mathcal{O}_H}^{X,Y} N$ has twisted diagonal vertices.

Proof (a) This can easily be seen by using a tensor product construction on bisets. We give a different proof to illustrate the method from 7.1. Since $\operatorname{Res}_{\tilde{X}}^{X}(M)$ is again a *p*-permutation module, we may assume that $p_2(X) \leq p_1(Y)$. Next, since $\bigotimes_{\mathcal{O}H}^{X,Y}$ respects direct sums, we may assume that $M = \operatorname{Ind}_{X'}^{X}(\mathcal{O}_{X'})$ for some subgroup X' of X. Using Lemma 6.5(a), we obtain $M \bigotimes_{\mathcal{O}H}^{\otimes} N \cong \operatorname{Ind}_{X'*Y}^{X*Y}(\mathcal{O}_{X'} \bigotimes_{\mathcal{O}H}^{\otimes} N)$. Since the class of *p*-permutation modules is stable under induction, we may assume that X = X' and $M = \mathcal{O}_X$. Similar arguments for Y and N reduce further to the case that $N = \mathcal{O}_Y$. But in this case we have $M \bigotimes_{\mathcal{O}H}^{X,Y} N = \mathcal{O}_{X*Y}$, the trivial module, which is a *p*-permutation module.

(b) Since $\bigotimes_{\substack{\mathcal{O}H\\\mathcal{O}H}}^{X,Y}$ respects direct sums and by 3.2(d), we may use 7.1 to reduce to the case where $p_2(X) \leq p_1(Y)$. Since $\bigotimes_{\substack{\mathcal{O}H\\\mathcal{O}H}}^{X,Y}$ respects direct sums, we may also assume that $M = \operatorname{Ind}_{X'}^X(M')$ for some twisted diagonal

subgroup X' of X and some indecomposable module $M' \in _{\mathcal{O}X'}$ mod. Lemma 6.5(a) now implies that $M \bigotimes_{\mathcal{O}H}^{X,Y} N \cong$ Ind $_{X'*Y}^{X*Y}(M' \bigotimes_{\mathcal{O}H}^{X',Y} N)$, and by 3.2(d) we may assume that X is twisted diagonal. Similar considerations for Y and N reduce further to the situation that also Y is twisted diagonal. But then also X * Y and its subgroups are twisted diagonal. This completes the proof.

Next we recall and improve Theorem 3.3 from [BD12]. For a fixed twisted diagonal *p*-subgroup $\Delta(P, \sigma, R) \leq G \times K$ we denote by $\Gamma = \Gamma_H(P, \sigma, R)$ the set of triples (ϕ, Q, ψ) , where $Q \leq H$ and $\psi \colon R \xrightarrow{\sim} Q$ and $\phi \colon Q \xrightarrow{\sim} P$ are isomorphisms with $\phi \circ \psi = \sigma$. The group *H* acts on $\Gamma_H(P, \sigma, R)$ by ${}^{h}(\phi, Q, \psi) \coloneqq (\phi c_h^{-1}, {}^{h}Q, c_h\psi)$. Note that stab_{*H*} $(\phi, Q, \psi) = C_H(Q)$. For $M \in {}_{OG}\mathsf{mod}_{\mathcal{O}H}$, $N \in {}_{OH}\mathsf{mod}_{\mathcal{O}K}$, and $(\phi, Q, \psi) \in \Gamma_H(P, \sigma, Q)$, one has an $(F[C_G(P)], F[C_K(R)])$ -bimodule homomorphism

$$\Phi_{(\phi,Q,\psi)} \colon M(\Delta(P,\phi,Q)) \otimes_{F[C_H(Q)]} N(\Delta(Q,\psi,R)) \to (M \otimes_{\mathcal{O}H} N)(\Delta(P,\sigma,R)),$$

$$\operatorname{Br}_{\Delta(P,\phi,Q)}(m) \otimes \operatorname{Br}_{\Delta(Q,\psi,R)}(n) \mapsto \operatorname{Br}_{\Delta(P,\sigma,R)}(m \otimes n),$$
(5)

see [BD12, 3.1(h)], which is natural in M and N. The following theorem describes $(M \otimes_{\mathcal{O}H} N)(\Delta(P, \sigma, R))$ as $(F[C_G(P)], F[C_K(R)])$ -bimodule, for a particular class of modules M and N.

7.3 Theorem ([BD12, Theorem 3.3]) Let $\Delta(P, \sigma, R) \leq G \times K$ be a twisted diagonal *p*-subgroup, let $\Gamma = \Gamma_H(P, \sigma, R)$ be as above and let $\widetilde{\Gamma} \subseteq \Gamma$ be a set of representatives of the *H*-orbits of Γ . Furthermore let $M \in {}_{\mathcal{O}G}$ triv_{$\mathcal{O}H$} and $N \in {}_{\mathcal{O}H}$ triv_{$\mathcal{O}K$} be *p*-permutation bimodules all of whose indecomposable direct summands have twisted diagonal vertices. Then the direct sum of the homomorphisms $\Phi_{(\phi,Q,\psi)}$, $(\phi,Q,\psi) \in \widetilde{\Gamma}$ yields an isomorphism

$$\Phi: \bigoplus_{(\phi,Q,\psi)\in\widetilde{\Gamma}} M(\Delta(P,\phi,Q)) \otimes_{F[C_H(Q)]} N(\Delta(Q,\psi,R)) \to (M \otimes_{\mathcal{O}H} N)(\Delta(P,\sigma,R))$$
(6)

of $(F[C_G(P)], F[C_K(R)])$ -bimodules which is natural in M and N.

Note that not only $C_G(P) \times C_K(R)$, but also the bigger group $N_{G \times K}(\Delta(P, \sigma, R))$ acts on the right hand side of the isomorphism (6). The corollary below describes the $F[N_{G \times K}(\Delta(P, \sigma, R))]$ -module structure of $(M \otimes_{\mathcal{O}H} N)(\Delta(P, \sigma, R))$. First note that the domain of the homomorphism $\Phi_{(\phi,Q,\psi)}$ in (5) carries an $F[N_{G \times H}(\Delta(P, \phi, Q)) * N_{H \times K}(\Delta(Q, \psi, R))]$ -module structure via the extended tensor product construction in 6.1. This module structure extends the $F[C_G(P)] \times F[C_K(R)]$ -module structure from (5), since $k_1(N_{G \times H}(\Delta(P, \phi, Q))) = C_G(P), \ k_2(N_{G \times H}(\Delta(P, \phi, Q))) = C_H(Q) = k_1(N_{H \times K}(\Delta(Q, \psi, R)))$ and $k_2(N_{H \times K}(\Delta(Q, \psi, R))) = C_K(R)$ by Proposition 2.4(b). It is a straightforward verification that $\Phi_{(\phi,Q,\psi)}$ in (5) actually defines a homomorphism

$$\Phi_{(\phi,Q,\psi)}: M(\Delta(P,\phi,Q)) \xrightarrow{N_{G \times H}(\Delta(P,\phi,Q)), N_{H \times K}(\Delta(Q,\psi,R))}_{FH} \to (M \otimes_{\mathcal{O}H} N)(\Delta(P,\sigma,R))$$
(7)

of $F[N_{G \times H}(\Delta(P, \phi, Q)) * N_{H \times K}(\Delta(Q, \psi, R))]$ -modules.

7.4 Corollary Let $M \in {}_{\mathcal{O}G}\mathsf{triv}_{\mathcal{O}H}, N \in {}_{\mathcal{O}H}\mathsf{triv}_{\mathcal{O}K}, (P, \sigma, R) \leq G \times K$, and $\widetilde{\Gamma} \subseteq \Gamma = \Gamma_H(P, \sigma, R)$ be as in Theorem 7.3.

(a) The group $N_{G \times K}(\Delta(P, \sigma, R)) \times H$ acts on Γ via $((g,k),h)(\phi, Q, \psi) := (c_g \phi c_h^{-1}, {}^hQ, c_h \psi c_k^{-1})$. For the induced action of $N_{G \times K}(\Delta(P, \sigma, R))$ on the H-orbits $[\phi, Q, \psi]_H$ of Γ one has $\operatorname{stab}_{N_{G \times K}(\Delta(P, \sigma, R))}([\phi, Q, \psi]_H) = N_{G \times H}(\Delta(P, \phi, Q)) * N_{H \times K}(\Delta(Q, \psi, R))$, for each $(\phi, Q, \psi) \in \Gamma$.

(b) Let $\widehat{\Gamma} \subseteq \widetilde{\Gamma}$ be a set of representatives of the $N_{G \times K}(\Delta(P, \sigma, R)) \times H$ -orbits of Γ . The homomorphisms $\Phi_{(\phi, Q, \psi)}, (\phi, Q, \psi) \in \widehat{\Gamma}$, in (7) induce an isomorphism

$$\Phi: \bigoplus_{\gamma=(\phi,Q,\psi)\in\widehat{\Gamma}} \operatorname{Ind}_{X(\gamma)*Y(\gamma)}^{N_{G\times K}(\Delta(P,\sigma,R))} \Big(M(\Delta(P,\phi,Q)) \overset{X(\gamma),Y(\gamma)}{\underset{FH}{\otimes}} N(\Delta(Q,\psi,R)) \Big) \xrightarrow{\sim} (M \otimes_{\mathcal{O}H} N)(\Delta(P,\sigma,R))$$

of $F[N_{G \times K}(\Delta(P, \sigma, R))]$ -modules which is natural in M and N, with $X(\gamma) := N_{G \times H}(\Delta(P, \phi, Q))$ and $Y(\gamma) := N_{H \times K}(\Delta(Q, \psi, R))$, for $\gamma = (\phi, Q, \psi) \in \Gamma$.

Proof (a) The first statement is clear and the second statement is a straightforward verification.

(b) First note that, for $(\phi, Q, \psi) \in \Gamma$, one has $X(\gamma) * Y(\gamma) \leq N_{G \times K}(\Delta(P, \sigma, R))$, by Part (a). Now let $(\phi, Q, \psi) \in \widetilde{\Gamma}, (g, k) \in N_{G \times K}(\Delta(P, \sigma, R)), m \in M^{\Delta(P, \phi, Q)}$, and $n \in N^{\Delta(P, \psi, R)}$. Then there exists $(\phi', Q', \psi') \in \widetilde{\Gamma}$ and $h \in H$ such that $((g, k), h)(\phi, Q, \psi) = (\phi', Q', \psi')$, and we have

$$\begin{aligned} &(g,k) \cdot \Phi_{(\phi,Q,\psi)} \big(\operatorname{Br}_{\Delta(P,\phi,Q)}(m) \otimes Br_{\Delta(Q,\psi,R)}(n) \big) = (g,k) \cdot \operatorname{Br}_{\Delta(P,\sigma,R)}(m \otimes n) = \operatorname{Br}_{\Delta(P,\sigma,R)}(gm \otimes nk^{-1}) \\ &= \operatorname{Br}_{\Delta(P,\sigma,R)}(gmh^{-1} \otimes hnk^{-1}) = \Phi_{(\phi',Q',\psi')} \big(\operatorname{Br}_{\Delta(P,\phi',Q')}(gmh^{-1}) \otimes \operatorname{Br}_{\Delta(Q',\psi',R)}(hnk^{-1}) \big) \,. \end{aligned}$$

This equation shows that if one transports the $F[N_{G \times K}(\Delta(P, \sigma, R))]$ -module structure from the right hand side of the isomorphism Φ in (6) via Φ^{-1} to the left hand side, then this action permutes the direct summands according to the action of $N_{G \times K}(\Delta(P, \sigma, R))$ on $H \setminus \Gamma \cong \widetilde{\Gamma}$. Moreover, the stabilizer of the (ϕ, Q, ψ) -component equals $N_{G \times H}(\Delta(P, \phi, Q)) * N_{H \times K}(\Delta(Q, \psi, R))$ by Part (a). Thus, the isomorphism in (6) defines the desired $F[N_{G \times K}(\Delta(P, \sigma, R))]$ -module isomorphism of Part (b).

In the next theorem we will give block-wise versions of Theorem 7.3 and Corollary 7.4. Note that for a twisted diagonal p-subgroup $\Delta(P, \phi, Q)$ of $G \times H$ one has $C_{G \times H}(\Delta(P, \phi, Q)) = C_G(P) \times C_H(Q)$ and that, for Brauer pairs (P, e) and (Q, f) of FG and FH, respectively, the pair $(\Delta(P, \phi, Q), e \otimes f^*)$ is a Brauer pair of $F[G \times H] \cong FG \otimes_F FH$.

7.5 Theorem Let (P, e) be a Brauer pair of FG and let (R, d) be a Brauer pair of FK. Suppose that $\sigma \colon R \xrightarrow{\sim} P$ is an isomorphism and let $C_G(P) \leq S \leq N_G(P, e)$ and $C_K(R) \leq T \leq N_K(R, d)$ be intermediate subgroups. Furthermore, let $\Omega := \Omega_H((P, e), \sigma, (R, d))$ denote the set of triples $(\phi, (Q, f), \psi)$, where (Q, f) is a Brauer pair of FH and $\psi \colon R \xrightarrow{\sim} Q$ and $\phi \colon Q \xrightarrow{\sim} P$ are isomorphisms such that $\sigma = \phi \circ \psi$. Finally, let $M \in \mathcal{O}_G \operatorname{triv}_{\mathcal{O}H}$ and $N \in \mathcal{O}_H \operatorname{triv}_{\mathcal{O}K}$ be p-permutation modules all of whose indecomposable direct summands have twisted diagonal vertices.

(a) The group $N_{G \times K}(\Delta(P, \sigma, R)) \times H$ acts on Ω via $((g,k),h)(\phi, (Q, f), \psi) = (c_g \phi c_h^{-1}, h(Q, f), c_h \psi c_k^{-1})$ and $\operatorname{stab}_H(\phi, (Q, f), \psi) = C_H(Q).$

(b) Let $\Omega \subseteq \Omega$ be a set of representatives of the *H*-orbits of Ω . One has an isomorphism. The restrictions $\Phi_{(\phi,(Q,f),\psi)}$ of $\Phi_{(\phi,Q,\psi)}$ in (5) to $eM(\Delta(P,\phi,Q))f \otimes_{F[C_H(Q)]} fN(\Delta(Q,\psi,R))d$ define an isomorphism

$$\Phi: \bigoplus_{(\phi,(Q,f),\psi)\in\widetilde{\Omega}} eM(\Delta(P,\phi,Q))f \otimes_{F[C_H(Q)]} fN(\Delta(Q,\psi,R))d \xrightarrow{\sim} e(M \otimes_{\mathcal{O}H} N)(\Delta(P,\sigma,R))d$$
(8)

of $(F[C_G(P)]e, F[C_K(R)]d)$ -bimodules, which is natural in M and N.

(c) Let $\widehat{\Omega} \subseteq \widetilde{\Omega} \subseteq \Omega$ be a set of representatives of the $N_{S\times T}(\Delta(P,\sigma,R)) \times H$ -orbits of Ω . Then the homomorphisms $\Phi_{(\phi,(Q,f),\psi)} : eM(\Delta(P,\phi,Q))f \otimes_{F[C_H(Q)]} fN(\Delta(Q,\psi,R))d \to e(M \otimes_{\mathcal{O}H} N)(\Delta(P,\sigma,R))d,$ $(\phi,(Q,f),\psi) \in \widehat{\Omega}$, induce an isomorphism

$$\Phi: \bigoplus_{\omega \in \widehat{\Omega}} \operatorname{Ind}_{X(\omega)*Y(\omega)}^{N_{S \times T}(\Delta(P,\sigma,R))} \left(M(\omega) \overset{X(\omega),Y(\omega)}{\underset{FH}{\otimes}} N(\omega) \right) \xrightarrow{\sim} e\left(M \otimes_{\mathcal{O}H} N \right) \left(\Delta(P,\sigma,R) \right) d \tag{9}$$

of $N_{S\times T}(\Delta(P,\sigma,R))$ -modules which is natural in M and N. Here, for $\omega = (\phi, (Q, f), \psi) \in \widehat{\Omega}$, we set $X(\omega) := N_{S\times H}(\Delta(P,\phi,Q), e \otimes f^*), Y(\omega) := N_{H\times T}(\Delta(Q,\psi,R), f \otimes d^*), M(\omega) := eM(\Delta(P,\phi,Q))f \in _{FX(\omega)}$ triv, $N(\omega) := fN(\Delta(Q,\psi,R))d \in _{FY(\omega)}$ triv.

(d) Assume that G = K, (P, e) = (R, d), S = T, and $\sigma = id_P$. Let $\Lambda := \Lambda_H(P)$ be the set of pairs $(\phi, (Q, f))$, where (Q, f) is a Brauer pair of FH and $\phi: Q \xrightarrow{\sim} P$ is an isomorphism. The group $S \times H$ acts on Λ via ${}^{(g,h)}(\phi, (Q, f)) = (c_g \phi c_h^{-1}, {}^h(Q, f))$, for $(g, h) \in S \times H$ and $(\phi, (Q, f)) \in \Lambda$. Let $\widehat{\Lambda} \subseteq \widetilde{\Lambda} \subseteq \Lambda$ be such that $\widetilde{\Lambda}$ (resp. $\widehat{\Lambda}$) is a set of representatives of the H-orbits (resp. $S \times H$ -orbits) of Λ . Then one has an isomorphism

$$e(M \otimes_{\mathcal{O}H} N)(\Delta(P))e \cong \bigoplus_{(\phi,(Q,f))\in\widetilde{\Lambda}} eM(\Delta(P,\phi,Q))f \otimes_{F[C_H(Q)]} fN(\Delta(Q,\phi^{-1},P))e$$
(10)

of $(F[C_G(P)]e, F[C_G(P)]e)$ -bimodules, and an isomorphism

$$e(M \otimes_{\mathcal{O}H} N)(\Delta(P))e \cong \bigoplus_{\lambda \in \widehat{\Lambda}} \operatorname{Ind}_{\Delta(I(\lambda))(C_G(P) \times \{1\})}^{N_{S \times S}(\Delta(P))} \left(M(\lambda) \bigotimes_{FH}^{X(\lambda), Y(\lambda)} N(\lambda)\right)$$
(11)

of $F[N_{S\times S}(\Delta(P))](e\otimes e^*)$ -modules. Here, for $\lambda = (\phi, (Q, f)) \in \Lambda$, we set $X(\lambda) := N_{S\times H}(\Delta(P, \phi, Q), e\otimes f^*)$, $Y(\lambda) := N_{H\times S}(\Delta(Q, \phi^{-1}, P), f\otimes e^*)$, $M(\lambda) := eM(\Delta(P, \phi, Q))f \in {}_{FX(\lambda)}$ triv, $N(\lambda) := fN(\Delta(Q, \phi^{-1}, P)e \in {}_{FY(\lambda)}$ triv, and $I(\lambda) := N_{(S,\phi,N_H(Q,f))}$.

Proof (a) This follows immediately from the definitions.

(b) Multiplication of the isomorphism in (6) with e from the left and d from the right yields again an isomorphism of $(F[C_G(P)]e, F[C_K(R)]d)$ -bimodules. For $(\phi, Q, \psi) \in \widetilde{\Gamma}$, the corresponding summand on the left hand side of this isomorphism can be written as the obvious direct sum over the block idempotents f of $F[C_H(Q)]$. Moreover, if (ϕ, Q, ψ) runs through $\widetilde{\Gamma}$ and, for each such (ϕ, Q, ψ) , f runs through the block idempotents of $FC_H(Q)$, then $(\phi, (Q, f), \psi)$ runs through a set of representatives of the H-orbits of Ω . This proves the claim for this particular set of representatives $\widetilde{\Omega}$ derived from $\widetilde{\Gamma}$. If $\widetilde{\widetilde{\Omega}}$ is an arbitrary set of representatives of the H-orbits of Ω , then for each $(\phi, (Q, f), \psi) \in \widetilde{\Omega}$ there exists a unique $(\phi', (Q', f'), \psi') \in \widetilde{\widetilde{\Omega}}$ and an element $h \in H$ such that ${}^{h}(\phi, (Q, f), \psi) = (\phi', (Q', f'), \psi')$. One obtains a well-defined isomorphism

$$\zeta_h \colon eM(\Delta(P,\phi,Q))f \otimes_{F[C_H(Q)]} fN(\Delta(Q,\psi,R))d \xrightarrow{\sim} eM(\Delta(P,\phi',Q'))f' \otimes_{F[C_H(Q')]} f'N(\Delta(Q',\psi',R))d$$

by mapping $\operatorname{Br}_{\Delta(P,\phi,Q)}(m) \otimes \operatorname{Br}_{\Delta(Q,\psi,R)}(n)$ to $\operatorname{Br}_{\Delta(P,\phi',Q')}(mh^{-1}) \otimes \operatorname{Br}_{\Delta(Q',\psi',R)}(hn)$, which is independent of the choice of h, such that $\Phi_{(\phi',(Q',f'),\psi')} \circ \zeta_h = \Phi_{(\phi,(Q,f),\psi)}$. This implies the statement in Part (b).

(c) Note that the right hand side of (8) is an $F[N_{S\times T}(\Delta(P,\sigma,R))]$ -module in a natural way. For $\omega = (\phi, (Q, f), \psi) \in \Omega$, we set $X'(\omega) := \Delta(P, \phi, Q)$ and $Y'(\omega) := \Delta(Q, \psi, R)$. Let $(g, k) \in N_{S\times T}(\Delta(P, \sigma, R))$ and $\omega = (\phi, (Q, f), \psi) \in \widetilde{\Omega}$. Then there exists a unique $\omega' = (\phi', (Q', f'), \psi') \in \widetilde{\Omega}$ and an element $h \in H$ such that ${}^{((g,k),h)}\omega =: \omega' \in \widetilde{\Omega}$. One verifies as in the proof of Corollary 7.4(b) that

$$(g,k)\Phi_{\omega}\left(e\mathrm{Br}^{M}_{X'(\omega)}(m)f\otimes f\mathrm{Br}^{N}_{Y'(\omega)}(n)d\right) = \Phi_{\omega'}\left(e\mathrm{Br}^{M}_{X'(\omega')}(gmh^{-1})f'\otimes f'\mathrm{Br}^{N}_{X'(\omega')}(hnk^{-1})d\right),\tag{12}$$

for $m \in M^{X'(\omega)}$ and $n \in N^{Y'(\omega)}$. This implies that if one transports the $FN_{S\times T}(\Delta(P,\sigma,R))$ -module structure of the right hand side of (8) via Φ^{-1} to the left hand side then $N_{S\times T}(\Delta(P,\sigma,R))$ permutes the components of the left hand side according to its action on the *H*-orbits of Ω . Moreover, it is straightforward to verify that $(g,k) \in N_{S\times T}(\Delta(P,\sigma,R))$ stabilizes the *H*-orbit of $\omega = (\phi, (Q,f), \psi)$ if and only if $(g,k) \in X(\omega) * Y(\omega)$. Equation (12) also implies that, for $\omega \in \tilde{\Omega}$, the $F[X(\omega) * Y(\omega)]$ -module structure of the ω -component of the left hand side of (8) coincides with the extended tensor product structure introduced in 6.1(a).

(d) Consider the bijection $\alpha \colon \Omega = \Omega_H((P, e), \operatorname{id}_P, (P, e)) \to \Lambda_H(P), (\phi, (Q, f), \phi^{-1}) \mapsto (\phi, (Q, f))$ and the group homomorphism $\kappa \colon N_{S \times S}(\Delta(P)) \times H \mapsto S \times H, ((s_1, s_2), h) \mapsto (s_1, h)$. Recall from Proposition 2.4(a) that $N_{S \times S}(\Delta(P)) = \Delta(S) \cdot (\{1\} \times C_G(P))$ and note that $\{1\} \times C_G(P)$ acts trivially on Ω . Therefore, one has

$$\alpha({}^{((s_1,s_2),h)}\!\omega) = (c_{s_1}\phi c_h^{-1}, {}^h\!(Q,f)) = {}^{\kappa((s_1,s_2),h)}\!\alpha(\omega) \,,$$

for $\omega = (\phi, (Q, f), \phi^{-1}) \in \Omega$ and $((s_1, s_2), h) \in N_{S \times S}(\Delta(P)) \times H$. Thus, the bijection α maps $\widetilde{\Omega}$ (resp. $\widehat{\Omega}$) to a set of representatives of the *H*-orbits (resp. $S \times H$ -orbits) of Λ . Now, the isomorphisms in (10) and (11) are immediate consequences of the isomorphisms in (8) and (9), after noting that $X(\lambda) * Y(\lambda) = \Delta(I(\lambda))(C_G(P) \times \{1\})$, since $Y(\lambda) = X(\lambda)^\circ$, see Proposition 2.4(c) and Lemma 2.2(a).

8 Character groups and perfect isometries

Throughout this section, G, H, K denote finite groups. We assume that the *p*-modular system (\mathbb{K} , \mathcal{O} , F) is large enough for G, H, K, and the groups H_1, \ldots, H_n appearing in Lemma 8.6 and Corollary 8.8. In this section, we recall and introduce notation, concepts, and basic results related to character groups and perfect isometries and we prove some results on perfect isometries that will be used in later sections.

For more details on the character group concepts of this section we refer the reader to [NT89, Section 3.6].

8.1 Notation (a) Let A be a finite-dimensional algebra over a field k. Recall that the *Grothendieck group* R(A), with respect to short exact sequences, is a free abelian group with \mathbb{Z} -basis given by elements [S], where S runs through a set of representatives of the isomorphism classes of simple left A-modules. For any $M \in {}_{A}$ mod one sets $[M] := [S_1] + \cdots + [S_n]$, where S_1, \ldots, S_n are the composition factors of M, (repeated according to their

multiplicities). If also B is a finite-dimensional algebra over the same field then we set $R(A, B) := R(A \otimes B^{\circ})$, where B° denotes the opposite algebra of B. This notation is motivated by the canonical category isomorphism ${}_{A}\mathsf{mod}_{B} \cong {}_{A \otimes B^{\circ}}\mathsf{mod}$. Thus, each $M \in {}_{A}\mathsf{mod}_{B}$ defines an element $[M] \in R(A, B)$. If B is a group algebra $\Bbbk H$, then we always identify $(\Bbbk H)^{\circ}$ with $\Bbbk H$ using the isomorphism $h^{\circ} \mapsto h^{-1}$. If additionally $A = \Bbbk G$ is a group algebra, we consequently identify $\Bbbk G \otimes (\Bbbk H)^{\circ}$ with $\Bbbk [G \times H]$ via $g \otimes h^{\circ} \mapsto (g, h^{-1})$.

(b) For an idempotent $e \in Z(\mathbb{K}G)$ we identify $R(\mathbb{K}Ge)$ with the virtual character group of $\mathbb{K}Ge$, the free \mathbb{Z} -span of the irreducible characters $\operatorname{Irr}(\mathbb{K}Ge)$ of $\mathbb{K}Ge$. This way, $R(\mathbb{K}Ge) \subseteq R(\mathbb{K}G)$. Similarly, if e is an idempotent in Z(FG) we identify R(FGe) with the group of virtual Brauer characters belonging to FGe. This way, $R(FGe) \subseteq R(FG)$. For convenience, we view Brauer characters throughout as class functions on G (rather than on $G_{p'}$) with values in \mathbb{K} that vanish on $G \smallsetminus G_{p'}$. Here, $G_{p'}$ denotes the set of p'-elements of G, i.e., elements whose order is not divisible by p. By scalar extension from \mathbb{Z} to \mathbb{K} we view these Grothendieck groups also as embedded into \mathbb{K} -vector spaces, denoted by $\mathbb{K}R(\mathbb{K}G)$, etc., and we identify $\mathbb{K}R(\mathbb{K}G)$ with the \mathbb{K} -vector space of \mathbb{K} -valued class functions on G, or by linear extension also as subspace of the space of \mathbb{K} -linear functions from $\mathbb{K}G$ to \mathbb{K} . In particular, we identify $\mathbb{K}R(FG)$ with the space of \mathbb{K} -valued class functions on G which vanish on $G \smallsetminus G_{p'}$. Note that for any idempotent $e \in Z(\mathcal{O}G)$ one has $\mathbb{K}R(FG\overline{e}) \subseteq \mathbb{K}R(\mathbb{K}Ge)$ as \mathbb{K} -vector spaces of function on G. In fact, since the determinant of the Cartan matrix of $FG\overline{e}$ is non-zero, each irreducible Brauer character in $FG\overline{e}$ is a \mathbb{Q} -linear combination of projective indecomposable characters of $\mathbb{K}Ge$.

If $e \in Z(\mathbb{K}G)$ and $f \in Z(\mathbb{K}H)$ are idempotents then, with the convention in (a), one has a group $R(\mathbb{K}Ge, \mathbb{K}Hf) := R(\mathbb{K}[G \times H](e \otimes f^*))$. Multiplication of $(\mathbb{K}G, \mathbb{K}H)$ -bimodules with e from the left and f from the right induces a projection map $R(\mathbb{K}G, \mathbb{K}H) \to R(\mathbb{K}Ge, \mathbb{K}Hf)$ which we denote by $\mu \mapsto e\mu f$. Taking \mathbb{K} -duals defines a map $R(\mathbb{K}Ge, \mathbb{K}Hf) \to R(\mathbb{K}Hf, \mathbb{K}Ge), \mu \mapsto \mu^{\circ}$, for idempotents $e \in Z(\mathbb{K}G)$ and $f \in Z(\mathbb{K}H)$. Similar notations apply with \mathbb{K} replaced by F.

(c) Tensor products of bimodules induce bilinear maps

$$R(\mathbb{K}G,\mathbb{K}H)\times R(\mathbb{K}H,\mathbb{K}K)\to R(\mathbb{K}G,\mathbb{K}K)\,,\quad (\mu,\nu)\mapsto \mu\underset{H}{\cdot}\nu\,,$$

and extended tensor products (see 6.1(a)) induce bilinear maps

$$R(\mathbb{K}X) \times R(\mathbb{K}Y) \to R(\mathbb{K}[X*Y]) \,, \quad (\mu,\nu) \mapsto \mu \stackrel{X,Y}{\stackrel{}{\overset{}_{H}}}_{\stackrel{}{H}} \nu \,,$$

for $X \leq G \times H$ and $Y \leq H \times K$. Each $\mu \in R(\mathbb{K}G, \mathbb{K}H)$, induces a group homomorphism

$$I_{\mu} \colon R(\mathbb{K}H) \to R(\mathbb{K}G), \quad \psi \mapsto \mu \underset{H}{\cdot} \psi,$$

using the special case $K = \{1\}$ from the beginning of Part (c). Note that similar constructions do not work for (bi-)modules over F, since an (FG, FH)-bimodule is not necessarily flat as right FH-module.

(d) If e is an idempotent in $Z(\mathcal{O}G)$ then the decomposition map $d_G^e \colon R(\mathbb{K}G) \to R(FG\overline{e}) \subseteq \mathbb{K}R(\mathbb{K}Ge)$ is given by

$$(d_G^e(\chi))(g) = \begin{cases} \chi(ge), & \text{if } g \in G_{p'}, \\ 0, & \text{otherwise}, \end{cases}$$

for $\chi \in R(\mathbb{K}G)$ and $g \in G$. If e = 1, one obtains the usual decomposition map $d_G \colon R(\mathbb{K}G) \to R(FG)$.

More generally, for a *p*-element $u \in G$ and an idempotent $e \in Z(\mathcal{O}[C_G(u)])$, the generalized decomposition map $d_G^{(u,e)} \colon \mathbb{K}R(\mathbb{K}G) \to \mathbb{K}R(FC_G(u)\overline{e})$ is given by

$$\left(d_G^{(u,e)}(\chi)\right)(g) = \begin{cases} \chi(uge) \,, & \text{if } g \in C_G(u)_{p'}, \\ 0 \,, & \text{otherwise,} \end{cases}$$

for $\chi \in \mathbb{K}R(\mathbb{K}G)$ and $g \in C_G(u)$. If $\chi \in \operatorname{Irr}(\mathbb{K}G)$ belongs to a sum of blocks A of $\mathcal{O}G$ and e is a primitive idempotent in $Z(\mathcal{O}[C_G(u)])$ then Brauer's second main theorem (see [NT89, Theorem 5.4.2]) implies that $d_G^{(u,e)}(\chi) = 0$ unless $(\langle u \rangle, e)$ is an A-Brauer pair. For u = 1 one recovers the decomposition map d_G^e from above.

8.2 Remark (a) Let $M \in {}_{\mathbb{K}G} \mathsf{mod}_{\mathbb{K}H}$, $N \in {}_{\mathbb{K}H} \mathsf{mod}_{\mathbb{K}K}$, and let $\mu \in R(\mathbb{K}G, \mathbb{K}H)$ and $\nu \in R(\mathbb{K}H, \mathbb{K}K)$ denote their respective characters as left modules for $\mathbb{K}[G \times H]$ and $\mathbb{K}[H \times K]$. Then the character $\mu \underset{H}{\cdot} \nu \in R(\mathbb{K}G, \mathbb{K}K)$ of $M \otimes_{\mathbb{K}H} N$ viewed as left $\mathbb{K}[G \times K]$ -module is given by

$$(\mu_{H}^{\cdot}\nu)(g,k) = \frac{1}{|H|} \sum_{h \in H} \mu(g,h)\nu(h,k), \qquad (13)$$

for $(g,k) \in G \times K$. In the special case that $M = V \otimes_{\mathbb{K}} W$ and $N = W' \otimes_{\mathbb{K}} U$ with irreducible modules $V \in_{\mathbb{K}G} \mathsf{mod}$, $W, W' \in_{\mathbb{K}H} \mathsf{mod}$, and $U \in_{\mathbb{K}K} \mathsf{mod}$, one has $M \otimes_{\mathbb{K}H} N \cong V \otimes U \in_{\mathbb{K}[G \times K]} \mathsf{mod}$ if $W^{\circ} \cong W'$, and $M \otimes_{\mathbb{K}H} N = \{0\}$ if $W^{\circ} \ncong W'$. Thus,

$$(\chi_V \times \chi_W)_{H} (\chi_{W'} \times \chi_U) = \begin{cases} \chi_V \times \chi_U, & \text{if } \chi_W^\circ = \chi_{W'}, \\ 0, & \text{otherwise.} \end{cases}$$
(14)

(b) If $e \in Z(\mathbb{K}G)$ is an idempotent then the character of $\mathbb{K}Ge$, viewed as element in $R(\mathbb{K}Ge, \mathbb{K}Ge) \subseteq R(\mathbb{K}[G \times G])$ is given by $\sum_{\chi \in Irr(\mathbb{K}Ge)} \chi \times \chi^{\circ}$.

Parts (a) and (c) of the following definition are due to Broué, see [Br90].

8.3 Definition Let $\mu \in R(\mathbb{K}G, \mathbb{K}H)$.

- (a) The virtual character μ is called *perfect* if it satisfies the following two conditions:
 - (i) For all $(g,h) \in G \times H$, one has $\mu(g,h) \in |C_G(g)| \mathcal{O} \cap |C_H(h)| \mathcal{O}$.
 - (ii) If $(g,h) \in G \times H$ is such that $\mu(g,h) \neq 0$, then g is a p'-element if and only if h is a p'-element.
- (b) We call the virtual character μ quasi-perfect if it satisfies condition (ii) in Part (a).

(c) Assume that $e \in Z(\mathbb{K}G)$ and $f \in Z(\mathbb{K}H)$ are idempotents and that $\mu \in R(\mathbb{K}Ge, \mathbb{K}Hf)$. One calls μ an *isometry* between $\mathbb{K}Ge$ and $\mathbb{K}Hf$ if the map $I_{\mu} \colon R(\mathbb{K}Hf) \to R(\mathbb{K}Ge)$ is bijective and satisfies $(I_{\mu}(\psi), I_{\mu}(\psi'))_G = (\psi, \psi')_H$, for all $\psi, \psi' \in R(\mathbb{K}Hf)$. If additionally μ is perfect, then μ is called a *perfect isometry* between $\mathbb{K}Ge$ and $\mathbb{K}Hf$.

Broué's abelian defect group conjecture in its weakest form states that if $\mathcal{O}Ge$ is a block of $\mathcal{O}G$ (e its block idempotent) with abelian defect group D and $\mathcal{O}[N_G(D)]f$ the block of $\mathcal{O}[N_G(D)]$ which is in Brauer correspondence with $\mathcal{O}Ge$, i.e., $\operatorname{br}_D(e) = \overline{f}$, then there exists a perfect isometry $\mu \in R(\mathbb{K}Ge, \mathbb{K}N_G(D)f)$.

8.4 Remark Let $e \in Z(\mathbb{K}G)$ and $f \in Z(\mathbb{K}H)$ be idempotents and let $\mu \in R(\mathbb{K}Ge, \mathbb{K}Hf)$. The first two of the following statements are quick consequences of (14).

- (a) The following are equivalent:
- (i) μ is an isometry between $\mathbb{K}Ge$ and $\mathbb{K}Hf$.
- (ii) $\mu_{\stackrel{\cdot}{H}}\mu^{\circ} = [\mathbb{K}Ge] \in R(\mathbb{K}Ge, \mathbb{K}Ge) \text{ and } \mu^{\circ}_{\stackrel{\cdot}{C}}\mu = [\mathbb{K}Hf] \in R(\mathbb{K}Hf, \mathbb{K}Hf).$

(iii) There exists a bijection $\operatorname{Irr}(\mathbb{K}Hf) \xrightarrow{\sim} \operatorname{Irr}(\mathbb{K}Ge), \ \psi \mapsto \chi_{\psi}$, and elements $\varepsilon_{\psi} \in \{\pm 1\}$, for $\psi \in \operatorname{Irr}(\mathbb{K}Hf)$, such that $\mu = \sum_{\psi \in \operatorname{Irr}(\mathbb{K}Hf)} \varepsilon_{\psi} \cdot \chi_{\psi} \times \psi^{\circ}$.

(b) One has $\mu \neq 0$ if and only if $\mu \underset{tt}{\cdot} \mu^{\circ} \neq 0$ in $R(\mathbb{K}Ge, \mathbb{K}Ge)$.

(c) The elements in $R(\mathbb{K}Ge, \mathbb{K}Hf)$ that satisfy Condition (i) (resp. Condition (ii)) in Definition 8.3(a) form a subgroup of $R(\mathbb{K}Ge, \mathbb{K}Hf)$.

(d) If $e \in Z(\mathcal{O}G)$, $f \in Z(\mathcal{O}H)$, and μ is the character of an indecomposable module $M \in \mathcal{O}Ge^{\mathsf{triv}}\mathcal{O}Hf}$ with twisted diagonal vertex then μ satisfies Conditions (i) and (ii) in Definition 8.3(a); see [Br90, Proposition 1.2].

(e) If μ is quasi-perfect then the K-linear extension $\mathbb{K}R(\mathbb{K}Hf) \to \mathbb{K}R(\mathbb{K}Ge)$ of I_{μ} restricts to a map $\mathbb{K}R(FHf) \to \mathbb{K}R(FGe)$. In fact, this follows immediately from the formula in (13) in the special case that $K = \{1\}$.

8.5 Proposition Let $\mu \in R(\mathbb{K}Ge, \mathbb{K}Hf)$. The following are equivalent:

- (i) The virtual character μ is quasi-perfect.
- (ii) One has $d_G \circ I_\mu = I_\mu \circ d_H$ as maps $\mathbb{K}R(\mathbb{K}H) \to \mathbb{K}R(\mathbb{K}G)$.
- (iii) One has $d_H \circ I_{\mu^\circ} = I_{\mu^\circ} \circ d_G$ as maps $\mathbb{K}R(\mathbb{K}G) \to \mathbb{K}R(\mathbb{K}H)$.

Proof Clearly, μ is quasi-perfect if and only if μ° is quasi-perfect. Thus it suffices to show the equivalence of (i) and (ii).

Assume first that (i) holds and let $\psi \in \mathbb{K}R(\mathbb{K}H)$ and $g \in G$. Consider the case that $g \notin G_{p'}$. By Equation (13), we have $I_{\mu}(d_H(\psi))(g) = |H|^{-1} \sum_{h \in H} \mu(g,h)(d_H(\psi))(h)$. By our assumptions, $\mu(g,h) = 0$ for

every $h \in H_{p'}$. On the other hand, if $h \in H \setminus H_{p'}$ then $(d_H(\psi))(h) = 0$. Thus, we obtain $I_{\mu}(d_H(\psi))(g) = 0 = d_G(I_{\mu}(\psi))(g)$. Now consider the case that $g \in G_{p'}$. Then, again by (13), we have

$$|H| \cdot d_G(I_{\mu}(\psi))(g) = |H| \cdot I_{\mu}(\psi)(g) = \sum_{h \in H} \mu(g, h)\psi(h) = \sum_{h \in H_{p'}} \mu(g, h)\psi(h) = |H| \cdot I_{\mu}(d_H(\psi))(g)$$

and (ii) holds.

Now assume that (ii) holds. Let $(g,h) \in G \times H$ and assume that $\mu(g,h) \neq 0$. Let $\psi \in \mathbb{K}R(\mathbb{K}H)$ denote the characteristic function on the conjugacy class of h. If $g \in G_{p'}$ and $h \notin H_{p'}$ then $d_H(\psi) = 0$ and (13) implies the contradiction $0 = I_{\mu}(d_H(\psi))(g) = d_G(I_{\mu}(\psi))(g) = I_{\mu}(\psi)(g) = |C_H(h)|^{-1} \cdot \mu(g,h) \neq 0$. And if $g \notin G_{p'}$ and $h \in H_{p'}$ then we obtain the contradiction $0 = d_G(I_{\mu}(\psi))(g) = I_{\mu}(d_H(\psi))(g) = |C_H(h)|^{-1} \cdot \mu(g,h) \neq 0$. Thus, (i) holds.

8.6 Lemma Let H_1, \ldots, H_n be finite groups and let $\mu_i \in R(\mathbb{K}G, \mathbb{K}H_i), i = 1, \ldots, n$, be virtual characters such that $\sum_{i=1}^n \mu_i \underset{H_i}{\cdot} \mu_i^{\circ} = \sum_{\chi \in \Omega} \chi \times \chi^{\circ}$, for some subset $\Omega \subseteq \operatorname{Irr}(\mathbb{K}G)$. Then Ω is the disjoint union of subsets Ω_i , $i = 1, \ldots, n$, with the property that $\mu_i \underset{H_i}{\cdot} \mu_i^{\circ} = \sum_{\chi \in \Omega_i} \chi \times \chi^{\circ}$.

Proof For each i = 1, ..., n we write $\mu_i = \sum_{\chi \in \operatorname{Irr}(\mathbb{K}G)} \chi \times \psi_{i,\chi}$ with $\psi_{i,\chi} \in R(\mathbb{K}H_i)$. Equation (14) implies that, for each $\chi \in \operatorname{Irr}(\mathbb{K}G)$, the coefficient of $\chi \times \chi^\circ$ in $\sum_{i=1}^n \mu_i \underset{H_i}{\cdot} \mu_i^\circ$ is equal to $\sum_{i=1}^n (\psi_{i,\chi}, \psi_{i,\chi})_{H_i}$. Thus, $\sum_{i=1}^n (\psi_{i,\chi}, \psi_{i,\chi})_{H_i}$ is equal to 1 if $\chi \in \Omega$ and equal to 0, if $\chi \notin \Omega$. This implies $\psi_{i,\chi} = 0$ for all i = 1, ..., n if $\chi \notin \Omega$. Moreover, if $\chi \in \Omega$, then there exists a unique $i \in \{1, ..., n\}$ such that $\psi_{i,\chi} \neq 0$. For i = 1, ..., n we define Ω_i as the set of those $\chi \in \Omega$ with $\psi_{i,\chi} \neq 0$. Now the lemma follows.

8.7 Lemma Let *e* be a block idempotent of $\mathcal{O}G$, *f* a block idempotent of $\mathcal{O}H$ and suppose that $\mu \in R(\mathbb{K}Ge, \mathbb{K}Hf)$ is a quasi-perfect virtual character such that there exists a non-empty subset Ω of $\operatorname{Irr}(\mathbb{K}Ge)$ with $\mu_{\mu} \mu^{\circ} = \sum_{\chi \in \Omega} \chi \times \chi^{\circ}$. Then $\Omega = \operatorname{Irr}(\mathbb{K}Ge)$ and μ is an isometry between $\mathbb{K}Ge$ and $\mathbb{K}Hf$.

Proof After writing μ as a \mathbb{Z} -linear combination of the basis elements $\chi \times \psi^{\circ}$, $(\chi, \psi) \in \operatorname{Irr}(\mathbb{K}Ge) \times \operatorname{Irr}(\mathbb{K}Hf)$, and using Equation (14), the hypothesis $\mu_{H}^{\cdot} \mu^{\circ} = \sum_{\chi \in \Omega} \chi \times \chi^{\circ}$ implies that there exists a subset Λ of $\operatorname{Irr}(\mathbb{K}Hf)$ and a bijection $\alpha \colon \Omega \to \Lambda$ such that $\mu = \sum_{\chi \in \Omega} \varepsilon_{\chi} \cdot \chi \times \alpha(\chi)^{\circ}$, with $\varepsilon_{\chi} \in \{\pm 1\}$ for $\chi \in \Omega$. This implies $\mu^{\circ}_{G} \mu = \sum_{\psi \in \Lambda} \psi \times \psi^{\circ}$. As μ is quasi-perfect, so is μ° . Thus, by symmetry, it suffices now to show that $\Omega = \operatorname{Irr}(\mathbb{K}Ge)$.

For $\chi, \chi' \in \operatorname{Irr}(\mathbb{K}Ge)$ set $m_{\chi,\chi'} := (d_G(\chi), \chi')_G \in \mathbb{K}$. Then $m_{\chi',\chi} = m_{\chi,\chi'}$ and if χ has height 0 then $m_{\chi,\chi'} \neq 0$ for all $\chi' \in \operatorname{Irr}(\mathbb{K}Ge)$, see [NT89, Lemma 3.6.34(ii)]. Thus, to complete the proof it suffices to show that if $\chi \in \Omega$ and $\chi' \in \operatorname{Irr}(\mathbb{K}Ge)$ with $m_{\chi,\chi'} \neq 0$ then also $\chi' \in \Omega$. But this holds if and only if $d_G(\Omega) \subseteq \langle \Omega \rangle_{\mathbb{K}}$. So let $\chi \in \Omega$ and set $\psi := \alpha(\chi)$. Then $I_{\mu}(\psi) = \varepsilon_{\chi} \cdot \chi$. Since μ is quasi-perfect, Proposition 8.5 implies that

$$d_G(\chi) = \varepsilon_{\chi} \cdot d_G(I_{\mu}(\psi)) = \varepsilon_{\chi} \cdot I_{\mu}(d_H(\psi)) \in I_{\mu}(\mathbb{K}R(\mathbb{K}Hf)) \subseteq \langle \Omega \rangle_{\mathbb{K}},$$

and the proof is complete.

8.8 Corollary Let e be a block idempotent of $\mathcal{O}G$, let H_1, \ldots, H_n be finite groups, and, for each $i = 1, \ldots, n$, let f_i be a block idempotent of $\mathcal{O}H_i$. Furthermore, for $i = 1, \ldots, n$, let $\mu_i \in R(\mathbb{K}Ge, \mathbb{K}H_if_i)$ be a quasiperfect virtual character such that $\sum_{i=1}^{n} \mu_i \stackrel{\cdot}{\underset{H_i}{}} \mu_i^{\circ} = \sum_{\chi \in \Omega} \chi \times \chi^{\circ}$ in $R(\mathbb{K}Ge, \mathbb{K}Ge)$ for some non-empty subset $\Omega \subseteq \operatorname{Irr}(\mathbb{K}Ge)$. Then there exists a unique $i \in \{1, \ldots, n\}$ such that $\mu_i \neq 0$. Moreover, $\Omega = \operatorname{Irr}(\mathbb{K}Ge)$ and μ_i is an isometry between $\mathbb{K}Ge$ and $\mathbb{K}H_if_i$.

Proof Applying Lemma 8.6, we see that Ω is a disjoint union of subsets Ω_i such that $\mu_i \underset{H_i}{\cdot} \mu_i^\circ = \sum_{\chi \in \Omega_i} \chi \times \chi^\circ$. Note that $\mu_i \neq 0$ if and only if $\Omega_i \neq \emptyset$. Choose $i \in \{1, \ldots, n\}$ such that Ω_i is non-empty. Then Lemma 8.7 implies that $\Omega_i = \operatorname{Irr}(\mathbb{K}Ge)$ and that μ_i is an isometry between $\mathbb{K}Ge$ and $\mathbb{K}H_if_i$. This also implies that $\Omega_j = \emptyset$ for all $j \neq i$ in $\{1, \ldots, n\}$ and therefore $\mu_j = 0$ for all $j \neq i$ in $\{1, \ldots, n\}$.

8.9 Corollary Let e be a block idempotent of \mathcal{OG} , let $f \in Z(\mathcal{OH})$ be an idempotent, and let $\mu \in R(\mathbb{K}Ge, \mathbb{K}Hf)$ be a quasi-perfect virtual character satisfying $\mu \cdot \mu^{\circ} = [\mathbb{K}Ge]$ in $R(\mathbb{K}Ge, \mathbb{K}Ge)$. Then there exists a unique primitive idempotent f' of $Z(\mathcal{OH}f)$ such that $\mu = \mu \cdot f'$. Furthermore, μ is an isometry between $\mathbb{K}Ge$ and $\mathbb{K}Hf'$.

Proof Let f'_1, \ldots, f'_n denote the primitive idempotents of $Z(\mathcal{O}Hf)$, and for each $i = 1, \ldots, n$, set $\mu_i := \mu \cdot f'_i \in R(\mathbb{K}Ge, \mathbb{K}Hf'_i)$. Then $\mu = \sum_{i=1}^n \mu_i$ and

$$\sum_{i=1}^{n} \mu_i \underset{H}{\cdot} \mu_i^{\circ} = \mu \underset{H}{\cdot} \mu^{\circ} = [\mathbb{K}Ge] = \sum_{\chi \in \operatorname{Irr}(\mathbb{K}Ge)} \chi \times \chi^{\circ}.$$

Proposition 8.5(ii), together with the fact that d_H respects the block decomposition, implies that with μ also μ_i is quasi-perfect. Now Corollary 8.8 applies and the proof is complete.

The following corollary is an immediate consequence of Corollary 8.9. It also slightly generalizes a result in [Br90, Théorème 1.5(2)].

8.10 Corollary Let $e \in Z(\mathcal{O}G)$ and $f \in Z(\mathcal{O}H)$ be idempotents. Let \mathcal{I} denote the set of primitive idempotents e' of $Z(\mathcal{O}Ge)$ with e'e = e' and let \mathcal{J} denote the set of primitive idempotents f' of $Z(\mathcal{O}Hf)$ with f'f = f'. Suppose that $\mu \in R(\mathbb{K}Ge, \mathbb{K}Hf)$ is a quasi-perfect isometry between $\mathbb{K}Ge$ and $\mathbb{K}Hf$. Then, for each $e' \in \mathcal{I}$ there exists a unique $f' \in \mathcal{J}$ such that $e'\mu f' \neq 0$. Conversely, for each $f' \in \mathcal{J}$ there exists a unique $e' \in \mathcal{I}$ such that $e'\mu f' \neq 0$. These conditions define inverse bijections between \mathcal{I} and \mathcal{J} . Moreover, if $e' \in \mathcal{I}$ and $f' \in \mathcal{J}$ satisfy $e'\mu f' \neq 0$ then $e'\mu f'$ is an isometry between $\mathbb{K}Ge'$ and $\mathbb{K}Hf'$.

Proof First note that by Proposition 8.5(ii), with μ also $e'\mu f'$ is a quasi-perfect character for every $e' \in \mathcal{I}$ and $f' \in \mathcal{J}$. Next let $e' \in \mathcal{I}$. Then, Corollary 8.9 applied to e' and f and the quasi-perfect virtual characters $e'\mu f'$, $f' \in \mathcal{J}$, implies that there exists a unique $f' \in \mathcal{J}$ with $e'\mu f' \neq 0$ and that $e'\mu f'$ is an isometry between $\mathbb{K}Ge'$ and $\mathbb{K}Hf'$. This proves the first statement. Symmetrically, fixing $f' \in \mathcal{J}$ and using μ° , we obtain the second statement. The remaining statements are clear from the above.

The following Lemma will be used in Section 10.

8.11 Lemma Let $e \in Z(\mathbb{K}G)$ and $f \in Z(\mathbb{K}H)$ be idempotents and let $\mu \in R(\mathbb{K}Ge, \mathbb{K}Hf)$ be quasi-perfect such that $d_G \circ I_{\mu} \colon R(\mathbb{K}Hf) \to \mathbb{K}R(\mathbb{K}Ge)$ is non-zero. Then $d_{G \times H}(\mu) \neq 0$. In particular, if $e \neq 0 \neq f$ and μ is a quasi-perfect isometry between $\mathbb{K}Ge$ and $\mathbb{K}Hf$, then $d_{G \times H}(\mu) \neq 0$.

Proof Since μ is quasi-perfect, we have $d_G \circ I_\mu = I_\mu \circ d_H$ as maps from $\mathbb{K}R(\mathbb{K}Hf)$ to $\mathbb{K}R(\mathbb{K}Ge)$ by Lemma 8.5. Since μ is quasi-perfect, it follows from Equation (13) and the definition of quasi-perfect that $I_\mu \circ d_H = I_{d_G \times H(\mu)} \circ d_H$ as functions from $\mathbb{K}R(\mathbb{K}Hf)$ to $\mathbb{K}R(\mathbb{K}Ge)$. Thus, we have $I_{d_G \times H(\mu)} \circ d_H = d_G \circ I_\mu \neq 0$ and hence $d_{G \times H}(\mu) \neq 0$.

The following lemma is an immediate consequence of the definition of $\bigotimes_{\mathbb{K}H}^{X,Y}$ and will be used in Section 10.

8.12 Lemma Let $X \leq G \times H$ and $Y \leq H \times K$ be subgroups satisfying $k_1(Y) \leq k_2(X)$. Further let $\mu \in R(\mathbb{K}X)$ and $\nu \in R(\mathbb{K}Y)$ be such that $\mu \stackrel{X,Y}{\stackrel{\cdot}{H}} \nu \in R(\mathbb{K}[X * Y])$ is equal to [M] or to -[M] for some non-zero module $M \in \mathbb{K}[X*Y]$ mod. Then $\operatorname{res}_{k_1(Y) \times k_2(Y)}^Y(\nu) \neq 0$.

Proof Clearly, $\operatorname{res}_{k_1(X) \times k_1(Y)}^X(\mu) \underset{k_1(Y)}{\cdot} \operatorname{res}_{k_1(Y) \times k_2(Y)}^Y(\nu) = \operatorname{res}_{k_1(X) \times k_2(Y)}^{X,Y}(\mu \underset{H}{\overset{X,Y}{\cdot}} \nu)$ and the latter is equal to $\operatorname{res}_{k_1(X) \times k_2(Y)}^{X,Y}([M])$ or its negative, and therefore non-zero. The result now follows.

9 Grothendieck groups of *p*-permutation modules and *p*-permutation equivalences

We assume again that G and H are finite groups and that the p-modular system $(\mathbb{K}, \mathcal{O}, F)$ is large enough for G and H.

9.1 Grothendieck groups of p-permutation modules. (a) For an idempotent $e \in Z(\mathcal{O}G)$, we denote by $T(\mathcal{O}Ge)$ the Grothendieck group of the category \mathcal{O}_{Ge} triv with respect to direct sums. The group $T(\mathcal{O}Ge)$ is free as abelian group with standard basis given by the elements [M], where M runs through a set of representatives of the isomorphism classes of indecomposable p-permutation $\mathcal{O}Ge$ -modules. For an arbitrary module $M \in \mathcal{O}_{Ge}$ triv we write $[M] = [M_1] + \cdots + [M_r] \in T(\mathcal{O}Ge)$ if $M = M_1 \oplus \cdots \oplus M_r$ is a decomposition of M into indecomposable submodules. We always view $T(\mathcal{O}Ge)$ as a subgroup of $T(\mathcal{O}G)$ in the natural way. Moreover, we say that an indecomposable module $M \in \mathcal{O}_{Ge}$ triv appears in an element $\omega \in T(\mathcal{O}G)$, if [M] occurs with non-zero coefficient in ω with respect to the above standard basis. Note that multiplying a p-permutation $\mathcal{O}G$ -modules with e defines a projection map $T(\mathcal{O}G) \to T(\mathcal{O}Ge)$, $\omega \mapsto e\omega$. Similarly, we define the Grothendieck group T(FGe). If additionally $f \in Z(\mathcal{O}H)$ is an idempotent then we define $T(\mathcal{O}Ge, \mathcal{O}Hf) := T(\mathcal{O}[G \times H](e \otimes f^*))$. If $M \in \mathcal{O}_{Ge}$ triv \mathcal{O}_{Hf} we denote by [M] the corresponding element in $T(\mathcal{O}Ge, \mathcal{O}Hf)$. Similar notations will be used over F. The \mathbb{Z} -span of the elements $[M] \in T(\mathcal{O}Ge)$, where M is an indecomposable projective $\mathcal{O}Ge$ -module will be denoted by $Pr(\mathcal{O}Ge)$. We also use the notations $Pr(\mathcal{O}Ge, \mathcal{O}Hf)$, Pr(FGe) and Pr(FGe, FHf) with obvious meanings.

(b) Tensor products of bimodules and generalized tensor products as introduced in Section 6 induce maps on Grothendieck group levels that we denote again by \vdots_{H} and $\overset{X,Y}{\underset{H}{\mapsto}}$, as in 8.1. Similarly, the Brauer construction with respect to a *p*-subgroup *P* of *G* induces a homomorphism $T(\mathcal{O}G) \to T(F[N_G(P)/P]), \ \omega \mapsto \omega(P)$. Often we will also consider $\omega(P)$ as element of $T(F[N_G(P)])$ after applying inflation. Note that for a Brauer pair (P, e) of *FG*, one obtains a homomorphism $-(P, e): T(FG) \to T(FIe), \ \omega \mapsto \omega(P, e) = e\omega(P)$, where $I = N_G(P, e)$. Similarly, one obtains a homomorphism $-(P, e): T(\mathcal{O}G) \to T(FIe)$.

(c) For each idempotent $e \in Z(\mathcal{O}G)$ we have a commutative diagram

$$\begin{array}{cccc} Pr(\mathcal{O}Ge) & \subseteq & T(\mathcal{O}Ge) & \xrightarrow{\kappa_G} & R(\mathbb{K}Ge) \\ & & & & & \downarrow \\ & & & & \downarrow \\ Pr(FGe) & \subseteq & T(FGe) & \xrightarrow{\eta_G} & R(FGe) \end{array}$$

whose top horizontal map κ_G is induced by the scalar extension functors $\mathbb{K} \otimes_{\mathcal{O}} -$, whose left vertical maps are induced by the scalar extension functor $F \otimes_{\mathcal{O}} -$, whose right vertical map is the decomposition map, and whose bottom horizontal map η_G sends [M] to [M] for any $M \in {}_{FGe}$ triv. In other words, if M is indecomposable (i.e., $[M] \in T(FGe)$ a standard basis element) then [M] is mapped to the sum of its composition factors (in terms of the standard basis in R(FGe)). Recall from Proposition 3.3(b) that the left vertical maps are indeed isomorphisms preserving the standard basis elements and vertices. Recall also from [NT89, Theorem 3.6.15(i)] that the map κ_G is injective on $Pr(\mathcal{O}Ge)$.

For an element $\omega \in T(FG\overline{e})$ we will denote the image under κ_G of the corresponding element in $T(\mathcal{O}Ge)$ by $\omega^{\mathbb{K}} \in R(\mathbb{K}Ge)$.

The following proposition is well-known to specialists. We state it for easy reference.

9.2 Proposition Let A be a block of $\mathcal{O}G$ and let $\mathcal{BP}(A)$ denote a set of representatives of the G-orbits of A-Brauer pairs.

(a) The map

$$T(A) \mapsto \prod_{(P,e)\in\widetilde{\mathcal{BP}}(A)} R(F[N_G(P,e)]e), \quad \omega \mapsto \left(\eta_{N_G(P,e)}(\omega(P,e))\right)_{(P,e)\in\widetilde{\mathcal{BP}}(A)},$$

is an injective group homomorphism and has finite cokernel.

(b) The map

$$T(A) \mapsto \prod_{(P,e)\in\widetilde{\mathcal{BP}}(A)} R(\mathbb{K}[N_G(P,e)]e), \quad \omega \mapsto \left((\omega(P,e)^{\mathbb{K}}) \right)_{(P,e)\in\widetilde{\mathcal{BP}}(A)}$$

is an injective group homomorphism.

Proof (a) By Conlon's Theorem (see [Be98, Theorem 5.5.4]), one has an injective map

$$T(\mathcal{O}G) \mapsto \prod_{P} R(F[N_G(P)]), \quad \omega \mapsto (\eta_{N_G(P)}(\omega(P)))_{P}$$

where P runs through a set of representatives of the G-conjugacy classes of p-subgroups of G. By Proposition 3.3(c) it has finite cokernel. By Lemma 3.7, this map splits into a direct sum of maps with respect to each block A of $\mathcal{O}G$. Further, the Morita equivalence in 5.2 gives the statement of Part (a).

(b) This follows from Part (a) and the commutativity of the diagram in 9.1(c).

The following Lemma will be used in the proof of Lemma 10.7.

9.3 Lemma Let $0 \neq \omega \in T(FG)$ and let (P, e) be a Brauer pair of FG which is maximal among all the Brauer pairs (Q, f) of FG satisfying $M(Q, f) \neq \{0\}$ for some indecomposable FG-module M appearing in ω .

(a) Let M be an indecomposable p-permutation FG-module appearing in ω and satisfying $M(P, e) \neq \{0\}$. Then (P, e) is a maximal M-Brauer pair, M(P, e) is an indecomposable p-permutation $F[N_G(P, e)]$ -module, and the coefficient of [M] in $\omega \in T(FG)$ equals the coefficient of [M(P, e)] in $\omega(P, e) \in T(F[N_G(P, e)])$.

(b) One has $0 \neq \omega(P, e) \in Pr(F[N_G(P, e)/P])$.

Proof (a) The maximality of (P, e) and Proposition 5.3 imply that P is a vertex of M and that (P, e) is a maximal M-Brauer pair. The rest follows from Proposition 5.4.

(b) This follows immediately from Part (a) and Proposition 3.3.

9.4 Remark We will need to use the following result from [BX08, Corollary 2.6], stating that the generalized decomposition map on characters of *p*-permutation $\mathcal{O}G$ -modules is an element of the Brauer character ring (without extending scalars) and can be expressed via the Brauer construction: Let $M \in \mathcal{O}G$ triv and let $u \in G$ be a *p*-element. Then

$$d_G^u(\kappa_G([M])) = \eta_{C_G(u)}(\operatorname{res}_{C_G(u)}^{N_G(\langle u \rangle)}([M(\langle u \rangle)]))$$
(15)

in $R(F[C_G(u)])$.

9.5 Lemma Let $\omega \in T(FG)$. For every Brauer pair (P, e) of FG set $\chi_{(P,e)} := (\omega(P, e))^{\mathbb{K}} \in R(\mathbb{K}[N_G(P, e)])$ and $\psi_{(P,e)} := \operatorname{res}_{C_G(P)}^{N_G(P,e)}(\chi_{(P,e)}) \in R(\mathbb{K}[C_G(P)])$. Assume that, for every Brauer pair (P, e) of FG, the following two conditions are satisfied:

- (i) If $\chi_{(P,e)} \neq 0$ in $R(\mathbb{K}[N_G(P,e)])$ then $\psi_{(P,e)} \neq 0$ in $R(\mathbb{K}[C_G(P)])$.
- (ii) If $\psi_{(P,e)} \neq 0$ in $R(\mathbb{K}[C_G(P)])$ then $d_{C_G(P)}(\psi_{(P,e)}) \neq 0$ in $R(F[C_G(P)])$.

Then, for any two Brauer pairs $(P, e) \leq (Q, f)$ of $\mathcal{O}G$, one has: If $\psi_{(Q,f)} \neq 0$ in $R(\mathbb{K}[C_G(Q)])$ then $\psi_{(P,e)} \neq 0$ in $R(\mathbb{K}[C_G(P)])$.

Proof Arguing by induction on [Q : P] we may assume that $(P, e) \leq (Q, f)$ and that Q/P is cyclic. Thus, there exists $u \in Q$ such that $P\langle u \rangle = Q$. Assume that $\psi_{(P,e)} = 0$ in $R(\mathbb{K}[C_G(P)])$ and set $I := N_G(P, e)$. Then, by (i), also $\chi_{(P,e)} = 0$ in $R(\mathbb{K}I)$. By Lemma 3.7 applied to the *I*-stable idempotent *e*, the element $\omega(P) \in T(FN_G(P))$ and the *p*-subgroup $\langle u \rangle \leq I$, we obtain $(e\omega(P))(\langle u \rangle) = br_{\langle u \rangle}(e)(\omega(P)(\langle u \rangle))$ in $T(N_I(\langle u \rangle))$. Note that $br_{\langle u \rangle}(e) = br_Q(e)$. Thus, by Proposition 3.5(b), after further restriction (omitted in the notation) we obtain

$$(e\omega(P))(\langle u\rangle) = br_Q(e)\omega(Q)$$

in $T(F[N_I(\langle u \rangle) \cap N_G(Q)])$. Applying Equation (15) to $e\omega(P) \in T(FI)$ and noting that $C_I(u) \leq N_I(\langle u \rangle) \cap N_G(Q)$, we obtain further

$$\eta_{C_I(u)}(\operatorname{br}_Q(e)\operatorname{res}_{C_I(u)}^{N_G(Q)}(\omega(Q))) = \eta_{C_I(u)}\operatorname{res}_{C_I(u)}^{N_I(\langle u \rangle)}((e\omega(P))(\langle u \rangle)) = d_I^u((e\omega(P))^{\mathbb{K}}) = d_I^u(\chi_{(P,e)}) = 0$$

Restricting further to $C_G(Q)$, multiplying by f, and using that $f \operatorname{br}_Q(e) = f$, we finally have

$$0 = \eta_{C_G(Q)} \left(f \operatorname{br}_Q(e) \operatorname{res}_{C_G(Q)}^{N_G(Q)}(\omega(Q)) \right) = \eta_{C_G(Q)} \left(\operatorname{res}_{C_G(Q)}^{N_G(Q,f)}(f\omega(Q)) \right)$$

= $d_{C_G(Q)} \left(\operatorname{res}_{C_G(Q)}^{N_G(Q,f)}(f\omega(Q))^{\mathbb{K}} \right) = d_{C_G(Q)} (\psi_{(Q,f)}).$

Condition (ii) now implies that $\psi_{(Q,f)} = 0$, and the result follows.

The following definition is similar to Definition 5.1. It will be used extensively in Section 10.

9.6 Definition Let $\omega \in T(FG)$ or $\omega \in T(\mathcal{O}G)$. A Brauer pair (P, e) of FG is called an ω -Brauer pair if $\omega(P, e) = e\omega(P) \neq 0$ in $T(F[N_G(P, e)])$. The set of ω -Brauer pairs is denoted by $\mathcal{BP}(\omega)$. Its corresponding set of Brauer pairs over \mathcal{O} is denoted by $\mathcal{BP}_{\mathcal{O}}(\omega)$.

9.7 Notation Let X be a subgroup of $G \times H$ and let $d \in Z(\mathcal{O}[G \times H])$. We denote by $T^{\Delta}(\mathcal{O}Xd)$ the subgroup of $T(\mathcal{O}Xd)$ which is spanned by all standard basis elements [M], where M is an indecomposable p-permutation $\mathcal{O}Xd$ -module with twisted diagonal vertices (as subgroups of $G \times H$). If $e \in Z(\mathcal{O}G)$ and $f \in Z(\mathcal{O}H)$ are idempotents then we set $T^{\Delta}(\mathcal{O}Ge, \mathcal{O}Hf) := T^{\Delta}(\mathcal{O}[G \times H](e \otimes f^*))$. Similarly we define groups $T^{\Delta}(FXd)$ and $T^{\Delta}(FGe, FHf)$.

9.8 Definition Let $e \in Z(\mathcal{O}G)$ and $f \in Z(\mathcal{O}H)$ be non-zero idempotents. A *p*-permutation equivalence between $\mathcal{O}Ge$ and $\mathcal{O}Hf$ is an element $\gamma \in T^{\Delta}(\mathcal{O}Ge, \mathcal{O}Hf)$ satisfying

$$\gamma \stackrel{\cdot}{}_{H} \gamma^{\circ} = [\mathcal{O}Ge] \text{ in } T^{\Delta}(\mathcal{O}Ge, \mathcal{O}Ge) \quad \text{and} \quad \gamma^{\circ} \stackrel{\cdot}{}_{G} \gamma = [\mathcal{O}Hf] \text{ in } T^{\Delta}(\mathcal{O}Hf, \mathcal{O}Hf).$$
(16)

The set of *p*-permutation equivalences between $\mathcal{O}Ge$ and $\mathcal{O}Hf$ will be denoted by $T_o^{\Delta}(\mathcal{O}Ge, \mathcal{O}Hf)$ ('o' for 'orthogonal'). Similarly, we define *p*-permutation equivalences between FGe and FHf, and denote the resulting set by $T_o^{\Delta}(FGe, FHf)$. Clearly, an element $\gamma \in T^{\Delta}(\mathcal{O}Ge, \mathcal{O}Hf)$ is a *p*-permutation equivalence between $\mathcal{O}Ge$ and $\mathcal{O}Hf$ if and only if $\overline{\gamma} \in T^{\Delta}(FGe, FHf)$ is a *p*-permutation equivalence between FGe and FHf. Moreover, we denote the set of elements $\gamma \in T^{\Delta}(\mathcal{O}Ge, \mathcal{O}Hf)$ satisfying the first (resp. second) equation in (16) by $T_l^{\Delta}(\mathcal{O}Ge, \mathcal{O}Hf)$ (resp. $T_r^{\Delta}(\mathcal{O}Ge, \mathcal{O}Hf)$) and call them left (resp. right) *p*-permutation equivalences between $\mathcal{O}Ge$ and $\mathcal{O}Hf$. We will see later that $T_l^{\Delta}(\mathcal{O}Ge, \mathcal{O}Hf) = T_o^{\Delta}(\mathcal{O}Ge, \mathcal{O}Hf) = T_r^{\Delta}(\mathcal{O}Ge, \mathcal{O}Hf)$.

9.9 Proposition Let $e \in Z(\mathcal{O}G)$ and $f \in Z(\mathcal{O}H)$ be non-zero idempotents and let $\gamma \in T_o^{\Delta}(\mathcal{O}Ge, \mathcal{O}Hf)$. Then $\mu := \kappa_{G \times H}(\gamma) \in R(\mathbb{K}Ge, \mathbb{K}Hf)$ is a perfect isometry between $\mathbb{K}Ge$ and $\mathbb{K}Hf$.

Proof The virtual character μ is perfect by Remark 8.4(c) and (d). Moreover, applying $\kappa_{G \times G}$ to the equation $\gamma \underset{H}{\cdot} \gamma^{\circ} = [\mathcal{O}Ge]$ in $T(\mathcal{O}Ge, \mathcal{O}Ge)$ implies $\mu \underset{H}{\cdot} \mu^{\circ} = [\mathbb{K}Ge] \in R(\mathbb{K}Ge, \mathbb{K}Ge)$. Similarly, $\gamma^{\circ} \underset{G}{\cdot} \gamma = [\mathcal{O}Hf]$ implies $\mu^{\circ} \underset{H}{\cdot} \mu = [\mathbb{K}Hf] \in R(\mathbb{K}Hf, \mathbb{K}Hf)$. By Remark 8.4(a), μ is an isometry between $\mathbb{K}Ge$ and $\mathbb{K}Hf$.

10 Brauer pairs of *p*-permutation equivalences

Throughout this section we assume that G and H are finite groups and that the *p*-modular system $(\mathbb{K}, \mathcal{O}, F)$ is large enough for G and H. Moreover, we fix a non-zero sum $A = \mathcal{O}Ge_A$ of blocks of $\mathcal{O}G$ and a non-zero sum $B = \mathcal{O}He_B$ of blocks of $\mathcal{O}H$, with e_A and e_B their respective identity elements. Furthermore, we assume throughout this section that $\gamma \in T_l^{\Delta}(A, B)$, i.e., $\gamma \in T^{\Delta}(A, B)$ and γ satisfies

$$\gamma_{H} \cdot \gamma^{\circ} = [A] \quad \text{in } T^{\Delta}(A, A).$$
⁽¹⁷⁾

Instead of requiring that γ is a *p*-permutation equivalence between *A* and *B*, we prove as much as we can under the weaker assumption in (17) and we will finally show in Section 12 that this implies that γ is a *p*-permutation equivalence between *A* and the sum of the blocks *B'* of $\mathcal{O}H$ that support γ from the right, i.e., the sum of those blocks *B'* with $\gamma e_{B'} \neq 0$ in $T(\mathcal{O}G, \mathcal{O}H)$. Note that Equation (17) implies that *A* is precisely the sum of those blocks of $\mathcal{O}G$ that support γ from the left.

The main result in this section is Theorem 10.11, especially Parts (a) and (b), which show that the γ -Brauer pairs behave exactly as the Brauer pairs of an indecomposable *p*-permutation module.

10.1 Notation Let $\Delta(P, \phi, Q)$ be a twisted diagonal *p*-subgroup of $G \times H$. We will abbreviate the element $\gamma(\Delta(P, \phi, Q)) \in T(F[N_{G \times H}(\Delta(P, \phi, Q))])$ by $\overline{\gamma}(P, \phi, Q)$ and will denote the resulting elements in the other representation groups of $N_{G \times H}(\Delta(P, \phi, Q))$ from the diagram in 9.1(c) by

$$\begin{array}{cccc} \gamma(P,\phi,Q) & \mapsto & \mu(P,\phi,Q) \\ & & \downarrow & & \downarrow \\ \hline \overline{\gamma}(P,\phi,Q) & \mapsto & \nu(P,\phi,Q) \end{array}$$

Via restriction, we will also view these elements in the corresponding Grothendieck groups of any subgroup of $N_{G\times H}(\Delta(P,\phi,Q))$, in particular of the subgroup $C_G(P) \times C_H(Q) = C_{G\times H}(\Delta(P,\phi,Q))$. Note that the four maps commute with restrictions, so that the restricted elements still are related through these maps.

10.2 Remark (a) Let $\Delta(P, \phi, Q)$ be a twisted diagonal *p*-subgroup of $G \times H$. If (P, e) is a Brauer pair of *FG* and (Q, e) is a Brauer pair of *FH* then $(\Delta(P, \phi, Q), e \otimes f^*)$ is a Brauer pair of $F[G \times H]$, using that $C_{G \times H}(\Delta(P, \phi, Q)) = C_G(P) \times C_H(Q)$. Conversely, if $(\Delta(P, \phi, Q), e \otimes f^*)$ is a Brauer pair of $F[G \times H]$ then (P, e)is a Brauer pair of *FG* and (Q, f) is a Brauer pair of *FH*. In this case, with $I := N_G(P, e)$ and $J := N_H(Q, f)$, one has $N_{G \times H}(\Delta(P, \phi, Q), e \otimes f^*) = N_{I \times J}(\Delta(P, \phi, Q))$ and $e\gamma(P, \phi, Q)f \in T^{\Delta}(\mathcal{O}[N_{I \times J}(\Delta(P, \phi, Q))](e \otimes f^*))$. Moreover, $(\Delta(P, \phi, Q), e \otimes f^*)$ is an $A \otimes B^*$ -Brauer pair if and only if (P, e) is an *A*-Brauer pair and (Q, f) is a *B*-Brauer pair. We will denote the set of $A \otimes B^*$ -Brauer pairs (X, d), where $X \leq G \times H$ is a twisted diagonal *p*-subgroup, by $\mathcal{BP}^{\Delta}(A, B)$, or $\mathcal{BP}^{\Delta}_{\mathcal{O}}(A, B)$ if lifted to \mathcal{O} .

(b) Note that, for every $(\Delta(P,\phi,Q), e \otimes f^*) \in \mathcal{BP}^{\Delta}_{\mathcal{O}}(\mathcal{O}G,\mathcal{O}H)$, the element $e\mu(P,\phi,Q)f \in R(\mathbb{K}[C_G(P)]e,\mathbb{K}[C_H(Q)]f)$ is perfect. In fact, $e\gamma(P,\phi,Q)f \in T^{\Delta}(\mathcal{O}[C_G(P)]e,\mathcal{O}[C_H(Q)]f)$, by Lemma 3.8(b) and then Remark 8.4(d) applies. Moreover, the restriction of $e\gamma(P,\phi,Q)f$ to $C_G(P) \times \{1\}$ and to $\{1\} \times C_H(Q)$ yields elements of $Pr(\mathcal{O}[C_G(Q)])$ and $Pr(\mathcal{O}[C_H(Q)])$, respectively, see Lemma 3.8(b). Note that $e\gamma(P,\phi,Q)f = 0$ unless (P,e) is an A-Brauer pair and (Q,f) is a B-Brauer pair. Since every module appearing in γ has twisted diagonal vertex, one has $\gamma(X) = 0 \in T(\mathcal{O}[N_{G \times H}(X)])$ for every p-subgroup $X \leq G \times H$ which is not twisted diagonal. Thus, every γ -Brauer pair has a twisted diagonal subgroup as first component.

(c) Let $\Delta(P', \phi', Q') \leq \Delta(P, \phi, Q)$ be twisted diagonal subgroups of $G \times H$, i.e., $Q' \leq Q$, $P' = \phi(Q) \leq P$, and $\phi' = \phi|_{Q'}$. Furthermore, let $(\Delta(P, \phi, Q), e \otimes f^*)$ and $(\Delta(P', \phi', Q'), e' \otimes f'^*)$ be $\mathcal{O}[G \times H]$ -Brauer pairs. Then $(\Delta(P', \phi', Q'), e' \otimes f'^*) \leq (\Delta(P, \phi, Q), e \otimes f^*)$ if and only if $(P', e') \leq (P, e)$ and $(Q', f') \leq (Q, f)$.

The proofs of the following two Lemmas use the two statements and the notation in Theorem 7.5(d).

10.3 Lemma Let $(P, e) \in \mathcal{BP}_{\mathcal{O}}(A)$. Consider the set $\Lambda_B \subseteq \Lambda$ of pairs $(\phi, (Q, f))$, where $(Q, f) \in \mathcal{BP}_{\mathcal{O}}(B)$ and $\phi: Q \xrightarrow{\sim} P$ is an isomorphism, together with its *H*-action from Theorem 7.5(d). There exists a unique *H*-orbit of pairs $(\phi, (Q, f)) \in \Lambda_B$ such that

$$e\mu(P,\phi,Q)f\neq 0$$
 in $R(\mathbb{K}[C_G(P)]e,\mathbb{K}[C_H(Q)]f)$.

Moreover, for each pair $(\phi, (Q, f)) \in \Lambda_B$ which satisfies this condition, the element $e\mu(P, \phi, Q)f$ is a perfect isometry between $\mathbb{K}[C_G(P)]e$ and $\mathbb{K}[C_H(Q)]f$, and $e\nu(P, \phi, Q)f \neq 0$ in $R(F[C_G(P)]e, F[C_H(Q)]f)$.

Proof We apply the Brauer construction with respect to $\Delta(P)$ to Equation (17). By Proposition 4.3(b), we have $eA(\Delta(P))e \cong F[C_G(P)]e$ as $(F[C_G(P)], F[C_G(P)])$ -bimodules. Thus, by Theorem 7.5(d), the following equation holds in $T^{\Delta}(F[C_G(P)]e, F[C_G(P)]e)$:

$$[F[C_G(P)]e] = [eA(\Delta(P))e] = e(\gamma \underset{H}{\cdot} \gamma^{\circ})(\Delta(P))e = \sum_{(\phi,(Q,f))\in\widetilde{\Lambda}} e\overline{\gamma}(P,\phi,Q)f \underset{C_H(Q)}{\cdot} f\overline{\gamma}^{\circ}(Q,\phi^{-1},P)e,$$

where Λ denotes a set of representatives of the *H*-orbits of Λ , as in Theorem 7.5(d). If $(Q, f) \in \mathcal{BP}_{\mathcal{O}}(\mathcal{O}G) \setminus \mathcal{BP}_{\mathcal{O}}(B)$ then $e\overline{\gamma}(P, \phi, Q)f = 0$, since $\gamma \in T(A, B)$. Thus, we may replace Λ in the above summation by $\Lambda_B := \Lambda_B \cap \Lambda$. Lifting the last equation from *F* to \mathcal{O} and extending scalars from \mathcal{O} to \mathbb{K} , we obtain the equation

$$[\mathbb{K}[C_G(P)e]] = \sum_{(\phi,(Q,f))\in\widetilde{\Lambda}_B} e\mu(P,\phi,Q)f \cdot_{C_H(Q)} (e\mu(P,\phi,Q)f)^{\circ}$$

in $R(\mathbb{K}[C_G(P)]e, \mathbb{K}[C_G(P)]e)$. Since each $e\mu(P, \phi, Q)f \in R(\mathbb{K}[C_G(P)]e, \mathbb{K}[C_H(Q)]f)$ is perfect, we can apply Corollary 8.8 and obtain that there exists a unique element $\lambda = (\phi, (Q, f)) \in \tilde{\Lambda}_B$ such that $e\mu(P, \phi, Q)f \neq 0$ in $R(\mathbb{K}[C_G(P)]e, \mathbb{K}[C_H(Q)]f)$ and that this element is a perfect isometry between $\mathbb{K}[C_G(P)]e$ and $\mathbb{K}[C_H(Q)]f$. The last statement of the lemma follows from Lemma 8.11.

10.4 Lemma Let $(P, e) \in \mathcal{BP}_{\mathcal{O}}(A)$ and set $I := N_G(P, e)$ and $X := N_{I \times I}(\Delta(P)) = \Delta(I)(C_G(P) \times \{1\})$. Consider the set Λ_B of pairs $(\phi, (Q, f))$, where $(Q, f) \in \mathcal{BP}_{\mathcal{O}}(B)$ and $\phi : Q \xrightarrow{\sim} P$ is an isomorphism, together with its $I \times H$ -action from Theorem 7.5(d). For each $\lambda = (\phi, (Q, f)) \in \Lambda_B$ we set

 $J(\lambda):=N_H(Q,f), \quad I(\lambda):=N_{(I,\phi,J(\lambda))}\leqslant I, \quad \text{and} \quad X(\lambda):=N_{I\times J(\lambda)}(\Delta(P,\phi,Q))\,.$

Then, $X * X(\lambda) = X(\lambda)$, and for each $\chi \in Irr(\mathbb{K}X(e \otimes e^*))$, there exists a unique $I \times H$ -orbit of pairs $\lambda = (\phi, (Q, f)) \in \Lambda_B$ such that

$$\chi \stackrel{X,X(\lambda)}{\underset{G}{\cdot}} e\mu(P,\phi,Q)f \neq 0 \quad in \ R(\mathbb{K}[X(\lambda)](e \otimes f^*)) \,.$$

Moreover, for each $\lambda = (\phi, (Q, f)) \in \Lambda_B$ satisfying this condition, one has

$$\chi_{\stackrel{i}{G}}^{X,X(\lambda)} e\mu(P,\phi,Q)f \in \pm \mathrm{Irr}(\mathbb{K}[X(\lambda)](e\otimes f^*))\,.$$

Proof First note that $X * X(\lambda) = X(\lambda)$ for each $\lambda \in \Lambda_B$, by Lemma 6.10(b) with S = I, $T = J(\lambda)$, and $Y = X(\lambda)$. Next, we apply the Brauer construction with respect to $\Delta(P)$ to Equation (17). By Proposition 4.3(b), we have $eA(\Delta(P))e \cong F[C_G(P)]e$ in F_X mod. Thus, by Theorem 7.5(d) applied to S = I, we have

$$[F[C_G(P)]e] = \sum_{\lambda = (\phi, (Q, f)) \in \widehat{\Lambda}_B} \operatorname{ind}_{X'(\lambda)}^X \left(e\overline{\gamma}(P, \phi, Q) f \stackrel{X(\lambda), X(\lambda)^{\circ}}{\overset{\cdot}{H}} f\overline{\gamma}^{\circ}(Q, \phi^{-1}, P)e \right)$$

in $T^{\Delta}(FX(e \otimes e^*))$, where $X'(\lambda) := X(\lambda) * X(\lambda)^{\circ} = \Delta(I(\lambda)) \cdot (C_G(P) \times \{1\}) \leq X$ and $\widehat{\Lambda}_B$ is a set of representatives of the $I \times H$ -orbits of Λ_B . Lifting this equation from F to \mathcal{O} and extending scalars from \mathcal{O} to \mathbb{K} , we obtain

$$[\mathbb{K}[C_G(P)]e] = \sum_{\lambda = (\phi, (Q, f)) \in \widehat{\Lambda}_B} \operatorname{ind}_{X'(\lambda)}^X \left(e\mu(P, \phi, Q) f \stackrel{X(\lambda), X(\lambda)^{\circ}}{\overset{\cdot}{H}} (e\mu(P, \phi, Q) f)^{\circ} \right)$$

in $R(\mathbb{K}X(e \otimes e^*))$. Now let $\chi \in Irr(\mathbb{K}X(e \otimes e^*))$ and apply the group homomorphism

$$\left(\chi \stackrel{X,X}{\stackrel{\cdot}{G}}_{G} -, \chi\right)_{X} \colon R(\mathbb{K}X(e \otimes e^{*})) \to \mathbb{Z}$$

to the last equation. We obtain

$$\left(\chi \overset{X,X}{\underset{G}{\cdot}} [\mathbb{K}C_G(P)e], \chi\right)_X = \sum \left(\chi \overset{X,X}{\underset{G}{\cdot}} \operatorname{ind}_{X'(\lambda)}^X \left(e\mu(P,\phi,Q)f \overset{X(\lambda),X(\lambda)^{\circ}}{\underset{H}{\cdot}} (e\mu(P,\phi,Q)f)^{\circ}\right), \ \chi\right)_X,$$

where the sum runs over elements $\lambda = (\phi, (Q, f)) \in \widehat{\Lambda}_B$. Applying Proposition 6.7(c) to the left hand side and Lemma 6.10(b) to the right hand side of the last equation, we obtain

$$1 = \sum_{\lambda = (\phi, (Q, f)) \in \widehat{\Lambda}_B} \left(\chi_{G}^{X, X(\lambda)} e\mu(P, \phi, Q) f, \chi_{G}^{X, X(\lambda)} e\mu(P, \phi, Q) f \right)_{X(\lambda)}$$

The statements in the lemma are now immediate.

Note that in general restrictions of virtual non-zero characters can vanish. However:

10.5 Corollary Let $(\Delta(P, \phi, Q), e \otimes f^*) \in \mathcal{BP}^{\Delta}_{\mathcal{O}}(A, B)$ and set $I := N_G(P, e)$ and $J := N_H(Q, f)$. Then $e\mu(P, \phi, Q)f \neq 0$ in $R(\mathbb{K}[N_{I \times J}(\Delta(P, \phi, Q))](e \otimes f^*))$

if and only if, after restriction,

$$e\mu(P,\phi,Q)f \neq 0$$
 in $R(\mathbb{K}[C_G(P)]e,\mathbb{K}[C_H(Q)]f)$

Proof Set $Y := N_{I \times J}(\Delta(P, \phi, Q))$ and $\mu := e\mu(P, \phi, Q)f \in R(\mathbb{K}Y(e \otimes f^*))$. We need to show that if $\mu \neq 0$ then also $\operatorname{res}_{C_G(P) \times C_H(Q)}^Y(\mu) \neq 0$. Set $X := N_{I \times I}(\Delta(P)) = \Delta(I) \cdot (C_G(P) \times \{1\})$. Then, following the construction in 7.1, we have $\tilde{X} = \Delta(N_{(I,\phi,J)}) \cdot (C_G(P) \times \{1\})$ (with respect to Y), since $p_1(Y) = N_{(I,\phi,J)}$ by Proposition 2.4(c). Moreover, Lemma 2.2(a) applied to Y implies $\tilde{X} = Y * Y^\circ$. By 7.1 and Proposition 6.7(c) we obtain

$$\left[\mathbb{K}[C_G(P)]e\right] \stackrel{X,Y}{\underset{H}{\cdot}} \mu = \operatorname{res}_{\tilde{X}}^X(\left[\mathbb{K}[C_G(P)]e\right]) \stackrel{\tilde{X},Y}{\underset{H}{\cdot}} \mu = \mu$$

in $R(\mathbb{K}Y(e \otimes f^*))$, since $X * Y = \tilde{X} * Y = Y * Y^{\circ} * Y = Y$. Therefore, $\mathbb{K}[C_G(P)]e$ has an irreducible constituent $\chi \in \operatorname{Irr}(\mathbb{K}X(e \otimes e^*))$ such that

$$\chi \stackrel{X,Y}{\underset{H}{\cdot}} \mu \neq 0 \quad \text{in } R(\mathbb{K}Y(e \otimes f^*)).$$

By Lemma 10.4, $\chi_{H}^{X,Y} \mu \in \pm \operatorname{Irr}(\mathbb{K}X(e \otimes e^*))$. Now, Lemma 8.12 implies that $\operatorname{res}_{C_G(P) \times C_H(Q)}^Y(\mu) \neq 0$.

10.6 Corollary Let $(\Delta(P', \phi', Q'), e' \otimes f'^*) \leq (\Delta(P, \phi, Q), e \otimes f^*) \in \mathcal{BP}^{\Delta}_{\mathcal{O}}(A, B)$. If $e\mu(P, \phi, Q)f \neq 0$ in $R(\mathbb{K}[C_G(P)]e, \mathbb{K}[C_H(Q)]f)$ then $e'\mu(P', \phi', Q')f' \neq 0$ in $R(\mathbb{K}[C_G(P')]e', \mathbb{K}[C_H(Q')]f')$.

Proof We apply Lemma 9.5 to $\overline{\gamma} \in T(F[G \times H])$ and the inclusion of $A \otimes B^*$ -Bauer pairs $(\Delta(P', \phi', Q'), e' \otimes f'^*) \leq (\Delta(P, \phi, Q), e \otimes f^*)$. Condition (i) in Lemma 9.5 is satisfied by Corollary 10.5. And Condition (ii) in Lemma 9.5 is satisfied by Lemma 10.3.

10.7 Lemma Let M be an indecomposable p-permutation (A, B)-bimodule that appears in γ and let $(\Delta(P, \phi, Q), e \otimes f^*) \in \mathcal{BP}_{\mathcal{O}}(M)$. Then $e\mu(P, \phi, Q)f \neq 0$ in $R(\mathbb{K}[C_G(P)]e, \mathbb{K}[C_H(Q)]f)$.

Proof We abbreviate $X := \Delta(P, \phi, Q)$. By Corollary 10.6 we may assume that the $A \otimes B^*$ -Brauer pair $(X, e \otimes f^*)$ is maximal with respect to the property that there exists an indecomposable *p*-permutation (A, B)-bimodule N appearing in γ such that $eN(X)f \neq \{0\}$. Then, by Lemma 9.3(b), we have $0 \neq e\gamma(X)f \in Pr(\mathcal{O}[N_{I \times J}(X)/X])$, where $I := N_G(P, e)$ and $J := N_H(Q, f)$. Since the map κ in 9.1(c) is injective on $Pr(\mathcal{O}[N_{I \times J}(X)/X])$ we have $e\mu(P, \phi, Q)f \neq 0$ in $R(\mathbb{K}[N_{I \times J}(X)/X])$. Since inflation is injective on character groups and commutes with the map κ , we obtain that $e\mu(P, \phi, Q)f \neq 0$ in $R(\mathbb{K}[C_G(P)]e, \mathbb{K}[C_H(Q)]f)$.

The following proposition gives convenient reformulations of being a γ -Brauer pair, see Definition 9.6.

10.8 Proposition Let $(\Delta(P, \phi, Q), e \otimes f^*) \in \mathcal{BP}^{\Delta}_{\mathcal{O}}(A, B)$ and set $I := N_G(P, e)$ and $J := N_H(Q, f)$. Then the following are equivalent:

- (i) $(\Delta(P,\phi,Q), e \otimes f^*) \in \mathcal{BP}_{\mathcal{O}}(\gamma)$, i.e., $e\overline{\gamma}(P,\phi,Q)f \neq 0$ in $T(F[N_{I \times J}(\Delta(P,\phi,Q))](e \otimes f^*))$.
- (ii) $e\overline{\gamma}(P,\phi,Q)f \neq 0$ in $T(F[C_G(P)]e,F[C_H(Q)]f)$.
- (iii) $e\mu(P,\phi,Q)f \neq 0$ in $R(\mathbb{K}[N_{I\times J}(\Delta(P,\phi,Q))](e\otimes f^*)).$
- (iv) $e\mu(P,\phi,Q)f \neq 0$ in $R(\mathbb{K}[C_G(P)]e,\mathbb{K}[C_H(Q)]f)$.
- (v) $e\nu(P,\phi,Q)f \neq 0$ in $R(F[N_{I\times J}(\Delta(P,\phi,Q))](e\otimes f^*)).$
- (vi) $e\nu(P,\phi,Q)f \neq 0$ in $R(F[C_G(P)]e,F[C_H(Q)]f)$.
- (vii) $(\Delta(P, \phi, Q), e \otimes f^*) \in \mathcal{BP}_{\mathcal{O}}(M)$ for some indecomposable module $M \in {}_A$ triv_B appearing in γ .

Proof Clearly, by the diagram in 10.1, each of the conditions (ii)–(vi) implies (i). Similarly, the condition in (vi) implies each of the conditions (i)–(v). Moreover, (iv) implies (vi) by Lemma 10.3. Finally, (vii) implies (iv) by Lemma 10.7, and clearly (i) implies (vii). This completes the proof of the lemma.

10.9 Corollary Let (P, e) be an A-Brauer pair and define Λ_B as in Lemma 10.3. Then there exists a unique H-orbit of pairs $(\phi, (Q, f)) \in \Lambda_B$ with the property that $(\Delta(P, \phi, Q), e \otimes f^*)$ is a γ -Brauer pair.

Proof This follows immediately from Lemma 10.3 and the equivalence of (i) and (iv) in Proposition 10.8.

The following Theorem shows that every p-permutation equivalence between A and B determines a bijection between the block direct summands of A and of B, and that it is the sum of p-permutation equivalences between corresponding blocks.

10.10 Theorem Assume that $\gamma \in T_l^{\Delta}(A, B)$. Let \mathcal{I} denote the set of primitive idempotents of Z(A) and let \mathcal{J} denote the set of primitive idempotents of Z(B). For each $e \in \mathcal{I}$ there exists a unique f in \mathcal{J} such that $e\gamma f \neq 0$ in $T^{\Delta}(\mathcal{O}Ge, \mathcal{O}Hf)$. If $e \in \mathcal{I}$ and $f \in \mathcal{J}$ satisfies $e\gamma f \neq 0$ then $e\gamma f \in T_l^{\Delta}(\mathcal{O}Ge, \mathcal{O}Hf)$.

In particular, if γ is a p-permutation equivalence between A and B, then the condition $e\gamma f \neq 0$ defines a bijection between \mathcal{I} and \mathcal{J} , and if $e\gamma f \neq 0$ then $e\gamma f$ is a p-permutation equivalence between $\mathcal{O}Ge$ and $\mathcal{O}Hf$. Moreover, $\gamma = \sum_{e \in \mathcal{I}} e\gamma f$, where $f \in \mathcal{J}$ corresponds to e.

Proof The first statement follows immediately from Corollary 10.9 applied to the A-Brauer pair ({1}, e). Thus, $\gamma = \sum_{e \in \mathcal{I}} e\gamma f$, where $f \in \mathcal{J}$ denotes the element corresponding to e. Suppose that $e\gamma f \neq 0$. Then $e\gamma = e\gamma f$ and multiplying the equation $\gamma \cdot_{H} \gamma^{\circ} = [A]$ with e from left and right yields $e\gamma f \cdot_{H} (e\gamma f)^{\circ} = [\mathcal{O}Ge]$. Thus, $e\gamma f \in T_{l}^{\Delta}(\mathcal{O}Ge, \mathcal{O}Hf)$.

We are now ready to prove the main theorem of this section. It shows that γ -Brauer pairs behave very similar to *M*-Brauer pairs of an indecomposable *p*-permutation module *M*, cf. Proposition 5.3(b).

10.11 Theorem (a) The set of γ -Brauer pairs form a $G \times H$ -stable ideal in the poset of $A \otimes B^*$ -Brauer pairs.

- (b) If A and B are blocks then any two maximal γ -Brauer pairs are $G \times H$ -conjugate.
- (c) For $(\Delta(P, \phi, Q), e \otimes f^*) \in \mathcal{BP}^{\Delta}_{\mathcal{O}}(\gamma)$ the following are equivalent:
 - (i) $(\Delta(P, \phi, Q), e \otimes f^*)$ is a maximal γ -Brauer pair;
 - (ii) (P, e) is a maximal A-Brauer pair;
 - (iii) (Q, f) is a maximal *B*-Brauer pair.

Proof (a) Clearly, the set of γ -Brauer pairs is closed under $G \times H$ -conjugation. Moreover, by Corollary 10.6 and Proposition 10.8, the set of γ -Brauer pairs is an ideal in the poset of $A \otimes B^*$ -Brauer pairs.

(b) Now assume that A and B are blocks. Let (D, e_D) be a maximal A-Brauer pair over \mathcal{O} . By Corollary 10.9 there exists a B-Brauer pair (E, f_E) over \mathcal{O} and an isomorphism $\psi \colon E \xrightarrow{\sim} D$ such that $(\Delta(D, \psi, E), e_D \otimes f_E^*)$ is a γ -Brauer pair. Let $(\Delta(P', \phi', Q'), e' \otimes f'^*)$ be an arbitrary γ -Brauer pair. We will first prove that

$$(\Delta(P',\phi',Q'),e'\otimes f'^*) \leqslant_{G\times H} (\Delta(D,\psi,E),e_D\otimes f_E^*).$$
(18)

Since any two maximal A-Brauer pairs are G-conjugate, we may assume that $(P', e') \leq (D, e_D)$. Set $R := \psi^{-1}(P') \leq \psi^{-1}(D) = E$ and let f_R denote the unique block idempotent of $\mathcal{O}[C_H(R)]$ such that $(R, f_R) \leq (E, f_E)$. Since $(\Delta(D, \psi, E), e_D \otimes f_E^*)$ is a γ -Brauer pair, Part (a) implies that also $(\Delta(P', \psi|_R, R), e' \otimes f_R^*)$ is a γ -Brauer pair. Since also $(\Delta(P', \phi', Q'), e' \otimes f'^*)$ is a γ -Brauer pair, Corollary 10.9 implies that $(\psi|_R, (R, f_R))$ and $(\phi', (Q', f'))$ are H-conjugate. Thus,

$$(\Delta(P',\phi',Q'),e'\otimes f'^*) =_{\{1\}\times H} (\Delta(P',\psi|_R,R),e'\otimes f^*_R) \leqslant (\Delta(D,\psi,E),e_D\otimes f^*_E)$$

and the claim is proven. This implies that $(\Delta(D, \psi, E), e_D \otimes f_E^*)$ is a maximal γ -Brauer pair, and also that every other maximal γ -Brauer pair is $G \times H$ -conjugate to $(\Delta(D, \psi, E), e_D \otimes f_E^*)$.

(c) Let A and B be again sums of blocks. First recall from Remark 10.2(b) that every γ -Brauer pair has a twisted diagonal subgroup as first component. Thus, (ii) implies (i) and (iii) implies (i), by the last statement in Remark 10.2(a).

In order to see that (i) implies (ii) and (iii) we claim that it suffices to show this in the situation where A and B are blocks. In fact, $(\Delta(P,\phi,Q), e \otimes f^*)$ is an $A' \otimes B'^*$ -Brauer pair for the unique block direct summands A' of A and B' of B, respectively, that satisfy $(\{1\}, e_{A'}) \leq (P, e)$ and $(\{1\}, e_{B'}) \leq (Q, f)$. Applying Corollary 10.9 to the A-Brauer pair $(\{1\}, e_{A'})$, we see that B' is uniquely determined by A'. Writing γ as the sum of the elements $e'\gamma f'$, where e' and f' run through the block idempotents of $\mathcal{O}G$ and $\mathcal{O}H$ with $e_A e' = e'$ and $e_B f' = f'$, then multiplying both sides of (17) by $e_{A'}$ from the left and right, we see that $(e_{A'}\gamma e_{B'}) \stackrel{\cdot}{\to} (e_{A'}\gamma e_{B'})^\circ = [A']$ in

 $T^{\Delta}(A', A')$. Since $(\Delta(P, \phi, Q), e \otimes f^*)$ is a maximal γ -Brauer pair, it is also a maximal $e_{A'}\gamma e_{B'}$ -Brauer pair.

Since maximal A'-Brauer pairs (resp. B'-Brauer pairs) are also maximal A-Brauer pairs (resp. B-Brauer pairs), we may assume from now on that A and B are blocks.

Now let (D, e_D) be a maximal A-Brauer pair and let the γ -Brauer pair $(\Delta(D, \psi, E), e_D \otimes f_E^*)$ be chosen as in the proof of Part (b). Then, by the claim proved there, we know that $(\Delta(D, \psi, E), e_D \otimes f_E^*)$ is a maximal γ -Brauer pair. Thus, by Part (b), $(\Delta(P, \phi, Q), e \otimes f^*) \leq_{G \times H} (\Delta(D, \psi, E), e_D \otimes f_E^*)$. This already shows that (i) implies (ii). Finally, in order to show that (i) implies (iii), it suffices to show that (E, f_E) is a maximal *B*-Brauer pair, i.e., that *E* is a defect group of *B*.

So let q denote the order of a defect group of B. It suffices to show that $q \leq |E|$, since (E, f_E) is a B-Brauer pair. Set $\mu := \kappa_{G \times H}(\gamma) \in R(\mathbb{K}G, \mathbb{K}H)$. Then $\mu \underset{H}{\cdot} \mu^{\circ} = [\mathbb{K}Ge_A]$ in $R(\mathbb{K}G, \mathbb{K}G)$. Lemma 8.7 now implies that μ is an isometry between $\mathbb{K}Ge_A$ and $\mathbb{K}He_B$. This implies further that

$$\operatorname{rk}_{\mathcal{O}}(B) = \dim_{\mathbb{K}}(\mathbb{K}He_B) = \dim_{\mathbb{K}}(\mu^{\circ} \cdot \mu) = \operatorname{rk}_{\mathcal{O}}(\gamma^{\circ} \cdot \gamma), \qquad (19)$$

where $\dim_{\mathbb{K}}$ and $\mathrm{rk}_{\mathcal{O}}$ also denote the integer-valued maps induced on the Grothendieck groups. Since every γ -Brauer pair is contained in a $G \times H$ -conjugate of $(\Delta(D, \psi, E), e_D \otimes f_E^*)$, Lemma 10.7 and Proposition 10.8 imply that every indecomposable $A \otimes B^*$ -module that appears in γ has a vertex contained in $\Delta(D, \psi, E)$. Therefore, every indecomposable $B \otimes A^*$ -module that appears in γ° has a vertex contained in $\Delta(E, \psi^{-1}, D)$. The Mackey formula for bimodules, see Theorem 6.2, implies that each indecomposable $B \otimes B^\circ$ -module that appears in $\gamma^\circ \cdot_G \gamma$ has a vertex of order dividing |E|. By Theorem 4.7.5 in [NT89], the *p*-part of the \mathcal{O} -rank of each such indecomposable $B \otimes B^\circ$ -module is a multiple of $|H|_p^2/|E|$. Thus, using Equation (19), $\mathrm{rk}_{\mathcal{O}}(B)_p$ is a multiple of $|H|_p^2/|E|$. On the other hand, by Theorem 5.10.1 in [NT89] we know that $\mathrm{rk}_{\mathcal{O}}(B)_p = |H|_p^2/q$. This implies that *q* divides |E| and the proof of Part (c) is complete.

We conclude this section by proving an inverse to the association constructed in Corollary 10.9. This will be used in Section 11 to show that if $\gamma \in T^{\Delta}(A, B)$ satisfies (17) then the fusion systems of A and B are isomorphic.

10.12 Lemma Let $(Q, f) \in \mathcal{BP}_{\mathcal{O}}(B')$ for a block B' of \mathcal{OH} satisfying $\gamma e_{B'} \neq 0$. Then there exists a unique G-conjugacy class of pairs $((P, e), \phi)$, where $(P, e) \in \mathcal{BP}_{\mathcal{O}}(A)$ and $\phi: Q \xrightarrow{\sim} P$ is an isomorphism, such that $(\Delta(P, \phi, Q), e \otimes f^*)$ is a γ -Brauer pair. Here, G acts on the set of such pairs via ${}^{g}((P, e), \phi) = (({}^{g}P, {}^{g}e), c_{g} \circ \phi)$.

Proof We first show the existence part of the lemma. There exists a block direct summand A' of A such that $e_{A'}\gamma e_{B'} \neq 0$ in $T(\mathcal{O}G, \mathcal{O}H)$. In other words, $(\{1\}, e_{A'} \otimes e_{B'}^*)$ is a γ -Brauer pair. Let $(\Delta(D, \psi, E), e_D \otimes f_E^*)$ be a maximal γ -Brauer pair containing $(\{1\}, e_{A'} \otimes e_{B'})$, then $(\Delta(D, \psi, E), e_D \otimes f_E^*)$ is also an $A' \otimes B'^*$ -Brauer pair and (E, f_E) is a maximal B'-Brauer pair by Theorem 10.11(c). After conjugating this maximal γ -Brauer pair by an element in $\{1\} \times H$, if necessary, we may assume that $(Q, f) \leq (E, f_E)$. Setting $P := \psi(Q)$ and $\phi := \psi|_Q : Q \xrightarrow{\sim} P$, there exists a unique primitive idempotent e of $Z(\mathcal{O}[C_G(P)])$ such that $(P, e) \leq (D, e_D)$. This implies that $(\Delta(P, \phi, Q), e \otimes f^*) \leq (\Delta(D, \psi, E), e_D \otimes f_E^*)$. Since the set of γ -Brauer pairs is an ideal, see Theorem 10.11(a), also $(\Delta(P, \phi, Q), e \otimes f^*)$ is a γ -Brauer pair.

Next, we show the uniqueness part. Consider $e\gamma(P,\phi,Q)f$ as element in $T^{\Delta}(\mathcal{O}[C_G(P)]e, \mathcal{O}[C_H(Q)]f)$ and $e\mu(P,\phi,Q)f$ as element in $R(\mathbb{K}[C_G(P)]e,\mathbb{K}[C_H(Q)]f)$. Since $\gamma \underset{H}{\cdot} \gamma^{\circ} = [A]$ in $T^{\Delta}(A,A)$, we obtain $\gamma \underset{H}{\cdot} \gamma^{\circ} \underset{G}{\cdot} \gamma = \gamma$ in $T^{\Delta}(A,B)$ and

$$e\gamma(P,\phi,Q)f = e(\gamma \underset{H}{\cdot} \gamma^{\circ} \underset{G}{\cdot} \gamma)(\Delta(P,\phi,Q))f = e\gamma(P,\phi,Q)f \underset{C_{H}(Q)}{\cdot} f(\gamma^{\circ} \underset{G}{\cdot} \gamma)(\Delta(Q))f,$$

in $T^{\Delta}(\mathcal{O}[C_G(P)]e, \mathcal{O}[C_H(Q)]f)$, where the last equation follows from Theorem 7.5(b) and Corollary 10.9. Extending scalars from \mathcal{O} to \mathbb{K} , the last equation implies

$$e\mu(P,\phi,Q)f = e\mu(P,\phi|_Q,Q)f \underset{C_H(Q)}{\cdot} \left(f(\gamma^{\circ} \underset{G}{\cdot} \gamma)(\Delta(Q))f\right)^{\mathbb{K}}$$

in $R(\mathbb{K}[C_G(P)]e, \mathbb{K}[C_H(Q)]f)$. By Lemma 10.3, $e\mu(P, \phi|_Q, Q)f$ is an isometry between $\mathbb{K}[C_G(P)]e$ and $\mathbb{K}[C_H(Q)]f$. Thus, the last equation implies

$$[\mathbb{K}[C_H(Q)]f] = \left(f(\gamma^\circ \mathop{\cdot}_G \gamma)(\Delta(Q))f\right)^{\mathbb{K}}$$

in $R(\mathbb{K}[C_H(Q)]f,\mathbb{K}[C_H(Q)]f)$. Using again Theorem 7.5(b), we can write

$$f(\gamma^{\circ} \underset{G}{\cdot} \gamma)(\Delta(Q))f = \sum_{((P',e'),\phi')} f\gamma^{\circ}(\Delta(Q,\phi'^{-1},P'))e' \underset{G}{\cdot} e'\gamma(P',\phi',Q)f$$

in $T^{\Delta}(\mathcal{O}[C_H(Q)]f, \mathcal{O}[C_H(Q)]f)$, where $((P', e'), \phi')$ runs through representatives of the *G*-conjugacy classes of pairs as described in the statement of the lemma. Thus,

$$[\mathbb{K}[C_H(Q)]f] = \sum_{((P',e'),\phi')} f\mu(P',\phi',Q)^{\circ}e' \mathop{\cdot}_{G} e'\mu(P',\phi',Q)f$$

in $R(\mathbb{K}[C_H(Q)]f,\mathbb{K}[C_H(Q)]f)$. Corollary 8.8 now implies that, up to *G*-conjugacy, there exists a unique pair $((P', e'), \phi')$ such that $e'\mu(P', \phi', Q)f \neq 0$ in $R(\mathbb{K}[C_G(P)]e', \mathbb{K}[C_H(Q)]f)$. Together with Proposition 10.8 this implies the desired uniqueness statement.

11 Fusion systems, local equivalences and finiteness

Throughout this section we assume again that G and H are finite groups, and that the *p*-modular system $(\mathbb{K}, \mathcal{O}, F)$ is large enough for G and H. Moreover, we assume that $A = \mathcal{O}Ge_A$ is a block of $\mathcal{O}G$, $B = \mathcal{O}He_B$ is a block of $\mathcal{O}H$, and that $\gamma \in T_l^{\Delta}(A, B)$ is a *left p*-permutation equivalence between A and B, i.e., $\gamma : \gamma^{\circ} = [A]$ in T(A, A). Furthermore, we assume the notation from 10.1 and that $(\Delta(D, \phi, E), e_D \otimes f_E^*)$ is a maximal γ -Brauer pair. Then, by Theorem 10.11(c), (D, e_D) is a maximal A-Brauer pair and (E, f_E) is a maximal B-Brauer pair. By \mathcal{A} we denote the fusion system of A associated with (D, e_D) and by \mathcal{B} we denote the fusion system of B associated with (E, f_E) .

In Theorem 11.2 we show that ϕ is an isomorphism between the fusion systems \mathcal{B} and \mathcal{A} . And in Theorem 11.4 we show that Brauer constructions with respect to subgroups of $\Delta(D, \phi, E)$ applied to γ yields again local *p*-permutation equivalences at various levels. Finally in Theorem 11.10 we show that there can only be finitely many *p*-permutation equivalences between given blocks *A* and *B*.

11.1 Proposition Let $(\Delta(P, \psi, Q), e \otimes f^*)$ be a γ -Brauer pair and set $I := N_G(P, e), J := N_H(Q, f)$. For every $g \in I$ there exists a unique element $hC_H(Q) \in J/C_H(Q)$ such that $c_g \circ \psi = \psi \circ c_h : Q \xrightarrow{\sim} P$. Similarly, for every $h \in J$ there exists a unique element $gC_G(P) \in I/C_G(P)$ such that $c_g \circ \psi = \psi \circ c_h$. These associations define mutually inverse group isomorphisms between $I/C_G(P)$ and $J/C_H(Q)$. The isomorphism $J/C_H(Q) \xrightarrow{\sim} I/C_G(P)$ restricts to the isomorphism $QC_H(Q)/C_H(Q) \xrightarrow{\sim} PC_G(P)/P$, $hC_H(Q) \mapsto \psi(h)C_G(P)$, for $h \in Q$. The group $Y := N_{G \times H}(\Delta(P, \psi, Q), e \otimes f^*)$ satisfies $p_1(Y) = I$, $p_2(Y) = J$, $k_1(Y) = C_G(P)$, $k_2(Y) = C_H(Q)$ and the resulting isomorphism $\eta_Y : J/C_H(Q) \xrightarrow{\sim} I/C_G(P)$ from 2.1(a) is equal to the one described above.

Proof Let $g \in I$. With $(\Delta(P, \psi, Q), e \otimes f^*)$ also ${}^{(g,1)}(\Delta(P, \psi, Q), e \otimes f^*) = (\Delta(P, c_g \circ \psi, Q), e \otimes f^*)$ is a γ -Brauer pair. Now, the uniqueness statement in Corollary 10.9 implies that there exists an element $h \in H$ such that ${}^{h}(c_g \circ \psi, (Q, f)) = (\psi, (Q, f))$. This implies that $h \in J$ and that $c_g \circ \psi = \psi \circ c_h$. Clearly, $h \in J$ is uniquely determined up to multiplication with elements of $C_H(Q) \leq J$ by this condition. It is easy to see that this defines a group homomorphism from $I/C_G(P)$ to $J/C_H(Q)$. A similar argument, now using Lemma 10.12, implies the second statement and defines a group homomorphism from $J/C_H(Q)$ to $I/C_G(P)$. Clearly, these two homomorphisms are mutually inverses. The following statement is clear, since $\psi c_h \psi^{-1} = c_{\psi(h)}$. The last statements about Y follow immediately from the above and Propostion 2.4(c).

Note that for the following theorem we only assume that $\gamma \in T_l^{\Delta}(A, B)$.

11.2 Theorem The isomorphism $\phi: E \xrightarrow{\sim} D$ is an isomorphism between the fusion systems \mathcal{B} and \mathcal{A} .

Proof Using Alperin's fusion theorem, see [AKO11, Theorem I.3.6], it suffices to show that for every subgroup $Q \leq E$ with $P := \phi(Q) \leq D$ and $\psi := \phi|_Q : Q \xrightarrow{\sim} P$, one has $\psi^{-1} \circ \operatorname{Hom}_{\mathcal{A}}(P,P) \circ \psi = \operatorname{Hom}_{\mathcal{B}}(Q,Q)$, an equation of sets of automorphisms of Q. Let $e_P \in Z(\mathcal{O}[C_G(P)])$ and $f_Q \in Z(\mathcal{O}[C_H(Q)])$ be the unique primitive idempotents such that $(P, e_P) \leq (D, e_D)$ and $(Q, f_Q) \leq (E, f_E)$, and set $I := N_G(P, e_P)$ and $J := N_H(Q, f_Q)$. Note that $I/C_G(P) \to \operatorname{Hom}_{\mathcal{A}}(P,P)$, $gC_G(P) \mapsto c_g$, and $J/C_H(Q) \to \operatorname{Hom}_{\mathcal{B}}(Q,Q)$, $hC_H(Q) \mapsto c_h$, are group isomorphisms. Now the claim follows immediately from Proposition 11.1, noting that $(\Delta(P, \psi, Q), e_P \otimes$

 f_Q^* $\leq (\Delta(D, \phi, E), e_D \otimes f_E^*)$, by Remark 10.2(c), so that also $(\Delta(P, \psi, Q), e_P \otimes f_Q^*)$ is a γ -Brauer pair by Theorem 10.11(a).

We can now improve the formulation of Lemma 10.4. Recall the definition of $\mu(P, \phi, Q)$ from 10.1.

11.3 Lemma Let $(\Delta(P, \psi, Q), e \otimes f^*) \in \mathcal{BP}^{\Delta}_{\mathcal{O}}(\gamma)$ and set $I := N_G(P, e), J := N_H(Q, f), X := N_{I \times I}(\Delta(P)) \leq G \times G$ and $Y := N_{I \times J}(\Delta(P, \psi, Q)) \leq G \times H$. Then X * Y = Y and for every $\chi \in \operatorname{Irr}(\mathbb{K}X(e \otimes e^*))$ one has

$$\chi \stackrel{X,Y}{\underset{G}{\cdot}} e\mu(P,\psi,Q)f \in \pm \operatorname{Irr}(\mathbb{K}Y(e \otimes f^*)) \,.$$

Proof By Proposition 11.1, we have $p_1(Y) = I$. Therefore, Lemma 2.2(c) implies that X * Y = Y. By Lemma 10.4 there exists a *B*-Brauer pair (Q', f') and an isomorphism $\psi' \colon Q' \xrightarrow{\sim} P$ such that $0 \neq \chi_{\dot{G}}^{X,Y'} e_{\mu}(P,\psi',Q')f' \in R(\mathbb{K}Y(e \otimes f'^*))$, where $Y' \coloneqq N_{I \times J'}(\Delta(P,\psi',Q'))$ with $J' \coloneqq N_H(Q',f')$. Therefore, by Proposition 10.8, $(\Delta(P,\psi',Q'), e \otimes f'^*)$ is a γ -Brauer pair. By Lemma 10.3 there exists $h \in H$ such that $(\Delta(P,\psi,Q), e \otimes f^*) = {}^{(1,h)}(\Delta(P,\psi',Q'), e \otimes f'^*)$. This, together with Equation (4), implies

$$\chi_{\dot{G}}^{X,Y} e\mu(P,\psi,Q)f = \chi_{\dot{G}}^{X,Y} {}^{(1,h)} (e\mu(P,\psi',Q')f') = {}^{(1,h)} (\chi_{\dot{G}}^{X,Y'} e\mu(P,\psi',Q')f') \neq 0$$

in $R(\mathbb{K}X(e \otimes f^*))$, since ${}^{(1,h)}Y' = Y$. By Lemma 10.4, this virtual character belongs to $\pm \operatorname{Irr}(\mathbb{K}Y(e \otimes f^*))$.

11.4 Theorem Let $(\Delta(P, \psi, Q), e \otimes f^*) \in \mathcal{BP}^{\Delta}_{\mathcal{O}}(\gamma)$ be a γ -Brauer pair and set $I := N_G(P, e)$ and $J := N_H(Q, f)$. Suppose that $C_G(P) \leq S \leq I$ and $C_H(Q) \leq T \leq J$ are intermediate groups related via the isomorphism in Proposition 11.1 and set $Y := N_{S \times H}(\Delta(P, \psi, Q), e \otimes f^*) = N_{S \times T}(\Delta(P, \psi, Q))$. Then, the element $e\gamma(P, \psi, Q)f \in T(\mathcal{OY}(e \otimes f^*))$, defined as in 10.1 and restricted to Y, satisfies

$$\mathcal{O}[C_G(P)]e] = e\gamma(P,\psi,Q)f \stackrel{Y,Y^\circ}{\stackrel{\cdot}{H}} f\gamma(P,\psi,Q)^\circ e \quad \text{in } T(\mathcal{O}[N_{S\times S}(\Delta(P))](e\otimes e^*)).$$
(20)

Moreover, the element

$$\tilde{\gamma} := \operatorname{ind}_Y^{S \times T} \left(e \gamma(P, \psi, Q) f \right) \in T^{\Delta}(\mathcal{O}Se, \mathcal{O}Tf)$$

satisfies $\tilde{\gamma}_{T} \tilde{\gamma}^{\circ} = [\mathcal{O}Se]$ in $T^{\Delta}(\mathcal{O}Se, \mathcal{O}Se)$. In particular, if γ is a *p*-permutation equivalence between A and B then $\tilde{\gamma}$ is a *p*-permutation equivalence between $\mathcal{O}Se$ and $\mathcal{O}Tf$.

Proof We apply the Brauer construction with respect to $\Delta(P)$ to Equation (17) and obtain after restriction

$$[\mathcal{O}[C_G(P)]e] = e(\gamma \underset{H}{\cdot} \gamma^{\circ})(\Delta(P))e \quad \text{in } T(\mathcal{O}[N_{S\times S}(\Delta(P))](e\otimes e^*)).$$

Next we apply Theorem 7.5(d) to the right hand side of the last equation. Note that $p_1(Y) = S$ and $p_2(Y) = T$ by Propositions 2.4(c) and 11.1. Thus, $Y * Y^\circ = \Delta(S)(C_G(P) \times \{1\}) = N_{S \times S}(\Delta(P))$ by Lemma 2.2 and Proposition 2.4. By Corollary 10.9, the pair $(\psi, (Q, f))$ is, up to *H*-conjugation, the only pair such that $e\gamma(P, \psi, Q)f \neq 0$. Thus, Theorem 7.5(d) implies Equation (20).

Next we apply $\operatorname{ind}_{N_{S\times S}(\Delta(P))}^{S\times S}$ to both sides of Equation (20). First we show that the left hand side yields $[\mathcal{O}Se] \in T^{\Delta}(\mathcal{O}Se, \mathcal{O}Se)$. In fact, $\mathcal{O}[C_G(P)]$ is an $\mathcal{O}[N_{S\times S}(\Delta(P))]$ -permutation module with $\Delta(S)$ as stabilizer of the standard basis element $1 \in C_G(P)$. Thus, $\mathcal{O}[C_G(P)] \cong \operatorname{Ind}_{\Delta(S)}^{N_{S\times S}(\Delta(P))}(\mathcal{O})$ in $T(\mathcal{O}[N_{S\times S}(\Delta(P))])$ and 3.1(f), applied to the central idempotent $e \otimes e^*$ of $\mathcal{O}[N_{S\times S}(\Delta(P))]$, implies that

$$\operatorname{Ind}_{N_{S\times S}(\Delta(P))}^{S\times S}(\mathcal{O}[C_G(P)]e) \cong e\left(\operatorname{Ind}_{N_{S\times S}(\Delta(P))}^{S\times S}(\mathcal{O}[C_G(P)])\right)e \cong e\operatorname{Ind}_{\Delta(S)}^{S\times S}(\mathcal{O})e \cong e\mathcal{O}Se = \mathcal{O}Se$$

as $(\mathcal{O}Se, \mathcal{O}Se)$ -bimodules. Next we show that applying $\operatorname{ind}_{N_{S\times S}(\Delta(P))}^{S\times S}$ to the right hand side of Equation (20) yields

$$\operatorname{ind}_{Y}^{S \times T}(e\gamma(P,\psi,Q)f) \underset{T}{\cdot} (\operatorname{ind}_{Y}^{S \times T}(e\gamma(P,\psi,Q)f))^{\circ}$$

$$\tag{21}$$

In fact, rewriting $(\operatorname{ind}_Y^{S \times T}(e\gamma(P,\psi,Q)f))^{\circ} = \operatorname{ind}_{Y^{\circ}}^{T \times S}(f\gamma(P,\psi,Q)^{\circ}e)$ in the above expression, then applying the Mackey formula in Theorem 6.2 to the resulting expression and noting that $p_2(Y) = T$ and $Y * Y^{\circ} = N_{S \times S}(\Delta(P))$, one obtains that the expression in (21) is equal to

$$\operatorname{ind}_{N_{S\times S}(\Delta(P))}^{S\times S} (e\gamma(P,\psi,Q) f \stackrel{Y,Y^{\circ}}{\underset{H}{\cdot}} f\gamma(P,\psi,Q)^{\circ} e \Big)$$

in $T(\mathcal{O}Se, \mathcal{O}Se)$ as desired.

The following proposition investigates maximal Brauer pairs of the local p-permutation equivalences from Theorem 11.4 for the two extreme choices of S and T. It will use the notions introduced in 4.4 and the results from Proposition 4.5.

11.5 Proposition For any subgroup P of D let (P, e_P) denote the unique A-Brauer pair with $(P, e_P) \leq (D, e_D)$ and for any subgroup Q of E let (Q, f_Q) denote the unique B-Brauer pair with $(Q, f_Q) \leq (E, f_E)$. Let $Q \leq E$ and set $P := \phi(Q)$. Then $(\Delta(P, \phi, Q), e_P \otimes f_Q^*)$ is a γ -Brauer pair (cf. Remark 10.2(c) and Theorem 10.11(a)).

(a) Set $\gamma' := e_P \gamma(P, \phi, Q) f_Q \in T_l^{\Delta}(\mathcal{O}[C_G(P)]e_P, \mathcal{O}[C_H(Q)]f_Q)$. The $\mathcal{O}[C_G(P)]e_P \otimes \mathcal{O}[C_H(Q)]f_Q^*$ -Brauer pair $(\Delta(C_D(P), \phi, C_E(Q)), e_{PC_D(P)} \otimes f_{QC_E(Q)}^*)$ is a γ' -Brauer pair. It is a maximal γ' -Brauer pair if and only if P is fully A-centralized. In particular, $(\Delta(Z(P), \phi, Z(Q)), e_P \otimes f_Q^*)$ is a maximal γ' -Brauer pair if and only if P is A-centric.

(b) Set $I := N_G(P, e_P)$, $J := N_H(Q, f_Q)$ and $\gamma'' := \operatorname{ind}_{N_I \times J}^{I \times J}(\Delta(P, \phi, Q))}(e_P \gamma(P, \phi, Q) f_Q) \in T_l^{\Delta}(\mathcal{O}Ie_P, \mathcal{O}Jf_Q)$. The $\mathcal{O}Ie_P \otimes \mathcal{O}Jf_Q^*$ -Brauer pair $(\Delta(N_D(P), \phi, N_E(Q)), e_{N_D(P)} \otimes f_{N_E(Q)}^*)$ is a γ'' -Brauer pair. It is a maximal γ'' -Brauer pair if and only if P is fully \mathcal{A} -normalized.

Proof By Theorem 11.4, we have $\gamma' \in T_l^{\Delta}(\mathcal{O}[C_G(P)]e_P, \mathcal{O}[C_H(Q)]f_Q)$ and $\gamma'' \in T_l^{\Delta}(\mathcal{O}Ie_P, \mathcal{O}Jf_Q)$, so that we can apply results from Section 10 to γ' and to γ'' .

(a) First note that $(\Delta(C_D(P), \phi, C_E(Q)), e_{PC_D(P)} \otimes f^*_{QC_E(Q)})$ is an $\mathcal{O}[C_G(P)]e_P \otimes \mathcal{O}[C_H(Q)]f^*_Q$ -Brauer pair, since $C_{C_G(P)}(C_D(P)) = C_G(P) \cap C_G(C_D(P)) = C_G(PC_D(P))$ and $C_{C_H(Q)}(C_E(Q)) = C_H(QC_E(Q))$. It is a γ' -Brauer pair, since

$$e_{PC_D(P)}\gamma'(\Delta(C_D(P),\phi,C_E(Q)))f_{QC_E(Q)} = e_{PC_D(P)}\gamma(\Delta(PC_D(P),\phi,EC_E(Q)))f_{QC_E(Q)} \neq 0$$

in $T^{\Delta}(\mathcal{O}[C_G(PC_D(P))]e_{PC_D(P)}, \mathcal{O}[C_H(QC_E(Q))]f_{QC_E(Q)})$ by Proposition 3.5(b) and Lemma 3.7, and since $(\Delta(PC_D(P), \phi, EC_E(Q)), e_{PC_D(P)} \otimes f^*_{QC_E(Q)})$ is a γ -Brauer pair by Theorem 10.11(a) and Proposition 10.8. By Theorem 10.11(c), $(\Delta(C_D(P), \phi, C_E(Q)), e_{PC_D(P)} \otimes f^*_{QC_E(Q)})$ is a maximal γ' -Brauer pair if and only if $C_D(P)$ is a defect group of $\mathcal{O}[C_G(P)]e_P$. But, by Proposition 4.5(a), this is equivalent to P being fully \mathcal{A} -centralized.

(b) Note that $(\Delta(N_D(P), \phi, N_E(Q)), e_{N_D(P)} \otimes f_{N_E(Q)}^*)$ is an $\mathcal{O}Ie_P \otimes \mathcal{O}Jf_Q^*$ -Brauer pair, since $C_I(N_D(P)) = C_G(N_D(P))$ and $C_J(N_E(Q)) = C_G(N_E(Q))$. It is a straightforward verification that the *p*-subgroups $\Delta(P, \phi, Q)$ and $\Delta(N_D(P), \phi, N_E(Q))$ of $I \times J$ satisfy the hypothesis of Lemma 3.9 (in place of the subgroups Q and P of G), since every indecomposable $\mathcal{O}[N_{I \times J}(\Delta(P, \phi, Q))]$ -module appearing in $e_P\gamma(P, \phi, Q)f_Q$ has a vertex contained in $\Delta(D, \phi, E)$ by Lemma 10.7, Proposition 10.8, and Theorem 10.11. Thus, Lemma 3.9 together with Lemma 3.7 implies

$$e_{N_D(P)}\gamma''(N_D(P),\phi,N_E(Q))f_{N_E(Q)} = e_{N_D(P)}\gamma(N_D(P),\phi,N_E(Q))f_{N_E(Q)}$$

in $T^{\Delta}(\mathcal{O}[C_G(N_D(P))]e_{N_D(P)}, \mathcal{O}[C_H(N_E(Q))]f_{N_H(Q)})$. Since $(\Delta(N_D(P), \phi, N_E(Q)), e_{N_D(P)} \otimes f^*_{N_E(Q)})$ is a γ -Brauer pair by Theorem 10.11(a), the element in the above equation is non-zero. This implies that $(\Delta(N_D(P), \phi, N_E(Q)), e_{N_D(P)} \otimes f^*_{N_E(Q)})$ is a γ'' -Brauer pair. By Theorem 10.11(c), it is a maximal γ'' -Brauer pair if and only if $N_D(P)$ is a defect group of $\mathcal{O}Ie_P$. But, by Proposition 4.5(b), this is equivalent to P being fully \mathcal{A} -normalized.

11.6 Schur classes. Let \Bbbk be a field. Recall from the proof of [NT89, Theorem 3.5.7] that whenever $N \leq G$ and V is an irreducible G-stable $\Bbbk N$ -module which is \Bbbk -split, i.e., $\operatorname{End}_{\Bbbk N}(V) = \Bbbk$, Schur assigned to these data a canonical cohomology class $\kappa \in H^2(G/N, \Bbbk^{\times})$. This construction has the following properties:

- (a) V extends to a &G-module if and only if $\kappa = 1$;
- (b) The class assigned to V° is κ^{-1} ;

(c) If $N \leq H \leq G$ then the canonical class assigned to $N \leq H$ and V is $\operatorname{res}_{H/N}^{G/N}(\kappa)$; and

(d) If κ_i is the class assigned to $N_i \leq G_i$ and V_i , for i = 1, 2, then $\kappa_1 \times \kappa_2 \in H^2((G_1 \times G_1)/(N_1 \times N_2), F^{\times})$ is the canonical class assigned to $N_1 \times N_2 \leq G_1 \times G_2$ and $V_1 \otimes_{\Bbbk} V_2$.

11.7 Lemma Let \Bbbk be a field, let G and H be finite groups, and let $Y \leq G \times H$ be such that $p_1(Y) = G$ and $p_2(Y) = H$. Set $M := k_1(Y) \trianglelefteq G$, $N := k_2(Y) \trianglelefteq H$ and let $\eta_Y : H/N \xrightarrow{\sim} G/M$ (see 2.1(a)) be the isomorphism induced by Y. Suppose that $V \in {}_{\Bbbk M}$ mod is irreducible, G-stable, and \Bbbk -split, and that $W \in {}_{\Bbbk N}$ mod is irreducible, H-stable and \Bbbk -split. Denote by $\kappa \in H^2(G/M, \mathbb{k}^{\times})$ and $\lambda \in H^2(H/N, \mathbb{k}^{\times})$ their respective Schur classes. Suppose further that there exists $U \in {}_{\Bbbk Y}$ mod with $\operatorname{Res}_{M \times N}^Y(U) = V \otimes_{\Bbbk} W^\circ$. Then

$$\lambda = \eta_Y^*(\kappa) \in H^2(H/N, \mathbb{k}^{\times}).$$

Proof The Schur class of $V \otimes_{\Bbbk} W^{\circ}$ with respect to $M \times N \leq Y$ is $\operatorname{res}_{Y/(M \times N)}^{(G \times H)/(M \times N)}(\kappa \times \lambda^{-1})$, by 11.6(b), (c), and (d). Since $V \otimes_{\Bbbk} W^{\circ}$ extends to Y, the latter class is trivial. Let $\overline{p}_2 \colon Y/(M \times N) \xrightarrow{\sim} H/N$ be the isomorphism induced by the projection $p_2 \colon G \times H \to H$. Then also $(\overline{p}_2^{-1})^* (\operatorname{res}_{Y/(M \times N)}^{(G \times H)/(M \times N)}(\kappa \times \lambda^{-1})) = 1$ in $H^2(H/N, \Bbbk^{\times})$. However, a straightforward cocycle computation shows that $(\overline{p}_2^{-1})^* (\operatorname{res}_{Y/(M \times N)}^{(G \times H)/(M \times N)}(\kappa \times \lambda^{-1})) = \eta^*(\kappa) \cdot \lambda^{-1}$, and the proof is complete.

The element μ in the next proposition will be specified in the follow-up proposition to $e\mu(P,\phi,Q)f$ with the notation as in 10.1.

11.8 Proposition Let G and H be finite groups, $Y \leq G \times H$ with $p_1(Y) = G$ and $p_2(Y) = H$. Set $M := k_1(Y) \leq G$ and $N := k_2(Y) \leq H$, and let $\eta_Y : H/N \xrightarrow{\sim} G/M$ denote the isomorphism induced by Y. Suppose that e is a G-stable idempotent of $Z(\mathbb{K}M)$, f is an H-stable idempotent of $Z(\mathbb{K}N)$, and that $\mu \in R(\mathbb{K}Y(e \otimes f^*))$ is a virtual character such that $\operatorname{res}_{M \times N}^Y(\mu)$ is an isometry between $\mathbb{K}Me$ and $\mathbb{K}Nf$.

(a) If α : Irr($\mathbb{K}Nf$) $\xrightarrow{\sim}$ Irr($\mathbb{K}Me$) denotes the bijection induced by res $_{M \times N}^{Y}(\mu)$ then $\alpha({}^{hN}\chi) = {}^{\eta_{Y}(hN)}\alpha(\chi)$, for all $\chi \in \operatorname{Irr}(\mathbb{K}Nf)$ and all $h \in H$. In particular, α induces a bijection $\overline{\alpha}$: Irr($\mathbb{K}Nf$)/ $H \xrightarrow{\sim}$ Irr($\mathbb{K}Me$)/G.

(b) The group homomorphisms $\mu \stackrel{Y,H}{\underset{H}{\cdot}}$ - and $\operatorname{ind}_{Y}^{G \times H}(\mu) \underset{H}{\cdot}$ - from $R(\mathbb{K}Hf)$ to $R(\mathbb{K}Ge)$ coincide.

(c) Set $X_1 := (M \times M)\Delta(G)$ and $X_2 := (N \times N)\Delta(H)$. Then $Y * Y^\circ = X_1$ and $Y^\circ * Y = X_2$. Moreover, the following are equivalent:

(i)
$$\mu \stackrel{Y,Y^{\circ}}{\stackrel{\cdot}{H}} \mu^{\circ} = [\mathbb{K}Me] \text{ in } R(\mathbb{K}X_1(e \otimes e^*)).$$

(ii) $\mu^{\circ} \stackrel{Y^{\circ},Y}{\stackrel{\cdot}{G}} \mu = [\mathbb{K}Nf] \text{ in } R(\mathbb{K}X_2(f \otimes f^*)).$
(iii) $(\mu,\mu)_Y = |\mathrm{Irr}(\mathbb{K}Me)/G|.$

(iv) μ has precisely $|\operatorname{Irr}(\mathbb{K}Me)/G|$ distinct irreducible constituents and each of them occurs with multiplicity ± 1 .

(d) Assume that the equivalent conditions (i)–(iv) in Part (c) hold and let \mathcal{J} be a set of representatives of the *H*-orbits of Irr($\mathbb{K}Nf$). Then there exist irreducible characters $\mu_{\chi} \in \text{Irr}(\mathbb{K}Y(e \otimes f^*))$ and signs $\varepsilon_{\chi} \in \{\pm 1\}$, for $\chi \in \mathcal{J}$, such that $\mu = \sum_{\chi \in \mathcal{J}} \varepsilon_{\chi} \cdot \mu_{\chi}$ and $\operatorname{res}_{M \times N}^{Y}(\mu_{\chi}) = \sum_{\chi' \in [\chi]_{H}} \alpha(\chi') \times \chi'^{\circ}$ for all $\chi \in \mathcal{J}$.

(e) Assume that the equivalent conditions (i)–(iv) in Part (c) hold. Let $\chi \in \operatorname{Irr}(\mathbb{K}Nf)$ and set $\zeta := \alpha(\chi) \in \operatorname{Irr}(\mathbb{K}Me)$, $G_1 := \operatorname{stab}_G(\zeta)$, and $H_1 := \operatorname{stab}_H(\chi)$. Then, $\eta_Y(H_1) = G_1$ by Part (a). If $\kappa \in H^2(G_1/M, \mathbb{K}^{\times})$ and $\lambda \in H^2(H/N, \mathbb{K}^{\times})$ denote the respective extension classes of ζ and χ then $\kappa = \eta_Y^*(\lambda)$.

(f) Assume that the equivalent conditions (i)–(iv) in Part (c) hold. Then $\operatorname{ind}_Y^{G \times H}(\mu) \in R(\mathbb{K}Ge, \mathbb{K}Hf)$ is an isometry between $\mathbb{K}Ge$ and $\mathbb{K}Hf$.

Proof (a) By Remark 8.4(a) we can write

$$\operatorname{res}_{M \times N}^{Y}(\mu) = \sum_{\chi \in \operatorname{Irr}(\mathbb{K}Nf)} \varepsilon_{\chi} \cdot \alpha(\chi) \times \chi^{\circ}$$
(22)

with $\varepsilon_{\chi} \in \{\pm 1\}$, for $\chi \in \operatorname{Irr}(\mathbb{K}Nf)$. Let $(g,h) \in Y$. Then ${}^{(g,h)}\mu = \mu$ and therefore, for every $\chi \in \operatorname{Irr}(\mathbb{K}Nf)$, $\varepsilon_{\chi} \cdot {}^{g}\alpha(\chi) \times {}^{h}\chi^{\circ}$ is again equal to one of the summands in the above sum. This implies ${}^{g}\alpha(\chi) = \alpha({}^{h}\chi)$ and

 $\varepsilon_{\chi} = \varepsilon_{h_{\chi}}$, and Part (a) is proved. For later use, let $\mathcal{J} \subseteq \operatorname{Irr}(\mathbb{K}Nf)$ denote a set of representatives of the *H*-orbits of $\operatorname{Irr}(\mathbb{K}Nf)$. Then the above also shows that $\chi \mapsto \alpha(\chi) \times \chi^{\circ}$ induces a bijection

$$\mathcal{J} \xrightarrow{\sim} \{\alpha(\chi) \times \chi^{\circ} \mid \chi \in \operatorname{Irr}(\mathbb{K}Nf)\}/Y.$$
(23)

(b) Note that identifying H with $H \times \{1\}$ and G with $G \times \{1\}$, the first map, given by the generalized tensor product, maps $R(\mathbb{K}Hf)$ to $R(\mathbb{K}Ge)$, since $Y * (H \times \{1\}) = G \times \{1\}$. The statement follows directly from Lemma 6.5(a) applied to $Y \leq G \times H$ and $H \times \{1\} \leq H \times \{1\}$.

(c) We will show that (i) \Rightarrow (iii) \Rightarrow (iv) \Rightarrow (i). Then also (ii) \iff (iii) by symmetry, since $(\mu^{\circ}, \mu^{\circ})_{Y^{\circ}} = (\mu, \mu)_{Y}$ and $|\operatorname{Irr}(\mathbb{K}Ge)/G| = |\operatorname{Irr}(\mathbb{K}Nf)/H|$, by Part (a). But first we establish some facts that hold without further hypotheses. Note that $Y * Y^{\circ} = X_1$ and $Y^{\circ} * Y = X_2$ by Lemma 2.2(a), since $p_1(Y) = G$ and $p_2(Y) = H$. Set $\theta_1 := [\mathbb{K}Me] \in R(\mathbb{K}X_1(e \otimes e^*))$. Then Proposition 6.7(c) and Corollary 6.9(b) imply

$$(\mu,\mu)_Y = (\mu,\theta_1 \frac{X_1,Y}{G} \mu)_Y = (\mu \frac{Y,Y^{\circ}}{H} \mu^{\circ},\theta_1)_{X_1}.$$
(24)

Now write $\mu = a_1 \mu_1 + \dots + a_r \mu_r$ with pairwise distinct $\mu_1, \dots, \mu_r \in \operatorname{Irr}(\mathbb{K}Y)$ and non-zero integers a_1, \dots, a_r . Since $\operatorname{res}_{M \times N}^Y(\mu_i)$ is a multiple of the sum of a Y-orbit of $\operatorname{Irr}(\mathbb{K}[M \times N])$, (22) and (23) imply that

$$(\mu,\mu)_Y = a_1^2 + \dots + a_r^2 \ge r \ge |\mathcal{J}| = |\mathrm{Irr}(\mathbb{K}Nf)/H|.$$
(25)

Moreover, one has $(\mu, \mu)_Y = |\operatorname{Irr}(\mathbb{K}Nf)/H|$, if and only if there is a bijection $\{1, \ldots, r\} \xrightarrow{\sim} \mathcal{J}, i \mapsto \chi_i$, such that

$$a_i = \varepsilon_{\chi_i}$$
 and $\operatorname{res}_{M \times N}^Y(\mu_i) = \sum_{\chi' \in [\chi_i]_H} \alpha(\chi') \times {\chi'}^\circ$, (26)

for all i = 1, ..., r, where $[\chi]_H$ denotes the *H*-orbit of $\chi \in Irr(\mathbb{K}Nf)$. This shows that (*iii*) implies (*iv*), since $|Irr(\mathbb{K}Ge)/G| = |Irr(\mathbb{K}Hf)/H|$ by Part (a).

The same considerations apply to θ_1 with $\operatorname{res}_{M \times M}^{X_1}(\theta_1) = \sum_{\zeta \in \operatorname{Irr}(\mathbb{K}Me)} \zeta \times \zeta^\circ$. Since θ_1 is the character of a $\mathbb{K}X_1$ -module, each *G*-orbit sum $\sum_{\zeta'} \zeta' \times \zeta'^\circ$ extends to an irreducible character and we obtain

$$(\theta_1, \theta_1)_{X_1} = |\operatorname{Irr}(\mathbb{K}Me)/G|.$$
(27)

Now, (i) implies (iii) by Equations (24) and (27).

Finally, we assume (iv) and aim to show (i). Since (iv) implies $(\mu, \mu)_Y = |\mathcal{J}|$, we obtain (26) and can write $\mu = \sum_{\chi \in \mathcal{J}} \varepsilon_{\chi} \mu_{\chi}$, with $\mu_{\chi} \in \operatorname{Irr}(\mathbb{K}Y)$ satisfying $\operatorname{res}_{M \times N}^Y(\mu_{\chi}) = \sum_{\chi' \in [\chi]_H} \alpha(\chi') \times \chi'^\circ$, for all $\chi \in \mathcal{J}$. If $\chi_1, \chi_2 \in \mathcal{J}$ are distinct then $\mu_{\chi_1} \stackrel{Y,Y^\circ}{\stackrel{\cdot}{H}} \mu_{\chi_2}^\circ = 0$, by the definition of the extended tensor product (taken over $\mathbb{K}N$ in this case). Thus

$$\mu_{\stackrel{\cdot}{H}}^{Y,Y^{\circ}}\mu^{\circ} = \sum_{\chi \in \mathcal{J}} \varepsilon_{\chi}\mu_{\chi} \stackrel{Y,Y^{\circ}}{\stackrel{\cdot}{H}} \varepsilon_{\chi}\mu_{\chi}^{\circ} = \sum_{\chi \in \mathcal{J}} \mu_{\chi} \stackrel{Y,Y^{\circ}}{\stackrel{\cdot}{H}} \mu_{\chi}^{\circ}.$$

Moreover, for each $\chi \in \mathcal{J}$, the character $\mu_{\chi} \stackrel{Y,Y^{\circ}}{\stackrel{H}{H}} \mu_{\chi}^{\circ}$ of X_1 is irreducible, since its restriction to $M \times M$ is the sum $\sum_{\zeta' \in [\alpha(\chi)]_G} \zeta' \times \zeta'^{\circ}$ is the sum of an X_1 -orbit of elements in $\operatorname{Irr}(\mathbb{K}[M \times M])$. Furthermore, the irreducible characters $\mu_{\chi} \stackrel{Y,Y^{\circ}}{\stackrel{H}{H}} \mu_{\chi}^{\circ}$, $\chi \in \mathcal{J}$, are pairwise orthogonal, since their restrictions to $M \times M$ are. Thus, both $\mu \stackrel{Y,Y^{\circ}}{\stackrel{H}{H}} \mu^{\circ}$ and θ_1 are multiplicity-free sums of $|\mathcal{J}|$ pairwise distinct irreducible characters of X_1 . But by Equation (24) and by (iv) we have $(\mu \stackrel{Y,Y^{\circ}}{\stackrel{H}{H}} \mu^{\circ}, \theta_1)_{X_1} = (\mu, \mu)_Y = |\mathcal{J}|$. This implies $\mu \stackrel{Y,Y^{\circ}}{\stackrel{H}{H}} \mu^{\circ} = \theta_1$ and the proof of Part (c) is complete.

- (d) This was shown in the proof of Part (c).
- (e) This follows immediately from Part (c) and Lemma 11.7.
- (f) By Theorem 6.2 we have

$$\left(\operatorname{ind}_{Y}^{G\times H}(\mu)\right)_{H}\left(\operatorname{ind}_{Y}^{G\times H}(\mu)\right)^{\circ} = \left(\operatorname{ind}_{Y}^{G\times H}(\mu)\right)_{H}\left(\operatorname{ind}_{Y^{\circ}}^{H\times G}(\mu^{\circ})\right) = \operatorname{ind}_{Y*Y^{\circ}}^{G\times G}\left(\mu^{\overset{\circ}{\cdot}}_{\overset{\circ}{H}}\mu^{\circ}\right),\tag{28}$$

since $p_2(Y) = p_1(Y^\circ) = H$. Moreover, $\mu \stackrel{Y,Y^\circ}{\underset{H}{\cdot}} \mu^\circ = [\mathbb{K}Me]$ in $R(\mathbb{K}X_1(e \otimes e^*))$ by hypothesis. Thus, the last expression in (28) equals $\operatorname{ind}_{X_1}^{G \times G}([\mathbb{K}Me])$ which in turn equals $[\mathbb{K}Ge] \in R(\mathbb{K}Ge, \mathbb{K}Ge)$. Similarly one shows that

$$(\mathrm{ind}_Y^{G\times H}(\mu))^\circ \underset{G}{\cdot} (\mathrm{ind}_Y^{G\times H}(\mu)) = [\mathbb{K}Hf]$$

in $R(\mathbb{K}Hf,\mathbb{K}Hf)$. Now Remark 8.4(a) implies the result.

11.9 Proposition Let A be a block of $\mathcal{O}G$, B a block of $\mathcal{O}H$, $\gamma \in T_l^{\Delta}(A, B)$, and let $(\Delta(P, \phi, Q), e \otimes f^*) \in \mathcal{BP}_{\mathcal{O}}(\gamma)$. Set $I := N_G(P, e)$, $J := N_H(Q, f)$, and $Y := N_{I \times J}(\Delta(P, \phi, Q))$. Then the element $\mu := e\mu(P, \phi, Q)f \in R(\mathbb{K}Y(e \otimes f^*))$ defined in 10.1 has the following properties:

(a) The restriction of μ to $C_G(P) \times C_H(Q)$ is a perfect isometry between $\mathbb{K}[C_G(P)]e$ and $\mathbb{K}[C_H(Q)]f$.

(b) The conditions (i)–(iv) in Part (c), and therefore Parts (d), (e), and (f) of Proposition 11.8 hold for the groups I and J, the subgroup $Y \leq I \times J$, their normal subgroups $C_G(P)$ and $C_H(Q)$, and the character μ .

Proof (a) This follows from Theorem 11.4 and Proposition 9.9.

(b) By Part (a) and Proposition 11.1, all the hypotheses of Proposition 11.8 are satisfied. Moreover, the condition in Proposition 11.8(c)(i) holds by Equation (20) in Theorem 11.4. Thus, the proof is complete.

As an immediate consequence of the above proposition, we obtain the following finiteness result.

11.10 Theorem Let A be a block of $\mathcal{O}G$ and B a block of $\mathcal{O}H$. The set $T_l^{\Delta}(A, B)$ of left p-permutation equivalences between A and B is finite. In particular, the set $T_o^{\Delta}(A, B)$ of p-permutation equivalences between A and B is finite and the group $T_o^{\Delta}(A, A)$ of p-permutation self equivalences of A is finite.

Proof By Proposition 11.9, for each $A \otimes B^*$ -Brauer pair $(Y, e \otimes f^*)$, there are only finitely many choices for the virtual character of $\gamma(Y, e \otimes f^*) \in T(\mathcal{O}Y(e \otimes f^*))$, since it has bounded norm by the condition in Proposition 11.8(c)(iii). On the other hand, Proposition 9.2 implies that these characters determine γ .

12 A character theoretic criterion and moving from left to right

Throughout this section we assume that G and H are finite groups and that the *p*-modular system ($\mathbb{K}, \mathcal{O}, F$) is large enough for G and H. Further, we assume that A is a block algebra of $\mathcal{O}G$ and that B is a block algebra of $\mathcal{O}H$.

In this section we will prove a character theoretic criterion for an element $\gamma \in T^{\Delta}(A, B)$ to be in $T_l^{\Delta}(A, B)$, see Theorem 12.2. Since one of the conditions in this criterion is symmetric, we can derive that $T_l^{\Delta}(A, B) = T_o^{\Delta}(A, B) = T_r^{\Delta}(A, B)$, see Theorem 12.3.

12.1 Lemma Suppose that (D, e_D) is a maximal A-Brauer pair, (E, f_E) is a maximal B-Brauer pair, and $\phi: E \xrightarrow{\sim} D$ is an isomorphism between the fusion systems \mathcal{B} and \mathcal{A} associated with (E, f_E) and (D, e_D) , respectively. Suppose further that one has A-Brauer pairs (P, e) and (R, d) and a B-Brauer pair (Q, f) with $(P, e) \leq (D, e_D), (R, d) \leq (D, e_D)$ and $(Q, f) \leq (E, f_E)$, and isomorphisms $\alpha: Q \xrightarrow{\sim} P$ and $\beta: Q \xrightarrow{\sim} R$ such that

$$(\Delta(P,\alpha,Q), e \otimes f^*) \leq_{G \times H} (\Delta(D,\phi,E), e_D \otimes f_E) \quad \text{and} \\ (\Delta(R,\beta,Q), d \otimes f^*) \leq_{G \times H} (\Delta(D,\phi,E), e_D \otimes f_E) \,.$$

Then $(\Delta(P, \alpha\beta^{-1}, R), e \otimes d^*) =_{\{1\} \times G} (\Delta(P), e \otimes e^*)$ and there exists $g \in G$ such that $c_g \circ \alpha = \beta \colon Q \xrightarrow{\sim} P$ and ${}^{g}(P, e) = (R, d)$.

Proof By assumption there exist $(g_1, h_1), (g_2, h_2) \in G \times H$ such that

$${}^{(g_1,h_1)}(\Delta(P,\alpha,Q), e \otimes f^*) \leq (\Delta(D,\phi,E), e_D \otimes f^*_E) \quad \text{and}$$
$${}^{(g_2,h_2)}(\Delta(R,\beta,Q), d \otimes f^*) \leq (\Delta(D,\phi,E), e_D \otimes f^*_E),$$

or equivalently,

$${}^{g_1}(P,e) \leqslant (D,e_D) \,, \quad {}^{h_1}(Q,f) \leqslant (E,f_E) \,, \quad {}^{g_2}(R,d) \leqslant (D,e_D) \,, \quad {}^{h_2}(Q,f) \leqslant (E,f_E) \,, \\ {}^{g_1}P = \phi(\,{}^{h_1}\!Q) \,, \quad c_{g_1}\alpha c_{h_1^{-1}} = \phi \colon \, {}^{h_1}\!Q \xrightarrow{\sim} {}^{g_1}\!P \,, \quad {}^{g_2}\!R = \phi(\,{}^{h_2}\!Q) \,, \quad c_{g_2}\beta c_{h_2^{-1}} = \phi \colon \, {}^{h_2}\!Q \xrightarrow{\sim} {}^{g_2}\!R \,.$$

Thus, $c_{h_2h_1^{-1}} \in \operatorname{Hom}_{\mathcal{B}}({}^{h_1}Q, {}^{h_2}Q)$. Since $\phi \colon E \xrightarrow{\sim} D$ is an isomorphism between \mathcal{B} and \mathcal{A} , there exists $g \in G$ such that

$$^{gg_1}(P,e) = {}^{g_2}(R,d) \text{ and } \phi c_{h_2h_1^{-1}} = c_g\phi \colon {}^{h_1}Q \xrightarrow{\sim} \phi({}^{h_2}Q) = {}^{g_2}R \,.$$

From this we obtain $g_2^{-1}gg_1(P,e) = (R,d), c_{g_2^{-1}gg_1}\alpha = \beta \colon Q \xrightarrow{\sim} R$, which implies

$${}^{(1,g_1^{-1}g^{-1}g_2)}(\Delta(P,\alpha\beta^{-1},R), e\otimes d^*) = (\Delta(P), e\otimes e^*)$$

and the proof of the lemma is complete.

12.2 Theorem Let $\gamma \in T^{\Delta}(A, B)$. Then γ belongs to $T_l^{\Delta}(A, B)$ if and only if the following hold:

(i) There exists a maximal element $(\Delta(D, \phi, E), e_D \otimes f_E^*)$ in $\mathcal{BP}_{\mathcal{O}}(\gamma)$ such that (D, e_D) is a maximal A-Brauer pair, (E, f_E) is a maximal B-Brauer pair and $\phi: E \to D$ is an isomorphism between the fusion systems \mathcal{B} and \mathcal{A} of B and A associated with (E, f_E) and (D, e_D) , respectively;

(ii) Any two maximal elements in $\mathcal{BP}_{\mathcal{O}}(\gamma)$ are $G \times H$ -conjugate; and

(iii) For every $(\Delta(P,\psi,Q), e \otimes f^*) \in \mathcal{BP}_{\mathcal{O}}(A \otimes B^*)$ with $(\Delta(P,\psi,Q), e \otimes f^*) \leq (\Delta(D,\phi,E), e_D \otimes f^*_E)$, setting $I := N_G(P,e), J := N_H(Q,f)$, and $Y := N_{I \times J}(\Delta(P,\psi,Q))$, the element $\mu := \gamma(\Delta(P,\psi,Q), e \otimes f^*)^{\mathbb{K}}$ in $R(\mathbb{K}Y(e \otimes f^*))$ satisfies

$$\mu \stackrel{Y,Y^{\circ}}{\underset{H}{\cdot}} \mu^{\circ} = [\mathbb{K}[C_G(P)]e] \quad in \ R(\mathbb{K}[N_{I \times I}(\Delta(P))]).$$

Proof First assume that $\gamma \in T_l^{\Delta}(A, B)$. Let $(\Delta(D, \phi, E), e_D \otimes f_E)$ be any maximal γ -Brauer pair. Then the conditions in (i) are satisfied by Theorem 10.11(c) and by Theorem 11.2. Moreover, Theorem 10.11(b), implies the condition in (ii), and Equation (20) in Theorem 11.4 implies the condition in (iii).

Next we assume that Conditions (i)–(iii) hold for $\gamma \in T^{\Delta}(A, B)$. We aim to show that $\gamma_{H} \cdot \gamma^{\circ} = [A]$ in $T^{\Delta}(A, A)$. By Proposition 9.2(b), it suffices to show that

$$\left((\gamma_{H} \cdot \gamma^{\circ})(X,c)\right)^{\mathbb{K}} = \left(A(X,c)\right)^{\mathbb{K}} \quad \text{in } T(\mathbb{K}[N_{G \times G}(X,c)]),$$
(29)

for all (X, c) running through a set of representatives of the $G \times G$ -orbits of $\mathcal{BP}(A \otimes A^*)$.

Claim 1: Both sides of (29) are 0 unless $(X,c) =_{G\times G} (\Delta(P), e \otimes e^*)$ for some Brauer pair (P,e) with $(P,e) \leq (D,e_D)$. This is clear for the right hand side, since $(\Delta(D),e_D \otimes e_D^*)$ is a maximal Brauer pair of the indecomposable *p*-permutation $A \otimes A^*$ -module A (see Proposition 5.3(c)). To prove this for the left hand side of (29), assume that $(X,c) \in \mathcal{BP}(A \otimes A^*)$ is such that the left hand side of (29) is non-zero. Then also $(\gamma \cdot_H \gamma^\circ)(X,c) \neq 0$ in $T(F[N_{G\times G}(X,c)])$. Since every indecomposable *p*-permutation $\mathcal{O}[G \times G]$ -module M appearing in $\gamma \cdot_H \gamma^\circ$ has twisted diagonal vertices (see Lemma 7.2(b)) and since every M-Brauer pair is also an $A \otimes A^*$ -Brauer pair (see Proposition 5.3(a)), we can conclude that (X,c) is $G \times G$ -conjugate to a Brauer pair of the form $(\Delta(P,\sigma,R), e \otimes d^*)$ with $(P,e) \leq (D,e_D)$ and $(R,d) \leq (D,e_D)$. Using Theorem 7.5(c) with $I := N_G(P,e)$ for S and $K := N_G(R,d)$ for T, this implies that there exists a Brauer pair (Q,f) of $\mathcal{O}H$ and isomorphisms $\alpha: Q \xrightarrow{\sim} P$ and $\beta: R \to Q$ such that $\alpha\beta = \sigma, \gamma(\Delta(P,\alpha,Q), e \otimes f^*) \neq 0$ in $T(F[N_{I\times J}(\Delta(P,\alpha,Q))])$, and $\gamma(\Delta(R,\beta^{-1},Q), d \otimes f^*) \neq 0$ in $T(F[N_{K\times J}(\Delta(R,\alpha,Q)]))$, where $J := N_H(Q,f)$. By Condition (ii) this implies $(\Delta(P,\alpha,Q), e \otimes f^*) \leq_{G\times H} (\Delta(D,\phi,E), e_D \otimes f_E^*)$ and $(\Delta(R,\beta^{-1},Q), d \otimes f^*) \leq_{G\times H} (\Delta(D,\phi,E), e_D \otimes f_E^*)$ and $(\Delta(R,\beta^{-1},Q), d \otimes f^*) \leq_{G\times H} (\Delta(D,\phi,E), e_D \otimes f_E^*)$. Now Lemma 12.1 implies Claim 1.

With Claim 1 it now suffices to show that

$$\left((\gamma_{H} \cdot \gamma^{\circ})(\Delta(P), e \otimes e^{*}) \right)^{\mathbb{K}} = \left([A](\Delta(P), e \otimes e^{*}) \right)^{\mathbb{K}} \quad \text{in } R(\mathbb{K}[N_{I \times I}(\Delta(P))]),$$
(30)

for all A-Brauer pairs (P, e) with $(P, e) \leq (D, e_D)$, where $I := N_G(P, e)$. By Proposition 4.3(b), the right hand side of (30) equals $[\mathbb{K}[C_G(P)]e]$. We use the second part of Theorem 7.5(d) with S = I to compute

1.0		

the left hand side of (30). For $Q \leq E$ let f_Q denote the unique block idempotent of $\mathcal{O}[C_H(Q)]$ such that $(Q, f_Q) \leq (E, f_E)$ and set $Q_0 := \phi^{-1}(P)$ and $f_0 := f_{Q_0}$. Then, with the notation in Theorem 7.5(d), the summand for $\lambda_0 = (\phi, (Q_0, f_0))$ contributes the element

$$\operatorname{ind}_{\Delta(I(\lambda_0))(C_G(P)\times\{1\})}^{N_{I\times I}(\Delta(P))} \left(\gamma(\Delta(P,\phi,Q_0),e\otimes f_0^*) \stackrel{Y_0,Y_0^{\circ}}{\stackrel{\cdot}{H}} \gamma(\Delta(P,\phi,Q_0),e\otimes f_0^*)^{\circ}\right),$$

where $Y_0 = N_{G \times H}(\Delta(P\phi, Q_0), e \otimes f_0^*)$. Since ϕ is an isomorphism between the fusion systems \mathcal{B} and \mathcal{A} by Condition (i), we have $I(\lambda_0) = I$ and therefore $\Delta(I(\lambda_0))(C_G(P) \times \{1\}) = N_{I \times I}(\Delta(P))$. Thus, after applying $(-)^{\mathbb{K}}$, the contribution of the summand parametrized by λ_0 to the left hand side of (30) is equal to $\mu_0 \frac{Y_0, Y_0^\circ}{H} \mu_0^\circ$ with $\mu_0 := \gamma(\Delta(P, \phi, Q_0), e \otimes f_0^*)^{\mathbb{K}}$. Condition (iii) now implies that this expression is also equal to $[\mathbb{K}[C_G(P)]e] \in R(\mathbb{K}[N_{I \times I}(\Delta(P))])$. Thus, it suffices to show *Claim 2: Let* (Q, f) *be an OH-Brauer pair and* $\alpha: Q \xrightarrow{\rightarrow} P$ *an isomorphism with* $\gamma(\Delta(P, \alpha, Q), e \otimes f^*) \neq 0$ *in* $T(F[N_{G \times H}(\Delta(P, \alpha, Q), e \otimes f^*)])$ *then* $\lambda = (\alpha, (Q, f))$ *belongs to the same* $I \times H$ -orbit as λ_0 . In order to prove this claim, note first that $\gamma(\Delta(P, \alpha, Q), e \otimes f^*) \neq 0$ implies that (Q, f) is a *B*-Brauer pair. Without loss of generality we may assume that $(Q, f) \leq (E, f_E)$. Condition (ii) now implies that $(\Delta(P, \alpha, Q), e \otimes f^*) \neq 0$ implies that $(\Delta(P, \alpha, Q), e \otimes f^*) \leq G_{XH} (\Delta(D, \phi, E), e_D \otimes f_E^*)$. Applying the symmetric statement of Lemma 12.1 to $(\Delta(P, \alpha, Q), e \otimes f^*)$ and $(\Delta(P, \phi, Q_0), e \otimes f_0^*)$, we obtain $(\alpha, (Q, f)) =_H (\phi, (Q_0, f_0))$. This completes the proof of the theorem.

12.3 Theorem Let A be a block of $\mathcal{O}G$, B a block of $\mathcal{O}H$ and $\gamma \in T^{\Delta}(A, B)$. The following are equivalent:

- (i) $\gamma \in T_l^{\Delta}(A, B)$.
- (ii) $\gamma \in T_r^{\Delta}(A, B)$.
- (iii) $\gamma \in T_{\alpha}^{\Delta}(A, B).$
- (iv) γ satisfies the conditions (i)–(iii) in Theorem 12.2.

Proof By Theorem 12.2, (i) is equivalent with (iv). Morover, clearly (iii) implies (i) and (ii). Next we show that (i) implies (iii). In fact, assuming (i), Theorem 12.2 shows that the conditions (i)–(iii) in Theorem 12.2 are satisfied. Since conditions (i) and (ii) are symmetric and since Condition (iii) for γ is equivalent to Condition (iii) for γ° , by Proposition 11.9 and the equivalence of (i) and (ii) in Proposition 11.8(c), we see that the conditions (i)–(iii) also hold for $\gamma^{\circ} \in T^{\Delta}(B, A)$. Now Theorem 12.2 implies that $\gamma^{\circ} \in T^{\Delta}_{l}(B, A)$. But this is equivalent to $\gamma \in T^{\Delta}_{r}(A, B)$. Thus, (i) implies (iii). Finally, if (ii) holds then $\gamma^{\circ} \in T^{\Delta}_{l}(B, A)$, and, since (i) implies (iii), we obtain $\gamma^{\circ} \in T^{\Delta}_{o}(B, A)$ and $\gamma \in T^{\Delta}_{o}(A, B)$. Thus, (ii) implies (iii) implies (iii), and the proof is complete.

13 Külshammer-Puig classes

Throughout this section we assume that G and H are finite groups and that the *p*-modular system $(\mathbb{K}, \mathcal{O}, F)$ is large enough for G and H. Further, we assume that A is a block algebra of $\mathcal{O}G$, that B is a block algebra of $\mathcal{O}H$, and that $\gamma \in T^{\Delta}(A, B)$ is a *p*-permutation equivalence.

The main goal of this section is to show that Külshammer-Puig classes are 'preserved' by γ , see Theorem 13.4 for the precise statement. In order to prove this we need to take a closer look at the elements $e\gamma(P,\psi,Q)f$, for a γ -Brauer pair ($\Delta(P,\psi,Q), e \otimes f^*$) with P and Q centric in the associated fusion systems. This is done in Proposition 13.2.

13.1 Lemma Let $(\Delta(D, \phi, E), e \otimes f^*)$ be a maximal γ -Brauer pair and let \mathcal{F} denote the fusion system associated to the maximal $A \otimes B^*$ -Brauer pair $(D \times E, e \otimes f^*)$. Then the subgroup $\Delta(D, \phi, E)$ of $D \times E$ is fully \mathcal{F} -normalized. Moreover, setting $Y := N_{G \times H}(\Delta(D, \phi, E), e \otimes f^*)$, the block $\mathcal{O}Y(e \otimes f^*)$ has the normal subgroup $(Z(D) \times Z(E)) \cdot \Delta(D, \phi, E)$ of Y as defect group.

Proof Any $G \times H$ -conjugate subgroup of $\Delta(D, \phi, E)$ which is contained in $D \times E$ must be again a twisted diagonal subgroup and therefore of the form $\Delta(D, \psi, E)$ for some isomorphism $\psi \colon E \xrightarrow{\sim} D$. By Lemma 2.4(b), one has $N_{D \times E}(\Delta(D, \psi, E)) = (Z(D) \times Z(E)) \cdot \Delta(D, \psi, E)$, whose order is independent of ψ . This proves the first statement. The second statement follows immediately from Proposition 4.5(b).

13.2 Proposition Let $(\Delta(P, \psi, Q), e \otimes f^*) \in \mathcal{BP}_{\mathcal{O}}(\gamma)$ and suppose that Z(P) is a defect group of $\mathcal{O}[C_G(P)]e$. Set

$$I := N_G(P, e), \quad J := N_H(Q, f), \quad Y := N_{I \times J}(\Delta(P, \psi, Q)), \quad C := C_G(P) \times C_H(Q),$$
$$Z := Z(P) \times Z(Q), \quad \text{and} \quad Z' := \Delta(Z(P), \psi, Z(Q)).$$

(a) One has

$$\operatorname{res}_{C}^{Y}(e\gamma(P,\psi,Q)f) = \varepsilon[N'] \quad \text{in } T(\mathcal{O}C(e\otimes f^{*})),$$

where $\varepsilon \in \{\pm 1\}$ and N' is the unique indecomposable p-permutation $\mathcal{O}C(e \otimes f^*)$ -module with vertex Z' (cf. Proposition 4.8(b)). Moreover, the p-permutation $\mathcal{O}C(e \otimes f^*)$ -module V' := $\operatorname{Inf}_{C/Z}^C \operatorname{Def}_{C/Z}^C(N')$ satisfies:

(i) $F \otimes_{\mathcal{O}} V'$ is the unique simple $FC(e \otimes f^*)$ -module;

(ii) $\mathbb{K} \otimes_{\mathcal{O}} V'$ is a simple $\mathbb{K}C(e \otimes f^*)$ -module.

(b) Set $\zeta := \inf_{Y/Z}^{Y} \operatorname{def}_{Y/Z}^{Y}(e\gamma(P,\psi,Q)f) \in T(\mathcal{O}Y(e \otimes f^*))$. There exists a simple $\mathbb{K}Y(e \otimes f^*)$ -module Wand a simple $FY(e \otimes f^*)$ -module \overline{W} such that $d_Y([W]) = [\overline{W}]$ in $R(FY(e \otimes f^*))$ and such that the following hold with ε from Part (a):

(i)
$$\kappa_Y(\zeta) = \varepsilon[W]$$
 in $R(\mathbb{K}Y(e \otimes f^*))$ and $\operatorname{Res}_C^Y(W) \cong \mathbb{K} \otimes_{\mathcal{O}} V';$

(ii) $\eta_Y(\overline{\zeta}) = \varepsilon[\overline{W}]$ in $R(FY(e \otimes f^*))$ and $\operatorname{Res}_C^Y(\overline{W}) \cong F \otimes_{\mathcal{O}} V'$.

(c) If $(\Delta(P, \psi, Q), e \otimes f^*)$ is a maximal γ -Brauer pair then there exists an indecomposable p-permutation $\mathcal{O}Y(e \otimes f^*)$ -module N such that

$$e\gamma(P,\psi,Q)f = \varepsilon[N] \text{ in } T(\mathcal{O}Y(e\otimes f^*)) \quad \text{and} \quad \operatorname{Res}^Y_C(N) \cong N'\,,$$

where ε is from Part (a).

We visualize the situation via the diagrams of elements

$$\begin{aligned} \zeta & \mapsto \varepsilon[W] & \varepsilon[V'] & \mapsto \varepsilon[\mathbb{K} \otimes_{\mathcal{O}} V'] \\ \downarrow & \downarrow & \text{and} & \downarrow & \downarrow \\ \overline{\zeta} & \mapsto \varepsilon[\overline{W}] & \varepsilon[F \otimes_{\mathcal{O}} V'] & \mapsto \varepsilon[F \otimes_{\mathcal{O}} V'] \end{aligned} \tag{31}$$

in the diagrams of groups

$$\begin{array}{cccc} T(\mathcal{O}Y(e\otimes f^*)) & \xrightarrow{\kappa_Y} & R(\mathbb{K}Y(e\otimes f^*)) & & T(\mathcal{O}C(e\otimes f^*)) & \xrightarrow{\kappa_C} & R(\mathbb{K}C(e\otimes f^*)) \\ & \downarrow & & \downarrow \\ T(FY(e\otimes f^*)) & \xrightarrow{\eta_Y} & R(FY(e\otimes f^*)) & & T(FC(e\otimes f^*)) & \xrightarrow{\eta_C} & R(FC(e\otimes f^*)) , \end{array}$$

where the right diagram of elements is the image of the left one under the restriction maps res_C^Y , as shown in the proof. Before we prove the proposition, we mention the following negative result:

13.3 Remark Assume the situation in Proposition 13.2. In general, the element $e\gamma(P, \psi, Q)f \in T(\mathcal{O}Y(e \otimes f^*))$ is not plus or minus the class of an indecomposable *p*-permutation $\mathcal{O}Y$ -module. In fact, let G = H be the dihedral group of order 8, P the cyclic subgroup of order 4 and $\gamma := [\mathcal{O}[G \times G/\Delta(G)]] - [\mathcal{O}[G \times G/\Delta(P)]] \in T^{\Delta}(\mathcal{O}G, \mathcal{O}G)$. Then γ is a *p*-permutation equivalence, $P = Z(P) = C_G(P)$ is a defect group of $\mathcal{O}[C_G(P)]$, but $\gamma(\Delta(P)) = [\mathcal{O}[Y/\Delta(G)]] - [\mathcal{O}[Y/\Delta(P)]]$ in $T(\mathcal{O}Y)$.

Proof of Proposition 13.2. We set $\omega := e\gamma(P, \psi, Q)f \in T(\mathcal{O}Y(e \otimes f^*))$ and $\omega' := \operatorname{res}_C^Y(\omega) \in T(\mathcal{O}C(e \otimes f^*)).$

(a) Since Z(P) is a defect group of $\mathcal{O}[C_G(P)]e$, Z(Q) is a defect group of $\mathcal{O}[C_H(Q)]f$. In fact, we can choose a maximal γ -Brauer pair $(\Delta(D, \phi, E), e' \otimes f'^*)$ such that $(\Delta(P, \psi, Q), e \otimes f^*) \leq (\Delta(D, \phi, E), e' \otimes f'^*)$. Then, ψ is the restriction of ϕ and $\phi \colon E \xrightarrow{\sim} D$ is an isomorphism between the fusion systems \mathcal{A} and \mathcal{B} associated with (D, e') and (E, f'), cf. Theorem 11.2, and since P is centric in the fusion system \mathcal{A} , Q must be centric in the fusion system \mathcal{B} . Applying now Proposition 4.5 the claim follows. Thus, Z is a defect group of $\mathcal{O}C(e \otimes f^*)$. Let U be an indecomposable $\mathcal{O}C$ -module appearing in ω' . We will show that Z' is a vertex of U. In fact, let $X \leq C$ denote a vertex of U. Since $\Delta(P, \psi, Q)$ acts trivially on every indecomposable $\mathcal{O}Y$ -module appearing in ω , the group Z' acts trivially on U. Thus, $Z' \leq X$. Moreover, X must be a twisted diagonal subgroup of Z, since Z is the unique defect group of $\mathcal{O}C(e \otimes f^*)$. This implies X = Z'.

Since $\mathcal{O}C(e \otimes f^*)$ has the central defect group Z, Proposition 4.8(b) implies that there exists a unique *p*-permutation $\mathcal{O}C(e \otimes f^*)$ -module N' with vertex Z'. Thus, by the previous paragraph, there exists $\varepsilon \in \mathbb{Z}$ with $\omega' = \varepsilon[N'] \in T(\mathcal{O}C)$. Since Lemma 10.3 implies that $\kappa_C(\omega') = \varepsilon[\mathbb{K} \otimes_{\mathcal{O}} N']$ is an isometry between $\mathbb{K}[C_G(P)]e$ and $\mathbb{K}[C_H(E)]f$, we obtain $\varepsilon \in \{\pm 1\}$. Moreover, by Proposition 4.8(b), the *p*-permutation $\mathcal{O}C(e \otimes f^*)$ -module $V' := \operatorname{Inf}_{C/Z}^C \operatorname{Def}_{C/Z}^C(N')$ has the property that $F \otimes_{\mathcal{O}} V'$ is the unique simple $FC(e \otimes f^*)$ -module. Note also that $F \otimes_{\mathcal{O}} V'$ can be viewed as the unique simple $F[C/Z]\pi(e \otimes f^*)$ -module, where $\pi : \mathcal{O}C \to \mathcal{O}[C/Z]$ denotes the canonical map $\mathcal{O}C \to \mathcal{O}[C/Z]$. Since $\mathcal{O}[C/Z]\pi(e \otimes f^*)$ is a block of defect 0 (see Proposition 4.8(a)), the $\mathbb{K}C(e \otimes f^*)$ -module $\mathbb{K} \otimes_{\mathcal{O}} V'$ is simple, namely the inflation of the unique simple $\mathbb{K}[C/Z]\pi(e \otimes f^*)$ -module. This establishes all the statements in Part (a).

(b) Since $Z' \leq \Delta(P, \psi, Q)$, Z' acts trivially on any indecomposable $\mathcal{O}Y$ -module appearing in ω . Using $Z = Z' \cdot (Z(P) \times \{1\})$ and Proposition 6.7(d), we obtain

$$\zeta = \inf_{Y/Z}^{Y} \operatorname{def}_{Y/Z}^{Y}(\omega) = \inf_{Y/(Z(P) \times \{1\})}^{Y} \operatorname{def}_{Y/(Z(P) \times \{1\})}^{Y}(\omega) = [\mathcal{O}[C_G(P)/Z(P)]\overline{e}] \xrightarrow{Y*Y^\circ, Y}_{G} \omega$$
(32)

in $T(\mathcal{O}Y(e \otimes f^*))$, where \overline{e} denotes the image of e under the canonical map $\mathcal{O}[C_G(P)] \to \mathcal{O}[C_G(P)/Z(P)]$. Note that $Y * Y^\circ = \Delta(I) \cdot (C_G(P) \times C_G(P))$ by Lemma 2.2(a) and Proposition 11.1. We claim that the character of the $\mathcal{O}[\Delta(I) \cdot (C_G(P) \times C_G(P))]$ -module $\mathcal{O}[C_G(P)/Z(P)]\overline{e}$ in (32) is irreducible. In fact, since $\mathcal{O}[C_G(P)]e$ has central defect group Z(P), $\mathcal{O}[C_G(P)/Z(P)]\overline{e}$ is a block of defect 0. Thus, $\mathbb{K}[C_G(P)/Z(P)]\overline{e}$ is an irreducible $(\mathbb{K}[C_G(P)/Z(P)]\overline{e}, \mathbb{K}[C_G(P)/Z(P)]\overline{e})$ -bimodule and via inflation an irreducible $\mathbb{K}[C_G(P) \times C_G(P)](e \otimes e^*)$ -module, which is a fortiori irreducible as $\mathbb{K}[\Delta(I) \cdot (C_G(P) \times C_G(P))]$ -module. Now, (32) and Lemma 11.3 imply that $\kappa_Y(\zeta) = \varepsilon' \cdot [W]$, for some $\varepsilon' \in \{\pm 1\}$ and some simple $\mathbb{K}Y(e \otimes f^*)$)-module W. Since clearly

$$\operatorname{res}_{C}^{Y} \operatorname{inf}_{Y/Z}^{Y} \operatorname{def}_{Y/Z}^{Y} = \operatorname{inf}_{C/Z}^{C} \operatorname{def}_{C/Z}^{C} \operatorname{res}_{C}^{Y} \colon T(\mathcal{O}Y) \to T(\mathcal{O}C) \,,$$

we have $\operatorname{res}_{C}^{Y}(\zeta) = \inf_{C/Z}^{C} \operatorname{def}_{C/Z}^{C} \operatorname{res}_{C}^{Y}(\omega) = \inf_{C/Z}^{C} \operatorname{def}_{C/Z}^{C}(\omega') = \varepsilon[V']$ and therefore $\operatorname{res}_{C}^{Y}(\varepsilon'[W]) = \operatorname{res}_{C}^{Y}(\kappa_{Y}(\zeta)) = \kappa_{C}(\operatorname{res}_{C}^{Y}(\zeta)) = \varepsilon \kappa_{C}[V'] = \varepsilon[\mathbb{K} \otimes_{\mathcal{O}} V']$. This implies that $\varepsilon' = \varepsilon$ and establishes Statement (i) of Part (b). Moreover, we have $\operatorname{res}_{C}^{Y}(d_{Y}([W])) = d_{C}(\operatorname{res}_{C}^{Y}([W])) = d_{C}([\mathbb{K} \otimes_{\mathcal{O}} V']) = [F \otimes_{\mathcal{O}} V'] \in R(FC)$. Since $F \otimes_{\mathcal{O}} V'$ is an irreducible $FC(e \otimes f^{*})$ -module, the previous equation implies that $d_{Y}([W]) = [\overline{W}]$, for some irreducible $FY(e \otimes f^{*})$ -module \overline{W} . This completes the proof of Part (b).

(c) By Lemma 13.1, the block algebra $\mathcal{O}Y(e \otimes f^*)$ has defect group $Z \cdot \Delta(P, \psi, Q)$, a normal subgroup of Y. Thus, by [NT89, Theorems V.8.7(ii) and V.8.10], $\mathcal{O}[Y/Z \cdot \Delta(P, \psi, Q)]\pi(e \otimes f^*)$ is a sum of blocks of defect 0, where $\pi : \mathcal{O}Y \to \mathcal{O}[Y/Z \cdot \Delta(P, \psi, Q)]$ denotes the canonical homomorphism. Since $\Delta(P, \psi, Q)$ acts trivially on every indecomposable $\mathcal{O}Y$ -module appearing in ω , we have

$$\overline{\zeta} = \inf_{Y/Z}^{Y} \operatorname{def}_{Y/Z}^{Y}(\overline{\omega}) = \inf_{Y/Z \cdot \Delta(P,\psi,Q)}^{Y} \operatorname{def}_{Y/Z \cdot \Delta(P,\psi,Q)}^{Y}(\overline{\omega}),$$

where $\operatorname{def}_{Y/Z \cdot \Delta(P,\psi,Q)}^{Y}(\overline{\omega}) \in T(F[Y/Z \cdot \Delta(P,\psi,Q)])$ is a \mathbb{Z} -linear combination of classes of simple *p*-permutation modules. This property is preserved under inflation, so that also $\overline{\zeta}$ is a \mathbb{Z} -linear combination of classes of simple *p*-permutation *FY*-modules. But since $\eta_Y(\overline{\zeta}) = \varepsilon[\overline{W}]$, we obtain that \overline{W} is a simple *p*-permutation *FY*($e \otimes f^*$)module and that $\overline{\zeta} = \varepsilon[\overline{W}]$ in $T(FY(e \otimes f^*))$.

Since every indecomposable FY-module appearing in $\overline{\omega} \in T(FY)$ has vertex $\Delta(P, \psi, Q)$ and $\inf_{Y/Z}^{Y} \det_{Y/Z}^{Y}(\overline{\omega}) = \overline{\zeta} = \varepsilon[\overline{W}] \in T(FY)$, Proposition 4.7(b) implies that $\overline{\omega} = \varepsilon[\overline{N}] \in T(FY(e \otimes f^*))$ for an indecomposable *p*-permutation $FY(e \otimes f^*)$ -module \overline{N} with vertex $\Delta(P, \psi, Q)$. Let N be the indecomposable *p*-permutation $\mathcal{O}Y(e \otimes f^*)$ -module corresponding to \overline{N} . Then $\omega = \varepsilon[N]$ and $N' \cong \operatorname{Res}_{C}^{Y}(N)$, since $\omega' = \operatorname{res}_{C}^{Y}(\omega)$. This completes the proof of the proposition.

Next we show that *p*-permutation equivalences 'preserve' Külshammer-Puig classes.

13.4 Theorem Let A be a block of $\mathcal{O}G$, B a block of $\mathcal{O}H$, and $\gamma \in T_o^{\Delta}(A, B)$. Further, let $(\Delta(P, \psi, Q), e \otimes f^*) \in \mathcal{BP}_{\mathcal{O}}(\gamma)$ and suppose that Z(P) is a defect group of the block $\mathcal{O}[C_G(P)]e$. Then Z(Q) is a defect group of the block $\mathcal{O}[C_H(Q)]f$, by Theorem 11.2 and the preservation of centric subgroups. Set $I := N_G(P, e)$,

 $\overline{I} := I/PC_G(P), \ J := N_H(Q, f), \ \text{and} \ \overline{J} := J/QC_H(Q).$ Furthermore, let $\kappa \in H^2(\overline{I}, F^{\times})$ and $\lambda \in H^2(\overline{J}, F^{\times})$ denote the Külshammer-Puig classes of (P, \overline{e}) and (Q, \overline{f}) , respectively. Then

$$\lambda = \overline{\eta}^*(\kappa) \in H^2(\overline{J}, F^{\times}),$$

where $\overline{\eta} \colon \overline{J} \xrightarrow{\sim} \overline{I}$ is induced by the isomorphisms in Proposition 11.1 and $\overline{\eta}^* := H^2(\overline{\eta}, F^{\times})$.

Proof Let M denote the unique simple $F[PC_G(P)]e$ -module and let N denote the unique simple $F[QC_H(Q)]f^*$ module. Then, by 4.6, κ is the Schur class of M with respect to $PC_G(P) \leq I$, and λ^{-1} is the Schur class of Nwith respect to $QC_H(Q) \leq J$. Set $C := C_G(P) \times C_H(Q)$, $Y := N_{I \times J}(\Delta(P, \psi, Q))$ and $\tilde{Y} := Y(P \times Q) \leq I \times J$. By Proposition 11.1 and Lemma 11.7, it suffices to show that the irreducible $F[C(P \times Q)]$ -module $M \otimes_F N$ can be extended to \tilde{Y} .

By Proposition 13.2(b), there exists an irreducible $FY(e \otimes f^*)$ -module W (it is denoted by \overline{W} there) such that $\operatorname{Res}_C^Y(W) \cong M \otimes N$. Consider the canonical isomorphism $\tilde{Y}/(P \times Q) \cong Y/Y \cap (P \times Q)$ and note that $Y \cap (P \times Q) = (Z(P) \times Z(Q))\Delta(P, \psi, Q)$ is a normal p-subgroup of Y and therefore acts trivially on W. Using the above isomorphism we see that W extends to an $F\tilde{Y}$ -module \tilde{W} . Then, $\operatorname{Res}_{C(P \times Q)}^{\tilde{Y}}(\tilde{W}) \cong M \otimes N$, since $P \times Q$ acts trivially on \tilde{W} and on $M \otimes N$ and since $\operatorname{Res}_C^{\tilde{Y}}(\tilde{W}) = \operatorname{Res}_C^Y(W) \cong M \otimes N$. Thus, the $I \times J$ -stable simple $F[C(P \times Q)]$ -module $M \otimes N$ can be extended to \tilde{Y} and the proof is complete.

14 The maximal module of a *p*-permutation equivalence

Throughout this section we assume that G and H are finite groups and that the *p*-modular system $(\mathbb{K}, \mathcal{O}, F)$ is large enough for G and H. Further, we assume that A is a block algebra of $\mathcal{O}G$, that B is a block algebra of $\mathcal{O}H$, and that $\gamma \in T_o^{\Delta}(A, B)$ is a *p*-permutation equivalence.

We will take a closer look at the Brauer pairs (and therefore vertices) of the indecomposable modules appearing in γ . We establish that there is a unique indecomposable module M appearing in γ such that $\mathcal{BP}_{\mathcal{O}}(M) = \mathcal{BP}_{\mathcal{O}}(\gamma)$ and $\mathcal{BP}_{\mathcal{O}}(M) \supset \mathcal{BP}_{\mathcal{O}}(N)$ for all other indecomposable modules N appearing in γ . Moreover, we show that the Brauer construction of M with respect to a vertex of M yields a p-permutation bimodule that induces a Morita equivalences between the Brauer correspondents of A and B.

All results and definitions in this section have obvious analogues if A and B are block algebras over F.

14.1 Theorem Up to isomorphism, there exists a unique indecomposable (A, B)-bimodule M appearing in γ whose vertex is of the form $\Delta(D, \phi, E)$ where D is a defect group of A. Moreover, the multiplicity of [M] in γ equals 1 or -1.

Proof Existence: Let $(\Delta(D, \varphi, E), e \otimes f^*) \in \mathcal{BP}_{\mathcal{O}}(\gamma)$ be a maximal γ -Brauer pair. Then, by Theorem 10.11(c), D is a defect group of A. Since $e\gamma(\Delta(D, \phi, E))f \neq 0$, there exists an indecomposable (A, B)-bimodule M appearing in γ with $eM(\Delta(D, \phi, E))f \neq \{0\}$. Thus, $(\Delta(D, \phi, E), e \otimes f^*) \in \mathcal{BP}_{\mathcal{O}}(M)$ and $\Delta(D, \phi, E)$ is contained in a vertex of M, by Proposition 5.3. Since M belongs to the block $A \otimes B^*$ with defect group $D \times E$ and M has twisted diagonal vertex, $\Delta(D, \phi, E)$ must be a vertex of M.

Uniqueness: Let N be an indecomposable (A, B)-bimodule appearing in γ which has a vertex of the form $\Delta(\tilde{D}, \tilde{\phi}, \tilde{E})$, where \tilde{D} is a defect group of A. We will show that N is isomorphic to M from above. Choose block idempotents \tilde{e} and \tilde{f} of $\mathcal{O}C_G(\tilde{D})$ and $\mathcal{O}C_H(\tilde{E})$, respectively, such that $(\Delta(\tilde{D}, \tilde{\phi}, \tilde{E}), \tilde{e} \otimes \tilde{f}^*)$ is a maximal N-Brauer pair. By Lemma 10.7 and Proposition 10.8(i) \iff (iv), $(\Delta(\tilde{D}, \tilde{\phi}, \tilde{E}), \tilde{e} \otimes \tilde{f}^*)$ is a γ -Brauer pair. Moreover, (\tilde{D}, \tilde{e}) is a maximal A-Brauer pair and Theorem 10.11(c) implies that $(\Delta(\tilde{D}, \tilde{\phi}, \tilde{E}), \tilde{e} \otimes \tilde{f}^*)$ is a maximal γ -Brauer pair. By Theorem 10.11(b) and Proposition 5.3(c), we may choose $(\Delta(\tilde{D}, \tilde{\phi}, \tilde{E}), \tilde{e} \otimes \tilde{f}^*)$ to be equal to $(\Delta(D, \phi, E), e \otimes f^*)$. By Lemma 9.3(a), the coefficients of [M] (resp. [N]) in γ equals the coefficient of $[eM(\Delta(D, \phi, E), e \otimes f^*)]$ (resp. $eN(\Delta(D, \phi, E))f$) in $e\overline{\gamma}(\Delta(D, \phi, E))f \in T(FN_{G \times H}(\Delta(D, \phi, E), e \otimes f^*))$). But, by Proposition 13.2(c), the element $e\overline{\gamma}(D, \phi, E)f \in T(FN_{G \times H}(\Delta(D, \phi, E), e \otimes f^*))$ is of the form $\varepsilon \cdot [L]$ for some $\varepsilon \in \{\pm 1\}$ and some indecomposable $FN_{G \times H}(\Delta(D, \phi, E), e \otimes f^*)$)-module L. Thus, since both $eM(\Delta(D, \phi, E))f$ and $eN(\Delta(D, \phi, E))f$ appear in $e\overline{\gamma}(\Delta(D, \phi, E))f$, we have

$$eM(\Delta(D,\phi,E))f \cong L \cong eN(\Delta(D,\phi,E))f$$
.

Now, Proposition 5.4 implies that $M \cong N$. Finally, the above statement about multiplicities implies that the multiplicity of M in γ is equal to ε .

14.2 Definition The module M from Theorem 14.1 is called the *maximal module* of γ and its multiplicity $\varepsilon \in \{\pm 1\}$ in γ is called the *sign* of γ .

14.3 Theorem Let N be an indecomposable (A, B)-bimodule appearing in γ and let M be the maximal module of γ . Then $\mathcal{BP}_{\mathcal{O}}(N) \subseteq \mathcal{BP}_{\mathcal{O}}(M) = \mathcal{BP}_{\mathcal{O}}(\gamma)$. If $M \not\cong N$ then $\mathcal{BP}_{\mathcal{O}}(N) \subset \mathcal{BP}_{\mathcal{O}}(M)$. In particular, every vertex of N is contained in a vertex of M, with strict containment if $N \not\cong M$.

Proof By Lemma 10.7, we have $\mathcal{BP}_{\mathcal{O}}(N) \subseteq \mathcal{BP}_{\mathcal{O}}(\gamma)$. To see that $\mathcal{BP}_{\mathcal{O}}(M) = \mathcal{BP}_{\mathcal{O}}(\gamma)$, it suffices to show that the maximal γ -Brauer pairs and the maximal M-Brauer pairs coincide, see Proposition 5.3(b) and Theorem 10.11(a). If $(\Delta(D, \phi, E), e \otimes f^*)$ is a maximal M-Brauer pair then it is a γ -Brauer pair by the first statement, and even a maximal γ -Brauer pair by Theorem 10.11(c). Conversely, if $(\Delta(D, \phi, E), e \otimes f^*)$ is a maximal γ -Brauer pair then by the existence part of the proof of Theorem 14.1, it is also an M-Brauer pair, and by the already established inclusion also a maximal M-Brauer pair. Finally, if $M \ncong N$, then, by the definition of M, the vertices of N have smaller order than the vertices of M so that the inclusion $\mathcal{BP}_{\mathcal{O}}(N) \subset \mathcal{BP}_{\mathcal{O}}(M)$ is proper.

14.4 Proposition Let $(\Delta(D, \phi, E), e \otimes f^*)$ be a maximal γ -Brauer pair, let M be the maximal module of γ , and let $C_G(D) \leq S \leq N_G(D, e)$ and $C_H(E) \leq T \leq N_H(E, f)$ be intermediate groups that correspond under the isomorphism in Proposition 11.1. Let $L \in \mathcal{O}[N_{G \times H}(\Delta(D, \phi, E))]$ triv be the Green correspondent of M. Then the p-permutation (FSe, FTf)-bimodule

$$\operatorname{Ind}_{N_{S\times T}(\Delta(D,\phi,E))}^{S\times T}((e\otimes f^{*})L)$$

induces a Morita equivalence between OSe and OTf.

Proof This follows immediately from Theorem 11.4 noting that $e\gamma(\Delta(D, \phi, E))f = \pm [(e \otimes f^*)L] \in T(\mathcal{O}N_{G \times H}(\Delta(D, \phi, E), e \otimes f^*))$ by Proposition 3.3 (c), Theorem 14.1, and Theorem 14.3.

14.5 Theorem Let $\Delta(D, \phi, E)$ be a vertex of the maximal module M of γ and let A' (resp. B') be the block algebra of $\mathcal{O}N_G(D)$ (resp. $\mathcal{O}N_H(E)$) that is in Brauer correspondence with A (resp. B) via Brauer's First Main Theorem. Furthermore, let $L \in \mathcal{O}[N_{G \times H}(\Delta(D, \phi, E))]$ triv be the Green correspondent of M. Then the p-permutation (A', B')-bimodule

$$\operatorname{Ind}_{N_G \times H}^{N_G(D) \times N_H(E)}(L) \tag{33}$$

induces a Morita equivalence between A' and B'.

Proof There exist block idempotents e of $\mathcal{O}C_G(D)$ and f of $\mathcal{O}C_H(E)$ such that $(\Delta(D, \phi, E), e \otimes f^*)$ is a maximal M-Brauer pair and therefore a maximal γ -Brauer pair by Theorem 14.3. Set $I := N_G(D, e)$ and $J := N_H(E, f)$. By Proposition 14.4, the p-permutation $(\mathcal{O}N_G(D, e), \mathcal{O}N_H(E, f))$ -bimodule $\operatorname{Ind}_{N_{I\times J}(\Delta(D, \phi, E))}^{I\times J}((e \otimes f^*)L)$ induces a Morita equivalence between $\mathcal{O}Ie$ and $\mathcal{O}Jf$. Moreover, if e' (resp. f') denotes the identity element of A' (resp. B') then the $(\mathcal{O}N_G(D)e', \mathcal{O}Ie)$ -bimodule $\mathcal{O}N_G(D)e = e'\mathcal{O}N_G(D)e$ (resp. the $(\mathcal{O}Jf, \mathcal{O}N_H(E)f')$ -bimodule $f\mathcal{O}N_H(E) = f\mathcal{O}N_H(E)f'$) induces a Morita equivalence between $\mathcal{O}N_G(D)e' = A'$ and $\mathcal{O}Ie$ (resp. between $\mathcal{O}Jf$ and $\mathcal{O}N_H(E)f' = B'$). Thus, the (A', B')-bimodule

$$\mathcal{O}N_G(D)e \otimes_{\mathcal{O}Ie} \operatorname{Ind}_{N_{I\times J}(\Delta(D,\phi,E))}^{I\times J} ((e\otimes f^*)L) \otimes_{\mathcal{O}Jf} f\mathcal{O}N_H(E)$$

induces a Morita equivalence between A' and B'. However, the latter (A', B')-bimodule is isomorphic to

$$\mathrm{Ind}_{N_{I\times J}(\Delta(D,\phi,E))}^{N_{G}(D)\times N_{H}(E)}((e\otimes f^{*})L)\cong\mathrm{Ind}_{N_{G\times H}(\Delta(D,\phi,E))}^{N_{G}(D)\times N_{H}(E)}\mathrm{Ind}_{N_{I\times J}(\Delta(D,\phi,E))}^{N_{G\times H}(\Delta(D,\phi,E))}((e\otimes f^{*})L)$$

which is isomorphic to the (A', B')-bimodule in (33), since $\operatorname{Ind}_{N_{I \times J}(\Delta(D, \phi, E))}^{N_{G \times H}(\Delta(D, \phi, E))}((e \otimes f^*)L) \cong L.$

15 Connection with isotypies and splendid Rickard equivalences

Throughout this section we assume that G and H are finite groups and that the *p*-modular system ($\mathbb{K}, \mathcal{O}, F$) is large enough for G and H. Further, we assume that A is a block algebra of $\mathcal{O}G$, that B is a block algebra of $\mathcal{O}H$. We will establish that a splendid Rickard equivalence between A and B induces a *p*-permutation equivalence between A and B and that a *p*-permutation equivalence between A and B induces an isotypy between A and B.

15.1 Definition A splendid Rickard equivalence between A and B is a bounded chain complex C_{\bullet} of p-permutation (A, B)-bimodules satisfying the following properties:

(i) For every $n \in \mathbb{Z}$, the vertices of indecomposable direct summands of C_n are twisted diagonal subgroups of $G \times H$.

(ii) One has $C_{\bullet} \otimes_B C_{\bullet}^{\circ} \cong A$ in the homotopy category of chain complexes of (A, A)-bimodules and $C_{\bullet}^{\circ} \otimes_A C_{\bullet} \cong B$ in the homotopy category of chain complexes of (B, B)-bimodules. Here, A denotes the chain complex with only one non-zero term A in degree 0, and C_{\bullet}° denotes the \mathcal{O} -dual of the chain complex C_{\bullet} .

The proof of the following theorem can be easily adapted from the proof of [BX08, Theorem 1.5].

15.2 Theorem If C_{\bullet} is a splendid Rickard equivalence between A and B then

$$\gamma := \sum_{n \in \mathbb{Z}} (-1)^n [C_n] \in T^{\Delta}(A, B)$$

is a p-permutation equivalence between A and B.

Next we will show that *p*-permutation equivalences induce isotypies. The following definition is due to Broué, cf. Definition 4.6 and the subsequent Remark 2 in [Br90] and Definition 2.1 in [Br95].

15.3 Definition An *isotypy* between A and B consists of the following data:

• Maximal Brauer pairs $(D, e) \in \mathcal{BP}_{\mathcal{O}}(A)$ and $(E, f) \in \mathcal{BP}_{\mathcal{O}}(B)$;

• an isomorphism $\phi: E \xrightarrow{\sim} D$ which is also an isomorphism between the fusion system \mathcal{B} of B associated to (E, f) and the fusion system \mathcal{A} of A associated to (D, e); and

• a family of perfect isometries $\mu_Q \in R(\mathbb{K}C_G(\phi(Q))e_{\phi(Q)}, \mathbb{K}C_H(Q)f_Q), Q \leq E$, where f_Q denotes the unique block idempotent of $\mathcal{O}C_H(Q)$ with $(Q, f_Q) \leq (E, f)$ and, for $P \leq D$, e_P denotes the unique block idempotent of $\mathcal{O}C_G(P)$ with $(P, e_P) \leq (D, e)$.

These data are subject to the following conditions:

(i) (Equivariance) For every Brauer pair $(\Delta(P, \psi, Q), e_P \otimes f_Q^*) \leq (\Delta(D, \phi, E), e \otimes f^*)$ and every $(g, h) \in G \times H$ such that also ${}^{(g,h)}(\Delta(P, \psi, Q), e_P \otimes f_Q^*) \leq (\Delta(D, \phi, E), e \otimes f^*)$, one has ${}^{(g,h)}\mu_Q = \mu_{{}^{(h_Q)}}$.

(ii) (Compatibility) For every $Q \leq E$ and every $y \in C_E(Q)$, setting $P := \phi(Q)$, $x := \phi(y)$, $Q' := Q\langle y \rangle$, and $P' := P\langle x \rangle$, the diagram

commutes, where I_Q denotes the K-linear extension of the group homomorphism $I_{\mu_Q} = \mu_Q \cdot -: R(\mathbb{K}C_H(Q)f_Q) \to R(\mathbb{K}C_G(P)e_P)$; cf. 8.1(b),(c),(d) for notation and Remark 8.4(e) for the bottom map in the above diagram.

15.4 Theorem Let $\gamma \in T^{\Delta}(A, B)$ be a *p*-permutation equivalence between A and B and let $(\Delta(D, \phi, E), e \otimes f^*) \in \mathcal{BP}_{\mathcal{O}}(\gamma)$ be a maximal γ -Brauer pair. For every $P \leq D$ (resp. $Q \leq E$) let $(P, e_P) \in \mathcal{BP}_{\mathcal{O}}(A)$ (resp. $(Q, f_Q) \in \mathcal{BP}_{\mathcal{O}}(B)$) denote the unique Brauer pair with $(P, e_P) \leq (D, e)$ (resp. $(Q, f_Q) \leq (E, f)$).

Then the data (D, e), (E, f), ϕ , and, for $Q \leq E$, the restriction μ_Q to $C_G(\phi(Q)) \times C_H(Q)$ of the element

$$e_{\phi(Q)}\mu(\Delta(\phi(Q),\phi,Q))f_Q \in R(\mathbb{K}N_{G\times H}(\Delta(\phi(Q),\phi,Q),e_{\phi(Q)}\otimes f_Q^*))$$

form an isotypy between A and B.

Proof By Theorem 10.11(c), (D, e) is a maximal A-Brauer pair and (E, f) is a maximal B-Brauer pair. By Theorem 11.2 the isomorphism $\phi: E \xrightarrow{\sim} D$ is an isomorphism between the fusion system \mathcal{B} of B associated to (E, f) and the fusion system \mathcal{A} of A associated to (D, e). Let $Q \leq E$ and set $P := \phi(Q)$. Then $(\Delta(P, \phi, Q), e_P \otimes f_Q^*) \leq (\Delta(D, \phi, E), e \otimes f^*)$ (see Remark 10.2(c)) and, by Theorem 10.11(a), $(\Delta(P, \phi, Q), e_P \otimes f_Q^*)$ is a γ -Brauer pair. Now, Proposition 11.9(a), implies that μ_Q is a perfect isometry between $\mathbb{K}C_G(P)e_P$ and $\mathbb{K}C_H(Q)f_Q$. Therefore the data have the required properties.

To see that the perfect isometries μ_Q , $Q \leq E$, satisfy the equivariance axiom (i), let $Q \leq E$, $P := \phi(Q)$, and $(g,h) \in G \times H$ such that

$${}^{(g,h)}(\Delta(P,\phi,Q), e_P \otimes f_Q^*) \leqslant (\Delta(D,\phi,E), e \otimes f^*).$$

$$(34)$$

In order to prove that ${}^{(g,h)}\mu_Q = \mu_{({}^hQ)}$, it suffices to show that

$$g^{(h)}\overline{\gamma}(\Delta(P,\phi,Q),e_P\otimes f_Q^*) = \overline{\gamma}(\Delta(\phi({}^hQ),\phi,{}^hQ),e_{\phi({}^hQ)}\otimes f^*_{({}^hQ)}).$$
(35)

Since the Brauer construction commutes with conjugation, the left hand side is equal to $\overline{\gamma}({}^{(g,h)}(\Delta(P,\phi,Q),e_P\otimes f_Q^*))$ and ${}^{(g,h)}(\Delta(P,\phi,Q),e_P\otimes f_Q^*) = (\Delta({}^{g}P,c_g\phi c_h^{-1}, {}^{h}Q), {}^{g}e_P\otimes {}^{h}(f_Q^*))$. Moreover, by (34) and Remark 10.2(c), we obtain $\Delta({}^{g}P,c_g\phi c_h^{-1}, {}^{h}Q) = \Delta(\phi({}^{h}Q),\phi,{}^{h}Q), ({}^{g}P,{}^{g}e_P) \leq (D,e)$, and $({}^{h}Q,{}^{h}f_Q) \leq (E,f)$. Thus, ${}^{g}P = \phi({}^{h}Q), {}^{g}e_P = e_{({}^{g}P)} = e_{\phi({}^{h}Q)}$, and ${}^{h}f_Q = f_{({}^{h}Q)}$. This establishes Equation (35).

Finally, we show that the perfect isometries μ_Q , $Q \leq E$, satisfy the compatibility axiom (ii). Let $Q \leq E$ and $y \in C_E(Q)$ and set $P := \phi(Q)$, $x := \phi(y)$, $Q' := Q\langle y \rangle$, and $P' := P\langle x \rangle = \phi(Q')$.

Let $\gamma_Q \in T^{\Delta}(\mathcal{O}C_G(P)e_P, \mathcal{O}C_H(Q)f_Q)$ denote the restriction of $e_P\gamma(\Delta(P,\phi,Q))f_Q$ to $C_G(P) \times C_H(Q)$ and let $\gamma_{Q'}$ be defined similarly. Recall from [BX08, 2.1] the definition of a linear source $\mathcal{O}G$ -module and its associated representation group $L(\mathcal{O}G)$. Moreover, recall from [BX08, 2.3] the map $-(\langle x \rangle, x) \colon L(\mathcal{O}C_G(P)e_P) \to$ $L(\mathcal{O}C_G(P')\mathrm{br}_{\langle x \rangle}(e_P))$. The map $-(\langle x \rangle, x)e_{P'}$ in the diagram below is defined as the composition of the map $-(\langle x \rangle, x)$ with the natural projection from $\mathbb{K}L(\mathcal{O}C_G(P')\mathrm{br}_{\langle x \rangle}(e_P)) \to \mathbb{K}L(\mathcal{O}C_G(P')e_{P'})$, noting that $\mathrm{br}_{\langle x \rangle}(e_P)e_{P'} = e_{P'}$. Similarly, we define the map $-(\langle y \rangle, y)f_{Q'}$. For any finite group X let $\pi_X \colon L(\mathcal{O}X) \to$ $T(FX) \to R(FX)$ denote the composition of the homomorphism $L(\mathcal{O}X) \to T(FX)$ induced by the functor $F \otimes_{\mathcal{O}} -: \mathcal{O}_X \operatorname{mod} \to F_X \operatorname{mod}$ and the map $\eta_X \colon T(FX) \to R(FX)$ from the diagram in 9.1(c). Moreover, $\kappa_X \colon L(\mathcal{O}X) \to R(\mathbb{K}X)$ will denote the homomorphism induced by the functor $\mathbb{K} \otimes_{\mathcal{O}} -: \mathcal{O}_X \operatorname{mod} \to \mathbb{K}_X \operatorname{mod}$. Now consider the following diagram:



Note that the diagram in the compatibility axiom equals the front square diagram involving the two generalized decomposition maps in the cube-like diagram. By Brauer's induction theorem, the map $\kappa_{C_H(Q)}: L(\mathcal{O}C_H(Q)f_Q) \to R(\mathbb{K}C_H(Q)f_Q)$ is surjective. Therefore, it suffices to show that the front square diagram commutes after precomposing with $\kappa_{C_H(Q)}$. It follows that the front square diagram commutes if the left, right and ceiling square diagram in the cube-like diagram commute and the concatenation of the rear and floor square diagrams commute. Clearly, the ceiling square of the cube (involving the κ -maps) commutes. Moreover, the left and right squares commute by Theorem 2.4 in [BX08]. Thus, it suffices now to show that the foor square commutes.

So let V be an indecomposable linear source $\mathcal{O}C_H(Q)f_Q$ -module. We need to show that

$$e_{P'}\big(\overline{\gamma_{Q}}_{C_{H}(Q)}[V]\big)(\langle x\rangle, x) = \overline{\gamma_{Q'}}_{C_{H}(Q')}(f_{Q'}[\overline{V}(\langle y\rangle, y)]) \in \mathbb{K}R(FC_{G}(P')e_{P'}),$$

in the notation of [BX08]. By the definition of $-(\langle x \rangle, x)$ and $-(\langle y \rangle, y)$ in [BX08, 2.3] it suffices to show that for every $\theta \in \text{Hom}(\langle x \rangle, \mathcal{O}^{\times})$, one has

$$e_{P'}(\overline{\gamma_Q}_{C_H(Q)}[V])(\langle x \rangle, \theta) = \overline{\gamma_{Q'}}_{C_H(Q')}[f_{Q'}\overline{V}(\langle y \rangle, \theta \circ \phi)].$$
(36)

By a slight variation of Lemma 3.5(d) in [BX08], identifying $\langle y \rangle$ and $\langle x \rangle$ via ϕ , we see that the left hand side of Equation (36) is equal to

$$\sum_{\substack{\rho,\sigma\in\mathrm{Hom}(\langle x\rangle,\mathcal{O}^{\times})\\ \sigma\sigma\sigma=\mathcal{A}}} e_{P'}\overline{\gamma_Q}(\Delta(\langle x\rangle,\phi,\langle y\rangle),\tilde{\rho}) \stackrel{\cdot}{\underset{FC_H(Q')}{\longrightarrow}} \overline{V}(\langle y\rangle,\sigma\circ\phi) \in \mathbb{K}R(FC_G(P')e_{P'}),$$

where $\tilde{\rho}$ is defined as $\rho \circ p_1$ on $\Delta(\langle x \rangle, \phi, \langle y \rangle)$. But since γ_Q is a virtual *p*-permutation module, one has $\overline{\gamma_Q}(\Delta(\langle x \rangle, \phi, \langle y \rangle), \tilde{\rho}) = 0$, unless $\rho = 1$ is the trivial homomorphism, and in this case one has $\overline{\gamma_Q}(\Delta(\langle x \rangle, \phi, \langle y \rangle), 1) = \overline{\gamma_Q}(\Delta(\langle x \rangle, \phi, \langle y \rangle))$, the usual Brauer construction. Thus, the left hand side of Equation (36) is equal to

$$e_{P'}\overline{\gamma_Q}(\Delta(\langle x \rangle, \phi, \langle y \rangle)) : \underset{C_H(Q')}{\cdot} [\overline{V}(\langle y \rangle, \theta \circ \phi)].$$

Since $\gamma_Q = e_P \gamma(\Delta(P, \phi, Q)) f_Q$, Proposition 3.5(b) and Lemma 3.7 imply that

$$e_{P'}\overline{\gamma_Q}(\Delta(\langle x\rangle,\phi,\langle y\rangle)) = e_{P'}\mathrm{br}_{\langle x\rangle}(e_P)\overline{\gamma}(\Delta(P',\phi,Q'))\mathrm{br}_{\langle y\rangle}(f_Q) = e_{P'}\overline{\gamma}(\Delta(P',\phi,Q'))\mathrm{br}_{Q'}(f_Q).$$
(37)

Here we used that $e_{P'} \operatorname{br}_{\langle x \rangle}(e_P) = e_{P'} \operatorname{br}_{P'}(e_P) = e_{P'}$, since $P \leq P'$ and $(P, e_P) \leq (P', e_{P'})$, see Proposition 4.2(b). We claim that the last expression in (37) is equal to $e_{P'}\overline{\gamma}(\Delta(P', \phi, Q'))f_{Q'}$. In fact, assume that f' is a primitive idempotent of $Z(\mathcal{O}C_H(Q'))$ such that $e_{P'}\overline{\gamma}(\Delta(P', \phi, Q'))f' \neq 0$. Then Corollary 10.9 implies that there exists $h \in H$ such that $(\phi, (Q', f_{Q'})) = {}^{h}(\phi, (Q', f')) = (\phi \circ c_h^{-1}, ({}^{h}Q', {}^{h}f'))$. But $Q' = {}^{h}Q'$ and $\phi = \phi \circ c_h^{-1}$ imply that $h \in C_H(Q')$, and we obtain $f_{Q'} = {}^{h}f' = f'$. Moreover, since $\operatorname{br}_{Q'}(f_Q)f_{Q'} = f_{Q'}$, the claim is proved. Thus, the left hand side of Equation (36) is equal to

$$e_{P'}\overline{\gamma}(\Delta(P',\phi,Q'))f_{Q'} : \overline{C_H(Q')}[\overline{V}(\langle y \rangle, \theta \circ \phi)],$$

as desired, and the proof is complete.

Acknowledgment. The first author would like to express his gratitude to the BERNOULLI CENTER at the EPFL, where some part of this research was achieved during his stay in November and December 2016 for the program 'Local representation theory and simple groups'.

References

[AB79] J. ALPERIN, M. BROUÉ: Local methods in block theory. Ann. of Math. (2) 110 (1979), 143–157.

[AKO11] M. ASCHBACHER, R. KESSAR, B. OLIVER: Fusion systems in algebra and topology. London Mathematical Society Lecture Note Series, 391. Cambridge University Press, Cambridge, 2011.

- [Be98] D.J. BENSON: Representations and cohomology. I. Basic representation theory of finite groups and associative algebras. Cambridge University Press, Cambridge, 1998.
- [BD12] R. BOLTJE, S. DANZ: A ghost ring for the left-free double Burnside ring and an application to fusion systems. Adv. Math. 229 (2012), 1688–1733.
- [BX08] R. BOLTJE, B. XU: On *p*-permutation equivalences: between Rickard equivalences and isotypies. Trans. Amer. Math. Soc. **360** (2008), 5067–5087.
- [Bc10a] S. BOUC: Biset functors for finite groups, Lecture Notes in Mathematics, 1990. Springer-Verlag, Berlin, 2010
- [Bc10b] S. BOUC: Bisets as categories and tensor product of induced bimodules. Appl. Categ. Structures 18 (2010), 517–521.
- [Br85] M. BROUÉ: On Scott modules and *p*-permutation modules: an approach through the Brauer morphism. Proc. Amer. Math. Soc. **93** (1985), 401–408.
- [Br90] M. BROUÉ: Isométries parfaites, types de blocs, catégories dérivées. Astrisque No. **181–182** (1990), 61–92.
- [Br95] M. BROUÉ: Rickard equivalences and block theory. Groups '93 Galway/St. Andrews, Vol. 1 (Galway, 1993), 58–79, London Math. Soc. Lecture Note Ser., 211, Cambridge Univ. Press, Cambridge, 1995
- [BrP80] M. BROUÉ, L. PUIG: Characters and local structure in G-algebras. J. Algebra 63 (1980), 306-317.
- [K07] R. KESSAR: Introduction to block theory. In: Group representation theory, 47–77, EPFL Press, Lausanne, 2007.
- [L06] M. LINCKELMANN: Simple fusion systems and the Solomon 2-local groups. J. Algebra 296 (2006), 385–401.
- [L09] M. LINCKELMANN: Trivial source bimodule rings for blocks and p-permutation equivalences. Trans. Amer. Math. Soc. **361** (2009), 1279–1316.
- [L18] M. LINCKELMANN: The block theory of finite group algebras. Vol. I and II. London Mathematical Society. Cambridge University Press, Cambridge, 2018.
- [NT89] H. NAGAO, Y. TSUSHIMA: Representations of finite groups. Academic Press, Boston, 1989.
- [R96] J. RICKARD: Splendid equivalences: derived categories and permutation modules. Proc. London Math. Soc. (3) 72 (1996), 331–358.
- [P99] L. PUIG: On the local structure of Morita and Rickard equivalences between Brauer blocks. Progress in Mathematics, 178. Birkhäuser Verlag, Basel, 1999.
- [S90] D. SIBLEY: Vertices, blocks and virtual characters. J. Algebra 132 (1990), 501–507.