APPLICATIONS OF A CHANGE OF MEASURES TECHNIQUE FOR COMPOUND MIXED RENEWAL PROCESSES TO THE RUIN PROBLEM

S. M. TZANINIS

ABSTRACT. In the present paper the change of measures technique for compound mixed renewal processes, developed in Tzaninis & Macheras [24], is applied to the ruin problem in order to compute the ruin probability and to find upper and lower bounds for it.

Key Words: Compound mixed renewal process, Change of measures, Progressively equivalent measures, Regular conditional probabilities, Ruin probability.

AMS Subject Classification (2010): Primary 60G55, 91B30 ; secondary 28A35, 60A10, 60G44, 60K05.

1. INTRODUCTION

The change of measures technique has been successfully applied to various theories such as queues and fluid flows (Asmussen [1],1994, [2] 1995, Palmowski and Rolski [16] 1996; [17] 1998), ruin theory (Dassios and Embrechts [6] 1989, Asmussen [1] 1996, Asmussen and Albrecher [3] 2010, Schmidli [20] 1996, [21] 1997, [22] 2017), simulation (Boogaert and De Waegenaere [4] 1990, Ridder [18] 1996) and pricing of insurance risks (premium calculation principles) (Delbaen and Haezendonck [7] 1989, Lyberopoulos and Macheras [12] 2019, Macheras and Tzaninis [15] 2020, [24] 2020). The process of interest is usually Markovian and, under a suitably chosen new probability measure, it is again a Markov process with some "nicer" desired properties.

In [24], the same problem was investigated for the class of compound mixed renewal processes, which are not, in general, Markov ones. In the same paper, given a compound mixed renewal process S under P, a full characterization of all probability measures Q, which are progressively equivalent to P and preserve the type of S, but with some better desired properties, was provided, see [24], Theorem 4.5 and Corollary 4.8 as well as Proposition 4.15. Note that the martingales L^r and the probability measures Q^r appearing in [22] are special instances of the martingales $\widetilde{M}^{(\gamma)}(\theta)$ and of the probability measures Q_{θ} of Theorem 4.5 from [24].

Part of [24], Proposition 4.15, is Proposition 3.5, formulated for the purposes of the present paper and being the starting point for applications to the ruin problem. Proposition 3.5, as well as Proposition 3.7, extend the corresponding results for the renewal risk model (see e.g. [22], Lemmas 8.4 and 8.6, respectively) to the compound mixed renewal processes.

A consequence of Proposition 3.5 is Theorem 4.1, where it is proven, that if the net profit condition is fulfilled under an original measure P, and S is a compound mixed renewal process under P then under the new measure resulting from Proposition 3.5 the process S will be of the same type, except that the net profit condition will no longer be fulfilled and ruin will always occur within finite time.

Date: July 21, 2020.

Thus, the ruin problem becomes easier to handle, since by Theorem 4.1 under the new measure the probability of ruin is equal to 1, giving us the opportunity to express the ruin probability under P as a quantity under the new measure, see Proposition 4.3, and to find upper and lower bounds for it, see Proposition 4.4.

2. Preliminaries

Throughout this paper, unless stated otherwise, (Ω, Σ, P) is a fixed but arbitrary probability space. The symbol $\mathcal{L}^1(P)$ stands for the family of all real-valued P-integrable functions on Ω . Functions that are *P*-a.s. equal are not identified. We denote by $\sigma(\mathcal{G})$ the σ -algebra generated by a family \mathcal{G} of subsets of Ω . Given a topology \mathfrak{T} on Ω we write $\mathfrak{B}(\Omega)$ for its **Borel** σ -algebra on Ω , i.e. the σ -algebra generated by \mathfrak{T} . Our measure theoretic terminology is standard and generally follows [5]. For the definitions of real-valued random variables and random variables we refer to [5], p. 308. We apply the notation $P_X := P_X(\theta) := \mathbf{K}(\theta)$ to mean that X is distributed according to the law $\mathbf{K}(\theta)$, where $\theta \in D \subseteq \mathbb{R}^d$ $(d \in \mathbb{N})$ is the parameter of the distribution. We denote again by $\mathbf{K}(\theta)$ the distribution function induced by the probability distribution $\mathbf{K}(\theta)$. Notation $\mathbf{Ga}(b, a)$, where $a, b \in (0, \infty)$, stands for the law of gamma distribution (cf. e.g. [23], p. 180). In particular, $\mathbf{Ga}(b, 1) = \mathbf{Exp}(b)$ stands for the law of exponential distribution. For two real-valued random variables X and Y we write X = Y P-a.s. if $\{X \neq Y\}$ is a P-null set. If $A \subseteq \Omega$, then $A^c := \Omega \setminus A$, while χ_A denotes the indicator (or characteristic) function of the set A. For a map $f: D \longrightarrow E$ and for a non-empty set $A \subseteq D$ we denote by $f \upharpoonright A$ the restriction of f to A. We write $\mathbb{E}_{P}[X \mid \mathcal{F}]$ for a version of a conditional expectation (under P) of $X \in \mathcal{L}^1(P)$ given a σ -subalgebra \mathcal{F} of Σ . For $X := \chi_E \in \mathcal{L}^1(P)$ with $E \in \Sigma$ we set $P(E \mid \mathcal{F}) := \mathbb{E}_P[\chi_E \mid \mathcal{F}]$. For the unexplained terminology of Probability and Risk Theory we refer to [23].

Given two measurable spaces (Ω, Σ) and (Υ, H) , a function k from $\Omega \times H$ into [0, 1] is a Σ -H-Markov kernel if it has the following properties:

(k1) The set-function $B \mapsto k(\omega, B)$ is a probability measure on H for any fixed $\omega \in \Omega$. (k2) The function $\omega \mapsto k(\omega, B)$ is Σ -measurable for any fixed $B \in H$.

In particular, given a real-valued random variable X on Ω and a d-dimensional random vector Θ on Ω , a **conditional distribution of** X **over** Θ is a $\sigma(\Theta)$ - \mathfrak{B} -Markov kernel denoted by $P_{X|\Theta} := P_{X|\sigma(\Theta)}$ and satisfying for each $B \in \mathfrak{B}$ condition

$$P_{X|\Theta}(\bullet, B) = P(X^{-1}[B] \mid \sigma(\Theta))(\bullet) \quad P \upharpoonright \sigma(\Theta) - a.s..$$

Clearly, for every \mathfrak{B}_d - \mathfrak{B} -Markov kernel k, the map $K(\Theta)$ from $\Omega \times \mathfrak{B}$ into [0,1] defined by means of

$$K(\Theta)(\omega, B) := (k(\bullet, B) \circ \Theta)(\omega) \text{ for any } (\omega, B) \in \Omega \times \mathfrak{B}$$

is a $\sigma(\Theta)$ - \mathfrak{B} -Markov kernel. Then for $\theta = \Theta(\omega)$ with $\omega \in \Omega$ the probability measures $k(\theta, \bullet)$ are distributions on \mathfrak{B} and so we may write $\mathbf{K}(\theta)(\bullet)$ instead of $k(\theta, \bullet)$. Consequently, in this case $K(\Theta)$ will be denoted by $\mathbf{K}(\Theta)$.

For any real-valued random variables X, Y on Ω we say that $P_{X|\Theta}$ and $P_{Y|\Theta}$ are $P \upharpoonright \sigma(\Theta)$ -equivalent and we write $P_{X|\Theta} = P_{Y|\Theta} P \upharpoonright \sigma(\Theta)$ -a.s., if there exists a *P*-null set $M \in \sigma(\Theta)$ such that for any $\omega \notin M$ and $B \in \mathfrak{B}$ the equality $P_{X|\Theta}(\omega, B) = P_{Y|\Theta}(\omega, B)$ holds true.

For the definition of a *P*-conditionally (stochastically) independent process over $\sigma(\Theta)$ as well as of a *P*-conditionally identically distributed process over $\sigma(\Theta)$ we refer to [24]. We

say that a process is *P*-conditionally (stochastically) independent or identically distributed given Θ , if it is conditionally independent or identically distributed over the σ -algebra $\sigma(\Theta)$.

For the rest of the paper we simply write "conditionally" in the place of "conditionally given Θ whenever conditioning refers to Θ .

Henceforth, unless stated otherwise, $(\Upsilon, H) := ((0, \infty), \mathfrak{B}(\Upsilon))$ and Θ is a d-dimensional random vector on Ω with values on $D \subseteq \mathbb{R}^d$ $(d \in \mathbb{N})$.

3. A Change of Measures Technique for Compound Mixed Renewal Processes

We first recall some additional background material, needed in this section.

A family $N := \{N_t\}_{t \in \mathbb{R}_+}$ of random variables from (Ω, Σ) into $(\mathbb{R}, \mathfrak{B}(\mathbb{R}))$ is called a **counting** (or **claim number**) **process**, if there exists a P-null set $\Omega_N \in \Sigma$ such that the process N restricted on $\Omega \setminus \Omega_N$ takes values in $\mathbb{N}_0 \cup \{\infty\}$, has right-continuous paths, presents jumps of size (at most) one, vanishes at t = 0 and increases to infinity. Without loss of generality we may and do assume, that $\Omega_N = \emptyset$. Denote by $T := \{T_n\}_{n \in \mathbb{N}_0}$ and $W := \{W_n\}_{n \in \mathbb{N}}$ the (claim) arrival process and (claim) interarrival process, respectively (cf. e.g. [23], Section 1.1, page 6 for the definitions) associated with N. Note also that every arrival process induces a counting process, and vice versa (cf. e.g. [23], Theorem 2.1.1).

Furthermore, let $X := \{X_n\}_{n \in \mathbb{N}}$ be a sequence of positive real-valued random variables on Ω , and for any $t \ge 0$ define

$$S_t := \begin{cases} \sum_{k=1}^{N_t} X_k & \text{if } t > 0; \\ 0 & \text{if } t = 0. \end{cases}$$

Accordingly, the sequence X is said to be the claim size process, and the family $S := \{S_t\}_{t \in \mathbb{R}_+}$ of real-valued random variables on Ω is said to be the **aggregate claims process** induced by the pair (N, X). Recall that a pair (N, X) is called a **risk process**, if N is a counting process, X is P-i.i.d. and the processes N and X are P-independent (see [23], Chapter 6, Section 6.1).

The following definition has been introduced in [11], Definition 3.1, see also [14], Definition 3.2(b).

Definition 3.1. A counting process N is a P-mixed renewal process with mixing parameter Θ and interarrival time conditional distribution $\mathbf{K}(\Theta)$ (written P-MRP($\mathbf{K}(\Theta)$) for short), if the induced interarrival process W is P-conditionally independent and

$$\forall n \in \mathbb{N} \qquad [P_{W_n | \Theta} = \mathbf{K}(\Theta) \quad \mathbf{P} \upharpoonright \sigma(\Theta) \text{-a.s.}].$$

In particular, if the distribution P_{Θ} of Θ is degenerate at some point $\theta_0 \in D$, then the counting process N becomes a *P*-renewal process with interarrival time distribution $\mathbf{K}(\theta_0)$ (written P-RP($\mathbf{K}(\theta_0)$) for short).

Accordingly, an aggregate claims process S induced by a P-risk process (N, X) such that N is a P-MRP $(\mathbf{K}(\Theta))$ is called a **compound mixed renewal process with parameters K** (Θ) and P_{X_1} (P-CMRP $(\mathbf{K}(\Theta), P_{X_1})$ for short). In particular, if P_{Θ} is degenerate at $\theta_0 \in D$, then S is called a **compound renewal process with parameters K** (θ_0) and P_{X_1} (P-CRP $(\mathbf{K}(\theta_0), P_{X_1})$ for short).

Throughout what follows we denote again by $\mathbf{K}(\Theta)$ and $\mathbf{K}(\theta)$ the conditional distribution function and the distribution function induced by the conditional probability distribution $\mathbf{K}(\Theta)$ and the probability distribution $\mathbf{K}(\theta)$, respectively.

The following conditions will be useful for our investigations:

- (a1) the pair (W, X) is *P*-conditionally independent;
- (a2) the random vector Θ and the process X are P-(unconditionally) independent.

Next, whenever condition (a1) and (a2) holds true we shall write that the quadruplet (P, W, X, Θ) or (if no confusion arises) the probability measure P satisfies (a1) and (a2), respectively.

Since conditioning is involved in the definition of (compound) mixed renewal processes, it is natural to expect that regular conditional probabilities (or disintegrations) will play a key. To this purpose we recall the following definition.

Definition 3.2. Let (Υ, H, R) be a probability probability space. A family $\{P_y\}_{y \in \Upsilon}$ of probability measures on Σ is called a **regular conditional probability** (rcp for short) of P over R if

(d1) for each $E \in \Sigma$ the map $y \mapsto P_y(E)$ is *H*-measurable;

(d2) $\int P_y(E) R(dy) = P(E)$ for each $E \in \Sigma$.

If $f : \Omega \longrightarrow \Upsilon$ is an inverse-measure-preserving function (i.e. $P(f^{-1}(B)) = R(B)$ for each $B \in H$), a rcp $\{P_y\}_{y \in \Upsilon}$ of P over R is called **consistent** with f if, for each $B \in H$, the equality $P_y(f^{-1}(B)) = 1$ holds for R-almost every $y \in B$.

From now on, unless stated otherwise, the family $\{P_{\theta}\}_{\theta \in D}$ is a rcp of P over P_{Θ} consistent with Θ .

Regular conditional probabilities seem to have a bad reputation when it comes to applications, and that is probably due to the fact that their own existence is not always guaranteed (see [13], Examples 4 and 5). Nevertheless, as the spaces used in applied Probability Theory are mainly Polish ones, such rcps always exist (see [8], Theorem 6), and in fact they can be explicitly constructed for the class of (compound) mixed renewal processes (see [24], Proposition 4.1).

We write $\mathcal{F} := {\mathcal{F}_t}_{t\in\mathbb{R}_+}$, where $\mathcal{F}_t := \sigma(\mathcal{F}_t^S \cup \sigma(\Theta))$, for the canonical filtration generated by S and Θ , $\mathcal{F}_{\infty}^S := \sigma(\bigcup_{t\in\mathbb{R}_+} \mathcal{F}_t^S)$ and $\mathcal{F}_{\infty} := \sigma(\mathcal{F}_{\infty}^S \cup \sigma(\Theta))$. For the definition of a (P, \mathcal{Z}) -martingale in $\mathcal{L}^1(P)$, where $\mathcal{Z} = {\mathcal{Z}_t}_{t\in\mathbb{R}_+}$ is a filtration for (Ω, Σ) , we refer to [23], p. 25. A (P, \mathcal{Z}) -martingale ${Z_t}_{t\in\mathbb{R}_+}$ in $\mathcal{L}^1(P)$ is P-a.s. positive, if Z_t is P-a.s. positive for each $t \ge 0$. For $\mathcal{Z} = \mathcal{F}$ we write "martingale in $\mathcal{L}^1(P)$ " instead of " (P, \mathcal{Z}) -martingale in $\mathcal{L}^1(P)$ ", for simplicity.

Notations 3.3. (a) The class of all real-valued $\mathfrak{B}(\Upsilon)$ -measurable functions γ such that $\mathbb{E}_P\left[e^{\gamma(X_1)}\right] = 1$ will be denoted by $\mathcal{F}_P := \mathcal{F}_{P,X_1,\ln}$. The class of all real-valued $\mathfrak{B}(D)$ -measurable functions ξ on D such that $P_{\Theta}(\{\xi > 0\}) = 1$ and $\mathbb{E}_P[\xi(\Theta)] = 1$ is denoted by $\mathcal{R}_+(D) := \mathcal{R}_+(D, \mathfrak{B}(D), P_{\Theta})$.

(b) Denote by $\mathfrak{M}^k(D)$ $(k \in \mathbb{N})$ the class of all $\mathfrak{B}(D)-\mathfrak{B}(\mathbb{R}^k)$ -measurable functions on D. For each $\rho \in \mathfrak{M}^k(D)$ the class of all probability measures Q on Σ satisfying (a1) and (a2), are **progressively equivalent** to P, i.e. $Q \upharpoonright \mathcal{F}_t \sim P \upharpoonright \mathcal{F}_t$ for any $t \ge 0$ (in the sense of absolute continuity), and such that S is a Q-CMRP $(\Lambda(\rho(\Theta)), Q_{X_1})$ is denoted by $\mathcal{M}_{S,\Lambda(\rho(\Theta))} := \mathcal{M}_{S,\Lambda(\rho(\Theta)),P,X_1}$. In the special case d = k and $\rho := id_D$ we write $\mathcal{M}_{S,\Lambda(\Theta)} := \mathcal{M}_{S,\Lambda(\rho(\Theta))}$ for simplicity.

(c) For given $\rho \in \mathfrak{M}^{k}(D)$ and $\theta \in D$, denote by $\mathcal{M}_{S,\Lambda(\rho(\theta))}$ the class of all probability measures Q_{θ} on Σ , such that $Q_{\theta} \upharpoonright \mathcal{F}_{t} \sim P_{\theta} \upharpoonright \mathcal{F}_{t}$ for any $t \in \mathbb{R}_{+}$ and S is a Q_{θ} -CRP $(\Lambda(\rho(\theta)), (Q_{\theta})_{X_{1}})$.

From now on, unless stated otherwise, $P \in \mathcal{M}_{S,\mathbf{K}(\Theta)}$ is the initial probability measure under which S is a $P-CMRP(\mathbf{K}(\Theta), P_{X_1})$.

It follows a result to show that the martingales $L^r := \{L^r_t\}_{t \in \mathbb{R}_+}$ and the measures Q^r appearing in [22], Chapter 8, Section 8.3, are special instances of the martingales $\widetilde{M}^{(\gamma)}(\theta)$ and the measures Q_{θ} , respectively, of the main result of [24], i.e. Theorem 4.5.

Remark 3.4. For any $r \in \mathbb{R}_+$ such that $\mathbb{E}_P[e^{rX_1}] < \infty$ and for any $\theta \in L_P^c$, where L_P is the P_{Θ} -null appearing in [24], Proposition 3.3, let $\kappa_{\theta}(r)$ be the unique solution to the equation

(1)
$$M_{X_1}(r) \cdot (M_{\theta})_{W_1} \left(-\kappa_{\theta}(r) - c(\theta) \cdot r \right) = 1,$$

where M_{X_1} and $(M_{\theta})_{W_1}$ are the moment generating function of X_1 and W_1 under the measures P and P_{θ} , respectively. (Such a solution exists by e.g. [19], Lemma 11.5.1(a)). Define the function $\kappa : D \times \mathbb{R}_+ \longrightarrow \mathbb{R}$ by means of

$$\kappa(\theta, r) := \kappa_{\theta}(r) \text{ for any } (\theta, r) \in D \times \mathbb{R}_+$$

and for fixed $r \in \mathbb{R}_+$ denote by κ_{Θ} the random variable defined by the formula

 $\kappa_{\Theta}(r)(\omega) := \kappa_{\Theta(\omega)}(r) \text{ for any } \omega \in \Omega.$

Then, due to [24], Lemma 4.13, $\kappa_{\Theta}(r)$ is the $P \upharpoonright \sigma(\Theta)$ -a.s. unique solution to the equation

(2)
$$M_{X_1}(r) \cdot \mathbb{E}_P\left[e^{-\left(\kappa_{\Theta}(r) + c(\Theta) \cdot r\right)W_1} \mid \Theta\right] = 1 \quad P \upharpoonright \sigma(\Theta) - \text{a.s.}.$$

The following proposition is a part of Proposition 4.15 from [24]. Since it is the basic tool for the proofs of our results, we restate it exactly in the form needed for our purposes.

Proposition 3.5. For any $r \in \mathbb{R}_+$ such that $\mathbb{E}_P[e^{rX_1}] < \infty$, and for any $\theta \notin L_P$, let $\kappa_{\theta}(r)$ be the unique solution to the equation (1), and let $\kappa_{\Theta}(r)$ be as in Remark 3.4. Fix on arbitrary $r \in \mathbb{R}_+$ as above and let $\rho \in \mathfrak{M}^k(D)$ be given.

For each pair $(\gamma, \xi) \in \mathcal{F}_P \times \mathcal{R}_+(D)$ with $\gamma(x) := r \cdot x - \ln \mathbb{E}_P[e^{r \cdot X_1}]$ for any $x \in \Upsilon$, there exists a unique probability measure $Q := Q^r \in \mathcal{M}_{S, \Lambda(\rho(\Theta))}$, where

$$\boldsymbol{\Lambda}(\rho(\Theta))(B) := \frac{\mathbb{E}_{P}[\chi_{W_{1}^{-1}[B_{2}]} \cdot e^{-(\kappa_{\Theta}(r) + c(\Theta) \cdot r) \cdot W_{1}} \mid \Theta]}{\mathbb{E}_{P}[e^{-(\kappa_{\Theta}(r) + c(\Theta) \cdot r) \cdot W_{1}} \mid \Theta]} \quad P \upharpoonright \sigma(\Theta) - a.s.$$

for any $B \in \mathfrak{B}(\Upsilon)$, determined by condition

$$(RRM_{\xi}) \qquad \qquad Q(A) = \int_{A} M_{t}^{(\gamma)}(\Theta) dP \quad for \ all \quad 0 \leq u \leq t \quad and \quad A \in \mathcal{F}_{u}$$

with the martingale $M^{(\gamma)}(\Theta)$ in $\mathcal{L}^1(P)$ fulfilling condition

$$M_t^{(\gamma)}(\Theta) = \xi(\Theta) \cdot \widetilde{M}_t^{(\gamma)}(\Theta) \quad P \upharpoonright \sigma(\Theta) - a.s..$$

Moreover, there exist an essentially unique rcp $\{Q_{\theta}\}_{\theta \in D} := \{Q_{\theta}^{r}\}_{\theta \in D}$ of Q over Q_{Θ} consistent with Θ and a P_{Θ} -null set $L_{**} \in \mathfrak{B}(D)$, satisfying for any $\theta \notin L_{**}$ conditions $Q_{\theta} \in \mathcal{M}_{S, \mathbf{\Lambda}(\rho(\theta))}$ and

$$(RRM_{\theta}) \qquad \qquad Q_{\theta}(A) = \int_{A} \widetilde{M}_{t}^{(\gamma)}(\theta) \, dP_{\theta} \quad \text{for all } 0 \leq u \leq t \text{ and } A \in \mathcal{F}_{u}$$
5

with the martingale $\widetilde{M}_t^{(\gamma)}(\theta)$ in $\mathcal{L}^1(P_{\theta})$ fulfilling condition

$$\widetilde{M}_{t}^{(\gamma)}(\theta) = e^{r \cdot S_{t} - (\kappa_{\theta}(r) + c(\theta) \cdot r) \cdot T_{N_{t}} + \ln \mathbb{E}_{P}[e^{r \cdot X_{1}}]} \cdot \frac{\int_{t-T_{N_{t}}}^{\infty} e^{-(\kappa_{\theta}(r) + c(\theta) \cdot r) \cdot w} (P_{\theta})_{W_{1}}(dw)}{1 - \mathbf{K}(\theta)(t - T_{N_{t}})}$$

In particular, if P_{W_1} is absolutely continuous with respect to the Lebesgue measure λ restricted to $\mathfrak{B}([0,1])$, then the martingale $L^r(\theta) := \{L_t^r(\theta)\}_{t \in \mathbb{R}_+}$ for $r \in \mathbb{R}_+$, appearing in [22], Lemma 8.4, coincides with the martingale $\widetilde{M}^{(\gamma)}(\theta)$ in $\mathcal{L}^1(P_\theta)$ for any $\theta \notin L_{**}$, and for any $t \in \mathbb{R}_+$ condition

$$M_t^{(\gamma)}(\Theta) = \xi(\Theta) \cdot L_t^r(\Theta)$$

holds $P \upharpoonright \sigma(\Theta)$ -a.s. true.

Lemma 3.6. For any $r \in \mathbb{R}_+$, $\theta \notin L_{**}$, Q_{θ}^r and $\kappa_{\theta}(r)$ as in Proposition 3.5 condition

$$\kappa_{\theta}'(r) = \frac{\mathbb{E}_{Q_{\theta}^r}[X_1]}{\mathbb{E}_{Q_{\theta}^r}[W_1]} - c(\theta),$$

holds true.

Proof. Fix on arbitrary $r \in \mathbb{R}_+$ and $\theta \notin L_{**}$ as in Proposition 3.5.

Since $L_P \subseteq L_{**}$ by [24], Theorem 4.5, it follows by [24], Proposition 3.3, that we can rewrite condition (1) in the form

(3)
$$(M_{\theta})_{X_1}(r) \cdot (M_{\theta})_{W_1}(-c(\theta) \cdot r - \kappa_{\theta}(r)) = 1.$$

Differentiation with respect to r gives

(4)
$$\begin{pmatrix} (M_{\theta})_{X_1}(r))' \cdot (M_{\theta})_{W_1}(-c(\theta) \cdot r - \kappa_{\theta}(r)) \\ + (M_{\theta})_{X_1}(r) \cdot ((M_{\theta})_{W_1}(-c(\theta) \cdot r - \kappa_{\theta}(r)))' \cdot (-c(\theta) - \kappa_{\theta}'(r)) = 0 \end{cases}$$

for all r in a neighbourhood of 0. The expectations $\mathbb{E}_{Q_{\theta}}^{r}[X_{1}]$ and $\mathbb{E}_{Q_{\theta}}^{r}[W_{1}]$ are given by

$$\mathbb{E}_{Q_{\theta}^{r}}[X_{1}] = \mathbb{E}_{P_{\theta}}\left[X_{1} \cdot \frac{e^{r \cdot X_{1}}}{\mathbb{E}_{P_{\theta}}[e^{r \cdot X_{1}}]}\right] = \frac{\mathbb{E}_{P_{\theta}}\left[X_{1} \cdot e^{r \cdot X_{1}}\right]}{\mathbb{E}_{P_{\theta}}[e^{r \cdot X_{1}}]} = \frac{(M_{X_{1}}^{\theta}(r))'}{M_{X_{1}}^{\theta}(r)}$$

and

$$\mathbb{E}_{Q_{\theta}^{r}}[W_{1}] = \mathbb{E}_{P_{\theta}}\left[W_{1} \cdot \frac{e^{-(r \cdot c(\theta) + \kappa_{\theta}(r)) \cdot W_{1}}}{\mathbb{E}_{P_{\theta}}[e^{-(r \cdot (\theta)c + \kappa_{\theta}(r)) \cdot W_{1}}]}\right] = \frac{\mathbb{E}_{P_{\theta}}\left[W_{1} \cdot e^{-(r \cdot c(\theta) + \kappa_{\theta}(r)) \cdot W_{1}}\right]}{\mathbb{E}_{P_{\theta}}[e^{-(r \cdot c(\theta) + \kappa_{\theta}(r)) \cdot W_{1}}]}$$
$$= \frac{\left((M_{\theta})_{W_{1}}(-r \cdot c(\theta) - \kappa_{\theta}(r))\right)'}{\left((M_{\theta})_{W_{1}}(-r \cdot c(\theta) - \kappa_{\theta}(r))\right)},$$

respectively, implying along with condition (4) that

$$\mathbb{E}_{Q_{\theta}^{r}}[X_{1}] \cdot (M_{\theta})_{X_{1}}(r) \cdot (M_{\theta})_{W_{1}}(-c(\theta) \cdot r - \kappa_{\theta}(r)) + (M_{\theta})_{X_{1}}(r) \cdot \mathbb{E}_{Q_{\theta}^{r}}[W_{1}] \cdot (M_{\theta})_{W_{1}}(-c(\theta) \cdot r - \kappa_{\theta}(r)) \cdot (-c(\theta) - \kappa_{\theta}'(r)) = 0.$$

The latter together with condition (3) gives

$$\mathbb{E}_{Q_{\theta}^{r}}[X_{1}] + \mathbb{E}_{Q_{\theta}^{r}}[W_{1}] \cdot (-c(\theta) - \kappa_{\theta}'(r)) = 0,$$

completing the proof.

Let S be the aggregate claims process induced by the counting process N and the claim size process X. Fix on arbitrary $u \in \Upsilon$ and $t \in \mathbb{R}_+$, and define the function $r_t^u : \Omega \times D \longrightarrow \mathbb{R}$ by means of $r_t^u(\omega, \theta) := u + c(\theta) \cdot t - S_t(\omega)$ for any $(\omega, \theta) \in \Omega \times D$, where c is a positive $\mathfrak{B}(D)$ measurable function. For arbitrary but fixed $\theta \in D$, the process $r^u(\theta) := \{r^u_t(\theta)\}_{t \in \mathbb{R}_+}$ defined by $r_t^u(\theta) := r_t^u(\omega, \theta)$ for any $\omega \in \Omega$, is called the **reserve process** induced by the initial reserve u, the premium intensity or premium rate $c(\theta)$ and the aggregate claims process S (see [23], Section 7.1, pages 155-156 for the definition). The function ψ_{θ} defined by $\psi_{\theta}(u) := P_{\theta}(\{\inf r_t^u(\theta) < 0\})$ is called the **probability of ruin** for the reserve process $r^u(\theta)$ with respect to P_{θ} (see [23], Section 7.1, page 158 for the definition).

Define the real-valued functions $r_t^u(\Theta)$ and R_t^u on Ω by means of $r_t^u(\Theta)(\omega) := r_t^u(\omega, \Theta(\omega))$ for any $\omega \in \Omega$, and $R_t^u := r_t^u \circ (id_\Omega \times \Theta)$, respectively. The process $R^u := \{R_t^u\}_{t \in \mathbb{R}_+}$ is called the reserve process induced by the initial reserve u, the stochastic premium intensity or stochastic premium rate $c(\Theta)$ and the aggregate claims process S. The function ψ defined by $\psi(u) := P(\{\inf R_t^u < 0\})$ is called the **probability of ruin** for the reserve process R^u with respect to P.

The following proposition extends Lemma 8.6 of [22].

Proposition 3.7. For any $r \in \mathbb{R}_+$, $\theta \notin L_{**}$, Q_{θ}^r and $\kappa_{\theta}(r)$ as in Proposition 3.5, the following statements hold true:

- (i) $\lim_{t\to\infty} \frac{r_t^u(\theta)-u}{t} = -\kappa_{\theta}'(r)$ $Q_{\theta}^r a.s.;$ (ii) $\lim_{t\to\infty} \frac{R_t^u-u}{t} = -\kappa_{\Theta}'(r)$ $Q^r a.s..$
- (iii) if there exists a P_{Θ} -null set \hat{L}_1 in $\mathfrak{B}(D)$ such that $P_{\theta} = Q_{\theta}^r$ for any $\theta \notin \hat{L}_1$, then the measures P and Q^r are equivalent on \mathcal{F}_{∞} ;
- (iv) if there exists a P_{Θ} -null set \hat{L}_2 in $\mathfrak{B}(D)$ such that $P_{\theta} \neq Q_{\theta}^r$ for any $\theta \notin \hat{L}_2$, then the measures P and Q^r are singular on \mathcal{F}_{∞} , i.e. there exists a set $E \in \mathcal{F}_{\infty}$ such that P(E) = 0 if and only if $Q^{r}(E) = 1$.

Proof. Fix on arbitrary $r \in \mathbb{R}_+$ as in Proposition 3.5.

Ad (i): Let us fix on arbitrary $\theta \notin L_{**}$, and note that $L_P \subseteq L_{**}$ by [24], Theorem 4.5. Since S is a Q_{θ}^{r} -CRP by [24], Proposition 3.3, we get by the strong law of large numbers that

$$\lim_{t \to \infty} \frac{S_t}{t} = \frac{\mathbb{E}_{Q_{\theta}^r}[X_1]}{\mathbb{E}_{Q_{\theta}^r}[W_1]} \quad Q_{\theta}^r \text{-a.s.}$$

(cf. e.g. [9], Section 1.2, Theorem 2.3), or equivalently that

$$\lim_{t \to \infty} \frac{r_t^u(\theta) - u}{t} = \lim_{t \to \infty} \frac{c(\theta) \cdot t - S_t}{t} = c(\theta) - \lim_{t \to \infty} \frac{S_t}{t} = c(\theta) - \frac{\mathbb{E}_{Q_\theta^r}[X_1]}{\mathbb{E}_{Q_\theta^r}[W_1]} \qquad Q_\theta^r \text{-a.s.},$$

implying along with Lemma 3.6, assertion (i).

Ad (ii): Consider the function $v := \chi_{\{\lim_{t \to \infty} \frac{r_t^u - u}{t} = -\kappa'_{\bullet}(r)\}} : \Omega \times D \longrightarrow [0, 1] \text{ and put } g :=$ $v \circ (id_{\Omega} \times \Theta) = \chi_{\left\{\lim_{t \to \infty} \frac{r_t^u(\Theta) - u}{t} = -\kappa'_{\Theta}(r)\right\}}.$ Since $v \in \mathcal{L}^1(M)$, where $M := P \circ (id_{\Omega} \times \Theta)^{-1}$, we may apply [10], Proposition 3.8(i), to get that

$$\mathbb{E}_{Q^r}\left[g \mid \Theta\right] = \mathbb{E}_{Q^r_{\bullet}}\left[v^{\bullet}\right] \circ \Theta \quad Q^r \upharpoonright \sigma(\Theta) \text{-a.s.}$$

or equivalently

$$Q^r\left(\left\{\lim_{t\to\infty}\frac{r_t^u(\Theta)-u}{t}=-\kappa'_{\Theta}(r)\right\}\mid\Theta\right)=Q^r_{\bullet}\left(\left\{\lim_{t\to\infty}\frac{r_t^u(\bullet)-u}{t}=-\kappa'_{\bullet}(r)\right\}\right)\circ\Theta\quad Q^r\upharpoonright\sigma(\Theta)-\text{a.s.}$$

Then for any $F \in \mathfrak{B}(D)$ we get

$$\begin{split} &\int_{\Theta^{-1}[F]} Q^r \left(\left\{ \lim_{t \to \infty} \frac{r_t^u(\Theta) - u}{t} = -\kappa'_{\Theta}(r) \right\} \mid \Theta \right) \, dQ^r \\ &= \int_{\Theta^{-1}(F)} Q^r_{\bullet} \left(\left\{ \lim_{t \to \infty} \frac{r_t^u(\bullet) - u}{t} = -\kappa'_{\bullet}(r) \right\} \right) \circ \Theta \, dQ^r \\ &= \int_F Q^r_{\theta} \left(\left\{ \lim_{t \to \infty} \frac{r_t^u(\theta) - u}{t} = -\kappa'_{\theta}(r) \right\} \right) \, Q^r_{\Theta}(d\theta) \\ &= \int_{F \cap L^c_{**}} Q^r_{\theta} \left(\left\{ \lim_{t \to \infty} \frac{r_t^u(\theta) - u}{t} = -\kappa'_{\theta}(r) \right\} \right) \, Q^r_{\Theta}(d\theta) \\ &= \int_{\Theta^{-1}(F)} \, dQ^r, \end{split}$$

where the last equality follows by (i); hence

$$Q^r\left(\left\{\lim_{t\to\infty}\frac{r_t^u(\Theta)-u}{t}=-\kappa'_{\Theta}(r)\right\}\mid\Theta\right)=1\quad Q^r\upharpoonright\sigma(\Theta)\text{-a.s.},$$

implying

$$Q^r\left(\left\{\lim_{t\to\infty}\frac{r_t^u(\Theta)-u}{t}=-\kappa'_{\Theta}(r)\right\}\right)=\int Q^r\left(\left\{\lim_{t\to\infty}\frac{r_t^u(\Theta)-u}{t}=-\kappa'_{\Theta}(r)\right\}\mid\Theta\right)\,dQ^r=1,$$

that is assertion (ii) holds true.

The proof of the statements (iii) and (iv) follow by Proposition 3.5 together with [24], Proposition 3.11.

4. Applications to the Ruin Problem

In this section we present the main result of the paper, where an explicit formula for the probability of ruin for the reserve process in the case of compound mixed renewal processes is proven. Before we formulate it, we need to establish the validity of the following theorem, which is a consequence of Proposition 3.5, and allows us to construct a probability measure Q^{R^*} , being singular to the original probability measure P and such that ruin occurs Q^{R^*} -a.s..

Theorem 4.1. Let $r \in \mathbb{R}_+$ and $\theta \notin L_{**}$ be as in Proposition 3.5. If for any $\theta \notin L_{**}$ the net profit condition is fulfilled with respect to P_{θ} , i.e.

$$c(\theta) > \frac{\mathbb{E}_{P_{\theta}}[X_1]}{\mathbb{E}_{P_{\theta}}[W_1]},$$

then there exists an adjustment coefficient $R(\theta) \in \Upsilon$ with respect to P_{θ} . In particular, if the $\sup_{\theta \in L_{**}^c} R(\theta) =: R^*$ exists in Υ , $\mathbb{E}_P[e^{R^* \cdot X_1}] < \infty$ and P_{W_1} is absolutely continuous with respect to the Lebesgue measure λ restricted to $\mathfrak{B}([0,1])$, then for any pair (γ,ξ) as in Proposition 3.5 there exist a unique probability measure $Q^{R^*} \in \mathcal{M}_{S,\Lambda(\rho(\theta))}$ determined by condition (RRM_{ξ}) , and a rcp $\{Q_{\theta}^{R^*}\}_{\theta \in D}$ of Q^{R^*} over $Q_{\theta}^{R^*}$ consistent with Θ satisfying conditions $Q_{\theta}^{R^*} \in \mathcal{M}_{S,\Lambda(\rho(\theta))}$ and (RRM_{θ}) for any $\theta \notin L_{**}$, and such that for any u > 0the probabilities of ruin $\psi_{\theta}^{R^*}(u)$ and $\psi^{R^*}(u)$ with respect to $Q_{\theta}^{R^*}$ and Q^{R^*} , respectively, are equal to 1. *Proof.* Fix on arbitrary $\theta \notin L_{**}$ and assume that $c(\theta) > \frac{\mathbb{E}_{P_{\theta}}[X_1]}{\mathbb{E}_{P_{\theta}}[W_1]}$. It then follows by e.g. [22], page 133, that there exists an adjustment coefficient $R(\theta)$ with respect to P_{θ} .

In particular, assume that $R^* \in \Upsilon$, $\mathbb{E}_P[e^{R^* \cdot X_1}] < \infty$ and that P_{W_1} is absolutely continuous with respect to the Lebesgue measure λ restricted to $\mathfrak{B}([0,1])$. By Proposition 3.5 there exist a unique probability measure $Q^{R^*} \in \mathcal{M}_{S,\Lambda(\rho(\Theta))}$ determined by condition (RRM_{ξ}) , and a rcp $\{Q_{\theta}^{R^*}\}_{\theta\in D}$ of Q^{R^*} over $Q_{\Theta}^{R^*}$ consistent with Θ satisfying conditions $Q_{\theta}^{R^*} \in \mathcal{M}_{S,\Lambda(\rho(\theta))}$ and (RRM_{θ}) . Because $\kappa''_{\theta}(r) > 0$ by e.g. [22], p. 133, we get that the function κ_{θ} is strictly convex, or equivalently that κ'_{θ} is strictly increasing. Thus, since by e.g. [22], page 133, we have that $\kappa_{\theta}(0) = \kappa_{\theta}(R(\theta)) = 0$ and $\kappa'_{\theta}(0) < 0$, it follows that there exists a point $r_0 \in (0, R(\theta))$ such that $\kappa'_{\theta}(r_0) = 0$; hence $\kappa'_{\theta}(r) > 0$ for any $r > r_0$. Because $r_0 < R(\theta) \leq R^*$ we deduce that $\kappa'_{\theta}(R^*) > 0$. The latter, along with Lemma 3.6, yields that

$$0 < \frac{\mathbb{E}_{Q_{\theta}^{R*}}[X_1]}{\mathbb{E}_{Q_{\theta}^{R*}}[W_1]} - c(\theta) \Longleftrightarrow c(\theta) < \frac{\mathbb{E}_{Q_{\theta}^{R*}}[X_1]}{\mathbb{E}_{Q_{\theta}^{R*}}[W_1]},$$

implying that the net profit condition is violated with respect to $Q_{\theta}^{R^*}$; hence by [23], Corollary 7.1.4, we obtain

$$\psi_{\theta}^{R^*}(u) = Q_{\theta}^{R^*}(\{\inf_{t \in \mathbb{R}_+} r_t^u(\theta) < 0\}) = 1 \quad \text{for any} \ u > 0,$$

implying along with [24], Remark 3.4(b) that

$$\psi^{R^*}(u) = Q^{R^*}(\{\inf_{t \in \mathbb{R}_+} R^u_t < 0\}) = \int_D \psi^{R^*}_{\theta}(u) Q^{R^*}_{\Theta}(d\theta) = 1 \quad \text{for any} \ u > 0,$$

completing the whole proof.

It follows an example where the assumptions $R^* < \infty$ and $\mathbb{E}_P[e^{R^*X_1}] < \infty$ of Theorem 4.1 hold.

Example 4.2. Assume that S is a P-CMPP(Θ) such that $P_{X_1} = \mathbf{Exp}(\eta), \eta \in \Upsilon$, and $P_{\Theta} = \mathbf{Beta}(a, b), a, b \in (0, \infty)$ i.e.

$$\mathbf{Beta}(a,b)(B) := \int_B \frac{1}{B(a,b)} \cdot \theta^{a-1} \cdot (1-\theta)^{b-1} \lambda(d\theta) \quad \text{for any } B \in \mathfrak{B}((0,1)).$$

According to [24], Proposition 3.3, there exists a P_{Θ} -null set $L_P \in \mathfrak{B}((0,1))$ such that S is a P_{θ} -CPP(θ) for any $\theta \notin L_P$. Fix on an arbitrary $\theta \notin L_P$ and assume that $c(\theta) = \frac{2\cdot\theta}{\eta\cdot(1+\theta)}$. Applying [23], Theorem 7.4.5, we get that $R(\theta) = \eta - \frac{\theta}{c(\theta)} = \frac{\eta\cdot(1-\theta)}{2} \in (0,\eta)$ is an adjustment coefficient with respect to P_{θ} . Since $\sup_{\theta \in L_P^c} R(\theta) = \sup_{\theta \in L_P^c} \frac{\eta\cdot(1-\theta)}{2} = \frac{\eta}{2}$, we obtain $R^* \in \Upsilon$. Furthermore, $\mathbb{E}_P[e^{R^*X_1}] = \frac{\eta}{\eta-R^*} < \infty$.

Denote by τ the **ruin time** of the portfolio of an insurance company (cf. e.g. [22], p. 84 for the definition). The next proposition is a consequence of Theorem 4.1.

Proposition 4.3. Let $r \in \mathbb{R}_+$ be as in Proposition 3.5, $u \in \Upsilon$ and $\theta \notin L_{**}$. Denote by $\psi_{\theta}(u)$ and $\psi(u)$ the probabilities of ruin with respect to the probability measures P_{θ} and P, respectively. Under the assumptions of Theorem 4.1 the following hold:

(i)
$$\psi_{\theta}(u) = \mathbb{E}_{Q_{\theta}^{R*}} \left[e^{R^* r_{\tau}^u(\theta) + \kappa_{\theta}(R^*) \cdot \tau} \right] \cdot e^{-R^* u};$$

(ii) $\psi(u) = \mathbb{E}_{Q^{R*}} \left[\frac{e^{R^* \cdot R_{\tau}^u + \kappa_{\Theta}(R^*) \cdot \tau}}{\xi(\Theta)} \right] \cdot e^{-R^* \cdot u}.$

Proof. Fix on arbitrary $u \in \Upsilon$

Ad (i): Let $\theta \notin L_{**}$ be arbitrary but fixed. Since $L^{R^*}(\theta)$ is an a.s. positive martingale in $\mathcal{L}^1(P_{\theta})$ by Proposition 3.5, and τ is a stopping time for \mathcal{F} , we may apply [22], Lemma 8.1, to get

 $\psi_{\theta}(u)$

$$= \int_{\{\tau < \infty\}} \frac{1}{L_{\tau}^{R^*}(\theta)} dQ_{\theta}^{R^*}$$

$$= \int_{\{\tau < \infty\}} \frac{e^{R^* \cdot (r_{\tau}^u(\theta) - u) - (\kappa_{\theta}(R^*) + c(\theta) \cdot R^*) \cdot (\tau - T_{N_{\tau}}) + \kappa_{\theta}(R^*) \cdot \tau - \ln \mathbb{E}_{P}[e^{R^* \cdot X_{1}}] \cdot (1 - \mathbf{K}(\theta)(\tau - T_{N_{\tau}}))}{\int_{\tau - T_{N_{\tau}}}^{\infty} e^{-(\kappa_{\theta}(R^*) + c(\theta) \cdot R^*) \cdot w} (P_{\theta})_{W_{1}}(dw)} dQ_{\theta}^{R^*}$$

$$= \int_{\{\tau < \infty\}} e^{R^* \cdot (r_{\tau}^u(\theta) - u) + \kappa_{\theta}(R^*) \cdot \tau} dQ_{\theta}^{R^*};$$

where the last equality follows from condition (3) for $r = R^*$ and the fact that $\tau - T_{N_{\tau}} = 0$; hence

$$\psi_{\theta}(u) = \mathbb{E}_{Q_{\theta}^{R^{*}}} \left[\chi_{\{\tau < \infty\}} \cdot e^{R^{*} \cdot r_{\tau}^{u}(\theta) + \kappa_{\theta}(R^{*}) \cdot \tau} \right] \cdot e^{-R^{*} \cdot u}.$$

Because the probability of ruin with respect to $Q_{\theta}^{R^*}$ is equal to 1, the previous condition yields

$$\psi_{\theta}(u) = \mathbb{E}_{Q_{\theta}^{R^*}} \left[e^{R^* \cdot r_{\tau}^u(\theta) + \kappa_{\theta}(R^*) \cdot \tau} \right] \cdot e^{-R^* \cdot u},$$

that is assertion (i) holds true.

Ad (ii): Assertion (i) together with [24], Remark 3.4(b), implies

$$\begin{split} \psi(u) &= \int \psi_{\theta}(u) P_{\Theta}(d\theta) \\ &= \int \mathbb{E}_{Q_{\theta}^{R*}} \left[\frac{e^{R^{*} \cdot r_{\tau}^{u}(\theta) + \kappa_{\theta}(R^{*}) \cdot \tau}}{\xi(\theta)} \right] Q_{\Theta}^{R*}(d\theta) \cdot e^{-R^{*} \cdot u} \\ &= \int \mathbb{E}_{Q^{R*}} \left[\frac{e^{R^{*} \cdot R_{\tau}^{u} + \kappa_{\Theta}(R^{*}) \cdot \tau}}{\xi(\Theta)} \mid \Theta \right] dQ^{R*} \cdot e^{-R^{*} \cdot u}, \end{split}$$

where the last equality follows from [10], Proposition 3.8; hence

$$\psi(u) = \mathbb{E}_{Q^{R^*}}\left[\frac{e^{R^* \cdot R^u_\tau + \kappa_\Theta(R^*) \cdot \tau}}{\xi(\Theta)}\right] \cdot e^{-R^* \cdot u},$$

that is assertion (ii) holds true.

The following result shows that Proposition 4.3 along with Proposition 3.5 give us the opportunity to find upper and lower bounds of the probability of ruin under P.

Proposition 4.4. In the situation of Proposition 4.3 the following holds true:

(i)
$$\psi_{\theta}(u) \ge \mathbb{E}_{Q_{\theta}^{R^{*}}} \left[e^{R^{*} \cdot r_{\tau}^{u}(\theta)} \right] \cdot e^{-R^{*} \cdot u};$$

(ii) $\psi(u) \ge \mathbb{E}_{Q^{R^{*}}} \left[\frac{e^{R^{*} \cdot R_{\tau}^{u}}}{\xi(\Theta)} \right] \cdot e^{-R^{*} \cdot u}.$

In particular, if condition $\mathbb{E}_P[e^{R^*\Theta}] < \infty$ holds and if the function $\xi : D \longrightarrow \mathbb{R}$ is defined by means of

$$\xi(\theta) := \frac{e^{R^*\theta}}{\mathbb{E}_P[e^{R^*\Theta}]} \quad \text{for any} \quad \theta \in D,$$

then there exist a unique probability measure $\nu^{R^*} \in \mathcal{M}_{S,\Lambda(\rho(\Theta))}$ determined by condition (RRM_{ξ}) with R^* in the place of r, and a rcp $\{\nu_{\theta}^{R^*}\}_{\theta\in D}$ of ν^{R^*} over $\nu_{\Theta}^{R^*}$ consistent with Θ satisfying conditions $\nu_{\theta}^{R^*} \in \mathcal{M}_{S,\Lambda(\rho(\theta))}$ and (RRM_{θ}) for any $\theta \notin L_{**}$, and such that

$$\psi(u) \leqslant \mathbb{E}_P[e^{R^*\Theta}] \cdot \mathbb{E}_{\nu^{R^*}}\left[e^{R^* \cdot R^u_\tau + \kappa_\Theta(R^*) \cdot \tau}\right] \cdot e^{-R^* \cdot u}.$$

Proof. Because $\kappa_{\theta}(R^* > 0)$, statements (i) and (ii) follow by statements (i) and (ii) of Proposition 4.3, respectively.

In particular, if for any $r \in \mathbb{R}_+$ condition $\mathbb{E}_P[e^{R^*\Theta}] < \infty$ holds and ξ is defined as above, the $\xi \in \mathcal{R}_+(D)$ and, due to Proposition 3.5, there exist a unique probability measure $\nu^{R^*} \in \mathcal{M}_{S,\Lambda(\rho(\Theta))}$ determined by condition (RRM_{ξ}) and a rcp $\{\nu_{\theta}^{R^*}\}_{\theta\in D}$ of ν^{R^*} over $\nu_{\Theta}^{R^*}$ consistent with Θ satisfying conditions $\nu_{\theta}^{R^*} \in \mathcal{M}_{S,\Lambda(\rho(\theta))}$ and (RRM_{θ}) for any $\theta \notin L_{**}$. By Proposition 4.3 we have

$$\psi(u) = \mathbb{E}_{\nu^{R^*}} \left[\frac{e^{R^* \cdot R^u_\tau + \kappa_\Theta(R^*) \cdot \tau}}{\xi(\Theta)} \right] \cdot e^{-R^* \cdot u}$$

$$\leq \mathbb{E}_P \left[e^{R^*\Theta} \right] \cdot \mathbb{E}_{\nu^{R^*}} \left[e^{R^* \cdot R^u_\tau + \kappa_\Theta(R^*) \cdot \tau} \right] \cdot e^{-R^* \cdot u},$$

completing the whole proof.

It is worth noting that, in the Cramér-Lundberg risk model one can construct exponential martingales, and using the stopping theorem one is able to prove upper bounds for the ruin probabilities. However, this technique does not allow us to prove a lower bound. A method to find also lower bounds for the ruin probabilities is the "change of measure technique" for a compound mixed renewal process S developed above.

Example 4.5. Take D := (1, 2), let Θ be a real-valued random variable on Ω , and assume that $P \in \mathcal{M}_{S,\mathbf{G}(\Theta,2)}$, such that $P_{X_1} = \mathbf{Ga}(2,2)$ and $P_{\Theta} = \mathbf{U}(1,2)$. Since conditions (a1) and (a2) hold true, it follows by [24], Proposition 3.3, that there exists a P_{Θ} -null set $L_P \in \mathfrak{B}((1,2))$ such that $P_{\theta} \in \mathcal{M}_{S,\mathbf{Ga}(\theta,2)}$ with $P_{X_1} = (P_{\theta})_{X_1}$ for any $\theta \notin L_P$, implying that

$$\frac{\mathbb{E}_{P_{\theta}}[X_1]}{\mathbb{E}_{P_{\theta}}[W_1]} = \frac{1}{\frac{2}{\theta}} = \frac{\theta}{2} \quad \text{for any } \theta \notin L_P$$

Put $c(\theta) := \theta + 1$ for any $\theta \in D$. As a first step we are going to explicitly determine the function κ_{θ} . For any $r \in (0, 2)$ and $\theta \notin L_{**}$ applying condition (1) and an easy computation we get

$$M_{X_1}(r) \cdot (M_{\theta})_{W_1} \left(-\kappa_{\theta}(r) - c(\theta) \cdot r \right) = 1$$

$$\iff \left(\frac{2}{2-r} \right)^2 \cdot \left(\frac{\theta}{\theta + \kappa_{\theta}(r) + c(\theta) \cdot r} \right)^2 = 1$$

$$\iff \kappa_{\theta}(r) = \frac{r^2 \cdot c(\theta) + r \cdot \theta - 2 \cdot c(\theta) \cdot r}{2-r} \text{ or } \kappa_{\theta}(r) = \frac{r^2 \cdot c(\theta) + r \cdot (\theta - 2 \cdot c(\theta)) - 4\theta}{2-r},$$

equivalently

(5)
$$\kappa_{\theta}(r) = \frac{r \cdot (r \cdot \theta + r - \theta - 2)}{2 - r}$$

or

(6)
$$\kappa_{\theta}(r) = \frac{r^2(\theta+1) - r(\theta+2) - 4\theta}{2 - r},$$

\square			

respectively. By Theorem 4.1 that there exists an adjustment coefficient $R(\theta) \in (0,2)$ with respect to P_{θ} , for any $\theta \notin L_{**}$, while $R(\theta)$ is the positive solution to the equation $\kappa_{\theta}(r) = 0$ by e.g. [22], page 133. The latter along with equations (5) and (6) yields $R(\theta) = \frac{\theta+2}{\theta+1} \in (0,2)$ and

$$R(\theta) = \frac{\theta + 2 + (17\theta^2 + 20\theta + 4)^{\frac{1}{2}}}{2(\theta + 1)} > 2 \text{ or } R(\theta) = \frac{\theta + 2 - (17\theta^2 + 20\theta + 4)^{\frac{1}{2}}}{2(\theta + 1)} < 0,$$

respectively; hence $R(\theta) = \frac{\theta+2}{\theta+1}$ is the solution to (5) in (0,2). But since $R(\theta)$ is a strictly decreasing function of θ we get that $R^* = \sup_{\theta \in L_{**}^c} R(\theta) =$ $\frac{3}{2} \in (0,2)$, implying that $\mathbb{E}_P[e^{R^* \cdot X_1}] = \frac{2}{2-\frac{3}{2}} = 4 < \infty$ as well as that

$$\kappa_{\theta}(R^*) = \frac{3}{2} \cdot (\theta - 1) \quad \text{for any } \theta \notin L_{**}.$$

Put $\gamma(x) := R^* \cdot x - \ln \mathbb{E}_P[e^{R^* \cdot X_1}]$ for any $x \in \Upsilon$. By Theorem 4.1, for any $\xi \in \mathcal{R}_+(D)$ there exist a unique probability measure $Q^{R^*} \in \mathcal{M}_{S,\Lambda(\rho(\Theta))}$ determined by condition (RRM_{ξ}) , and a rcp $\{Q_{\theta}^{R^*}\}_{\theta\in D}$ of Q^{R^*} over $Q_{\Theta}^{R^*}$ consistent with Θ satisfying conditions $Q_{\theta}^{R^*} \in \mathcal{M}_{S, \mathbf{\Lambda}(\rho(\theta))}$ and (RRM_{θ}) for any $\theta \notin L_{**}$, and such that for any u > 0 the probabil-ities of ruin $\psi_{\theta}^{R^*}(u)$ and $\psi^{R^*}(u)$ with respect to $Q_{\theta}^{R^*}$ and Q^{R^*} , respectively, are equal to 1. It then follows by Proposition 4.3 that for any u > 0 and $\theta \notin L_{**}$, the ruin probabilities $\psi(u)$ and $\psi_{\theta}(u)$ satisfy conditions

$$\psi_{\theta}(u) = \mathbb{E}_{Q_{\theta}^{R^*}} \left[e^{R^* r_{\tau}^u(\theta) + \kappa_{\theta}(R^*) \cdot \tau} \right] \cdot e^{-R^* u} = \mathbb{E}_{Q_{\theta}^{R^*}} \left[e^{\frac{3}{2} \cdot r_{\tau}^u(\theta) + \frac{3}{2} \cdot (\theta - 1) \cdot \tau} \right] \cdot e^{-\frac{3}{2} \cdot u}$$

and

$$\psi(u) = \mathbb{E}_{Q^{R^*}}\left[\frac{e^{R^* \cdot R^u_\tau + \kappa_\Theta(R^*) \cdot \tau}}{\xi(\Theta)}\right] \cdot e^{-R^* \cdot u} = \mathbb{E}_{Q^{R^*}}\left[\frac{e^{\frac{3}{2} \cdot R^u_\tau + \frac{3}{2} \cdot (\Theta-1) \cdot \tau}}{\xi(\Theta)}\right] \cdot e^{-\frac{3}{2} \cdot u}$$

Example 4.6. Take $D := \Upsilon$, let Θ be a real-valued random variable on Ω , and assume that $P \in \mathcal{M}_{S,\mathbf{G}(\Theta,2)}$, such that $P_{X_1} = \mathbf{Ga}(2,2)$ and $P_{\Theta} = \mathbf{Ga}(b,a)$, where $(b,a) \in \Upsilon^2$. Since conditions (a1) and (a2) hold true, it follows by [24], Proposition 3.3, that there exists a P_{Θ} -null set $L_P \in \mathfrak{B}(D)$ such that $P_{\theta} \in \mathcal{M}_{S,\mathbf{Ga}(\theta,2)}$ with $P_{X_1} = (P_{\theta})_{X_1}$ for any $\theta \notin L_P$, implying that

$$\frac{\mathbb{E}_{P_{\theta}}[X_1]}{\mathbb{E}_{P_{\theta}}[W_1]} = \frac{1}{\frac{2}{\theta}} = \frac{\theta}{2} \quad \text{for any } \theta \notin L_P$$

Put $c(\theta) := \theta$ for any $\theta \in D$. For any $r \in (0,2)$ and $\theta \notin L_{**}$ we get as in Example 4.5 that there exists an adjustment coefficient $R(\theta) \in (0,2)$ with respect to P_{θ} being the solution to the equation

(7)
$$\kappa_{\theta}(r) = \frac{r \cdot \theta \cdot (r-1)}{2-r} = 0$$

for any $\theta \notin L_{**}$. Thus, we get $R^* = \sup_{\theta \in L_{**}^c} R(\theta) = 1 \in (0,2)$, implying that $\mathbb{E}_P[e^{R^* \cdot X_1}] =$ $\frac{2}{2-1} = 2 < \infty.$

Put $\gamma(x) := R^* \cdot x - \ln \mathbb{E}_P[e^{R^* \cdot X_1}]$ for any $x \in \Upsilon$. By Theorem 4.1 for any $\xi \in \mathcal{R}_+(D)$ there exist a unique probability measure $Q^{R^*} \in \mathcal{M}_{S,\Lambda(\rho(\Theta))}$ determined by condition (RRM_{ξ}) , and a rcp $\{Q_{\theta}^{R^*}\}_{\theta \in D}$ of Q^{R^*} over $Q_{\theta}^{R^*}$ consistent with Θ satisfying conditions $Q_{\theta}^{R^*} \in \mathcal{M}_{S, \mathbf{\Lambda}(\rho(\theta))}$ and (RRM_{θ}) for any $\theta \notin L_{**}$, and such that for any u > 0 the probabilities of ruin $\psi_{\theta}^{R^*}(u)$ and $\psi^{R^*}(u)$ with respect to $Q_{\theta}^{R^*}$ and Q^{R^*} , respectively, are equal to 1. It then follows by Proposition 4.3 that for any u > 0 and $\theta \notin L_{**}$, the ruin probabilities $\psi(u)$ and $\psi_{\theta}(u)$ satisfy conditions

$$\psi_{\theta}(u) = \mathbb{E}_{Q_{\theta}^{R*}} \left[e^{R^* r_{\tau}^u(\theta) + \kappa_{\theta}(R^*) \cdot \tau} \right] \cdot e^{-R^* u} = \mathbb{E}_{Q_{\theta}^{R*}} \left[e^{r_{\tau}^u(\theta)} \right] \cdot e^{-u} \leqslant e^{-u}$$

and

$$\psi(u) = \mathbb{E}_{Q^{R^*}}\left[\frac{e^{R^* \cdot R^u_\tau + \kappa_\Theta(R^*) \cdot \tau}}{\xi(\Theta)}\right] \cdot e^{-R^* \cdot u} = \mathbb{E}_{Q^{R^*}}\left[\frac{e^{R^u_\tau}}{\xi(\Theta)}\right] \cdot e^{-u} \leqslant e^{-u}$$

where the inequalities follow by Theorem 4.1.

References

- Asmussen, S. (1994) Busy period analysis, rare events and transient behaviour in fluid models, J. Appl. Math. Stochastic Anal. 7(3), 269–299.
- [2] Asmussen, S. (1995) Stationary distribution for fluid models with or without Brownian noise, Stochastic Models 11, 21–49.
- [3] Asmussen, S. and Albrecher, H. (2010) *Ruin Probabilities*, 2nd edn. World Scientific Publishing, London.
- Boogaert, P. and De Waegenaere, A. (1990) Simulation of ruin probabilities. Insur. Math. Econ., 9, 95–99.
- [5] Cohn, D.L. (2013) Measure Theory, 2nd ed. Birkhäuser Advanced Texts.
- [6] Dassios, A. and Embrechts, P. (1989) Martingales and insurance risk, Stochastic Models, 5, 181-217.
- [7] Delbaen, F. and Haezendock, J. (1989) A martingale approach to premium calculation principles in an arbitrage free market, *Insur. Math. Econ.*, 8, 269-277.
- [8] Faden, A. M. (1985) The existence of regular conditional probabilities: Necessary and sufficient conditions, Ann. Probab., 13, 288–298.
- [9] Gut, A. (2009) Stopped Random Walks: Limit Theorems and Applications, 2nd ed. Springer-Verlag New York.
- [10] Lyberopoulos, D. P. and Macheras, N. D. (2012) Some characterizations of mixed Poisson processes, Sankhyā A, 74, 57–79.
- [11] Lyberopoulos, D. P. and Macheras, N. D. (2014) Some characterizations of mixed renewal processes, arXiv: 1205.4441.
- [12] Lyberopoulos, D.P. and Macheras, N.D. (2019) A characterization of martingale-equivalent compound mixed Poisson processes, arXiv:1905.07629.
- [13] Lyberopoulos, D.P., Macheras, N.D. and Tzaninis S. M. (2019) On the equivalence of various definitions of mixed Poisson processes, *Math. Slovaca*, 69, 453-468.
- [14] Macheras, N. D. and Tzaninis, S. M. (2018) Some characterizations for Markov processes as mixed renewal processes, *Math. Slovaca*, 68, 1477-1494.
- [15] Macheras, N. D. and Tzaninis, S. M. (2020) A characterization of equivalent martingale measures in a renewal risk model with applications to premium calculation principles, *Mod. Stoch.: Theory Appl.*, 7, 43–60.
- [16] Palmowski, Z. and Rolski, T. (1996) A note on martingale inequalities for fluid models, Statist. Appl. Probab. Lett., 31, 13–21.
- [17] Palmowski, Z. and Rolski, T. (1998) Superposition of alternating on -off flows and a fluid model, Ann. Appl. Probab., 8, 524–541.
- [18] Ridder, A. (1996) Fast simulation of Markov fluid models, J. Appl. Probab., 33, 786–804.
- [19] Rolski, T., Schmidli, H., Schmidt, V. and Teugels, J. L. (1999) Stochastic Processes for Insurance and Finance. Wiley, Chichester.
- [20] Schmidli, H. (1996) Lundberg inequalities for a Cox model with a piecewise constant intensity, J. Appl. Probab., 33, 196–210.
- [21] Schmidli, H. (1997) An extension to the renewal theorem and an application to risk theory, Ann. Appl. Probab., 7, 121–133.
- [22] Schmidli, H. (2017) Risk Theory. Springer-Verlag.

- [23] Schmidt, K. D. (1996) Lectures on Risk Theory. B.G. Teubner, Stuttgart.
- [24] Tzaninis, S. M. and Macheras, N. D. (2020) A characterization of progressively equivalent probability measures preserving the structure of a compound mixed renewal process, arXiv:2007.05289.

DEPARTMENT OF STATISTICS AND INSURANCE SCIENCE, UNIVERSITY OF PIRAEUS, 80 KARAOLI AND DIMITRIOU STR., 185 34 PIRAEUS, GREECE

E-mail address: stzaninis@unipi.gr