LONG-TIME BEHAVIORS OF MEAN-FIELD INTERACTING PARTICLE SYSTEMS RELATED TO MCKEAN-VLASOV EQUATIONS

WEI LIU, LIMING WU, AND CHAOEN ZHANG

ABSTRACT. In this paper, we investigate gradient estimate of the Poisson equation and the exponential convergence in the Wasserstein metric $W_{1,d_{l^1}}$, uniform in the number of particles, and uniform-in-time propagation of chaos for the mean-field weakly interacting particle system related to McKean-Vlasov equation. By means of the known approximate componentwise reflection coupling and with the help of some new cost function, we obtain explicit estimates for those three problems, avoiding the technical conditions in the known results. Our results apply when the confinement potential V has many wells, the interaction potential W has bounded second mixed derivative $\nabla_{xy}^2 W$ which should be not too big so that there is no phase transition. As an application, we obtain the concentration inequality of the mean-field interacting particle system with explicit and sharp constants, uniform in time. Several examples are provided to illustrate the results.

AMS 2010 Subject classifications. 82C22, 60J60, 60E15, 65C35, 65C05.

Key words and Phrases. McKean-Vlasov equation, interacting particle system, propagation of chaos, exponential convergence, concentration inequality.

1. INTRODUCTION

In this paper, we consider the following nonlinear McKean-Vlasov equation with initial condition u_0

$$\partial_t u_t = \nabla \cdot [\nabla u_t + u_t \nabla V + u_t (\nabla_x W \circledast u_t)], \tag{1.1}$$

where the unknown u_t is a time dependent probability density on \mathbb{R}^d $(d \ge 1), V : \mathbb{R}^d \to \mathbb{R}$ is a confinement potential and $W : \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$ is an interaction potential. Here ∇ and $\nabla \cdot$ (applied to a vector field) denote the gradient operator and the divergence operator respectively, while $\nabla_x W$ stands for the gradient of W with respect to (w.r.t. in short) the first variable, and

$$\nabla_x W \circledast u_t(x) := \int_{\mathbb{R}^d} \nabla_x W(x, y) u_t(y) dy.$$

Date: July 21, 2020.

The first author is supported by NSFC (11731009), the Fundamental Research Funds for the Central Universities 2042020kf0217 and 2042020kf0031, and CSC.

When $W(x, y) = W_0(x - y)$ for some even potential W_0 , $\nabla_x W \circledast u = \nabla W_0 \ast u$ (the usual convolution).

The probabilistic equivalent version of (1.1) is the following self-interacting stochastic differential equation (SDE in short):

$$\begin{cases} dX_t = \sqrt{2}dB_t - \nabla V(X_t)dt - \nabla_x W \circledast \mu_t(X_t)dt, \\ X_0 \stackrel{law}{=} u_0(x)dx, \end{cases}$$
(1.2)

where μ_t is the law of X_t . The density u_t of the law μ_t of X_t at time t is the solution of the McKean-Vlasov equation (1.1) and vice versa. The existence and uniqueness of the solution of the SDE (1.2) and the McKean-Vlasov equation (1.1) have been extensively studied. The reader is referred to [28, 17, 27, 32] and recent works [31, 7, 19] as well as the references therein. For the convergence to equilibrium of solution μ_t as $t \to +\infty$, it is worth mentioning that Carrillo, McCann and Villani [8] obtained the explicit exponential convergence in entropy under various kinds of convexity conditions on the potentials V and W, via their enlightening idea of interpreting the McKean-Vlasov equation as the gradient descent flow of the free energy on the space of probability measures equipped with the L^2 -Wasserstein metric. Eberle et al. [15] got the quantitative bounds on the exponential convergence in some appropriate transport cost to equilibrium for McKean-Vlasov equations by using Lyapunov condition and reflection coupling. Eberle [14] showed the exponential contractivity for diffusion semigroups w.r.t. Kantorovich distance by using componentwise reflection coupling methods and choosing appropriate distance functions. The reader is referred also to Luo and Wang [23] for the exponential convergence of diffusion semigroups w.r.t. the L^p -Wasserstein distance for all p > 1.

The McKean-Vlasov equation (1.1) or (1.2) is the idealization of the following interacting particle system of mean-field type when the number N of particles goes to infinity:

$$\begin{cases} dX_t^{i,N} = \sqrt{2} dB_t^i - \nabla V(X_t^{i,N}) dt - \frac{1}{N-1} \sum_{j: j \neq i, 1 \le j \le N} \nabla_x W(X_t^{i,N}, X_t^{j,N}) dt, \\ X_0^{i,N} = X_0^i, \ i = 1, \cdots, N, \end{cases}$$
(1.3)

where the initial values X_0^1, \dots, X_0^N are i.i.d. random variables with common law $\mu_0(dx) = u_0(x)dx$, and $B_t^1 \dots, B_t^N$ are N independent Brownian motions taking values in \mathbb{R}^d , independent of $X_0^i, 1 \leq i \leq N$. In fact this is the goal of the so-called propagation of chaos: when the number N of particles goes to infinity, the empirical measures $\frac{1}{N} \sum_{i=1}^N \delta_{X_t^{i,N}}$ of the particle system (1.3) (or the law of a single particle) converge weakly to the solution μ_t of the self-interacting diffusion (1.2).

The propagation of chaos for the mean-field interacting particle systems has been widely studied during the last forty years. The early studies were concentrated on the propagation of chaos in bounded time intervals, see [24, 32, 27] and the references therein. The study on the propagation of chaos in the whole time interval \mathbb{R}^+ is much more difficult and recent. When the confinement potential V is strictly convex and the interaction potential $W(x, y) = W_0(x - y)$ with W_0 strictly convex, Malrieu [25] showed the uniform in time propagation of chaos by applying the logarithmic Sobolev inequality. In the case that there is no confinement (i.e. $V \equiv 0$) and the interaction potential W_0 is strictly convex, Benachour et al. [1, 2] proved propagation of chaos (but not uniform in time) and polynomial convergence to equilibrium; Malrieu [26] obtained the uniform in time propagation of chaos and exponential convergence to equilibrium for the particle system viewed from the center, by using functional inequalities. When W_0 is degenerately convex and V = 0, Cattiaux et al. [9] showed the uniform in time propagation of chaos and exponential convergence to equilibrium by using synchronous coupling.

Without the convexity of V and W_0 , recently Durmus, Eberle, Guillin and Zimmer [13] use the componentwise reflection coupling introduced in [14] to prove the exponential convergence in some Wasserstein metric and uniform in time propagation of chaos for weakly interacting mean-field particle system. For more results about propagation of chaos, we refer the reader to [12, 20, 21, 22, 29, 30] and the references therein.

The main purpose of this paper is to investigate the exponential convergence in L^1 -Wasserstein metric in the purpose of refining the previous results in [14, 13], the concentration inequalities and the propagation of chaos of the mean-field weakly interacting particle system. Although we use the same approximate componentwise reflection coupling ([14]), our next approach will be quite different from [14, 13]:

- (1) our starting point is some explicit gradient estimate of the Poisson equation, which implies moreover the concentration inequalities of the empirical mean of the interacting particle system, useful for numerical computation of solution μ_t of the McKean-Vlasov equation;
- (2) we will choose a different metric from that in [14, 13], which allows us to obtain some explicit and almost sharp estimate of the exponential rate in the convergence of the interacting particles system to its equilibrium in the W_1 -metric, uniform in the number N of the particles.
- (3) As a by-product, we obtain some explicit estimate on the propagation of chaos, uniform in time.

The paper is organized as follows. In the next section, we will present our framework and main results. The proofs are provided in Section 3 and section 4. The applications to concentration inequalities are given in the last section.

2. Main resutls

2.1. Framework: notations and conditions.

2.1.1. Conditions on the dissipativity rate of a single particle. First we introduce the dissipative rate $b_0(r)$ of the drift of one single particle in (1.3) at distance r > 0,

$$\langle x - y, -[\nabla V(x) - \nabla V(y)] - [\nabla_x W(x, z) - \nabla_x W(y, z)] \rangle \le b_0(r)|x - y|$$
(2.1)

holds for any $x, y, z \in \mathbb{R}^d$ with |x - y| = r. Throughout this paper we assume that $b_0(r)$ is a continuous function on $(0, +\infty)$ satisfying

$$\limsup_{r \to +\infty} \frac{b_0(r)}{r} < 0, \tag{2.2}$$

i.e. the drift of one particle is dissipative at infinity.

We also assume that

$$\lim_{r \to 0+} b_0^+(r) = 0. \tag{2.3}$$

Next we introduce an important reference function h which enables us to obtain some new results, avoiding the technical parameters in [14, 13]. For any function $f \in C^2(0, +\infty)$ and r > 0, let \mathcal{L}_{ref} be the generator defined by

$$\mathcal{L}_{ref}f(r) := 4f''(r) + b_0(r)f'(r).$$
(2.4)

Let $h: \mathbb{R}^+ \to \mathbb{R}^+$ be the function determined by: h(0) = 0 and

$$h'(r) = \frac{1}{4} \exp\left(-\frac{1}{4} \int_0^r b_0(s) ds\right) \int_r^{+\infty} s \cdot \exp\left(\frac{1}{4} \int_0^s b_0(u) du\right) ds.$$
(2.5)

h is a well defined C^2 function by the dissipative condition (2.2), and it is a solution of the one-dimensional Poisson equation

$$\mathcal{L}_{ref}h(r) = 4h''(r) + b_0(r)h'(r) = -r, \ r > 0$$
(2.6)

with h(0) = 0. This function was used by the second named author [33] for functional and isoperimetric inequalities on Riemmanian manifolds.

2.1.2. Kantorovich-Wasserstein W_1 -metric. For the configuration space $(\mathbb{R}^d)^N$, instead of the usual Euclidean metric, we will use the l^1 -metric (generalized Hamming metric)

$$d_{l^1}(x,y) = \sum_{i=1}^N |x^i - y^i|, \ x = (x^1, \cdots, x^N), \ y = (y^1, \cdots, y^N) \in (\mathbb{R}^d)^N.$$

We consider the Kantorovich-Wasserstein distance w.r.t. d_{l^1} metric on $(\mathbb{R}^d)^N$, i.e., for any two probability measures μ and ν on $(\mathbb{R}^d)^N$,

$$W_{1,d_{l^{1}}}(\mu,\nu) = \inf_{P \in \Pi(\mu,\nu)} \iint_{(\mathbb{R}^{d})^{N} \times (\mathbb{R}^{d})^{N}} d_{l^{1}}(x,y) P(dx,dy)$$

where $\Pi(\mu, \nu)$ is the set of all couplings of μ, ν , i.e. the set of all probability measures on $(\mathbb{R}^d)^N \times (\mathbb{R}^d)^N$ whose marginal distributions of x and y are respectively μ and ν .

Notice that for a C^1 -function g on $(\mathbb{R}^d)^N$, its Lipschitzian norm $||g||_{Lip(d_{l^1})}$ w.r.t. d_{l^1} coincides with $\max_{1 \le i \le N} ||\nabla_i g||_{\infty}$ where ∇_i is the gradient w.r.t. x_i . By Kantorovich-Rubinstein duality relation,

$$W_{d_{l^1}}(\mu,\nu) = \sup_{g \in C_b^1((\mathbb{R}^d)^N): \max_{1 \le i \le N} \|\nabla_i g\|_{\infty} \le 1} \left(\int g d\mu - \int g d\nu \right)$$

When N = 1, we write simply W_1 for W_{1,d_n} .

We notice that for two probability measures μ, ν on $(\mathbb{R}^d)^N$,

$$\sum_{i=1}^{N} W_1(\mu^i, \nu^i) \le W_{d_{l^1}}(\mu, \nu)$$
(2.7)

and the equality holds when $\mu = \bigotimes_{i=1}^{N} \mu^{i}$, $\nu = \bigotimes_{i=1}^{N} \nu^{i}$ are product measures, where μ^{i} (resp. ν^{i}) is the marginal distribution of x_{i} of μ (resp. ν). In fact if $X = (X^{1}, \dots, X^{N}), Y = (Y^{1}, \dots, Y^{N})$ are two random vectors such that the law of (X, Y) is

an optimal coupling of (μ, ν) in $W_{1,d_{l^1}}$, then for each *i*, the law of (X^i, Y^i) is a coupling of (μ^i, ν^i) , so

$$W_{d_{l^1}}(\mu,\nu) = \mathbb{E}d_{l^1}(X,Y) = \sum_{i=1}^N \mathbb{E}|X^i - Y^i| \ge \sum_{i=1}^N W_1(\mu^i,\nu^i).$$

When μ , ν are product measures, let (X^i, Y^i) (or its joint law) be an optimal coupling of (μ^i, ν^i) for $W_1(\mu^i, \nu^i)$ so that $(X^1, Y^1), \dots, (X^N, Y^N)$ are independent. Then $(X = (X^i)_{1 \le i \le N}, Y = (Y^i)_{1 \le i \le N})$ is a coupling of (μ, ν) , so we get

$$\sum_{i=1}^{N} W_1(\mu^i, \nu^i) = \sum_{i=1}^{N} \mathbb{E}|X^i - Y^i| = \mathbb{E}d_{l^1}(X, Y) \ge W_{d_{l^1}}(\mu, \nu)$$

i.e. the equality in (2.7) holds in the product measures case. (This is well known.)

2.2. An explicit gradient estimate of the Poisson equation and its applications in concentration inequalities. Let $\{P_t^{(N)}\}_{t\geq 0}$ be the transition semigroup of the mean-field interacting particle system (1.3), whose generator is given by

$$\mathcal{L}^{(N)}f(x^1,\cdots,x^N) = \sum_{i=1}^N \left(\Delta_i f - \nabla V(x^i) \cdot \nabla_i f - \frac{1}{N-1} \sum_{j \neq i} \nabla_x W(x^i,x^j) \cdot \nabla_i f \right).$$

Its unique invariant probability measure is the mean-field Gibbs measure, given by

$$\mu^{(N)}(dx^{1}, \cdots, dx^{N}) = \frac{1}{C_{N}} \exp\left(-\sum_{i=1}^{N} V(x^{i}) - \frac{1}{N-1} \sum_{1 \le i < j \le N} W(x^{i}, x^{j})\right) dx^{1} \cdots dx^{N},$$

where C_N is the normalization constant.

We introduce the following key assumption on the interaction potential:

(**H**): $\|\nabla_{xy}^2 W\|_{\infty} \|h'\|_{\infty} < 1$

where h is given by (2.5), $\|h'\|_{\infty} := \sup_{r \ge 0} h'(r)$, and $\nabla_{xy}^2 W = (\frac{\partial^2}{\partial x_i \partial y_j} W)_{1 \le i,j \le d}$,

$$\|\nabla_{xy}^2 W\|_{\infty} := \sup_{x,y \in \mathbb{R}^d} \sup_{z \in \mathbb{R}^d, |z|=1} |\nabla_{xy}^2 W(x,y)z|.$$

Notice that when the dissipativity at infinity condition (2.2) is satisfied, $b_0(r)$ can be taken as $-c_1r + c_2$ (with $c_1, c_2 > 0$), so $||h'||_{\infty} := \sup_{r>0} h'(r) < +\infty$.

This assumption is a translation of Dobrushin-Zegarlinski's uniqueness condition in the framework of mean field, and it implies that the mean field has no phase of transition (see [18]).

Notice that under the assumption (**H**) and (2.2), both the equations (1.2) and (1.3) have unique strong solutions. On the space of continuous paths $C([0, T], (\mathbb{R}^d)^N)$ where $T \in (0, +\infty]$, we consider the L^1 -metric

$$d_{L^{1}[0,T]}(\gamma_{1},\gamma_{2}) := \int_{0}^{T} d_{l^{1}}(\gamma_{1}(t),\gamma_{2}(t))dt.$$
(2.8)

Given the starting point $x \in (\mathbb{R}^d)^N$, let \mathbb{P}_x be the law of $X^{(N)} = (X_t^{(N)})_{t \ge 0}$ with $X_0^{(N)} = x$.

Theorem 2.1. Assume (2.2), (2.3) and (**H**). For any $x_0 = (x_0^1, \dots, x_0^N) \in (\mathbb{R}^d)^N$ and $y_0 = (y_0^1, \dots, y_0^N) \in (\mathbb{R}^d)^N$, we have

$$\int_{0}^{+\infty} W_{d_{l^{1}}}(P_{t}^{(N)}(x_{0},\cdot),P_{t}^{(N)}(y_{0},\cdot))dt \leq W_{1,d_{L^{1}[0,\infty]}}(\mathbb{P}_{x_{0}},\mathbb{P}_{y_{0}})$$

$$\leq \frac{1}{1-\|\nabla_{xy}^{2}W\|_{\infty}\|h'\|_{\infty}}\sum_{i=1}^{N}h(|x_{0}^{i}-y_{0}^{i}|).$$
(2.9)

In particular for any $g \in C_b^1((\mathbb{R}^d)^N)$ with $\mu^{(N)}(g) = 0$, the solution G of the Poisson equation $-\mathcal{L}^{(N)}G = g$ with $\mu(G) = 0$ satisfies

$$\|\nabla_i G\|_{\infty} \le c_{Lip} \cdot \max_{1 \le j \le N} \|\nabla_j g\|_{\infty}, 1 \le i \le N,$$
(2.10)

where

$$c_{Lip} := \frac{h'(0)}{1 - \|\nabla_{xy}^2 W\|_{\infty} \|h'\|_{\infty}}$$
(2.11)

and

$$h'(0) = \frac{1}{4} \int_0^{+\infty} s \cdot \exp\left(\frac{1}{4} \int_0^s b_0(u) du\right) ds.$$

By the theorem above we can immediately obtain the following result about the nonlinear McKean-Vlasov equation (1.1).

Corollary 2.2. Under the same assumptions as in Theorem 2.1, for any two solutions μ_t, ν_t of the self-interacting diffusion (1.2) with the initial distributions μ_0, ν_0 with finite second moment respectively, we have

$$\int_{0}^{\infty} W_{1}(\mu_{t},\nu_{t})dt \leq \frac{\|h'\|_{\infty}}{1-\|\nabla_{xy}^{2}W\|_{\infty}\|h'\|_{\infty}}W_{1}(\mu_{0},\nu_{0}).$$
(2.12)

Proof. By (2.9) in Theorem 2.1 and the fact that

$$h(r) \le h(0) + ||h'||_{\infty} \cdot r = ||h'||_{\infty} \cdot r, \ \forall r \ge 0$$

we have

$$\int_{0}^{\infty} W_{1,d_{l^{1}}}(\mu_{0}^{\otimes N}P_{t}^{(N)},\nu_{0}^{\otimes N}P_{t}^{(N)})dt \leq \frac{\|h'\|_{\infty}}{1-\|\nabla_{xy}^{2}W\|_{\infty}\|h'\|_{\infty}}W_{1,d_{l^{1}}}(\mu_{0}^{\otimes N},\nu_{0}^{\otimes N}).$$
(2.13)

Notice that $\mu_t^{(N)} := \mu_0^{\otimes N} P_t^{(N)}$ and $\nu_t^{(N)} := \nu_0^{\otimes N} P_t^{(N)}$ are symmetric probability measures on $(\mathbb{R}^d)^N$ and their marginal distributions $\mu_t^{(i,N)}$, $\nu_t^{(i,N)}$ of x_i converge weakly to μ_t, ν_t (respectively) by the finite time propagation of chaos. By using (2.7) we have

$$NW_1(\mu_t^{(1,N)},\nu_t^{(1,N)}) = \sum_{i=1}^N W_1(\mu_t^{(i,N)},\nu_t^{(i,N)}) \le W_{1,d_{l^1}}(\mu_t^{(N)},\nu_t^{(N)})$$

and then by the lower semi-continuity of W_1 in the weak convergence topology,

$$W_1(\mu_t, \nu_t) \le \liminf_{N \to +\infty} W_1(\mu_t^{(1,N)}, \nu_t^{(1,N)}) \le \liminf_{N \to +\infty} \frac{1}{N} W_{1,d_{l^1}}(\mu_t^{(N)}, \nu_t^{(N)}).$$
(2.14)

Combining (2.13) and (2.14) together, we obtain by Fatou's lemma,

$$\int_{0}^{\infty} W_{1}(\mu_{t},\nu_{t})dt \leq \liminf_{N \to +\infty} \frac{1}{N} \int_{0}^{\infty} W_{1,d_{l}1}(\mu_{t}^{(N)},\nu_{t}^{(N)})dt$$
$$\leq \frac{\|h'\|_{\infty}}{1-\|\nabla_{xy}^{2}W\|_{\infty}\|h'\|_{\infty}} \liminf_{N \to +\infty} \frac{1}{N} W_{1,d_{l}1}(\mu_{0}^{\otimes N},\nu_{0}^{\otimes N})$$
$$= \frac{\|h'\|_{\infty}}{1-\|\nabla_{xy}^{2}W\|_{\infty}\|h'\|_{\infty}} W_{1}(\mu_{0},\nu_{0})$$

where the last equality follows by (2.7). That completes the proof.

As an application of Theorem 2.1 to the concentration inequality, we have the following result about the Gaussian concentration of the U-statistics, which is a straightforward application of a general result in Proposition 5.3. The proofs are given in the last section §5.

For any $1 \leq m \leq N$, let $f_m : (\mathbb{R}^d)^m \to \mathbb{R}$ be a measurable and symmetric function. The U-statistic of order m with kernel f_m is defined by

$$U_N(f_m)(x^1, \cdots, x^N) = \frac{1}{|I_N^m|} \sum_{(i_1, \cdots, i_m) \in I_N^m} f_m(x^{i_1}, \cdots, x^{i_m}), \ \forall (x^1, \cdots, x^N) \in (\mathbb{R}^d)^N,$$
(2.15)

where

$$I_N^m := \{ (i_1, \cdots, i_m) \in \mathbb{N}^k | i_1, \cdots, i_m \text{ are different }, 1 \le i_1, \cdots, i_m \le N \}$$
(2.16)

and $|I_N^m|$ denotes the number of elements in I_N^m (equal to N!/(N-m)!).

Next we introduce the following *Gaussian integrability* assumption of the initial distribution μ_0 :

$$\int_{\mathbb{R}^d} e^{\lambda_0 |x|^2} \mu_0(dx) < +\infty, \text{ for some } \lambda_0 > 0$$
(2.17)

which is equivalent to say that there is some Gaussian concentration constant $c_G(\mu_0) > 0$ such that

$$\int_{\mathbb{R}} e^{f(x) - \mu_0(f)} d\mu_0(x) \le \exp\left(\frac{c_G(\mu_0)}{2} \|f\|_{Lip}^2\right)$$
(2.18)

for all Lipschitzian functions f on \mathbb{R}^d (w.r.t. the usual Euclidean distance).

Remark 2.3. The equivalence between the *Gaussian integrability* (2.17) and the *Gaussian concentration inequality* (2.18) was established by H. Djellout, A. Guillin and the second named author [10], and (2.18) is the famous characterization of Bobkov-Götze [3] of the transport-entropy inequality. By the tensorization of the transport-entropy inequality for product measure, (2.18) implies that for any $N \ge 1$,

$$\int_{(\mathbb{R}^d)^N} e^{g(x) - \mu_0^{\otimes N}(g)} d\mu_0^{\otimes N}(x) \le \exp\left(\frac{N}{2} c_G(\mu_0) \|g\|_{Lip(d_{l^1})}^2\right)$$
(2.19)

for all Lipschitzian functions g on $(\mathbb{R}^d)^N$.

Corollary 2.4. Assume the conditions in Theorem 2.1 and the Gaussian integrability (2.17) of the initial distribution μ_0 . Let $f_m \in C^2((\mathbb{R}^d)^m, \mathbb{R})$ be measurable, symmetric, and 1-Lipschitz w.r.t. the d_{l^1} -metric on $(\mathbb{R}^d)^m$, i.e. $\max_i \|\nabla_i f\|_{\infty} \leq 1$. Then for any $\lambda, T > 0$, we have

$$\mathbb{E}\exp\left(\frac{\lambda}{T}\left[\int_{0}^{T}U_{N}(f_{m})(X_{t}^{1,N},\cdots,X_{t}^{N,N})dt-\int_{0}^{T}\mathbb{E}f_{m}(X_{t}^{1,N},\cdots,X_{t}^{m,N})dt\right]\right) \\
\leq \exp\left(\frac{m^{2}\lambda^{2}c_{Lip}^{2}}{2NT}\left(1+\frac{c_{G}(\mu_{0})}{T}\right)\right),$$
(2.20)

where c_{Lip} is the same as given in (2.11). In particular we have for any $\delta > 0$

$$\mathbb{P}\left\{\frac{1}{T}\int_{0}^{T}U_{N}(f_{m})(X_{t}^{1,N},\cdots,X_{t}^{N,N})dt - \frac{1}{T}\int_{0}^{T}\mathbb{E}f_{m}(X_{t}^{1,N},\cdots,X_{t}^{m,N})dt > \delta\right\} \\
\leq \exp\left(-\frac{(1-\|\nabla_{xy}^{2}W\|_{\infty}\|h'\|_{\infty})^{2}}{2m^{2}(h'(0))^{2}(1+c_{G}(\mu_{0})/T)}NT\delta^{2}\right).$$
(2.21)

The concentration inequality (2.21) is sharp when V is quadratic and W = 0, see Example 2.12 for explicit expression of all involved constants in the Gaussian case.

2.3. Exponential convergence of the particle system in the W_{1,d_1} -metric.

Theorem 2.5. Assume (2.2) and (**H**). Suppose that there exists a constant $M \in \mathbb{R}$ such that

$$b_0(r) \le rM, \forall r > 0 \tag{2.22}$$

(this condition is stronger than (2.3)), then for any $\varepsilon > 0$ such that

$$K_{\varepsilon} := \frac{1 - \|\nabla_{xy}^2 W\|_{\infty} \|h'\|_{\infty} - \varepsilon (M + \|\nabla_{xy}^2 W\|_{\infty})}{\|h'\|_{\infty} + \varepsilon} > 0, \qquad (2.23)$$

we have for any $x_0, y_0 \in (\mathbb{R}^d)^N$

$$W_{d_{l^1}}(P_t^{(N)}(x_0, \cdot), P_t^{(N)}(y_0, \cdot)) \le A_{\varepsilon} e^{-K_{\varepsilon} t} d_{l^1}(x_0, y_0), \ \forall t \ge 0,$$
(2.24)

where

$$A_{\varepsilon} = \sup_{r>0} \frac{r}{h(r) + \varepsilon r} \cdot \sup_{r>0} \frac{h(r) + \varepsilon r}{r}.$$
(2.25)

Remark 2.6. An easy estimate of A_{ε} is $A_{\varepsilon} \leq \frac{\sup_{r\geq 0} h'(r) + \varepsilon}{\inf_{r\geq 0} h'(r) + \varepsilon}$ (since h(0) = 0). Note that when $M + \|\nabla_{xy}^2 W\|_{\infty} > 0$, the exponential rate K_{ε} increases (then better and better) as ε decreases to 0, but A_{ε} may explode once if $\inf_{r>0} h'(r) = 0$.

Remark 2.7. Notice that (2.22) is equivalent to say that

$$\nabla^2 V(x) + \nabla^2_{xx} W(x, y) \ge -MI, \ x, y \in \mathbb{R}^d.$$

When $\kappa := -M - \|\nabla_{xy}^2 W\|_{\infty} > 0$, we see that the Hessian of the Hamiltonian

$$H(x^{1}, \cdots, x^{N}) = \sum_{i=1}^{N} V(x^{i}) + \frac{1}{N-1} \sum_{1 \le i < j \le N} W(x^{i}, x^{j})$$

is bounded from below by κI (this estimate of the lower bound of the Bakry-Emery curvature is sharp if $\nabla_{xy}^2 W$ is constant and definitely nonnegative). Notice that when M < 0, we can take $b_0(r) = Mr$, so h'(r) = -1/M. Then $\kappa > 0$ if and only if **(H)** is satisfied. The advantage of our condition **(H)** (w.r.t. the positive curvature condition) is: it does not depend on the curvature but on the dissipativity, it holds even if V has many wells once if the interaction is weak enough.

In the positive curvature $\kappa>0$ case we have by Bakry-Emery's curvature characterization

$$W_1(P_t^{(N)}(x,\cdot), P_t^{(N)}(y,\cdot)) \le e^{-\kappa t} |x-y|$$

in the Euclidean metric on $(\mathbb{R}^d)^N$. On the other hand as above $b_0(r) = Mr, h(r) = -r/M$, we see that $K_{\varepsilon} \to -M - \|\nabla^2_{xy}W\|_{\infty} = \kappa$ as $\varepsilon \to +\infty$, and $A_{\varepsilon} \equiv 1$, so (2.24) yields

 $W_{1,d_{l^1}}(P_t^{(N)}(x,\cdot), P_t^{(N)}(y,\cdot)) \le e^{-\kappa t} d_{l^1}(x,y),$ (2.26)

a new but not at all surprising observation.

Theorem 2.5 above will give us an explicit exponential convergence in W_1 of the nonlinear McKean-Vlasov equation (1.1). For the exponential convergence in entropy of the nonlinear McKean-Vlasov equation (1.1) under the condition (**H**), see Guillin *et al.* [18].

Corollary 2.8. Under the same assumptions as in Theorem 2.5, for any $\varepsilon > 0$ so that $K_{\varepsilon} > 0$ (i.e. (2.23)), we have for the solutions μ_t, ν_t of the self-interacting diffusion (1.2) with the initial distributions μ_0, ν_0 which have finite second moments respectively,

$$W_1(\mu_t, \nu_t) \le A_{\varepsilon} e^{-K_{\varepsilon} t} W_1(\mu_0, \nu_0), \ \forall t \ge 0,$$
 (2.27)

where K_{ε} and A_{ε} are given by (2.23) and (2.25) respectively.

Proof. The proof of this corollary is similar to that of Corollary 2.2, and we utilize the same notations as in the Corollary 2.2. First by Theorem 2.5, we have for any $t \ge 0$

$$W_{1,d_{l^{1}}}(\mu_{0}^{\otimes N}P_{t}^{(N)},\nu_{0}^{\otimes N}P_{t}^{(N)}) \leq A_{\varepsilon}e^{-K_{\varepsilon}t}W_{1,d_{l^{1}}}(\mu_{0}^{\otimes N},\nu_{0}^{\otimes N}).$$

Combining the inequality above with (2.14), we obtain

$$W_1(\mu_t, \nu_t) \le A_{\varepsilon} e^{-K_{\varepsilon}t} \liminf_{N \to +\infty} \frac{1}{N} W_{1, d_{l^1}}(\mu_0^{\otimes N}, \nu_0^{\otimes N})$$
$$= A_{\varepsilon} e^{-K_{\varepsilon}t} W_1(\mu_0, \nu_0)$$

the desired result.

2.4. **Propagation of chaos in large time.** We have the following uniform in time propagation of chaos.

Theorem 2.9. Assume (2.2), (2.22) and (**H**). Suppose that there exist some positive constants c_1, c_2, c_3 such that

$$\langle x, \nabla V(x) \rangle \ge c_1 |x|^2 - c_2, \forall x \in \mathbb{R}^d$$
 (2.28)

and

$$\langle z, \nabla_{xx}^2 W(x, y) z \rangle \ge -c_3 |z|^2, \forall x, y, z \in \mathbb{R}^d.$$
 (2.29)

Assume

$$c_1 - c_3 - \|\nabla_{xy}^2 W\|_{\infty} > 0.$$
(2.30)

Then for any $\varepsilon > 0$ such that $K_{\varepsilon} > 0$, and $\tilde{\varepsilon} \in (0, c_1 - c_3 - \|\nabla_{xy}^2 W\|_{\infty})$, the following estimates of propagation of chaos hold for the mean-field interacting particle system (1.3) with any initial probability measure μ_0 having finite second moment:

(a) (path-type propagation of chaos) for any $T > 0, 1 \le k \le N$, denote $\mathbb{P}_{\nu}(\cdot) = \int_{(\mathbb{R}^d)^N} \mathbb{P}_x(\cdot) d\nu(x)$ the law of $(X_t^{(N)})_{t\ge 0}$ with the initial distribution ν , $\mathbb{P}_{\nu}^{[1,k],N}|_{[0,T]}$ the joint law of paths of the k particles $((X_t^{i,N})_{t\in[0,T]}, 1\le i\le k)$ in time interval [0,T], and \mathbb{Q}_{μ_0} the law of the self-interacting diffusion $(X_t)_{t\ge 0}$ with the initial distribution μ_0 . We have

$$\frac{1}{k}W_{1,d_{L^{1}[0,T]}}(\mathbb{P}_{\mu_{0}^{\otimes N}}^{[1,k],N}|_{[0,T]},\mathbb{Q}_{\mu_{0}}^{\otimes k}|_{[0,T]}) \leq \frac{T}{\sqrt{N-1}} \frac{\|\nabla_{xy}^{2}W\|_{\infty}\|h'\|_{\infty}}{1-\|\nabla_{xy}^{2}W\|_{\infty}\|h'\|_{\infty}} \cdot \max\{m_{2}(\mu_{0}),\hat{c}(\varepsilon)\}$$
(2.31)

where

$$m_{2}(\mu_{0}) = \left(\int_{\mathbb{R}^{d}} |x|^{2} d\mu_{0}(x)\right)^{\frac{1}{2}},$$

$$\hat{c}(\varepsilon) = \left(\frac{d + c_{2} + \frac{1}{4\varepsilon} |\nabla_{x} W(0, 0)|^{2}}{c_{1} - c_{3} - \|\nabla_{xy}^{2} W\|_{\infty} - \tilde{\varepsilon}}\right)^{\frac{1}{2}}.$$
(2.32)

(b) (Uniform in time propagation of chaos) for all time t > 0 and any $1 \le k \le N$:

$$W_{1,d_{l^{1}}}(\mu_{t}^{[1,k],N},\mu_{t}^{\otimes k}) \leq \frac{k}{\sqrt{N-1}} \frac{A_{\varepsilon}}{K_{\varepsilon}} \|\nabla_{xy}^{2}W\|_{\infty} \max\{m_{2}(\mu_{0}), \hat{c}(\varepsilon)\}$$
(2.33)

where $\mu_t = u_t dx$ is the solution of the McKean-Vlasov equation (1.1), and $\mu_t^{[1,k],N}$ is the joint law of the k particles $(X_t^{i,N}, 1 \leq i \leq k)$ in the mean-field system (1.3) of interacting particles $(X_t^{i,N})_{1\leq i\leq N}$ with $X_0^{i,N}, 1\leq i\leq N$ i.i.d. of law μ_0 (independent of $(B_t^{i,N})_{1\leq i\leq N,t\geq 0}$), and the constants K_{ε} , A_{ε} , $m_2(\mu_0)$ and $\hat{c}(\varepsilon)$ are given in (2.23), (2.25) and (2.32) respectively.

Remark 2.10. The time-uniform propagation of chaos is much more difficult than the bounded time propagation of chaos, accomplished in the 80-90's of the last century. The physical reason is that the time-uniform propagation of chaos fails in the regime of phase transition. That is why we impose the condition **(H)**, which excludes the phase transition.

The reader is referred to [5, 9, 12, 13, 22] and the references therein for recent studies and progresses on this subject. The main new point here is that our estimate (2.33) is explicit and relatively neat.

Remark 2.11. All the results presented in this paper can be extended to more general case:

$$dX_t = \sqrt{2}dB_t + b(X_t, \mu_t)dt$$

where μ_t is the law of X_t , if *b* satisfies some dissipative condition in *x* (uniformly in μ) and a Lipchitzian condition in μ with sufficiently small Lipschitzian constant. For the sake of clarity, we deal only with the case of $b(X_t, \mu_t) = -\nabla V(X_t) - \nabla_x W \circledast \mu_t(X_t)$ in this paper.

2.5. Examples. We first present the Gaussian model for which the constants in Theorem 2.1 and Theorem 2.5 become exact, showing their sharpness.

Example 2.12. (Gaussian model) Let d = 1, and

$$V(x) = \beta \frac{x^2}{2}, \ W(x,y) = -\beta Kxy$$

where $\beta > 0$ is the inverse temperature, $K \ge 0$.

For this model, by some simple calculations we have

$$b_0(r) = -\beta r, \ \forall r > 0.$$

and

$$h'(r) \equiv \beta^{-1}, \ \forall r \ge 0.$$

It is obvious that conditions (2.2) and (2.3) hold, and the assumption (\mathbf{H}) holds once if

$$K < 1. \tag{2.34}$$

But this condition is equivalent to say that the matrix $A = (a_{ij})_{1 \le i,j \le N}$ is positively definite, where

$$a_{ii} = \beta, a_{ij} = \frac{-\beta K}{N-1}, i \neq j$$

A must be the inverse of the covariance matrix of the Gaussian measure $\mu^{(N)}$. In other words **(H)** is equivalent to well defining the equilibrium probability measure $\mu^{(N)}$.

Note that $\|\nabla_{xy}^2 W\|_{\infty} = \beta K$, so we have $c_{Lip} = \frac{1}{\beta(1-K)}$ under (2.34). Moreover (2.22) is satisfied with $M = -\beta$.

• Sharpness of Theorem 2.1. The gradient estimate (2.10) in Theorem 2.1 tells us: if $-\mathcal{L}^{(N)}G = g$, then

$$\|\nabla_i G\|_{\infty} \le \frac{1}{\beta(1-K)} \max_i \|\nabla_i g\|_{\infty}.$$

Let us show that it becomes equality for $g(x^1, \dots, x^N) = \sum_{i=1}^N x^i$. In fact

$$\mathcal{L}^{(N)}g(x^{1},\cdots,x^{N}) = -\sum_{i}\beta x^{i} + \sum_{i}\frac{1}{N-1}\sum_{j\neq i}\beta Kx^{j} = -\beta(1-K)g.$$

In other words $G = \frac{1}{\beta(1-K)}g$ for which the gradient estimate above becomes equality. As the gradient estimate (2.10) comes from (2.9), the process level $W_{1,d_{L^1}}$ estimate (2.9) is sharp too.

• Sharpness of Theorem 2.5. As $\varepsilon \to +\infty$ in (2.23), we have by Theorem 2.5

$$W_{1,d_{l^1}}(P_t^{(N)}(x_0,\cdot),P_t^{(N)}(y_0,\cdot)) \le e^{-\beta(1-K)t}d_{l^1}(x_0,y_0).$$

This is equivalent to say that

$$\max \|\nabla_i P_t^{(N)} g\|_{\infty} \le e^{-\beta(1-K)t} \max \|\nabla_i g\|_{\infty}$$

But it becomes equality for $g = \sum_{i=1}^{N} x^i$: in fact as $\mathcal{L}^{(N)}g = -\beta(1-K)g$, $P_t^{(N)}g = e^{-\beta(1-K)t}g$.

Hence the exponential convergence result (2.24) in Theorem 2.5 is sharp.

Of course for this Gaussian model all results in Theorems 2.1 and 2.5 can be derived easily by using the synchronous coupling, or from the commutativity relation

$$\nabla P_t^{(N)}g = e^{-At}P_t^{(N)}\nabla g$$

which is one of the origins of the Bakry-Emery curvature.

Next we give another two typical models to illustrate our results.

Example 2.13. (Curie-Weiss mean-field lattice model) Let d = 1, and

$$V(x) = \beta(x^4/4 - x^2/2), \ W(x,y) = -\beta Kxy$$

where $\beta = \frac{1}{\kappa T} > 0$ (κ is the Boltzmann constant) is the inverse temperature, $K \in \mathbb{R}^*$. This model is ferromagnetic or anti-ferromagnetic according to K > 0 or K < 0.

By an elementary calculation, we get

$$b_0(r) = \beta r(1 - r^2/4), \ \forall r > 0.$$

It is obvious that conditions (2.2) and (2.3) are satisfied and (2.22) holds with $M = \beta$.

For the assumption (**H**), first notice that $\|\nabla_{xy}^2 W\|_{\infty} = |K|\beta$. Next we estimate $\|h'\|_{\infty}$. By (2.5) and some calculations, we have for any $r \ge 0$

$$h'(r) = \frac{1}{4} \exp(\beta (r^4 - 8r^2)/64) \int_r^{+\infty} s \cdot \exp(\beta (8s^2 - s^4)/64) ds$$
$$= \frac{1}{4} e^{\beta/4} \exp(\beta (r^4 - 8r^2)/64) \int_{r^2/2}^{+\infty} \exp(-\beta (u - 2)^2/16) du.$$

When $\frac{r^2}{2} > 2$, i.e. r > 2, we have

$$h'(r) \le \frac{1}{4}e^{\beta/4}\exp(\beta(r^4 - 8r^2)/64)\sqrt{2\pi\frac{8}{\beta}}\exp(-\beta(\frac{r^2}{2} - 2)^2/16) = \frac{\sqrt{\pi}}{\sqrt{\beta}}.$$
 (2.35)

When $0 \le r \le 2$, by (2.6) we have

$$4h''(r) = -r - \beta r(1 - r^2/4)h'(r) \le 0,$$

hence

$$h'(r) \le h'(0) = \frac{1}{4} e^{\beta/4} \int_0^{+\infty} \exp(-\beta(u-2)^2/16) du < e^{\beta/4} \frac{\sqrt{\pi}}{\sqrt{\beta}}.$$
 (2.36)

Combining (2.35) and (2.36), we obtain $||h'||_{\infty} < e^{\beta/4} \frac{\sqrt{\pi}}{\sqrt{\beta}}$. Thus assumption (**H**) holds once if

$$|K|\sqrt{\pi\beta}e^{\beta/4} \le 1 \tag{2.37}$$

and then the conclusions of Theorem 2.1 and Theorem 2.5 hold under (2.37).

For the result of propagation of chaos, we can take $c_1 = |K|\beta + \varepsilon'$, $c_2 = \frac{\beta}{4}(1+|K|+\frac{\varepsilon'}{\beta})^2$ for any $\varepsilon' > 0$, and $c_3 = 0$. Then condition (2.30) is satisfied and then the conclusion of Theorem 2.9 holds under (2.37).

Example 2.14. (Double-Well confinement potential and quadratic interaction) Let d = 1, and

$$V(x) = \beta(x^4/4 - x^2/2), \ W(x,y) = \beta K(x-y)^2$$

where $\beta > 0$ is the inverse temperature, $K \in \mathbb{R}$. This model has the double-well confinement potential and quadratic interaction potential.

For this model, we have

$$b_0(r) = \beta r(1 - 2K - r^2/4), \ \forall r > 0.$$

So conditions (2.2) and (2.3) are satisfied. By the similar calculations as in Example 2.13, we get

$$\|h'\|_{\infty} < \begin{cases} e^{(1-2K)^2\beta/4} \frac{\sqrt{\pi}}{\sqrt{\beta}}, & \text{if } K \le \frac{1}{2}, \\ \frac{\sqrt{\pi}}{\sqrt{\beta}}, & \text{if } K > \frac{1}{2}. \end{cases}$$

Since $\|\nabla_{xy}^2 W\|_{\infty} = 2|K|\beta$, assumption (**H**) holds once if

$$\begin{cases} 2|K|\sqrt{\pi\beta}e^{(1-2K)^2\beta/4} \le 1, & \text{if } K \le \frac{1}{2} \\ 2|K|\sqrt{\pi\beta} \le 1, & \text{if } K > \frac{1}{2}, \end{cases}$$
(2.38)

and then the conclusion of Theorem 2.1 holds under (2.38).

Furthermore, note that (2.22) holds with $M = \beta(1 - 2K)$, and

$$M + \|\nabla_{xy}^2 W\|_{\infty} = \begin{cases} \beta, & \text{if } K \ge 0, \\ \beta(1 - 4K), & \text{if } K < 0 \end{cases}$$

which is strictly positive. Then the conclusion of Theorem 2.5 holds.

For the result of propagation of chaos in Theorem 2.9, we can take $c_3 = 0$ when $K \ge 0$, and $c_3 = -2K\beta$ when K < 0. To ensure that conditions (2.28) and (2.30) are satisfied, one can take $c_1 = 2|K|\beta + \varepsilon'$, $c_2 = \frac{\beta}{4}(1+2|K| + \frac{\varepsilon'}{\beta})^2$ in the case of K > 0 and $c_1 = -4K\beta + \varepsilon'$, $c_2 = \frac{\beta}{4}(1-4K + \frac{\varepsilon'}{\beta})^2$ in the case of K < 0, for any $\varepsilon' > 0$.

WEI LIU, LIMING WU, AND CHAOEN ZHANG

3. Proofs of Theorems 2.1 and 2.5

3.1. Coupling. We first introduce the approximate componentwise reflection coupling by following A. Eberle [14]. Given $\delta > 0$, let $\lambda_{\delta}, \pi_{\delta} : \mathbb{R}^+ \to [0, 1]$ be two Lipschitz continuous functions such that

$$\lambda_{\delta}(r)^2 + \pi_{\delta}(r)^2 = 1, \ \forall r \in \mathbb{R}^+$$
(3.1)

and

$$\lambda_{\delta}(r) = \begin{cases} 1, & \text{if } r \ge \delta, \\ 0, & \text{if } r \le \delta/2. \end{cases}$$
(3.2)

Then a coupling of two solutions of the mean-field interacting particle system (1.3)with initial values $x_0, y_0 \in (\mathbb{R}^d)^N$ is given by a strong solution of the system

$$dX_{t}^{i,N} = \sqrt{2} [\lambda_{\delta}(|Z_{t}^{i}|) dB_{t}^{1,i} + \pi_{\delta}(|Z_{t}^{i}|) dB_{t}^{2,i}] - \nabla V(X_{t}^{i,N}) dt - \frac{1}{N-1} \sum_{j:j \neq i, 1 \leq j \leq N} \nabla_{x} W(X_{t}^{i,N}, X_{t}^{j,N}) dt, dY_{t}^{i,N} = \sqrt{2} [\lambda_{\delta}(|Z_{t}^{i}|) R_{t}^{i} dB_{t}^{1,i} + \pi_{\delta}(|Z_{t}^{i}|) dB_{t}^{2,i}] - \nabla V(Y_{t}^{i,N}) dt - \frac{1}{N-1} \sum_{j:j \neq i, 1 \leq j \leq N} \nabla_{x} W(Y_{t}^{i,N}, Y_{t}^{j,N}) dt,$$
(3.3)

 $1 \leq i \leq N$. Here $Z_t^i := X_t^{i,N} - Y_t^{i,N}$ and $R_t^i := I_d - 2e_t^i(e_t^i)^T$, where I_d is the *d*-dimensional unit matrix and $e_t^i(e_t^i)^T$ is the orthogonal projection onto the unit vector $e_t^i := Z_t^i/|Z_t^i|$ if $|Z_t^i| \neq 0$. $B_t^{1,i}$ and $B_t^{2,i}$, $1 \leq i \leq N$, are independent standard Brownian motions taking values in \mathbb{R}^d . We will denote $X_t^{(N)} = (X_t^{1,N}, \cdots, X_t^{N,N}), Y_t^{(N)} = (Y_t^{1,N}, \cdots, Y_t^{N,N})$ and $Z_t^{(N)} = V_t^{(N)}$ $Z_t^{(N)} := X_t^{(N)} - Y_t^{(N)}.$ To see that $(X_t^{(N)}, Y_t^{(N)})$ is a coupling process, it is enough to notice that

$$\hat{B}_{t}^{i} := \int_{0}^{t} \lambda_{\delta}(|Z_{s}^{i}|) dB_{s}^{1,i} + \int_{0}^{t} \pi_{\delta}(|Z_{s}^{i}|) dB_{s}^{2,i}
\check{B}_{t}^{i} := \int_{0}^{t} \lambda_{\delta}(|Z_{s}^{i}|) R_{t}^{i} dB_{s}^{1,i} + \int_{0}^{t} \pi_{\delta}(|Z_{s}^{i}|) dB_{s}^{2,i}, 1 \le i \le N,$$
(3.4)

are standard Brownian motions on $(\mathbb{R}^d)^N$.

- **Remark 3.1.** (1) The coupling (3.3) behaves as a reflection coupling when the distance between the two particles $X_t^{i,N}$ and $Y_t^{i,N}$ are larger than δ . When the particles are very close (with distance less than $\frac{1}{2}\delta$), they are driven by the same Brownian motion, i.e., it is a synchronous coupling. And when the distance is between $\frac{1}{2}\delta$ and δ , it is a mixture of reflection coupling and synchronous coupling. The aim is to make λ_{δ} and π_{δ} globally Lipschitz continuous, so that the coupling SDE has a unique strong solution, given the independent Brownian motions $B_t^{1,i}, B_t^{2,i}, 1 \le i \le N$.
- (2) If one adopts the componentwise reflection coupling (i.e. the limit coupling when $\delta \to 0$), since $X^{i,N}$, $Y^{i,N}$ will separate after the time that they meet (i.e. $X_t^{i,N} =$ $Y_t^{i,N}$), the local times will appear when Itô's formula is applied for $|X_t^{i,N} - Y_t^{i,N}|$.

This makes the control of $\sum_{i=1}^{N} |X_t^{i,N} - Y_t^{i,N}|$ difficult to deal with. That is the reason why A. Eberle [14] introduced the synchronous coupling when $|X_t^{i,N} - Y_t^{i,N}|$ is small.

3.2. Proofs of Theorem 2.1.

Proof of Theorem 2.1. **1). Proof of (2.9).** The first inequality in (2.9) is trivial, and next we prove the second inequality. By doing subtraction of the equations in (3.3), we have

$$dZ_{t}^{i} = 2\sqrt{2}\lambda_{\delta}(|Z_{t}^{i}|)e_{t}^{i}d\tilde{B}_{t}^{i} - [\nabla V(X_{t}^{i,N}) - \nabla V(Y_{t}^{i,N})]dt - \frac{1}{N-1}\sum_{j:j\neq i,1\leq j\leq N} [\nabla_{x}W(X_{t}^{i,N}, X_{t}^{j,N}) - \nabla_{x}W(Y_{t}^{i,N}, Y_{t}^{j,N})]dt, \quad (3.5)$$
$$Z_{0}^{i} = x_{0}^{i} - y_{0}^{i},$$

where the processes $\tilde{B}_t^i = \int_0^t (e_s^i)^T dB_s^{1,i}, 1 \leq i \leq N$, are one-dimensional standard Brownian motions such that $\langle \tilde{B}^i, \tilde{B}^j \rangle_t = 0$ for $i \neq j$.

Let $r_t^i = |Z_t^i|, 1 \le i \le N$. By applying Itô's formula, we have

$$dr_{t}^{i} = 1_{\{r_{t}^{i}\neq0\}} 2\sqrt{2}\lambda_{\delta}(r_{t}^{i})d\tilde{B}_{t}^{i} - 1_{\{r_{t}^{i}\neq0\}}\langle e_{t}^{i}, \nabla V(X_{t}^{i,N}) - \nabla V(Y_{t}^{i,N})\rangle dt - 1_{\{r_{t}^{i}\neq0\}}\langle e_{t}^{i}, \frac{1}{N-1}\sum_{j:j\neq i,1\leq j\leq N} [\nabla_{x}W(X_{t}^{i,N}, X_{t}^{j,N}) - \nabla_{x}W(Y_{t}^{i,N}, Y_{t}^{j,N})]\rangle dt + 1_{\{r_{t}^{i}\neq0\}}\sum_{k,l=1}^{d} [1_{\{k=l\}}(r_{t}^{i})^{-1} - (X_{t}^{i,N,k} - Y_{t}^{i,N,k})(X_{t}^{i,N,l} - Y_{t}^{i,N,l})(r_{t}^{i})^{-3}]\lambda_{\delta}(r_{t}^{i})^{2}(I_{d} - R_{t}^{i})_{kl}^{2}dt,$$

$$(3.6)$$

where $X_t^{i,N,k}$ and $Y_t^{i,N,k}$ denote the k-th coordinate of $X_t^{i,N}$ and $Y_t^{i,N}$ respectively, $1 \le k \le d$. Notice that the last term in the right hand side of the above equation equals to 0 by an easy calculation. Hence we get

$$dr_{t}^{i} = 1_{\{r_{t}^{i}\neq0\}} 2\sqrt{2}\lambda_{\delta}(r_{t}^{i})d\tilde{B}_{t}^{i} - 1_{\{r_{t}^{i}\neq0\}}\langle e_{t}^{i}, \frac{1}{N-1}\sum_{j:j\neq i,1\leq j\leq N} [\nabla_{x}W(X_{t}^{i,N}, X_{t}^{j,N}) - \nabla_{x}W(X_{t}^{i,N}, Y_{t}^{j,N})]\rangle dt \\ - 1_{\{r_{t}^{i}\neq0\}}\langle e_{t}^{i}, \nabla V(X_{t}^{i,N}) - \nabla V(Y_{t}^{i,N}) + \frac{1}{N-1}\sum_{j:j\neq i,1\leq j\leq N} [\nabla_{x}W(X_{t}^{i,N}, Y_{t}^{j,N}) - \nabla_{x}W(Y_{t}^{i,N}, Y_{t}^{j,N})]\rangle dt \\ \leq 1_{\{r_{t}^{i}\neq0\}} 2\sqrt{2}\lambda_{\delta}(r_{t}^{i})d\tilde{B}_{t}^{i} + \frac{1}{N-1} \|\nabla_{xy}^{2}W\|_{\infty}\sum_{j:j\neq i,1\leq j\leq N} r_{t}^{j}dt + 1_{\{r_{t}^{i}\neq0\}}b_{0}(r_{t}^{i})dt,$$

$$(3.7)$$

where we use the definition (2.1) of b_0 in the last inequality. Here $d\xi_t \leq d\eta_t$ means that $\eta_t - \xi_t$ is a non-decreasing process.

Let $L_{\lambda_{\delta}}$ be the generator defined by for any function $f \in C^2(0, +\infty)$ and r > 0,

$$L_{\lambda_{\delta}}f(r) := 4\lambda_{\delta}^{2}(r)f''(r) + b_{0}(r)f'(r).$$
(3.8)

Note that $L_{\lambda_{\delta}}$ equals \mathcal{L}_{ref} when $\lambda_{\delta} \equiv 1$.

Applying Itô's formula to the function $h(r_t^i)$ and using (3.7) and the fact that h'(r) > 0, we get for any t > 0 and $i = 1, \dots, N$,

$$dh(r_{t}^{i}) \leq 2\sqrt{2}\lambda_{\delta}(r_{t}^{i})h'(r_{t}^{i})d\tilde{B}_{t}^{i} + h'(r_{t}^{i})b_{0}(r_{t}^{i})dt + 4h''\lambda_{\delta}(r_{t}^{i})^{2}dt + \frac{1}{N-1} \|\nabla_{xy}^{2}W\|_{\infty}h'(r_{t}^{i})\sum_{j:j\neq i,1\leq j\leq N} r_{t}^{j}dt = 2\sqrt{2}\lambda_{\delta}(r_{t}^{i})h'(r_{t}^{i})d\tilde{B}_{t}^{i} + L_{\lambda_{\delta}}h(r_{t}^{i})dt + \frac{1}{N-1} \|\nabla_{xy}^{2}W\|_{\infty}h'(r_{t}^{i})\sum_{j:j\neq i,1\leq j\leq N} r_{t}^{j}dt.$$

$$(3.9)$$

Notice that by the definition of $L_{\lambda_{\delta}}$ and the Poisson equation (2.6),

$$L_{\lambda_{\delta}}h(r) = \mathcal{L}_{ref}h(r) + 4(\lambda_{\delta}^2 - 1)h''(r) = -r + (1 - \lambda_{\delta}^2)(r + b_0(r)h'(r)).$$
(3.10)

Then

$$-\sum_{i=1}^{N} \left(L_{\lambda_{\delta}} h(r_{t}^{i}) + \frac{1}{N-1} \| \nabla_{xy}^{2} W \|_{\infty} h'(r_{t}^{i}) \sum_{j: j \neq i, 1 \leq j \leq N} r_{t}^{j} \right)$$

$$\geq (1 - \| \nabla_{xy}^{2} W \|_{\infty} \| h' \|_{\infty}) \sum_{i=1}^{N} r_{t}^{i} - \sum_{i=1}^{N} (1 - \lambda_{\delta}(r_{t}^{i})^{2}) (r_{t}^{i} + b_{0}(r_{t}^{i})h'(r_{t}^{i}))$$

which is bounded from below by $-N(\delta + \sup_{r \in (0,\delta)} b_0^+(r) ||h'||_{\infty})$ according to the conditions **(H)** and (2.3). By integrating from 0 to T and taking expectation in the previous inequality (3.9) for $dh(r_t^i)$ and using Fatou's lemma, we have for any T > 0,

$$\mathbb{E} \int_{0}^{T} \left\{ (1 - \|\nabla_{xy}^{2}W\|_{\infty} \|h'\|_{\infty}) \sum_{i=1}^{N} r_{t}^{i} - \sum_{i=1}^{N} (1 - \lambda_{\delta}(r_{t}^{i})^{2})(r_{t}^{i} + b_{0}(r_{t}^{i})h'(r_{t}^{i})) \right\} dt \\
\leq \sum_{i=1}^{N} h(|x_{0}^{i} - y_{0}^{i}|).$$
(3.11)

Letting $\mathbb{P}_{x}|_{[0,T]}$ be the law of $(X_{t}^{(N)})_{t\in[0,T]}$, we obtain by assumption (**H**) and (3.11)

$$W_{1,d_{L^{1}[0,T]}}(\mathbb{P}_{x_{0}}|_{[0,T]},\mathbb{P}_{y_{0}}|_{[0,T]}) \leq \mathbb{E}\int_{0}^{T} d_{l^{1}}(X_{t}^{(N)},Y_{t}^{(N)})dt = \mathbb{E}\int_{0}^{T}\sum_{i=1}^{N}r_{t}^{i}dt$$

$$\leq \frac{1}{1-\|\nabla_{xy}^{2}W\|_{\infty}\|h'\|_{\infty}}\left\{\sum_{i=1}^{N}h(|x_{0}^{i}-y_{0}^{i}|) + \sum_{i=1}^{N}\mathbb{E}\int_{0}^{T}(1-\lambda_{\delta}(r_{t}^{i})^{2})(r_{t}^{i}+b_{0}^{+}(r_{t}^{i})h'(r_{t}^{i}))dt\right\}.$$
(3.12)

By the definition of λ_{δ} and the assumption $\lim_{r\to 0} b_0^+(r) = 0$, the second term in the right hand side of the inequality above converges to 0, a.s., as $\delta \downarrow 0$. Hence

$$W_{1,d_{L^{1}[0,T]}}(\mathbb{P}_{x_{0}}|_{[0,T]},\mathbb{P}_{y_{0}}|_{[0,T]}) \leq \frac{1}{1 - \|\nabla_{xy}^{2}W\|_{\infty}\|h'\|_{\infty}} \sum_{i=1}^{N} h(|x_{0}^{i} - y_{0}^{i}|).$$
(3.13)

Let Q_n be an optimization coupling of $(\mathbb{P}_{x_0}|_{[0,n]}, \mathbb{P}_{y_0}|_{[0,n]})$ for $W_{1,d_{L^1[0,n]}}(\mathbb{P}_{x_0}|_{[0,n]}, \mathbb{P}_{y_0}|_{[0,n]})$. Then $\{Q_n|_{[0,T]}; n \geq T\}$ is tight for any finite time T (because their marginal distributions are respectively $\mathbb{P}_{x_0}|_{[0,T]}$ and $\mathbb{P}_{y_0}|_{[0,T]}$), hence one can find a probability measure Q on $C(\mathbb{R}^+, (\mathbb{R}^d)^N)^2$ such that $Q_n|_{[0,T]} \to Q|_{[0,T]}$ weakly for all T > 0. Thus

$$W_{1,d_{L^{1}[0,\infty]}}(\mathbb{P}_{x_{0}},\mathbb{P}_{y_{0}}) \leq \mathbb{E}^{Q} \int_{0}^{\infty} d_{l^{1}}(\gamma_{1}(t),\gamma_{2}(t))dt$$

$$= \lim_{T \to +\infty} \mathbb{E}^{Q} \int_{0}^{T} d_{l^{1}}(\gamma_{1}(t),\gamma_{2}(t))dt$$

$$\leq \lim_{T \to \infty} \lim_{T \leq n \to +\infty} \mathbb{E}^{Q_{n}} \int_{0}^{T} d_{l^{1}}(\gamma_{1}(t),\gamma_{2}(t))dt$$

$$\leq \lim_{n \to \infty} W_{1,d_{L^{1}[0,n]}}(\mathbb{P}_{x_{0}}|_{[0,n]},\mathbb{P}_{y_{0}}|_{[0,n]}).$$

The converse inequality is evident. Therefore we have

$$W_{1,d_{L^1}}(\mathbb{P}_{x_0},\mathbb{P}_{y_0}) = \lim_{n \to \infty} W_{1,d_{L^1[0,n]}}(\mathbb{P}_{x_0}|_{[0,n]},\mathbb{P}_{y_0}|_{[0,n]})$$

From this and (3.13) we obtain (2.9).

2). Proof of (2.10). Note that for any Lipschitzian function g w.r.t the d_{l^1} -metric on $(\mathbb{R}^d)^N$, g is $\mu^{(N)}$ -integrable because $\int \sum_{i=1}^N |x^i| d\mu^{(N)}(x) < +\infty$. So we can assume $\mu^{(N)}(g) = 0$ without loss of generality. Moreover we have

$$\begin{split} \int_{0}^{+\infty} |P_{t}^{(N)}g(x)|dt &= \int_{0}^{+\infty} |P_{t}^{(N)}g(x) - \int P_{t}^{(N)}g(y)d\mu^{(N)}(y)|dt \\ &\leq \|g\|_{Lip(d_{l^{1}})} \int_{(\mathbb{R}^{d})^{N}} \int_{0}^{+\infty} W_{d_{l^{1}}}(P_{t}^{(N)}(x,\cdot), P_{t}^{(N)}(y,\cdot))dtd\mu^{(N)}(y) \\ &\leq \frac{\|g\|_{Lip(d_{l^{1}})}}{1 - \|\nabla_{xy}^{2}W\|_{\infty}} \|h'\|_{\infty} \int_{(\mathbb{R}^{d})^{N}} \sum_{i=1}^{N} h(|x^{i} - y^{i}|)d\mu^{(N)}(y) \\ &< +\infty, \end{split}$$

then the unique solution of the Poisson equation $-\mathcal{L}^{(N)}G = g$ with $\mu^{(N)}(G) = 0$ is given by $G(x) = \int_0^{+\infty} P_t^{(N)}g(x)dt, \forall x \in (\mathbb{R}^d)^N.$ For each $1 \leq i \leq N$, letting $\tilde{x}^i \neq x^i$ and $\tilde{x} \in (\mathbb{R}^d)^N$ so that $(\tilde{x})^j = x^j$ for $j \neq i$ and $(\tilde{x})^i = \tilde{x}^i$, we have

$$\begin{aligned} |\nabla_{i}G(x)| &\leq \limsup_{\tilde{x}^{i} \to x^{i}} \frac{|G(x) - G(\tilde{x})|}{|x^{i} - \tilde{x}^{i}|} \\ &\leq \limsup_{\tilde{x}^{i} \to x^{i}} \frac{1}{|x^{i} - \tilde{x}^{i}|} \int_{0}^{+\infty} |P_{t}^{(N)}g(x) - P_{t}^{(N)}g(\tilde{x})|dt \\ &\leq \limsup_{\tilde{x}^{i} \to x^{i}} \frac{1}{|x^{i} - \tilde{x}^{i}|} \|g\|_{Lip(d_{l^{1}})} \int_{0}^{+\infty} W_{d_{l^{1}}}(P_{t}^{(N)}(x, \cdot), P_{t}^{(N)}(\tilde{x}, \cdot))dt \quad (3.14) \\ &\leq \frac{1}{1 - \|\nabla_{xy}^{2}W\|_{\infty} \|h'\|_{\infty}} \|g\|_{Lip(d_{l^{1}})} \lim_{\tilde{x}^{i} \to x^{i}} \frac{h(|x^{i} - \tilde{x}^{i}|)}{|x^{i} - y^{i}|} \\ &= \frac{h'(0)}{1 - \|\nabla_{xy}^{2}W\|_{\infty} \|h'\|_{\infty}} \|g\|_{Lip(d_{l^{1}})}, \end{aligned}$$

where the fourth inequality follows from (3.13).

3.3. Proof of Theorem 2.5.

Proof. Here we also adopt the coupling (3.3) . Let h be defined as in (2.5). Define for any $\varepsilon > 0$,

$$h_{\varepsilon}(r) := h(r) + \varepsilon r, \forall r \ge 0, \tag{3.15}$$

and

$$H_t^{\varepsilon} := e^{K_{\varepsilon}t} \sum_{i=1}^N h_{\varepsilon}(r_t^i),$$

where $r_t^i = |X_t^{i,N} - Y_t^{i,N}|, 1 \le i \le N$, as in the proof of Theorem 2.1. By using Ito's formula and (3.7), we get for any $t \ge 0$,

$$dH_t^{\varepsilon} \le 2\sqrt{2}e^{K_{\varepsilon}t} \sum_{i=1}^N \lambda_{\delta}(r_t^i) d\tilde{B}_t^i + K_{\varepsilon}H_t^{\varepsilon}dt + e^{K_{\varepsilon}t} \sum_{i=1}^N (L_{\lambda_{\delta}}h(r_t^i) + \varepsilon b_0(r_t^i))dt + e^{K_{\varepsilon}t} \sum_{i=1}^N (h'(r_t^i) + \varepsilon) \sum_{j:j \neq i} \frac{1}{N-1} \|\nabla_{xy}^2 W\|_{\infty} r_t^j dt$$
(3.16)

Let

$$D_t^{\varepsilon} := K_{\varepsilon} H_t^{\varepsilon} + e^{K_{\varepsilon} t} \sum_{i=1}^N (L_{\lambda_{\delta}} h(r_t^i) + \varepsilon b_0(r_t^i)) + \frac{1}{N-1} e^{K_{\varepsilon} t} \|\nabla_{xy}^2 W\|_{\infty} \sum_{i \neq j, 1 \le i, j \le N} (h'(r_t^i) + \varepsilon) r_t^j$$

$$(3.17)$$

be the drift term at the right hand side above. Calculating as in the proof of Theorem 2.1, we have

$$D_{t}^{\varepsilon} \leq e^{K_{\varepsilon}t} \sum_{i=1}^{N} [1 - \lambda_{\delta}(r_{t}^{i})^{2}][r_{t}^{i} + b_{0}(r_{t}^{i})h'(r_{t}^{i})] \\ + e^{K_{\varepsilon}t} \sum_{i=1}^{N} \{K_{\varepsilon}h_{\varepsilon}(r_{t}^{i}) - [1 - (\|h'\|_{\infty} + \varepsilon)\|\nabla_{xy}^{2}W\|_{\infty}]r_{t}^{i} + \varepsilon b_{0}(r_{t}^{i})\} \\ \leq e^{K_{\varepsilon}t} \sum_{i=1}^{N} [1 - \lambda_{\delta}(r_{t}^{i})^{2}][r_{t}^{i} + b_{0}(r_{t}^{i})h'(r_{t}^{i})] \\ + e^{K_{\varepsilon}t} \sum_{i=1}^{N} \{K_{\varepsilon}(\|h'\|_{\infty} + \varepsilon) + \varepsilon M - [1 - (\|h'\|_{\infty} + \varepsilon)\|\nabla_{xy}^{2}W\|_{\infty}]\}r_{t}^{i},$$
(3.18)

where we use the assumption $b_0(r) \leq Mr, \forall r > 0$.

By taking

$$K_{\varepsilon} = \frac{1 - \|\nabla_{xy}^2 W\|_{\infty} \|h'\|_{\infty} - \varepsilon (M + \|\nabla_{xy}^2 W\|_{\infty})}{\|h'\|_{\infty} + \varepsilon},$$
(3.19)

the second term in the right hand of the inequality above vanishes. Then by taking expectation in (3.16) and the localization stopping time technique, we have for any $t \ge 0$,

$$\mathbb{E}e^{K_{\varepsilon}t}\sum_{i=1}^{N}h_{\varepsilon}(r_{t}^{i}) \leq \sum_{i=1}^{N}h_{\varepsilon}(|x_{0}^{i}-y_{0}^{i}|) + \mathbb{E}\int_{0}^{t}e^{K_{\varepsilon}s}[1-\lambda_{\delta}(r_{t}^{i})^{2}][r_{t}^{i}+b_{0}^{+}(r_{t}^{i})h'(r_{t}^{i})]ds.$$
(3.20)

Note that the second term in the right hand side of the above inequality converges to 0 as $\delta \downarrow 0$, under the assumption (2.3). Therefore we get

$$W_{1,d_{l^{1}}}(P_{t}^{(N)}(x_{0},\cdot),P_{t}^{(N)}(y_{0},\cdot)) \leq \lim_{\delta \to 0} \mathbb{E} \sum_{i=1}^{N} r_{t}^{i}$$

$$\leq \sup_{r>0} \frac{r}{h(r) + \varepsilon r} \lim_{\delta \to 0} \mathbb{E} \sum_{i=1}^{N} h_{\varepsilon}(r_{t}^{i})$$

$$\leq \sup_{r>0} \frac{r}{h(r) + \varepsilon r} e^{-K_{\varepsilon}t} \sum_{i=1}^{N} h_{\varepsilon}(|x_{0}^{i} - y_{0}^{i}|)$$

$$\leq \sup_{r>0} \frac{r}{h(r) + \varepsilon r} \cdot \sup_{r>0} \frac{h(r) + \varepsilon r}{r} e^{-K_{\varepsilon}t} \sum_{i=1}^{N} |x_{0}^{i} - y_{0}^{i}|$$
(3.21)

where the third inequality above follows by (3.20).

WEI LIU, LIMING WU, AND CHAOEN ZHANG

4. PROPAGATION OF CHAOS

We begin with a uniform in time control of the second moment, which is more or less known, see e.g. Cattiaux *et al.* [9].

Lemma 4.1. Suppose that there exist some positive constants c_1, c_2, c_3 such that

$$\langle x, \nabla V(x) \rangle \ge c_1 |x|^2 - c_2, \forall x \in \mathbb{R}^d$$

$$(4.1)$$

and

$$\langle z, \nabla_{xx}^2 W(x, y) z \rangle \ge -c_3 |z|^2, \forall x, y, z \in \mathbb{R}^d.$$
 (4.2)

Assume

$$c_1 - c_3 - \|\nabla_{xy}^2 W\|_{\infty} > 0.$$
(4.3)

Let X_t be a solution of (1.2) with $\mathbb{E}|X_0|^2 < \infty$, then for any $\varepsilon \in (0, c_1 - c_3 - \|\nabla_{xy}^2 W\|_{\infty})$,

$$\sup_{t \ge 0} \mathbb{E}(|X_t|^2)^{\frac{1}{2}} \le \max\{m_2(\mu_0), \hat{c}(\varepsilon)\},\tag{4.4}$$

where $m_2(\mu_0)$ and $\hat{c}(\varepsilon)$ are given in (2.32).

Proof. By Itô's formula, we have

$$d|X_t|^2 = -2\langle X_t, \nabla V(X_t)\rangle dt - 2\langle X_t, \nabla_x W \circledast \mu_t(X_t)\rangle dt + 2d \cdot dt + 2\sqrt{2}\langle X_t, dB_t\rangle$$

Notice that for any $x \in \mathbb{R}^d$, we have

$$\langle x, \nabla_x W \circledast \mu_t(x) - \nabla_x W \circledast \mu_t(0) \rangle = \langle x, \int_0^1 \frac{d}{ds} \nabla_x W \circledast \mu_t(sx) ds \rangle$$

$$= \langle x, \int_0^1 \frac{d}{ds} \int_{\mathbb{R}^d} \nabla_x W(sx, y) \mu_t(dy) ds \rangle$$

$$= \int_0^1 \int_{\mathbb{R}^d} \langle x, \nabla_{xx}^2 W(sx, y) x \rangle \mu_t(dy) ds$$

$$\ge -c_3 |x|^2,$$

$$(4.5)$$

where the last inequality follows from (4.2).

On the other hand,

$$\begin{aligned} |\nabla_x W \circledast \mu_t(0)\rangle| &\leq |\nabla_x W(0,0)| + \int_{\mathbb{R}^d} |\nabla_x W(0,y) - \nabla_x W(0,0)|\mu_t(dy) \\ &\leq |\nabla_x W(0,0)| + \|\nabla_{xy}^2 W\|_{\infty} \mathbb{E}|X_t|. \end{aligned}$$
(4.6)

Therefore we have

$$\begin{aligned} d|X_t|^2 &\leq 2\left(c_3|X_t|^2 + \|\nabla_{xy}^2 W\|_{\infty} |X_t| \mathbb{E}|X_t| + |\nabla_x W(0,0)| |X_t|\right) dt \\ &+ 2(-c_1|X_t|^2 + c_2 + d) dt + 2\sqrt{2} \langle X_t, dB_t \rangle \\ &\leq -2(c_1 - c_3 - \varepsilon) |X_t|^2 dt + 2\|\nabla_{xy}^2 W\|_{\infty} |X_t| \mathbb{E}|X_t| dt \\ &+ 2(d + c_2 + \frac{1}{4\varepsilon} |\nabla_x W(0,0)|^2) dt + 2\sqrt{2} \langle X_t, dB_t \rangle \end{aligned}$$

where $0 < \varepsilon < c_1 - c_3 - \|\nabla_{xy}^2 W\|_{\infty}$. By the previous stochastic differential inequality, $|X_t|^2 + \int_0^t [2(c_1 - c_3 - \varepsilon)|X_s|^2 - 2\|\nabla_{xy}^2 W\|_{\infty}|X_s|\mathbb{E}|X_s|]ds - 2t(d + c_2 + \frac{1}{4\varepsilon}|\nabla_x W(0, 0)|^2)$

is a local supermartingale, then a supermartingale by Fatou's lemma. Then for any T > 0, we have

$$\mathbb{E}|X_{0}|^{2} \geq \mathbb{E}|X_{T}|^{2} + 2(c_{1} - c_{3} - \varepsilon) \int_{0}^{T} \mathbb{E}|X_{s}|^{2}ds - 2\|\nabla_{xy}^{2}W\|_{\infty} \int_{0}^{T} (\mathbb{E}|X_{s}|)^{2}ds - 2T(d + c_{2} + \frac{1}{4\varepsilon}|\nabla_{x}W(0, 0)|^{2}) \geq 2(c_{1} - c_{3} - \varepsilon - \|\nabla_{xy}^{2}W\|_{\infty}) \int_{0}^{T} \mathbb{E}|X_{s}|^{2}ds - 2T(d + c_{2} + \frac{1}{4\varepsilon}|\nabla_{x}W(0, 0)|^{2}),$$

which implies $\mathbb{E} \int_0^T |X_s|^2 ds < +\infty$. In other words $\int_0^t 2\sqrt{2} \langle X_s, dB_s \rangle$ is a L^2 -martingale. By taking expectation in (4.1) we obtain by (4.5) and (4.6),

$$\frac{d}{dt}\mathbb{E}|X_t|^2 \leq -2c_1\mathbb{E}|X_t|^2 + 2[c_3\mathbb{E}|X_t|^2 + \|\nabla_{xy}^2W\|_{\infty}(\mathbb{E}|X_t|)^2 + |\nabla_xW(0,0)|\mathbb{E}|X_t|] + 2(d+c_2) \\
\leq -2(c_1 - c_3 - \|\nabla_{xy}^2W\|_{\infty} - \varepsilon)\mathbb{E}|X_t|^2 + 2(d+c_2 + \frac{1}{4\varepsilon}|\nabla_xW(0,0)|^2)$$
(4.7)

where $0 < \varepsilon < c_1 - c_3 - \|\nabla_{xy}^2 W\|_{\infty}$. By Gronwall's lemma we get for any $t \ge 0$

$$\begin{split} \mathbb{E}|X_t|^2 &\leq e^{-2(c_1 - c_3 - \|\nabla_{xy}^2 W\|_{\infty} - \varepsilon)t} \left(\mathbb{E}|X_0|^2 - \frac{d + c_2 + \frac{1}{4\varepsilon} |\nabla_x W(0, 0)|^2}{c_1 - c_3 - \|\nabla_{xy}^2 W\|_{\infty} - \varepsilon} \right) \\ &+ \frac{d + c_2 + \frac{1}{4\varepsilon} |\nabla_x W(0, 0)|^2}{c_1 - c_3 - \|\nabla_{xy}^2 W\|_{\infty} - \varepsilon} \\ &\leq \max\left\{ \mathbb{E}|X_0|^2, \frac{d + c_2 + \frac{1}{4\varepsilon} |\nabla_x W(0, 0)|^2}{c_1 - c_3 - \|\nabla_{xy}^2 W\|_{\infty} - \varepsilon} \right\} \end{split}$$

the desired result.

Following the proof above we have the much stronger uniform Gaussian integrability for X_t , which should be of independent interest.

Lemma 4.2. Assume (4.1), (4.2) and (4.3). Let X_t be a solution of (1.2) with

 $\mathbb{E}\exp\left(\lambda_0|X_0|^2\right) < \infty, \quad for \ some \ \lambda_0 > 0.$

If

$$0 < \lambda \le \min\{\lambda_0; \frac{1}{2}(c_1 - c_3 - \|\nabla_{xy}^2 W\|_{\infty} - \varepsilon)\}$$

for some $\varepsilon > 0$, then

(

$$\sup_{t \ge 0} \mathbb{E} \exp(\lambda |X_t|^2) < +\infty.$$

Proof. By Itô's formula, we have by the estimates leading to (4.7) in the proof of Lemma 4.1,

$$d \exp(\lambda |X_t|^2)$$

$$= \lambda \exp(\lambda |X_t|^2) \left([2d - 2\langle X_t, \nabla V(X_t) + \nabla_x W \circledast \mu_t(X_t) \rangle] dt + 2\sqrt{2} \langle X_t, dB_t \rangle \right)$$

$$+ 4\lambda^2 |X_t|^2 \exp(\lambda |X_t|^2) dt$$

$$\leq \lambda \exp(\lambda |X_t|^2) \left[-2(c_1 - c_3 - \|\nabla_{xy}^2 W\|_{\infty} - \varepsilon - 2\lambda) |X_t|^2 + 2(d + c_2 + \frac{1}{4\varepsilon} |\nabla_x W(0, 0)|) \right] dt$$

$$+ \lambda \exp(\lambda |X_t|^2) 2\sqrt{2} \langle X_t, dB_t \rangle$$

where $\varepsilon > 0, \lambda > 0$ verify $c_1 - c_3 - \|\nabla_{xy}^2 W\|_{\infty} - \varepsilon - 2\lambda > 0$. Taking L > 0 large sufficient so that

$$c_5 := 2(c_1 - c_3 - \|\nabla_{xy}^2 W\|_{\infty} - \varepsilon - 2\lambda)L^2 - 2(d + c_2 + \frac{1}{4\varepsilon}|\nabla_x W(0,0)|) > 0,$$

and noting that

$$-ax^{2} + b \leq -(aL^{2} - b) + aL^{2}1_{|x| \leq L}, \ \forall a > 0, \forall x \in \mathbb{R}$$

we obtain by following the same argument as in Lemma 4.1

$$\frac{d}{dt}\mathbb{E}\exp(\lambda|X_t|^2) \le -\lambda c_5\mathbb{E}\exp(\lambda|X_t|^2) + 2(c_1 - c_3 - \|\nabla_{xy}^2 W\|_{\infty} - \varepsilon - 2\lambda)L^2\lambda e^{\lambda L^2}.$$

Therefore by Gronwall's lemma

$$\sup_{t\geq 0} \mathbb{E} \exp(\lambda |X_t|^2) < +\infty.$$

Next we present the proof of Theorem 2.9, which is quite close to those of Theorems 2.1 and 2.5 .

Proof of Theorem 2.9. Let λ_{δ} and π_{δ} be defined as in Section 3.1. Consider the following coupling between the independent copies \bar{X}_t^i , $1 \leq i \leq N$ of the nonlinear diffusion processes (1.2) and the mean-field interacting particle system (1.3):

$$d\bar{X}_{t}^{i} = \sqrt{2} [\lambda_{\delta}(|Z_{t}^{i}|) dB_{t}^{1,i} + \pi_{\delta}(|Z_{t}^{i}|) dB_{t}^{2,i}] - \nabla V(\bar{X}_{t}^{i}) dt - \nabla_{x} W \circledast \mu_{t}(\bar{X}_{t}^{i}) dt,$$

$$dX_{t}^{i,N} = \sqrt{2} [\lambda_{\delta}(|Z_{t}^{i}|) R_{t}^{i} dB_{t}^{1,i} + \pi_{\delta}(|Z_{t}^{i}|) dB_{t}^{2,i}] - \nabla V(X_{t}^{i,N}) dt - \frac{1}{N-1} \sum_{j: j \neq i, 1 \leq j \leq N} \nabla_{x} W(X_{t}^{i,N}, X_{t}^{j,N}) dt.$$
(4.8)

Here $Z_t^i := \bar{X}_t^i - X_t^{i,N}$ and $R_t^i := I_d - 2e_t^i e_t^{i,T}$, where I_d is the *d*-dimensional unit matrix and $e_t^i e_t^{i,T}$ is the orthogonal projection onto the unit vector $e_t^i := Z_t^i / |Z_t^i|$ if $|Z_t^i| \neq 0$. $B_t^{1,i}$ and $B_t^{2,i}$, $1 \le i \le N$, are independent standard Brownian motions in \mathbb{R}^d . We assume that \bar{X}_t^i and $X_t^{i,N}$, $1 \le i \le N$ have the same starting points X_0^i , $1 \le i \le N$, i.i.d. of law μ_0 . The independence of \bar{X}_t^i , $1 \le i \le N$ comes from the fact that the Brownian motions $\{\int_0^t \lambda_{\delta}(|Z_s^i|) dB_s^{1,i} + \int_0^t \pi_{\delta}(|Z_s^i|) dB_s^{2,i}, 1 \le i \le N\}$ are independent because their inter-brackets are zero.

By doing subtraction of the equations in (4.8), we have

$$\begin{split} dZ_t^i &= 2\sqrt{2}\lambda_{\delta}(|Z_t^i|)e_t^id\tilde{B}_t^i - [\nabla V(\bar{X}_t^i) - \nabla V(X_t^{i,N})]dt - \nabla_x W \circledast \mu_t(\bar{X}_t^i)dt \\ &+ \frac{1}{N-1}\sum_{j:j\neq i, 1\leq j\leq N} \nabla_x W(X_t^{i,N}, X_t^{j,N})dt, \end{split}$$

where the processes $\tilde{B}_t^i = \int_0^t (e_s^i)^T dB_s^{1,i}, 1 \leq i \leq N$, are one-dimensional standard Brownian motions such that $\langle \tilde{B}^i, \tilde{B}^j \rangle_t = 0$ for $i \neq j$.

Let $r_t^i = |Z_t^i|, 1 \le i \le N$. By applying Itô's formula, we have

$$\begin{split} dr_t^i &= \mathbf{1}_{\{r_t^i \neq 0\}} 2\sqrt{2} \lambda_{\delta}(r_t^i) d\tilde{B}_t^i - \mathbf{1}_{\{r_t^i \neq 0\}} \langle e_t^i, \nabla V(\bar{X}_t^i) - \nabla V(X_t^{i,N}) \rangle dt \\ &\quad - \mathbf{1}_{\{r_t^i \neq 0\}} \langle e_t^i, \nabla_x W \circledast \mu_t(\bar{X}_t^i) - \frac{1}{N-1} \sum_{j:j \neq i, 1 \leq j \leq N} \nabla_x W(X_t^{i,N}, X_t^{j,N}) \rangle dt \\ &= \mathbf{1}_{\{r_t^i \neq 0\}} 2\sqrt{2} \lambda_{\delta}(r_t^i) d\tilde{B}_t^i \\ &\quad - \mathbf{1}_{\{r_t^i \neq 0\}} \langle e_t^i, \nabla V(\bar{X}_t^i) - \nabla V(X_t^{i,N}) \rangle dt \\ &\quad - \mathbf{1}_{\{r_t^i \neq 0\}} \langle e_t^i, \nabla_x W \circledast \mu_t(\bar{X}_t^i) - \frac{1}{N-1} \sum_{j:j \neq i, 1 \leq j \leq N} \nabla_x W(\bar{X}_t^i, \bar{X}_t^j) \rangle dt \\ &\quad - \mathbf{1}_{\{r_t^i \neq 0\}} \langle e_t^i, \frac{1}{N-1} \sum_{j:j \neq i, 1 \leq j \leq N} [\nabla_x W(\bar{X}_t^i, \bar{X}_t^j) - \nabla_x W(\bar{X}_t^i, X_t^{j,N})] \rangle dt \\ &\quad - \mathbf{1}_{\{r_t^i \neq 0\}} \langle e_t^i, \frac{1}{N-1} \sum_{j:j \neq i, 1 \leq j \leq N} [\nabla_x W(\bar{X}_t^i, X_t^{j,N}) - \nabla_x W(X_t^{i,N}, X_t^{j,N})] \rangle dt. \end{split}$$

Remark that the sum of the first and the fourth drift terms above is $\leq b_0(r_t^i)dt$, the third drift term above is $\leq \frac{1}{N-1} \|\nabla_{xy}^2 W\|_{\infty} \sum_{j: j \neq i, 1 \leq j \leq N} r_t^j dt$, and the second drift term is bounded by $I_t^i dt$, where

$$I_t^i := |\nabla_x W \circledast \mu_t(\bar{X}_t^i) - \frac{1}{N-1} \sum_{j: j \neq i, 1 \le j \le N} \nabla_x W(\bar{X}_t^i, \bar{X}_t^j)|.$$
(4.9)

Therefore we obtain

$$dr_t^i \le 2\sqrt{2\lambda_\delta}(r_t^i)d\tilde{B}_t^i + b_0(r_t^i)dt + \frac{1}{N-1} \|\nabla_{xy}^2 W\|_{\infty} \sum_{j:j \ne i, 1 \le j \le N} r_t^j dt + I_t^i dt.$$
(4.10)

Recall that for any $\varepsilon \ge 0$, $h_{\varepsilon}(r) = h(r) + \varepsilon r$, $\forall r \ge 0$. By using (4.10) and Itô's formula again, we get

$$\begin{aligned} dh_{\varepsilon}(r_{t}^{i}) &\leq 2\sqrt{2}\lambda_{\delta}(r_{t}^{i})h_{\varepsilon}'(r_{t}^{i})d\tilde{B}_{t}^{i} + 4\lambda_{\delta}^{2}(r_{t}^{i})h_{\varepsilon}''(r_{t}^{i})dt + b_{0}(r_{t}^{i})h_{\varepsilon}'(r_{t}^{i})dt \\ &+ \frac{1}{N-1} \|\nabla_{xy}^{2}W\|_{\infty}h_{\varepsilon}'(r_{t}^{i})\sum_{j:j\neq i,1\leq j\leq N} r_{t}^{j}dt + h_{\varepsilon}'(r_{t}^{i})I_{t}^{i}dt \\ &= 2\sqrt{2}\lambda_{\delta}(r_{t}^{i})h_{\varepsilon}'(r_{t}^{i})d\tilde{B}_{t}^{i} + [4\lambda_{\delta}^{2}(r_{t}^{i})h''(r_{t}^{i}) + b_{0}(r_{t}^{i})h'(r_{t}^{i})]dt + \varepsilon b_{0}(r_{t}^{i})dt \\ &+ \frac{1}{N-1} \|\nabla_{xy}^{2}W\|_{\infty}(h'(r_{t}^{i}) + \varepsilon)\sum_{j:j\neq i,1\leq j\leq N} r_{t}^{j}dt + (h'(r_{t}^{i}) + \varepsilon)I_{t}^{i}dt \\ &\leq 2\sqrt{2}\lambda_{\delta}(r_{t}^{i})h_{\varepsilon}'(r_{t}^{i})d\tilde{B}_{t}^{i} + [1 - \lambda_{\delta}^{2}(r_{t}^{i})][r_{t}^{i} + b_{0}(r_{t}^{i})h'(r_{t}^{i})]dt - (1 - \varepsilon M)r_{t}^{i}dt \\ &+ \frac{1}{N-1} \|\nabla_{xy}^{2}W\|_{\infty}(\|h'\|_{\infty} + \varepsilon)\sum_{j:j\neq i,1\leq j\leq N} r_{t}^{j}dt + (\|h'\|_{\infty} + \varepsilon)I_{t}^{i}dt, \end{aligned}$$

$$(4.11)$$

where the last inequality follows from (3.8), (3.10) and (2.22).

Taking expectation in the inequality above and using the fact that $r_t^i, 1 \leq i \leq N$ have the same law, and setting

$$c_{\varepsilon} := 1 - \|\nabla_{xy}^2 W\|_{\infty} \|h'\|_{\infty} - \varepsilon (M + \|\nabla_{xy}^2 W\|_{\infty}),$$

we have

$$d\mathbb{E}h_{\varepsilon}(r_t^1) \leq \mathbb{E}[1 - \lambda_{\delta}(r_t^1)^2][r_t^1 + b_0^+(r_t^1)h'(r_t^1)]dt + (\|h'\|_{\infty} + \varepsilon)\mathbb{E}I_t^1dt - c_{\varepsilon}\mathbb{E}r_t^1dt$$

$$(4.12)$$
Proof of part (a). Choose $\varepsilon = 0, c_0 = 1 - \|\nabla_{xy}^2 W\|_{\infty} \|h'\|_{\infty}$. For any $1 \leq k \leq N$, by (4.12) we have

$$\begin{split} \frac{1}{k} W_{1,d_{L^{1}[0,T]}}(\mathbb{P}_{\mu_{0}^{\otimes N}}^{[1,k],N}|_{[0,T]}, \mathbb{Q}_{\mu_{0}}^{\otimes k}|_{[0,T]}) &\leq \frac{1}{k} \mathbb{E} \int_{0}^{T} \sum_{i=1}^{k} r_{t}^{i} dt = \int_{0}^{T} \mathbb{E} r_{t}^{1} dt \\ &\leq \frac{1}{c_{0}} \|h'\|_{\infty} \int_{0}^{T} \mathbb{E} I_{t}^{1} dt dt + \frac{1}{c_{0}} \mathbb{E} \int_{0}^{T} [1 - \lambda_{\delta}(r_{t}^{1})^{2}] [r_{t}^{1} + b_{0}^{+}(r_{t}^{1})h'(r_{t}^{1})] dt. \end{split}$$

Letting $\delta \rightarrow 0+,$ the last term tends to zero. Hence

$$\frac{1}{k}W_{1,d_{L^{1}[0,T]}}(\mathbb{P}_{\mu_{0}^{\otimes N}}^{[1,k],N}|_{[0,T]},\mathbb{Q}_{\mu_{0}}^{\otimes k}|_{[0,T]}) \leq \frac{\|h'\|_{\infty}}{c_{0}} \int_{0}^{T} \mathbb{E}I_{t}^{1}dt.$$

$$(4.13)$$

Next we estimate $\mathbb{E}I_t^1$, which is the only new point w.r.t. the proofs in Theorems 2.1 and 2.5. Note that $\bar{X}_t^j, 2 \leq j \leq N$ are independent copies of \bar{X}_t^1 , and

$$\mathbb{E}[\nabla_x W(\bar{X}^1_t, \bar{X}^j_t) | \bar{X}^1_t] = \nabla_x W \circledast \mu_t(\bar{X}^1_t).$$

Thus by using Cauchy-Schwartz inequality, we get

$$\begin{split} \mathbb{E}I_{t}^{1} &\leq \left(\mathbb{E}\left\{\mathbb{E}\left[|\nabla_{x}W \circledast \mu_{t}(\bar{X}_{t}^{1}) - \frac{1}{N-1}\sum_{2 \leq j \leq N} \nabla_{x}W(\bar{X}_{t}^{1}, \bar{X}_{t}^{j})|^{2}|\bar{X}_{t}^{1}\right]\right\}\right)^{\frac{1}{2}} \\ &= \left(\mathbb{E}\frac{1}{N-1}\int |\nabla_{x}W(\bar{X}_{t}^{1}, y) - \nabla_{x}W \ast \mu_{t}(\bar{X}_{t}^{1})|^{2}d\mu_{t}(y)\right)^{\frac{1}{2}} \\ &\leq \frac{1}{\sqrt{N-1}}\|\nabla_{xy}^{2}W\|_{\infty}\left(\int_{x \in \mathbb{R}^{d}} |x - \mu_{t}(\bar{X}_{t}^{1})|^{2}\mu_{t}(dx)\right)^{\frac{1}{2}} \\ &\leq \frac{1}{\sqrt{N-1}}\|\nabla_{xy}^{2}W\|_{\infty}\sup_{t \geq 0}(\mathbb{E}|X_{t}|^{2})^{\frac{1}{2}}. \end{split}$$
(4.14)

Plugging (4.14) into (4.13), we get

$$\frac{1}{k}W_{1,d_{L^{1}[0,T]}}(\mathbb{P}_{\mu_{0}^{\otimes N}}^{[1,k],N}|_{[0,T]},\mathbb{Q}_{\mu_{0}}^{\otimes k}|_{[0,T]}) \leq \frac{T}{\sqrt{N-1}}\frac{\|\nabla_{xy}^{2}W\|_{\infty}\|h'\|_{\infty}}{1-\|\nabla_{xy}^{2}W\|_{\infty}\|h'\|_{\infty}}\sup_{t\geq 0}(\mathbb{E}|X_{t}|^{2})^{\frac{1}{2}}.$$

Then by using Lemma 4.1, we obtain the desired result (2.31).

Proof of part (b). For any
$$\varepsilon > 0$$
, by (4.12) we have
 $d\mathbb{E}h_{\varepsilon}(r_t^1) \leq \mathbb{E}[1-\lambda_{\delta}(r_t^1)^2][r_t^1+b_0^+(r_t^1)h'(r_t^1)]dt + (\|h'\|_{\infty}+\varepsilon)\mathbb{E}I_t^1dt - c_{\varepsilon} \cdot \inf_{r>0} \frac{r}{h(r)+\varepsilon r} \mathbb{E}h_{\varepsilon}(r_t^1)dt$
(4.15)

Plugging (4.14) into (4.15), we obtain by Gronwall's inequality that for any $\varepsilon > 0$ so that $\beta = c_{\varepsilon} \cdot \inf_{r>0} \frac{r}{h(r) + \varepsilon r} > 0$ (i.e. $K_{\varepsilon} > 0$),

$$\inf_{r>0} \frac{h_{\varepsilon}(r)}{r} \cdot \mathbb{E}|\bar{X}_{t}^{1} - X_{t}^{1,N}|) \leq \mathbb{E}h_{\varepsilon}(|\bar{X}_{t}^{1} - X_{t}^{1,N}|) \\
\leq \int_{0}^{t} e^{-\beta(t-s)} \frac{1}{\sqrt{N-1}} (\|h'\|_{\infty} + \varepsilon) \|\nabla_{xy}^{2} W\|_{\infty} \sup_{t\geq 0} (\mathbb{E}|X_{t}|^{2})^{\frac{1}{2}} ds \qquad (4.16) \\
+ \int_{0}^{t} e^{-\beta(t-s)} \mathbb{E}[1 - \lambda_{\delta}(r_{s}^{1})^{2}][r_{s}^{1} + b_{0}^{+}(r_{s}^{1})h'(r_{s}^{1})] ds.$$

By letting $\delta \to 0+$, the last term tends to zero. As the joint law of $(\bar{X}_t^i, 1 \le i \le k)$ is $\mu_t^{\otimes k}$, we get for any $1 \le k \le N$,

$$W_{1,d_{l^{1}}}(\mu_{t}^{\otimes k},\mu_{t}^{[1,k],N}) \leq \limsup_{\delta \to 0} \mathbb{E} \sum_{i=1}^{k} |\bar{X}_{t}^{i} - X_{t}^{i,N}| = k \cdot \limsup_{\delta \to 0} \mathbb{E} |\bar{X}_{t}^{1} - X_{t}^{1,N}|$$
$$\leq k \cdot \sup_{r>0} \frac{r}{h(r) + \varepsilon r} \frac{1}{\beta \sqrt{N-1}} (\|h'\|_{\infty} + \varepsilon) \|\nabla_{xy}^{2} W\|_{\infty} \sup_{t\geq 0} (\mathbb{E} |X_{t}|^{2})^{\frac{1}{2}}$$
$$= \frac{k}{\sqrt{N-1}} \frac{A_{\varepsilon}}{K_{\varepsilon}} \|\nabla_{xy}^{2} W\|_{\infty} \sup_{t\geq 0} (\mathbb{E} |X_{t}|^{2})^{\frac{1}{2}},$$
(4.17)

which completes the proof by using Lemma 4.1.

The proof above yields

Proposition 4.3. Under the conditions of Theorem 2.9, we have

$$\mathbb{E}W_{1}\left(\frac{1}{N}\sum_{i=1}^{N}\delta_{X_{t}^{i,N}}, \frac{1}{N}\sum_{i=1}^{N}\delta_{\bar{X}_{t}^{i}}\right) \leq \frac{1}{\sqrt{N-1}}\frac{A_{\varepsilon}}{K_{\varepsilon}}\|\nabla_{xy}^{2}W\|_{\infty}\sup_{t\geq0}(\mathbb{E}|X_{t}|^{2})^{\frac{1}{2}}$$
(4.18)

where $(\bar{X}_t^i)_{t\geq 0}$, $i \geq 1$ are independent copies of the solution $(X_t)_{t\geq 0}$ of the McKean-Vlasov equation (1.2), and $X_t^{i,N}$, $1 \leq i \leq N$ are defined as in (1.3).

Proof. Notice that

$$\mathbb{E}W_1\left(\frac{1}{N}\sum_{i=1}^N \delta_{X_t^{i,N}}, \frac{1}{N}\sum_{i=1}^N \delta_{\bar{X}_t^i}\right) \le \mathbb{E}[\frac{1}{N}\sum_{i=1}^N |X_t^{i,N} - \bar{X}_t^i|] = \mathbb{E}\frac{1}{N}\sum_{i=1}^N r_t^i,$$

where $r_t^i, 1 \leq i \leq N$ are the same as defined in the proof of Theorem 2.9. And by (4.17), we have

$$\limsup_{\delta \to 0} \mathbb{E} \frac{1}{N} \sum_{i=1}^{n} r_t^i = \limsup_{\delta \to 0} \mathbb{E} r_t^1 \le \frac{1}{\sqrt{N-1}} \frac{A_{\varepsilon}}{K_{\varepsilon}} \|\nabla_{xy}^2 W\|_{\infty} \sup_{t \ge 0} (\mathbb{E} |X_t|^2)^{\frac{1}{2}}.$$

Therefore we obtain (4.18).

Remark 4.4. In one-dimensional case, i.e. d = 1, it is well known that

$$W_1\left(\frac{1}{N}\sum_{i=1}^N \delta_{\bar{X}_t^i}, \mu_t\right) = \int_{-\infty}^\infty |\frac{1}{N}\sum_{i=1}^N \mathbf{1}_{(-\infty,x]}(\bar{X}_t^i) - \mu_t(-\infty,x]|dx.$$

Then letting $F_t(x) = \mu_t(-\infty, x]$ (the cumulative distribution function), we have by the Cauchy-Schwarz inequality,

$$\mathbb{E}W_1\left(\frac{1}{N}\sum_{i=1}^N \delta_{\bar{X}_t^i}, \mu_t\right) \le \int_{\mathbb{R}} \sqrt{\operatorname{Var}\left(\frac{1}{N}\sum_{i=1}^N \mathbb{1}_{(-\infty,x]}(\bar{X}_t^i)\right)} dx$$
$$= \frac{1}{\sqrt{N}} \int_{\mathbb{R}} \sqrt{F_t(x)(1-F_t(x))} dx,$$

where the last factor is uniformly bounded in time t > 0 by some constant K once if $\sup_{t\geq 0} \mathbb{E}|X_t|^{2+\varepsilon} < +\infty$ for some $\varepsilon > 0$. The latter uniform $2 + \varepsilon$ -moment condition is verified once if μ_0 has the $2 + \varepsilon$ -moment by the arguments in Lemmas 4.1 and 4.2. In other words if μ_0 has the $2 + \varepsilon$ -moment, there is some constant K > 0 such that

$$\mathbb{E}W_1\left(\frac{1}{N}\sum_{i=1}^N \delta_{\bar{X}_t^i}, \mu_t\right) \le \frac{K}{\sqrt{N}}, \ \forall t > 0$$
(4.19)

and then the same type bound holds for $\mathbb{E}W_1\left(\frac{1}{N}\sum_{i=1}^N \delta_{X_t^{i,N}}, \mu_t\right)$, by Proposition 4.3 and the triangular inequality.

But (4.19) does not hold in the multi-dimensional (d > 1) case, see Fournier and Guillin [16].

Remark 4.5. A consequence of Proposition 4.3 is on the bias of $\frac{1}{N} \sum_{i=1}^{N} f(X_t^{i,N})$ from $\mu_t(f)$: if f is Lipschitzian on \mathbb{R}^d ,

$$bias_{t}(f) := |\mathbb{E}\frac{1}{N}\sum_{i=1}^{N} f(X_{t}^{i,N}) - \mu_{t}(f)| = |\mathbb{E}\frac{1}{N}\sum_{i=1}^{N} f(X_{t}^{i,N}) - \mathbb{E}\frac{1}{N}\sum_{i=1}^{N} f(\bar{X}_{t}^{i})|$$

$$\leq ||f||_{Lip}\mathbb{E}W_{1}\left(\frac{1}{N}\sum_{i=1}^{N}\delta_{X_{t}^{i,N}}, \frac{1}{N}\sum_{i=1}^{N}\delta_{\bar{X}_{t}^{i}}\right)$$

$$\leq \frac{||f||_{Lip}}{\sqrt{N-1}}\frac{A_{\varepsilon}}{K_{\varepsilon}}||\nabla_{xy}^{2}W||_{\infty}\sup_{t\geq 0}(\mathbb{E}|X_{t}|^{2})^{\frac{1}{2}}.$$

It is expected that the bias is of order O(1/N), which remains an open question.

5. QUANTITATIVE CONCENTRATION INEQUALITIES

This section is devoted to the concentration inequalities of the mean-field interaction particle system (1.3), as applications of our main theorems. This kind of concentration estimate are useful to numerical simulations and mean-field limit. Under the conditions that V is uniformly convex and W is convex, Malrieu [25] established logarithmic Sobolev inequality and then used its connection with optimal transport and concentration of measure to get the following non-asymptotic bounds on the deviation of the empirical mean of an observable f from its true mean,

$$\sup_{\|f\|_{Lip} \le 1} \mathbb{P}\left\{ \left| \frac{1}{N} \sum_{i=1}^{N} f(X_t^{i,N}) - \mu_t(f) \right| > \frac{A}{\sqrt{N}} + \delta \right\} \le 2e^{-\lambda N \delta^2}, \ t > 0, \ \delta \ge 0$$
(5.1)

where A and λ are positive constants depending on the particle system.

As pointed out in [6], this approach can lead to nice bounds but it is limited to a finite number of observables. Bolley-Guillin-Villani [6, Theorem 2.9] obtained for any t > 0 fixed and $\delta > 0$

$$\mathbb{P}\left\{\sup_{\|f\|_{Lip} \le 1} \left|\frac{1}{N} \sum_{i=1}^{N} f(X_t^{i,N}) - \mu_t(f)\right| > \delta\right\} \le C(1 + t\delta^{-2})e^{-K(t)N\delta^2}, \tag{5.2}$$

for all N big enough (quantifiable), where K(t) depending on t is some explicitly computable constant. Furthermore, Bolley [4] got quantitative concentration inequalities on the sample path space with uniform norm, on a given time interval [0, T], which implies (5.2) by projection at time $t \in [0, T]$.

5.1. Uniform in time concentration inequality. Our previous general results allow us to generalize (5.1) and to reinforce (5.2) (under stronger conditions of course).

Proposition 5.1. Assume **(H)**, (2.2) and (2.22). Let $X_t^{(N)} = (X_t^{1,N}, \dots, X_t^{N,N}), \forall t \geq 0$, then for any Lipschitzian function F on $(\mathbb{R}^d)^N$, we have for any lower bounded convex function φ on \mathbb{R} ,

$$\mathbb{E}_{x}\varphi\left(F(X_{T}^{(N)}) - \mathbb{E}_{x}F(X_{T}^{(N)})\right) \leq \mathbb{E}\varphi\left(\alpha A_{\varepsilon}\sqrt{\frac{N}{2K_{\varepsilon}}}\xi\right), \ \forall x \in (\mathbb{R}^{d})^{N}, \ \forall T > 0$$
(5.3)

where ξ is some standard real Gaussian random variable of law $\mathcal{N}(0, 1)$, $\alpha := ||F||_{Lip(d_{l^1})} = \max_{1 \le i \le N} ||\nabla_i F||_{\infty}$, A_{ε} and K_{ε} are given in Theorem 2.5.

In particular for any initial distribution μ_0 satisfying the Gaussian integrability assumption on \mathbb{R}^d , we have for any $\delta, T > 0$

$$\mathbb{P}_{\mu_0^{\otimes N}}\left\{F(X_T^{(N)}) - \mathbb{E}_{\mu_0^{\otimes N}}F(X_T^{(N)}) > \delta\right\} \le \exp\left(-\frac{K_{\varepsilon}\delta^2}{N\alpha^2 A_{\varepsilon}^2 \left[1 + 2c_G(\mu_0)K_{\varepsilon}e^{-2K_{\varepsilon}T}\right]}\right).$$
(5.4)

Proof. Without loss of generality we may assume that $\alpha = \max_{1 \le i \le N} \|\nabla_i F\|_{\infty} = 1$.

By approximation we may assume that F is C^2 -smooth with bounded derivatives of the first and the second order. For any initial position $x \in (\mathbb{R}^d)^N$, let $M_t = \mathbb{E}_x(F(X_T^{(N)})|\mathcal{F}_t), 0 \le t \le T$. Then by applying Itô's formula to $u(t,x) = P_{T-t}F(x)$, we have

$$F(X_T^{(N)}) - \mathbb{E}_x F(X_T^{(N)}) = M_T - M_0 = \sum_{i=1}^N \int_0^T \nabla_i P_{T-t} F(X_t^{(N)}) dB_t^i, \qquad (5.5)$$

Note that by Theorem 2.5, for any $\varepsilon > 0$ such that $K_{\varepsilon} > 0$, we have

$$W_{d_{l^{1}}}(P_{t}^{(N)}(x,\cdot),P_{t}^{(N)}(y,\cdot)) \le A_{\varepsilon}e^{-K_{\varepsilon}t}d_{l^{1}}(x,y), \forall x,y \in (\mathbb{R}^{d})^{N},$$
(5.6)

which implies that

$$|\nabla_i P_{T-t}F| \le A_{\varepsilon} e^{-K_{\varepsilon}(T-t)}, \ 1 \le i \le N,$$
(5.7)

where A_{ε} and K_{ε} are the same as given in Theorem 2.5.

Since $M_t = \xi_{\tau_t}$ where (ξ_t) is a real valued Brownian motion w.r.t. some new filtration $(\tilde{\mathcal{F}}_t)$ and

$$\tau_t = \langle M \rangle_t = \int_0^t \sum_{i=1}^N |\nabla_i P_{t-s} F(X_s^{(N)})|^2 ds \le \frac{A_{\varepsilon}^2}{2K_{\varepsilon}} N =: CN$$

is a stopping time w.r.t. $(\tilde{\mathcal{F}}_t)$, we obtain

$$\mathbb{E}_{x}\varphi\left(F(X_{T}^{(N)}) - \mathbb{E}_{x}F(X_{T}^{(N)})\right) = \mathbb{E}\varphi\left(M_{T} - M_{0}\right) = \mathbb{E}\varphi(\xi_{\tau_{T}})$$
$$= \mathbb{E}\varphi\left(\mathbb{E}(\xi_{CN}|\tilde{\mathcal{F}}_{\tau_{T}})\right)$$
$$\leq \mathbb{E}\varphi\left(\xi_{CN}\right) \quad \text{(by Jensen's inequality)}$$
$$= \mathbb{E}\varphi\left(A_{\varepsilon}\sqrt{\frac{N}{2K_{\varepsilon}}}\xi_{1}\right)$$
(5.8)

the desired result (5.3).

Letting $g(x) := \mathbb{E}_x F(X_T^{(N)}), \forall x \in (\mathbb{R}^d)^N$. By (5.7) we have

$$\|g\|_{Lip(d_{l^1})} = \max_{1 \le i \le N} \|\nabla_i g\|_{\infty} \le A_{\varepsilon} e^{-K_{\varepsilon} T}.$$
(5.9)

Applying (5.3) to $\varphi(z) = e^{\lambda z}$ ($\lambda \in \mathbb{R}$), we get

$$\begin{split} & \mathbb{E}_{\mu_0^{\otimes N}} \exp\left(\lambda[F(X_T^{(N)}) - \mathbb{E}_{\mu_0^{\otimes N}}F(X_T^{(N)})])\right) \\ &= \int_{(\mathbb{R}^d)^N} \mathbb{E}_x \exp\left(\lambda[F(X_T^{(N)}) - \mathbb{E}_x F(X_T^{(N)})]\right) \cdot \exp\left(\lambda[g(x) - \mu_0^{\otimes N}(g)]\right) d\mu_0^{\otimes N}(x) \\ &\leq \int_{(\mathbb{R}^d)^N} \mathbb{E} \exp\left(\lambda A_{\varepsilon} \sqrt{\frac{N}{2K_{\varepsilon}}} \xi_1\right) \cdot \exp\left(\lambda[g(x) - \mu_0^{\otimes N}(g)]\right) d\mu_0^{\otimes N}(x) \\ &\leq \exp\left(\frac{NA_{\varepsilon}^2 \lambda^2}{4K_{\varepsilon}}\right) \exp\left(\frac{\lambda^2}{2} N c_G(\mu_0) \|g\|_{Lip(d_{l^1})}^2\right) \\ &\leq \exp\left(\frac{N\lambda^2 A_{\varepsilon}^2}{2} \left[\frac{1}{2K_{\varepsilon}} + c_G(\mu_0) e^{-2K_{\varepsilon}T}\right]\right) \end{split}$$

where the third and the last inequality follows from the Gaussian concentration condition on the initial distribution μ_0 (see (2.19) in Remark 2.3) and (5.9) respectively.

Finally the concentration inequality (5.10) is derived from the above inequality by the standard procedure of Chebyshev's inequality and optimization over λ .

Example 5.2. Given a Lipschitzian observable $f : \mathbb{R}^d \to \mathbb{R}$ with $||f||_{Lip} = 1$ and $N \ge 2$, let $F(x) = \frac{1}{N} \sum_{i=1}^{N} f(x^i)$. For any T > 0,

$$F(X_T^{(N)}) = \frac{1}{N} \sum_{i=1}^N f(X_T^{i,N})$$

is the empirical mean of f at time T. Since

$$||F||_{Lip_{(d_{l^1})}} = \frac{1}{N} ||f||_{Lip} = \frac{1}{N}$$

we obtain by (5.10) for any $\delta > 0$,

$$\mathbb{P}_{\mu_0^{\otimes N}}\left\{\frac{1}{N}\sum_{i=1}^N f(X_T^{i,N}) - \mathbb{E}_{\mu_0^{\otimes N}}f(X_T^{1,N}) > \delta\right\} \le \exp\left(-\frac{NK_{\varepsilon}\delta^2}{A_{\varepsilon}^2\left[1 + 2c_G(\mu_0)K_{\varepsilon}e^{-2K_{\varepsilon}T}\right]}\right).$$
(5.10)

As the absolute value of the bias $|\mathbb{E}_{\mu_0^{\otimes N}} f(X_T^{1,N}) - \mu_T(f)| \leq A/\sqrt{N}$ by Remark 4.5, our result above generalizes Malrieu's result (5.1) to the case that V may have many wells.

5.2. Concentration for time average. The counterpart of Proposition 5.1 for the empirical time average is presented in the following result.

Proposition 5.3. Assume **(H)**, (2.2) and (2.3). Given any $T \in (0, +\infty]$, let F be any $d_{L^1[0,T]}$ -Lipschitzian continuous function on $C([0,T], (\mathbb{R}^d)^N)$, given by

$$F(X_{[0,T]}^{(N)}) := G\left(\int_0^T g_1(X_t^{(N)})dt, \cdots, \int_0^T g_n(X_t^{(N)})dt\right),$$

where $G \in C^2(\mathbb{R}^n)$, $g_i \in C^2((\mathbb{R}^d)^N, \mathbb{R})$, $1 \leq i \leq n$. Then for any convex function φ on \mathbb{R} and any starting point $X_0^{(N)} = x \in (\mathbb{R}^d)^N$, we have

$$\mathbb{E}_{x}\varphi\left(F(X_{[0,T]}^{(N)}) - \mathbb{E}_{x}F(X_{[0,T]}^{(N)})\right) \leq \mathbb{E}\varphi\left(\sqrt{NT}\|F\|_{Lip(d_{L^{1}[0,T]})}c_{Lip}\xi\right),$$
(5.11)

where ξ is some standard real Gaussian random variable of law $\mathcal{N}(0,1)$, and

$$c_{Lip} = \frac{h'(0)}{1 - \|\nabla_{xy}^2 W\|_{\infty} \|h'\|_{\infty}}.$$

Proof. Let $\{\mathcal{F}_t\}_{t\geq 0}$ be the filtration generated by the process $(X_t^{(N)})_{t\geq 0}$ and

$$M_t = \mathbb{E}(F(X_{[0,T]}^{(N)})|\mathcal{F}_t), \ 0 \le t \le T.$$

Then by the martingale representation theorem, we have

$$F(X_{[0,T]}^{(N)}) - \mathbb{E}F(X_{[0,T]}^{(N)}) = M_T - M_0 = \sum_{i=1}^N \int_0^T \beta_t^i dB_t^i,$$
(5.12)

where $\beta_t^i, 1 \leq i \leq N$ are adapted processes w.r.t. \mathcal{F}_t , and $B_t^i, 1 \leq i \leq N$ are N independent standard Brownian motions on \mathbb{R}^d .

Let $A_t^k = \int_0^t g_k(X_s^{(N)}) ds$, $1 \le k \le n$, and $A_t = (A_t^1, \cdots, A_t^n)$. Note that

$$M_t = \phi(A_t, X_t^{(N)})$$

where

$$\phi(a,x) := \mathbb{E}\left(G\left(a_1 + \int_t^T g_1(X_s^{(N)})ds, \cdots, a_n + \int_t^T g_N(X_s^{(N)})ds\right) | X_t^{(N)} = x\right),$$

for $a = (a_1, \dots, a_n) \in \mathbb{R}^n, x \in (\mathbb{R}^d)^N$. Since φ is C^2 (for V, W are C^2), we can apply Itô's formula to obtain that

$$\beta_t^i = \partial_{x_i} \varphi(A_t, X_t^{(N)}).$$

For any $x = (x^1, \dots, x^i, \dots, x^N) \in (\mathbb{R}^d)^N$, denote $y = (x^1, \dots, y^i, \dots, x^N)$ which only differs from x at the *i*-th coordinate. Let $(X_t^{(N)})_{t\geq 0}, (Y_t^{(N)})_{t\geq 0}$ be an optimal coupling of $\mathbb{P}_x, \mathbb{P}_y$ for $W_{1,d_{L^1[0,T]}}(\mathbb{P}_x, \mathbb{P}_y)$ (this optimal coupling exists because $d_{L^1[0,T]}$ is lower semi-continuous from $(C(\mathbb{R}^+, (\mathbb{R}^d)^N))^2$ to $[0, +\infty]$). Then for any $0 \leq t \leq T$ and $i = 1, \cdots, N$, we have

$$\begin{aligned} |\partial_{x_{i}}\phi(a,x)| &\leq \limsup_{y^{i} \to x^{i}} \frac{|\phi(a,x) - \phi(a,y)|}{|x^{i} - y^{i}|} |\\ &= \limsup_{y^{i} \to x^{i}} \frac{1}{|x^{i} - y^{i}|} |\mathbb{E}[G(a + \int_{0}^{T-t} g(X_{s}^{(N)}ds)] - \mathbb{E}[G(a + \int_{0}^{T-t} g(Y_{s}^{(N)}ds)]| \\ &\leq \limsup_{y^{i} \to x^{i}} \frac{\|F\|_{Lip(d_{L^{1}[0,T]})}}{|x^{i} - y^{i}|} \mathbb{E} \int_{0}^{\infty} d_{l^{1}}(X_{s}^{(N)}, Y_{s}^{(N)}) ds \end{aligned}$$
(5.13)
$$&= \|F\|_{Lip(d_{L^{1}[0,T]})} \limsup_{y^{i} \to x^{i}} \frac{W_{1,d_{L^{1}}}(\mathbb{P}_{x}, \mathbb{P}_{y})}{|x^{i} - y^{i}|} \\ &\leq \|F\|_{Lip(d_{L^{1}[0,T]})} \cdot c_{Lip} \end{aligned}$$

where the last inequality follows by Theorem 2.1.

We now repeat the argument in the proof of Proposition 5.1. Since $\sum_{i=1}^{N} \int_{0}^{T} \beta_{t}^{i} dB_{t}^{i} = \xi_{\tau_{T}}$ where (ξ_{t}) is a real valued Brownian motion w.r.t. some new filtration $(\tilde{\mathcal{F}}_{t})$ and $\tau_{T} = \int_{0}^{T} \sum_{i=1}^{N} |\beta_{t}^{i}|^{2} dt \leq ||F||_{Lip(d_{L^{1}[0,T]})}^{2} c_{Lip}^{2} NT =: CNT$ is a stopping time w.r.t. $(\tilde{\mathcal{F}}_{t})$, we obtain

$$\mathbb{E}_{x}\varphi\left(F(X_{[0,T]}^{(N)}) - \mathbb{E}F(X_{[0,T]}^{(N)})\right) = \mathbb{E}\varphi\left(\sum_{i=1}^{N} \int_{0}^{T} \beta_{t}^{i} dB_{t}^{i}\right) = \mathbb{E}\varphi(\xi_{\tau_{T}})$$

$$= \mathbb{E}\varphi\left(\mathbb{E}(\xi_{CNT} | \tilde{\mathcal{F}}_{\tau_{T}})\right)$$

$$\leq \mathbb{E}\varphi\left(\xi_{CNT}\right) \quad \text{(by Jensen's inequality)}$$

$$= \mathbb{E}\varphi\left(\sqrt{NT} ||F||_{Lip(d_{L^{1}[0,T]})}c_{Lip}\xi_{1}\right)$$
(5.14)

the desired result.

Next we give the proof of Corollary 2.4.

Proof of Corollary 2.4. For any given $\lambda, T > 0$, let

$$F(X_{[0,T]}^{(N)}) = \frac{1}{T} \int_0^T U_N(f_m)(X_t^{(N)}) dt.$$

Since f_m is 1-Lipschitzian w.r.t. the d_{l^1} -metric on $(\mathbb{R}^d)^m$, by an easy calculation we have

$$||F||_{Lip_{(d_{L^1}[0,T])}} \le \frac{m}{NT}.$$

Let $g(x) = \mathbb{E}_x F$, $\forall x \in (\mathbb{R}^d)^N$. For any fixed initial value $x \in (\mathbb{R}^d)^N$, by applying Proposition 5.3 with $\varphi(z) = e^{\lambda z}$, we get

$$\mathbb{E}_{x} \exp\left(\lambda \left[\frac{1}{T} \int_{0}^{T} U_{N}(f_{m})(X_{t}^{(N)}) dt - g(x)\right]\right)$$

$$\leq \mathbb{E} \exp\left(\frac{m\lambda}{\sqrt{NT}} c_{Lip}\xi_{1}\right) = \exp\left(\frac{m^{2}\lambda^{2}c_{Lip}^{2}}{2NT}\right).$$
(5.15)

31

By the proof of Proposition 5.3,

$$||g||_{Lip(d_{l^1})} \le c_{Lip} ||F||_{d_{L^1[0,T]}} \le \frac{mc_{Lip}}{NT}.$$

By the condition (2.18) and its consequence (2.19), the product measure $\mu_0^{\otimes N}$ satisfies

$$\int_{(\mathbb{R}^d)^N} e^{\lambda(g-\mu_0^{\otimes N}(g))} d\mu_0^{\otimes N} \leq \exp\left(\frac{1}{2}Nc_G(\mu_0)\lambda^2 \|g\|_{Lip(d_{l^1})}^2\right)$$

$$\leq \exp\left(\frac{1}{2NT^2}c_G(\mu_0)\lambda^2 m^2 c_{Lip}^2\right).$$
(5.16)

Hence for the i.i.d. initial values $X_0^{1,N}, \dots, X_0^{N,N}$ with the common law μ_0 , noting that

$$\mathbb{E}\frac{1}{T}\int_0^T U_N(f_m)(X_t^{(N)})dt = \mu_0^{\otimes N}(g)$$

we have

$$\mathbb{E} \exp\left(\lambda \left[\frac{1}{T} \int_{0}^{T} U_{N}(f_{m})(X_{t}^{(N)}) dt - \mathbb{E}\frac{1}{T} \int_{0}^{T} U_{N}(f_{m})(X_{t}^{(N)}) dt\right]\right)$$

$$= \int_{(\mathbb{R}^{d})^{N}} \mathbb{E}_{x} \left[\exp\left(\lambda \left[\frac{1}{T} \int_{0}^{T} U_{N}(f_{m})(X_{t}^{(N)}) dt - g(x)\right]\right)\right] e^{\lambda(g(x) - \mu_{0}^{\otimes N}(g))} d\mu_{0}^{\otimes N}(x)$$

$$\leq \exp\left(\frac{m^{2}\lambda^{2}c_{Lip}^{2}}{2NT}\right) \int_{(\mathbb{R}^{d})^{N}} e^{\lambda(g(x) - \mu_{0}^{\otimes N}(g))} d\mu_{0}^{\otimes N}(x)$$

$$\leq \exp\left(\frac{m^{2}\lambda^{2}c_{Lip}^{2}}{2NT} \left(1 + \frac{c_{G}(\mu_{0})}{T}\right)\right)$$
(5.17)

where the second inequality follows from (5.15), and the last inequality is a consequence of (5.16). This gives us (2.20). Finally (2.21) follows from (2.20), by the standard procedure of Chebyshev's inequality and optimization over $\lambda > 0$.

Remark 5.4. The time-particle average $\frac{1}{NT} \int_0^T f(X_t^{i,N}) dt$ is used to approximate $\mu_{\infty}(f)$ where $\mu_{\infty} = \lim_{t \to +\infty} \mu_t$ is the unique equilibrium state of the McKean-Vlasov equation (a consequence of Corollary 2.2 by Banach's fixed point theorem). For applying

Corollary 2.4, it remains to bound the bias

$$\begin{split} & \|\mathbb{E}_{\mu_{0}^{\otimes N}} \frac{1}{NT} \int_{0}^{T} \sum_{i=1}^{N} f(X_{t}^{i,N}) dt - \mu_{\infty}(f) \| \\ & \leq \|\mathbb{E}_{\mu_{0}^{\otimes N}} \frac{1}{T} \int_{0}^{T} [f(X_{t}^{1,N}) - \mu_{t}(f)] dt \| + \frac{1}{T} \int_{0}^{T} |\mu_{t}(f) - \mu_{\infty}(f)| dt \\ & \leq \frac{1}{T} \int_{0}^{T} |\mu_{t}^{1,N}(f) - \mu_{t}(f)| dt + \|f\|_{Lip} \frac{1}{T} \int_{0}^{T} W_{1}(\mu_{t},\mu_{\infty}) dt \\ & \leq \|f\|_{Lip} \left(\sup_{t \geq 0} W_{1}(\mu_{t}^{1,N},\mu_{t}) + \frac{1}{T} \int_{0}^{T} W_{1}(\mu_{t},\mu_{\infty}) dt \right) \\ & \leq \|f\|_{Lip} \left(\frac{A}{\sqrt{N}} + \frac{B}{T} \right) \end{split}$$

where in the last inequality, the first term comes from the uniform in time propagation of chaos (2.33) in Theorem 2.9, and the second follows by (2.12) in Corollary 2.2. We believe that the bias should be of order O(1/N + 1/T), but we do not know how to prove it.

References

- S. Benachour, B. Roynette, D. Talay, and P. Vallois, Nonlinear self-stabilizing processes. I. Existence, invariant probability, propagation of chaos, *Stoch. Proc. Appl.* 75 (2) (1998), 173-201.
- [2] S. Benachour, B. Roynette, and P. Vallois, Nonlinear self-stabilizing processes. II. Convergence to invariant probability. Stoch. Proc. Appl. 75 (2) (1998), 203-224.
- [3] S. Bobkov and F. Götze, Exponential integrability and transportation cost related to logarithmic Sobolev inequalities J. Funct. Anal. 163 (1999), 1-28.
- [4] F. Bolley, Quantitative concentration inequalities on sample path space for mean field interaction. ESAIM Probab. Stat. 14 (2010), 192-209.
- [5] F. Bolley, A. Guillin and F. Malrieu, Trend to equilibrium and particle approximation for a weakly selfconsistent Vlasov-Fokker-Planck equation. ESAIM: Mathematical Modelling and Numerical Analysis 44 (2010), 867-884.
- [6] F. Bolley, A. Guillin and C. Villani, Quantitative concentration inequalities for empirical measures on non-compcact spaces. *Probab. Theory Relat. Fields* 137 (2007), 541-593.
- [7] R. Buckdahn, J. Li, S. Peng and C. Rainer, Mean-field stochastic differential equations and associated PDEs. Ann. Probab. 45 (2017) 824-878.
- [8] J.A. Carrillo, R.J. McCann, and C. Villani, Kinetic equilibration rates for granular media and related equations: entropy dissipation and mass transportation estimates. *Rev. Mat. Iberoam.* 19 (3) (2003), 971-1018.
- [9] P. Cattiaux, A. Guillin, and F. Malrieu. Probabilistic approach for granular media equations in the non-uniformly convex case. *Probab. Theory Relat. Fields* 140 (2008) 19-40.
- [10] H. Djellout, A. Guillin and L. Wu. Transportation cost-information inequalities for random dynamical systems and diffusions. Ann. Probab. 32 (3B) (2004) 2702-2732.
- [11] H. Djellout, L. Wu, Lipschitzian norm estimate of one-dimensional Poisson equations. Ann. Inst. Henri Poincaré (B) Prob. and Stat., 47 (2) (2011), 450-465.
- [12] P. Del Moral, J. Tugaut, Uniform propagation of chaos and creation of chaos for a class of nonlinear diffusions. To appear in Stochastic Analysis and Applications.
- [13] A. Durmus, A. Eberle, A. Guillin and R. Zimmer, An elementary approach to uniform in Time propagation of chaos, to appear in Proceedings of the American Mathematical Society.

- [14] A. Eberle, Reflection couplings and contraction rates for diffusions. Probab. Theory Relat. Fields 166 (2016), 851-886.
- [15] A. Eberle, A. Guillin, and R. Zimmer, Quantitative Harris type theorems for diffusions and McKean-Vlasov processes. Published electronically in Trans. Am. Math. Soc., on September 28, 2018.
- [16] N. Fournier and A. Guillin, On the rate of convergence in Wasserstein distance of the empirical measure. Probab. Theory Relat. Fields 162 (2015), 707-738.
- [17] J. Gärtner, On the McKean-Vlasov limit for interacting diffusions. *Mathematische Nachrichten* 137 (1) (1988), 197-248.
- [18] A. Guillin, W. Liu, L. Wu and C. Zhang, Poincaré and logarithmic Sobolev inequalities for particles in mean field interactions. https://arxiv.org/abs/1909.07051, 2019.
- [19] W. Hammersley, D. Siska, and L. Szpruch, McKean-Vlasov SDEs under measure dependent Lyapunov conditions. https://arxiv.org/abs/1802.03974, 2018.
- [20] M. Hauray and S. Mischler, On Kacs chaos and related problems. J. Funct. Anal. 266 (10) (2014), 6055-6157.
- [21] P. E. Jabin and Z. Wang, Quantitative estimates of propagation of chaos for stochastic systems with $W^{-1,\infty}$ kernels. *Invent. Math.* 214 (2018), 523-591.
- [22] D. Lacker, On a strong form of propagation of chaos for McKean-Vlasov equations. *Electron. Commun. Probab.* 23 (45) (2018), 1C11.
- [23] D. Luo and J. Wang, Exponential convergence in L^p-Wasserstein distance for diffusion processes without uniformly dissipative drift, Math. Nachr. 289 (2016), 1909C1926.
- [24] M. Kac, Foundations of kinetic theory. In Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, 1954-1955, vol. III, pages 171-197. University of California Press, Berkeley and Los Angeles, 1956.
- [25] F. Malrieu, Logarithmic sobolev inequalities for some nonlinear PDE's. Stoch. Proc. Appl. 95 (2001), 109-132.
- [26] F. Malrieu, Convergence to equilibrium for granular media equations and their Euler schemes. Ann. Appl. Probab. 13 (2) (2003), 540-560.
- [27] S. Méléard, Asymptotic behaviour of some interacting particle systems; McKean-Vlasov and Boltzmann models. In D. Talay and L. Tubaro, editors, Probabilistic Models for Nonlinear Partial Differential Equations, Montecatini Terme, 1995, Lecture Notes in Mathematics, vol. 1627, pp. 42-95, Springer-Verlag, 1996.
- [28] H.P. McKean, A class of Markov processes associated with nonlinear parabolic equations. Proceedings of the National Academy of Sciences 56 (6) (1966), 1907C1911.
- [29] S. Mischler and C. Mouhot, Kacs program in kinetic theory. Invent. Math. 193 (1) (2013), 1-147.
- [30] S. Mischler, C. Mouhot, and B. Wennberg, A new approach to quantitative propagation of chaos for drift, diffusion and jump processes. *Probab. Theory Relat. Fields* 161 (1) (2015), 1-59.
- [31] Y.S. Mishura and A.Y. Veretennikov, Existence and uniqueness theorems for solutions of McKean– Vlasov stochastic equations. https://arxiv.org/abs/1603.02212, 2016.
- [32] A.S. Sznitman, Topics in propagation of chaos. École d'Été de Probabilités de Saint-Flour XIX, Lecture Notes in Math., vol. 1464, pp. 165-251, 1991.
- [33] L. Wu, Gradient estimates of Poisson equations on Riemannian manifolds and applications. J. Funct. Anal. 257 (2009), 4015-4033.

WEI LIU, SCHOOL OF MATHEMATICS AND STATISTICS, WUHAN UNIVERSITY, WUHAN, HUBEI 430072, PR CHINA; HUBEI KEY LABORATORY OF COMPUTATIONAL SCIENCE, WUHAN UNIVERSITY, WUHAN, HUBEI 430072, PR CHINA.

E-mail address: wliu.math@whu.edu.cn

LIMING WU. LABORATOIRE DE MATHÉMATIQUES BLAISE PASCAL, CNRS-UMR 6620, UNIVER-SITÉ CLERMONT-AUVERGNE (UCA), CAMPUS UNIVERSITAIRE DES CEZEAUX, 3 PLACE VASARELY, 63178 AUBIÈRE, FRANCE.

E-mail address: Li-Ming.Wu@math.univ-bpclermont.fr

CHAOEN ZHANG. LABORATOIRE DE MATHÉMATIQUES BLAISE PASCAL, CNRS-UMR 6620, UNIVERSITÉ CLERMONT-AUVERGNE (UCA), CAMPUS UNIVERSITAIRE DES CEZEAUX, 3 PLACE VASARELY, 63178 AUBIÈRE, FRANCE.

E-mail address: chaoenz@gmail.com