

INSTABILITY INTERVALS OF THE WHITTAKER-HILL OPERATOR

XU-DAN LUO

ABSTRACT. The Hill operator admits a band gap structure. As a special case, like the Mathieu operator, one has only open gaps, however, the instability intervals of the Whittaker-Hill operator may be open or closed. In 2007, P. Djakov and B. Mityagin gave the asymptotics of band gaps for a special Whittaker-Hill operator [P. Djakov and B. Mityagin, *J. Funct. Anal.*, 242, 157-194 (2007)]. In this paper, a more general Whittaker-Hill operator is considered and the asymptotics of the instability intervals are studied.

1. INTRODUCTION AND MAIN RESULTS

The Floquet (Bloch) theory indicates that the spectrum of the Schrödinger operator

$$(1.1) \quad Lf := -f''(x) + \nu(x)f(x), \quad x \in \mathbb{R}$$

with a smooth real-valued periodic potential $\nu(x)$ has a band gap structure. If we further assume that $\nu(x)$ is of periodic π and set

$$\nu(x) = -\sum_{n=1}^{\infty} \theta_n \cos(2nx) - \sum_{m=1}^{\infty} \phi_m \sin(2mx),$$

where θ_n and ϕ_m are real, then (1.1) can be written as:

$$(1.2) \quad Lf := -f''(x) - \left[\sum_{n=1}^{\infty} \theta_n \cos(2nx) + \sum_{m=1}^{\infty} \phi_m \sin(2mx) \right] f(x).$$

Moreover, there are [3] two monotonically increasing infinite sequence of real numbers

$$\lambda_0^+, \lambda_1^+, \lambda_2^+, \dots$$

and

$$\lambda_1^-, \lambda_2^-, \lambda_3^-, \dots$$

such that the Hill equation

$$(1.3) \quad Lf = \lambda f$$

has a solution of period π if and only if $\lambda = \lambda_n^+$, $n = 0, 1, 2, \dots$, and a solution of semi-period π (i.e., $f(x + \pi) = -f(x)$) if and only if $\lambda = \lambda_n^-$, $n = 1, 2, 3, \dots$. The λ_n^+ and λ_n^- satisfy the inequalities

$$\lambda_0^+ < \lambda_1^- \leq \lambda_2^- \leq \lambda_1^+ \leq \lambda_2^+ < \lambda_3^- \leq \lambda_4^- < \lambda_3^+ \leq \lambda_4^+ < \dots$$

2010 *Mathematics Subject Classification.* Primary : 34L15; 41A60; 47E05.

Key words and phrases. The Whittaker-Hill operator; Instability intervals; Asymptotics.

and the relations

$$\lim_{n \rightarrow \infty} \lambda_n^+ = \infty, \quad \lim_{n \rightarrow \infty} \lambda_n^- = \infty.$$

Besides, $\gamma_n := (\lambda_{n+1}^- - \lambda_n^-)$ for odd n and $\gamma_n := (\lambda_n^+ - \lambda_{n-1}^+)$ for even n are referred to as band gaps or instability intervals, where $n \geq 1$.

It is well-known that there is an extensive theory for the Mathieu operator, where the potential $\nu(x)$ is a single trigonometric function, i.e.,

$$(1.4) \quad \nu(x) = -B \cos 2x.$$

Ince [5] proved that all instability intervals of the Mathieu operator are open, i.e., no closed gaps for the Mathieu operator. In 1963, Levy and Keller [6] gave the asymptotics of $\gamma_n = \gamma_n(B)$, i.e., for fixed n and real nonzero number B , when $B \rightarrow 0$,

$$(1.5) \quad \gamma_n = \frac{8}{[(n-1)!]^2} \cdot \left(\frac{B}{8}\right)^n (1 + O(B)).$$

18 years later, Harrell [4] gave the asymptotics of the band gaps of the Mathieu operator for fixed B and $n \rightarrow \infty$, i.e.,

$$(1.6) \quad \gamma_n = \lambda_n^+ - \lambda_n^- = \frac{8}{[(n-1)!]^2} \cdot \left(\frac{|B|}{8}\right)^n \left(1 + O\left(\frac{1}{n^2}\right)\right).$$

Compared with the Mathieu potential, the band gaps for the Whittaker-Hill potential

$$(1.7) \quad \nu(x) = -(B \cos 2x + C \cos 4x)$$

may be open or closed. Specifically, if $B = 4\alpha t$ and $C = 2\alpha^2$, for any real α and natural number t , it is already known that for odd $t = 2m + 1$, all the even gaps are closed except the first m , but no odd gap disappears; similarly, for even $t = 2m$, except for the first m , all the odd gaps are closed, but even gaps remain open (see Theorem 11, [1] and Theorem 7.9, [7]).

In 2007, P. Djakov and B. Mityagin (see [2]) gave the asymptotics of the instability intervals for the above special Whittaker-Hill potential, namely, for real $B, C \neq 0$, $B = 4\alpha t$ and $C = 2\alpha^2$, they have the following results, where either both α and t are real numbers if $C > 0$ or both α and t are pure imaginary numbers if $C < 0$.

Theorem 1.1 ([2]). *Let γ_n be the n -th band gap of the Whittaker-Hill operator*

$$(1.8) \quad Lf = -f'' - [4\alpha t \cos 2x + 2\alpha^2 \cos 4x]f,$$

where either both α and t are real, or both are pure imaginary numbers. If t is fixed and $\alpha \rightarrow 0$, then for even n

$$(1.9) \quad \gamma_n = \left| \frac{8\alpha^n}{2^n [(n-1)!]^2} \prod_{k=1}^{n/2} (t^2 - (2k-1)^2) \right| (1 + O(\alpha)),$$

and for odd n

$$(1.10) \quad \gamma_n = \left| \frac{8\alpha^{nt}}{2^n [(n-1)!]^2} \prod_{k=1}^{(n-1)/2} (t^2 - (2k)^2) \right| (1 + O(\alpha)).$$

Theorem 1.2 ([2]). *Let γ_n be the n -th band gap of the Whittaker-Hill operator*

$$(1.11) \quad Lf = -f'' - [4\alpha t \cos 2x + 2\alpha^2 \cos 4x]f,$$

where either both α and $t \neq 0$ are real, or both are pure imaginary numbers. Then the following asymptotic formulae hold for fixed α , t and $n \rightarrow \infty$:

for even n

$$(1.12) \quad \gamma_n = \frac{8|\alpha|^n}{2^n[(n-2)!!]^2} \left| \cos\left(\frac{\pi}{2}t\right) \right| \left[1 + O\left(\frac{\log n}{n}\right) \right],$$

and for odd n

$$(1.13) \quad \gamma_n = \frac{8|\alpha|^n}{2^n[(n-2)!!]^2} \frac{2}{\pi} \left| \sin\left(\frac{\pi}{2}t\right) \right| \left[1 + O\left(\frac{\log n}{n}\right) \right],$$

where $(2m-1)!! = 1 \cdot 3 \cdots (2m-1)$, $(2m)!! = 2 \cdot 4 \cdots (2m)$.

In this paper, a more general Whittaker-Hill operator

$$(1.14) \quad L = -D^2 + (bq^{m_1} \cos 2x + cq^{m_2} \cos 4x)$$

is considered and the asymptotics of the instability intervals are derived, where b , c , q , m_1 and m_2 are real. Our theorems are stated as follows, in particular, we can deduce P. Djakov and B. Mityagin's results by choosing $m_1 = 1$, $m_2 = 2$, $b = -4\alpha t$ and $c = -2\alpha^2$.

Theorem 1.3. *Let the Whittaker-Hill operator be*

$$(1.15) \quad Ly = -y'' + (bq^{m_1} \cos 2x + cq^{m_2} \cos 4x)y,$$

where b , c and q are real. If $q \rightarrow 0$ and m_1 , m_2 are positive real parameters, then one of the following results holds:

- (1) *when $m_1 > \frac{m_2}{2}$,*
(i)

$$(1.16) \quad \gamma_{2m} = \left| \frac{32 \cdot \left(\frac{c}{2}\right)^m \cdot q^{m_2 m}}{2^{4m}[(m-1)!!]^2} \right| + O\left(q^{m_2(m-\frac{1}{2})+m_1}\right),$$

(ii)

$$(1.17) \quad \gamma_1 = |bq^{m_1}| + O(q^{2m_1-\frac{m_2}{2}}), \quad \gamma_3 = \frac{|bcq^{m_1+m_2}|}{8} + O(q^{2m_1+\frac{m_2}{2}}),$$

(1.18)

$$\gamma_{2m-1} = \left| \left(\frac{c}{2}\right)^{m-1} \cdot b \cdot q^{m_2(m-1)+m_1} \cdot \frac{8}{2^{3m}} \cdot \left\{ \frac{1}{[(2m-3)!!]^2} \cdot \sum_{i=1}^{m-2} \frac{(2m-2i-3)!! \cdot (2i-1)!!}{i! \cdot (m-1-i)!} + \frac{2}{(2m-3)!! \cdot (m-1)!} \right\} \right| + O\left(q^{m_2(m-\frac{3}{2})+2m_1}\right) \quad \text{for } m \geq 3;$$

- (2) *when $m_1 < \frac{m_2}{2}$,*

$$(1.19) \quad \gamma_1 = |bq^{m_1}| + O(q^{m_2-m_1}), \quad \gamma_2 = \left| cq^{m_2} + \frac{b^2 q^{2m_1}}{8} \right| + O(q^{m_2}),$$

$$(1.20) \quad \gamma_n = \left| \frac{8 \cdot b^n \cdot q^{m_1 n}}{2^{3n} \cdot [(n-1)!!]^2} \right| + O(q^{m_1 n + m_2 - 2m_1}) \quad \text{for } n \geq 3;$$

(3) when $m_1 = \frac{m_2}{2}$ and $c < 0$,
(i)

$$(1.21) \quad \gamma_1 = |bq^{m_1}| + O(q^{3m_1}), \quad \gamma_2 = \left| cq^{m_2} + \frac{b^2 q^{2m_1}}{8} \right| + O(q^{4m_1}),$$

(ii)

$$(1.22) \quad \gamma_{2m} = 8 \left| \frac{\prod_{k=1}^m \left(\left(\frac{b}{2} \right)^2 + 8c \left(k - \frac{1}{2} \right)^2 \right) \cdot q^{2m_1 \cdot m}}{2^{4m} \cdot [(2m-1)!]^2} \right| + O(q^{2m_1(m+1)}) \quad \text{for } m \geq 2,$$

(iii)

$$(1.23) \quad \gamma_{2m-1} = 32 \left| \frac{\frac{b}{2} \prod_{k=1}^{m-1} \left(\left(\frac{b}{2} \right)^2 + 8ck^2 \right) \cdot q^{m_1 \cdot (2m-1)}}{2^{4m} \cdot [(2m-2)!]^2} \right| + O(q^{m_1(2m+1)}) \quad \text{for } m \geq 2.$$

Here, m is a positive integer and γ_n is the n -th instability interval.

Theorem 1.4. Let the Whittaker-Hill operator be

$$(1.24) \quad Ly = -y'' + (bq^{m_1} \cos 2x + cq^{2m_1} \cos 4x)y,$$

where b, q are real, $c < 0$ and $m_1 > 0$. Then the following asymptotic formula holds for fixed b, c, q and $n \rightarrow \infty$.

$$(1.25) \quad \gamma_{2m} = \frac{q^{2m_1 \cdot m} \cdot |c|^m}{2^{3m-3} \cdot [(2m-2)!!]^2} \cdot \left| \cos \left(\frac{b\pi}{4\sqrt{-2c}} \right) \right| \cdot \left[1 + O \left(\frac{\log m}{m} \right) \right],$$

$$(1.26) \quad \gamma_{2m-1} = \frac{q^{m_1(2m-1)} \cdot |c|^{m-1} \cdot \sqrt{-2c}}{2^{3m-5} \cdot [(2m-3)!!]^2 \cdot \pi} \cdot \left| \sin \left(\frac{b\pi}{4\sqrt{-2c}} \right) \right| \cdot \left[1 + O \left(\frac{\log m}{m} \right) \right].$$

Here, m is a positive integer and γ_n is the n -th instability interval.

2. SOME LEMMAS

Lemma 2.1 ([2]). Let the Schrödinger operator

$$(2.1) \quad Ly = -y'' + v(x)y$$

be defined on \mathbb{R} , with a real-valued periodic $L^2([0, \pi])$ -potential $v(x)$, where $v(x) = \sum_{m \in \mathbb{Z}} V(m) \exp(imx)$, $V(m) = 0$ for odd m , then $\|v\|^2 = \sum |V(m)|^2$.

(a) If $\|v\| < \frac{1}{9}$, then for each $n = 1, 2, \dots$, there exists $z = z_n$ such that $|z| \leq 4\|v\|$, and

$$(2.2) \quad 2|\beta_n(z)|(1 - 3\|v\|^2/n^2) \leq \gamma_n \leq 2|\beta_n(z)|(1 + 3\|v\|^2/n^2),$$

where

$$(2.3) \quad \beta_n(z) = V(2n) + \sum_{k=1}^{\infty} \sum_{j_1, \dots, j_k \neq \pm n} \frac{V(n-j_1)V(j_1-j_2) \cdots V(j_{k-1}-j_k)V(j_k+n)}{(n^2-j_1^2+z) \cdots (n^2-j_k^2+z)}$$

converges absolutely and uniformly for $|z| \leq 1$, and γ_n is the n -th instability interval.

(b) If $V(0) = \frac{1}{\pi} \int_0^\pi v(x) dx = 0$, then there is $N_0 = N_0(v)$ such that (2.2) holds for $n \geq N_0$ with $z = z_n$, $|z_n| < 1$.

Lemma 2.2 ([8]). *The Ince equation*

$$(2.4) \quad (1 + a \cos 2t)y''(t) + b(\sin 2t)y'(t) + (c + d \cos 2t)y(t) = 0$$

can be transformed into the Whittaker-Hill equation by assuming

$$(2.5) \quad a = 0, \quad b = -4q, \quad c = \lambda + 2q^2, \quad d = 4(m-1)q.$$

Moreover,

$$(2.6) \quad \text{sign}(\alpha_{2n} - \beta_{2n}) = \text{sign } q^2 \cdot \text{sign} \prod_{p=-n}^{n-1} (2p - m + 1)$$

and

$$(2.7) \quad \text{sign}(\alpha_{2n+1} - \beta_{2n+1}) = \text{sign } q \cdot \text{sign} \prod_{p=-n}^{n-1} (2p - m),$$

where a , b and d are real; α_{2n} and β_{2n+2} are defined by the eigenvalues corresponding to non-trivial even and odd solutions with period π , respectively; and α_{2n+1} and β_{2n+1} are defined by the eigenvalues corresponding to non-trivial even and odd solutions with semi-period π , respectively.

Lemma 2.3 ([7]). *The Whittaker-Hill equation*

$$(2.8) \quad f'' + [\lambda + 4mq \cos 2x + 2q^2 \cos 4x]f = 0$$

can have two linearly independent solutions of period π or 2π if and only if m is an integer. If $m = 2l$ is even, then the odd intervals of instability on the λ axis disappear, with at most $|l| + 1$ exceptions, but no even interval of instability disappears. If $m = 2l + 1$ is odd, then at most $|l| + 1$ even intervals of instability remain but no interval of instability disappears.

Lemma 2.4. *The Whittaker-Hill operator*

$$(2.9) \quad L = -D^2 - (B \cos 2x + C \cos 4x)$$

admits all even gaps closed except the first n when $\pm \frac{B}{4\sqrt{2C}} = -n - \frac{1}{2}$, $n \in \mathbb{Z}_{\geq 0}$; and all odd gaps closed except the first $n + 1$ when $\pm \frac{B}{4\sqrt{2C}} = -n - 1$, $n \in \mathbb{Z}_{\geq 0}$.

Proof. By Lemma 2.3, we obtain that the Whittaker-Hill equation

$$(2.10) \quad f''(x) + (A + B \cos 2x + C \cos 4x)f(x) = 0$$

have two linearly independent solutions of period or semi-periodic π if and only if $\frac{B}{2\sqrt{2C}} \in \mathbb{Z}$. Moreover, we transform (2.10) into the Ince equation

$$(2.11) \quad g''(x) + 4\sqrt{\frac{C}{2}} \sin 2x \cdot g'(x) + \left[(A + C) + \left(B + 4\sqrt{\frac{C}{2}} \right) \cos 2x \right] g(x) = 0.$$

via $f(x) = e^{-\sqrt{\frac{C}{2}} \cos 2x} \cdot g(x)$. From Lemma 2.2, we can write the parameters q , λ and m of equation (2.4) in terms of A , B and C , i.e.,

$$q = -\sqrt{\frac{C}{2}}, \quad \lambda = A, \quad m = -\frac{B}{2\sqrt{2C}}.$$

(1) If $m = 2n + 1$, $n \in \mathbb{N}^+ \cup \{0\}$, i.e., $\frac{B}{2\sqrt{2C}} = -2n - 1$, and the solutions satisfy the periodic boundary conditions, then we deduce from Lemma 2.2 that the first $2n + 1$ eigenvalues are simple, and others are double.

(2) If $m = 2n + 2$, $n \in \mathbb{N}^+ \cup \{0\}$, i.e., $\frac{B}{2\sqrt{2C}} = -2n - 2$, and the solutions satisfy the semi-periodic boundary conditions, then we also deduce from Lemma 2.2 that the first $2n + 2$ eigenvalues are simple, and others are double.

Besides, we can also transform (2.10) into the Ince equation

$$(2.12) \quad g''(x) - 4\sqrt{\frac{C}{2}} \sin 2x \cdot g'(x) + \left[(A + C) + \left(B - 4\sqrt{\frac{C}{2}} \right) \cos 2x \right] g(x) = 0.$$

via $f(x) = e^{\sqrt{\frac{C}{2}} \cos 2x} \cdot g(x)$. Thus, we have similar conclusions. \square

In order to prove our results, let us consider all possible walks from $-n$ to n . Each such walk is determined by the sequence of its steps

$$(2.13) \quad x = (x_1, \dots, x_{\nu+1}),$$

or by its vertices

$$(2.14) \quad j_s = -n + \sum_{k=1}^s x_k, \quad s = 1, \dots, \nu.$$

The relationship between steps and vertices are given by the formula

$$(2.15) \quad x_1 = n + j_1; \quad x_k = j_k - j_{k-1}, \quad k = 2, \dots, \nu; \quad x_{\nu+1} = n - j_{\nu}.$$

Definition 2.5. Let X denote the set of all walks from $-n$ to n with steps ± 2 or ± 4 . For each $x = (x_s)_{s=1}^{\nu+1} \in X$ and each $z \in \mathbb{R}$, we define

$$(2.16) \quad B_n(x, z) = \frac{V(x_1) \cdots V(x_{\nu+1})}{(n^2 - j_1^2 + z) \cdots (n^2 - j_{\nu}^2 + z)}.$$

Definition 2.6. Let X^+ denote the set of all walks from $-n$ to n with positive steps equal to 2 or 4. For each $\xi \in X^+$, let X_{ξ} denote the set of all walks $x \in X \setminus X^+$ such that each vertex of ξ is a vertex of x also. For each $\xi \in X^+$ and $\mu \in \mathbb{N}$, let $X_{\xi, \mu}$ be the set of all $x \in X_{\xi}$ such that x has μ more vertices than ξ . Moreover, for each μ -tuple (i_1, \dots, i_{μ}) of integers in $I_n = (n + 2\mathbb{Z}) \setminus \{\pm n\}$, we define $X_{\xi}(i_1, \dots, i_{\mu})$ as the set of all walks x with $\nu + 1 + \mu$ steps such that (i_1, \dots, i_{μ}) and the sequence of the vertices of ξ are complementary subsequences of the sequence of the vertices of x .

From Definition 2.6, we deduce

$$(2.17) \quad X_{\xi, \mu} = \bigcup_{(i_1, \dots, i_{\mu}) \in (I_n)^{\mu}} X_{\xi}(i_1, \dots, i_{\mu}).$$

Lemma 2.7 ([2]). *If $\xi \in X^+$ and $n \geq 3$, then for $z \in [0, 1)$*

$$(2.18) \quad 1 - z \frac{\log n}{n} \leq \frac{B_n(\xi, z)}{B_n(\xi, 0)} \leq 1 - z \frac{\log n}{4n},$$

and for $z \in (-1, 0]$

$$(2.19) \quad 1 + |z| \frac{\log n}{2n} \leq \frac{B_n(\xi, z)}{B_n(\xi, 0)} \leq 1 + |z| \frac{2 \log n}{n}.$$

Lemma 2.8 ([2]). *For each walk $\xi \in X^+$ and each μ -tuple $(i_1, \dots, i_{\mu}) \in (I_n)^{\mu}$,*

$$(2.20) \quad \#X_{\xi}(i_1, \dots, i_{\mu}) \leq 5^{\mu}.$$

Lemma 2.9. *If $\xi \in X^+$ and $|z| \leq 1$, then there exists n_1 such that for $n \geq n_1$,*

$$(2.21) \quad \sum_{x \in X_\xi} |B_n(x, z)| \leq |B_n(\xi, z)| \cdot \frac{K \log n}{n},$$

where $K = 40 \left(\left| \frac{b}{2} q^{m_1} \right| + \left| \frac{c}{2} q^{m_2} \right| + \left| \frac{b^2}{2c} q^{2m_1 - m_2} \right| \right)$.

Proof. By Definition 2.6, we have

$$(2.22) \quad \sum_{x \in X_\xi} |B_n(x, z)| = \sum_{\mu=1}^{\infty} \sum_{x \in X_{\xi, \mu}} |B_n(x, z)|.$$

Moreover,

$$(2.23) \quad \sum_{x \in X_{\xi, \mu}} |B_n(x, z)| \leq \sum_{(i_1, \dots, i_\mu)} \sum_{X_\xi(i_1, \dots, i_\mu)} |B_n(x, z)|,$$

where the first sum on the right is taken over all μ -tuples (i_1, \dots, i_μ) of integers $i_s \in n + 2\mathbb{Z}$ such that $i_s \neq \pm n$.

Fix (i_1, \dots, i_μ) , if $x \in X_\xi(i_1, \dots, i_\mu)$, then

$$(2.24) \quad \frac{B_n(x, z)}{B_n(\xi, z)} = \frac{\prod_k V(x_k)}{\prod_s V(\xi_s)} \cdot \frac{1}{(n^2 - i_1^2 + z) \cdots (n^2 - i_\mu^2 + z)}.$$

Note that $V(\pm 2) = \frac{b}{2} q^{m_1}$ and $V(\pm 4) = \frac{c}{2} q^{m_2}$. If each step of ξ is a step of x , then

$$(2.25) \quad \frac{\prod_k |V(x_k)|}{\prod_s |V(\xi_s)|} \leq C^\mu,$$

where $C := \left| \frac{b}{2} q^{m_1} \right| + \left| \frac{c}{2} q^{m_2} \right| + \left| \frac{b^2}{2c} q^{2m_1 - m_2} \right|$. For the general case, let $(j_s)_{s=1}^\nu$ be the vertices of ξ , and let us put for convenience $j_0 = -n$ and $j_{\nu+1} = n$. Since each vertex of ξ is a vertex of x , for each s , $1 \leq s \leq \nu + 1$,

$$(2.26) \quad \xi_s = j_s - j_{s-1} = \sum_{k \in J_s} x_k,$$

where x_k , $k \in J_s$, are the steps of x between the vertices j_{s-1} and j_s . Fix an s , $1 \leq s \leq \nu + 1$. If $\xi_s = 2$, then there is a step x_{k^*} , $k^* \in J_s$ such that $|x_{k^*}| = 2$, otherwise, ξ_s would be a multiple of 4. Hence, $|V(\xi_s)| = |V(x_{k^*})|$, which implies that

$$(2.27) \quad \frac{\prod_{J_s} |V(x_k)|}{|V(\xi_s)|} \leq C^{b_s - 1},$$

where b_s is the cardinality of J_s .

If $\xi_s = 4$, there are two possibilities. (1) If there is $k_* \in J_s$ with $|x_{k_*}| = 4$, then $|V(\xi_s)| = |V(x_{k_*})|$, so the above inequality holds. (2) There are $k', k'' \in J_s$ such that $|x_{k'}| = |x_{k''}| = 2$, hence,

$$(2.28) \quad \frac{|V(x_{k'})V(x_{k''})|}{|V(\xi_s)|} = \left| \frac{b^2}{2c} q^{2m_1 - m_2} \right| \leq C,$$

so the above inequality also holds. Note that

$$(2.29) \quad \sum_s (b_s - 1) = \mu,$$

we get

$$(2.30) \quad \frac{\prod_k V(x_k)}{\prod_s V(\xi_s)} \leq C^\mu$$

holds for the general case.

By

$$(2.31) \quad \frac{1}{|n^2 - i^2 + z|} \leq \frac{2}{|n^2 - i^2|},$$

where $i \neq \pm n$, $|z| \leq 1$, we have

$$(2.32) \quad \frac{|B_n(x, z)|}{|B_n(\xi, z)|} \leq \frac{(2C)^\mu}{|n^2 - i_1^2| \cdots |n^2 - i_\mu^2|}, \quad x \in X_\xi(i_1, \dots, i_\mu).$$

By Lemma 2.8, we derive

$$(2.33) \quad \sum_{x \in X_\xi(i_1, \dots, i_\mu)} \frac{|B_n(x, z)|}{|B_n(\xi, z)|} \leq \frac{(10C)^\mu}{|n^2 - i_1^2| \cdots |n^2 - i_\mu^2|}.$$

Combining Lemma 2.7, it yields

$$(2.34) \quad \begin{aligned} \sum_{x \in X_{\xi, \mu}} \frac{|B_n(x, z)|}{|B_n(\xi, z)|} &\leq \sum_{(i_1, \dots, i_\mu)} \frac{(10C)^\mu}{|n^2 - i_1^2| \cdots |n^2 - i_\mu^2|} \leq \left(\sum_{i \in (n+2\mathbb{Z}) \setminus \{\pm n\}} \frac{10C}{|n^2 - i^2|} \right)^\mu \\ &\leq (10C)^\mu \left(\frac{1 + \log n}{n} \right)^\mu \leq \left(\frac{20C \log n}{n} \right)^\mu. \end{aligned}$$

Thus,

$$(2.35) \quad \sum_{x \in X_{\xi, \mu}} |B_n(x, z)| \leq |B_n(\xi, z)| \cdot \left(\frac{20C \log n}{n} \right)^\mu.$$

Hence,

$$(2.36) \quad \sum_{x \in X_\xi} \frac{|B_n(x, z)|}{|B_n(\xi, z)|} \leq \sum_{\mu=1}^{\infty} \left(\frac{20C \log n}{n} \right)^\mu.$$

We can choose $n_1 \in \mathbb{N}^+$ such that $\frac{20C \log n}{n} \leq \frac{1}{2}$ for $n \geq n_1$. Then

$$(2.37) \quad \sum_{x \in X_\xi} \frac{|B_n(x, z)|}{|B_n(\xi, z)|} \leq \frac{40C \log n}{n}.$$

Therefore, there exists n_1 such that for $n \geq n_1$,

$$(2.38) \quad \sum_{x \in X_\xi} |B_n(x, z)| \leq |B_n(\xi, z)| \cdot \frac{K \log n}{n},$$

where $K := 40C = 40 \left(\left| \frac{b}{2} q^{m_1} \right| + \left| \frac{c}{2} q^{m_2} \right| + \left| \frac{b^2}{2c} q^{2m_1 - m_2} \right| \right)$. □

3. PROOF OF THEOREM 1.3

Note that

$$(3.1) \quad V(\pm 2) = \frac{bq^{m_1}}{2}, \quad V(\pm 4) = \frac{cq^{m_2}}{2}$$

and

$$(3.2) \quad \|v\|^2 = \frac{1}{2} \left(b^2 q^{2m_1} + c^2 q^{2m_2} \right),$$

by Lemma 2.1, we have

$$(3.3) \quad \gamma_n = \pm 2 \left(V(2n) + \sum_{k=1}^{\infty} \beta_k(n, z) \right) \left(1 + O(q^{2 \cdot \min\{m_1, m_2\}}) \right),$$

where

$$(3.4) \quad \begin{aligned} \beta_k(n, z) &= \sum_{j_1, \dots, j_k \neq \pm n} \frac{V(n - j_1)V(j_1 - j_2) \cdots V(j_{k-1} - j_k)V(j_k + n)}{(n^2 - j_1^2 + z) \cdots (n^2 - j_k^2 + z)} \\ &= \sum_{j_1, \dots, j_k \neq \pm n} \frac{V(n + j_1)V(j_2 - j_1) \cdots V(j_k - j_{k-1})V(n - j_k)}{(n^2 - j_1^2 + z) \cdots (n^2 - j_k^2 + z)} \end{aligned}$$

and $z = O(q)$. Moreover, all series converge absolutely and uniformly for sufficiently small q .

Note that

$$(3.5) \quad (n + j_1) + (j_2 - j_1) + \cdots + (j_k - j_{k-1}) + (n - j_k) = 2n,$$

and

$$(3.6) \quad \frac{V(n + j_1)V(j_2 - j_1) \cdots V(j_k - j_{k-1})V(n - j_k)}{(n^2 - j_1^2 + z) \cdots (n^2 - j_k^2 + z)} \neq 0$$

when

$$(3.7) \quad (n + j_1), (j_2 - j_1), \dots, (j_k - j_{k-1}), (n - j_k) \in \{\pm 2, \pm 4\}.$$

We distinguish three cases to discuss.

Case 1. If $m_1 > \frac{m_2}{2}$, then

$$(3.8) \quad V(n + j_1)V(j_2 - j_1) \cdots V(j_k - j_{k-1})V(n - j_k)$$

is a monomial in q of degree at least

$$(3.9) \quad \frac{m_2}{4} \cdot \left[|n + j_1| + |j_2 - j_1| + \cdots + |j_k - j_{k-1}| + |n - j_k| \right].$$

The minimum case occurs when

$$(3.10) \quad (n + j_1), (j_2 - j_1), \dots, (j_k - j_{k-1}), (n - j_k) \in \{\pm 4\},$$

then

$$(3.11) \quad (n + j_1) + (j_2 - j_1) + \cdots + (j_k - j_{k-1}) + (n - j_k) \in 4\mathbb{Z},$$

while

$$(3.12) \quad (n + j_1) + (j_2 - j_1) + \cdots + (j_k - j_{k-1}) + (n - j_k) = 2n.$$

If n is even, i.e., $n = 2m$, $m \in \mathbb{Z}_{>0}$, since

$$(3.13) \quad \begin{aligned} & |n + j_1| + |j_2 - j_1| + \cdots + |j_k - j_{k-1}| + |n - j_k| \\ & \geq (n + j_1) + (j_2 - j_1) + \cdots + (j_k - j_{k-1}) + (n - j_k) \\ & = 2n = 4m, \end{aligned}$$

we obtain

$$(3.14) \quad V(n + j_1)V(j_2 - j_1) \cdots V(j_k - j_{k-1})V(n - j_k)$$

is a monomial in q of degree at least $m_2 \cdot m$. Such monomial in q of degree $m_2 \cdot m$ corresponds to a walk from $-n$ to n with vertices $j_1, j_2, \dots, j_k \neq \pm n$ and positive steps of length 4. Thus,

$$(3.15) \quad \gamma_{2m} = \pm P_{2m}(t)q^{m_2 m} + O\left(q^{m_2(m-\frac{1}{2})+m_1}\right),$$

where

$$(3.16) \quad P_{2m}(t)q^{m_2 m} = 2\left(V(4m) + \sum_{k=1}^{\infty} \beta_k(2m, z)\right).$$

We have

$$(3.17) \quad P_2(t)q^{m_2} = 2V(4) = cq^{m_2}$$

and

$$(3.18) \quad \begin{aligned} P_{2m}(t)q^{m_2 m} &= 2 \sum_{k=1}^{\infty} \beta_k(2m, z) \\ &= 2 \cdot \left(\frac{c}{2}\right)^m \cdot q^{m_2 m} \cdot \prod_{j=1}^{m-1} \left((4m^2 - (-2m + 4j)^2)\right)^{-1} \\ &= \frac{32 \cdot \left(\frac{c}{2}\right)^m \cdot q^{m_2 m}}{2^{4m} [(m-1)!]^2} \end{aligned}$$

for $m \geq 2$. Therefore,

$$(3.19) \quad \gamma_{2m} = \left| \frac{32 \cdot \left(\frac{c}{2}\right)^m \cdot q^{m_2 m}}{2^{4m} [(m-1)!]^2} \right| + O\left(q^{m_2(m-\frac{1}{2})+m_1}\right).$$

If n is odd, i.e., $n = 2m - 1$, $m \in \mathbb{Z}_{>0}$, since

$$(3.20) \quad (n + j_1) + (j_2 - j_1) + \cdots + (j_k - j_{k-1}) + (n - j_k) = 2n = 4m - 2,$$

then

$$(3.21) \quad V(n + j_1)V(j_2 - j_1) \cdots V(j_k - j_{k-1})V(n - j_k)$$

is a monomial in q of degree at least

$$(3.22) \quad \begin{aligned} & \frac{m_2}{4} \cdot \left[|n + j_1| + |j_2 - j_1| + \cdots + |j_k - j_{k-1}| + |n - j_k| \right] + m_1 - \frac{m_2}{2} \\ & \geq \frac{m_2}{4} \cdot (4m - 2) + m_1 - \frac{m_2}{2} \\ & = m_2(m - 1) + m_1. \end{aligned}$$

Such monomial in q of degree $m_2(m-1) + m_1$ corresponds to a walk from $-n$ to n with vertices $j_1, j_2, \dots, j_k \neq \pm n$ and positive steps. Specifically, except for one step with length 2, the others are of length 4. Thus,

$$(3.23) \quad \gamma_{2m-1} = \pm P_{2m-1}(t)q^{m_2(m-1)+m_1} + O\left(q^{m_2(m-\frac{3}{2})+2m_1}\right),$$

where

$$(3.24) \quad P_{2m-1}(t)q^{m_2(m-1)+m_1} = 2\left(V(4m-2) + \sum_{k=1}^{\infty} \beta_k(2m-1, z)\right).$$

We obtain

$$(3.25) \quad P_1(t)q^{m_1} = 2V(2) = bq^{m_1},$$

$$(3.26) \quad P_3(t)q^{m_1+m_2} = 2 \sum_{k=1}^{\infty} \beta_k(3, z) = 2 \left(\frac{bq^{m_1}}{2}\right) \left(\frac{cq^{m_2}}{2}\right) \left(\frac{1}{3^2-1^2} + \frac{1}{3^2-1^2}\right) = \frac{bcq^{m_1+m_2}}{8},$$

$$(3.27) \quad P_5(t)q^{m_1+2m_2} = 2 \sum_{k=1}^{\infty} \beta_k(5, z) \\ = 2 \left(\frac{bq^{m_1}}{2}\right) \left(\frac{cq^{m_2}}{2}\right)^2 \left[\frac{1}{(5^2-3^2)(5^2-1^2)} + \frac{1}{(5^2-1^2)(5^2-1^2)} + \frac{1}{(5^2-3^2)(5^2-1^2)}\right] \\ = \frac{bc^2q^{m_1+2m_2}}{3^2 \cdot 2^6},$$

$$(3.28) \quad P_{2m-1}(t)q^{m_2(m-1)+m_1} = 2 \sum_{k=1}^{\infty} \beta_k(2m-1, z) \\ = 2 \left(\frac{c}{2}\right)^{m-1} \cdot \left(\frac{b}{2}\right) \cdot q^{m_2(m-1)+m_1} \cdot \left\{ \sum_{i=1}^{m-2} \prod_{j=1}^i \left[(2m-1)^2 - (-2m+1+4j)^2\right]^{-1} \right. \\ \cdot \prod_{j=i}^{m-2} \left[(2m-1)^2 - (-2m+3+4j)^2\right]^{-1} + \prod_{j=0}^{m-2} \left[(2m-1)^2 - (-2m+3+4j)^2\right]^{-1} \\ \left. + \prod_{j=1}^{m-1} \left[(2m-1)^2 - (-2m+1+4j)^2\right]^{-1} \right\} \\ = 2 \left(\frac{c}{2}\right)^{m-1} \cdot \left(\frac{b}{2}\right) \cdot q^{m_2(m-1)+m_1} \cdot \frac{8}{2^{3m}} \cdot \left\{ \frac{1}{[(2m-3)!!]^2} \cdot \sum_{i=1}^{m-2} \frac{(2m-2i-3)!! \cdot (2i-1)!!}{i! \cdot (m-1-i)!} \right. \\ \left. + \frac{2}{(2m-3)!! \cdot (m-1)!} \right\}.$$

Hence,

$$(3.29) \quad \gamma_1 = |bq^{m_1}| + O(q^{2m_1 - \frac{m_2}{2}}), \quad \gamma_3 = \frac{|bcq^{m_1+m_2}|}{8} + O(q^{2m_1 + \frac{m_2}{2}}),$$

(3.30)

$$\begin{aligned} \gamma_{2m-1} = & \left| \left(\frac{c}{2}\right)^{m-1} \cdot b \cdot q^{m_2(m-1)+m_1} \cdot \frac{8}{2^{3m}} \cdot \left\{ \frac{1}{[(2m-3)!!]^2} \right. \right. \\ & \left. \left. \cdot \sum_{i=1}^{m-2} \frac{(2m-2i-3)!! \cdot (2i-1)!!}{i! \cdot (m-1-i)!} + \frac{2}{(2m-3)!! \cdot (m-1)!} \right\} \right| + O\left(q^{m_2(m-\frac{3}{2})+2m_1}\right) \end{aligned}$$

for $m \geq 3$.**Case 2.** If $m_1 < \frac{m_2}{2}$, then

(3.31)
$$V(n+j_1)V(j_2-j_1)\cdots V(j_k-j_{k-1})V(n-j_k)$$

is a monomial in q of degree at least

(3.32)
$$\frac{m_1}{2} \cdot \left[|n+j_1| + |j_2-j_1| + \cdots + |j_k-j_{k-1}| + |n-j_k| \right].$$

The minimum case occurs when

(3.33)
$$(n+j_1), (j_2-j_1), \cdots, (j_k-j_{k-1}), (n-j_k) \in \{\pm 2\},$$

then

(3.34)
$$(n+j_1) + (j_2-j_1) + \cdots + (j_k-j_{k-1}) + (n-j_k) \in 2\mathbb{Z},$$

while

(3.35)
$$(n+j_1) + (j_2-j_1) + \cdots + (j_k-j_{k-1}) + (n-j_k) = 2n.$$

Since

(3.36)
$$\begin{aligned} & |n+j_1| + |j_2-j_1| + \cdots + |j_k-j_{k-1}| + |n-j_k| \\ & \geq (n+j_1) + (j_2-j_1) + \cdots + (j_k-j_{k-1}) + (n-j_k) \\ & = 2n, \end{aligned}$$

we have

(3.37)
$$V(n+j_1)V(j_2-j_1)\cdots V(j_k-j_{k-1})V(n-j_k)$$

is a monomial in q of degree at least $m_1 \cdot n$. Such monomial in q of degree $m_1 \cdot n$ corresponds to a walk from $-n$ to n with vertices $j_1, j_2, \cdots, j_k \neq \pm n$ and positive steps of length 2. Thus,

(3.38)
$$\gamma_n = \pm P_n(t)q^{m_1 n} + O(q^{m_1 n + m_2 - 2m_1}),$$

where

(3.39)
$$P_n(t)q^{m_1 n} = 2 \left(V(2n) + \sum_{k=1}^{\infty} \beta_k(n, z) \right).$$

We deduce

(3.40)

$$P_1(t)q^{m_1} = 2V(2) = bq^{m_1}, \quad P_2(t)q^{2m_1} = 2 \left(V(4) + \frac{\left(\frac{bq^{m_1}}{2}\right)^2}{2^2} \right) = cq^{m_2} + \frac{b^2 q^{2m_1}}{8},$$

$$\begin{aligned}
P_n(t)q^{m_1 n} &= 2 \sum_{k=1}^{\infty} \beta_k(n, z) \\
&= 2 \cdot \left(\frac{b}{2}\right)^n \cdot q^{m_1 \cdot n} \cdot \prod_{j=1}^{n-1} (n^2 - (-n + 2j)^2)^{-1} \\
(3.41) \quad &= 2 \cdot \left(\frac{b}{2}\right)^n \cdot q^{m_1 \cdot n} \cdot \prod_{j=1}^{n-1} (n^2 - (-n + 2j)^2)^{-1} \\
&= 2 \cdot \frac{\left(\frac{b}{2}\right)^n \cdot q^{m_1 \cdot n}}{4^{n-1} \cdot [(n-1)!]^2} \\
&= \frac{8 \cdot b^n \cdot q^{m_1 n}}{2^{3n} \cdot [(n-1)!]^2}
\end{aligned}$$

for $n \geq 3$. Therefore,

$$(3.42) \quad \gamma_1 = |bq^{m_1}| + O(q^{m_2 - m_1}), \quad \gamma_2 = \left| cq^{m_2} + \frac{b^2 q^{2m_1}}{8} \right| + O(q^{m_2}),$$

$$(3.43) \quad \gamma_n = \left| \frac{8 \cdot b^n \cdot q^{m_1 n}}{2^{3n} \cdot [(n-1)!]^2} \right| + O(q^{m_1 n + m_2 - 2m_1})$$

for $n \geq 3$.

Case 3. If $m_1 = \frac{m_2}{2}$, then

$$(3.44) \quad V(n + j_1)V(j_2 - j_1) \cdots V(j_k - j_{k-1})V(n - j_k)$$

is a monomial in q of degree

$$(3.45) \quad \frac{m_1}{2} \cdot \left[|n + j_1| + |j_2 - j_1| + \cdots + |j_k - j_{k-1}| + |n - j_k| \right].$$

Since

$$\begin{aligned}
(3.46) \quad &|n + j_1| + |j_2 - j_1| + \cdots + |j_k - j_{k-1}| + |n - j_k| \\
&\geq (n + j_1) + (j_2 - j_1) + \cdots + (j_k - j_{k-1}) + (n - j_k) \\
&= 2n,
\end{aligned}$$

we have

$$(3.47) \quad V(n + j_1)V(j_2 - j_1) \cdots V(j_k - j_{k-1})V(n - j_k)$$

is a monomial in q of degree at least $m_1 \cdot n$, and each such monomial of degree $m_1 \cdot n$ corresponds to a walk from $-n$ to n with vertices $j_1, j_2, \dots, j_k \neq \pm n$ and positive steps of length 2 or 4. The minimum case occurs when $n + j_1, j_2 - j_1, \dots, j_k - j_{k-1}$ and $n - j_k$ are of the same sign, while the second smallest degree is for one step of length 2 with opposite sign. Thus,

$$(3.48) \quad \gamma_n = \pm P_n(t)q^{m_1 n} + O(q^{m_1(n+2)}),$$

where

$$(3.49) \quad P_n(t)q^{m_1 n} = 2 \left(V(2n) + \sum_{k=1}^{\infty} \beta_k(n, z) \right).$$

We obtain

(3.50)

$$P_1(t)q^{m_1} = 2V(2) = bq^{m_1}, \quad P_2(t)q^{2m_1} = 2 \left(V(4) + \frac{\left(\frac{bq^{m_1}}{2}\right)^2}{2^2} \right) = cq^{m_2} + \frac{b^2q^{2m_1}}{8},$$

$$\begin{aligned} P_n(t)q^{m_1n} &= 2 \sum_{k=1}^{\infty} \beta_k(n, z) \\ &= 2 \cdot P_n\left(\frac{b}{2}\right) \cdot q^{m_1 \cdot n} \cdot \prod_{j=1}^{n-1} (n^2 - (-n + 2j)^2)^{-1} \\ (3.51) \quad &= 2 \cdot P_n\left(\frac{b}{2}\right) \cdot q^{m_1 \cdot n} \cdot \prod_{j=1}^{n-1} (n^2 - (-n + 2j)^2)^{-1} \\ &= 8 \cdot \frac{P_n\left(\frac{b}{2}\right) \cdot q^{m_1 \cdot n}}{2^{2n} \cdot [(n-1)!]^2} \end{aligned}$$

for $n \geq 3$. Therefore,

$$(3.52) \quad \gamma_1 = |bq^{m_1}| + O(q^{3m_1}), \quad \gamma_2 = \left| cq^{m_2} + \frac{b^2q^{2m_1}}{8} \right| + O(q^{4m_1}),$$

$$(3.53) \quad \gamma_n = \left| 8 \cdot \frac{P_n\left(\frac{b}{2}\right) \cdot q^{m_1 \cdot n}}{2^{2n} \cdot [(n-1)!]^2} \right| + O(q^{m_1(n+2)})$$

for $n \geq 3$, where $P_n\left(\frac{b}{2}\right)$ is a polynomial of $\frac{b}{2}$ with degree n and leading coefficient 1.

Specifically, if n is even, i.e., $n = 2m$, $m \in \mathbb{Z}_{>0}$, then

$$(3.54) \quad (n + j_1) + (j_2 - j_1) + \cdots + (j_k - j_{k-1}) + (n - j_k) = 4m,$$

which implies that each walk from $-2m$ to $2m$ has even number of steps with length 2. We have

$$(3.55) \quad P_{2m}\left(\frac{b}{2}\right) = \prod_{k=1}^m \left(\left(\frac{b}{2}\right)^2 - x_k \right),$$

where x_k , $k = 1, \dots, m$, depend on m . By Lemma 2.4, we obtain all even gaps closed except the first k if $\left(\frac{b}{2}\right)^2 = -8c\left(k + \frac{1}{2}\right)^2$, which yields

$$(3.56) \quad P_{2m}\left(\frac{b}{2}\right) = \prod_{k=1}^m \left(\left(\frac{b}{2}\right)^2 + 8c\left(k - \frac{1}{2}\right)^2 \right).$$

Hence,

$$(3.57) \quad \gamma_{2m} = 8 \left| \frac{\prod_{k=1}^m \left(\left(\frac{b}{2}\right)^2 + 8c\left(k - \frac{1}{2}\right)^2 \right) \cdot q^{2m_1 \cdot m}}{2^{4m} \cdot [(2m-1)!]^2} \right| + O(q^{2m_1(m+1)})$$

for $m \geq 2$. If n is odd, i.e., $n = 2m - 1$, $m \in \mathbb{Z}_{>0}$, then

$$(3.58) \quad (n + j_1) + (j_2 - j_1) + \cdots + (j_k - j_{k-1}) + (n - j_k) = 2n = 4m - 2,$$

which implies that each walk from $-2m$ to $2m$ has odd number of steps with length 2. We have

$$(3.59) \quad P_{2m-1}\left(\frac{b}{2}\right) = \frac{b}{2} \prod_{k=1}^{m-1} \left(\left(\frac{b}{2}\right)^2 - y_k \right),$$

where y_k , $k = 1, \dots, m-1$, depend on m . By Lemma 2.4, we deduce

$$(3.60) \quad P_{2m-1}\left(\frac{b}{2}\right) = \frac{b}{2} \prod_{k=1}^{m-1} \left(\left(\frac{b}{2}\right)^2 + 8ck^2 \right).$$

Hence,

$$(3.61) \quad \gamma_{2m-1} = 32 \left| \frac{\frac{b}{2} \prod_{k=1}^{m-1} \left(\left(\frac{b}{2}\right)^2 + 8ck^2 \right) \cdot q^{m_1 \cdot (2m-1)}}{2^{4m} \cdot [(2m-2)!]^2} \right| + O(q^{m_1(2m+1)})$$

for $m \geq 2$.

4. PROOF OF THEOREM 1.4

Since $V(\pm 2) = \frac{b}{2}q^{m_1}$ and $V(\pm 4) = \frac{c}{2}q^{2m_1}$, thus,

$$(4.1) \quad \|v\|^2 = \frac{1}{2} \left(b^2 q^{2m_1} + c^2 q^{4m_1} \right).$$

By Lemma 2.1, we get

$$(4.2) \quad \gamma_n = 2 \left| \sum_{x \in X} B_n(x, z) \right| \left(1 + O\left(\frac{1}{n^2}\right) \right),$$

where $z = z_n$ depends on n , but $|z| < 1$.

Set $\sigma_n = \sum_{\xi \in X^+} B_n(\xi, 0) := \sigma_n^+ + \sigma_n^-$, where $\sigma_n^\pm := \sum_{\xi: B_n(\xi, 0) \gtrless 0} B_n(\xi, 0)$. When $\xi \in X^+$,

$$(4.3) \quad B_n(\xi, 0) = \frac{V(x_1) \cdots V(x_{\nu+1})}{(n^2 - j_1^2) \cdots (n^2 - j_\nu^2)},$$

where $x_i = 2$ or 4 for $i = 1, \dots, \nu + 1$.

Note that $X \setminus X^+ = \bigcup_{\xi \in X^+} X_\xi$, we choose disjoint sets $X'_\xi \subset X_\xi$ so that

$$(4.4) \quad X \setminus X^+ = \bigcup_{\xi \in X^+} X'_\xi.$$

Then

$$(4.5) \quad \sum_{x \in X \setminus X^+} B_n(x, z) = \sum_{\xi \in X^+} \left(\sum_{x \in X'_\xi} B_n(x, z) \right),$$

therefore, we have

$$\begin{aligned}
(4.6) \quad \sum_{x \in X} B_n(x, z) &= \sum_{\xi \in X^+} \left(B_n(\xi, z) + \sum_{x \in X'_\xi} B_n(x, z) \right) \\
&= \sum_{\xi: B_n(\xi, 0) > 0} \left(B_n(\xi, z) + \sum_{x \in X'_\xi} B_n(x, z) \right) + \sum_{\xi: B_n(\xi, 0) < 0} \left(B_n(\xi, z) + \sum_{x \in X'_\xi} B_n(x, z) \right) \\
&: \Sigma = \Sigma^+ + \Sigma^-,
\end{aligned}$$

where $\Sigma^\pm := \sum_{\xi: B_n(\xi, 0) \gtrless 0} \left(B_n(\xi, z) + \sum_{x \in X'_\xi} B_n(x, z) \right)$.

By Lemma 2.7 and Lemma 2.9, we get there exists a constant $C_1 > 0$ such that

$$(4.7) \quad \left[1 \mp C_1 \frac{\log n}{n} \right] \sigma_n^\pm \leq \Sigma^\pm \leq \left[1 \pm C_1 \frac{\log n}{n} \right] \sigma_n^\pm,$$

which is followed by

$$(4.8) \quad \left| \frac{\Sigma}{\sigma_n} - 1 \right| \leq C_1 \frac{|\sigma_n^-| + \sigma_n^+}{|\sigma_n|} \cdot \frac{\log n}{n}.$$

If $\xi \in X^+$, then $V(x_1) \cdots V(x_{\nu+1})$ is a monomial in q of degree $\frac{m_1}{2} \cdot (x_1 + \cdots + x_{\nu+1}) = m_1 \cdot n$. From Case 3 of Theorem 1.3, we have

$$(4.9) \quad \sigma_{2m} = \sum_{\xi \in X^+} B_{2m}(\xi, 0) = \frac{q^{2m_1 \cdot m}}{4^{2m-1} \cdot [(2m-1)!]^2} \cdot \prod_{k=1}^m \left(\left(\frac{b}{2} \right)^2 + 8c \left(k - \frac{1}{2} \right)^2 \right)$$

and

$$(4.10) \quad \sigma_{2m-1} = \sum_{\xi \in X^+} B_{2m-1}(\xi, 0) = \frac{q^{m_1(2m-1)}}{4^{2m-2} \cdot [(2m-2)!]^2} \cdot \frac{b}{2} \cdot \prod_{k=1}^{m-1} \left(\left(\frac{b}{2} \right)^2 + 8ck^2 \right).$$

Moreover, $\sigma_{2m} \neq 0$ when $\frac{b}{2} \neq 2\sqrt{-2c} \cdot (k - \frac{1}{2})$ and $\sigma_{2m-1} \neq 0$ when $\frac{b}{2} \neq 2\sqrt{-2c} \cdot k$, where $c < 0$. So

$$(4.11) \quad \frac{|\sigma_{2m}^-| + \sigma_{2m}^+}{|\sigma_{2m}|} = \frac{\prod_{k=1}^m \left(1 - \frac{b^2}{8c(2k-1)^2} \right)}{\prod_{k=1}^m \left| 1 + \frac{b^2}{8c(2k-1)^2} \right|} \leq \frac{\prod_{k=1}^\infty \left(1 - \frac{b^2}{8c(2k-1)^2} \right)}{\prod_{k=1}^\infty \left| 1 + \frac{b^2}{8c(2k-1)^2} \right|} = \left| \frac{\cosh\left(\frac{b\pi}{4\sqrt{-2c}}\right)}{\cos\left(\frac{b\pi}{4\sqrt{-2c}}\right)} \right|.$$

Similarly, we have

$$(4.12) \quad \frac{|\sigma_{2m-1}^-| + \sigma_{2m-1}^+}{|\sigma_{2m-1}|} \leq \left| \frac{\sinh\left(\frac{b\pi}{4\sqrt{-2c}}\right)}{\sin\left(\frac{b\pi}{4\sqrt{-2c}}\right)} \right|.$$

By (4.8), we obtain

$$(4.13) \quad \sum_{x \in X} B_{2m}(x, z) = \sigma_{2m} \left[1 + O\left(\frac{\log m}{m}\right) \right] = \left(\sum_{\xi \in X^+} B_{2m}(\xi, 0) \right) \left[1 + O\left(\frac{\log m}{m}\right) \right]$$

and

(4.14)

$$\sum_{x \in X} B_{2m-1}(x, z) = \sigma_{2m-1} \left[1 + O\left(\frac{\log m}{m}\right) \right] = \left(\sum_{\xi \in X^+} B_{2m-1}(\xi, 0) \right) \left[1 + O\left(\frac{\log m}{m}\right) \right].$$

Notice that

$$(4.15) \quad \cos\left(\frac{b\pi}{4\sqrt{-2c}}\right) = \prod_{k=1}^{\infty} \left(1 + \frac{b^2}{8c(2k-1)^2} \right)$$

and

$$(4.16) \quad \sin\left(\frac{b\pi}{4\sqrt{-2c}}\right) = \frac{b\pi}{4\sqrt{-2c}} \prod_{k=1}^{\infty} \left(1 + \frac{b^2}{8c(2k)^2} \right),$$

then

$$(4.17) \quad \cos\left(\frac{b\pi}{4\sqrt{-2c}}\right) = \prod_{k=1}^m \left(1 + \frac{b^2}{8c(2k-1)^2} \right) \left[1 + O\left(\frac{1}{m}\right) \right]$$

and

$$(4.18) \quad \sin\left(\frac{b\pi}{4\sqrt{-2c}}\right) = \frac{b\pi}{4\sqrt{-2c}} \prod_{k=1}^{m-1} \left(1 + \frac{b^2}{8c(2k)^2} \right) \left[1 + O\left(\frac{1}{m}\right) \right].$$

Hence,

$$(4.19) \quad \sum_{\xi \in X^+} B_{2m}(\xi, 0) = \frac{q^{2m_1 \cdot m} \cdot (-1)^m \cdot c^m}{2^{3m-2} \cdot [(2m-2)!!]^2} \cdot \cos\left(\frac{b\pi}{4\sqrt{-2c}}\right) \cdot \left[1 + O\left(\frac{1}{m}\right) \right]$$

and

(4.20)

$$\sum_{\xi \in X^+} B_{2m-1}(\xi, 0) = \frac{q^{m_1(2m-1)} \cdot (-1)^{m-1} \cdot c^{m-1} \cdot \sqrt{-2c}}{2^{3m-4} \cdot [(2m-3)!!]^2 \cdot \pi} \cdot \sin\left(\frac{b\pi}{4\sqrt{-2c}}\right) \cdot \left[1 + O\left(\frac{1}{m}\right) \right].$$

Combining (4.8), (4.11) and (4.12), we deduce

$$(4.21) \quad \begin{aligned} \sum_{x \in X} B_{2m}(x, z) &= \left(\sum_{\xi \in X^+} B_{2m}(\xi, 0) \right) \left[1 + O\left(\frac{\log m}{m}\right) \right] \\ &= \frac{q^{2m_1 \cdot m} \cdot (-1)^m \cdot c^m}{2^{3m-2} \cdot [(2m-2)!!]^2} \cdot \cos\left(\frac{b\pi}{4\sqrt{-2c}}\right) \cdot \left[1 + O\left(\frac{\log m}{m}\right) \right] \end{aligned}$$

and

(4.22)

$$\begin{aligned} \sum_{x \in X} B_{2m-1}(x, z) &= \left(\sum_{\xi \in X^+} B_{2m-1}(\xi, 0) \right) \left[1 + O\left(\frac{\log m}{m}\right) \right] \\ &= \frac{q^{m_1(2m-1)} \cdot (-1)^{m-1} \cdot c^{m-1} \cdot \sqrt{-2c}}{2^{3m-4} \cdot [(2m-3)!!]^2 \cdot \pi} \cdot \sin\left(\frac{b\pi}{4\sqrt{-2c}}\right) \cdot \left[1 + O\left(\frac{\log m}{m}\right) \right]. \end{aligned}$$

Therefore,

$$(4.23) \quad \gamma_{2m} = \frac{q^{2m_1 \cdot m} \cdot |c|^m}{2^{3m-3} \cdot [(2m-2)!!]^2} \cdot \left| \cos\left(\frac{b\pi}{4\sqrt{-2c}}\right) \right| \cdot \left[1 + O\left(\frac{\log m}{m}\right) \right]$$

and

$$(4.24) \quad \gamma_{2m-1} = \frac{q^{m_1(2m-1)} \cdot |c|^{m-1} \cdot \sqrt{-2c}}{2^{3m-5} \cdot [(2m-3)!!]^2 \cdot \pi} \cdot \left| \sin \left(\frac{b\pi}{4\sqrt{-2c}} \right) \right| \cdot \left[1 + O \left(\frac{\log m}{m} \right) \right].$$

REFERENCES

1. P. Djakov and B. Mityagin, *Simple and double eigenvalues of the Hill operator with a two-term potential*, Journal of Approximation Theory **135**, 70-104 (2005).
2. P. Djakov and B. Mityagin, *Asymptotics of instability zones of the Hill operator with a two term potential*, Journal of Functional Analysis **242**, 157-194 (2007).
3. M.S.P. Eastham, *The spectral theory of periodic differential equations*, Scottish academic press, 1973.
4. E. Harrell, *On the effect of the boundary conditions on the eigenvalues of ordinary differential equations*, in: Suppl. Amer. J. Math., dedicated to P. Hartman, Johns Hopkins Univ. Press, Baltimore, MD, 139-150 (1981).
5. E. L. Ince, *A proof of the impossibility of the coexistence of two Mathieu functions*, Proc. Cambridge Philos. Soc. **21** 117-120 (1922).
6. D. M. Levy and J. B. Keller, *Instability Intervals of Hill's Equation*, Communications on Pure and Applied Mathematics, **XVI** 469-476 (1963).
7. W. Magnus and Winkler, *Hill's equation*, InterScience publisher, 1966.
8. H. Volkmer, *Coexistence of periodic solutions of Ince's equation*, Analysis(Munich) **23**, 97-105 (2003).

ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA.

E-mail address: `1xd@amss.ac.cn`